

the response to PEG-IFN and RBV combination therapy for CH-C.²⁶⁻²⁸ In Japanese patients, in particular, a single nucleotide polymorphism (SNP rs8099917) near the IL28B gene on chromosome 19 was shown to be strongly associated with non-virological response in genotypes 1.²⁷ However, in genotype 2, this association was weak especially in genotype 2a.^{29,30} A recent paper from Japan reported that CH-C patients infected with genotype 1b who showed hepatic steatosis >10% were likely to harbor IL28B minor allele (TG or GG genotype).¹⁸ In our preliminary retrospective data, two out of five (40%) non-SVR patients had the TG genotype and ten out of ten (100%) of SVR patients had the TT genotype (Table 1). The relationship between hepatic steatosis and IL28B polymorphism may be an interesting subject for further investigation.

In conclusion, CH-C patients infected with genotype 2 with hepatic steatosis >10% showed resistance to PEG-IFN α 2b and RBV combination therapy. Considering cost minimization, a variable-duration regimen is recommended.³¹ We advocate that CH-C patients infected with genotype 2 should be evaluated for hepatic steatosis by biopsied liver specimens before starting PEG-IFN and RBV combination therapy. In combination with other factors such as mutations in HCV and IL28B gene polymorphism,³⁰ evaluation of hepatic steatosis will enable us to more accurately predict the outcome of CH-C patients infected with genotype 2 in PEG-IFN and RBV combination therapy.

ACKNOWLEDGMENTS

WE ARE GRATEFUL to the members of the Hepatology Research Unit in Molecular Gastroenterology and Hepatology, Kyoto Prefectural University of Medicine, Kyoto, Japan, and related hospitals for their contribution to this work.

REFERENCES

- Manns MP, McHutchinson JG, Gordon SC *et al.* Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomised trial. *Lancet* 2001; 358: 958–65.
- Fried MW, Shiffman ML, Reddy KR *et al.* Peginterferon alfa2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 2002; 347: 975–82.
- Akuta N, Suzuki F, Tsubota A *et al.* Efficacy of interferon monotherapy to 394 consecutive naive cases infected with hepatitis C virus genotype 2a in Japan: therapy efficacy as consequence of tripartite interaction of viral, host and interferon treatment-related factors. *J Hepatol* 2002; 37: 831–6.
- Akuta N, Suzuki F, Suzuki Y *et al.* Hepatocyte steatosis is an important predictor of response to interferon (IFN) monotherapy in Japanese patients infected with HCV genotype 2a: virological features of IFN-resistant cases with hepatocyte steatosis. *J Med Virol* 2005; 75: 550–8.
- Poynard T, Ratziu V, McHutchison J *et al.* Effect of treatment with peginterferon or interferon alfa-2b and ribavirin on steatosis in patients infected with hepatitis C. *Hepatology* 2003; 38: 75–85.
- Poustchi H, Negro F, Hui J *et al.* Insulin resistance and response to therapy in patients with chronic hepatitis C virus genotype 2 and 3. *J Hepatol* 2008; 48: 28–34.
- Rodriguez-Torres M, Govindarajan S, Diago M *et al.* Hepatic steatosis in patients with chronic hepatitis C virus genotype 2 or 3 does not affect viral response in patients treated with peginterferon alpha-2a (40KD) (PEGASYS) plus ribavirin (COPEGUS) for 16 or 24 weeks. *Liver Int* 2009; 29: 237–41.
- Mangia A, Santoro R, Minerva N *et al.* Peginterferon alfa-2b and ribavirin for 12 vs. 24 weeks in HCV genotype 2 or 3. *N Engl J Med* 2005; 352: 2609–17.
- von Wagner M, Huber M, Berg T *et al.* Peginterferon-alpha-2a (40KD) and ribavirin for 16 or 24 weeks in patients with genotype 2 or 3 chronic hepatitis C. *Gastroenterology* 2005; 129: 522–7.
- Shiffman ML, Suter F, Bacon BR *et al.* Peginterferon alfa-2a and ribavirin for 16 or 24 weeks in HCV genotype 2 or 3. *N Engl J Med* 2007; 357: 124–34.
- Dalgard O, Bjørø K, Ring-Larsen H *et al.* Pegylated interferon alfa and ribavirin for 14 versus 24 weeks in patients with hepatitis C virus genotype 2 or 3 and rapid virological response. *Hepatology* 2008; 47: 35–42.
- Lagging M, Langeland N, Pedersen C *et al.* Randomized comparison of 12 or 24 weeks of peginterferon alpha-2a and ribavirin in chronic hepatitis C virus genotype 2/3 infection. *Hepatology* 2008; 47: 1837–45.
- Yu ML, Dai CY, Huang JF *et al.* A randomised study of peginterferon and ribavirin for 16 versus 24 weeks in patients with genotype 2 chronic hepatitis C. *Gut* 2007; 56: 553–9.
- Mangia A, Minerva N, Bacca D *et al.* Determinants of relapse after a short (12 weeks) course of antiviral therapy and re-treatment efficacy of a prolonged course in patients with chronic hepatitis C virus genotype 2 or 3 infection. *Hepatology* 2009; 49: 358–63.
- Rodriguez Torres M, Rios Bedoya CF, Ortiz-Lasanta G, Purcell-Arevalo D, Marxuach-Cuetara A, Jimenez-Rivera J. Weight affect relapse rates in latinos with genotype 2/3 chronic hepatitis C (CHC) treated with peg IFN alfa-2a (Pegasys) 180mcg/week and 800 mg daily of ribavirin for 24 weeks. *J Med Virol* 2008; 80: 1576–80.

- 16 Ichida F, Tsuji T, Omata M *et al.* New Inuyama classification; new criteria for histological assessment of chronic hepatitis. *Int Hepatol Commun* 1996; 6: 112–19.
- 17 Inoue Y, Hiramatsu N, Oze T *et al.* Factors affecting efficacy in patients with genotype 2 chronic hepatitis C treated by pegylated interferon alpha-2b and ribavirin: reducing drug doses has no impact on rapid and sustained virological responses. *J Viral Hepat* 2010; 17: 336–44.
- 18 Kurosaki M, Tanaka Y, Nishida N *et al.* Pre-treatment prediction of response to pegylated-interferon plus ribavirin for chronic hepatitis C using genetic polymorphism in IL28B and viral factors. *J Hepatol* 2011; 54: 439–48.
- 19 Leandro G, Mangia A, Hui J *et al.* Relationship between steatosis, inflammation, and fibrosis in chronic hepatitis C: a meta-analysis of individual patient data. *Gastroenterology* 2006; 130: 1636–42.
- 20 Patton HM, Patel K, Behling C *et al.* The impact of steatosis on disease progression and early and sustained treatment response in chronic hepatitis C patients. *J Hepatol* 2004; 40: 484–90.
- 21 Lok AS, Everhart JE, Chung RT *et al.* Hepatic steatosis in hepatitis C: comparison of diabetic and nondiabetic patients in the hepatitis C antiviral long-term treatment against cirrhosis trial. *Clin Gastroenterol Hepatol* 2007; 5: 245–54.
- 22 Lerat H, Kammoun HL, Hainault I *et al.* Hepatitis C virus proteins induce lipogenesis and defective triglyceride secretion in transgenic mice. *J Biol Chem* 2009; 284: 33466–74.
- 23 Vanni E, Abate ML, Gentilcore E *et al.* Site and mechanisms of insulin resistance in nonobese, nondiabetic patients with chronic hepatitis. *Hepatology* 2009; 50: 697–706.
- 24 Miyaaki H, Ichikawa T, Nakao K *et al.* Predictive value of suppressor of cytokine signal 3(SOCS3) in the outcome of interferon therapy in chronic hepatitis C. *Hepatol Res* 2009; 39: 850–5.
- 25 Tillmann HL. Does the presence of hepatic steatosis influence the efficacy of antiviral treatment in patients with hepatitis C? *Nat Clin Pract* 2007; 4: 548–49.
- 26 Ge D, Fellay J, Thompson AJ *et al.* Genetic variation in IL28B predict hepatitis C treatment-induced viral clearance. *Nature* 2009; 461: 399–401.
- 27 Tanaka Y, Nishida N, Sugiyama M *et al.* Genome-wide association of IL28B with response to pegylated interferon-alpha and ribavirin therapy for chronic hepatitis C. *Nat Genet* 2009; 41: 1105–9.
- 28 Suppiah V, Moldovan M, Ahlenstiel G *et al.* IL28B is associated with response to chronic hepatitis C interferon-alpha and ribavirin therapy. *Nat Genet* 2009; 41: 1100–4.
- 29 Sakamoto N, Nakagawa M, Tanaka Y *et al.* Association of IL28B variants with response to pegylated-interferon alpha plus ribavirin combination therapy reveals intersubgenotypic differences between genotype 2a and 2b. *J Med Virol* 2011; 83: 871–8.
- 30 Kawaoka T, Hayes CN, Ohishi W *et al.* Predictive value of the IL28B polymorphism on the effect of interferon therapy in chronic hepatitis C patients with genotype 2a and 2b. *J Hepatol* 2011; 54: 408–14.
- 31 De Compadri P, Koleva D, Mangia A, Motterlini L, Garattini L. Cost minimisation analysis of 12 or 24 weeks of peginterferon alfa2b + ribavirin for hepatitis C virus. *J Med Econ* 2008; 11: 151–63.

Characterization of naturally occurring protease inhibitor-resistance mutations in genotype 1b hepatitis C virus patients

Hiroko Shindo · Shinya Maekawa · Kazuki Komase · Ryota Sueki · Mika Miura · Makoto Kadokura · Kuniaki Shindo · Fumitake Amemiya · Takatoshi Kitamura · Yasuhiro Nakayama · Taisuke Inoue · Minoru Sakamoto · Shun-ichi Okada · Yasuhiro Asahina · Namiki Izumi · Masao Honda · Shuichi Kaneko · Nobuyuki Enomoto

Received: 21 April 2011 / Accepted: 22 July 2011
© Asian Pacific Association for the Study of the Liver 2011

Abstract

Background and aims Protease inhibitor (PI)-resistant hepatitis C virus (HCV) variants may be present in substantial numbers in PI-untreated patients according to recent reports. However, influence of these viruses in the clinical course of chronic hepatitis C has not been well characterized.

Methods The dominant HCV nonstructural 3 (NS3) amino acid sequences were determined in 261 HCV genotype 1b-infected Japanese patients before pegylated interferon plus ribavirin (PEG-IFN/RBV) therapy, and investigated the patients' clinical characteristics as well as treatment responses including sustained virological response (SVR) rate. HCV-NS3 sequences were also determined in 39 non-SVR patients after completion of the therapy.

Results Four single mutations (T54S, Q80K, I153V, and D168E) known to confer PI resistance were found in 35 of 261 patients (13.4%), and double mutations (I153V plus

T54S/D168E) were found in 6 patients (2.3%). Responses to PEG-IFN/RBV therapy did not differ between patients with and without PI-resistance mutations (mutation group, SVR 48%; wild-type group, SVR 40%; $P = 0.38$). On the other hand, two mutations appeared in two non-SVR patients after PEG-IFN/RBV therapy (I153V and E168D, 5.1%).

Conclusions PI-resistance-associated NS3 mutations exist in a substantial proportion of untreated HCV-1b-infected patients. The impact of these mutations in the treatment of PIs is unclear, but clinicians should pay attention to avoid further development of PI resistance.

Keywords HCV · Protease inhibitor · Naturally occurring viral resistance mutations

Introduction

Hepatitis C virus (HCV) infects more than 170 million persons worldwide and thus represents a global health problem. At least 130 million infected individuals are chronic carriers of HCV and are at significant risk of developing liver cirrhosis and hepatocellular carcinoma [1]. The current standard treatment with pegylated interferon plus ribavirin (PEG-IFN/RBV) is complicated by frequent adverse reactions, and a sustained virologic response (SVR) can be achieved only in 50% of patients infected with the most prevalent genotype 1 [2]. In Japan, since 70% of patients are infected with intractable genotype 1b HCV, more effective treatments are urgently required.

A promising approach is the development of specifically targeted antiviral therapies for hepatitis C (STAT-C). HCV-specific protease inhibitors (PIs) target an essential step in HCV replication by blocking the nonstructural 3/4A (NS3/4A) protease-dependent cleavage of the HCV polyprotein

Electronic supplementary material The online version of this article (doi:10.1007/s12072-011-9306-7) contains supplementary material, which is available to authorized users.

H. Shindo · S. Maekawa (✉) · K. Komase · R. Sueki · M. Miura · M. Kadokura · K. Shindo · F. Amemiya · T. Kitamura · Y. Nakayama · T. Inoue · M. Sakamoto · S. Okada · N. Enomoto

First Department of Internal Medicine, University of Yamanashi, 1110, Shimokato, Chuo, Yamanashi 409-3898, Japan
e-mail: maekawa@yamanashi.ac.jp

Y. Asahina · N. Izumi
Division of Gastroenterology and Hepatology,
Musashino Red Cross Hospital, Tokyo, Japan

M. Honda · S. Kaneko
Department of Gastroenterology, Kanazawa University Graduate
School of Medicine, Kanazawa, Japan

[1]. Among these NS3/4A PIs, telaprevir, boceprevir, SCH446211, danoprevir (ITMN-191), naldaprevir (SCH900 518), and TMC435 are now under clinical trials [1, 3–7]. In PROVE1 and PROVE2 studies [3, 4] undertaken in North America and Europe, the SVR rate was favorable (67 and 69%, respectively) in a triple therapy regimen including telaprevir. In addition, some studies have suggested that shortening of treatment duration may be possible for patients who achieve a rapid virologic response (RVR) [8, 9].

However the sole use of STAT-C drugs, such as PIs, promotes production and selection of drug-resistant variants in patients experiencing viral rebound during treatment [3, 10, 11] as well as in HCV replicon experiments [11, 12]. Therefore, these drugs should be used in combination with the PEG-IFN/RBV to prevent the appearance of drug-resistant variants. However, Kuntzen et al. [13] demonstrated the presence of these drug-resistant variants in high frequencies (8.6–16.2%) by population-based sequencing in patients not treated with the drugs [1, 13]. Gaudieri et al. [14] have suggested that regions of NS3 protease and NS5B polymerase are likely to be under HLA immune pressure and therapeutic selection, and that drug-resistant variants may occur naturally to escape the immune system. These observations seem quite astonishing and troubling, since a substantial number of patients may not respond to the new therapies such as STAT-C drugs.

In the present study, to assess the prevalence of NS3 mutations conferring PI resistance in HCV genotype 1b-infected Japanese patients who had not been previously treated with PIs, as well as to assess the influence of those mutations in response to PEG-IFN/RBV therapy, the dominant HCV-NS3 sequences were determined in 261 HCV-1b patients before starting the PEG-IFN/RBV therapy.

Methods

Patients

Serum samples were acquired from 261 HCV genotype 1b-infected adult Japanese patients before combination therapy with PEG-IFN (PEGINTRON[®], Schering-Plough, Tokyo, Japan) plus RBV (REBETOL[®], Schering-Plough) between 2004 and 2008 at the University of Yamanashi, Musashino Red Cross Hospital and Kanazawa University. The therapy was administered according to the standard PEG-IFN/RBV treatment protocol established for Japanese patients by a hepatitis study group of the Ministry of Health, Labor, and Welfare, Japan. Specifically, the patients were subcutaneously administered PEG-IFN α -2b, 1.5 μ g/kg body weight, once weekly and RBV 600–800 mg daily per os for 48 weeks. These patients were not infected with human immunodeficiency virus (HIV). The study was

approved by the ethics committees of all participating universities and the hospital, and the protocol conformed to the ethical guidelines of the 1975 Declaration of Helsinki as reflected in a priori approval by the Institutional Review Board at Massachusetts General Hospital. Written informed consent was obtained from each study participant.

Amplification and sequencing of full-length HCV genomes

Viral loads were determined using the Amplicor HCV RNA kit, version 2.0 (Roche Diagnostics, Tokyo, Japan) or the Cobas TaqMan test (Roche Diagnostics). HCV RNA was extracted from pretreatment serum samples by the AGPC method using Isogen (Wako, Osaka, Japan) according to the manufacturer's protocol. Complementary DNA was synthesised using Superscript II (Invitrogen, Tokyo, Japan) and random primers (Invitrogen), and then amplified by two-step nested PCR using the primers listed in Supplementary Table 1. All samples were initially denatured at 95°C for 7 min, followed by 40 cycles of amplification with denaturation at 95°C for 15 s, annealing at 55°C for 15 s, and extension at 72°C for 45 s using the BD Advantage[™] 2 PCR Enzyme system (BD Biosciences Clontech, CA, USA). PCR amplicons were directly sequenced using BigDye Terminator version 3.1 (ABI, Tokyo, Japan) and universal M13 forward/reverse primers using an ABI prism 3130 sequencer (ABI).

Sequence alignment and analysis

Sequences were determined in both directions, particularly for the ambiguous stretches, were assembled using the Vector NTI software (Invitrogen), and base-calling errors were corrected following the inspection of chromatograms. If mixed bases were detected as two different chromatogram peaks at the same residue, only the dominant base was called after evaluation of all overlapping fragments. A consensus sequence was generated from the alignment on the basis of the most common amino acid at each site.

Determination of PI resistance mutations

Multiple viral NS3 mutations were observed in amino acid positions reported to confer PI resistance among 261 patients: V36, Q41, F43, T54, V55, Q80, R109, I153, R155, A156, D168, V170, and M175. NS3 amino acid mutations with proven PI resistance in previously published studies (Table 1) were designated as resistance proven mutations (e.g., V36M/A). Mutations in the PI-resistance site not known to confer drug resistance were designated resistance unproven mutations (e.g., V36I). Patients were allocated to two groups according to the presence of PI-resistance

mutations (including resistance unproven mutations), and clinical characteristics including HCV RNA levels and responses to PEG-IFN/RBV therapy were compared. To assess the influence of PEG-IFN/RBV therapy on NS3 mutational status, posttreatment HCV-NS3 sequences in 39 of 58 non-SVR patients were also examined.

Statistical analysis

Statistical differences in the data, including all available patients' demographic, biochemic, hematologic, and virologic data such as sequence variation factors, were determined among the various groups by Student's *t* test or Mann-Whitney *U* test for numerical variables and Fisher's exact probability test for categorical variables.

Results

Prevalence of dominant PI-resistance-associated nonstructural 3 mutations in untreated patients

Figure 1 shows the frequency of substitutions in 261 patients for each of 181 NS3 protease amino acid residues

compared to the consensus sequence. A total of 41 resistance proven mutations were detected in 35 (13.4%) patients: T54S (14 patients, 5.4%), Q80K (1 patient, 0.4%), I153V (22 patients, 8.4%), D168E (4 patients, 1.5%), T54S plus I153V double mutation (4 patients, 1.5%), and I153V plus D168E double mutation (2 patients, 0.8%). The mutation number increased to 54 in 47 (18.0%) patients when resistance unproven mutations were included: V36I (2 patients, 0.8%), I153L (11 patients, 4.2%), and I153V plus V36I double mutation (2 patients, 1.5%). Double mutations were found in 7 patients (2.7%) (Table 1). Q80L was observed in 47 (18%) patients but these were excluded from consideration because a previous study demonstrated that this mutation does not confer resistance [15]. All mutations observed in this study would confer low- to moderate-level PI resistance according to previous studies [6, 15–19]. No mutations conferring high-level resistance such as R155 or A156 [11, 17, 19–22] were observed.

Clinical characteristics of patients with PI-resistance mutations

Table 2 presents the characteristics of patients classified according to the presence of PI-resistance mutations

Table 1 Prevalence of PI-resistance-associated NS3 mutations

Drug-resistance mutations described in the literature				References	Detected resistance mutations Genotype 1b (<i>N</i> = 261), (%)
NS3 residue	Resistance mutations	Drugs			
V36	A, M, L, G, C	Telaprevir, Boceprevir	[1, 3, 4, 10, 11, 19, 31, 37]	I × 2 (0.8)	
Q41	R	ITMN-191, Boceprevir	[19]		
F43	S, C	ITMN-191, Boceprevir, Telaprevir, TMC435	[15, 19]	S × 14 (5.4)	
T54	A, S	Telaprevir, Boceprevir, SCH900518	[1, 3, 10, 11, 19, 20, 31, 38]		
V55	A	Boceprevir	[1]	K × 1 (0.4)	
Q80	R, K	TMC435	[6, 15]		
R109	K	SCH446211	[17]	V × 22 (8.4), L × 11 (4.2)	
I153	V	SCH446211	[17]		
R155	K, T, I, M, G, L, S, Q	Telaprevir, Boceprevir, ITMN-191, BILN2061, TMC435	[1, 3, 4, 6, 10, 11, 15, 19, 20]	E × 4 (1.5)	
A156	S, T, V, I, G	Telaprevir, Boceprevir, ITMN-191, BILN2061, SCH446211, TMC435, SCH900518	[1, 3, 4, 10, 11, 15, 17, 19, 20, 38]		
D168	A, V, E, N, T, H	BILN2061, ITMN-191, TMC435	[6, 15, 20]	35 (13.4)	
V170	A	Telaprevir, Boceprevir	[1, 19, 20]		
M175	L	Boceprevir	[39]	47 (18.0)	
Total number (%) of patients with resistance proven mutations					
Total number (%) of patients with resistance proven and unproven mutations					

Amino acid mutations conferring PI resistance in the literatures and those observed in PI-treatment-naive patients in this study are indicated. Bold indicates resistance proven mutations, and the others indicate resistance unproven mutations

Double mutations found were as follows: V36I and I153V × 1, T54S and I153V × 4, I153V and D168E × 2

(including resistance unproven mutations). Age, sex ratio, body mass index, alanine aminotransferase (ALT) levels, serum albumin, platelet count, and fibrosis stage did not differ between the NS3 mutation and wild-type groups. No significant difference was observed between the two groups in the parameters of PEG-IFN/RBV treatment response, HCV sequence variations in interferon sensitivity determining region (ISDR), Core 70, interferon plus ribavirin resistance-determining region (IRRDR), or interleukin 28B (IL28B) single nucleotide polymorphism (SNP) (rs8099917; T/G and G/G vs. T/T) [23–30]. These clinical variables were also compared between the mutation group defined as resistance proven mutations and the wild-type group, but no notable differences were observed.

Unimpaired in vivo fitness of viral strains with resistance mutations

Because most PI-resistance mutations described till date have been associated with reduced replicative capacity of varying degrees [1, 10, 11, 13, 17, 20–22, 31, 32], we examined viral replication levels in patients with drug-resistance mutations (Fig. 2). The estimated *P* value indicated no significant difference between the mutation (median 1,500 KIU/ml) and wild-type (median 1,800 KIU/ml) groups (*P* = 0.69). The results indicate that drug-resistant HCVs were not necessarily impaired in their ability to replicate in vivo. However, patients with double mutations (*N* = 7) tended to have low viral loads (median 1,200 KIU/ml) (*P* = 0.09).

Resistance mutations and virologic response to PEG-IFN/RBV therapy

To determine the difference in virologic response to PEG-IFN/RBV therapy according to the PI mutation, frequency of HCV RNA levels below detection at 4 weeks (rapid viral response, RVR) and 12 weeks (complete early viral response, cEVR), and SVR rate (%) were investigated in

each group. The frequency of HCV RNA levels below detection at 4 and 12 weeks was 14 and 50%, respectively, in the mutation group, and was 11 and 46%, respectively, in the wild-type group. The SVR rate was 48 and 40% in the mutation and wild-type groups, respectively (*P* = 0.38). No significant difference was observed between the two groups in any of the indexes investigated (Table 2). The time-dependent viral clearance rate during PEG-IFN/RBV therapy was estimated in 133 patients including 25 patients (19%) with PI-resistance mutations available for the analysis. Kaplan–Meier analysis demonstrated that HCV clearance did not differ between the two groups with and without resistance mutations (log-rank test, *P* = 0.30) (Fig. 3).

Changes in nonstructural 3 amino acid sequence diversity during PEG-IFN/RBV therapy

Full-length NS3 protease sequences were determined in 39 non-SVR patients after PEG-IFN/RBV therapy. A single amino acid change at resistance-associated sites in two patients was observed. In one patient, isoleucine (Ile) at position 153 changed to valine (Val), and glutamic acid (Glu) changed to aspartic acid (Asp) at position 168 in the second (Fig. 4). At the nucleotide level, ATC (Ile) changed to GTC (Val) in I153V, and GAA (Glu) changed to GAC (Asp) in E168D. Both mutations were caused by one nucleotide exchange. No other changes were observed in the other 37 patients.

Discussion

Here we report that in 18% (47/261) HCV genotype 1b-infected patients who had not been previously treated with NS3 PIs, the viral genome contained dominant amino acid mutations within the NS3 PI-resistance sites. Even after confining the data to established PI-resistance mutations, the mutation rate was still significant in 13.4% (35/261). No clinical differences were observed between patients




Fig. 1 Frequency of polymorphic mutations for each of the 181 NS3 protease amino acid residues in 261 patients. *Arrowheads* indicate the sites reported to confer PI resistance. *Dark bars* denote the amino acid

variations at the resistant sites in this study. *80, we detected one resistant mutation (Q80K) and 47 (18%) non-resistant variations (Q80L) at the 80th residue

Table 2 Characteristics of patients with or without HCV genomes harboring drug-resistance mutations

Characteristics	Mutation type (<i>N</i> = 47)	Wild-type (<i>N</i> = 214)	<i>P</i> value
Patients' characteristics			
Age, median (range)	59 (46–72)	57 (19–77)	0.17
Male, no. (%)	26 (55)	112 (52)	0.70
BMI, median (range)	23.2 (15.5–31.9)	22.8 (16.1–31.9)	0.41
ALT IU/ml	81.3 ± 72.6 ^a	74.8 ± 51.9	0.93
Serum albumin g/dl	4.00 ± 0.37	4.01 ± 0.36	0.81
Platelet count × 10 ⁴ /μl	15.8 ± 4.3	14.5 ± 4.8	0.18
HCV RNA KIU/ml, median (range)	1,500 (58–6,310)	1800 (28–15,849)	0.69
Fibrosis, no. (%)			0.97
F0	0 (0)	7 (3)	
F1	23 (50)	89 (42)	
F2	9 (20)	52 (24)	
F3	9 (20)	40 (19)	
F4	5 (11)	26 (12)	
IFN pre-treatment no. (%)	15/40 (38) ^b	66/172 (38)	1.00
IL28B (rs8099917) T/G or G/G no. (%)	6/20 (30)	19/67 (28)	1.00
Response to PEG-IFN/RBV therapy			
SVR total cases no. (%)	22/46 (48)	83/210 (40)	0.38
RVR in total cases no. (%)	6/44 (14)	22/195 (11)	0.83
cEVR in total cases no. (%)	22/44 (50)	92/200 (46)	0.75
SVR 48w treatment no. (%)	16/29 (55)	55/130 (42)	0.29
End of treatment response no. (%)	26/41 (63)	123/202 (61)	0.91
HCV genome sequence variation			
ISDR mutation ≤1 no. (%)	32/46 (70)	167/210 (80)	0.21
Core70 R no. (%)	26/44 (59)	136/210 (65)	0.56
IRRDR mutation >3 no. (%)	25/38 (66)	107/190 (56)	0.34

^a Mean ± SD^b Number/total number (%)

harboring viruses with and without these mutations. Moreover, no differences were observed in the responses of either group to PEG-IFN/RBV therapy.

Recent studies reported that significant number of patients who were never treated with PI possess viral sequences with PI-resistance-associated NS3 mutations. In these studies, the prevalence of PI-resistance mutations was determined to be 8.6–16.2% [13, 14], in HCV genotype 1- and 3-infected patients in European–American populations. These patients were often coinfecting with HIV. Analysis of the public HCV databases (EuHCVdb and Los Alamos) also reported the presence of naturally occurring PI-resistance-associated NS3 mutations in worldwide isolates [33]. However, *in vivo* and *in vitro* studies demonstrated that most of the mutations observed conferred only low- to moderate-level PI resistance [7, 13, 14, 34, 35]. Regarding viral fitness, PI-resistant HCVs show lower fitness at varying degrees as revealed by *in vitro* studies [1, 10, 11, 17, 20–22, 31, 32], but HCV RNA levels in a clinical study did not differ significantly. The response to PEG-IFN/RBV therapy was almost comparable to that in HCV-infected patients without PI-resistance mutations either in HCV replicon experiments or in a clinical study of small number of treated patients [34].

The prevalence of 13.4% for PI-resistance-proven patients observed in the present study was almost comparable to the results of previous studies. Although HIV is known to increase HCV replication in coinfection with HCV [36], and HIV patients are often treated with the HIV-specific PIs, the HIV infection might not affect the natural occurrence of HCV-specific PI-resistance mutations since our studied patients were all proven to be free from coinfection with HIV infection. As shown in Table 1 and Fig. 1, I153 V (22/261, 8.4%), T54S (14/261, 5.4%), and D168E (4/261, 1.5%) were among the most prevalent PI-resistance-proven mutations in the present study. The most frequent mutation detected in our study I153V was reported to appear secondarily to the occurrence of R109K mutations in a HCV replicon system [17]. Although the role of this mutation is not understood, the I153V mutation on its own conferred SCH446211 resistance to the HCV replicon to a lesser degree [17]. Interestingly, I153V was often found in double mutations in our study, as shown in Fig. 2. This suggests analogy between *in vitro* and *in vivo* data. T54S and D168E, the other frequent mutations, have been also reported to occur as single dominant mutations in previous *in vitro* or *in vivo* studies in HCV genotype 1

Fig. 2 In vivo fitness of HCV with PI-resistance-associated NS3 mutations. HCV RNA levels were compared between patients with and without NS3 PI-resistance-associated mutations (a) and between patients with each resistance mutation (b). The estimated *P* value (Mann–Whitney *U* test) indicates no significant difference between the wild-type and other groups (wild-type vs. mutation type, wild-type vs. single mutation type, and wild-type vs. double mutation type). (Wild-type, *N* = 214; mutation type, *N* = 47; single mutation type, *N* = 40; double mutation type, *N* = 7; V36I, *N* = 2; T54S, *N* = 14; Q80K, *N* = 1; I153L, *N* = 11; I153V, *N* = 22; D168E, *N* = 4; E176A, *N* = 1; V36I + I153V, *N* = 1; T54S + I153V, *N* = 4, and I153V + D168E, *N* = 2)





Fig. 3 Comparison of virologic response to PEG-IFN/RBV therapy between HCV-infected patients with and without PI-resistance-associated NS3 mutations. Time-dependent HCV clearance rate analysis was based on serum HCV RNA positivity during PEG-IFN/RBV therapy for HCV isolates with resistance mutations or wild-


type sequences. A total of 133 patients for whom the limit of viral genome detection could be determined were analyzed. Among this group, NS3 mutations were detected in 25 patients (19%). The estimated *P* value (log-rank test) shows no significant difference between the two groups (*P* = 0.30)

infections showing moderate degrees of resistance [16, 18, 19].

Most PI-resistance mutations described to date have been associated with varying degrees of reduced replicative

capacity [10, 11, 17, 20–22, 31, 32]. In the present study, HCV RNA levels of those patients with low- to moderate-level resistance mutations were similar to those in patients in the wild-type groups, suggesting that in vitro viral fitness

Fig. 4 Appearance of PI-resistance-associated NS3 mutations during the PEG-IFN/RBV therapy. Chromatograms show part of the HCV NS3 sequence demonstrating PI-resistance mutations in two patients receiving therapy. **a** Site 153 isoleucine (Ile) (ATC) changed to valine (Val) (GTC), **b** Site 168 glutamic acid (Glu) (GAA) changed to aspartic acid (Asp) (GAC)



does not necessarily reflect *in vivo* viral fitness. This, however, does not rule out the possibility that some unknown compensatory viral mutations might have resulted in upregulation of reduced viral fitness. Interestingly, although the replicative capacity conferred by a single mutation seemed to be the same, the HCV RNA levels of double mutations were frequently low, suggesting that double mutations might weaken viral fitness.

In previous studies, clinical characteristics representing the state of liver disease other than HCV RNA levels were not studied in patients with PI-resistance mutations. In this study, we show that those clinical characteristics did not differ according to the presence of viral NS3 mutations. As shown in Table 2, age, sex ratio, fibrosis stage, ALT levels, serum albumin, platelet count, and past history of IFN pretreatment did not differ according to the presence of NS3 mutations. These results suggest that NS3 mutations occur independently of disease progression. Moreover, no evident differences were observed between viral and host factors known to affect IFN-based treatment responses. However, viral amino acid variations in the core and NS5A or the allelic frequency of IL28B SNPs, which were recently reported for the close relationship of responses to PEG-IFN/RBV therapy, did not differ between the two groups.

A significant outcome of the present study is the demonstration that PI-resistance mutations might not affect responses to PEG-IFN/RBV therapy. Previous *in vitro* studies demonstrated that HCV replicons harboring PI-resistance mutations were also sensitive to IFN treatment [31]. In addition, recent clinical studies also indicated that PI-resistance mutations were sensitive to the PEG-IFN/RBV [10, 34]. However, our analysis was more comprehensive because viral and host factors that contribute to treatment responses were simultaneously analyzed. A unique aspect of the present study is that we investigated the influence of the PEG-IFN/RBV treatment on the occurrence of new PI mutations by direct nucleotide sequencing, and were able to show that the PEG-IFN/RBV might not induce amino acid mutations.

Will the pre-existence of naturally occurring PI-resistance mutations have an influence on future treatment of HCV infections? Since new PIs are on the verge of clinical use, all clinicians should bear in mind the substantial numbers of HCV-infected patients with PI-resistance mutations. Although the degree of resistance is considered to be low or moderate in untreated patients, weak resistance might progress to more potent resistance with additional mutations, when PIs become widely used. Therefore, all clinicians need to be sufficiently prepared for the possibility of later onset of PI-resistance mutations that confer greater drug resistance and concomitant poorer responses to therapy. In SPRINT-1 study, the lead-in therapy was associated with a modestly lower rate of breakthrough than with no lead in [7]. Considering that PEG-IFN/RBV was equally effective for PI-resistant viruses, sufficient “lead-in” therapy before the administration of PIs could be an option in the forthcoming triple therapy modality.

In conclusion, we demonstrate here that PI-resistance-associated NS3 mutations exist in a substantial proportion of untreated HCV-1b-infected patients. Although the degree of resistance might not be strong, clinicians will need to consider this upon the introduction of triple therapy.

Acknowledgements This work was supported in part by a grant-in-aid scientific research fund from the Ministry of Education, Science, Sports, and Culture [grant number 20390206] and in part by a grant-in-aid from the Ministry of Health, Labor, and Welfare of Japan [grant number H19-kanen-002].

References

1. Susser S, Welsch C, Wang Y, et al. Characterization of resistance to the protease inhibitor boceprevir in hepatitis C virus-infected patients. *Hepatology* 2009;50:1709–1718
2. Hadziyannis SJ, Sette H Jr, Morgan TR, et al. Peginterferon-alpha2a and ribavirin combination therapy in chronic hepatitis C: a randomized study of treatment duration and ribavirin dose. *Ann Intern Med* 2004;140:346–355

3. Hezode C, Forestier N, Dusheiko G, et al. Telaprevir and peginterferon with or without ribavirin for chronic HCV infection. *N Engl J Med* 2009;360:1839–1850
4. McHutchison JG, Everson GT, Gordon SC, et al. Telaprevir with peginterferon and ribavirin for chronic HCV genotype 1 infection. *N Engl J Med* 2009;360:1827–1838
5. McHutchison JG, Manns MP, Muir AJ, et al. Telaprevir for previously treated chronic HCV infection. *N Engl J Med* 2010;362:1292–1303
6. Reesink HW, Fanning GC, Farha KA, et al. Rapid HCV-RNA decline with once daily TMC435: a phase I study in healthy volunteers and hepatitis C patients. *Gastroenterology* 2010;138:913–921
7. Kwo PY, Lawitz EJ, McCone J, et al. Efficacy of boceprevir, an NS3 protease inhibitor, in combination with peginterferon alfa-2b and ribavirin in treatment-naïve patients with genotype 1 hepatitis C infection (SPRINT-1): an open-label, randomised, multicentre phase 2 trial. *Lancet* 2010;376:705–716
8. Forestier N, Reesink HW, Weegink CJ, et al. Antiviral activity of telaprevir (VX-950) and peginterferon alfa-2a in patients with hepatitis C. *Hepatology* 2007;46:640–648
9. Suzuki F, Akuta N, Suzuki Y, et al. Rapid loss of hepatitis C virus genotype 1b from serum in patients receiving a triple treatment with telaprevir (MP-424), pegylated interferon and ribavirin for 12 weeks. *Hepatol Res* 2009;39:1056–1063
10. Kieffer TL, Sarrazin C, Miller JS, et al. Telaprevir and pegylated interferon-alpha-2a inhibit wild-type and resistant genotype 1 hepatitis C virus replication in patients. *Hepatology* 2007;46:631–639
11. Sarrazin C, Kieffer TL, Bartels D, et al. Dynamic hepatitis C virus genotypic and phenotypic changes in patients treated with the protease inhibitor telaprevir. *Gastroenterology* 2007;132:1767–1777
12. Lin C, Lin K, Luong YP, et al. In vitro resistance studies of hepatitis C virus serine protease inhibitors, VX-950 and BILN 2061: structural analysis indicates different resistance mechanisms. *J Biol Chem* 2004;279:17508–17514
13. Kuntzen T, Timm J, Berical A, et al. Naturally occurring dominant resistance mutations to hepatitis C virus protease and polymerase inhibitors in treatment-naïve patients. *Hepatology* 2008;48:1769–1778
14. Gaudieri S, Rauch A, Pfafferoth K, et al. Hepatitis C virus drug resistance and immune-driven adaptations: relevance to new antiviral therapy. *Hepatology* 2009;49:1069–1082
15. Lenz O, Verbinen T, Lin TI, et al. (2010) In vitro resistance profile of the hepatitis C virus NS3/4A protease inhibitor TMC435. *Antimicrob Agents Chemother* 54:1878–1887
16. Kieffer TL, Kwong AD, Picchio GR. Viral resistance to specifically targeted antiviral therapies for hepatitis C (STAT-Cs). *J Antimicrob Chemother* 2010;65:202–212
17. Yi M, Tong X, Skelton A, et al. Mutations conferring resistance to SCH6, a novel hepatitis C virus NS3/4A protease inhibitor. Reduced RNA replication fitness and partial rescue by second-site mutations. *J Biol Chem* 2006;281:8205–8215
18. Welsch C, Domingues FS, Susser S, et al. Molecular basis of telaprevir resistance due to V36 and T54 mutations in the NS3-4A protease of the hepatitis C virus. *Genome Biol* 2008;9:R16
19. Tong X, Bogen S, Chase R, et al. Characterization of resistance mutations against HCV ketoamide protease inhibitors. *Antiviral Res* 2008;77:177–185
20. He Y, King MS, Kempf DJ, et al. Relative replication capacity and selective advantage profiles of protease inhibitor-resistant hepatitis C virus (HCV) NS3 protease mutants in the HCV genotype 1b replicon system. *Antimicrob Agents Chemother* 2008;52:1101–1110
21. Tong X, Chase R, Skelton A, Chen T, Wright-Minogue J, Malcolm BA. Identification and analysis of fitness of resistance mutations against the HCV protease inhibitor SCH 503034. *Antiviral Res* 2006;70:28–38
22. Lu L, Pilot-Matias TJ, Stewart KD, et al. Mutations conferring resistance to a potent hepatitis C virus serine protease inhibitor in vitro. *Antimicrob Agents Chemother* 2004;48:2260–2266
23. Akuta N, Suzuki F, Hirakawa M, et al. Amino acid substitutions in the hepatitis C virus core region of genotype 1b affect very early viral dynamics during treatment with telaprevir, peginterferon, and ribavirin. *J Med Virol* 2010;82:575–582
24. Ge D, Fellay J, Thompson AJ, et al. Genetic variation in IL28B predicts hepatitis C treatment-induced viral clearance. *Nature* 2009;461:399–401
25. Tanaka Y, Nishida N, Sugiyama M, et al. Genome-wide association of IL28B with response to pegylated interferon-alpha and ribavirin therapy for chronic hepatitis C. *Nat Genet* 2009;41:1105–1109
26. Suppiah V, Moldovan M, Ahlenstiel G, et al. IL28B is associated with response to chronic hepatitis C interferon-alpha and ribavirin therapy. *Nat Genet* 2009;41:1100–1104
27. El-Shamy A, Nagano-Fujii M, Sasase N, Imoto S, Kim SR, Hotta H. Sequence variation in hepatitis C virus nonstructural protein 5A predicts clinical outcome of pegylated interferon/ribavirin combination therapy. *Hepatology* 2008;48:38–47
28. Enomoto N, Sakuma I, Asahina Y, et al. Mutations in the non-structural protein 5A gene and response to interferon in patients with chronic hepatitis C virus 1b infection. *N Engl J Med* 1996;334:77–81
29. Akuta N, Suzuki F, Kawamura Y, et al. Predictive factors of early and sustained responses to peginterferon plus ribavirin combination therapy in Japanese patients infected with hepatitis C virus genotype 1b: amino acid substitutions in the core region and low-density lipoprotein cholesterol levels. *J Hepatol* 2007;46:403–410
30. Akuta N, Suzuki F, Sezaki H, et al. Predictive factors of virological non-response to interferon-ribavirin combination therapy for patients infected with hepatitis C virus of genotype 1b and high viral load. *J Med Virol* 2006;78:83–90
31. Zhou Y, Bartels DJ, Hanzelka BL, et al. Phenotypic characterization of resistant Val36 variants of hepatitis C virus NS3-4A serine protease. *Antimicrob Agents Chemother* 2008;52:110–120
32. Mo H, Lu L, Pilot-Matias T, et al. Mutations conferring resistance to a hepatitis C virus (HCV) RNA-dependent RNA polymerase inhibitor alone or in combination with an HCV serine protease inhibitor in vitro. *Antimicrob Agents Chemother* 2005;49:4305–4314
33. Lopez-Labrador FX, Moya A, Gonzalez-Candelas F. Mapping natural polymorphisms of hepatitis C virus NS3/4A protease and antiviral resistance to inhibitors in worldwide isolates. *Antivir Ther* 2008;13:481–494
34. Bartels DJ, Zhou Y, Zhang EZ, et al. Natural prevalence of hepatitis C virus variants with decreased sensitivity to NS3.4A protease inhibitors in treatment-naïve subjects. *J Infect Dis* 2008;198:800–807
35. Cubero M, Esteban JI, Otero T, et al. Naturally occurring NS3-protease-inhibitor resistant mutant A156T in the liver of an untreated chronic hepatitis C patient. *Virology* 2008;370:237–245
36. Soriano V, Puoti M, Sulkowski M, et al. Care of patients with hepatitis C and HIV co-infection. *Aids* 2004;18:1–12
37. Barbotte L, Ahmed-Belkacem A, Chevaliez S, et al. Characterization of V36c, a novel amino acid substitution conferring hepatitis C virus (HCV) resistance to telaprevir, a potent peptidomimetic inhibitor of HCV protease. *Antimicrob Agents Chemother* 2010;54:2681–2683

38. Tong X, Arasappan A, Bennett F, et al. Preclinical characterization of the antiviral activity of SCH 900518 (Narlaprevir), a novel mechanism-based inhibitor of hepatitis C virus NS3 protease. *Antimicrob Agents Chemother* 2010;54:2365–2370
39. Chase R, Skelton A, Xia E, et al. A novel HCV NS3 protease mutation selected by combination treatment of the protease inhibitor boceprevir and NS5B polymerase inhibitors. *Antiviral Res* 2009;84:178–184

Comparative Analysis of Various Tumor-Associated Antigen-Specific T-Cell Responses in Patients with Hepatocellular Carcinoma

Eishiro Mizukoshi, Yasunari Nakamoto, Kuniaki Arai, Tatsuya Yamashita, Akito Sakai, Yoshio Sakai, Takashi Kagaya, Taro Yamashita, Masao Honda, and Shuichi Kaneko

Many tumor-associated antigens (TAAs) recognized by cytotoxic T cells (CTLs) have been identified during the last two decades and some of them have been used in clinical trials. However, there are very few in the field of immunotherapy for hepatocellular carcinoma (HCC) because there have not been comparative data regarding CTL responses to various TAAs. In the present study, using 27 peptides derived from 14 different TAAs, we performed comparative analysis of various TAA-specific T-cell responses in 31 HCC patients to select useful antigens for immunotherapy and examined the factors that affect the immune responses to determine a strategy for more effective therapy. Twenty-four of 31 (77.4%) HCC patients showed positive responses to at least one TAA-derived peptide in enzyme-linked immunospot assay. The TAAs consisting of cyclophilin B, squamous cell carcinoma antigen recognized by T cells (SART) 2, SART3, p53, multidrug resistance-associated protein (MRP) 3, alpha-fetoprotein (AFP) and human telomerase reverse transcriptase (hTERT) were frequently recognized by T cells and these TAA-derived peptides were capable of generating peptide-specific CTLs in HCC patients, which suggested that these TAAs are immunogenic. HCC treatments enhanced TAA-specific immune responses with an increased number of memory T cells and induced *de novo* T-cell responses to lymphocyte-specific protein tyrosine kinase, human epidermal growth factor receptor type 2, p53, and hTERT. Blocking cytotoxic T-lymphocyte antigen-4 (CTLA-4) resulted in unmasking of TAA-specific immune responses by changing cytokine and chemokine profiles of peripheral blood mononuclear cells stimulated by TAA-derived peptides. **Conclusion:** Cyclophilin B, SART2, SART3, p53, MRP3, AFP, and hTERT were immunogenic targets for HCC immunotherapy. TAA-specific immunotherapy combined with HCC treatments and anti-CTLA-4 antibody has the possibility to produce stronger tumor-specific immune responses. (HEPATOLOGY 2011;53:1206-1216)

Hepatocellular carcinoma (HCC) is the most common primary malignancy of the liver and becoming an important public health concern.^{1,2} Although many kinds of treatments have

been performed for HCC, their effects are limited because the recurrence rate of HCC is very high; therefore, the development of new therapeutic options to prevent recurrence is necessary.^{3,4}

To protect against recurrence, tumor antigen-specific immunotherapy is an attractive strategy. Many tumor-associated antigens (TAAs) and their epitopes recognized by cytotoxic T cells (CTLs) have been identified during the last two decades and some of them have been used in clinical trials for several cancers.⁵⁻²¹ The epitopes have been under investigation for the treatment of cancer, with major clinical responses in some trials.^{22,23} With regard to immunotherapy for HCC, few kinds of TAAs and their epitopes have been used and only clinical data of α -fetoprotein (AFP) have been reported.^{24,25} In human trials targeting AFP, it is possible to raise an AFP-specific T-cell response using AFP-derived peptides, but this has shown little

Abbreviations: AFP, alpha-fetoprotein; CTL, cytotoxic T cell; ELISPOT, enzyme-linked immunospot; HCC, hepatocellular carcinoma; HCV, hepatitis C virus; HLA, human leukocyte antigen; hTERT, human telomerase reverse transcriptase; IFN, interferon; Lck, lymphocyte-specific protein tyrosine kinase; MRP, multidrug resistance-associated protein; PBMC, peripheral blood mononuclear cell; TAA, tumor-associated antigen.

From the Department of Gastroenterology, Graduate School of Medicine, Kanazawa University, Kanazawa, Ishikawa, Japan.

Received August 6, 2010; accepted December 14, 2010.

Address reprint requests to: Shuichi Kaneko, M.D., Department of Gastroenterology, Graduate School of Medicine, Kanazawa University, Kanazawa, Ishikawa 920-8641, Japan. E-mail: skaneko@m-kanazawa.jp; fax: 81-76-234-4250.

Copyright © 2011 by the American Association for the Study of Liver Diseases.

View this article online at wileyonlinelibrary.com.

DOI 10.1002/hep.24149

Potential conflict of interest: Nothing to report.

antitumor effect. On the other hand, immunotherapy trials using autologous tumor lysate or dendritic cells have shown statistically significant improvements in the risk of HCC recurrence and recurrence-free survival.²⁶ These reports suggest that tumor antigen-specific immunotherapy is effective to reduce the recurrence rate after HCC treatment; therefore, it is necessary to find immunogenic antigens or their epitopes to develop more effective immunotherapy.

In addition, in the field of molecular targeting therapies, developments of monoclonal antibodies targeting immunomodulatory molecules to enhance anti-tumor immunity are progressing and some of these are under clinical trial.²⁷ In particular, clinical data of anti-cytotoxic T-lymphocyte antigen-4 (anti-CTLA-4) antibody have shown durable objective response and stable disease in melanoma patients.²⁸

In the present study we performed comparative analysis of various TAA-specific T-cell responses in patients with HCC and examined the factors that affect the immune responses, including anti-CTLA-4 antibody. This approach offers useful information to select immunogenic TAAs and to develop a new strategy for HCC immunotherapy.

Patients and Methods

Patients and Laboratory Testing. In this study we examined 31 human leukocyte antigen (HLA)-A24-positive patients with HCC, 29 chronic hepatitis C patients without HCC, who were diagnosed by liver biopsy, and 11 healthy blood donors who did not have a history of cancer and were negative for hepatitis B surface antigen and anti-hepatitis C virus (HCV) antibody (Ab). The diagnosis of HCC was histologically confirmed in 21 patients. For the remaining 10 patients the diagnosis was based on typical hypervascular tumor staining on angiography in addition to typical findings, which showed hyperattenuated areas in the early phase and hypoattenuation in the late phase on dynamic computed tomography (CT).²⁹

HLA-based typing of peripheral blood mononuclear cells (PBMCs) from patients and normal blood donors was performed as described.¹⁹ The pathological grading of tumor cell differentiation was assessed according to the general rules for the clinical and pathological study of primary liver cancer.³⁰ The severity of liver disease was evaluated according to the criteria of Desmet et al.³¹ using biopsy specimens of liver tissue.

All patients gave written informed consent to participate in the study in accordance with the Helsinki Declaration and this study was approved by the re-

Table 1. Peptides

Peptide No.	Peptide Name	Source	Reference	Amino Acid Sequence	Number of Specific Spots in Normal Donors (Mean SD)
1	ART1 ₁₈₈	ART1	5	EYCLKFKL	0.9 ± 1.1
2	ART4 ₁₆₁	ART4	6	AFLRHAAL	0.3 ± 0.5
3	ART4 ₈₉₉	ART4	6	DYPSLSATDI	0.6 ± 1.0
4	Cyp-B ₁₀₉	Cyp-B	7	KFHRVVKDF	0.5 ± 0.9
5	Cyp-B ₃₁₅	Cyp-B	7	DFMIQGGDF	1.2 ± 1.7
6	Lck ₂₀₈	Lck	8	HYTNASDGL	0.3 ± 0.6
7	Lck ₄₈₆	Lck	8	TFDYLRSLV	0.2 ± 0.8
8	Lck ₄₈₈	Lck	8	DYLRSLVLEDF	0.9 ± 1.5
9	MAGE1 ₁₃₅	MAGE-A1	9	NYKHCPEI	1.0 ± 0.9
10	MAGE3 ₁₉₅	MAGE-A3	10	IMPKAGLLI	1.4 ± 1.7
11	SART1 ₁₆₉₀	SART1	11	EYRGFTQDF	0.9 ± 1.3
12	SART2 ₈₉₉	SART2	12	SYTRLFLIL	1.0 ± 1.4
13	SART3 ₁₀₉	SART3	13	VYDYNCHVDL	2.1 ± 1.9
14	Her-2/neu ₈	Her-2/neu	14	RWGLLLALL	1.4 ± 2.0
15	p53 ₁₂₅	p53	15	TYSPALNKMF	1.4 ± 1.5
16	p53 ₁₆₁	p53	16	AIYQSQHM	0.4 ± 0.6
17	p53 ₂₀₄	p53	17	EYLDNRNTF	1.1 ± 1.5
18	p53 ₂₁₁	p53	17	TFRHSVW	0.9 ± 1.9
19	p53 ₂₃₅	p53	17	NYMCNSSCM	2.1 ± 2.6
20	MRP3 ₅₀₃	MRP3	18	LYAWEPSFL	0.2 ± 0.5
21	MRP3 ₆₉₂	MRP3	18	AVPQQAWI	1.5 ± 2.1
22	MRP3 ₇₆₅	MRP3	18	VYSDADIFL	0.9 ± 1.0
23	AFP ₃₅₇	AFP	19	EYSRRHPQL	1.8 ± 2.0
24	AFP ₄₀₃	AFP	19	KYIQESQAL	1.1 ± 1.5
25	AFP ₄₃₄	AFP	19	AYTKKAPQL	0.8 ± 1.1
26	hTERT ₁₆₇	hTERT	20	AYQVCGPPL	0.8 ± 1.1
27	hTERT ₃₂₄	hTERT	20	VYAETKHL	0.5 ± 0.7
28	HIV env ₅₈₄	HIV env	32	RYLRDQQLL	1.3 ± 2.0
29	HCV NS3 ₁₀₃₁	HCV NS3	33	AYSQTRGL	ND
30	CMV pp65 ₃₂₈	CMV pp65	34	QYDPVAALF	13.3 ± 15.7

ND, not determined.

gional ethics committee (Medical Ethics Committee of Kanazawa University, No. 829).

Peptides, Cell Lines, and Preparation of PBMCs. Twenty-seven peptides derived from 14 different TAAs (Table 1), human immunodeficiency virus (HIV) envelope-derived peptide (HIVenv₅₈₄),³² HCV NS3-derived peptide (HCVNS3₁₀₃₁),³³ and cytomegalovirus (CMV) pp65-derived peptide (CMVpp65₃₂₈),³⁴ which were identified as HLA-A24 restricted CTL epitopes in previous studies, were used. Peptides were synthesized at Mimotope (Melbourne, Australia) and Sumitomo Pharmaceuticals (Osaka, Japan). They were identified using mass spectrometry and their purities were determined to be >80% by analytical high-performance liquid chromatography (HPLC). The HLA-A*2402 gene-transfected C1R cell line (C1R-A24) was cultured in RPMI 1640 medium containing 10% fetal calf serum (FCS) and 500 μg/mL hygromycin B (Sigma, St. Louis, MO), and K562 was cultured in RPMI 1640 medium containing 10% FCS.³⁵ PBMCs were isolated before HCC treatments as described.²⁰ In 12 patients their PBMCs were also obtained 4 weeks after treatments.

Table 2. Characteristics of the Patients Studied

Clinical Diagnosis	No. of		Age (yr) Mean ± SD	ALT (IU/L) Mean ± SD	AFP (ng/ml) Mean ± SD	Child Pugh (A/B/C)	Diff. Degree* (wel/mod/ por/ND)	Tumor Size† (large/small)	Tumor Multiplicity (multiple/solitary)	Vascular Invasion (+/-)	TNM Stage (I/II/IIIa/IIIB/ IIIC/IV)
	Patients	Sex M/F									
Normal donors	11	8/3	35 ± 2	ND	ND	ND	ND	ND	ND	ND	ND
Chronic hepatitis	29	16/13	59 ± 10	92 ± 94	31 ± 87	27/2/0	ND	ND	ND	ND	ND
HCC	31	23/8	71 ± 4	74 ± 33	1768 ± 9103	20/10/1	11/10/0/10	22/9	20/11	9/22	10/12/3/1/2/3

*Histological degree of HCC; wel; well differentiated, mod: moderately differentiated, por: poorly differentiated, ND: not determined.

†Tumor size was divided into either "small" (≤ 2 cm) or "large" (> 2 cm).

CTL Induction and Cytotoxicity Assay. CTL induction and cytotoxicity assays were performed as described.²⁰ Briefly, stimulated PBMCs were added at effector to target ratios of 100:1, 50:1, 25:1, 13:1, 6:1, and 3:1. In cases where the number of CTLs was insufficient, cytotoxicity assays were performed at effector to target ratios less than 100:1.

Interferon Gamma IFN- γ Enzyme-Linked Immunospot (ELISPOT) Assay. IFN- γ ELISPOT assays were performed as reported.²⁰ Responses to TAA-derived peptides were considered positive if more than 10 specific spots were detected, which is greater than the mean plus 3 standard deviations (SDs) of the baseline response detected in 11 normal blood donors (Table 1), and if the number of spots in the presence of an antigen was at least 2-fold that in its absence. Responses to HIV-, HCV-, and CMV-derived peptides were considered positive if more than 10 specific spots were detected and if the number of spots in the presence of an antigen was at least 2-fold that in its absence. In ELISPOT assay with blocking CTLA-4, anti-human CTLA-4 (eBioscience, Tokyo, Japan) was added at a final concentration of 50 μ g/mL, which has been described to have maximum effect in *in vitro* cultures.³⁶ As a control, functional grade mouse immunoglobulin G (IgG)2a isotype control was used. The assay with blocking CTLA-4 was performed in triplicate and the results were statistically analyzed using the unpaired Student's *t* test.

Cytokine and Chemokine Profiling. The effect of CTLA-4 antibody on TAA-specific T-cell responses was also analyzed by cytokine and chemokine profiling. Cytokine and chemokine levels in the medium of ELISPOT assay were measured using the Bio-plex assay (Bio-Rad, Hercules, CA). These included interleukin (IL)-1 β , IL-1Ra, IL-2, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-12p70, IL-13, IL-15, IL-17, basic fibroblast growth factor (FGF), eotaxin, G-CSF, GM-CSF, IFN- γ , IP-10, MCP-1, macrophage inflammatory protein (MIP)-1 α , MIP-1 β , platelet-derived growth factor (PDGF)-BB, RANTES, tumor necrosis factor alpha (TNF- α), and vascular endothelial growth

factor (VEGF). Eight standards (ranging from 2 to 32,000 pg/mL) were used to generate calibration curves for each cytokine. Data acquisition and analysis were carried out using Bio-plex Manager software v. 4.1.1.

Cytokine Secretion Assay. TAA-specific IFN- γ -producing T cells were also analyzed by cytokine secretion assay. The assay was performed with the MACS cytokine secretion assay (Miltenyi Biotec K.K., Tokyo, Japan), in accordance with the manufacturer's instructions. Briefly, 5,000,000 PBMCs were pulsed with TAA-derived peptides for 16 hours and then incubated with 20 μ L of IFN- γ detection antibody, 10 μ L of anti-CD8-APC Ab (Becton Dickinson, Tokyo, Japan), 10 μ L of anti-CCR7-FITC Ab (eBioscience, Tokyo, Japan), and 10 μ L of anti-CD45RA-PerCP-Cy5.5 Ab (eBioscience, Tokyo, Japan) for 10 minutes at 4°C. After washing with a cold buffer (phosphate-buffered saline/0.5% bovine serum albumin with 2 mM EDTA), the cells were resuspended with 500 μ L of cold buffer and analyzed using FACSCalibur (Becton Dickinson, Tokyo, Japan). As a positive control, CMVpp65₃₂₈-specific IFN- γ -producing T cells were also analyzed by the same methods. The number of IFN- γ -producing T cells was calculated from the results of FACS analysis and is shown as a number per 300,000 PBMCs.

Results

Patient Profile. The clinical profiles of the 11 healthy blood donors, 29 patients with chronic hepatitis C, and 31 patients with HCV-related HCC analyzed in the present study are shown in Table 2 and Fig. 1. Using TNM staging of the Union Internationale Contre Le Cancer (UICC) system (6th v.), 10, 12, 3, 1, 2, and 3 patients were classified as having stage I, II, IIIA, IIIB, IIIC, and IV tumors, respectively.

Detection of TAA-Specific T Cells in HCC Patients. First we examined the frequency of cells that specifically reacted with TAA-derived and control peptides in HCC patients. Fifty-one responses in total were observed against TAA-derived peptides. Twenty-




Fig. 1. TAA-, HIV-, HCV-, and CMV-derived peptide-specific T-cell responses. Results of all HCC patients examined are shown. The T-cell responses were examined by IFN- γ ELISPOT assay. Responses to peptides were considered positive if more than 10 specific spots per 300,000 PBMCs were detected and if the number of spots in the presence of an antigen was at least 2-fold that in its absence. Black boxes indicate the presence of a significant IFN- γ T-cell response to peptides. Peptide sequences are described in Table 1 and characteristics of patients in Table 2.

four of 31 (77.4%) patients showed positive responses to at least one TAA-derived peptide and most of them showed responses to 1 to 4 kinds of TAA-derived peptide. Twenty-three of 27 (85.2%) TAA-derived peptides were recognized by T cells of at least one patient. Peptides 4, 12, 13, 16, 17, 22, 24, and 27 were recognized in more than two patients, suggesting that these peptides were immunogenic. Peptides 28 (HIV env₅₈₄), 29 (HCV 1031), and 30 (CMV pp65₃₂₈) were recognized by 0 (0%), 5 (16%), and 13 (42%) patients, respectively.

The magnitude of TAA-specific T-cell responses was assessed by the frequency of peptide-specific IFN- γ -producing T cells in the PBMC population (Fig. 2A). The range of TAA-derived peptide-specific T-cell frequency was 10-60.5 cells/300,000 PBMCs. Those specific to peptides 13 and 16 numbered more than 30 cells/300,000 PBMCs, suggesting that these peptides were immunogenic. The frequencies of T cells specific to HCV- and CMV-derived peptides were 12-22 cells and 12-92/300,000 PBMCs, respectively.

Whether these TAA-derived peptides were capable of generating peptide-specific CTLs from PBMCs was investigated in HCC patients. The seven peptides were selected according to the magnitude of TAA-specific T-cell responses determined by the fre-

quency of T cells with a positive response. The CTLs generated with these peptides were cytotoxic to C1RA24 cells pulsed with the corresponding peptides (Fig. 2B).

Comparison of TAA-Specific T-Cell Responses Between the Patient Groups With and Without HCC. To characterize the immunogenicity and specificity of TAA-derived peptides, we compared T-cell responses to the peptides derived from TAA, HIV, HCV, and CMV among three groups consisting of normal blood donors, patients with chronic hepatitis C, and patients with HCV-related HCC. A significant TAA-specific T-cell response was not detected in normal blood donors (Fig. 3A). A response was detected in both chronic hepatitis C and HCC patient groups, but it was more frequently observed in HCC patients. HIV-specific T-cell response was not detected in any group. HCV-specific T-cell response rate was not different between the groups with chronic hepatitis C and HCC. CMV-specific T-cell response rates were similar among the three groups. Similar tendencies were observed in the analysis of individual peptides (Fig. 3B). We also examined the frequency of T cells responsive to peptides among the three groups. The mean frequency of TAA-specific T cells without *in vitro* expansion was higher in HCC patients than in




Fig. 2. Vigor of TAA-, HCV-, and CMV-derived peptide-specific T-cell responses. (A) The frequency of TAA-specific IFN- γ -producing T cells was analyzed by ELISPOT assay. Only positive responses are shown. Black bars indicate the response of one patient. *1, *2, *3, *4, and *5 denote 33, 60.5, 44, 92, and 67.5 specific spots, respectively. (B) Representative TAA-specific T-cell responses were also analyzed by CTL assay. T cell lines were generated from PBMC of the HLA-A24-positive HCC patients by stimulation with TAA-derived peptides (peptides 4, 12, 13, 16, 22, 24, and 27) (see Table 1). Expanded T cell lines were then tested for specific cytotoxicity against the corresponding peptides in a standard ⁵¹Cr release assay at the indicated E:T ratios.

patients with chronic hepatitis C for 14 of 27 TAA-derived peptides (peptides 1, 2, 3, 4, 12, 16, 18, 19, 20, 21, 22, 24, 25 and 27) (Fig. 3C).

Enhancement of TAA-Specific T-Cell Responses After HCC Treatments. Several studies including our own have clarified that HCC treatments enhanced HCC-specific immune responses (19, 37, 38). In this study, we examined whether the enhancement was observed equally in all kinds of TAAs or specifically in some TAAs. For this purpose we measured the frequency of TAA-specific T cells before and after HCC treatment by ELISPOT assay in 12 cases who received transcatheter arterial embolization (TAE), radiofrequency ablation (RFA), or chemotherapy. The frequency of TAA-specific T cells increased in all patients and it was observed for 23 of 27 TAA-derived peptides (Fig. 4A). The enhancement was observed in the

patients who received TAE, RFA, or chemotherapy and even in the patients without an increase in the frequency of CMV-specific T cells. Peptides 7, 14, 15, and 26, which were not recognized by T cells in all HCC patients before treatments (Fig. 1), were recognized by T cells in 1, 4, 1, and 5, respectively, of 12 patients after treatments. Representative results of enhancement of TAA-specific immune responses are shown in Fig. 4B. The frequency of TAA-specific T cells increased to 11-80 cells/300,000 PBMCs after treatments.

The enhancement of TAA-specific immune responses was also confirmed by cytokine secretion assay. Representative results are shown in Fig. 4C. In this patient (patient 25) the frequency of TAA-specific IFN- γ -producing CD8⁺ T cells was increased from 0.4% to 1.4% of CD8⁺ T cells after HCC treatment.




Fig. 3. Comparative analysis of TAA-, HIV-, HCV-, and CMV-derived peptide-specific T-cell responses among three groups of subjects: normal donors, patients with chronic hepatitis C not complicated by HCC, and HCC patients. (A) Summary of the number of patients with a significant IFN- γ T-cell response to tumor-associated, HIV, HCV, and CMV antigens in each group. (B) Graph shows the percentage of patients in each group who showed a significant IFN- γ T-cell response to individual peptides. Peptide sequences are described in Table 1. (C) Mean frequency of peptide-specific IFN- γ -producing T cells in each group. The frequency of IFN- γ -producing T cells was analyzed by ELISPOT assay.

In this assay we also examined the naïve/effector/memory phenotype of these cells by the criterion of CD45RA/CCR7 expression.³⁹ Phenotypic analysis of TAA-specific, IFN- γ -producing memory CD8⁺ T cells before and after treatment showed that the frequency of CD45RA⁻/CCR7⁺ central memory T cells was the highest, indicating that the posttherapeutic increase in these T cells is due to the increase in cells with this phenotype (Fig. 4D). In this patient the number of T cells with the CD45RA⁻/CCR7⁺ phenotype increased from 73 cells/300,000 PBMCs before treatment to 316 cells/300,000 PBMCs after treatment. Similar results were noted in five patients.

Blocking CTLA-4 Restores TAA-Specific T-Cell Responses. In previous studies including our own,^{19,20,24} the CTL epitopes that correlate with the prevention of tumor progression or prognosis of HCC patients have not been identified. One of the reasons for this is considered to be that the naturally occurring

T-cell responses to the epitopes are weak; therefore, recent tumor immunotherapeutic studies are moving toward modulation of T-cell responses.

CTLA-4 is recognized as a critical negative regulator of immune response; therefore, its blockade has been considered to contribute to antitumor activity.²⁷ In a recent study it was reported that blocking of CTLA-4 on both effector and regulatory T cell compartments contributes to the antitumor activity of CTLA-4 antibodies.⁴⁰ To examine whether similar occurs for immune response in HCC patients, we analyzed 32 separate TAA-specific T-cell responses in 15 HCC patients using 13 TAA-derived peptides. Incubation of T cells with CTLA-4 antibodies resulted in an increase of the number of TAA-specific T cells in 18 of 32 (56%) responses and in 9 of 15 (60%) patients (Fig. 5A). Fourteen and four patients showed increases of 1-10 and more than 10 TAA-specific T cells, respectively. Representative results of six patients are shown




Fig. 4. Enhancement of TAA-specific T-cell responses in HCC patients after treatments. (A) Summary of patients and peptides with a significant increase of the number of IFN- γ -producing T cells (black boxes). A significant change in the IFN- γ response was defined as a more than 2-fold increase and the presence of more than 10 specific spots in ELISPOT assay after HCC treatments. The assays were performed in 12 HCC patients using 27 TAA-, HIV-, and CMV-derived peptides. Goal* shows the goal of HCC treatment. C and P denote "curative intention" and "palliative intention," respectively. (B) Representative results of ELISPOT assay are shown. White and black bars indicate the frequency of T cells before and after HCC treatments, respectively. *1, *2, *3, and *4 denote 53, 60, 80, and 121 specific spots, respectively. (C) Enhancement of TAA-specific T-cell responses was also analyzed by cytokine secretion assay. Representative results are shown (patient 25). PBMCs were pulsed with TAA-derived peptides (peptides 14, 21, and 24) for 16 hours and then analyzed for IFN- γ production. (D) IFN- γ -producing T cells were also examined for naïve/effector/memory phenotype by the criterion of CD45RA/CCR7 expression. The number of cells was calculated from the results of FACS analysis and is shown as a number per 300,000 PBMCs. White and black bars indicate the frequency of TAA-specific IFN- γ -producing T cells before and after HCC treatments, respectively. The experiments were performed in five patients and similar results were observed.

in Fig. 5B. The magnitude of TAA-specific T-cell increase was statistically significant in four patients.

To examine the effect of CTLA-4 antibodies for production of other cytokines by T cells, we measured 27 kinds of human cytokines and chemokines in the medium of ELISPOT assay. Figure 5C shows the results of cytokine production in the well with positive T-cell responses against TAA-derived peptides. The various cytokines consisting of IL-1 β , IL-4, IL-6, IL-10, IL-17, eotaxin, G-CSF, GM-CSF, IFN- γ , MIP-1 α , MIP-1 β , RANTES, and TNF- α were increased in the medium with CTLA-4 antibodies compared with that without CTLA-4 antibodies. In contrast, increased

production of these cytokines in the well without positive T-cell responses against TAA-derived peptides was not observed in medium either with or without CTLA-4 antibodies (Fig. 5D).

Discussion

In recent years, specific TAAs and their CTL epitopes have been identified in many tumors.²¹ Several TAAs and their CTL epitopes, such as AFP, MAGE, and human telomerase reverse transcriptase (hTERT) have also been reported in HCC.^{19,20,24,41} Although AFP-targeting immunotherapy could induce TAA-




Fig. 5. Enhancement of TAA-specific T-cell responses in HCC patients by CTLA-4 antibodies. (A) Summary of patients and peptides with an increase of the number of IFN- γ -producing T cells. Black, gray, white, and hatched boxes indicate the immune responses with an increase of more than 10 specific spots, an increase of 1-10 specific spots, without change and a decrease of 1-10 specific spots, respectively. (B) Representative results of six patients are shown. Black and white bars indicate the results of assays incubated with CTLA-4 antibodies and mouse IgG2a isotype control, respectively. Data are expressed as the mean \pm SD of specific spots, except for patients 14 and 31. (C) Effects of CTLA-4 antibodies on production of cytokine and chemokine. Cytokine and chemokine levels in the medium of ELISPOT assay were measured using the Bio-plex assay. The graphs indicate the concentrations of cytokine and chemokine in the medium of ELISPOT assay using PBMCs of patient 31 and peptide 13 (medium in ELISPOT assay with enhancement of T-cell response) (see A,B). The increase of cytokines and chemokines after incubation with anti-CTLA-4 antibodies was confirmed in another three experiments using PBMCs of three other patients. (D) The graphs indicate the concentrations of cytokine and chemokine in the medium of ELISPOT assay using PBMCs of patient 31 and peptide 22 (medium in ELISPOT assay without enhancement of T-cell response) (see A).

specific CTLs, no patients achieved an objective tumor response; therefore, the search for TAAs as suitable targets for HCC immunotherapy and identification of their epitopes are important issues in therapy development. However, to date, T-cell responses to previously identified TAAs or their epitopes have been measured simultaneously and comparatively in only one study involving several patients with HBV-related HCC,⁴² but no T-cell responses to the many other TAAs or their epitopes have been evaluated.

In this study we performed a simultaneous, comparative analysis of immune responses to 27 different CTL epitopes derived from 14 previously reported TAAs in the peripheral blood lymphocytes of 31 HCV-related HCC patients. We noted immune responses to epitopes (peptides 4, 12, 13, 16, 17, 22, 24, and 27) derived from CypB, SART2, SART3,

p53, MRP3, AFP, and hTERT in more than two patients (Fig. 1). These findings suggest the immunogenicity of these TAAs and their epitopes. In addition, the frequencies of peripheral blood CTLs specific to epitopes (peptides 4, 13, 16, 22, and 24) derived from CypB, SART3, p53, MRP3, and AFP, as detected by the ELISPOT assay, were high (≥ 20 specific spots/300,000 PBMCs), suggesting the high immunogenicity of these TAAs and their epitopes.

Among these immunogenic antigens the expression of p53, MRP3, AFP, and hTERT was reported in HCC.^{18,19,43,44} We also previously confirmed that the expression of SART2 and SART3 was observed in 100% of human HCC tissue (data not shown). As for CypB, this protein is well known to be widely expressed in normal and malignant tissue⁷; therefore, it is considered to be expressed in HCC.

Regarding tumor immunotherapy, it has recently been reported that strong immune responses can be induced at an earlier postvaccination time using, as peptide vaccines, epitopes that frequently occur in peripheral blood CTL precursors.²³ The epitopes (peptides 4, 12, 13, 16, 22, 24, and 27) that were derived from CypB, SART2, SART3, p53, MRP3, AFP, and hTERT and considered to be highly immunogenic in this study were capable of inducing epitope-specific CTLs from the PBMCs of HCC patients, suggesting that these epitopes can be candidates for peptide vaccines.

Next, TAA-specific immune responses were compared among three groups of subjects: HCC patients, normal blood donors, and patients with chronic hepatitis C not complicated by HCC. The results showed that there were no differences in the positive rate of immune responses to CMV among the three groups and no difference in the positive rate of immune responses to HCV between chronic hepatitis C patients with and without HCC. However, TAA-specific immune responses were observed frequently only in HCC patients, indicating that these immune responses are specific to HCC.

In the present study we also analyzed factors influencing host immune responses to these TAA-derived epitopes. Previous studies have reported that treatments, such as RFA and TAE, enhance HCC-specific T-cell responses.^{19,37,38} However, TAAs and their epitopes, to which these enhanced immune responses occur, have not been identified. Thus, we simultaneously measured immune responses to 27 different epitopes derived from 14 TAAs in 12 patients who were available for analysis before and after treatment. The results showed that the antigens and their epitopes to which treatment-enhanced T-cell responses occur were diverse and some of them were newly induced after HCC treatment, suggesting that HCC treatments could induce *de novo* T-cell responses and these TAAs and their epitopes can be candidates as targets for HCC immunotherapy.

Furthermore, it became clear that enhanced immune responses to TAAs were induced not only by previously reported RFA and TAE, but also by cytotoxic drug chemotherapy. The patients who received chemotherapy showed partial responses after the treatment; therefore, we considered that it induced release of TAA into the tumor environment by tumor necrosis and/or apoptosis such as the mechanism reported in RFA or TAE.^{19,37,38} Thus, our findings suggest that combined cancer chemotherapy and immunotherapy is useful as a treatment for HCC.

Analysis of the memory phenotypes of the T cells thus induced showed that the phenotypes of T cells whose frequency increased were mostly CD45RA⁻/CCR7⁺ T cells (central memory T cells). Previous studies have reported that T cells with this phenotype differentiate into effector memory T cells and effector T cells, and that they require secondary stimulation by antigen to exert stronger antitumor effects.³⁹ Therefore, our findings suggest that the antitumor effect of tumor-specific T cells induced by HCC treatment is insufficient, and a booster with TAAs or epitope-containing peptides is a suitable method to further enhance antitumor effects.

Finally, we investigated the effect of anti-CTLA-4 antibodies, which have recently been in clinical trials as drugs enhancing antitumor immunity, on the host immune response to HCC. Regarding the mechanism of the antitumor activity of anti-CTLA-4 antibodies, it has been reported that they maximize the antitumor effect by blocking CTLA-4 on the surface of effector and regulatory T cells.⁴⁰ Because the number of peripheral blood regulatory T cells has been reported to increase in HCC patients,⁴⁵ TAA-specific CTLs that should be present but may not be detected by the ELISPOT assay. Therefore, in this study anti-CTLA-4 antibodies were added along with peptides to examine their effect on the ELISPOT assay.

The addition of anti-CTLA-4 antibodies resulted in an increase in the frequency of TAA-specific T cells in 60% of HCC patients. Although most patients showed an increase of only 1-10 TAA-specific T cells, the increased number of T cells was statistically significant. In addition, an increase of more than 10 TAA-specific T cells and a conversion from a negative to a positive response were observed in four patients. These results suggested that the anti-CTLA-4 antibodies unmasked IFN- γ production by CTLs. However, the function might be limited because the number of TAA-specific T cells was not changed and even decreased in some patients.

The cytokine and chemokine profiling showed that the addition of anti-CTLA-4 antibodies increased the production of not only IFN- γ but also cytokines, such as TNF- α , IL-1, and IL-6, and chemokines such as MIP-1; therefore, we speculate that the increased production of these antitumor immunity substances also plays a role in the unmasking of TAA-specific CTLs by anti-CTLA-4 antibodies. These results suggest that anti-CTLA-4 antibody is promising as a drug to enhance antitumor immunity, and that the ELISPOT assay with this antibody may serve as a more appropriate test tool to detect more HCC-specific TAAs or their epitopes.