

**Table 1.** Clinical characteristics of patients in this study

	GWAS (n = 303)	Replication (n = 391)
Age	57.4 (9.7)	56.8 (9.9)
Sex (M/F)	151/152	209/182
Weight (kg)	60.6 (10.4)	61.3 (10.7)
Body mass index	23.5 (3.1)	23.7 (4.1)
Baseline Hb (g/dl)	14.1 (1.4)	14.1 (1.4)
Baseline platelet count (10 <sup>9</sup> /l)	151.3 (54.3)	159.7 (55.0)
Baseline ALT (IU/l)	83.5 (79.4)	86.8 (71.9)
Baseline creatinine (mg/dl)	0.70 (0.15)	0.72 (0.16)
Baseline liver fibrosis (F0–2/F3–4/ ND)	153/77/73	175/59/43
rs8099917: TT/non-TT	165/138	296/95
rs1127354: AA/CA/CC	4/79/220	6/101/284
Week 4 Hb (g/dl)	11.8 (1.7)	11.9 (1.5)
Week 4 platelet count (10 <sup>9</sup> /l)	127.6 (48.2)	132.4 (51.0)
Hb reduction at week 4	–2.3 (1.4)	–2.2 (1.4)
Platelet reduction at week 4	–22.2 (38.4)	–24.7 (30.4)

located on *DDRGKI* gene and *ITPA* gene on chromosome 20 showed strong associations in the allele frequency model ( $P = 3.29 \times 10^{-10}$  and  $P = 2.56 \times 10^{-9}$ ) with Hb reduction in response to PEG-IFN plus RBV treatment (Table 3).

The above 22 SNPs were selected for the replication study and fine mapping, including rs1127354, which was reported by the US group (22) to be strongly associated with Hb reduction (Supplementary Material, Table S2). All SNPs were genotyped using the DigiTag2 assay in an independent set of 391 Japanese HCV patients with quantitative change in Hb in response to PEG-IFN/RBV treatment [137 patients with Hb reduction versus 254 patients without Hb reduction (Table 3)]. The associations of the original SNPs were replicated in the second set of 391 patients in the minor-allele-dominant model ( $P = 3.86 \times 10^{-16}$ , OR = 0.02 for rs11697186;  $P = 6.90 \times 10^{-18}$ , OR = 0.03 for rs6139030, Table 3). The combined  $P$ -values for both stages reached  $9.43 \times 10^{-25}$  (OR = 0.03; 95% CI = 0.01–0.08) and  $2.12 \times 10^{-25}$  (OR = 0.04; 95% CI = 0.02–0.09), respectively (Table 3). The rs1127354 was also strongly associated with a quantitative change in Hb in response to PEG-IFN/RBV treatment in a set of 694 Japanese HCV patients (303 patients from the GWAS stage plus the second set of 391 patients) with and without Hb reduction ( $P = 4.58 \times 10^{-26}$ , OR = 0.03; 95% CI = 0.01–0.08).

Fine mapping with 22 SNPs around *DDRGKI* and *ITPA* genes showed that four significant SNPs (rs11697186, rs6139030, rs1127354 and rs13830) at the GWAS stage had a strong linkage disequilibrium (LD) ( $r^2 > 0.86$ ) within the 22.7 kb region (Fig. 2). As the rs1127354 is known as a functional variant in the *ITPA* gene that caused ITPase deficiency and protected against RBV-induced HA (22,25), the representative SNP was applied for the following detailed studies.

#### *ITPA/DDRGKI* variants reflect anemia and reactive increase of the platelet count


The mean quantitative reduction of blood cells from the baseline according to the *ITPA* rs1127354 genotypes is shown in Figure 3. Patients with the rs1127354 genotypes AA and CA showed lower degree of Hb reduction at weeks 2, 4, 8 and

12 during therapy compared with those with the CC genotype ( $P < 0.0001$  for weeks 2, 4, 8 and 12 in Fig. 3A). The most difference of mean Hb reduction was found at week 4 (AA/CA –1.14 versus CC –2.72). These results show that the AA and CA genotypes are significantly associated with less absolute reduction in Hb levels, especially during the early weeks of therapy, and protect against the development of severe anemia. Interestingly, the CC genotype had significantly less reduction in the mean platelet count compared with the AA/CA genotype ( $P < 0.0001$  for weeks 2, 4, 8;  $P = 0.019$  for week 12 in Fig. 3B), due to a reactive increase of platelet count through weeks 1–4. The most difference of mean platelet reduction was found at week 4 [AA/CA –41.2 versus CC –18.0 (10<sup>9</sup>/l)]. There was no difference in the neutrophil leukocyte count between genotypes (Fig. 3C). We then compared the percentage of patients with platelet count reduction in the *ITPA* rs1127354 genotypes at week 4 of PEG-IFN/RBV therapy (Fig. 4). The percentage of patients with a platelet count reduction of  $<30$  (10<sup>9</sup>/l) at week 4 was significantly higher in the rs1127354 genotypes CC ( $P < 0.0001$ ), indicating that the degree of platelet count reduction was less in patients with the rs1127354 genotype CC. A multivariate analysis for factors associated with a platelet reduction  $>30$  (10<sup>9</sup>/l) at week 4 showed that lower platelet count at the baseline and the rs1127354 genotypes AA/CA were independently associated with platelet reduction (OR = 1.15; 95% CI = 1.11–1.20;  $P < 0.0001$ , OR = 5.92; 95% CI = 3.82–9.17;  $P < 0.0001$ , respectively).

Figure 5 showed reactive increase of the platelet count through weeks 1–4 of PEG-IFN/RBV therapy. Patients with anemia (Hb reduction  $\geq 3.0$  g/dl) at week 4 had a significantly higher degree of the reactive increase of the platelet count than those without anemia ( $P < 0.0001$  in Fig. 5A). Within a subgroup of patients with the rs1127354 genotypes CC, patients with anemia still had a significantly higher degree of reactive increase of the platelet count than those without anemia ( $P = 0.004$  in Fig. 5B). On the other hand, patients with the rs1127354 genotypes CC had a significantly higher degree of the reactive increase of the platelet count than those with genotypes AA/CA ( $P < 0.0001$  in Fig. 5C), and a similar result was obtained in a subgroup of patients without anemia (Fig. 5D). To elucidate the significant factors associated with the rs1127354 genotypes by multivariate analysis, the rs1127354 genotypes AA/CA were independently associated with protection against the reduction in Hb and more reduction in platelet counts at week 4 due to a lower degree of the reactive increase of the platelet count (OR = 0.029; 95% CI = 0.009–0.092;  $P < 0.0001$ , OR = 4.73; 95% CI = 3.04–7.37;  $P < 0.0001$ , respectively). Indeed, the reactive increase of the platelet count through weeks 1–4 was positively correlated with a high platelet count at the baseline and anemia (Hb reduction  $\geq 3.0$  g/dl) at week 4, but was negatively correlated with rs1127354 genotypes AA/CA and a platelet count reduction of  $\geq 30$  (10<sup>9</sup>/l) at week 4 (Table 4).

#### Relationship between *ITPA* rs1127354 genotypes and treatment outcome due to dose reduction of PEG-IFN or RBV

In this population, a multivariate analysis showed that SVR was significantly associated with *IL28B* TT-genotype [OR



**Figure 1.** Genome-wide association results in 303 Japanese HCV patients with the decrease of platelets in response to PEG-IFN plus RBV treatment (107 patients with the decrease of PLT and 196 patients without the decrease of PLT).  $P$ -values were calculated using the  $\chi^2$  test for allele frequencies. Dots with arrow on chromosome 20 showed a significant SNP ( $P = 8.17 \times 10^{-9}$  for rs11697186) and a candidate SNP with a marginal significance ( $P = 4.30 \times 10^{-7}$  for rs6139030) associated with the decrease of PLT with response to PEG-IFN/RBV treatment. The dotted line indicates a genome-wide significance ( $P < 8.40 \times 10^{-8}$ ).

**Table 2.** Two SNPs (rs11697186 and rs6139030) significantly associated with the decrease of PLT in response to PEG-IFN/RBV treatment

dbSNP rsID	Nearest gene	MAF <sup>a</sup> (allele)	Allele (1/2)	Stage	Patients with the decrease of PLT			Patients without the decrease of PLT			OR (95% CI) <sup>b</sup>	$P$ -value <sup>c</sup>
					11	12	22	11	12	22		
rs11697186	<i>DDRGK1</i>	0.15 (T)	T/A	GWAS	3 (2.8)	48 (44.9)	56 (52.3)	0 (0.0)	32 (16.6)	161 (83.4)	4.6 (2.7–7.8)	$8.17 \times 10^{-9}$
				Replication	3 (1.8)	65 (39.9)	95 (58.3)	3 (1.4)	25 (12.0)	181 (86.6)	4.6 (2.8–7.7)	$5.88 \times 10^{-10}$
				Combined	6 (2.2)	113 (41.9)	151 (55.9)	3 (0.7)	57 (14.2)	342 (85.1)	4.5 (3.1–6.5)	$5.29 \times 10^{-17}$
rs6139030	<i>ITPA</i>	0.17 (C)	T/C	GWAS	56 (52.3)	48 (44.9)	3 (2.8)	157 (80.1)	38 (19.4)	1 (0.5)	3.7 (2.2–6.1)	$4.30 \times 10^{-7}$
				Replication	96 (54.9)	74 (42.3)	5 (2.9)	181 (83.8)	32 (14.8)	3 (1.4)	4.3 (2.7–6.8)	$3.83 \times 10^{-10}$
				Combined	152 (53.9)	122 (43.3)	8 (2.8)	338 (82.0)	70 (17.0)	4 (1.0)	3.9 (2.8–5.5)	$1.33 \times 10^{-15}$

<sup>a</sup>Minor allele frequency and minor allele in 184 healthy Japanese individuals.

<sup>b</sup>OR for the minor allele in a dominant model.

<sup>c</sup> $P$ -value by  $\chi^2$  test for the minor allele dominant model.

6.12 (2.78–13.46),  $P < 0.0001$ ] as well as platelet counts [OR 1.18 (1.11–1.26),  $P < 0.00001$ ]. We analyzed whether the rs1127354 genotype could influence the treatment outcome by PEG-IFN/RBV therapy. When analyzed in the patients available for treatment outcome (172 with *ITPA*-AA/CA and 450 with *ITPA*-CC), the percentage of patients receiving  $>80\%$  of the expected PEG-IFN and RBV dose at baseline and week 4 was not significantly different among the rs1127354 genotypes. However, the rate of SVR tended to be higher in patients with *ITPA*-AA/CA genotype than those with *ITPA*-CC (48.8 versus 37.3%), because the relapse rate was lower in patients with *ITPA*-AA/CA. To investigate the influence on treatment outcome by dose reduction of PEG-IFN, in a subgroup of patients with low platelet counts ( $<10$ ) at baseline (19 with *ITPA*-AA/CA and 53 with *ITPA*-CC) we analyzed the treatment outcome according to

rs1127354 genotypes. The SVR rate was very low in each group (21.1% in *ITPA*-AA/CA and 17.0% in *ITPA*-CC), because many patients had the initial dose reduction of PEG-IFN ( $<80\%$  of standard dose)—36.8% of patients with *ITPA*-AA/CA and 44.6% of patients with *ITPA*-CC genotype. Further prospective studies are required among the pre-cirrhotic or cirrhotic patients with low platelet counts.

## DISCUSSION

Recent genome-wide association studies, including our study on HCV infection, have identified two important host genetic variants: the SNP in *IL28B* gene, which is strongly associated with response to therapy for chronic genotype 1 HCV infection (16–21), and the SNP in *ITPA* gene, which precisely predicts RBV-induced anemia in


**Table 3.** Two SNPs (rs11697186 and rs6139030) significantly associated with quantitative change in Hb levels from baseline to week 4 of PEG-IFN/RBV treatment

dbSNP rsID	Nearest gene	MAF <sup>a</sup> (allele)	Allele (1/2)	Stage	Patients with quantitative change in Hb			Patients without quantitative change in Hb			OR (95% CI) <sup>b</sup>	P-value <sup>c</sup>
					11	12	22	11	12	22		
rs11697186	<i>DDRGK1</i>	0.15 (T)	T/A	GWAS	0 (0.0)	3 (3.3)	89 (96.7)	3 (1.5)	77 (37.0)	128 (61.5)	0.06 (0.02–0.16)	$3.29 \times 10^{-10}$
				Replication	0 (0.0)	2 (1.5)	134 (98.5)	6 (2.5)	88 (37.3)	142 (60.2)	0.02 (0.01–0.09)	$3.86 \times 10^{-16}$
				Combined	0 (0.0)	5 (2.2)	223 (97.8)	9 (2.0)	165 (37.2)	270 (60.8)	0.03 (0.01–0.08)	$9.43 \times 10^{-25}$
rs6139030	<i>ITPA</i>	0.17 (C)	T/C	GWAS	88 (93.6)	6 (6.4)	0 (0.0)	125 (59.8)	80 (38.3)	4 (1.9)	0.08 (0.03–0.22)	$2.56 \times 10^{-9}$
				Replication	134 (97.8)	3 (2.2)	0 (0.0)	143 (56.3)	103 (40.6)	8 (3.1)	0.03 (0.01–0.08)	$6.90 \times 10^{-18}$
				Combined	222 (96.1)	9 (3.9)	0 (0.0)	268 (57.9)	183 (39.5)	12 (2.6)	0.04 (0.02–0.09)	$2.12 \times 10^{-25}$

<sup>a</sup>Minor allele frequency and minor allele in 184 healthy Japanese individuals.

<sup>b</sup>OR for the minor allele in a dominant model.


<sup>c</sup>P-value by  $\chi^2$  square test for the minor allele dominant model.




**Figure 2.** Pairwise LD ( $r^2$ ) diagrams for *DDRGK1* and *ITPA*. Lower panel shows estimates of pairwise  $r^2$  for 22 SNPs selected in the replication study using the second set of 391 Japanese HCV patients with and without quantitative change in PLT levels from baseline to week 4 of PEG-IFN/RBV treatment.

European-American population (22) and Japanese population (26). The genetic variation of *ITPA* causing an accumulation of inosine triphosphate (ITP) has been shown to protect patients against RBV-induced anemia during treatment for

CHC infection. A recent report showed the biologic mechanism that ITP confers protection against RBV-induced ATP reduction by substituting for erythrocyte GTP, which is depleted by RBV, in the biosynthesis of ATP (25).



**Figure 3.** *ITPA* rs1127354 genotypes and the quantitative reduction of blood cells from baseline. Mean reduction of (A) Hb levels, (B) platelet counts and (C) neutrophil leukocyte counts during treatment according to rs1127354 genotype is shown. Solid and dotted lines indicate patients with CC and AA/CA genotypes, respectively. Error bars indicate standard error. CC genotype had more reduction in mean Hb levels during therapy compared with the AA/CA genotype ( $*P < 0.0001$  for weeks 2, 4, 8, 12). CC genotype had less of a reduction in mean platelet counts ( $*P < 0.0001$  for weeks 2, 4, 8, and  $**P = 0.019$  for week 12), and showed a reactive increase of platelet counts through weeks 1–4.




**Figure 4.** *ITPA* rs1127354 genotypes and reduction of platelet counts at week 4 of PEG-IFN/RBV therapy. The percentage of patients with platelet count reduction of  $>50$  ( $10^9/l$ ) (black bar),  $30-50$  ( $10^9/l$ ) (gray bar) and  $<30$  ( $10^9/l$ ) (white bar) at week 4 is shown for rs1127354 genotypes. The incidence of platelet count reduction of  $>50$  and  $<30$  was significantly lower in patients with the rs1127354 genotypes CC compared with AA/CA genotypes: 12 versus 35%,  $P < 0.0001$ , and 69 versus 31%,  $P < 0.0001$ , respectively.

In this study, two SNPs, rs11697186 and rs6139030, which were within and around *DDRGK1* gene on chromosome 20, were strongly associated with thrombocytopenia as well as

with Hb reduction at week 4. In clinical practice, the positive predictive value and negative predictive value by rs11697186 genotypes were 66.5 and 69.4% for thrombocytopenia, as well as 97.2 and 45% for RBV-induced anemia at week 4. As previously reported (22,26), a functional SNP (rs1127354) in the *ITPA* locus, which is in strong LD with rs11697186, was the most significant SNP associated with RBV-induced anemia and, in this study, IFN-induced thrombocytopenia in Japanese genetic populations. Note that severe Hb decline, which is mainly found in *ITPA*-CC patients, was inversely correlated with platelet reduction. This would contribute to an association between severe anemia and relative reactive increase of platelet count in this population, which attenuated the IFN effect on the platelet count. Our data supported a previous report which described that the current use of RBV, inducing severe anemia, might blunt the thrombocytopenic effect of IFNs as a result of reactive increase of platelet counts (27).

A previous paper showed hematological and bone marrow effects of RBV in rhesus monkeys (28). Hb values decreased significantly during RBV administration due to dose-related erythroid hypoplasia in bone marrow and returned to normal following withdrawal. On the other hand, increase of the platelet count occurred in both low- and high-dose treatment groups during RBV administration, with a fall of the platelet count to normal after drug withdrawal. The effect on platelet count was clearly dose related, with maximum counts rising to twice and three times above baseline levels in the low- and high-dose groups, respectively. This caused a significant increase of



**Figure 5.** Reactive increase of platelet counts through weeks 1–4. Box plots of reactive increase of platelet count through weeks 1–4 according to the degree of anemia at week 4 are shown for all patients (A) and a subgroup of patients with the rs1127354 genotypes CC (B). Patients with anemia (Hb reduction  $\geq 3.0$  g/dl) at week 4 had a significantly higher degree of reactive increase of platelet count than those without anemia ( $P < 0.0001$ ). Box plots of reactive increase of platelet counts according to the rs1127354 genotype CC are shown for all patients (C) and a subgroup of patients without anemia (D) (Hb reduction  $< 3.0$  g/dl) at week 4. Patients with the rs1127354 genotypes CC had a significantly high degree of reactive increase of platelet counts compared with those with genotypes AA/CA ( $P < 0.0001$ ).

**Table 4.** Multivariate analysis of factors associated with reactive increase of platelets  $\geq 20$  ( $10^9/l$ ) through weeks 1–4

	OR	95% CI	P-value
Baseline platelet counts	1.168	1.101–1.239	$< 0.0001$
ITPA AA/CA	0.379	0.168–0.856	0.0196
Platelet reduction $\geq 30$ ( $10^9/l$ ) at week 4	0.051	0.021–0.120	$< 0.0001$
Hb reduction $\geq 3.0$ g/dl at week 4	1.602	0.914–2.809	0.0996

the platelet count associated with increased numbers of megakaryocytes. Additionally, the sequence homology of thrombopoietin (TPO) and erythropoietin (EPO) may explain the synergy of the physiologic role of TPO and EPO in platelet production. When EPO is elevated, as in iron deficiency anemia, an amino acid sequence similar to TPO may increase the platelet count (29).

Another possibility is a direct association between *ITPA* SNPs or the related SNPs with a strong LD and IFN-induced thrombocytopenia. *DDRGI1* (DDRGI1 domain-containing protein 1) is a novel C53/LZAP-interacting protein. C53/LZAP (also named as Cdk5rap3) is a putative tumor suppressor that plays important roles in multiple cell signaling pathways, including DNA damage response and NF-kappaB signaling (30); however, it remains largely unknown how the function of *DDRGI1* variants is regulated. Further studies are required to elucidate the possible association between *DDRGI1* variants and thrombocytopenia.

Multivariate analysis demonstrated that rs1127354 in the *ITPA* gene was independently associated with RBV-induced severe anemia and IFN-induced thrombocytopenia. This finding suggests that rs1127354 would be a useful marker to predict these hematological side effects by PEG-IFN/RBV therapy, indicating that genetic testing of *ITPA* variant might be applied to establish personalized dosages of PEG-IFN/RBV therapy. The rate of SVR tended to be higher in patients with *ITPA*-AA/CA genotype than those

with *ITPA*-CC in this population. This might reflect decreased treatment efficacy (higher relapse rate) due to dose reduction of RBV in patients with *ITPA*-CC genotype. Our recent paper also demonstrated that the incidence of early dose reduction was significantly higher in *ITPA*-major (CC) patients as expected and, more importantly, that a significantly higher SVR rate was achieved in *ITPA*-hetero/minor (CA/AA) patients with HCV non-1b or low viral load strains (31) and in a subset of Japanese patients with the favorable TT genotype at rs8099917 of *IL28B* (32). Taken together, our results indicate that the *ITPA* minor variant A is not only a protective allele against PEG-IFN and RBV treatment-associated anemia in Japanese population, but also a significant predictor of SVR in certain HCV strains that show good response to IFN. The possible mechanism of protection against RBV-induced hemolysis is that ITP deficiency or low-activity variants (*ITPA* minor variant A) in turn lead to the accumulation of ITP in red blood cells (33,34), and the ITP confers protection against RBV-induced ATP reduction by substituting for erythrocyte GTP (25). On the other hand, half of the *ITPA*-major (CC) patients did not develop a significant Hb decline. This finding suggests other low-frequency *ITPA* variants or SNPs in other enzymes that are involved in erythrocyte purine nucleoside metabolism.

In Japan, the older HCV-infected patients developing liver fibrosis have been prevalent (mean age 62 years) (9). Thrombocytopenia by PEG-IFN/RBV therapy could lead to poor treatment efficiency among such Japanese patients with LC due to the initial or early dose reduction of PEG-IFN. In fact,  $\sim 40\%$  of such population in this study had the initial dose reduction of PEG-IFN, resulting in a low SVR rate. Splenectomy or embolization of the splenic artery might be one of the options to increase the SVR rate, but a sufficient treatment outcome had not been obtained at present (35). Based on the recently accumulated SNP data, if patients had favorable *IL28B* genotype and *ITPA*-CC (lower reduction of platelet counts), a standard dose of PEG-IFN might be available for

the patients with lower platelet counts and the SVR rate might be increased due to sufficient dose of PEG-IFN.

Several STAT-C agents (specifically targeted antiviral therapies for hepatitis C) are being tested for clinical efficacy against hepatitis C (12,13,15,16). Most experts believe that when new drugs are approved to treat hepatitis C, they will be used in combination with PEG-IFN and RBV. Moreover, recent clinical trials, including NS3 protease inhibitors, have shown that PEG-IFN plus RBV would be necessary to achieve optimal treatment responses (12,13). Our present results may provide a valuable pharmacogenetic diagnostic tool for tailoring PEG-IFN and RBV dosing to minimize drug-induced adverse events and for further optimization of clinical anti-HCV chemotherapeutics.

## MATERIALS AND METHODS

### Patients

From April 2007 to April 2010, samples were obtained from 303 patients with chronic HCV (genotype 1) infection who were treated at 14 multi-center hospitals (liver units with hepatologists) throughout Japan. Each patient was treated with PEG-IFN- $\alpha$ 2b (1.5  $\mu$ g/kg body weight, subcutaneously once a week) or PEG-IFN- $\alpha$ 2a (180  $\mu$ g once a week) plus RBV (600–1000 mg daily according to body weight) for 48 weeks. Treatment duration was extended in some patients up to 72 weeks, according to the physicians' preferences. The dose of PEG-IFN or RBV was reduced according to the recommendations on the package inserts or the clinical conditions of the individual patients. EPO or other growth factors were not given. Written informed consent was obtained from each patient and the study protocol conformed to the ethics guidelines of the Declaration of Helsinki and was approved by the institutional ethics review committees. HBsAg-positive and/or anti-HIV-positive patients were excluded from this study.

In the following stage of replication study, SNP genotyping in an independent set of 391 Japanese HCV patients treated with PEG-IFN plus RBV treatment was completed using the DigiTag2 or TaqMan assay (ABI) following the manufacturer's protocol. The characteristics of patients for each GWAS stage and replication stage are summarized in Table 1.

### SNP genotyping and data cleaning

In the GWAS stage, we genotyped 303 Japanese HCV patients with and without the decrease of platelet counts from baseline to week 4 of PEG-IFN/RBV treatment [107 patients with a decrease of  $>30$  ( $10^9/l$ ) in platelet counts and 196 patients without a decrease of  $>30$  ( $10^9/l$ ) in platelet counts], using the Affymetrix Genome-Wide Human SNP Array 6.0 according to the manufacturer's instructions. The cut-off value was calculated to maximize the difference, which was also close to the median change. The average overall call rate of patients with and without the decrease of PLT reached 98.69 and 98.72%, respectively. We then applied the following thresholds for SNP QC in data cleaning: SNP call rate  $\geq 95\%$  for all samples, MAF  $\geq 1\%$  for all samples. A total of 595 052 SNPs on autosomal chromosomes passed the QC filters and were used for association analysis. All cluster

plots of SNPs showing  $P < 0.0001$  in association analyses by comparing allele frequencies in both groups with and without the decrease of PLT were checked by visual inspection, and SNPs with ambiguous genotype calls were excluded.

In the following stage of the replication study and high-density association mapping, we selected 23 tag SNPs from the 44.7 kb region, including *DDRGK1* gene and *ITPA* gene by analyzing LD and haplotype structure based on the HapMap data of Japanese, using the Haploview software. Of these tag SNPs, rs1127354 within the *ITPA* gene, which was associated with RBV-induced anemia (22), was included; however, rs7270101 was excluded because recent papers studying Japanese patients showed no variants in rs7270101 (26,31,32). The SNP genotyping in an independent set of 391 Japanese HCV patients with and without quantitative change in PLT levels from baseline to week 4 of PEG-IFN/RBV treatment (175 patients with quantitative change in PLT and 216 patients without quantitative change in PLT) was completed using the DigiTag2 assay (36). Twenty-two of the 23 SNPs were successfully analyzed and were used for SNP genotyping and data cleaning. All 22 SNPs in the replication study cleared HWE  $P$ -value  $> 0.001$ .

Based on the above SNPs data obtained from 303 Japanese HCV patients, using the Affymetrix Genome-Wide Human SNP Array 6.0, we also performed GWAS between 94 patients with a quantitative change of  $>3$  g of reduction in Hb and 209 patients without quantitative change in Hb levels from baseline to week 4 of PEG-IFN/RBV treatment. SNP genotyping in an independent set of 391 Japanese HCV patients with and without quantitative change in Hb levels from baseline to week 4 of PEG-IFN/RBV treatment (137 patients with quantitative change in Hb and 254 patients without quantitative change in Hb) was also completed using the DigiTag2 assay (36). Twenty-two of the 23 SNPs were successfully analyzed and were used for SNP genotyping and data cleaning.

An application of the Cochran–Armitage test on all the SNPs showed the genetic inflation factor  $\lambda = 1.000$  for thrombocytopenia and  $\lambda = 1.006$  for anemia in the GWAS stage (Supplementary Material, Figs S1 and S2). In addition, principal component analysis was performed in 303 samples for the GWAS stage together with the HapMap samples (CEU, YRI, CHB and JPT) (Supplementary Material, Fig. S3). These results implied that the effect of population stratification was negligible, except one sample, which was excluded from further analysis.

### Laboratory and histological tests

Blood samples were obtained at baseline, 1, 2, 4, 8 and 12 weeks after the start of therapy and for hematologic tests after the start of therapy and for hematologic tests, blood chemistry and HCV-RNA. Genetic polymorphism in the *IL28B* gene (rs8099917) was determined using the ABI TaqMan assay (Applied Biosystems, Carlsbad, CA, USA). Fibrosis was evaluated on a scale of 0–4 according to the METAVIR scoring system. The SVR was defined as an undetectable HCV-RNA level by qualitative PCR with a lower detection limit of 50 IU/ml (Amplicor, Roche Diagnostic Systems, CA, USA) or by Cobas Ampliprep/Cobas TaqMan assay (CAP/CTM) with a lower detection limit of



15 IU/ml (Roche Diagnostic Systems) 24 weeks after the completion of therapy.

### Statistical analysis

The observed association between an SNP and the decrease of platelets/quantitative change in Hb levels with response to PEG-IFN plus RBV treatment was assessed by  $\chi^2$  test with a two-by-two contingency table in three genetic models: allele frequency model, dominant-effect model and recessive-effect model. SNPs on chromosome X were removed because gender was not matched between groups with and without the decrease of PLT and quantitative change in Hb levels. A total of 595 052 SNPs passed the quality control filters in the GWAS stage; therefore, significance levels after Bonferroni correction for multiple testing were  $P = 8.40 \times 10^{-8}$  (0.05/595052) in the GWAS stage and  $P = 2.27 \times 10^{-3}$  (0.05/22) in the replication stage.

The association between an SNP of the *ITPA* gene (rs1127354) and the incidence of platelet reduction at week 4 was analyzed by Fisher's exact test. The association between *ITPA* polymorphisms and the degree of reduction in platelet counts and Hb levels at each time point during therapy were analyzed by Mann-Whitney *U* test. Multivariable regression analysis was used to analyze the factors associated with *ITPA*, the rs1127354 genotype, factors associated with platelet count reductions and factors associated with the reactive increase in platelet counts. IBM-SPSS software v.15.0 (SPSS, Inc., Chicago, IL, USA) was used for these analyses.

Possible heterogeneity in allele frequencies at rs1127354 was assessed by Tarone's test. The association between the SNP and thrombocytopenia/anemia were analyzed by the Cochran-Mantel-Haenszel test. Both analyses were performed using the R (version 2.9.0) software (Supplementary Material, Table S3).

### AUTHORS' CONTRIBUTIONS

Drafting of the paper, statistical analysis and approval of the final draft submitted: M.M.; drafting of the paper, statistical analysis, collecting samples and clinical data and approval of the final draft submitted: Y.T. and M.K.; statistical analysis and approval of the final draft submitted: N.N., M.S. and K.T.; collecting samples and clinical data and approval of the final draft submitted: K.M., N.S., N.E., H.Y., S.N., K.H., S.H., Y.I., E.T., S.M., M.H., Y.H., F.S., S.K. and N.I.

### SUPPLEMENTARY MATERIAL

Supplementary Material is available at *HMG* online.

### ACKNOWLEDGEMENTS

This study is based on 14 multi-center hospitals throughout Japan: Hokkaido area (Hokkaido University Hospital), Kanto area (Saitama University Hospital; Konodai Hospital; Musashino Red Cross Hospital; Tokyo Medical and Dental University Hospital; Yamanashi University Hospital), Koshin area (Shinshu University Hospital; Kanazawa University

Hospital), Tokai area (Nagoya City University Hospital), Kinki area (Kyoto Prefectural University of Medicine Hospital; Hyogo College of Medicine Hospital), Chugoku/Shikoku area (Ehime University Hospital; Kawasaki Medical College Hospital) and Kyushu area (National Nagasaki Medical Center). We thank Ms Yasuka Uehara-Shibata, Yuko Ogasawara-Hirano, Yoshimi Ishibashi, Natsumi Baba and Megumi Yamaoka-Sageshima (Tokyo University) for technical assistance. We also thank Dr Masaaki Korenaga (Kawasaki), Dr Akihiro Matsumoto (Shinshu), Dr Kayoko Naiki (Saitama), Dr Takeshi Nishimura (Kyoto), Dr Hirayuki Enomoto (Hyogo), Dr Minako Nakagawa (Tokyo Medical and Dental University) and Ochanomizu Liver Conference Study Group for collecting samples, and Dr Mamoru Watanabe (Tokyo Medical and Dental University) and Dr Moriichi Onji (Ehime University) for their advice throughout the study.

*Conflict of Interest statement.* Y.T., E.T. and S.K. are currently conducting research sponsored by Merck Sharp & Dohme, Corp. and Chugai Pharmaceutical Co. Ltd. The other co-authors have no conflict of interest.

### FUNDING

This study was supported by a grant-in-aid from the Ministry of Health, Labour, and Welfare of Japan (H22-kannen-005), and the Ministry of Education, Culture, Sports, Science, and Technology.

### REFERENCES

1. Global Burden of Hepatitis C Working Group (2004) Global burden of disease (GBD) for hepatitis C. *J. Clin. Pharmacol.*, **44**, 20–29.
2. Shiratori, Y., Shiina, S., Imamura, M., Kato, N., Kanai, F., Okudaira, T., Teratani, T., Tohgo, G., Toda, N., Ohashi, M. *et al.* (1995) Characteristic difference of hepatocellular carcinoma between hepatitis B- and C-viral infection in Japan. *Hepatology*, **22**, 1027–1033.
3. Yoshida, H., Tateishi, R., Arakawa, Y., Sata, M., Fujiyama, S., Nishiguchi, S., Ishibashi, H., Yamada, G., Yokosuka, O., Shiratori, Y. *et al.* (2004) Benefit of interferon therapy in hepatocellular carcinoma prevention for individual patients with chronic hepatitis C. *Gut*, **53**, 425–430.
4. George, S.L., Bacon, B.R., Brunt, E.M., Mihindukulasuriya, K.L., Hoffmann, J. and Di Bisceglie, A.M. (2009) Clinical, virologic, histologic, and biochemical outcomes after successful HCV therapy: a 5-year follow-up of 150 patients. *Hepatology*, **49**, 729–738.
5. Fried, M.W., Shiffman, M.L., Reddy, K.R., Smith, C., Marinos, G., Goncalves, F.L. Jr, Haussinger, D., Diago, M., Carosi, G., Dhumeaux, D. *et al.* (2002) Peginterferon alpha-2a plus ribavirin for chronic hepatitis C virus infection. *N. Engl. J. Med.*, **347**, 975–982.
6. Manns, M.P., McHutchison, J.G., Gordon, S.C., Rustgi, V.K., Shiffman, M., Reindollar, R., Goodman, Z.D., Koury, K., Ling, M. and Albrecht, J.K. (2001) Peginterferon alpha-2b plus ribavirin compared with interferon alpha-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomised trial. *Lancet*, **358**, 958–965.
7. Hadziyannis, S.J., Sette, H. Jr, Morgan, T.R., Balan, V., Diago, M., Marcellin, P., Ramadori, G., Bodenheimer, H. Jr, Bernstein, D., Rizzetto, M. *et al.* (2004) Peginterferon-alpha2a and ribavirin combination therapy in chronic hepatitis C: a randomized study of treatment duration and ribavirin dose. *Ann. Intern. Med.*, **140**, 346–355.
8. Hiramatsu, N., Oze, T., Tsuda, N., Kurashige, N., Koga, K., Toyama, T., Yasumaru, M., Kanto, T., Takehara, T., Kasahara, A. *et al.* (2006) Should aged patients with chronic hepatitis C be treated with interferon and ribavirin combination therapy? *Hepatol. Res.*, **35**, 185–189.
9. Iwasaki, Y., Ikeda, H., Araki, Y., Osawa, T., Kita, K., Ando, M., Shimoe, T., Takaguchi, K., Hashimoto, N., Kobatake, T. *et al.* (2006) Limitation of

- combination therapy of interferon and ribavirin for older patients with chronic hepatitis C. *Hepatology*, **43**, 54–63.
10. Sezaki, H., Suzuki, F., Akuta, N., Yatsuji, H., Hosaka, T., Kobayashi, M., Suzuki, Y., Arase, Y., Ikeda, K., Miyakawa, Y. *et al.* (2009) An open pilot study exploring the efficacy of fluvastatin, pegylated interferon and ribavirin in patients with hepatitis C virus genotype 1b in high viral loads. *Intervirology*, **52**, 43–48.
  11. Bruno, R., Sacchi, P., Maiocchi, L., Patrino, S. and Filice, G. (2006) Hepatotoxicity and antiretroviral therapy with protease inhibitors: a review. *Dig. Liver Dis.*, **38**, 363–373.
  12. Hezode, C., Forestier, N., Dusheiko, G., Ferenci, P., Pol, S., Goeser, T., Bronowicki, J.P., Bourliere, M., Gharakhanian, S., Bengtsson, L. *et al.* (2009) Telaprevir and peginterferon with or without ribavirin for chronic HCV infection. *N. Engl. J. Med.*, **360**, 1839–1850.
  13. McHutchison, J.G., Everson, G.T., Gordon, S.C., Jacobson, I.M., Sulkowski, M., Kauffman, R., McNair, L., Alam, J. and Muir, A.J. (2009) Telaprevir with peginterferon and ribavirin for chronic HCV genotype 1 infection. *N. Engl. J. Med.*, **360**, 1827–1838.
  14. Suzuki, F., Akuta, N., Suzuki, Y., Sezaki, H., Yatsuji, H., Kawamura, Y., Hosaka, T., Kobayashi, M., Arase, Y., Ikeda, K. *et al.* (2009) Rapid loss of hepatitis C virus genotype 1b from serum in patients receiving a triple treatment with telaprevir (MP-424), pegylated interferon and ribavirin for 12 weeks. *Hepatol. Res.*, **39**, 1056–1063.
  15. Sakamoto, N. and Watanabe, M. (2009) New therapeutic approaches to hepatitis C virus. *J. Gastroenterol.*, **44**, 643–649.
  16. Afdhal, N.H., McHutchison, J.G., Zeuzem, S., Mangia, A., Pawlotsky, J.M., Murray, J.S., Shianna, K.V., Tanaka, Y., Thomas, D.L., Booth, D.R. *et al.* (2010) Hepatitis C pharmacogenetics: state of the art in 2010. *Hepatology*, **53**, 336–345.
  17. Tanaka, Y., Nishida, N., Sugiyama, M., Kurosaki, M., Matsuura, K., Sakamoto, N., Nakagawa, M., Korenaga, M., Hino, K., Hige, S. *et al.* (2009) Genome-wide association of IL28B with response to pegylated interferon-alpha and ribavirin therapy for chronic hepatitis C. *Nat. Genet.*, **41**, 1105–1109.
  18. Ge, D., Fellay, J., Thompson, A.J., Simon, J.S., Shianna, K.V., Urban, T.J., Heinzen, E.L., Qiu, P., Bertelsen, A.H., Muir, A.J. *et al.* (2009) Genetic variation in IL28B predicts hepatitis C treatment-induced viral clearance. *Nature*, **461**, 399–401.
  19. Suppiah, V., Moldovan, M., Ahlenstiel, G., Berg, T., Weltman, M., Abate, M.L., Bassendine, M., Spengler, U., Dore, G.J., Powell, E. *et al.* (2009) IL28B is associated with response to chronic hepatitis C interferon-alpha and ribavirin therapy. *Nat. Genet.*, **41**, 1100–1104.
  20. Thomas, D.L., Thio, C.L., Martin, M.P., Qi, Y., Ge, D., O’Huigin, C., Kidd, J., Kidd, K., Khakoo, S.I., Alexander, G. *et al.* (2009) Genetic variation in IL28B and spontaneous clearance of hepatitis C virus. *Nature*, **461**, 798–801.
  21. Tanaka, Y., Nishida, N., Sugiyama, M., Tokunaga, K. and Mizokami, M. (2010) lambda-Interferons and the single nucleotide polymorphisms: a milestone to tailor-made therapy for chronic hepatitis C. *Hepatol. Res.*, **40**, 449–460.
  22. Fellay, J., Thompson, A.J., Ge, D., Gumbs, C.E., Urban, T.J., Shianna, K.V., Little, L.D., Qiu, P., Bertelsen, A.H., Watson, M. *et al.* (2010) ITPA gene variants protect against anaemia in patients treated for chronic hepatitis C. *Nature*, **464**, 405–408.
  23. Afdhal, N., McHutchison, J., Brown, R., Jacobson, I., Manns, M., Poordad, F., Weksler, B. and Esteban, R. (2008) Thrombocytopenia associated with chronic liver disease. *J. Hepatol.*, **48**, 1000–1007.
  24. Wazny, L.D. and Ariano, R.E. (2000) Evaluation and management of drug-induced thrombocytopenia in the acutely ill patient. *Pharmacotherapy*, **20**, 292–307.
  25. Hitomi, Y., Cirulli, E.T., Fellay, J., McHutchison, J.G., Thompson, A.J., Gumbs, C.E., Shianna, K.V., Urban, T.J. and Goldstein, D.B. (2011) Inosine triphosphate protects against ribavirin-induced adenosine triphosphate loss by adenylosuccinate synthase function. *Gastroenterology*, **140**, 1314–1321.
  26. Ochi, H., Maekawa, T., Abe, H., Hayashida, Y., Nakano, R., Kubo, M., Tsunoda, T., Hayes, C.N., Kumada, H., Nakamura, Y. *et al.* (2010) ITPA polymorphism affects ribavirin-induced anemia and outcomes of therapy—a genome-wide study of Japanese HCV virus patients. *Gastroenterology*, **139**, 1190–1197.
  27. Ong, J.P. and Younossi, Z.M. (2004) Managing the hematologic side effects of antiviral therapy for chronic hepatitis C: anemia, neutropenia, and thrombocytopenia. *Cleve. Clin. J. Med.*, **71** (Suppl. 3), S17–S21.
  28. Canonico, P.G., Castello, M.D., Cosgriff, T.M., Donovan, J.C., Ross, P.E., Spears, C.T. and Stephen, E.L. (1984) Hematological and bone marrow effects of ribavirin in rhesus monkeys. *Toxicol. Appl. Pharmacol.*, **74**, 163–172.
  29. Akan, H., Guven, N., Aydogdu, I., Arat, M., Beksac, M. and Dalva, K. (2000) Thrombopoietic cytokines in patients with iron deficiency anemia with or without thrombocytosis. *Acta Haematol.*, **103**, 152–156.
  30. Wu, J., Lei, G., Mei, M., Tang, Y. and Li, H. (2010) A novel C53/LZAP-interacting protein regulates stability of C53/LZAP and DDRGK domain-containing protein 1 (DDRGK1) and modulates NF-kappaB signaling. *J. Biol. Chem.*, **285**, 15126–15136.
  31. Sakamoto, N., Tanaka, Y., Nakagawa, M., Yatsuhashi, H., Nishiguchi, S., Enomoto, N., Azuma, S., Nishimura-Sakurai, Y., Kakinuma, S., Nishida, N. *et al.* (2010) ITPA gene variant protects against anemia induced by pegylated interferon-alpha and ribavirin therapy for Japanese patients with chronic hepatitis C. *Hepatol. Res.*, **40**, 1063–1071.
  32. Kurosaki, M., Tanaka, Y., Tanaka, K., Suzuki, Y., Hoshioka, Y., Tamaki, N., Kato, T. and Yasui, Y. (2011) Analysis of the correlations between genetic polymorphisms of the ITPA gene and hemolytic anemia or outcome after treatment with pegylated-interferon and ribavirin in genotype 1b chronic hepatitis C. *Antivir. Ther.*, in press.
  33. Shipkova, M., Lorenz, K., Oellerich, M., Wieland, E. and von Ahsen, N. (2006) Measurement of erythrocyte inosine triphosphate pyrophosphohydrolase (ITPA) activity by HPLC and correlation of ITPA genotype-phenotype in a Caucasian population. *Clin. Chem.*, **52**, 240–247.
  34. Fraser, J.H., Meyers, H., Henderson, J.F., Brox, L.W. and McCoy, E.E. (1975) Individual variation in inosine triphosphate accumulation in human erythrocytes. *Clin. Biochem.*, **8**, 353–364.
  35. Kumada, H., Okanoue, T., Onji, M., Moriwaki, H., Izumi, N., Tanaka, E., Chayama, K., Sakisaka, S., Takehara, T., Oketani, M. *et al.* (2010) Guidelines for the treatment of chronic hepatitis and cirrhosis due to hepatitis C virus infection for the fiscal year 2008 in Japan. *Hepatol. Res.*, **40**, 8–13.
  36. Nishida, N., Tanabe, T., Takasu, M., Suyama, A. and Tokunaga, K. (2007) Further development of multiplex single nucleotide polymorphism typing method, the DigiTag2 assay. *Anal. Biochem.*, **364**, 78–85.



# Comparative Analysis of Various Tumor-Associated Antigen-Specific T-Cell Responses in Patients with Hepatocellular Carcinoma

Eishiro Mizukoshi, Yasunari Nakamoto, Kuniaki Arai, Tatsuya Yamashita, Akito Sakai, Yoshio Sakai, Takashi Kagaya, Taro Yamashita, Masao Honda, and Shuichi Kaneko

Many tumor-associated antigens (TAAs) recognized by cytotoxic T cells (CTLs) have been identified during the last two decades and some of them have been used in clinical trials. However, there are very few in the field of immunotherapy for hepatocellular carcinoma (HCC) because there have not been comparative data regarding CTL responses to various TAAs. In the present study, using 27 peptides derived from 14 different TAAs, we performed comparative analysis of various TAA-specific T-cell responses in 31 HCC patients to select useful antigens for immunotherapy and examined the factors that affect the immune responses to determine a strategy for more effective therapy. Twenty-four of 31 (77.4%) HCC patients showed positive responses to at least one TAA-derived peptide in enzyme-linked immunospot assay. The TAAs consisting of cyclophilin B, squamous cell carcinoma antigen recognized by T cells (SART) 2, SART3, p53, multidrug resistance-associated protein (MRP) 3, alpha-fetoprotein (AFP) and human telomerase reverse transcriptase (hTERT) were frequently recognized by T cells and these TAA-derived peptides were capable of generating peptide-specific CTLs in HCC patients, which suggested that these TAAs are immunogenic. HCC treatments enhanced TAA-specific immune responses with an increased number of memory T cells and induced *de novo* T-cell responses to lymphocyte-specific protein tyrosine kinase, human epidermal growth factor receptor type 2, p53, and hTERT. Blocking cytotoxic T-lymphocyte antigen-4 (CTLA-4) resulted in unmasking of TAA-specific immune responses by changing cytokine and chemokine profiles of peripheral blood mononuclear cells stimulated by TAA-derived peptides. **Conclusion:** Cyclophilin B, SART2, SART3, p53, MRP3, AFP, and hTERT were immunogenic targets for HCC immunotherapy. TAA-specific immunotherapy combined with HCC treatments and anti-CTLA-4 antibody has the possibility to produce stronger tumor-specific immune responses. (HEPATOLOGY 2011;53:1206-1216)

Hepatocellular carcinoma (HCC) is the most common primary malignancy of the liver and becoming an important public health concern.<sup>1,2</sup> Although many kinds of treatments have

been performed for HCC, their effects are limited because the recurrence rate of HCC is very high; therefore, the development of new therapeutic options to prevent recurrence is necessary.<sup>3,4</sup>

*Abbreviations:* AFP, alpha-fetoprotein; CTL, cytotoxic T cell; ELISPOT, enzyme-linked immunospot; HCC, hepatocellular carcinoma; HCV, hepatitis C virus; HLA, human leukocyte antigen; hTERT, human telomerase reverse transcriptase; IFN, interferon; Lck, lymphocyte-specific protein tyrosine kinase; MRP, multidrug resistance-associated protein; PBMC, peripheral blood mononuclear cell; TAA, tumor-associated antigen.

From the Department of Gastroenterology, Graduate School of Medicine, Kanazawa University, Kanazawa, Ishikawa, Japan.

Received August 6, 2010; accepted December 14, 2010.

Address reprint requests to: Shuichi Kaneko, M.D., Department of Gastroenterology, Graduate School of Medicine, Kanazawa University, Kanazawa, Ishikawa 920-8641, Japan. E-mail: skaneko@m-kanazawa.jp; fax: 81-76-234-4250.

Copyright © 2011 by the American Association for the Study of Liver Diseases.

View this article online at [wileyonlinelibrary.com](http://wileyonlinelibrary.com).

DOI 10.1002/hep.24149

Potential conflict of interest: Nothing to report.

To protect against recurrence, tumor antigen-specific immunotherapy is an attractive strategy. Many tumor-associated antigens (TAAs) and their epitopes recognized by cytotoxic T cells (CTLs) have been identified during the last two decades and some of them have been used in clinical trials for several cancers.<sup>5-21</sup> The epitopes have been under investigation for the treatment of cancer, with major clinical responses in some trials.<sup>22,23</sup> With regard to immunotherapy for HCC, few kinds of TAAs and their epitopes have been used and only clinical data of  $\alpha$ -fetoprotein (AFP) have been reported.<sup>24,25</sup> In human trials targeting AFP, it is possible to raise an AFP-specific T-cell response using AFP-derived peptides, but this has shown little

antitumor effect. On the other hand, immunotherapy trials using autologous tumor lysate or dendritic cells have shown statistically significant improvements in the risk of HCC recurrence and recurrence-free survival.<sup>26</sup> These reports suggest that tumor antigen-specific immunotherapy is effective to reduce the recurrence rate after HCC treatment; therefore, it is necessary to find immunogenic antigens or their epitopes to develop more effective immunotherapy.

In addition, in the field of molecular targeting therapies, developments of monoclonal antibodies targeting immunomodulatory molecules to enhance anti-tumor immunity are progressing and some of these are under clinical trial.<sup>27</sup> In particular, clinical data of anti-cytotoxic T-lymphocyte antigen-4 (anti-CTLA-4) antibody have shown durable objective response and stable disease in melanoma patients.<sup>28</sup>

In the present study we performed comparative analysis of various TAA-specific T-cell responses in patients with HCC and examined the factors that affect the immune responses, including anti-CTLA-4 antibody. This approach offers useful information to select immunogenic TAAs and to develop a new strategy for HCC immunotherapy.

## Patients and Methods

**Patients and Laboratory Testing.** In this study we examined 31 human leukocyte antigen (HLA)-A24-positive patients with HCC, 29 chronic hepatitis C patients without HCC, who were diagnosed by liver biopsy, and 11 healthy blood donors who did not have a history of cancer and were negative for hepatitis B surface antigen and anti-hepatitis C virus (HCV) antibody (Ab). The diagnosis of HCC was histologically confirmed in 21 patients. For the remaining 10 patients the diagnosis was based on typical hypervascular tumor staining on angiography in addition to typical findings, which showed hyperattenuated areas in the early phase and hypoattenuation in the late phase on dynamic computed tomography (CT).<sup>29</sup>

HLA-based typing of peripheral blood mononuclear cells (PBMCs) from patients and normal blood donors was performed as described.<sup>19</sup> The pathological grading of tumor cell differentiation was assessed according to the general rules for the clinical and pathological study of primary liver cancer.<sup>30</sup> The severity of liver disease was evaluated according to the criteria of Desmet et al.<sup>31</sup> using biopsy specimens of liver tissue.

All patients gave written informed consent to participate in the study in accordance with the Helsinki Declaration and this study was approved by the re-

**Table 1. Peptides**

Peptide No.	Peptide Name	Source	Reference	Amino Acid Sequence	Number of Specific Spots in Normal Donors (Mean SD)
1	ART1 <sub>188</sub>	ART1	5	EYCLKFTKL	0.9 ± 1.1
2	ART4 <sub>161</sub>	ART4	6	AFLRHAAL	0.3 ± 0.5
3	ART4 <sub>899</sub>	ART4	6	DYPSLSATDI	0.6 ± 1.0
4	Cyp-B <sub>109</sub>	Cyp-B	7	KFHRVIKDF	0.5 ± 0.9
5	Cyp-B <sub>315</sub>	Cyp-B	7	DFMIQGGDF	1.2 ± 1.7
6	Lck <sub>208</sub>	Lck	8	HYTNASDGL	0.3 ± 0.6
7	Lck <sub>486</sub>	Lck	8	TFDYLRSVL	0.2 ± 0.8
8	Lck <sub>488</sub>	Lck	8	DYLRVLEDF	0.9 ± 1.5
9	MAGE1 <sub>135</sub>	MAGE-A1	9	NYKHCPEI	1.0 ± 0.9
10	MAGE3 <sub>195</sub>	MAGE-A3	10	IMPKAGLLI	1.4 ± 1.7
11	SART1 <sub>1690</sub>	SART1	11	EYRGFTQDF	0.9 ± 1.3
12	SART2 <sub>899</sub>	SART2	12	SYTRLFLIL	1.0 ± 1.4
13	SART3 <sub>109</sub>	SART3	13	VDYNDCHVDL	2.1 ± 1.9
14	Her-2/neu <sub>8</sub>	Her-2/neu	14	RWGLLLALL	1.4 ± 2.0
15	p53 <sub>125</sub>	p53	15	TYPALNKMF	1.4 ± 1.5
16	p53 <sub>161</sub>	p53	16	AIYKQSQHM	0.4 ± 0.6
17	p53 <sub>204</sub>	p53	17	EYLDNRNTE	1.1 ± 1.5
18	p53 <sub>211</sub>	p53	17	TFRHSVVV	0.9 ± 1.9
19	p53 <sub>235</sub>	p53	17	NYMCNSSCM	2.1 ± 2.6
20	MRP3 <sub>503</sub>	MRP3	18	LYAWEPSFL	0.2 ± 0.5
21	MRP3 <sub>692</sub>	MRP3	18	AYVPPQAWI	1.5 ± 2.1
22	MRP3 <sub>765</sub>	MRP3	18	VYSDADIFL	0.9 ± 1.0
23	AFP <sub>357</sub>	AFP	19	EYSRRHPQL	1.8 ± 2.0
24	AFP <sub>403</sub>	AFP	19	KYIQESQAL	1.1 ± 1.5
25	AFP <sub>434</sub>	AFP	19	AYTKKAPQL	0.8 ± 1.1
26	hTERT <sub>167</sub>	hTERT	20	AYQVCGPPL	0.8 ± 1.1
27	hTERT <sub>324</sub>	hTERT	20	VYAETKHFL	0.5 ± 0.7
28	HIV env <sub>584</sub>	HIV env	32	RYLRDQQLL	1.3 ± 2.0
29	HCV NS3 <sub>1031</sub>	HCV NS3	33	AYSQQTIRGL	ND
30	CMV pp65 <sub>328</sub>	CMV pp65	34	QYDPVAALF	13.3 ± 15.7

ND, not determined.

gional ethics committee (Medical Ethics Committee of Kanazawa University, No. 829).

**Peptides, Cell Lines, and Preparation of PBMCs.** Twenty-seven peptides derived from 14 different TAAs (Table 1), human immunodeficiency virus (HIV) envelope-derived peptide (HIVenv<sub>584</sub>),<sup>32</sup> HCV NS3-derived peptide (HCVNS3<sub>1031</sub>),<sup>33</sup> and cytomegalovirus (CMV) pp65-derived peptide (CMVpp65<sub>328</sub>),<sup>34</sup> which were identified as HLA-A24 restricted CTL epitopes in previous studies, were used. Peptides were synthesized at Mimotope (Melbourne, Australia) and Sumitomo Pharmaceuticals (Osaka, Japan). They were identified using mass spectrometry and their purities were determined to be >80% by analytical high-performance liquid chromatography (HPLC). The HLA-A\*2402 gene-transfected C1R cell line (C1R-A24) was cultured in RPMI 1640 medium containing 10% fetal calf serum (FCS) and 500 µg/mL hygromycin B (Sigma, St. Louis, MO), and K562 was cultured in RPMI 1640 medium containing 10% FCS.<sup>35</sup> PBMCs were isolated before HCC treatments as described.<sup>20</sup> In 12 patients their PBMCs were also obtained 4 weeks after treatments.

**Table 2. Characteristics of the Patients Studied**

Clinical Diagnosis	No. of		Age (yr) Mean $\pm$ SD	ALT (IU/L) Mean $\pm$ SD	AFP (ng/ml) Mean $\pm$ SD	Child Pugh (A/B/C)	Diff. Degree* (wel/mod/ por/ND)	Tumor Size† (large/small)	Tumor Multiplicity (multiple/solitary)	Vascular Invasion (+/-)	TNM Stage (I/II/IIIa/IIIB/ IIIC/IV)
	Patients	Sex M/F									
Normal donors	11	8/3	35 $\pm$ 2	ND	ND	ND	ND	ND	ND	ND	ND
Chronic hepatitis	29	16/13	59 $\pm$ 10	92 $\pm$ 94	31 $\pm$ 87	27/2/0	ND	ND	ND	ND	ND
HCC	31	23/8	71 $\pm$ 4	74 $\pm$ 33	1768 $\pm$ 9103	20/10/1	11/10/0/10	22/9	20/11	9/22	10/12/3/1/2/3

\*Histological degree of HCC; wel: well differentiated, mod: moderately differentiated, por: poorly differentiated, ND: not determined.

†Tumor size was divided into either "small" ( $\leq 2$  cm) or "large" ( $> 2$  cm).

**CTL Induction and Cytotoxicity Assay.** CTL induction and cytotoxicity assays were performed as described.<sup>20</sup> Briefly, stimulated PBMCs were added at effector to target ratios of 100:1, 50:1, 25:1, 13:1, 6:1, and 3:1. In cases where the number of CTLs was insufficient, cytotoxicity assays were performed at effector to target ratios less than 100:1.

**Interferon Gamma IFN- $\gamma$  Enzyme-Linked Immunospot (ELISPOT) Assay.** IFN- $\gamma$  ELISPOT assays were performed as reported.<sup>20</sup> Responses to TAA-derived peptides were considered positive if more than 10 specific spots were detected, which is greater than the mean plus 3 standard deviations (SDs) of the baseline response detected in 11 normal blood donors (Table 1), and if the number of spots in the presence of an antigen was at least 2-fold that in its absence. Responses to HIV-, HCV-, and CMV-derived peptides were considered positive if more than 10 specific spots were detected and if the number of spots in the presence of an antigen was at least 2-fold that in its absence. In ELISPOT assay with blocking CTLA-4, anti-human CTLA-4 (eBioscience, Tokyo, Japan) was added at a final concentration of 50  $\mu$ g/mL, which has been described to have maximum effect in *in vitro* cultures.<sup>36</sup> As a control, functional grade mouse immunoglobulin G (IgG)2a isotype control was used. The assay with blocking CTLA-4 was performed in triplicate and the results were statistically analyzed using the unpaired Student's *t* test.

**Cytokine and Chemokine Profiling.** The effect of CTLA-4 antibody on TAA-specific T-cell responses was also analyzed by cytokine and chemokine profiling. Cytokine and chemokine levels in the medium of ELISPOT assay were measured using the Bio-plex assay (Bio-Rad, Hercules, CA). These included interleukin (IL)-1 $\beta$ , IL-1Ra, IL-2, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-12p70, IL-13, IL-15, IL-17, basic fibroblast growth factor (FGF), eotaxin, G-CSF, GM-CSF, IFN- $\gamma$ , IP-10, MCP-1, macrophage inflammatory protein (MIP)-1 $\alpha$ , MIP-1 $\beta$ , platelet-derived growth factor (PDGF)-BB, RANTES, tumor necrosis factor alpha (TNF- $\alpha$ ), and vascular endothelial growth

factor (VEGF). Eight standards (ranging from 2 to 32,000 pg/mL) were used to generate calibration curves for each cytokine. Data acquisition and analysis were carried out using Bio-plex Manager software v. 4.1.1.

**Cytokine Secretion Assay.** TAA-specific IFN- $\gamma$ -producing T cells were also analyzed by cytokine secretion assay. The assay was performed with the MACS cytokine secretion assay (Miltenyi Biotec K.K., Tokyo, Japan), in accordance with the manufacturer's instructions. Briefly, 5,000,000 PBMCs were pulsed with TAA-derived peptides for 16 hours and then incubated with 20  $\mu$ L of IFN- $\gamma$  detection antibody, 10  $\mu$ L of anti-CD8-APC Ab (Becton Dickinson, Tokyo, Japan), 10  $\mu$ L of anti-CCR7-FITC Ab (eBioscience, Tokyo, Japan), and 10  $\mu$ L of anti-CD45RA-PerCP-Cy5.5 Ab (eBioscience, Tokyo, Japan) for 10 minutes at 4°C. After washing with a cold buffer (phosphate-buffered saline/0.5% bovine serum albumin with 2 mM EDTA), the cells were resuspended with 500  $\mu$ L of cold buffer and analyzed using FACSCalibur (Becton Dickinson, Tokyo, Japan). As a positive control, CMVpp65<sub>328</sub>-specific IFN- $\gamma$ -producing T cells were also analyzed by the same methods. The number of IFN- $\gamma$ -producing T cells was calculated from the results of FACS analysis and is shown as a number per 300,000 PBMCs.

## Results

**Patient Profile.** The clinical profiles of the 11 healthy blood donors, 29 patients with chronic hepatitis C, and 31 patients with HCV-related HCC analyzed in the present study are shown in Table 2 and Fig. 1. Using TNM staging of the Union Internationale Contre Le Cancer (UICC) system (6th v.), 10, 12, 3, 1, 2, and 3 patients were classified as having stage I, II, IIIA, IIIB, IIIC, and IV tumors, respectively.

**Detection of TAA-Specific T Cells in HCC Patients.** First we examined the frequency of cells that specifically reacted with TAA-derived and control peptides in HCC patients. Fifty-one responses in total were observed against TAA-derived peptides. Twenty-




Fig. 1. TAA-, HIV-, HCV-, and CMV-derived peptide-specific T-cell responses. Results of all HCC patients examined are shown. The T-cell responses were examined by IFN- $\gamma$  ELISPOT assay. Responses to peptides were considered positive if more than 10 specific spots per 300,000 PBMCs were detected and if the number of spots in the presence of an antigen was at least 2-fold that in its absence. Black boxes indicate the presence of a significant IFN- $\gamma$  T-cell response to peptides. Peptide sequences are described in Table 1 and characteristics of patients in Table 2.

four of 31 (77.4%) patients showed positive responses to at least one TAA-derived peptide and most of them showed responses to 1 to 4 kinds of TAA-derived peptide. Twenty-three of 27 (85.2%) TAA-derived peptides were recognized by T cells of at least one patient. Peptides 4, 12, 13, 16, 17, 22, 24, and 27 were recognized in more than two patients, suggesting that these peptides were immunogenic. Peptides 28 (HIV env<sub>584</sub>), 29 (HCV<sub>1031</sub>), and 30 (CMV pp65<sub>328</sub>) were recognized by 0 (0%), 5 (16%), and 13 (42%) patients, respectively.

The magnitude of TAA-specific T-cell responses was assessed by the frequency of peptide-specific IFN- $\gamma$ -producing T cells in the PBMC population (Fig. 2A). The range of TAA-derived peptide-specific T-cell frequency was 10-60.5 cells/300,000 PBMCs. Those specific to peptides 13 and 16 numbered more than 30 cells/300,000 PBMCs, suggesting that these peptides were immunogenic. The frequencies of T cells specific to HCV- and CMV-derived peptides were 12-22 cells and 12-92/300,000 PBMCs, respectively.

Whether these TAA-derived peptides were capable of generating peptide-specific CTLs from PBMCs was investigated in HCC patients. The seven peptides were selected according to the magnitude of TAA-specific T-cell responses determined by the fre-

quency of T cells with a positive response. The CTLs generated with these peptides were cytotoxic to C1RA24 cells pulsed with the corresponding peptides (Fig. 2B).

**Comparison of TAA-Specific T-Cell Responses Between the Patient Groups With and Without HCC.** To characterize the immunogenicity and specificity of TAA-derived peptides, we compared T-cell responses to the peptides derived from TAA, HIV, HCV, and CMV among three groups consisting of normal blood donors, patients with chronic hepatitis C, and patients with HCV-related HCC. A significant TAA-specific T-cell response was not detected in normal blood donors (Fig. 3A). A response was detected in both chronic hepatitis C and HCC patient groups, but it was more frequently observed in HCC patients. HIV-specific T-cell response was not detected in any group. HCV-specific T-cell response rate was not different between the groups with chronic hepatitis C and HCC. CMV-specific T-cell response rates were similar among the three groups. Similar tendencies were observed in the analysis of individual peptides (Fig. 3B). We also examined the frequency of T cells responsive to peptides among the three groups. The mean frequency of TAA-specific T cells without *in vitro* expansion was higher in HCC patients than in




Fig. 2. Vigor of TAA-, HCV-, and CMV-derived peptide-specific T-cell responses. (A) The frequency of TAA-specific IFN- $\gamma$ -producing T cells was analyzed by ELISPOT assay. Only positive responses are shown. Black bars indicate the response of one patient. \*1, \*2, \*3, \*4, and \*5 denote 33, 60.5, 44, 92, and 67.5 specific spots, respectively. (B) Representative TAA-specific T-cell responses were also analyzed by CTL assay. T cell lines were generated from PBMC of the HLA-A24-positive HCC patients by stimulation with TAA-derived peptides (peptides 4, 12, 13, 16, 22, 24, and 27) (see Table 1). Expanded T cell lines were then tested for specific cytotoxicity against the corresponding peptides in a standard  $^{51}\text{Cr}$  release assay at the indicated E:T ratios.

patients with chronic hepatitis C for 14 of 27 TAA-derived peptides (peptides 1, 2, 3, 4, 12, 16, 18, 19, 20, 21, 22, 24, 25 and 27) (Fig. 3C).

**Enhancement of TAA-Specific T-Cell Responses After HCC Treatments.** Several studies including our own have clarified that HCC treatments enhanced HCC-specific immune responses (19, 37, 38). In this study, we examined whether the enhancement was observed equally in all kinds of TAAs or specifically in some TAAs. For this purpose we measured the frequency of TAA-specific T cells before and after HCC treatment by ELISPOT assay in 12 cases who received transcatheter arterial embolization (TAE), radiofrequency ablation (RFA), or chemotherapy. The frequency of TAA-specific T cells increased in all patients and it was observed for 23 of 27 TAA-derived peptides (Fig. 4A). The enhancement was observed in the

patients who received TAE, RFA, or chemotherapy and even in the patients without an increase in the frequency of CMV-specific T cells. Peptides 7, 14, 15, and 26, which were not recognized by T cells in all HCC patients before treatments (Fig. 1), were recognized by T cells in 1, 4, 1, and 5, respectively, of 12 patients after treatments. Representative results of enhancement of TAA-specific immune responses are shown in Fig. 4B. The frequency of TAA-specific T cells increased to 11-80 cells/300,000 PBMCs after treatments.

The enhancement of TAA-specific immune responses was also confirmed by cytokine secretion assay. Representative results are shown in Fig. 4C. In this patient (patient 25) the frequency of TAA-specific IFN- $\gamma$ -producing CD8 $^{+}$  T cells was increased from 0.4% to 1.4% of CD8 $^{+}$  T cells after HCC treatment.




Fig. 3. Comparative analysis of TAA-, HIV-, HCV-, and CMV-derived peptide-specific T-cell responses among three groups of subjects: normal donors, patients with chronic hepatitis C not complicated by HCC, and HCC patients. (A) Summary of the number of patients with a significant IFN- $\gamma$  T-cell response to tumor-associated, HIV, HCV, and CMV antigens in each group. (B) Graph shows the percentage of patients in each group who showed a significant IFN- $\gamma$  T-cell response to individual peptides. Peptide sequences are described in Table 1. (C) Mean frequency of peptide-specific IFN- $\gamma$ -producing T cells in each group. The frequency of IFN- $\gamma$ -producing T cells was analyzed by ELISPOT assay.

In this assay we also examined the naïve/effector/memory phenotype of these cells by the criterion of CD45RA/CCR7 expression.<sup>39</sup> Phenotypic analysis of TAA-specific, IFN- $\gamma$ -producing memory CD8<sup>+</sup> T cells before and after treatment showed that the frequency of CD45RA<sup>-</sup>/CCR7<sup>+</sup> central memory T cells was the highest, indicating that the posttherapeutic increase in these T cells is due to the increase in cells with this phenotype (Fig. 4D). In this patient the number of T cells with the CD45RA<sup>-</sup>/CCR7<sup>+</sup> phenotype increased from 73 cells/300,000 PBMCs before treatment to 316 cells/300,000 PBMCs after treatment. Similar results were noted in five patients.

**Blocking CTLA-4 Restores TAA-Specific T-Cell Responses.** In previous studies including our own,<sup>19,20,24</sup> the CTL epitopes that correlate with the prevention of tumor progression or prognosis of HCC patients have not been identified. One of the reasons for this is considered to be that the naturally occurring

T-cell responses to the epitopes are weak; therefore, recent tumor immunotherapeutic studies are moving toward modulation of T-cell responses.

CTLA-4 is recognized as a critical negative regulator of immune response; therefore, its blockade has been considered to contribute to antitumor activity.<sup>27</sup> In a recent study it was reported that blocking of CTLA-4 on both effector and regulatory T cell compartments contributes to the antitumor activity of CTLA-4 antibodies.<sup>40</sup> To examine whether similar occurs for immune response in HCC patients, we analyzed 32 separate TAA-specific T-cell responses in 15 HCC patients using 13 TAA-derived peptides. Incubation of T cells with CTLA-4 antibodies resulted in an increase of the number of TAA-specific T cells in 18 of 32 (56%) responses and in 9 of 15 (60%) patients (Fig. 5A). Fourteen and four patients showed increases of 1-10 and more than 10 TAA-specific T cells, respectively. Representative results of six patients are shown






Fig. 4. Enhancement of TAA-specific T-cell responses in HCC patients after treatments. (A) Summary of patients and peptides with a significant increase of the number of IFN- $\gamma$ -producing T cells (black boxes). A significant change in the IFN- $\gamma$  response was defined as a more than 2-fold increase and the presence of more than 10 specific spots in ELISpot assay after HCC treatments. The assays were performed in 12 HCC patients using 27 TAA-, HIV-, and CMV-derived peptides. Goal\* shows the goal of HCC treatment. C and P denote "curative intention" and "palliative intention," respectively. (B) Representative results of ELISpot assay are shown. White and black bars indicate the frequency of T cells before and after HCC treatments, respectively. \*1, \*2, \*3, and \*4 denote 53, 60, 80, and 121 specific spots, respectively. (C) Enhancement of TAA-specific T-cell responses was also analyzed by cytokine secretion assay. Representative results are shown (patient 25). PBMCs were pulsed with TAA-derived peptides (peptides 14, 21, and 24) for 16 hours and then analyzed for IFN- $\gamma$  production. (D) IFN- $\gamma$ -producing T cells were also examined for naive/effector/memory phenotype by the criterion of CD45RA/CCR7 expression. The number of cells was calculated from the results of FACS analysis and is shown as a number per 300,000 PBMCs. White and black bars indicate the frequency of TAA-specific IFN- $\gamma$ -producing T cells before and after HCC treatments, respectively. The experiments were performed in five patients and similar results were observed.

in Fig. 5B. The magnitude of TAA-specific T-cell increase was statistically significant in four patients.

To examine the effect of CTLA-4 antibodies for production of other cytokines by T cells, we measured 27 kinds of human cytokines and chemokines in the medium of ELISpot assay. Figure 5C shows the results of cytokine production in the well with positive T-cell responses against TAA-derived peptides. The various cytokines consisting of IL-1 $\beta$ , IL-4, IL-6, IL-10, IL-17, eotaxin, G-CSF, GM-CSF, IFN- $\gamma$ , MIP-1 $\alpha$ , MIP-1 $\beta$ , RANTES, and TNF- $\alpha$  were increased in the medium with CTLA-4 antibodies compared with that without CTLA-4 antibodies. In contrast, increased

production of these cytokines in the well without positive T-cell responses against TAA-derived peptides was not observed in medium either with or without CTLA-4 antibodies (Fig. 5D).

## Discussion

In recent years, specific TAAs and their CTL epitopes have been identified in many tumors.<sup>21</sup> Several TAAs and their CTL epitopes, such as AFP, MAGE, and human telomerase reverse transcriptase (hTERT) have also been reported in HCC.<sup>19,20,24,41</sup> Although AFP-targeting immunotherapy could induce TAA-




Fig. 5. Enhancement of TAA-specific T-cell responses in HCC patients by CTLA-4 antibodies. (A) Summary of patients and peptides with an increase of the number of IFN- $\gamma$ -producing T cells. Black, gray, white, and hatched boxes indicate the immune responses with an increase of more than 10 specific spots, an increase of 1-10 specific spots, without change and a decrease of 1-10 specific spots, respectively. (B) Representative results of six patients are shown. Black and white bars indicate the results of assays incubated with CTLA-4 antibodies and mouse IgG2a isotype control, respectively. Data are expressed as the mean  $\pm$  SD of specific spots, except for patients 14 and 31. (C) Effects of CTLA-4 antibodies on production of cytokine and chemokine. Cytokine and chemokine levels in the medium of ELISPOT assay were measured using the Bio-plex assay. The graphs indicate the concentrations of cytokine and chemokine in the medium of ELISPOT assay using PBMCs of patient 31 and peptide 13 (medium in ELISPOT assay with enhancement of T-cell response) (see A,B). The increase of cytokines and chemokines after incubation with anti-CTLA-4 antibodies was confirmed in another three experiments using PBMCs of three other patients. (D) The graphs indicate the concentrations of cytokine and chemokine in the medium of ELISPOT assay using PBMCs of patient 31 and peptide 22 (medium in ELISPOT assay without enhancement of T-cell response) (see A).

specific CTLs, no patients achieved an objective tumor response; therefore, the search for TAAs as suitable targets for HCC immunotherapy and identification of their epitopes are important issues in therapy development. However, to date, T-cell responses to previously identified TAAs or their epitopes have been measured simultaneously and comparatively in only one study involving several patients with HBV-related HCC,<sup>42</sup> but no T-cell responses to the many other TAAs or their epitopes have been evaluated.

In this study we performed a simultaneous, comparative analysis of immune responses to 27 different CTL epitopes derived from 14 previously reported TAAs in the peripheral blood lymphocytes of 31 HCV-related HCC patients. We noted immune responses to epitopes (peptides 4, 12, 13, 16, 17, 22, 24, and 27) derived from CypB, SART2, SART3,

p53, MRP3, AFP, and hTERT in more than two patients (Fig. 1). These findings suggest the immunogenicity of these TAAs and their epitopes. In addition, the frequencies of peripheral blood CTLs specific to epitopes (peptides 4, 13, 16, 22, and 24) derived from CypB, SART3, p53, MRP3, and AFP, as detected by the ELISPOT assay, were high ( $\geq 20$  specific spots/300,000 PBMCs), suggesting the high immunogenicity of these TAAs and their epitopes.

Among these immunogenic antigens the expression of p53, MRP3, AFP, and hTERT was reported in HCC.<sup>18,19,43,44</sup> We also previously confirmed that the expression of SART2 and SART3 was observed in 100% of human HCC tissue (data not shown). As for CypB, this protein is well known to be widely expressed in normal and malignant tissue<sup>7</sup>; therefore, it is considered to be expressed in HCC.

Regarding tumor immunotherapy, it has recently been reported that strong immune responses can be induced at an earlier postvaccination time using, as peptide vaccines, epitopes that frequently occur in peripheral blood CTL precursors.<sup>23</sup> The epitopes (peptides 4, 12, 13, 16, 22, 24, and 27) that were derived from CypB, SART2, SART3, p53, MRP3, AFP, and hTERT and considered to be highly immunogenic in this study were capable of inducing epitope-specific CTLs from the PBMCs of HCC patients, suggesting that these epitopes can be candidates for peptide vaccines.

Next, TAA-specific immune responses were compared among three groups of subjects: HCC patients, normal blood donors, and patients with chronic hepatitis C not complicated by HCC. The results showed that there were no differences in the positive rate of immune responses to CMV among the three groups and no difference in the positive rate of immune responses to HCV between chronic hepatitis C patients with and without HCC. However, TAA-specific immune responses were observed frequently only in HCC patients, indicating that these immune responses are specific to HCC.

In the present study we also analyzed factors influencing host immune responses to these TAA-derived epitopes. Previous studies have reported that treatments, such as RFA and TAE, enhance HCC-specific T-cell responses.<sup>19,37,38</sup> However, TAAs and their epitopes, to which these enhanced immune responses occur, have not been identified. Thus, we simultaneously measured immune responses to 27 different epitopes derived from 14 TAAs in 12 patients who were available for analysis before and after treatment. The results showed that the antigens and their epitopes to which treatment-enhanced T-cell responses occur were diverse and some of them were newly induced after HCC treatment, suggesting that HCC treatments could induce *de novo* T-cell responses and these TAAs and their epitopes can be candidates as targets for HCC immunotherapy.

Furthermore, it became clear that enhanced immune responses to TAAs were induced not only by previously reported RFA and TAE, but also by cytotoxic drug chemotherapy. The patients who received chemotherapy showed partial responses after the treatment; therefore, we considered that it induced release of TAA into the tumor environment by tumor necrosis and/or apoptosis such as the mechanism reported in RFA or TAE.<sup>19,37,38</sup> Thus, our findings suggest that combined cancer chemotherapy and immunotherapy is useful as a treatment for HCC.

Analysis of the memory phenotypes of the T cells thus induced showed that the phenotypes of T cells whose frequency increased were mostly CD45RA<sup>-</sup>/CCR7<sup>+</sup> T cells (central memory T cells). Previous studies have reported that T cells with this phenotype differentiate into effector memory T cells and effector T cells, and that they require secondary stimulation by antigen to exert stronger antitumor effects.<sup>39</sup> Therefore, our findings suggest that the antitumor effect of tumor-specific T cells induced by HCC treatment is insufficient, and a booster with TAAs or epitope-containing peptides is a suitable method to further enhance antitumor effects.

Finally, we investigated the effect of anti-CTLA-4 antibodies, which have recently been in clinical trials as drugs enhancing antitumor immunity, on the host immune response to HCC. Regarding the mechanism of the antitumor activity of anti-CTLA-4 antibodies, it has been reported that they maximize the antitumor effect by blocking CTLA-4 on the surface of effector and regulatory T cells.<sup>40</sup> Because the number of peripheral blood regulatory T cells has been reported to increase in HCC patients,<sup>45</sup> TAA-specific CTLs that should be present but may not be detected by the ELISPOT assay. Therefore, in this study anti-CTLA-4 antibodies were added along with peptides to examine their effect on the ELISPOT assay.

The addition of anti-CTLA-4 antibodies resulted in an increase in the frequency of TAA-specific T cells in 60% of HCC patients. Although most patients showed an increase of only 1-10 TAA-specific T cells, the increased number of T cells was statistically significant. In addition, an increase of more than 10 TAA-specific T cells and a conversion from a negative to a positive response were observed in four patients. These results suggested that the anti-CTLA-4 antibodies unmasked IFN- $\gamma$  production by CTLs. However, the function might be limited because the number of TAA-specific T cells was not changed and even decreased in some patients.

The cytokine and chemokine profiling showed that the addition of anti-CTLA-4 antibodies increased the production of not only IFN- $\gamma$  but also cytokines, such as TNF- $\alpha$ , IL-1, and IL-6, and chemokines such as MIP-1; therefore, we speculate that the increased production of these antitumor immunity substances also plays a role in the unmasking of TAA-specific CTLs by anti-CTLA-4 antibodies. These results suggest that anti-CTLA-4 antibody is promising as a drug to enhance antitumor immunity, and that the ELISPOT assay with this antibody may serve as a more appropriate test tool to detect more HCC-specific TAAs or their epitopes.

On the other hand, recent studies have shown the important role of CD4<sup>+</sup> helper T cells in optimal function and proliferation of CD8<sup>+</sup> T cells.<sup>46</sup> Therefore, the lack of CD4<sup>+</sup> helper T cells or anergic CD4<sup>+</sup> T cells may explain the limited TAA-specific CD8<sup>+</sup> T-cell responses in HCC. Further studies using CD4<sup>+</sup> T-cell-depleted PBMCs or CD8<sup>+</sup> T cells expanded with TAA-derived peptide may enable identification of more immunogenic HCC-specific TAAs and their epitopes.

In conclusion, the results of this study suggest that CypB, SART2, SART3, p53, MRP3, AFP, and hTERT are promising TAAs in HCC immunotherapy, that the administration of these TAAs or peptides containing their epitopes as vaccines after HCC treatment is likely to be effective, and that the concomitant use of anti-CTLA-4 antibodies may further increase antitumor immunity. We believe that the results of this study provide useful information for the development of immunotherapy for HCC.

*Acknowledgment:* The authors thank Kazumi Fushimi, Maki Kawamura, Masayo Baba and Nami Nishiyama for technical assistance.

## References

- Deuffic S, Poynard T, Buffat L, Valleron AJ. Trends in primary liver cancer. *Lancet* 1998;351:214-215.
- Parkin DM, Bray F, Ferlay J, Pisani P. Estimating the world cancer burden: Globocan 2000. *Int J Cancer* 2001;94:153-156.
- Lencioni R. Loco-regional treatment of hepatocellular carcinoma. *HEPATOLOGY* 2010;52:762-773.
- Okuwaki Y, Nakazawa T, Shibuya A, Ono K, Hidaka H, Watanabe M, et al. Intrahepatic distant recurrence after radiofrequency ablation for a single small hepatocellular carcinoma: risk factors and patterns. *J Gastroenterol* 2008;43:71-78.
- Nishizaka S, Gomi S, Harada K, Oizumi K, Itoh K, Shichijo S. A new tumor-rejection antigen recognized by cytotoxic T lymphocytes infiltrating into a lung adenocarcinoma. *Cancer Res* 2000;60:4830-4837.
- Kawano K, Gomi S, Tanaka K, Tsuda N, Kamura T, Itoh K, et al. Identification of a new endoplasmic reticulum-resident protein recognized by HLA-A24-restricted tumor-infiltrating lymphocytes of lung cancer. *Cancer Res* 2000;60:3550-3558.
- Gomi S, Nakao M, Niiya F, Imamura Y, Kawano K, Nishizaka S, et al. A cyclophilin B gene encodes antigenic epitopes recognized by HLA-A24-restricted and tumor-specific CTLs. *J Immunol* 1999;163:4994-5004.
- Harashima N, Tanaka K, Sasatomi T, Shimizu K, Miyagi Y, Yamada A, et al. Recognition of the Lck tyrosine kinase as a tumor antigen by cytotoxic T lymphocytes of cancer patients with distant metastases. *Eur J Immunol* 2001;31:323-332.
- Fujie T, Tahara K, Tanaka F, Mori M, Takesako K, Akiyoshi T. A MAGE-1-encoded HLA-A24-binding synthetic peptide induces specific anti-tumor cytotoxic T lymphocytes. *Int J Cancer* 1999;80:169-172.
- Nishiyama T, Tachibana M, Horiguchi Y, Nakamura K, Ikeda Y, Takesako K, et al. Immunotherapy of bladder cancer using autologous dendritic cells pulsed with human lymphocyte antigen-A24-specific MAGE-3 peptide. *Clin Cancer Res* 2001;7:23-31.
- Kikuchi M, Nakao M, Inoue Y, Matsunaga K, Shichijo S, Yamana H, et al. Identification of a SART-1-derived peptide capable of inducing HLA-A24-restricted and tumor-specific cytotoxic T lymphocytes. *Int J Cancer* 1999;81:459-466.
- Nakao M, Shichijo S, Imaizumi T, Inoue Y, Matsunaga K, Yamada A, et al. Identification of a gene coding for a new squamous cell carcinoma antigen recognized by the CTL. *J Immunol* 2000;164:2565-2574.
- Yang D, Nakao M, Shichijo S, Sasatomi T, Takasu H, Matsumoto H, et al. Identification of a gene coding for a protein possessing shared tumor epitopes capable of inducing HLA-A24-restricted cytotoxic T lymphocytes in cancer patients. *Cancer Res* 1999;59:4056-4063.
- Tanaka H, Tsunoda T, Nukaya I, Sette A, Matsuda K, Umamo Y, et al. Mapping the HLA-A24-restricted T-cell epitope peptide from a tumour-associated antigen HER2/neu: possible immunotherapy for colorectal carcinomas. *Br J Cancer* 2001;84:94-99.
- Eura M, Chikamatsu K, Katsura F, Obata A, Sobao Y, Takiguchi M, et al. A wild-type sequence p53 peptide presented by HLA-A24 induces cytotoxic T lymphocytes that recognize squamous cell carcinomas of the head and neck. *Clin Cancer Res* 2000;6:979-986.
- Umamo Y, Tsunoda T, Tanaka H, Matsuda K, Yamaue H, Tanimura H. Generation of cytotoxic T-cell responses to an HLA-A24 restricted epitope peptide derived from wild-type p53. *Br J Cancer* 2001;84:1052-1057.
- Ferries E, Connan F, Pages F, Gaston J, Hagnere AM, Vieillefond A, et al. Identification of p53 peptides recognized by CD8(+) T lymphocytes from patients with bladder cancer. *Hum Immunol* 2001;62:791-798.
- Yamada A, Kawano K, Koga M, Matsumoto T, Itoh K. Multidrug resistance-associated protein 3 is a tumor rejection antigen recognized by HLA-A2402-restricted cytotoxic T lymphocytes. *Cancer Res* 2001;61:6459-6466.
- Mizukoshi E, Nakamoto Y, Tsuji H, Yamashita T, Kaneko S. Identification of alpha-fetoprotein-derived peptides recognized by cytotoxic T lymphocytes in HLA-A24+ patients with hepatocellular carcinoma. *Int J Cancer* 2006;118:1194-1204.
- Mizukoshi E, Nakamoto Y, Marukawa Y, Arai K, Yamashita T, Tsuji H, et al. Cytotoxic T-cell responses to human telomerase reverse transcriptase in patients with hepatocellular carcinoma. *HEPATOLOGY* 2006;43:1284-1294.
- Ribas A, Butterfield LH, Glaspy JA, Economou JS. Current developments in cancer vaccines and cellular immunotherapy. *J Clin Oncol* 2003;21:2415-2432.
- Rosenberg SA, Yang JC, Schwartzentruber DJ, Hwu P, Marincola FM, Topalian SL, et al. Immunologic and therapeutic evaluation of a synthetic peptide vaccine for the treatment of patients with metastatic melanoma. *Nat Med* 1998;4:321-327.
- Itoh K, Yamada A. Personalized peptide vaccines: a new therapeutic modality for cancer. *Cancer Sci* 2006;97:970-976.
- Butterfield LH, Ribas A, Meng WS, Dissette VB, Amarnani S, Vu HT, et al. T-cell responses to HLA-A\*0201 immunodominant peptides derived from alpha-fetoprotein in patients with hepatocellular cancer. *Clin Cancer Res* 2003;9:5902-5908.
- Butterfield LH, Ribas A, Dissette VB, Lee Y, Yang JQ, De la Rocha P, et al. A phase I/II trial testing immunization of hepatocellular carcinoma patients with dendritic cells pulsed with four alpha-fetoprotein peptides. *Clin Cancer Res* 2006;12:2817-2825.
- Butterfield LH. Recent advances in immunotherapy for hepatocellular cancer. *Swiss Med Wkly* 2007;137:83-90.
- Dougan M, Dranoff G. Immune therapy for cancer. *Annu Rev Immunol* 2009;27:83-117.
- O'Day SJ, Maio M, Chiarion-Sileni V, Gajewski TF, Pehamberger H, Bondarenko IN, et al. Efficacy and safety of ipilimumab monotherapy in patients with pretreated advanced melanoma: a multicenter single-arm phase II study. *Ann Oncol* 2010;21:1712-1717.
- Araki T, Itai Y, Furui S, Tasaka A. Dynamic CT densitometry of hepatic tumors. *AJR Am J Roentgenol* 1980;135:1037-1043.

30. Japan. LCSCGo. Classification of primary liver cancer. English ed 2. Tokyo: Kanehara; 1997.
31. Desmet VJ, Gerber M, Hoofnagle JH, Manns M, Scheuer PJ. Classification of chronic hepatitis: diagnosis, grading and staging. *HEPATOLOGY* 1994;19:1513-1520.
32. Ikeda-Moore Y, Tomiyama H, Miwa K, Oka S, Iwamoto A, Kaneko Y, et al. Identification and characterization of multiple HLA-A24-restricted HIV-1 CTL epitopes: strong epitopes are derived from V regions of HIV-1. *J Immunol* 1997;159:6242-6252.
33. Kurokohchi K, Arima K, Nishioka M. A novel cytotoxic T-cell epitope presented by HLA-A24 molecule in hepatitis C virus infection. *J Hepatol* 2001;34:930-935.
34. Kuzushima K, Hayashi N, Kimura H, Tsurumi T. Efficient identification of HLA-A\*2402-restricted cytomegalovirus-specific CD8(+) T-cell epitopes by a computer algorithm and an enzyme-linked immunospot assay. *Blood* 2001;98:1872-1881.
35. Oiso M, Eura M, Katsura F, Takiguchi M, Sobao Y, Masuyama K, et al. A newly identified MAGE-3-derived epitope recognized by HLA-A24-restricted cytotoxic T lymphocytes. *Int J Cancer* 1999;81:387-394.
36. Takahashi T, Tagami T, Yamazaki S, Uede T, Shimizu J, Sakaguchi N, et al. Immunologic self-tolerance maintained by CD25(+)CD4(+) regulatory T cells constitutively expressing cytotoxic T lymphocyte-associated antigen 4. *J Exp Med* 2000;192:303-310.
37. Zerbini A, Pilli M, Penna A, Pelosi G, Schianchi C, Molinari A, et al. Radiofrequency thermal ablation of hepatocellular carcinoma liver nodules can activate and enhance tumor-specific T-cell responses. *Cancer Res* 2006;66:1139-1146.
38. Ayaru L, Pereira SP, Alisa A, Pathan AA, Williams R, Davidson B, et al. Unmasking of alpha-fetoprotein-specific CD4(+) T-cell responses in hepatocellular carcinoma patients undergoing embolization. *J Immunol* 2007;178:1914-1922.
39. Sallusto F, Lenig D, Forster R, Lipp M, Lanzavecchia A. Two subsets of memory T lymphocytes with distinct homing potentials and effector functions. *Nature* 1999;401:708-712.
40. Peggs KS, Quezada SA, Chambers CA, Korman AJ, Allison JP. Blockade of CTLA-4 on both effector and regulatory T cell compartments contributes to the antitumor activity of anti-CTLA-4 antibodies. *J Exp Med* 2009;206:1717-1725.
41. Zerbini A, Pilli M, Soliani P, Ziegler S, Pelosi G, Orlandini A, et al. Ex vivo characterization of tumor-derived melanoma antigen encoding gene-specific CD8+ cells in patients with hepatocellular carcinoma. *J Hepatol* 2004;40:102-109.
42. Gehring AJ, Ho ZZ, Tan AT, Aung MO, Lee KH, Tan KC, et al. Profile of tumor antigen-specific CD8 T cells in patients with hepatitis B virus-related hepatocellular carcinoma. *Gastroenterology* 2009;137:682-690.
43. Fujioka M, Nakashima Y, Nakashima O, Kojiro M. Immunohistologic study on the expressions of alpha-fetoprotein and protein induced by vitamin K absence or antagonist II in surgically resected small hepatocellular carcinoma. *HEPATOLOGY* 2001;34:1128-1134.
44. Hussain SP, Schwank J, Staib F, Wang XW, Harris CC. TP53 mutations and hepatocellular carcinoma: insights into the etiology and pathogenesis of liver cancer. *Oncogene* 2007;26:2166-2176.
45. Fu J, Xu D, Liu Z, Shi M, Zhao B, Fu B, et al. Increased regulatory T cells correlate with CD8 T-cell impairment and poor survival in hepatocellular carcinoma patients. *Gastroenterology* 2007;132:2328-2339.
46. Kennedy R, Celis E. Multiple roles for CD4+ T cells in anti-tumor immune responses. *Immunol Rev* 2008;222:129-144.

## Malnutrition Impairs Interferon Signaling Through mTOR and FoxO Pathways in Patients With Chronic Hepatitis C

MASAO HONDA,\*<sup>‡</sup> KENJI TAKEHANA,<sup>§</sup> AKITO SAKAI,\* YUSUKE TAGATA,<sup>§</sup> TAKAYOSHI SHIRASAKI,<sup>‡</sup> SHINOBU NISHITANI,<sup>§</sup> TAKAHIKO MURAMATSU,<sup>||</sup> TATSUYA YAMASHITA,\* YASUNARI NAKAMOTO,\* EISHIRO MIZUKOSHI,\* YOSHIO SAKAI,\* TARO YAMASHITA,\* MIKIKO NAKAMURA,\* TETSURO SHIMAKAMI,<sup>||</sup> MINKYUNG YI,<sup>#</sup> STANLEY M. LEMON,<sup>||</sup> TETSUO SUZUKI,<sup>\*\*</sup> TAKAJI WAKITA,<sup>\*\*</sup> SHUICHI KANEKO,\* and the Hokuriku Liver Study Group

\*Department of Gastroenterology, <sup>‡</sup>Department of Advanced Medical Technology, Kanazawa University Graduate School of Medicine, Kanazawa, Japan; <sup>§</sup>Exploratory Research Laboratories, Research Center, Ajinomoto Pharmaceuticals, Co, Ltd, Kanagawa, Japan; <sup>||</sup>Frontier Research Labs, Institute for Innovation, Ajinomoto, Co, Inc, Kanagawa, Japan; <sup>||</sup>Division of Infectious Diseases, School of Medicine, The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina; <sup>#</sup>Center for Hepatitis Research, Institute for Human Infections and Immunity, and Department of Microbiology and Immunology, University of Texas Medical Branch, Galveston, Texas; and <sup>\*\*</sup>Department of Virology II, National Institute of Infectious Diseases, Tokyo, Japan

CLINICAL LIVER

**BACKGROUND & AIMS:** Patients with advanced chronic hepatitis C (CH-C) often are malnourished, but the effects of malnutrition on interferon (IFN) signaling and response to treatment have not been determined. We assessed the importance of the nutritional state of the liver on IFN signaling and treatment response. **METHODS:** We studied data from 168 patients with CH-C who were treated with the combination of pegylated-IFN and ribavirin. Plasma concentrations of amino acids were measured by mass spectrometry. Liver gene expression profiles were obtained from 91 patients. Huh-7 cells were used to evaluate the IFN signaling pathway, mammalian target of rapamycin complex 1 (mTORC1), and forkhead box O (FoxO). Antiviral signaling induced by branched-chain amino acids (BCAAs) was determined using the in vitro hepatitis C virus replication system. **RESULTS:** Multivariate logistic regression analysis showed that Fischer's ratio was associated significantly with nonresponders, independent of interleukin-28B polymorphisms or the histologic stage of the liver. Fischer's ratio was correlated inversely with the expression of BCAA transaminase 1, and was affected by hepatic mTORC1 signaling. IFN stimulation was impaired substantially in Huh-7 cells grown in medium that was low in amino acid concentration, through repressed mTORC1 signaling, and increased Socs3 expression, which was regulated by Foxo3a. BCAA could restore impaired IFN signaling and inhibit hepatitis C virus replication under conditions of malnutrition. **CONCLUSIONS: Malnutrition impaired IFN signaling by inhibiting mTORC1 and activating Socs3 signaling through Foxo3a. Increasing BCAAs to up-regulate IFN signaling might be used as a new therapeutic approach for patients with advanced CH-C.**

*Keywords:* HCV; Liver Disease; Therapy; Diet.

Interferon (IFN) and ribavirin (RBV) combination therapy is a popular modality for treating patients with chronic hepatitis C (CH-C), but approximately 50% of patients usually relapse, particularly those with hepatitis C virus (HCV) genotype 1b and a high viral load.<sup>1</sup>

Recent landmark studies of genome-wide associations identified genomic loci associated with treatment responses to pegylated (Peg)-IFN and RBV combination therapy,<sup>2,3</sup> and a polymorphism in the interleukin (IL)-28B gene was found to predict hepatitis C treatment-induced viral clearance. Moreover, we previously showed that expression of hepatic IFN-stimulated genes (ISGs) was associated with the IL-28B polymorphism and might contribute to the treatment response.<sup>4</sup> In addition to the IL-28B polymorphism, host factors such as fibrosis stage and metabolic status of the liver might be associated with the treatment outcome<sup>4,5</sup>; however, the significance of these factors in conjunction with the IL-28B polymorphism has not been evaluated fully.

In CH-C livers, prolonged liver cell damage, fibrosis development, and microcirculation failure can lead to a state of malnutrition in hepatocytes, resulting in the impairment of multiple metabolic pathways. In patients with advanced stage CH-C, hypoalbuminemia and decreased plasma values for the Fischer's ratio of branched-amino acids (BCAA; leucine, isoleucine, and valine) to aromatic amino acids (tyrosine and phenylalanine) commonly are observed. BCAA are the essential amino acids necessary for ammonium metabolism in muscle when the liver is unable to perform this function. Recent reports have shown that BCAA activates albumin synthesis in rat

*Abbreviations used in this paper:* BCAA, branched-chain amino acid; BCAT1, branched chain amino-acid transaminase 1; CH-C, chronic hepatitis C; ChIP, chromatin immunoprecipitation; DMEM, Dulbecco's modified Eagle medium; FBE, Foxo binding element; FBEmut, Foxo binding element mutant; FoxO, forkhead box, subgroup O; GLuc, Gaussia luciferase; IFN, interferon; IL, interleukin; ISG, interferon-stimulated genes; mTOR, mammalian target of rapamycin; mTORC1, mammalian target of rapamycin complex 1; NR, no response; PCR, polymerase chain reaction; Peg, pegylated; p-mTOR, phosphorylated form of mammalian target of rapamycin; pS6K, phosphorylated form of p70 S6 protein kinase; pSTAT1, phosphorylated form of signal transducer and activator of transcription 1; Raptor, regulatory associated protein of mTOR; RBV, ribavirin; S6K, p70 S6 protein kinase; siRNA, small interfering RNA; SVR, sustained viral response; TR, transient response.

© 2011 by the AGA Institute

0016-5085/\$36.00

doi:10.1053/j.gastro.2011.03.051