

- radio-frequency thermal ablation versus percutaneous ethanol injection. *Radiology* 2003; 228 (1): 235–40.
- 66 Lin SM, Lin CJ, Lin CC, Hsu CW, Chen YC. Radiofrequency ablation improves prognosis compared with ethanol injection for hepatocellular carcinoma < or =4 cm. *Gastroenterology* 2004; 127 (6): 1714–23.
- 67 Lin SM, Lin CJ, Lin CC, Hsu CW, Chen YC. Randomised controlled trial comparing percutaneous radiofrequency thermal ablation, percutaneous ethanol injection, and percutaneous acetic acid injection to treat hepatocellular carcinoma of 3 cm or less. *Gut* 2005; 54: 1151–6.
- 68 Chen MS, Li JQ, Zheng Y *et al.* A prospective randomized trial comparing percutaneous local ablative therapy and partial hepatectomy for small hepatocellular carcinoma. *Ann Surg* 2006; 243: 321–8.
- 69 Llovet JM, Di Bisceglie AM, Bruix J *et al.* Design and endpoints of clinical trials in hepatocellular carcinoma. *J Natl Cancer Inst* 2008; 100: 698–711.
- 70 The Japanese Liver Transplantation Society. Liver transplantation in Japan in 2006 (part 2)-Registry by the Japanese Liver Transplantation Society. *Ishoku* 2008; 43: 45–55, (in Japanese).
- 71 Mazzaferro V, Regalia E, Doci R *et al.* Liver transplantation for the treatment of small hepatocellular carcinomas in patients with cirrhosis. *N Engl J Med* 1996; 334: 693–9.
- 72 Iwatsuki S, Starzl TE, Sheahan DG *et al.* Hepatic resection versus transplantation for hepatocellular carcinoma. *Ann Surg* 1991; 214: 221–8, discussion 228–229.
- 73 Penn I. Hepatic transplantation for primary and metastatic cancers of the liver. *Surgery* 1991; 110: 726–34, discussion 734–725.
- 74 Todo S, Furukawa H, Tada M. Extending indication: role of living donor liver transplantation for hepatocellular carcinoma. *Liver Transpl* 2007; 13 (11 Suppl 2): S48–54.
- 75 Mazzaferro V, Llovet JM, Miceli R *et al.* Predicting survival after liver transplantation in patients with hepatocellular carcinoma beyond the Milan criteria: a retrospective, exploratory analysis. *Lancet Oncol* 2009; 10 (1): 35–43.
- 76 Yao FY, Ferrell L, Bass NM *et al.* Liver transplantation for hepatocellular carcinoma: expansion of the tumor size limits does not adversely impact survival. *Hepatology* 2001; 33 (6): 1394–403.
- 77 Sugawara Y, Tamura S, Makuuchi M. Living donor liver transplantation for hepatocellular carcinoma: Tokyo University series. *Dig Dis* 2007; 25: 310–12.
- 78 Lee SG, Hwang S, Moon DB *et al.* Expanded indication criteria of living donor liver transplantation for hepatocellular carcinoma at one large-volume center. *Liver Transpl* 2008; 14: 935–45.
- 79 Ito T, Takada Y, Ueda M *et al.* Expansion of selection criteria for patients with hepatocellular carcinoma in living donor liver transplantation. *Liver Transpl* 2007; 13: 1637–44.
- 80 Duffy JP, Vardanian A, Benjamin E *et al.* Liver transplantation criteria for hepatocellular carcinoma should be expanded: a 22-year experience with 467 patients at UCLA. *Ann Surg* 2007; 246: 502–9, discussion 509–511.
- 81 Yamamoto J, Iwatsuki S, Kosuge T *et al.* Should hepatomas be treated with hepatic resection or transplantation? *Cancer* 1999; 86: 1151–8.
- 82 Figueras J, Jaurrieta E, Valls C *et al.* Resection or transplantation for hepatocellular carcinoma in cirrhotic patients: outcomes based on indicated treatment strategy. *J Am Coll Surg* 2000; 190: 580–7.
- 83 Llovet JM, Fuster J, Bruix J. Intention-to-treat analysis of surgical treatment for early hepatocellular carcinoma: resection versus transplantation. *Hepatology* 1999; 30 (6): 1434–40.
- 84 Llovet JM, Real MI, Montana X *et al.* Arterial embolisation or chemoembolisation versus symptomatic treatment in patients with unresectable hepatocellular carcinoma: a randomised controlled trial. *Lancet* 2002; 359 (9319): 1734–9.
- 85 Lo CM, Ngan H, Tso WK *et al.* Randomized controlled trial of transarterial lipiodol chemoembolization for unresectable hepatocellular carcinoma. *Hepatology* 2002; 35 (5): 1164–71.
- 86 Llovet JM, Bruix J. Systematic review of randomized trials for unresectable hepatocellular carcinoma: chemoembolization improves survival. *Hepatology* 2003; 37 (2): 429–42.
- 87 Bruix J, Sala M, Llovet JM. Chemoembolization for hepatocellular carcinoma. *Gastroenterology* 2004; 127 (5 Suppl 1): S179–88.
- 88 Makuuchi M, Kokudo N. Clinical practice guidelines for hepatocellular carcinoma: the first evidence based guidelines from Japan. *World J Gastroenterol* 2006; 12: 828–9.
- 89 Kudo M, Okanoue T. Management of hepatocellular carcinoma in Japan: consensus-based clinical practice manual proposed by the Japan Society of Hepatology. *Oncology* 2007; 72: S2–15.
- 90 Ikai I, Arii S, Okazaki M *et al.* Report of the 17th Nationwide Follow-up Survey of Primary Liver Cancer in Japan. *Hepatol Res* 2007; 37: 676–91.
- 91 Takayasu K, Arii S, Ikai I *et al.* Prospective cohort study of transarterial chemoembolization for unresectable hepatocellular carcinoma in 8510 patients. *Gastroenterology* 2006; 131 (2): 461–9.
- 92 Ueno S, Tanabe G, Nuruki K *et al.* Prognostic performance of the new classification of primary liver cancer of Japan (4th edition) for patients with hepatocellular carcinoma: a validation analysis. *Hepatol Res* 2002; 24: 395–403.
- 93 Marelli L, Stigliano R, Triantos C *et al.* Treatment outcomes for hepatocellular carcinoma using chemoembolization in combination with other therapies. *Cancer Treat Rev* 2006; 32: 594–606.
- 94 Veltri A, Moretto P, Doriguzzi A, Pagano E, Carrara G, Gandini G. Radiofrequency thermal ablation (RFA) after

- transarterial chemoembolization (TACE) as a combined therapy for unresectable non-early hepatocellular carcinoma (HCC). *Eur Radiol* 2006; 16 (3): 661–9.
- 95 Buscarini L, Buscarini E, Di Stasi M, Quaretti P, Zangrandi A. Percutaneous radiofrequency thermal ablation combined with transcatheter arterial embolization in the treatment of large hepatocellular carcinoma. *Ultraschall Med* 1999; 20 (2): 47–53.
- 96 Yamakado K, Nakatsuka A, Akeboshi M, Shiraki K, Nakano T, Takeda K. Combination therapy with radiofrequency ablation and transcatheter chemoembolization for the treatment of hepatocellular carcinoma: short-term recurrences and survival. *Oncol Rep* 2004; 11: 105–9.
- 97 Koda M, Ueki M, Maeda Y *et al.* The influence on liver parenchymal function and complications of radiofrequency ablation or the combination with transcatheter arterial embolization for hepatocellular carcinoma. *Hepatol Res* 2004; 29 (1): 18–23.
- 98 Obi S, Yoshida H, Toune R *et al.* Combination therapy of intraarterial 5-fluorouracil and systemic interferon-alpha for advanced hepatocellular carcinoma with portal venous invasion. *Cancer* 2006; 106 (9): 1990–7.
- 99 Ando E, Tanaka M, Yamashita F *et al.* Hepatic arterial infusion chemotherapy for advanced hepatocellular carcinoma with portal vein tumor thrombosis: analysis of 48 cases. *Cancer* 2002; 95: 588–95.
- 100 Lin CP, Yu HC, Cheng JS *et al.* Clinical effects of intraarterial infusion chemotherapy with cisplatin, mitomycin C, leucovorin and 5-fluorouracil for unresectable advanced hepatocellular carcinoma. *J Chin Med Assoc* 2004; 67: 602–10.
- 101 Sumie S, Yamashita F, Ando E *et al.* Interventional radiology for advanced hepatocellular carcinoma: comparison of hepatic artery infusion chemotherapy and transcatheter arterial lipiodol chemoembolization. *AJR Am J Roentgenol* 2003; 181 (5): 1327–34.
- 102 Tzoracoleftherakis EE, Spiliotis JD, Kyriakopoulou T, Kakkos SK. Intra-arterial versus systemic chemotherapy for non-operable hepatocellular carcinoma. *Hepatogastroenterology* 1999; 46 (26): 1122–5.
- 103 Court WS, Order SE, Siegel JA *et al.* Remission and survival following monthly intraarterial cisplatin in non-resectable hepatoma. *Cancer Invest* 2002; 20 (5–6): 613–25.
- 104 Okuda K, Tanaka M, Shibata J *et al.* Hepatic arterial infusion chemotherapy with continuous low dose administration of cisplatin and 5-fluorouracil for multiple recurrence of hepatocellular carcinoma after surgical treatment. *Oncol Rep* 1999; 6: 587–91.
- 105 Tanioka H, Tsuji A, Morita S *et al.* Combination chemotherapy with continuous 5-fluorouracil and low-dose cisplatin infusion for advanced hepatocellular carcinoma. *Anticancer Res* 2003; 23 (2C): 1891–7.
- 106 Park JY, Ahn SH, Yoon YJ *et al.* Repetitive short-course hepatic arterial infusion chemotherapy with high-dose 5-fluorouracil and cisplatin in patients with advanced hepatocellular carcinoma. *Cancer* 2007; 110: 129–37.
- 107 Chung YH, Song IH, Song BC *et al.* Combined therapy consisting of intraarterial cisplatin infusion and systemic interferon-alpha for hepatocellular carcinoma patients with major portal vein thrombosis or distant metastasis. *Cancer* 2000; 88 (9): 1986–91.
- 108 Kaneko S, Urabe T, Kobayashi K. Combination chemotherapy for advanced hepatocellular carcinoma complicated by major portal vein thrombosis. *Oncology* 2002; 62 (Suppl 1): 69–73.
- 109 Llovet JM, Ricci S, Mazzaferro V *et al.* Sorafenib in advanced hepatocellular carcinoma. *N Engl J Med* 2008; 359: 378–90.
- 110 Cheng AL, Kang YK, Chen Z *et al.* Efficacy and safety of sorafenib in patients in the Asia-Pacific region with advanced hepatocellular carcinoma: a phase III randomised, double-blind, placebo-controlled trial. *Lancet Oncol* 2009; 10 (1): 25–34.
- 111 Kudo M. Hepatocellular carcinoma 2009 and beyond: from the surveillance to molecular targeted therapy. *Oncology* 2008; 75: S1–12.
- 112 Kudo M. The 2008 Okuda lecture: management of hepatocellular carcinoma: from surveillance to molecular targeted therapy. *J Gastroenterol Hepatol* 2010; 25 (3): 439–52.

HCC develops even in the early stage of chronic liver disease in elderly patients with HCV infection

AKIO TAKATA, RYOKO KUROMATSU, EIJI ANDO, HIDEKI IWAMOTO, NOBUYOSHI FUKUSHIMA, SHUJI SUMIE, TAKUJI TORIMURA and MICHIO SATA

Division of Gastroenterology, Department of Medicine, Kurume University School of Medicine, Fukuoka 830-0011, Japan

Received February 11, 2010; Accepted April 9, 2010

DOI: 10.3892/ijmm_00000459

Abstract. In recent years, the number of elderly patients with hepatocellular carcinoma (HCC) has been increasing. The aim of this study was to compare the liver function and the background factors of HCC patients with hepatitis C virus (HCV) infection by generation and to examine the characteristics of this disease in the elderly. A total of 1096 patients (776 men and 320 women) diagnosed with HCV-related HCC at our institution from 1995 to 2006 were divided into 4 groups as follows: D group, 75 years of age or older; C group, 65-74 years of age; B group, 55-64 years of age; A group, 54 years of age or younger, and the liver function and other clinical characteristics were compared among these 4 groups. The average age at initial diagnosis of HCV-related HCC was 66.9 years of age. The A, B, C and D groups were comprised of 87, 363, 514 and 132 patients, respectively. The rate of Child-Pugh class A patients in the D group was significantly higher than that of the other groups ($P < 0.05$). The average levels of ALT, TB and PT-INR in the D group were significantly lower than the levels in the other groups ($P < 0.05$). The average Alb level in the D group was significantly higher than that in the other groups ($P < 0.05$). In conclusion, we found that HCV-related HCC in the elderly occurred against a background of chronic liver disease with mild inflammation and fibrosis.

Introduction

In recent years, the average age of the population of Japan has been increasing annually, and in 2006, elderly individuals over 65 years of age accounted for more than 20.8% of the entire population. The number of elderly, 65 years of age or older, who are afflicted with hepatocellular carcinoma (HCC) has also been increasing rapidly since 1990 (1,2). The average age at initial diagnosis of HCC at our institution from 1986 to

1993 was 60.9 years and 68.4 years in the period 2000 to 2006 and has been increasing annually (Fig. 1).

In Japan, more than 80% of HCC patients have hepatitis C virus (HCV) infection, and there are approximately 2 million patients with HCV-related chronic liver disease. It is estimated that there are 700,000 patients with undiagnosed HCV infection (3-5). The contributing factors include the spread of stimulant drugs, vaccinations and medical counter measures (injections, surgeries, blood transfusion) during the period after World War II when preventative measures against infection were inadequate. As a result, the incidence of HCC in patients with HCV who were infected during this period has been increasing in recent years (1,2,4,5).

Generally, 60-80% of patients develop chronic hepatitis after HCV infection, while many patients present with HCC approximately 30 years after HCV infection (6-9). On the other hand, it has been reported that the majority of patients with posttransfusion chronic HCV infection develop HCC after the age of 60 years regardless of when they acquired the HCV infection (10). Moreover, with the increase in fibrosis of transplanted livers in HCV-infected patients, it has been reported that the age of the donor is closely correlated with the rate of progression of fibrosis of the transplanted liver rather than the age of the recipient (11). Namely, the age of the patients who were infected with HCV is a more significant factor than the duration of the HCV infection in regards to the progression of HCV-persistent infection and fibrosis or carcinogenesis (10). Furthermore, in recent years the number of patients in which HCC does not occur before 60 years of age but thereafter develops in patients in their 70s or 80s has been increasing (12-16).

Therefore, it is assumed that age is the important factor contributing to hepatocarcinogenesis of HCV-related HCC. Therefore, we divided HCV-related HCC patients into 4 groups according to age. The aim of this study was to compare the liver function and the background features of the HCV-related HCC patients by generation, particularly focusing on the elderly group and examining the characteristics of this disease in the elderly.

Materials and methods

Patient population and experimental design. Among 1,404 patients with primary HCC consecutively diagnosed at our institution from January 1995 to December 2006, 1096 patients (776 men and 320 women) who were both HCV antibody-

Correspondence to: Dr Akio Takata, Division of Gastroenterology, Department of Medicine, Kurume University School of Medicine, 67 Asahi-machi, Kurume, Fukuoka 830-0011, Japan
E-mail: oika@med.kurume-u.ac.jp

Key words: elderly patients, HCV-related HCC, mild inflammation and fibrosis, age




Figure 1. Changes in the average age at initial diagnosis of HCC. The average age of onset was 60.9 years in 1986-1992, 64.9 years in 1993-1999, and 68.4 years in 2000-2006, with an annual increase in the average age.

positive and hepatitis B surface (HBs) antigen-negative were enrolled. The average age at initial diagnosis of HCC was 66 ± 9 years (range, 42-87). The clinical characteristics of the patients are documented in Table I. These patients were divided into 4 groups as follows: D group, patients ≥ 75 years of age; C group, patients ≥ 65 and < 75 years of age; B group, patients ≥ 55 and < 65 years of age, and A group, patients ≤ 54 years of age. Liver function and background factors were compared among the 4 groups. A liver biopsy was also performed in noncancerous areas in 18 patients in the D group targeted consecutively from 2005 to 2006.

HCC diagnosis. The diagnosis of HCC was based on hypervascularity, confirmed by dynamic computed tomography (CT), magnetic resonance imaging (MRI), angiography or CT angiography, when the serum levels of HCC-related tumor markers, such as α -fetoprotein (AFP) or des- γ carboxy prothrombin (DCP), were increased or a mass lesion was observed by ultrasonography. When a nodule was not proven to be hypervascular, percutaneous biopsy under ultrasonography was performed for confirmation of the diagnosis of HCC, and thereafter a pathological study using hematoxylin and eosin staining was carried out. The fibrosis staging scores and activity grades were assigned according to the criteria of Desmet and colleagues (17) and the French METAVIR Cooperative Study (18-20). Staging was defined as F0 (no fibrosis), F1 (mild fibrosis), F2 (moderate fibrosis), F3 (severe fibrosis), or F4 (cirrhosis), and grading was defined as A0 (no activity), A1 (mild activity), A2 (moderate activity), or A3 (severe activity). Three pathologists independently evaluated the disease stage and grade. Portal vein tumor thrombosis (PVTT) was defined as a protrusion of the tumor into the first and/or second branch, or into the main trunk of the portal vein. Tumor stage was classified according to the International Union Against Cancer (UICC) TNM classification.

Factors evaluated in the analysis. The following 19 factors were compared: gender, a history of blood transfusion, a history of habitual drinking, underlying diseases except for liver diseases, gastro-esophageal varices, Child-Pugh classification, serum albumin (Alb), serum total bilirubin (TB), prothrombin time-international normalized ratio (PT-INR),

Table I. Profile of the 1096 patients at initial diagnosis of HCV-related HCC.

Age (range)	66.9 \pm 7.6 (42-87)
Gender (male/female)	776/320
Alcohol consumption (none/light/heavy)	392/371/333
Underlying disease (none/single/multiple)	471/436/189
Gastro-esophageal varices (none/small/large) ^a	436/393/240
Blood transfusion (yes/no)	377/719
Child-Pugh classification (A/B/C)	751/290/55
Alb (g/dl)	3.48 \pm 0.47
PT-INR	1.17 \pm 0.15
TB (mg/dl)	1.17 \pm 0.66
ALT (U/l)	64.4 \pm 39.0
PLT ($\times 10^4/\mu$ l)	10.8 \pm 5.54
HA (ng/ml)	363 \pm 385
ICG R15 (%)	32.1 \pm 17.4

Date are expressed as the median range \pm standard deviation (SD). ^aTwenty seven patients did not undergo endoscopic examination. Alb, serum albumin; PT-INR, prothrombin time-international normalized ratio; TB, serum total bilirubin; ALT, serum alanine aminotransferase; PLT, platelet count; HA, hyaluronic acid; ICG R15, indocyanine green retention rate at 15 min.

serum alanine aminotransferase (ALT), platelet count (PLT), indocyanine green retention rate at 15 min (ICG R15), serum hyaluronic acid (HA), number of tumors, tumor size, portal vein tumor thrombosis (PVTT), tumor stage, α -fetoprotein (AFP) and extra-hepatic metastasis. The values at initial diagnosis of HCC were used for Alb, TB, PT-INR, ALT, PLT, ICG R15 and HA. Chronic hepatitis was included as Child-Pugh class A. Gastro-esophageal varices were classified into three types according to the endoscopic form factor grade: none, no varix; small, form 1 or 2; large, form 3, based on the classification of the Japanese Research Society for Portal Hypertension (21). The underlying diseases included hypertension, diabetes mellitus, cerebrovascular damage, cardiac disorder, renal disorder and cancer of other organs, and these were divided into three groups: none, no underlying disease; single, underlying disease by one disorder; multiple,




Figure 2. The age at initial diagnosis of HCC and background factors. The number of female patients, the patients without a history of habitual drinking and the patients with underlying diseases increased with age. (a) Concerning gender, the number of female patients in the D group was significantly higher than that in groups A and B (** $P < 0.01$). (b) The number of patients without a history of habitual drinking in the D group was significantly higher than that in the A and B groups (** $P < 0.01$). (c) The number of patients with underlying diseases in the D group was significantly higher than that in the A and B groups (** $P < 0.01$). (d) A history of blood transfusion was noted most frequently in the D group and was significantly higher than that in the A group (* $P < 0.05$). * $P < 0.05$, ** $P < 0.01$ between the indicated groups.

underlying diseases by more than two disorders. A history of habitual drinking was classified into three groups: none, non-drinker; light, < 84 g/day of ethanol; heavy, > 84 g/day of ethanol for > 5 years. The number of tumors was divided into two groups: solitary and multiple tumors. Tumor size was divided into two groups: those measuring ≤ 3 cm and > 3 cm in size. The AFP level was divided into three categories: ≤ 20 , 21-200 and > 200 ng/ml.

Statistical analysis. Data are expressed as the mean \pm standard deviation (SD). Data regarding Alb, TB, PT-INR, ALT, PLT, ICG R15 and HA levels were analyzed by one-way analysis of variance followed by the Dunnett's multiple comparison of means test. A Logistics regression analysis was used to analyze the effects of the background factors (gender, a history of blood transfusion, a history of habitual drinking, underlying diseases, degree of gastro-esophageal varices, Child-Pugh classification) and the tumor features (number of tumors, tumor size, PVTT, tumor stage, AFP, extrahepatic metastasis) on age at the initial diagnosis of HCC. The background factors and the tumor features were individually chosen as outcome variables, and the age at initial diagnosis of HCC divided into four groups were dummy coded and used as predictor variables. The statistical analysis was performed using SPSS for Windows (version 12.0). $P < 0.05$ was considered to be statistically significant.

Results

Comparison of background factors. Among the 1096 patients, 132 (12%) were classified in the D group (patients ≥ 75 years of age), with the oldest patient being 87 years of




Figure 3. The age at initial diagnosis of HCC and Child-Pugh classification. The percentages of Child-Pugh class A patients were 59.8, 64.5, 70.1 and 79.5% in the A, B, C and D groups, respectively. The rate of Child-Pugh class A patients increased with age, and the rate of Child-Pugh class A patients in the D group was significantly higher than the rate in the other groups (D vs. A and B, ** $P < 0.01$; D vs. C, * $P < 0.05$). * $P < 0.05$, ** $P < 0.01$ between the indicated groups.

age. The number of female patients increased with age, and the male:female patient ratio in the D group was 82:50. As a result, the number for female patients in the D group was significantly higher than that in the A and B groups (patients < 65 years of age) ($P < 0.01$) (Fig. 2a). The number of patients without a history of habitual drinking increased with age, and that of the D group was significantly higher than that in the A and B groups ($P < 0.01$) (Fig. 2b). The number of patients with underlying diseases increased with age, and that of the D group was significantly higher than that in the A and B groups ($P < 0.01$) (Fig. 2c). A history of blood transfusion was




Figure 4. A comparison of age at initial diagnosis of HCC and liver functions. The average values of (a) ALT, (b) TB and (c) PT-INR in the D group were significantly lower than the values in the other groups (ALT: D vs. A, B and C, ** $P < 0.01$; TB and PT-INR: D vs. A and B, ** $P < 0.01$, D vs. C, * $P < 0.05$). The average value of (d) Alb in the D group was significantly higher than that in the other groups (D vs. A and B, ** $P < 0.01$; D vs. C, * $P < 0.05$). The average value of (e) ICG R15 in the D group was significantly lower than that in the A and B groups (D vs. A and B, ** $P < 0.01$). The average value of (f) HA did not show a significant difference. * $P < 0.05$, ** $P < 0.01$ between the indicated groups.




Figure 5. Comparison of age at initial diagnosis of HCC and PLT, and PLT in Child-Pugh class A patients. (a) The average value of PLT in the D group was significantly higher than the value in the A (** $P < 0.01$) and B (* $P < 0.05$) groups. (b) The rate of 100,000 or more of PLT in Child-Pugh class A patients of the D group was significantly higher than that in the A (** $P < 0.01$) and B (* $P < 0.05$) groups. * $P < 0.05$, ** $P < 0.01$ between the indicated groups.




Figure 6. The age at initial diagnosis of HCC and the endoscopic form factor grade of varices. The number of patients with large varices was the lowest in the D group and the highest in the A group. The number of patients with large varices in the D group was significantly lower than that in the A (** $P < 0.01$) and B (* $P < 0.05$) groups. * $P < 0.05$, ** $P < 0.01$ between the indicated groups.

noted most frequently in the D group, and it was significantly higher than that in the A group ($P < 0.05$) (Fig. 2d).

Comparison of liver function. The percentages of Child-Pugh class A patients were 59.8, 64.5, 70.1 and 79.5%, in the A, B, C and D groups, respectively. Therefore, the percentage of Child-Pugh class A patients increased with age, and the percentage of Child-Pugh class A patients in the D group was higher than the other groups (D vs. A, B, $P < 0.01$; D vs. C, $P < 0.05$) (Fig. 3). The average values of ALT, TB and PT-INR were lower with age, and these average values in the D group were significantly lower than those in the other groups (ALT: D vs. A, B and C, $P < 0.01$; TB, PT-INR: D vs. A and B, $P < 0.01$; D vs. C, $P < 0.05$) (Fig. 4a-c). The average value of Alb in the D group was significantly higher than that in the other groups (D vs. A and B, $P < 0.01$; D vs. C, $P < 0.05$)

Table II. Characteristics of the HCC tumors of the 1096 patients at initial diagnosis of HCV-related HCC.

	A group (87) 42-54 years of age	B group (363) 55-64 years	C group (514) 65-74 years	D group (132) 75-87 years	P-value
Number of tumors					<0.05, D vs. C
Solitary	37	156	217	59	
Multiple	50	207	297	73	
Tumor size					<0.05, D vs. A and B
≤3 cm	61	238	312	73	
>3 cm	26	125	202	59	
PVTT					NS
Present	7	45	66	11	
Absent	80	318	448	121	
TNM staging					NS
I	14	88	106	20	
II	37	120	171	51	
III	30	107	174	49	
IV	6	48	63	12	
AFP (ng/ml)					NS
≤20	28	120	185	53	
21-200	41	158	183	42	
>200	18	85	146	37	
Extrahepatic metastasis					NS
Yes	2	20	18	3	
No	85	343	496	129	

PVTT, portal vein tumor thrombosis; AFP, α -fetoprotein.

(Fig. 4d). The average value of ICG R15 was the lowest in the D group, and the average value of ICG R15 in the D group was significantly lower than that in the A and B groups (D vs. A and B, $P<0.01$) (Fig. 4e). The average value of HA was the lowest in the C group, thus indicating a strong variation and no significant difference (Fig. 4f). The average value of PLT was the highest in the D group, and the average value of PLT in the D group was significantly higher than that in the A and B groups (D vs. A, $P<0.01$; D vs. B, $P<0.05$) (Fig. 5a). When the Child-Pugh class A patients were divided per 100,000/ μ l of PLT and compared, the rate of 100,000 or more in the D group increased with age, and it was higher than the A and B groups (D vs. A, $P<0.01$; D vs. B, $P<0.05$) (Fig. 5b). The number of patients with large varices decreased with age, and the number of patients with large varices in the D group was lower than that in the A and B groups (D vs. A, $P<0.01$; D vs. B, $P<0.05$) (Fig. 6).

Comparison of characteristics of the HCC tumors. The number of patients with small HCCs ≤ 3 cm in the D group was lower than that in the A and B groups (D vs. A and B, $P<0.05$), and the number of patients with solitary HCC tumor in the D group was higher than that in the C group (D vs. C, $P<0.05$) (Table II). There were no statistical differences in the prevalence of PVTT, tumor stage, AFP or extrahepatic metastasis.

Pathological findings of liver biopsies from noncancerous areas. The pathological findings of patients for whom liver biopsies from noncancerous areas were performed are documented in Table III. Four of the 18 patients showed liver cirrhosis with activity ≥ 2 and fibrosis 4 in the Desmet classification. Fourteen of the 18 patients showed chronic hepatitis that was milder than activity 2 and fibrosis 3 in the Desmet classification.

Discussion

An elder is generally defined as anyone 65 years of age or older, but this definition is not standardized worldwide. In Japan, the average age at initial diagnosis of HCC is 65 years or older (13,14,22). Therefore, it is not appropriate to draw the dividing line at 65 years of age in order to define the characteristics of HCC that presents at old age (23). On the other hand, it has been reported that HCV-related HCC which develops in individuals in their 40s exhibits different characteristics from HCV-related HCC which develops in individuals 50 years of age or older, since HCC in individuals in their 40s is independently related to heavy drinking and the presence of HBV coinfection (24). It is therefore not appropriate that younger patients in their 40s and older patients who are 65 years of age or older are treated as a similar group, even when HCC patients are divided by a line drawn at 70 years

Table III. Pathological findings of liver biopsies from non-cancerous areas and the liver function of 18 patients with HCC 75 years of age or older.

No.	Age	Gender	Histology Desmet classification	PLT ($\times 10^4/\mu\text{l}$)	ALT (U/l)	TB (mg/dl)	PT-INR	Alb (g/dl)	ICG R15 (%)	HA (ng/ml)
1	78	F	A1F1	9.4	87	1.10	1.09	3.6	30.6	287
2	80	M	A2F4	7.9	79	0.70	1.03	3.3	14.3	237
3	82	F	A2F2	37.5	37	0.50	0.97	4.1	9.7	123
4	85	F	A2F2	8.3	28	0.50	1.06	4.1	16.8	470
5	75	M	A2F2	13.8	44	0.90	1.03	3.5	28.9	265
6	77	M	A2F2	17.5	64	1.30	1.13	4.0	30.3	225
7	75	M	A2F1	12.5	22	0.90	0.97	4.1	14.2	62
8	80	F	A2F2	15.3	54	1.00	1.12	3.4	45.7	1340
9	76	M	A2F4	8.3	62	0.80	1.18	3.5	24.5	557
10	87	M	A2F2	8.1	43	0.90	1.05	3.8	22.1	503
11	78	M	A1F1	16.3	33	0.95	1.05	4.5	16.8	126
12	76	M	A2F4	13.9	77	1.20	1.16	3.8	16.2	405
13	75	M	A3F4	8.9	33	1.40	1.16	3.7	48.2	619
14	76	F	A1F2	8.0	52	0.90	1.08	3.7	18.0	260
15	82	F	A2F3	11.0	37	2.20	1.20	3.1	23.8	1360
16	76	F	A2F2	11.9	45	0.70	1.02	4.2	9.6	353
17	76	M	A1F2	13.3	40	0.40	1.04	3.4	25.9	732
18	77	M	A2F3	10.1	54	1.02	1.01	4.1	45.2	163
Average	78.3			12.9	49.5	0.97	1.08	3.77	24.5	449

PLT, platelet count; ALT, serum alanine aminotransferase; TB, serum total bilirubin; PT-INR, prothrombin time-international normalized ratio; Alb, serum albumin; ICG R15, indocyanine green retention rate at 15 min; HA, serum hyaluronic acid.

of age as reported in previous studies (14,15,25,26). We therefore divided the elderly into the C group (>65 years of age and <75 years) and the D group (≥ 75 years of age), and the non-elderly into the A group (<54 years of age) and the B group (>55 and <65 years of age) to create 4 groups. The clinical characteristics of HCC in each group were thus examined in further detail by comparing each group, while particularly focusing on the D group.

Consequently, the most significant characteristic in patients with HCV-related HCC in the D group was that hepatic reserve was maintained and HCC occurred against a background of liver disorder with mild inflammation and fibrosis. In our study, levels of ALT and TB, which are well-known markers of inflammatory necrosis in the liver, were significantly lower in the D group than in all of the other groups. Notably, some studies found that alcohol promotes the progression of background chronic liver disease and consequently enhances carcinogenesis of the liver or that alcohol directly promotes carcinogenesis of the liver (27-29). In our study, the A group of younger HCC patients included many heavy drinkers of alcohol with severe inflammation in addition to high values of ALT and TB, as was previously found by Shimauchi *et al* (24). This finding corroborates reports that HCC frequently occurs in association with severe inflammation (30). Conversely, it was assumed to be one of the main reasons that the D group had many patients with low values of ALT and TB, since the D group included many patients who did not habitually consume alcohol.

It has been reported that in chronic liver disease, changes in the platelet count correlate with the degree of fibrosis in liver histology (31-33). As a result, the platelet count is regarded as a marker for fibrosis. In our study, the platelet count in the D group was significantly higher than that in the A and B groups (younger than 65 years of age), and the D group in particular showed the highest platelet count of 100,000 or more in the Child-Pugh class A patients. The study of 18 patients in the D group was assessed by means of a liver biopsy, and the background liver showed chronic hepatitis of less than F3 except for 4 cases of F4, thus indicating that many patients with mild fibrosis had not advanced to liver cirrhosis.

Gastro-esophageal varices are the most common clinical manifestation of portal hypertension in patients with liver cirrhosis. Nakayama *et al* (34) reported that as the endoscopic form factor grade of varix increased, the incidence of the occurrence of HCC also increased, and in particular, a form 3 (large size) factor of varices was an independent predictor for HCC. In our study, many of the patients in the D group either did not have varices or only had small varices, and also the number of patients with large varices in the D group was significantly smaller than the non-elderly groups (younger than 65 years of age). Therefore, it was assumed that many of the patients in the D group had not advanced to liver cirrhosis, and many cases of HCC had thus occurred against a background of chronic liver disease that was not conducive to carcinogenesis.

Alb, PT-INR and ICG R15 values are well-known markers of hepatic reserve in the liver. In our study, the average values of Alb, PT-INR and ICG R15 in the D group were higher than in all of the other groups, showing the highest rate in the Child-Pugh class A patients. Generally, it has been reported that a decreased number of hepatocytes and decreased liver weight due to aging generally results in a decreased regeneration capacity of hepatocytes and a decreased hepatic blood flow (35-38). In addition, it has been suggested that the elderly often have a latent nutrient disturbance or metabolic anomalies, thus causing a declining trend in Alb and ICG R15 (39,40). However, in our study, many cases of HCC in the D group were present along with low values of ALT and TB, unlike the non-elderly group. Therefore, it appears that the progression of chronic liver disease was not promoted, and the hepatic reserve was consequently well-maintained in the elderly HCC group, despite the normal decline in the physiological liver function with age. As noted above, many cases of HCC in the elderly occur against a background of chronic liver disease with mild inflammation and fibrosis, and it is assumed that there are some characteristic factors which contribute to hepatocarcinogenesis in the elderly.

In recent years, age has become a point of focus in the progression of HCV-persistent infection and fibrosis or carcinogenesis. Mahmood *et al* (41) reported that age was the most important factor contributing to a high value of reactive oxygen metabolites in the blood associated with HCV-related chronic liver disease. Moreover, the involvement of oxidative stress has been previously reported in aging itself, which is believed to cause a decrease in various organ functions and immunity (42). The elderly are susceptible to oxidative stress due to a decrease in the SOD value related to the antioxidation mechanism and the decrease in NK cell activity (42,43). Therefore, hepatocarcinogenesis in the elderly may occur in association with only mild inflammation or fibrosis.

In addition, an increase in the rate of women is cited as one of the characteristics of HCC of the elderly. It has been reported that sex hormones such as estrogen and immune response play an important role in hepatocarcinogenesis (44,45). Since the levels of sex hormones decrease in ageing women, the risk of the incidence of HCC in elderly females increases.

In conclusion, we found that HCV-related HCC in the elderly occurred against a background of chronic liver disease with mild inflammation and fibrosis.

References

1. Tanaka H, Imai Y, Hiramatsu N, *et al*: Declining incidence of hepatocellular carcinoma in Osaka, Japan, from 1990 to 2003. *Ann Intern Med* 148: 820-826, 2008.
2. Taura N, Hamasaki K, Nakao K, *et al*: Aging of patients with hepatitis C virus-associated hepatocellular carcinoma: Long-term trends in Japan. *Oncol Rep* 16: 837-843, 2006.
3. Yano M, Yatsuhashi H, Inoue O, Inokuchi K and Koga M: Epidemiology and long term prognosis of hepatitis C virus infection in Japan. *Gut* 34: S13-S16, 1993.
4. Moriya T, Koyama T, Tanaka J, Mishiro S and Yoshizawa H: Epidemiology of hepatitis C virus in Japan. *Intervirology* 42: 153-158, 1999.
5. Tsukuma H, Tanaka H, Ajiki W and Oshima A: Liver cancer and its prevention. *Asian Pac J Cancer Prev* 6: 244-250, 2005.
6. Gerlach JT, Diepolder HM, Zachoval R, *et al*: Acute hepatitis C: high rate of both spontaneous and treatment-induced viral clearance. *Gastroenterology* 125: 80-88, 2003.
7. Ikeda K, Saitoh S, Koida I, *et al*: A multivariate analysis of risk factors for hepatocellular carcinogenesis: a prospective observation of 795 patients with viral and alcoholic cirrhosis. *Hepatology* 18: 47-53, 1993.
8. Kiyosawa K, Umemura T, Ichijo T, *et al*: Hepatocellular carcinoma: recent trends in Japan. *Gastroenterology* 127: S17-S26, 2004.
9. Yatsuhashi H and Yano M: Natural history of chronic hepatitis C. *J Gastroenterol Hepatol* 15: E111-E116, 2000.
10. Hamada H, Yatsuhashi H, Yano K, *et al*: Impact of aging on the development of hepatocellular carcinoma in patients with posttransfusion chronic hepatitis C. *Cancer* 95: 331-339, 2002.
11. Wali M, Harrison RF, Gow PJ and Mutimer D: Advancing donor liver age and rapid fibrosis progression following transplantation for hepatitis C. *Gut* 51: 248-252, 2002.
12. Dohmen K, Shirahama M, Shigematsu H, Irie K and Ishibashi H: Optimal treatment strategy for elderly patients with hepatocellular carcinoma. *J Gastroenterol Hepatol* 19: 859-865, 2004.
13. Ohishi W, Kitamoto M, Aikata H, *et al*: Impact of aging on the development of hepatocellular carcinoma in patients with hepatitis C virus infection in Japan. *Scand J Gastroenterol* 38: 894-900, 2003.
14. Teratani T, Ishikawa T, Shiratori Y, *et al*: Hepatocellular carcinoma in elderly patients: beneficial therapeutic efficacy using percutaneous ethanol injection therapy. *Cancer* 95: 816-823, 2002.
15. Ferrero A, Viganò L, Polastri R, *et al*: Hepatectomy as treatment of choice for hepatocellular carcinoma in elderly cirrhotic patients. *World J Surg* 29: 1101-1105, 2005.
16. Tsukioka G, Kakizaki S, Sohara N, *et al*: Hepatocellular carcinoma in extremely elderly patients: an analysis of clinical characteristics, prognosis and patient survival. *World J Gastroenterol* 12: 48-53, 2006.
17. Desmet VJ, Gerber M, Hoofnagle JH, Manns M and Scheuer PJ: Classification of chronic hepatitis: diagnosis, grading and staging. *Hepatology* 19: 1513-1520, 1994.
18. Poynard T, Bedossa P and Opolon P: Natural history of liver fibrosis progression in patients with chronic hepatitis C. The OBSVIRC, METAVIR, CLINIVIR, and DOSVIRC groups. *Lancet* 349: 825-832, 1997.
19. Bedossa P and Poynard T: An algorithm for the grading of activity in chronic hepatitis C. The METAVIR Cooperative Study Group. *Hepatology* 24: 289-293, 1996.
20. The French METAVIR Cooperative Study Group: Intraobserver and interobserver variations in liver biopsy interpretation in patients with chronic hepatitis C. *Hepatology* 20: 15-20, 1994.
21. The general rules for recording endoscopic findings on esophageal varices. *Jpn J Surg* 10: 84-87, 1980.
22. Dohmen K, Shigematsu H, Irie K and Ishibashi H: Trends in clinical characteristics, treatment and prognosis of hepatocellular carcinoma. *Hepatogastroenterology* 50: 1872-1877, 2003.
23. Zhou L, Rui JA, Wang SB, *et al*: Clinicopathological features, post-surgical survival and prognostic indicators of elderly patients with hepatocellular carcinoma. *Eur J Surg Oncol* 32: 767-772, 2006.
24. Shimauchi Y, Tanaka M, Koga K, *et al*: Clinical characteristics of patients in their 40s with HCV antibody-positive hepatocellular carcinoma. *Alcohol Clin Exp Res* 24: 64S-67S, 2000.
25. Nomura F, Ohnishi K, Honda M, Satomura Y, Nakai T and Okuda K: Clinical features of hepatocellular carcinoma in the elderly: a study of 91 patients older than 70 years. *Br J Cancer* 70: 690-693, 1994.
26. Hanazaki K, Kajikawa S, Shimozawa N, *et al*: Hepatic resection for hepatocellular carcinoma in the elderly. *J Am Coll Surg* 192: 38-46, 2001.
27. Yamagishi Y, Horie Y, Kajihara M, *et al*: Hepatocellular carcinoma in heavy drinkers with negative markers for viral hepatitis. *Hepatol Res* 28: 177-183, 2004.
28. Sata M, Fukuizumi K, Uchimura Y, *et al*: Hepatitis C virus infection in patients with clinically diagnosed alcoholic liver diseases. *J Viral Hepat* 3: 143-148, 1996.
29. Corrao G and Arico S: Independent and combined action of hepatitis C virus infection and alcohol consumption on the risk of symptomatic liver cirrhosis. *Hepatology* 27: 914-919, 1998.
30. Tarao K, Rino Y, Ohkawa S, *et al*: Association between high serum alanine aminotransferase levels and more rapid development and higher rate of incidence of hepatocellular carcinoma in patients with hepatitis C virus-associated cirrhosis. *Cancer* 86: 589-595, 1999.

31. Ono E, Shiratori Y, Okudaira T, *et al*: Platelet count reflects stage of chronic hepatitis C. *Hepato Res* 15: 192-200, 1999.
32. Matsumura H, Moriyama M, Goto I, Tanaka N, Okubo H and Arakawa Y: Natural course of progression of liver fibrosis in Japanese patients with chronic liver disease type C - a study of 527 patients at one establishment. *J Viral Hepat* 7: 268-275, 2000.
33. Karasu Z, Tekin F, Ersoz G, *et al*: Liver fibrosis is associated with decreased peripheral platelet count in patients with chronic hepatitis B and C. *Dig Dis Sci* 52: 1535-1539, 2007.
34. Nakayama H, Masuda H, Miyake H, Takayama T and Yokoyama E: Endoscopic prediction of hepatocellular carcinoma by evaluation of bleeding esophageal varices. *Digestion* 70: 233-239, 2004.
35. Okudaira M, Ikawa N, Yasuhara M, Kumagai T and Kurosu K: Liver weight of adult Japanese, especially recent weight values. *Hepato Res* 18: 95-103, 2000.
36. Zoli M, Iervese T, Abbati S, Bianchi GP, Marchesini G and Pisi E: Portal blood velocity and flow in aging man. *Gerontology* 35: 61-65, 1989.
37. Mooney H, Roberts R, Cooksley WG, Halliday JW and Powell LW: Alterations in the liver with ageing. *Clin Gastroenterol* 14: 757-771, 1985.
38. Jansen PL: Liver disease in the elderly. *Best Pract Res Clin Gastroenterol* 16: 149-158, 2002.
39. Wynne HA, Mutch E, Williams FM, James OF, Rawlins MD and Woodhouse KW: The relation of age to the acute effects of ethanol on acetanilide disposition. *Age Ageing* 18: 123-126, 1989.
40. Tietz NW, Shuey DF and Wekstein DR: Laboratory values in fit aging individuals - sexagenarians through centenarians. *Clin Chem* 38: 1167-1185, 1992.
41. Mahmood S, Kawanaka M, Kamei A, *et al*: Immunohistochemical evaluation of oxidative stress markers in chronic hepatitis C. *Antioxid Redox Signal* 6: 19-24, 2004.
42. Stadtman ER: Importance of individuality in oxidative stress and aging. *Free Radic Biol Med* 33: 597-604, 2002.
43. Hirokawa K, Utsuyama M, Zeng YX, Kurashima C and Michiyuki K: Immunological alterations with aging - laying a stress on recent progress in Japan. *Arch Gerontol Geriatr* 19: 171-183, 1994.
44. Shimizu I, Yasuda M, Mizobuchi Y, *et al*: Suppressive effect of oestradiol on chemical hepatocarcinogenesis in rats. *Gut* 42: 112-119, 1998.
45. Shimizu I, Inoue H, Yano M, *et al*: Estrogen receptor levels and lipid peroxidation in hepatocellular carcinoma with hepatitis C virus infection. *Liver* 21: 342-349, 2001.

Original Article

Adipocytokine involvement in hepatocellular carcinoma after sustained response to interferon for chronic hepatitis C

Nobuyoshi Fukushima,^{1,2} Ryoko Kuromatsu,¹ Teruko Arinaga-Hino,¹ Eiji Ando,¹ Akio Takata,¹ Shuji Sumie,¹ Masahito Nakano,¹ Takumi Kawaguchi,¹ Tatsuya Ide,¹ Takuji Torimura¹ and Michio Sata¹

¹Division of Gastroenterology, Department of Medicine, Kurume University School of Medicine, Kurume, and

²Department of Gastroenterology, Clinical Research Institute, National Hospital Organization Kyushu Medical Center, Fukuoka, Japan

Aim: Interferon (IFN) dramatically reduces the risk of hepatocellular carcinoma (HCC) after a sustained virological response (SVR) to chronic hepatitis C (CH-C). However, HCC still develops in some patients after SVR. To evaluate metabolic factors in patients with HCC occurring after SVR and to determine whether insulin resistance and adipocytokines were involved in this etiology.

Methods: We examined clinical and biochemical features, histological findings and serum levels of adipocytokine prior to IFN therapy and at the detection of HCC in nine patients who were diagnosed with HCC. As controls, 27 patients were included who showed SVR but had not been diagnosed with HCC for at least 5 years after SVR.

Results: Three of four patients who developed HCC within 5 years after SVR showed liver cirrhosis when HCC was diagnosed. Prior to IFN therapy, four of nine HCC patients were

diagnosed as having type 2 diabetes mellitus. Serum levels of leptin and insulin, Homeostatic Model of Assessment of Insulin Resistance and body mass index (BMI) were significantly higher and serum adiponectin was significantly lower in HCC patients at the time of HCC detection than in control patients more than 5 years after SVR. Six HCC patients had increased BMI and one HCC patient had a decreased BMI during the observation period.

Conclusion: Hepatic fibrosis may be tightly related to the emergence of HCC after SVR. Insulin resistance and adipocytokine disorders may be implicated in hepatocarcinogenesis after SVR, in part by promoting hepatic fibrosis.

Key words: adipocytokine, adiponectin, hepatocellular carcinoma, insulin resistance, leptin, sustained virological response.

INTRODUCTION

HEPATOCELLULAR CARCINOMA (HCC) is one of the most common cancers worldwide, especially in South-East Asia.¹ Hepatitis C virus (HCV) infection is a major risk factor for HCC. Seventy to eighty percent of Japanese patients with HCC are infected with HCV.² Interferon (IFN) is an antiviral agent against HCV that can eradicate the virus and which improves hepatic

inflammation and fibrosis,³ and is also believed to prevent clinical complications, including the development of HCC. IFN therapy for HCV was proven to lower the incidence of HCC, especially in patients who are treated successfully with IFN and show a sustained virological response (SVR) to the therapy.⁴ However, previous studies have revealed that HCC still develops in 1.5–4.0% of patients with SVR.^{5–11} Although advanced age,^{6–11} male sex,^{6,8} an advanced histological stage of hepatic damage,^{6–10} higher aspartate aminotransferase levels,¹¹ lower platelet counts,¹¹ alcohol intake^{7,9} and hepatic steatosis¹⁰ before IFN therapy are thought to be risk factors for HCC, the mechanism of carcinogenesis after SVR is not yet fully understood.

The liver is one of the major organs regulating glucose metabolism. Patients with chronic liver disease

Correspondence: Dr Nobuyoshi Fukushima, Division of Gastroenterology, Department of Medicine, Kurume University School of Medicine, 67 Asahi-machi, Kurume, Fukuoka 830-0011, Japan.
Email: kingsno1@med.kurume-u.ac.jp

Received 26 January 2010; revision 31 March 2010; accepted 5 June 2010.

frequently present with glucose intolerance which is referred to as hepatogenous diabetes with insulin resistance and hyperinsulinemia. The prevalence of glucose intolerance in HCV-related chronic liver disease is higher than in other liver diseases¹² and it has been suggested that HCV directly causes hepatic insulin resistance and hyperinsulinemia.^{13,14} Previous studies have shown that the incidence of HCC has been increased 2–4-fold in patients with diabetes mellitus (DM)^{15,16} and some reports have indicated that insulin resistance may be involved in cell growth and the carcinogenesis of HCC.^{17,18} Meanwhile, several recent studies have reported that obesity increases the risk of malignancy,¹⁹ including HCC.²⁰ In human obesity, intra-abdominal visceral fat content is strongly correlated with glucose intolerance.²¹ In the obese state, adipose tissue promotes inflammation and macrophages migrating into adipose tissue secrete adipocytokines, including tumor necrosis factor- α (TNF- α), interleukin-6 (IL-6) and monocyte chemoattractant protein-1 (MCP-1). Enlarged adipose cells also secrete these adipocytokines and free fatty acids (FFA). These adipocytokines and FFA can induce insulin resistance in the whole body.^{22–24}

Adiponectin is an adipose tissue-specific plasma protein and is specifically secreted from fat tissue. Hypoadiponectinemia may be implicated in the pathogenesis of the various disorders which comprise metabolic disease,²⁵ and may be related to the pathogenesis of liver disease through cells expressing the adiponectin receptor in the liver.²⁶ Leptin is another circulating hormone secreted by adipocytes which acts as an important signaling molecule in energy regulation and food intake,²⁷ and modulates multiple physiological and pathophysiological states.^{27,28}

Meanwhile, adiponectin is thought to be involved in the carcinogenesis of several cancers.^{29,30} Adachi *et al.* showed that an adiponectin receptor, T-cadherin, was selectively expressed in intratumoral capillary endothelial cells in HCC, suggesting a positive role for T-cadherin in mediating angiogenesis in HCC.³¹ In addition, leptin may also play an important role in the processes of initiation and progression of various human cancers.^{32,33} Chen *et al.* reported that leptin induced proliferation and inhibited apoptosis in human HCC cells.³⁴ Zhou *et al.* showed that leptin stimulated Hep G2 cell proliferation through increased DNA synthesis and the enhancement of mitotic activity.³⁵ Ribatti *et al.* have reported an involvement of leptin and the leptin receptor in angiogenesis in human HCC.³⁶

In this study, we investigated whether obesity, insulin resistance, leptin and adiponectin were involved in the development of HCC in patients after SVR undergoing IFN treatment for chronic hepatitis C (CH-C).

METHODS

Patients

THE SUBJECTS IN this study were nine patients who were diagnosed with HCC at Kurume University Hospital after being treated with IFN therapy for CH-C between 1993 and 2003 and exhibiting SVR. Seven patients were treated with IFN at Kurume University Hospital or affiliated hospitals, and two were treated with IFN at other hospitals. The diagnosis of HCC was based on histological examination for eight patients and on computed tomography (CT), magnetic resonance imaging (MRI) and elevation of tumor markers in one patient. One thousand one hundred and forty-three patients received IFN therapy for CH-C and 308 patients achieved SVR between 1993 and 2003 at Kurume University Hospital. Out of these patients, we selected 27 patients who met the following requirements as control: (i) periodically observed at our hospital after SVR and had no evidence of HCC for over 5 years from the time of SVR until the last outpatient visit; (ii) blood samples before IFN treatment and at following outpatient visit were stored; and (iii) hepatitis B virus surface antigen (HBsAg) was negative in the serum. All control patients and six HCC patients were followed up after SVR. These patients received ultrasonography (US), computed tomography (CT) and examination of tumor markers at least once per year. HCC were detected 1 year or later after SVR regarding six patients developing HCC with periodic observation. Another three patients developing HCC did not undergo periodic observation from the time of SVR to the detection of HCC. Intervals from SVR until detection of HCC regarding these three patients were 106, 156 and 108 months, respectively. SVR was defined as being serum HCV RNA negative for more than 6 months after the termination of IFN therapy. Serum HCV RNA was measured using the COBAS AmpliPrep/COBAS TaqMan HCV Test (Roche Molecular Systems, Branchburg, NJ, US). This newly established technique has excellent sensitivity for the detection of HCV RNA and is more sensitive than conventional methods of detection.³⁷

We had less information prior to their IFN therapy about the two HCC patients treated with IFN at other hospitals. Upon detection of their HCC, both patients

were seropositive for anti-HCV antibodies, seronegative for HBsAg and negative for serum HCV RNA as determined by the COBAS AmpliPrep/COBAS TaqMan HCV Test. The other seven HCC patients and 27 control patients were all seropositive for anti-HCV antibodies, positive for serum HCV RNA and seronegative for hepatitis B antigen prior to IFN therapy. The HCV RNA load was quantitated by competitive reverse-transcription polymerase chain reaction, a branched-DNA probe assay, or by the Amplicor-HCV monitor assay.^{38–40} HCV genotype was determined according to previously described methods.⁴¹ We excluded patients who had a coexisting liver disease, such as autoimmune hepatitis or primary biliary cirrhosis. Past occurrence of HCC and treatment history for HCC were also part of the exclusion criteria. Informed consent was obtained from each patient and the study was approved by the Ethical Committee of Kurume University. The study was carried out according to the ethical guidelines of the 1975 Declaration of Helsinki.

Metabolic parameters

We compared the clinical characteristics of the nine HCC patients and 27 control patients just before the start of IFN therapy. Body mass index (BMI) was calculated as weight in kilograms/height in meters squared. We obtained BMI values prior to IFN therapy from seven HCC patients treated with IFN in Kurume University Hospital or affiliated hospitals, and from 23 out of 27 control patients. We obtained BMI values for nine HCC patients at the time of HCC diagnosis, and for 22 of the control patients more than 5 years after SVR. The high molecular weight (HMW) form of adiponectin, leptin, insulin and glucose were measured from blood samples stored at -80°C . Blood samples were obtained from five HCC patients and all control patients within 9 months before the start of IFN therapy. Additional samples were obtained within 9 months after the time of SVR in three patients who developed HCC and from all control patients. Blood was also collected later as the last time point sample at the time of HCC diagnosis for nine HCC patients and more than 5 years after SVR for all the control patients. All samples were obtained at fasting time.

Serum insulin levels were measured using an enzyme-linked immunoassay kit (Lumipulse Presto Insulin; Fujirebio, Tokyo, Japan). Plasma glucose levels were measured using the hexokinase-glucose-6-phosphate dehydrogenase method. Insulin resistance was calculated from fasting levels of glucose and insulin using the

Homeostatic Model of Assessment (HOMA) method. The formula for the HOMA model is: insulin resistance (HOMA-IR) = fasting glucose (mg/dL) \times fasting insulin ($\mu\text{U/mL}$) / 405. Adiponectin is present in serum in a number of forms, and often exists as a trimer or hexamer of HMW.⁴² Adiponectin in the HMW form has been reported to play a primary role in hepatic and whole-body insulin sensitivity and to have significant anti-inflammatory effects.⁴³ For this reason, we measured serum HMW adiponectin in this study. The levels of HMW adiponectin were determined using a HMW adiponectin assay kit (Fujirebio). Leptin levels were determined using a human leptin RIA kit (Linco Research, St Charles, MO, USA). Values of each metabolic parameter at each time point were compared between the HCC patients and controls.

Histopathology

We compared the histological features of liver tissue before IFN therapy in HCC patients and controls and also compared these with liver tissue at the time of detection of HCC. Liver biopsies were performed on 32 patients treated with IFN at Kurume University Hospital or at affiliated hospitals less than 1 year before the start of IFN therapy. One HCC patient underwent a biopsy 8 years before IFN therapy and showed cirrhosis at that point, and no biopsy was performed on one of the control patients. Assessment of the histological diagnosis was made according to the classifications of Desmet *et al.*⁴⁴ Hepatic steatosis was also graded by a modification of the Brunt scoring system as described^{45,46} using the following categories: steatosis less than 5%, 5–33%, 33–66% and patients with over 66% steatosis were categorized as grades 0, 1, 2 and 3, respectively. All tissue samples were evaluated by two independent hepatologists. We also examined non-cancerous hepatic lesions resected with HCC from seven patients. These seven cases were assessed for histological tumor differentiation⁴⁷ in the resected specimen and one other case was examined using a specimen obtained by aspiration tumor biopsy.

Statistical analysis

Statistical analysis was performed using SPSS ver. 12.0J. Fisher's exact probability test was used to compare categorical data. Differences between two groups were measured using the Mann-Whitney *U*-test. A relationship between different continuous variables was measured by linear regression analysis. $P < 0.05$ was considered statistically significant.

Table 1 Characteristics of nine patients with HCC after sustained virological response

Age at detection of HCC (years)†	61 (54–75)
Maximal tumor diameter (mm)†	26.8 (16–43)
No. of tumors (1/2/3)	7/1/1
Tumor stage (I/II/III)‡	1/5/3
Differentiation (well/moderately)	3/5§
Child–Pugh score (at detection of HCC)¶	A5 for all
Therapy (resection/RFA/radiation)	7/1/1
Periodic observation	
Before detection of HCC (yes/no)	6/3

†Mean (range).

‡Based on the tumor–node–metastasis (TNM) classification.⁴⁸

§Tumor histology was not available for one case.

¶Classified according to Child–Pugh classification.⁴⁹

HCC, hepatocellular carcinoma; RFA, radiofrequency ablation.

RESULTS

Characteristics of nine patients with HCC after SVR

THE CHARACTERISTICS OF nine SVR patients who developed HCC are summarized in Table 1. Eight patients underwent curative therapy, including resection for seven patients and percutaneous radiofrequency ablation for one patient. Another patient was evaluated for resection, but ultimately received radiation therapy due to dementia.

Table 2 shows the clinical and histological profiles of HCC patients. Histological fibrosis worsened in three patients and did not change in one patient out of the six in whom we could compare liver histology at both time points. Four out of five alcohol consumers with HCC continued drinking alcohol after IFN therapy, and only patient 6 stopped drinking after IFN therapy. Four patients developed HCC within 5 years after SVR and five patients developed HCC more than 5 years after SVR. Three of four patients in the former group showed liver cirrhosis when HCC was diagnosed, and no patient showed liver cirrhosis in the latter group although histological data of two patients were not available.

Comparison of baseline characteristics of SVR patients with and without HCC after SVR

A comparison of the baseline characteristics of SVR patients with and without HCC is summarized in Table 3. Patients with regular smoking were significantly more in the HCC group than in the control group ($P = 0.046$). Before the start of IFN therapy, four out of nine HCC patients and none of the controls were diagnosed with type 2 DM, a significant difference between the two groups ($P = 0.002$). The average observation period from the time of SVR to the detection of HCC for the HCC group was 81.7 months (range

Table 2 Clinical and histological profiles of SVR patients with HCC

	Before IFN therapy			At the detection of HCC			DM	Alcohol (23 g/day)	Interval (months)§
	G/S†	Steatosis‡	BMI	G/S†	Steatosis‡	BMI			
Patient 1	2/2	1	23.4	1/4	0	23.7	–	–	22
Patient 2	2/4	2	26.5	1/4	1	28.9	+	–	38
Patient 3	2/1	0	19.3	1/4	1	23.1	–	–	42
Patient 4¶	2/2	1	27.8	1/1	0	29.6	–	+	48
Patient 5	2/2	0	20.2	N/A	NA	24.1	–	+	80
Patient 6	2/2	1	29.8	2/1	0	27.6	+	+	106
Patient 7¶	NA	NA	NA	NA	NA	24.5	+	–	108
Patient 8¶	1/1	2	23.3	1/2	1	24.9	–	+	135
Patient 9¶	NA	NA	NA	1/1	1	26.9	+	+	156

†G and S refer to hepatic inflammatory grading (A0–3) and fibrotic staging (F0–4), respectively.

‡Graded 0–3.

§Interval from SVR until detection of HCC.

¶Adiponectin, leptin, insulin and HOMA-IR prior to IFN therapy could not be acquired for patients 4 and 8. Detailed information for IFN therapy, laboratory and histological data, and metabolic parameters prior to IFN therapy could not be acquired for patients 7 and 9.

BMI, body mass index; DM, type 2 diabetes mellitus; HOMA-IR, Homeostatic Model of Assessment of Insulin Resistance; HCC, hepatocellular carcinoma; IFN, interferon; NA, not available; SVR, sustained virological response.

Table 3 Comparison of baseline characteristics between SVR patients with and without HCC after SVR

	HCC group	Control group	P-value†
Age at IFN induction (years)‡	53.0 (44–65)	51.3 (23–74)	NS
Sex (male/female)	5/4	11/16	NS
Alcohol (≥ 23 g/day) (+/-)	5/4	6/21	NS
Smoking (+/-)	6/3	7/20	0.046
Type 2 diabetes mellitus (+/-)	4/5	0/27	0.002
HBcAb (+/-)	5/4	11/16	NS
HCV viral load before IFN (high/low)§	2/5	8/18	NS
Virus genotype (1/2)	3/3	15/12	NS
IFN (a/b/a + b/consensus)	5/2/0/0	19/4/2/2	NS
(with RBV¶; PEG IFN††)	(3:0)	(4:1)	
Total IFN amount ($\times 10^4$) (mean)	59 800	56 392	NS
Observation period after SVR (months)‡	81.7 (22–156)	102 (61–149)	NS
AST (IU/L)‡‡	75.9 \pm 49.9	62.2 \pm 44.7	NS
ALT (IU/L)‡‡	69.3 \pm 40.7	78.3 \pm 70.4	NS
Total bilirubin (mg/dL)‡‡	1.0 \pm 0.6	0.8 \pm 0.2	NS
Albumin (g/dL)‡‡	3.7 \pm 0.3	4.0 \pm 0.2	NS
Prothrombin time (%)‡‡	93.2 \pm 12.4	97.8 \pm 16.4	NS
Platelet ($\times 10^4$ /mL)‡‡	12.3 \pm 3.4	15.4 \pm 5.3	NS
Hyaluronic acid (ng/mL)‡‡	187.5 \pm 153.4	122.9 \pm 212.6	0.045
α -Fetoprotein (ng/mL)‡‡	42.4 \pm 81.4	11.0 \pm 29.2	NS
Child–Pugh score (5/6/7)	5/0/1	24/1/0	NS

All data were based on the patients from whom it was available for each characteristics.

†Comparison between patients with HCC (HCC group) and without HCC (control group) after SVR. *P*-values were calculated with Fisher's exact probability test and the Mann–Whitney *U*-test.

‡Mean (range).

§When the serum HCV RNA level was more than 1 Meq/mL by branched DNA assay, more than 10^6 copies/mL by competitive reverse transcription polymerase chain reaction, or more than 10^5 copies/mL by the Amplicor-HCV monitor assay, it was determined to be a high viral load.

¶Number of the patients in whom Ribavirin was used in combination.

††Number of patients in whom pegylated interferon was used.

‡‡Means \pm standard deviation.

ALT, alanine aminotransferase; AST, aspartate aminotransferase; HBcAb, hepatitis B core antigen; HCC, hepatocellular carcinoma; HCV, hepatitis C virus; IFN, interferon; NS, not significant.

22–156), and the average time from SVR to the outpatient visit when we obtained the last follow-up results for the control group was 102 months (range 61–149). There was no significant difference in the length of the observation period between the two groups. Serum levels of hyaluronic acid prior to IFN therapy were significantly higher in the HCC group than in the control group ($P = 0.045$).

Changes in adipocytokines during the observation period

Our findings on metabolic parameters are summarized in Table 4. Prior to IFN therapy and at the time of SVR, serum levels of adiponectin and leptin were not significantly different between the HCC group and the control group. However, adiponectin was significantly lower in

the HCC group upon HCC detection than in the controls at their follow-up visit more than 5 years after SVR ($P = 0.030$), and leptin was significantly higher in HCC patients than in controls at that time ($P = 0.036$). When we only analyzed patients for whom we had data both prior to IFN therapy and at the last time point, percentage of change per year in adiponectin level was significantly lower ($P = 0.046$) and that in leptin level was significantly higher ($P = 0.003$) in the HCC patients than in controls. Figure 1(A) shows changes in adiponectin and Figure 1(B) shows changes in leptin in both groups during the observation period.

In all cases in both the HCC group and control group, serum leptin levels were positively correlated with serum insulin levels and HOMA-IR but not with adiponectin levels or BMI at the last time point (data not

Table 4 Metabolic parameters of the study population

	Adiponectin ($\mu\text{g/mL}$)		
	Before IFN Tx	at SVR	Last Time point†
HCC group	10.8 \pm 8.6 (5)	8.8 \pm 4.0 (3)	5.0 \pm 2.7 (9)
Control group (27)‡	8.5 \pm 4.1	8.0 \pm 4.2	8.1 \pm 4.0
<i>P</i> -value§	NS	NS	0.03
	Leptin (ng/mL)		
	Before IFN Tx	At SVR	Last Time point†
HCC group	8.9 \pm 2.9 (5)	8.4 \pm 2.1 (3)	11.1 \pm 6.9 (9)
Control group (27)‡	6.4 \pm 4.7	6.5 \pm 5.7	7.0 \pm 6.8
<i>P</i> -value§	NS	NS	0.036
	BMI		
	Before IFN Tx		Last Time point†
HCC group		24.3 \pm 3.9 (7)	25.9 \pm 2.4 (9)
Control group		21.7 \pm 2.6 (23)	21.9 \pm 2.5 (22)
<i>P</i> -value§		NS	0.001
	Insulin ($\mu\text{U/mL}$)		
	Before IFN Tx	At SVR	Last Time point†
HCC group	16.4 \pm 10.6 (5)	9.3 \pm 1.3 (3)	11.5 \pm 6.9 (9)
Control group (27)‡	8.9 \pm 6.6	8.7 \pm 9.8	6.3 \pm 6.8
<i>P</i> -value§	0.022	NS	0.03
	HOMA-IR		
	Before IFN Tx	At SVR	Last Time point†
HCC group	5.0 \pm 4.9 (5)	1.6 \pm 0.6 (3)	2.9 \pm 2.1 (9)
Control group (27)‡	1.8 \pm 1.7	1.7 \pm 1.7	1.3 \pm 0.9
<i>P</i> -value§	0.014	NS	0.004

Data are shown as the means \pm standard deviation.

P-values were calculated with the Mann–Whitney *U*-test.

†Data at the time of HCC diagnosis in the HCC group and at an outpatient visit more than 5 years after SVR for control group.

Number in parenthesis means the number of cases involved in the analysis.

‡The number of control group cases was 27 except BMI analysis.

§Comparison between patients with HCC (HCC group) and without HCC (Control group) after SVR.

HCC, hepatocellular carcinoma; HOMA-IR, Homeostatic Model of Assessment of Insulin Resistance; IFN, interferon; NS, not significant; SVR, sustained virological response; Tx, therapy.

shown). In addition, adiponectin levels were inversely correlated with serum insulin and HOMA-IR but not with BMI at the last time point (data not shown). Prior to IFN therapy, leptin and adiponectin levels were not correlated with other metabolic parameters (data not shown). At the time of SVR, serum leptin levels were positively correlated with serum insulin levels and HOMA-IR, however, serum adiponectin levels were not correlated with these parameters and leptin levels (data not shown).

Variation in BMI and insulin resistance during the period of observation

Prior to IFN therapy, the BMI of HCC patients and control patients were not significantly different (Table 4). However, at the last time point, the BMI of HCC patients was significantly higher than that of control patients ($P = 0.001$). Figure 1(C) shows the changes in BMI during the period of observation. For six out of seven HCC patients, BMI was higher at the time of HCC detection compared to before IFN therapy. When we only analyzed patients for whom we had data both prior to IFN therapy and at the last time point, percentage of change per year in BMI values was significantly higher in the HCC patients than in controls ($P = 0.048$). Prior to IFN therapy, serum insulin and HOMA-IR were significantly higher in HCC patients than in controls ($P = 0.022$ and $P = 0.014$, respectively) (Table 4). In addition, serum insulin and HOMA-IR were significantly higher in HCC patients than in controls at the last time point ($P = 0.003$ and $P = 0.004$, respectively). Percentage of change per year in the insulin level or HOMA-IR in the HCC patients were not significantly different from those in the controls (data not shown). At the time of SVR, there was no significant difference in serum insulin and HOMA-IR between these two groups.

Histological features of the study population

Table 5 shows the histological features of the study population. When patients were divided into the hepatic inflammation groups A0–1 and A2–3, the histological activity grade at HCC detection was significantly improved compared to that before IFN therapy ($P = 0.029$). The histological activity grade prior to IFN therapy was not significantly different between the HCC group and the control group. When patients were divided into the fibrosis groups F0–1 and F2–4, the histological fibrotic stage at HCC detection was not significantly improved compared to that before IFN

therapy, and fibrotic stage prior to IFN therapy was not significantly different between the HCC group and the control group. When patients were divided into the hepatic steatosis groups grade 0–1 and grade 2–3, there was no significant difference in the two groups before IFN therapy. Hepatic steatosis at the time of HCC detection was not significantly different from before IFN therapy in the HCC group.

DISCUSSION

FOR OUR STUDY, we selected patients who had been followed for more than 5 years after SVR with no detectable HCC as a control group because we wanted to be sure to exclude patients who developed HCC from the control group as much as possible. When we divide the patients in the HCC group into two groups – one group in which patients developed HCC within 5 years after SVR and another group in which patients developed HCC more than 5 years after SVR – three of four patients in the former group showed liver cirrhosis when HCC was diagnosed. This observation suggests that liver cirrhosis is tightly related to the development of HCC with a relatively short interval after SVR.

Insulin is known to be an important factor not only for a variety of metabolic pathways, but also for cell proliferation.¹⁷ One of the major functions induced by elevated serum insulin is the activation of the mitogen-activated protein kinase cascade which has effects on cell proliferation.⁵⁰ Saito *et al.* reported that hyperinsulinemia activated the growth of human HCC cells from patients with liver cirrhosis.¹⁸ Insulin resistance has also been shown to induce fibrotic progression in the liver with CH-C.⁵¹ Advanced hepatic fibrosis is known to be a major risk factor for the occurrence of HCC in patients with CH-C,⁵² even after SVR.^{6–10} In our study, liver fibrosis was not significantly improved at the time of detection of HCC compared with before IFN therapy, and had progressed in three patients and was unchanged in another one out of the seven patients for whom we were able to compare the histological findings at both time points, although previous studies had shown an improvement of hepatic fibrosis after SVR to IFN therapy.⁵³ Kawaguchi *et al.* have reported that clearance of HCV improves insulin resistance.¹⁴ In this study, both the HCC group and control group also showed a decline in serum insulin levels and HOMA-IR after SVR, and significant differences in these parameters between the two groups were not found once at the time of SVR; however, the values in the HCC group were still relatively high at the time of detection of HCC (normal









Figure 1 Changes in metabolic parameters during the period of observation in sustained virological response (SVR) patients. Hepatocellular carcinoma (HCC) group (G1) comprised SVR patients who developed HCC, and control group (G2) comprised those who did not develop HCC. (A-a) Changes in the high molecular weight form of adiponectin in the serum (B-a) changes in serum leptin and (C-a) changes in body mass index (BMI). Serum samples were collected and BMI values were obtained before the start of interferon (IFN) therapy. Serum samples and BMI values were also obtained as the last time point sample when HCC was diagnosed in the HCC group and at an outpatient visit more than 5 years after SVR for the control group. Percentages of change per year in the level of the high molecular weight form of adiponectin (A-b), leptin level (B-b) and BMI value in each group were shown. We only included patients for whom both time points were available in this analysis. The data represent means ± standard deviation. *Statistically significant differences between the indicated groups ($P < 0.05$).

range: insulin 1.84–12.2 $\mu\text{U/mL}$, HOMA-IR ≤ 1.6) and were significantly different from those of the control group at the last time point. Hyperinsulinemia might be one of the reasons why hepatic fibrosis was not shown to improve in the SVR patients developing HCC.

Adiponectin has been reported to increase insulin sensitivity⁵⁴ and to inhibit hepatic fibrosis,⁵⁵ however, adiponectin paradoxically decreases with the accumulation of visceral fat.⁵⁶ Therefore, hypoadiponectinemia resulting from obesity can cause insulin resistance and accelerate hepatic fibrosis. Two out of three HCC patients in which histological fibrosis worsened showed very low ratio of adiponectin at the last time point to that before IFN therapy (0.2 and 0.51), and another one case that lacked the data before IFN therapy showed very low value of adiponectin at HCC detection (2.7 $\mu\text{g/mL}$).

Serum leptin levels are positively correlated with BMI as fat tissue increases.⁵⁷ Leptin has inhibitory effects on insulin resistance,⁵⁸ however, the risk of insulin resistance rises with obesity and in these patients, increased leptin may not be sufficient to improve insulin resistance. It is conceivable that the serum level of leptin is simply a reflection of the degree of insulin resistance.⁵⁹ Leptin facilitates hepatic fibrosis through the induction of TNF- α , the proliferation of hepatic stellate cells and

stimulation of the sympathetic nervous system.^{55,60} In addition, leptin have been reported to be associated with proliferation of HCC.^{31,34–36} Taken together, body-weight gain leads to insulin resistance, increased leptin and decreased adiponectin, and these metabolic alterations may induce the initiation and progression of HCC, in part by promoting hepatic fibrosis in the HCC group.

Serum leptin levels have been reported to decrease after the end of IFN therapy for hepatitis C and then to recover to pretreatment levels after a long follow up.⁶¹ One study reported that serum adiponectin was increased at 6 months after IFN therapy,⁶² but another reported decreased adiponectin at 12 weeks after IFN therapy.⁶³ Thus, the influence of IFN on the levels of serum adiponectin for an extended period is unclear.

Fibrotic stage prior to IFN therapy was not significantly different between the HCC group and the control group ($P = 0.106$), although the number of patients in both groups was very low. Serum levels of hyaluronic acid prior to IFN therapy were significantly higher in the HCC group than in the control group ($P = 0.045$), therefore, we could not disclaim an association between hepatic fibrosis prior to IFN therapy and the occurrence of HCC after SVR.

Table 5 Histological features of the study population

	Inflammatory grading A 0/1/2/3		Fibrotic staging F 0/1/2/3/4		Steatosis Grade 0/1/2/3	
	HCC group	Control group	HCC group	Control group	HCC group	Control group
Before IFN	0/1/6/0	2/11/13/0	0/2/4/0/1	3/14/7/2/0	2/3/2/0	11/13/1/1
At HCC detection	*0/6/1/0	–	0/3/1/0/3	–	3/4/0/0	–

* $P = 0.029$ when A0–1 and A2–3 were compared between the two time points of the HCC group using the Mann–Whitney U -test. Prior to IFN therapy, the activity grade was not significantly different between the HCC group and control group. When divided into F0–1 and F2–4 groups for hepatic fibrosis, and when divided into grade 0–1 and grade 2–3 for hepatic steatosis, there was no significant difference between the two groups before IFN therapy and between the two time points of HCC group. HCC, hepatocellular carcinoma; IFN, interferon.

An association between smoking and hepatic fibrosis, and also an association between smoking and HCC were reported previously.^{64,65} Ratio of patients with smoking was significantly higher in the HCC group than in the control group. Therefore, smoking might facilitate hepatic fibrosis and increase the risk of HCC synergistically with the other risk factors in our study.

It is possible that undetectable HCC may have already developed in patients in this study before IFN therapy, based on information about the estimated doubling time of HCC.⁵² Even so, abnormalities in metabolic factors prior to IFN therapy and alterations of these factors after IFN therapy might affect the progression of HCC.

This study was conducted at a single medical center and the number of enrolled patients was limited. As a result, the number of patients developing HCC after SVR and of controls was fairly low, and thus the accuracy of our statistical analysis was limited. A large-scale study and careful analysis are needed to confirm our results, which indicate the importance of metabolic factors in the hepatocarcinogenesis process after SVR.

In conclusion, hepatic fibrosis may be tightly related to the emergence of HCC after SVR, and insulin resistance and adipocytokine disorders may be implicated in hepatocarcinogenesis after SVR, in part by promoting hepatic fibrosis. This study is the first to report the correlation between the development of HCC after SVR and metabolic factors including insulin resistance, obesity and adipocytokine levels.

ACKNOWLEDGMENTS

WE THANK DR Kenji Hirai and Dr Shuichiro Nagata for providing us with the patient sample as well as extensive clinical, laboratory and histological data.

REFERENCES

- 1 Wong F, Choi T. Primary liver cancer. Asian experience. In: Blumgart LH, ed. *Surgery of Liver and Biliary Tract*. London: Churchill-Livingstone, 1988; 1135–51.
- 2 Yoshizawa H. Hepatocellular carcinoma associated with hepatitis C virus infection in Japan: projection to other countries in the foreseeable future. *Oncology* 2002; 62 (Suppl 1): 8–17.
- 3 Hiramatsu N, Hayashi N, Kasahara A *et al.* Improvement of liver fibrosis in chronic hepatitis C patients treated with natural interferon alpha. *J Hepatol* 1995; 22: 135–42.
- 4 Yoshida H, Shiratori Y, Moriyama M *et al.* Interferon therapy reduces the risk for hepatocellular carcinoma: national surveillance program of cirrhotic and noncirrhotic patients with chronic hepatitis C in Japan. IHIT Study Group. Inhibition of Hepatocarcinogenesis by Interferon Therapy. *Ann Intern Med* 1999; 131: 174–81.
- 5 Kasahara A, Hayashi N, Mochizuki K *et al.* Risk factors for hepatocellular carcinoma and its incidence after interferon treatment in patients with chronic hepatitis C. Osaka Liver Disease Study Group. *Hepatology* 1998; 27: 1394–402.
- 6 Makiyama A, Itoh Y, Kasahara A *et al.* Characteristics of patients with chronic hepatitis C who develop hepatocellular carcinoma after a sustained response to interferon therapy. *Cancer* 2004; 101: 1616–22.
- 7 Iwasaki Y, Takaguchi K, Ikeda H *et al.* Risk factors for hepatocellular carcinoma in Hepatitis C patients with sustained virologic response to interferon therapy. *Liver Int* 2004; 24: 603–10.
- 8 Kobayashi S, Takeda T, Enomoto M *et al.* Development of hepatocellular carcinoma in patients with chronic hepatitis C who had a sustained virological response to interferon therapy: a multicenter, retrospective cohort study of 1124 patients. *Liver Int* 2007; 27: 186–91.
- 9 Tokita H, Fukui H, Tanaka A *et al.* Risk factors for the development of hepatocellular carcinoma among patients with chronic hepatitis C who achieved a sustained virological response to interferon therapy. *J Gastroenterol Hepatol* 2005; 20: 752–8.
- 10 Tanaka A, Uegaki S, Kurihara H *et al.* Hepatic steatosis as a possible risk factor for the development of hepatocellular carcinoma after eradication of hepatitis C virus with antiviral therapy in patients with chronic hepatitis C. *World J Gastroenterol* 2007; 13: 5180–7.
- 11 Ikeda M, Fujiyama S, Tanaka M *et al.* Risk factors for development of hepatocellular carcinoma in patients with chronic hepatitis C after sustained response to interferon. *J Gastroenterol* 2005; 40: 148–56.
- 12 Mason AL, Lau JY, Hoang N *et al.* Association of diabetes mellitus and chronic hepatitis C virus infection. *Hepatology* 1999; 29: 328–33.
- 13 Shintani Y, Fujie H, Miyoshi H *et al.* Hepatitis C virus infection and diabetes: direct involvement of the virus in the development of insulin resistance. *Gastroenterology* 2004; 126: 840–8.
- 14 Kawaguchi T, Ide T, Taniguchi E *et al.* Clearance of HCV improves insulin resistance, beta-cell function, and hepatic expression of insulin receptor substrate 1 and 2. *Am J Gastroenterol* 2007; 102: 570–6.
- 15 Adami HO, Chow WH, Nyren O *et al.* Excess risk of primary liver cancer in patients with diabetes mellitus. *J Natl Cancer Inst* 1996; 88: 1472–7.
- 16 El-Serag HB, Tran T, Everhart JE. Diabetes increases the risk of chronic liver disease and hepatocellular carcinoma. *Gastroenterology* 2004; 126: 460–8.
- 17 Gruppuso PA, Boylan JM, Bienieki TC, Curran TR Jr. Evidence for a direct hepatotropic role for insulin in the fetal