

drug delivery was to use a synthetic peptide corresponding to the extracellular loop region of occludin *in vitro* [12]. The testes are rich in receptors for follicle-stimulating hormone. The effects of follicle-stimulating hormone-fused occludin peptide on the *in vivo* blood-testis barrier were investigated. The fusion protein modulated the blood-testis barrier, resulting in delivery of inulin into the testis [25]. Astrovirus infection causes diarrhea [26]. Moser and co-workers found that the astrovirus capsid disrupted occludin and increased the permeability of the TJ barrier without cytotoxicity in human intestinal cells [27]. A pro-inflammatory cytokine, IL-1 β causes a functional opening of the intestinal TJ barrier without induction of apoptosis [28,29]. The IL-1 β -induced enhancement of TJ permeability was mediated by downregulation of occludin through an increase in the myosin light chain kinase [29,30].

Thus, occludin peptides containing the ligand-targeting motifs and novel types of occludin modulators, such as the component capsid and the activator of myosin light chain kinase, may provide novel methods to deliver drugs into target tissues across endothelial cell sheets.

2.3 Ephrin

Ephrin-A2, a family of receptor tyrosine kinases, directly phosphorylates claudin-4 in epithelial cells, leading to the disruption of the epithelial barrier function [31]. Intravenous administration of ephrin-A2 ligand causes vascular permeability in the lungs, resulting in the leakage of albumin into the lungs of rats [32]. The ephrin-A2 ligand is altered in the disruption of the TJ barrier in the lungs of rats and in cultured lung vascular endothelial cells [32]. High levels of ephrin-A2 mRNA are also expressed in the intestine [33]. A modulator of the ephrin-A2 system will be a novel type of pulmonary and intestinal absorption enhancer.

2.4 Zonula occludens toxin

Zonula occludens toxin (Zot) is a 44.8-kDa envelope protein of *Vibrio cholera*, and zonulin is the intestinal Zot analogue that governs the permeability of intercellular TJs [34-36]. Zot and Zot derivatives are reversible TJ openers that enhance the delivery of drugs through the paracellular route without toxicity [35-40]. The Zots bind to a putative receptor on the apical surface of enterocytes, leading to protein kinase C-mediated polymerization of soluble G-actin and the subsequent loosening of TJs [38,41]. Zot enhanced the absorption of insulin in diabetic rats, and the bioavailability of oral insulin was sufficient to lower the serum glucose concentrations to an extent that was comparable to the parenteral injection of the hormone [35]. In 2001, an active fragment of Zot, Δ G with a molecular mass of 12 kDa, was identified [42]. In 2008, a hexapeptide derived from Zot, AT1002, was found to enhance absorption [43]. AT1002 increased permeability in human epithelial cell sheets without cytotoxicity and enhanced duodenal absorption of ciclosporin A.

2.5 Chitosan

Chitosan is derived from chitin, a polysaccharide found in the exoskeletons of insects, arachnids, and crustaceans. Chitosan is a nontoxic, biocompatible and mucoadhesive polymer that is a safe and efficient intestinal permeation enhancer for the absorption of drugs [44-46]. The chitosan-mediated activation of protein kinase C α is followed by the redistribution of ZO-1 and an increase in TJ permeability, suggesting that the protein kinase C α -dependent signal transduction pathway affects TJ integrity [47]. The oral administration of recently developed chitosan-coated nanoparticles containing insulin dramatically decreased blood glucose levels in diabetic rats [48].

2.6 HA, HAstV-1

Hemagglutinin (HA), a non-toxic component of the large 16S of the botulinum neurotoxin [49], and the human astrovirus serotype 1 (HAstV-1) capsid [27] may be a novel absorption enhancer via the paracellular route. The HA protein affected distribution of occludin, ZO-1, E-cadherin and β -catenin, and increased TJ permeability in human intestinal epithelial cells without cytotoxicity [49]. When HAstV-1 infected a Caco-2 cell monolayer from the apical side, the paracellular permeability was increased. UV-inactivated HAstV-1 also increased the permeability and disrupted occludin, indicating that the enhancement of the permeability was not dependent on viral replication [27]. Further analysis of the mode of action of these toxin- and virus-derived enhancers will lead to the development of novel intestinal absorption enhancers.

3. Physiological barriers modulated by TJ modulators

In the second section, we overviewed recent progress in TJ modulators with respect to the barrier separating different body compartments.

3.1 Blood-brain barrier

The blood-brain barrier, which comprises endothelial cell sheets with extremely tight junctions, limits the diffusion of hydrophilic molecules between the bloodstream and brain. Many pharmaceutical chemicals developed for the treatment of brain disorders cannot be applied in clinical therapy because they do not pass through the blood-brain barrier. Methods to open or reversibly regulate the blood-brain barrier have been investigated. Blood-brain barrier modulation based on the infection mechanisms of HIV has been proposed. Disruption of TJs occurs in the brains of HIV-infected patients [50-52], and tat protein, which is released from HIV-infected cells, decreases ZO-1 levels at the cell-cell borders in brain microvascular endothelial cells [53]. Tat treatment reduced expression of occludin, ZO-1, and ZO-2 in human brain microvascular endothelial cells via caveolin-1 and Ras signaling. Other HIV-1-derived proteins, gp120 and Nef,

Table 1. Candidates of absorption enhancer.

Target barrier	Candidates
Intestinal barrier	C-CPE
	AT1002
	Ephrin
	Chitosan and its derivatives
	Haemagglutinin
	HAstV-1 capsid
	Spermine
Blood-brain barrier	HIV-1 tat
	Sodium caprate
	Nitric oxide
Nasal barrier	AT1002
	Sperminated gelatin
	FDWITP
Blood-testis barrier	C-type natriuretic peptide domain I of laminin β 3

C-CPE: C-terminal of *Clostridium perfringens* enterotoxin; HAstV-1: Human astrovirus serotype 1; HIV: Human immunodeficiency virus.

can change the expression of TJ proteins *in vitro* [54]. Cocaine [55-56], sodium caprate [57] and nitric oxide [58] also modulate the blood–brain barrier.

3.2 Blood–testis barrier

Disruption of the blood–testis barrier affects spermatogenesis; thus, junctional proteins, such as occludin, ZO-1, and N-cadherin, could be the primary targets for testicular toxicants [59]. Monophthalates (mono-*n*-butyl phthalate and mono-2-ethylhexyl phthalate) were recently shown to disrupt the inter-Sertoli TJs in rat [60]. Phthalates are used as plasticizing and suspension agents in personal care products, plastics, paints, and pesticides. Monophthalates reduced the TJ barrier in Sertoli cells and induced the disappearance of ZO-1 and F-actin from around the cell periphery. The expression of occludin mRNA was also suppressed in a dose-dependent manner. C-type natriuretic peptide is a novel regulator of blood–testis barrier dynamics [61]. C-type natriuretic peptide regulates blood pressure, blood volume, fat metabolism, bone growth, and steroidogenesis in the testis and also reduces the expression of N-cadherin, occludin, and JAM-A [62,63]. Laminin fragments can also modulate the blood–testis barrier [64]. Treatment of primary Sertoli cells with domain I of laminin β 3 caused a dose-dependent reduction in β 1-integrin, occludin and ZO-1 and a decrease in the blood–testis barrier. Domain IV of laminin γ 3 also reduced the expression of β 1-integrin, occludin and JAM-A.

3.3 Epithelial barrier

Intranasal delivery is a convenient, reliable, rapid, and noninvasive delivery approach for low-molecular-weight

compounds, and intranasal absorption enhancers have been developed for improvement of the nasal absorption of therapeutic macromolecules. AT1002, a polypeptide derived from Zot, enhanced not only intestinal absorption, but also nasal absorption of hydrophilic markers, PEG4000 and inulin [65]. Sperminated gelatin is a nasal absorption enhancer of insulin; when intranasally delivered, it decreases the plasma glucose level [66]. Aminated gelatin enhanced absorption of protein drugs through mucosal membranes with negligible mucosal damage [67].

Phage display technology is a powerful method for the selection of peptide ligands [68,69]. Recently, novel TJ modulators were isolated by using a phage-display library [70]. TJ-bound peptides were screened by using confluent monolayer cell sheets that were treated with a calcium chelator, EGTA. The polypeptide FDFWITP was isolated as a TJ binder. FDFWITP and its derivative peptides modulated TJ barriers without cytotoxicity, and these TJ-modulating activities were reversible. Thus, the phage-display system is a promising and powerful tool for developing TJ modulators.

4. Expert opinion

Many TJ-associated integral proteins, including occludin, claudin, tricellulin, ZO-1, ZO-2 and ZO-3, have been identified. These proteins play pivotal roles in the regulation of solute movement via the paracellular route, indicating that TJ modulators can be promising methods to deliver drugs. Studies of claudin-deficient mice initially indicated the possibility of TJ component-based drug delivery. Claudin-1-deficient mice lose their epidermal barrier function against a tracer with a molecular weight of around 600 Da, [15], indicating that claudin-1 modulators can enhance the transdermal absorption of drugs. The transdermal route is an easy, painless, and noninvasive method for drug administration, and the claudin-1 modulators have been the subject of pharmaceutical research. Claudin-5-deficient mice lose their blood–brain barrier [16], and small molecules (< 800 Da) selectively passed across the blood–brain barrier. The claudin-5 modulator will be a candidate for the pharmaceutical therapy of brain diseases. We found that the intestinal absorption-enhancing effects of a claudin-4 modulator were 400-fold more potent than those of a clinically used absorption enhancer [18]. Disruption of occludin or tricellulin increases TJ permeability [12,25,71]. These findings strongly indicate that modulation of TJ is a promising method for drug delivery. Because TJ proteins are poor in antigenicity, it is difficult to develop antibodies against the extracellular domain, resulting in a severe delay in the development of TJ modulators. At this point, there have been two breakthroughs in the development of TJ modulators. The first breakthrough is the determination of the structure of the only known claudin modulator, C-CPE [21]. The second breakthrough is the establishment of an efficient phage-display method to isolate a novel peptide to bind TJ components [22]. We believe that the

development of a claudin modulator by using C-CPE as a prototype will be successful, and that a peptide type of TJ modulator will be prepared in the near future. We are also optimistic about the production of a novel TJ modulator based on fragments of toxins, viruses and natural products. These fragments appear to use a novel mechanism to modulate the TJ barrier, and further analysis of this novel type of TJ modulator may lead to the next generation of TJ modulators (Table 1).

Very recently, Lee and colleagues proposed the lipid-protein hybrid model for TJ that the TJ proteins by themselves, and in combination with the lipids, serve, in addition, essential roles in barrier function, indicating that a lipid modulator can

be a TJ modulator [72]. Glycosylated sphingosine, oxidized lipids and ether lipids were identified as TJ modulators, and the displacement of claudins and occludin from lipid raft was involved in the absorption-enhancing effect of sodium caprate [73,74]. Future investigation of the lipid-protein hybrid model for TJ may be the third breakthrough in the development of TJ modulators.

Declaration of interest

The authors state no conflict of interest and have received no payment in the preparation of this manuscript.

Bibliography

Papers of special note have been highlighted as either of interest (*) or of considerable interest (**) to readers.

- Majumdar S, Duvvuri S, Mitra AK. Membrane transporter/receptor-targeted prodrug design: strategies for human and veterinary drug development. *Adv Drug Deliv Rev* 2004;56:1437-52
- Mizuno N, Niwa T, Yotsumoto Y, Sugiyama Y. Impact of drug transporter studies on drug discovery and development. *Pharmacol Rev* 2003;55:425-61
- Inui KI, Masuda S, Saito H. Cellular and molecular aspects of drug transport in the kidney. *Kidney Int* 2000;58:944-58
- Koepsell H. Organic cation transporters in intestine, kidney, liver, and brain. *Annu Rev Physiol* 1998;60:243-66
- Meijer DK, Hooiveld GJ, Schinkel AH, et al. Transport mechanisms for cationic drugs and proteins in kidney, liver and intestine: implication for drug interactions and cell-specific drug delivery. *Nephrol Dial Transplant* 1999;4(Suppl 14):1-3
- Van Aubel RA, Masereeuw R, Russel FG. Molecular pharmacology of renal organic anion transporters. *Am J Physiol Renal Physiol* 2000;279:F216-32
- Anderson JM, Van Itallie CM. Tight junctions and the molecular basis for regulation of paracellular permeability. *Am J Physiol* 1995;269:G467-75
- Powell DW. Barrier function of epithelia. *Am J Physiol* 1981;241:G275-88
- Furuse M, Hirase T, Itoh M, et al. Occludin: a novel integral membrane protein localizing at tight junctions. *J Cell Biol* 1993;123:1777-88
- The first paper to identify a component of tight junction.**
- Furuse M, Fujita K, Hiiiragi T, et al. Claudin-1 and -2: novel integral membrane proteins localizing at tight junctions with no sequence similarity to occludin. *J Cell Biol* 1998;141:1539-50
- The first paper to identify a functional component of tight junction.**
- Furuse M, Tsukita S. Claudins in occluding junctions of humans and flies. *Trends Cell Biol* 2006;16(4):181-8
- Wong V, Gumbiner BM. A synthetic peptide corresponding to the extracellular domain of occludin perturbs the tight junction permeability barrier. *J Cell Biol* 1997;136(2):399-409
- Van Itallie CM, Anderson JM. Claudins and epithelial paracellular transport. *Annu Rev Physiol* 2006;68:403-29
- Morita K, Furuse M, Fujimoto K, Tsukita S. Claudin multigene family encoding four-transmembrane domain protein components of tight junction strands. *Proc Natl Acad Sci USA* 1999;96:511-6
- Furuse M, Hata M, Furuse K, et al. Claudin-based tight junctions are crucial for the mammalian epidermal barrier: a lesson from claudin-1-deficient mice. *J Cell Biol* 2002;156:1099-111
- Nitta T, Hata M, Gotoh S, et al. Size-selective loosening of the blood-brain barrier in claudin-5-deficient mice. *J Cell Biol* 2003;161:653-60
- Sonoda N, Furuse M, Sasaki H, et al. Clostridium perfringens enterotoxin fragment removes specific claudins from tight junction strands: evidence for direct involvement of claudins in tight junction barrier. *J Cell Biol* 1999;147:195-204
- Shows the first experimental evidence that claudin is responsible for tight junction barrier.**
- Kondoh M, Masuyama A, Takahashi A, et al. A novel strategy for the enhancement of drug absorption using a claudin modulator. *Mol Pharmacol* 2005;67:749-56
- Takahashi A, Komiya E, Kakutani H, et al. Domain mapping of a claudin-4 modulator, the C-terminal region of C-terminal fragment of Clostridium perfringens enterotoxin, by site-directed mutagenesis. *Biochem Pharmacol* 2008;75:1639-48
- Takahashi A, Kondoh M, Masuyama A, et al. Role of C-terminal regions of the C-terminal fragment of Clostridium perfringens enterotoxin in its interaction with claudin-4. *J Control Release* 2005;108:56-62
- Van Itallie CM, Betts L, Smedley JG III, et al. Structure of the claudin-binding domain of Clostridium perfringens enterotoxin. *J Biol Chem* 2008;283:268-74
- The first paper about the structure of C-CPE.**
- Ling J, Liao H, Clark R, et al. Structural constraints for the binding of short peptides to claudin-4 revealed by surface plasmon resonance. *J Biol Chem* 2008;283:30585-95
- Mrsny RJ, Brown GT, Gerner-smidt K, et al. A key claudin extracellular loop domain is crucial for epithelial barrier integrity. *Am J Pathol* 2008;172:905-15
- Furuse M, Sasaki H, Tsukita S. Manner of interaction of heterogeneous claudin species within and between tight junction strands. *J Cell Biol* 1999;147:891-903

Tight junction modulator and drug delivery

25. Wong CH, Mruk DD, Lee WM, Cheng CY. Targeted and reversible disruption of the blood-testis barrier by an FSH mutant-occludin peptide conjugate. *FASEB J* 2007;21:438-48
26. Madeley CR, Cosgrove BP. Letter: 28 nm particles in faeces in infantile gastroenteritis. *Lancet* 1975;2:451-2
27. Moser LA, Carter M, Schultz Cherry S. Astrovirus increases epithelial barrier permeability independently of viral replication. *J Virol* 2007;81:11937-45
28. Monteleone G, Fina D, Caruso R, Pallone F. New mediators of immunity and inflammation in inflammatory bowel disease. *Curr Opin Gastroenterol* 2006;22(4):361-4
29. Al Sadi RM, Ma TY. IL-1beta causes an increase in intestinal epithelial tight junction permeability. *J Immunol* 2007;178:4641-9
30. Al Sadi R, Ye D, Dokladny K, Ma TY. Mechanism of IL-1beta-induced increase in intestinal epithelial tight junction permeability. *J Immunol* 2008;180:5653-61
31. Tanaka M, Kamata R, Sakai R. EphA2 phosphorylates the cytoplasmic tail of Claudin-4 and mediates paracellular permeability. *J Biol Chem* 2005;280:42375-82
32. Larson J, Schomberg S, Schroeder W, Carpenter TC. Endothelial EphA receptor stimulation increases lung vascular permeability. *Am J Physiol* 2008;295:L431-9
33. Aasheim HC, Pedeutour F, Grosgeorge J, Logtenberg T. Cloning, chromosomal mapping, and tissue expression of the gene encoding the human Eph-family kinase ligand ephrin-A2. *Biochem Biophys Res Commun* 1998;252:378-82
34. Salama NN, Eddington ND, Fasano A. Tight junction modulation and its relationship to drug delivery. *Adv Drug Deliv Rev* 2006;58:15-28
35. Fasano A, Uzzau S. Modulation of intestinal tight junctions by Zonula occludens toxin permits enteral administration of insulin and other macromolecules in an animal model. *J Clin Invest* 1997;99:1158-64
36. Fasano A, Uzzau S, Fiore C, Margaretten K. The enterotoxic effect of Zonula occludens toxin on rabbit small intestine involves the paracellular pathway. *Gastroenterology* 1997;112:839-46
37. Fasano A, Baudry B, Pumplun DW, et al. Vibrio cholerae produces a second enterotoxin, which affects intestinal tight junctions. *Proc Natl Acad Sci USA* 1991;88:5242-6
38. Fasano A, Fiorentini C, Donelli G, et al. Zonula occludens toxin modulates tight junctions through protein kinase C-dependent actin reorganization, in vitro. *J Clin Invest* 1995;96:710-20
39. Cox DS, Gao H, Raje S, et al. Enhancing the permeation of marker compounds and enaminone anticonvulsants across Caco-2 monolayers by modulating tight junctions using zonula occludens toxin. *Eur J Pharm Biopharm* 2001;52:145-50
40. Cox DS, Raje S, Gao H, et al. Enhanced permeability of molecular weight markers and poorly bioavailable compounds across Caco-2 cell monolayers using the absorption enhancer, zonula occludens toxin. *Pharm Res* 2002;19:1680-8
41. Lee A, White N, Van der Walle CF. The intestinal zonula occludens toxin (ZOT) receptor recognises non-native ZOT conformers and localises to the intercellular contacts. *FEBS Lett* 2003;555:638-42
42. Di Pierro M, Lu R, Uzzau S, et al. Zonula occludens toxin structure-function analysis. Identification of the fragment biologically active on tight junctions and of the zonulin receptor binding domain. *J Biol Chem* 2001;276:19160-5
43. Song KH, Fasano A, Eddington ND. Effect of the six-mer synthetic peptide (AT1002) fragment of zonula occludens toxin on the intestinal absorption of cyclosporin A. *Int J Pharm* 2008;351:8-14
44. Thanou M, Verhoef JC, Junginger HE. Chitosan and its derivatives as intestinal absorption enhancers. *Adv Drug Deliv Rev* 2001;50(Suppl 1):S91-101
45. Kotze AF, Luessen HL, De leeuw BJ, et al. Comparison of the effect of different chitosan salts and N-trimethyl chitosan chloride on the permeability of intestinal epithelial cells (Caco-2). *J Control Release* 1998;51:35-46
46. Lee DW, Baney RH. Oligochitosan derivatives bearing electron-deficient aromatic rings for adsorption of amitriptyline: implications for drug detoxification. *Biomacromolecules* 2004;5:1310-5
47. Smith JM, Dornish M, Wood EJ. Involvement of protein kinase C in chitosan glutamate-mediated tight junction disruption. *Biomaterials* 2005;26:3269-76
48. Lin YH, Mi FL, Chen CT, et al. Preparation and characterization of nanoparticles shelled with chitosan for oral insulin delivery. *Biomacromolecules* 2007;8:146-52
49. Matsumura T, Jin Y, Kabumoto Y, et al. The HA proteins of Botulinum toxin disrupt intestinal epithelial intercellular junctions to increase toxin absorption. *Cell Microbiol* 2008;10:355-64
50. Boven LA, Middel J, Verhoef J, et al. Monocyte infiltration is highly associated with loss of the tight junction protein zonula occludens in HIV-1-associated dementia. *Neuropathol Appl Neurobiol* 2000;26:356-60
51. Dallasta LM, Pizarov LA, Esplen JE, et al. Blood-brain barrier tight junction disruption in human immunodeficiency virus-1 encephalitis. *Am J Pathol* 1999;155:1915-27
52. Persidsky Y, Heilman D, Haorah J, et al. Rho-mediated regulation of tight junctions during monocyte migration across the blood-brain barrier in HIV-1 encephalitis (HIVE). *Blood* 2006;107:4770-80
53. Zhong Y, Smart EJ, Weksler B, et al. Caveolin-1 regulates human immunodeficiency virus-1 Tat-induced alterations of tight junction protein expression via modulation of the Ras signaling. *J Neurosci* 2008;28:7788-96
54. Annunziata P. Blood-brain barrier changes during invasion of the central nervous system by HIV-1. Old and new insights into the mechanism. *J Neurol* 2003;250:901-6
55. Dhillon NK, Peng F, Bokhari S, et al. Cocaine-mediated alteration in tight junction protein expression and modulation of CCL2/CCR2 axis across the blood-brain barrier: implications for HIV-dementia. *J Neuroimmune Pharmacol* 2008;3:52-6
56. Zhang L, Looney D, Taub D, et al. Cocaine opens the blood-brain barrier to HIV-1 invasion. *J Neurovirol* 1998;4:619-26

57. Preston E, Slinn J, Vinokourov I, Stanimirovic D. Graded reversible opening of the rat blood-brain barrier by intracarotid infusion of sodium caprate. *J Neurosci Methods* 2008;168:443-9
58. Yamauchi A, Dohgu S, Nishioku T, et al. An inhibitory role of nitric oxide in the dynamic regulation of the blood-brain barrier function. *Cell Mol Neurobiol* 2007;27:263-70
59. Fiorini C, Tilloy Ellul A, Chevalier S, et al. Sertoli cell junctional proteins as early targets for different classes of reproductive toxicants. *Reprod Toxicol* 2004;18:413-21
60. Zhang YH, Lin L, Liu ZW, et al. Disruption effects of monophthalate exposures on inter-Sertoli tight junction in a two-compartment culture model. *Environ Toxicol* 2008;23:302-8
61. Xia W, Mruk DD, Cheng CY. C-type natriuretic peptide regulates blood-testis barrier dynamics in adult rat testes. *Proc Natl Acad Sci USA* 2007;104:3841-6
62. Potter LR, Abbey Hosch S, Dickey DM. Natriuretic peptides, their receptors, and cyclic guanosine monophosphate-dependent signaling functions. *Endocr Rev* 2006;27:47-72
63. Gnessi L, Fabbri A, Spera G. Gonadal peptides as mediators of development and functional control of the testis: an integrated system with hormones and local environment. *Endocr Rev* 1997;18:541-609
64. Yan HH, Mruk DD, Wong EW, et al. An autocrine axis in the testis that coordinates spermiation and blood-testis barrier restructuring during spermatogenesis. *Proc Natl Acad Sci USA* 2008;105:8950-5
65. Song KH, Fasano A, Eddington ND. Enhanced nasal absorption of hydrophilic markers after dosing with AT1002, a tight junction modulator. *Eur J Pharm Biopharm* 2008;69:231-7
66. Seki T, Kanbayashi H, Chono S, et al. Effects of a sperminated gelatin on the nasal absorption of insulin. *Int J Pharm* 2007;338:213-8
67. Seki T, Kanbayashi H, Nagao T, et al. Effect of aminated gelatin on the nasal absorption of insulin in rats. *Biol Pharm Bull* 2005;28:510-4
68. Lowman HB. Bacteriophage display and discovery of peptide leads for drug development. *Annu Rev Biophys Biomol Struct* 1997;26:401-24
69. Sidhu SS. Phage display in pharmaceutical biotechnology. *Curr Opin Biotechnol* 2000;11:610-6
70. Herman RE, Makienko EG, Prieve MG, et al. Phage display screening of epithelial cell monolayers treated with EGTA: identification of peptide FDFWITP that modulates tight junction activity. *J Biomol Screen* 2007;12:1092-101
- Provides a novel strategy for development of tight junction modulator using a phage display library.
71. Ikenouchi J, Furuse M, Furuse K, et al. Tricellulin constitutes a novel barrier at tricellular contacts of epithelial cells. *J Cell Biol* 2005;171:939-45
72. Lee DBN, Jangotchian N, Allen SG, et al. A lipid-protein hybrid model for tight junction. *Am J Physiol* 2008;295:F1601-12
73. Chen-Quay SC, Eting KT, Li awa, et al. Identification of tight junction modulating lipids. *J Pharm Sci* 2009;98:606-19
74. Sugibayashi K, Onuki Y, Takayama K. Displacement of tight junction proteins from detergent-resistant membrane domains by treatment with sodium caprate. *Eur J Pharm Sci* 2009;36:246-53

Affiliation

Koji Matsuhisa, Masuo Kondoh[†]
 Azusa Takahashi & Kiyohito Yagi
[†]Author for correspondence
 Osaka University,
 Graduate School of Pharmaceutical Sciences,
 Department of Bio Functional Molecular Chemistry,
 Suita, Osaka 565-0871, Japan
 Tel: +81 6 6879 8196; Fax: +81 6 6879 8199;
 E-mail: masuo@phs.osaka-u.ac.jp

A Novel Tumor-Targeted Therapy Using a Claudin-4-Targeting Molecule

Rie Saeki, Masuo Kondoh, Hideki Kakutani, Shin-ichi Tsunoda, Yasuhiro Mochizuki, Takao Hamakubo, Yasuo Tsutsumi, Yasuhiko Horiguchi, and Kiyohito Yagi

Laboratories of Bio-Functional Molecular Chemistry (R.S., M.K., H.K., K.Y.) and Toxicology (Y.T.), Graduate School of Pharmaceutical Sciences and Department of Molecular Bacteriology, Research Institute for Microbial Diseases (Y.H.), Osaka University, Suita, Japan; Laboratory of Pharmaceutical Proteomics, Division of Biomedical Research, National Institute of Biomedical Innovation, Ibaraki, Japan (S.T., Y.T.); and Department of Molecular Biology and Medicine, Research Center for Advanced Science and Technology, the University of Tokyo, Meguro, Japan (Y.M., T.H.)

Received June 8, 2009; accepted July 28, 2009

ABSTRACT

Carcinogenesis is often accompanied by dysfunctional tight junction (TJs), resulting in the loss of cellular polarity. Claudin, a tetra-transmembrane protein, plays a pivotal role in the barrier and fence functions of TJs. Claudin-4 is deregulated in various cancers, including breast, prostate, ovarian, and gastric cancer. Claudin-4 may be a promising target molecule for tumor therapy, but the claudin-targeting strategy has never been fully developed. In the present study, we prepared a claudin-4-targeting molecule by fusion of the C-terminal fragment of *Clostridium perfringens* enterotoxin (C-CPE) with the protein synthesis inhibitory factor (PSIF) derived from *Pseudomonas aeruginosa* exotoxin. PSIF was not cytotoxic to claudin-4-expressing cells, whereas C-CPE-

PSIF was cytotoxic. Cells that express claudin-1, -2, and -5 were less sensitive to C-CPE-PSIF. Pretreatment of the cells with C-CPE attenuated C-CPE-PSIF-induced cytotoxicity, and mutation of C-CPE in the claudin-4-binding residues attenuated the cytotoxicity of C-CPE-PSIF. TJ-undeveloped cells were more sensitive to C-CPE-PSIF than TJ-developed cells. It is noteworthy that polarized epithelial cells are sensitive to C-CPE-PSIF applied to the basal side, whereas the cells were less sensitive to C-CPE-PSIF applied to the apical side. Intratumoral injection of C-CPE-PSIF reduced tumor growth. This is the first report to indicate that a claudin-4-targeting strategy may be a promising method to overcome the malignant tumors.

The majority of lethal cancers are derived from epithelial tissues (Jemal et al., 2008), and various therapeutic strategies against epithelium-derived cancers have been developed. Targeted therapies that use differences between normal cells and cancer cells are promising antitumor therapies, and cellular surface proteins displayed on cancer cells are often targeted. Genetically modified toxins that target the

surface proteins have emerged as a promising treatment strategy for refractory cancers (Michl and Gress, 2004). However, malignant tumors are still a major cause of death; more than 7 million people worldwide die from cancer each year (Dunham, 2007). Thus, the development of a novel strategy for cancer-targeting therapy is needed.

A defining feature of epithelial cells is cellular polarity. Epithelial cells have tight junctions (TJs) on the membrane between adjacent cells. TJs seal the intercellular space between adjacent cells and regulate solute movement across epithelial cell sheets (Anderson and Van Itallie, 1995). In addition, TJs form the fence of the membrane that prevents lateral diffusion of membrane proteins and lipids, thereby maintaining the differential composition of the apical and basolateral domains. TJs also seem to be involved in the regulation of proliferation, differentiation, and other cellular

This work was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology, Japan; a Health and Labor Sciences Research Grants from the Ministry of Health, Labor and Welfare of Japan; Takeda Science Foundation; a Suzuken Memorial Foundation; and a Grant from Kansai Biomedical Cluster project in Saito, which is promoted by the Knowledge Cluster Initiative of the Ministry of Education, Culture, Sports, Science and Technology, Japan.

Article, publication date, and citation information can be found at <http://molpharm.aspetjournals.org>.
doi:10.1124/mol.109.058412.

ABBREVIATIONS: TJ, tight junction; C-CPE, C-terminal fragment of *Clostridium perfringens* enterotoxin from 194 to 319 amino acids; PSIF, protein synthesis inhibitory factor derived from *P. aeruginosa* exotoxin; CPE, *Clostridium perfringens* enterotoxin; PE, *P. aeruginosa* exotoxin; C-CPE-PSIF, C-terminal fragment of *Clostridium perfringens* enterotoxin-fused protein synthesis inhibitory factor derived from *P. aeruginosa* exotoxin; PAGE, polyacrylamide gel electrophoresis; BV, budded baculovirus; Ab, antibody; TER, transepithelial electric resistance; FBS, fetal bovine serum; MCS, multiple cloning sites; PCR, polymerase chain reaction; PBS, phosphate-buffered saline; BSA, bovine serum albumin; TBS, Tris-buffered saline; LDH, lactate dehydrogenase; aa, amino acid.

functions (Mitic and Anderson, 1998; Vermeer et al., 2003). Malignant tumor cells frequently exhibit abnormal TJ function, followed by the loss of cellular polarity and intercellular contact that commonly occurs in advanced tumors and early stages of carcinogenesis (Wodarz and Näthke, 2007). These findings indicate that TJ proteins, which are barely accessible in well structured normal epithelia but are exposed on malignant tumor cells, may be promising candidates for targeted therapy.

Freeze-fracture replica electron microscopy analysis reveals that TJs appear as a series of continuous, anastomotic, and intramembranous particle strands or fibrils (Tsukita and Furuse, 1999). The TJ strands consist of integral membrane proteins polymerizing linearly within a lipid layer of the cell membrane. Claudin, a ~24-kDa four-transmembrane protein that forms a family containing 24 members, is a crucial component of the TJ strand in forming the TJ fence (Furuse and Tsukita, 2006). The expression of claudins is altered in several cancers. In particular, claudin-4 is frequently up-regulated in breast, prostate, pancreatic, and ovarian cancers (Morin, 2005). Thus, the claudins are potential targets for antitumor therapy. Although antibodies to the extracellular region of claudins are promising molecules for tumor-targeted therapy, these antibodies have not yet been successfully prepared.

Clostridium perfringens enterotoxin (CPE) is a single polypeptide of 35-kDa that can cause food poisoning in humans. Functional domains of CPE are classified into the N-terminal cytotoxic region and the C-terminal receptor-binding region (Hanna et al., 1992). CPE binds to its receptor, claudin-4, followed by the formation of the complex on the membrane leading to cell damage (Katahira et al., 1997; Paperna et al., 1998; McClane and Chakrabarti, 2004). Prostate adenocarcinoma cells expressing claudin-4 are sensitive to CPE-mediated cytotoxicity (Long et al., 2001). Breast, ovarian, and pancreatic cancer cells expressing claudin-4 are also sensitive to CPE treatment (Michl et al., 2001; Kominsky et al., 2004; Santin et al., 2005). These findings suggest that the receptor-binding region of CPE is useful for targeting claudin-4-expressing cancer cells.

Pseudomonas aeruginosa exotoxin A (PE) is widely used in cancer-targeting therapy. PE binds to the cell surface and is internalized via an endocytotic pathway, followed by escape of the PE fragment (protein synthesis inhibitory factor, PSIF) from the endosome into the cytosol. The released PSIF inhibits protein synthesis by the inhibition of elongation factor 2 (Ogata et al., 1990). PSIF lacks the receptor binding domain of PE, and fusion of the ligand of tumor antigen with PSIF is a promising strategy for cancer-targeting therapy. For example, the fusion protein of PSIF with anti-interleukin-2 antibody has been used in clinical therapy (Kreitman and Pastan, 2006). In the present study, we genetically prepared the claudin-4-targeting molecule (C-CPE-PSIF) containing the claudin-4-binding region of CPE and PSIF, and we found that C-CPE-PSIF has in vivo antitumor activity.

Materials and Methods

Cell Culture. Mouse fibroblast cell line L cells and mouse claudin-expressing L cells (claudin-1/L, -2/L, -4/L, and -5/L cells), kindly provided by Dr. S. Tsukita (Kyoto University, Kyoto, Japan), were cultured in modified Eagle's medium supplemented with 10% fetal

bovine serum (FBS). Human hepatocarcinoma cell lines HepG2 cell and SK-HEP-1 cells were maintained in Dulbecco's modified Eagle's medium containing 10% FBS. The human intestinal cell line Caco-2 was maintained in Dulbecco's modified Eagle's medium containing 10% FBS and 1% nonessential amino acids. The murine mammary carcinoma cell line 4T1 was maintained in Dulbecco's modified Eagle's medium containing 10% FBS and 10 mM HEPES. The cells were maintained in a 5% CO₂ atmosphere at 37°C.

Preparation of C-CPE-PSIF. We prepared plasmids containing the C-terminal CPE-fused PSIF. In brief, we generated pET16b-MCS. Double-stranded oligonucleotides of MCS were prepared by annealing (heating at 95°C for 5 min and chilling at room temperature for 60 min) single-strand oligonucleotides, a forward oligonucleotide (5'-TCGAAGGTACCCGGGACTAGTTAATTAAG-3', XhoI binding site is underlined) and a reverse oligonucleotide (5'-TCGAACTTTTATAACTAGTCCGGGTCCAT-3', XhoI binding site is underlined). The annealed oligonucleotides were subcloned into the pET16b vector (Novagen, Darmstadt, Germany), resulting in pET16b-MCS. PSIF was amplified by polymerase chain reaction (PCR) using pPBV-PE40 as a template, a forward primer (5'-GATGATCTGAGCGGCCCAACCCGAGGGCGGCAG-3', NotI site is underlined), and a reverse primer (5'-TCCAGATCTTTACAGTTCGCTCTTTCTTCAGTCCCTC-3', BglII site is underlined). The resulting PCR fragments were subcloned into NotI/BamHI-digested pET16b-MCS to create pET-PSIF, and the sequence was confirmed. C-CPE and C-CPE_{Y306A/L315A} were amplified by PCR with pETH₁₀PER as a template. The common forward primer is (5'-GGAATTCATATGGATATAGAAAAAGAAATCCTTGATTTAGCTGCT-3', SpeI site is underlined), and the reverse primer for C-CPE is (5'-GGACTAGTAAATTTTTGAAATAATATTGAATAAGGGTAATTTCCACTATATG-3', NdeI site is underlined) or C-CPE_{Y306A/L315A} (5'-GGACTAGTAAATTTTTGCTATTGAATAAGGGTAATTTCCACTAGCTGATGAATTAGCTTTTCATTAC-3', NdeI site is underlined). The resulting PCR products were subcloned into SpeI/NdeI-digested pET-PSIF to create pET-C-CPEs-PSIF, and the sequence was confirmed. The plasmids, pET-PSIF and pET-C-CPEs-PSIF, were transfected into *Escherichia coli* BL21 (DE3) strains (Novagen), and the production of PSIF and C-CPE-PSIF was induced by the addition of 0.25 mM isopropyl-D-thiogalactopyranoside. The cells were harvested and then lysed in buffer A (10 mM Tris-HCl, pH 8.0, 400 mM NaCl, 5 mM MgCl₂, 0.1 mM phenylmethylsulfonyl fluoride, 1 mM 2-mercaptoethanol, and 10% glycerol). The lysates were centrifuged, and the resultant supernatant was applied to HiTrap Chelating HP (GE Healthcare, Chalfont St. Giles, Buckinghamshire, UK). The proteins were eluted by imidazole in buffer A. The buffer was exchanged with phosphate-buffered saline (PBS) by using a PD-10 column (GE Healthcare), and the purified protein was stored at -80°C until use. Protein was quantified by using a BCA protein assay kit (Pierce Chemical, Rockford, IL) with bovine serum albumin (BSA) as a standard.

Cytotoxic Activity. In the cytotoxic assay, L-cells were seeded onto a 96-well culture dish at 10⁴ cells/well. After 24 h of the culture, the cells were treated with PSIF or C-CPE-PSIF for 24 h at the indicated concentration. HepG2, SK-HEP-1, and 4T1 cells were seeded onto a 96-well culture dish at 10⁴ cells/well. After 24 h of culture, the cells were treated with PSIF or C-CPE-PSIF for 48 h at the indicated concentration. In a pre-confluent assay, Caco-2 cells were seeded onto a 96-well culture dish at 10⁴ cells/well. After 24 h of culture, the cells were treated with PSIF or C-CPE-PSIF for 48 h at the indicated concentration. In a postconfluent assay, Caco-2 cells were cultured in a 96-well culture dish for an additional 3 days after reaching a confluent condition. Then, the cells were treated with PSIF or C-CPE-PSIF for 48 h at the indicated concentration. The cytotoxicity was determined by a WST-8 kit according to the manufacturer's instructions (Nacalai Tesque, Kyoto, Japan).

Immunoblot Analysis. Cells were lysed in lysis buffer [20 mM Tris, pH 7.4, 150 mM NaCl, 1% Triton X-100, protease inhibitor cocktail (Sigma-Aldrich, St. Louis, MO), and phosphatase inhibitor cocktail (Nacalai)]. The cell lysates were subjected to SDS-polyacrylamide gel electrophoresis (PAGE). The separated proteins were

transferred onto a polyvinylidene difluoride membrane, followed by immunoblotting with anti-claudin-1, -2, -4, and -5 (Zymed Laboratories, South San Francisco, CA), anti-His-tag (Novagen), or anti- β -actin Ab (Sigma-Aldrich). The immunoreactive band was visualized by chemiluminescence reagents (GE Healthcare).

Competition Assay. After pretreatment of claudin-4/L (CL4/L) cells with C-CPE or BSA for 2 h, the cells were incubated with C-CPE-PSIF for 24 h. Then the cell viability was assayed by the cell counting kit, as described above.

Preparation of the Claudin-Displaying Budded Baculovirus. The cDNAs for mouse claudin-1 and claudin-4 were amplified by PCR from pGTCL-1 and pGTCL-4 (kindly provided by Dr. M. Furuse, Kobe University, Kobe, Japan). The DNA fragments were subcloned into the baculoviral transfer vector pFastBac1 (Invitrogen, Carlsbad, CA). Recombinant baculoviruses were generated by using the Bac-to-Bac system (Invitrogen) according to the manufacturer's instructions.

Sf9 cells were cultured in Grace's Insect medium (Invitrogen) containing 10% FBS at 27°C. Sf9 cells were infected with the recombinant baculovirus. Seventy-two hours after infection, the budded baculovirus (BV) fraction was isolated from the culture supernatant of infected Sf9 cells by centrifugation at 40,000g for 25 min. The pellets of the BV fraction were suspended in Tris-buffered saline (TBS) containing protease inhibitor cocktail and then stored at 4°C. The expression of claudins in the BV fraction was confirmed by SDS-PAGE and immunoblot analysis.

Enzyme-Linked Immunosorbent Assay. The BV-displaying claudins were diluted with TBS and adsorbed to the wells of 96-well immunoplates (Nunc, Roskilde, Denmark) overnight at 4°C. The wells were washed with PBS and blocked with TBS containing 1.6% BlockAce (Dainippon Sumitomo Pharma, Osaka, Japan) for 2 h at room temperature. C-CPEs-PSIF was added to the well and incubated for an additional 2 h at room temperature. The wells were washed with PBS and incubated with anti-His-tag antibody for 2 h at room temperature. The immunoreactive proteins were detected by horseradish peroxidase-labeled secondary antibody by using 3,3',5,5'-tetramethyl benzene as a substrate. The reaction was terminated by the addition of 0.2 M H₂SO₄, and the immunoreactive proteins were measured at 450 nm.

Measurement of Transepithelial Electric Resistance. Confluent monolayers of Caco-2 cells were grown in Transwell chambers (Corning Life Sciences, Lowell, MA). The formation of tight junction barriers in Caco-2 monolayers was monitored by measuring transepithelial electric resistance (TER) with a Millicell-ERS epithelial volt-ohmmeter (Millipore Corporation, Billerica, MA). After 7 to 10 days of culture, the TER values reached a plateau. Then, C-CPE-PSIF was added to the apical or basolateral compartment of the Transwell chamber, and the TER values were measured for 48 h. The TER values were normalized by the area of the Caco-2 monolayer. The background TER of a blank Transwell chamber was subtracted from the TER of cell monolayers.

L-Lactate Dehydrogenase Release Assay. The release of lactate dehydrogenase (LDH) from cells was analyzed by using a Cytotox-96 NonRadioactive Cytotoxicity Assay kit (Promega, Madison, WI) according to the manufacturer's instructions. LDH release was calculated by using the following equation: % maximal LDH release = LDH in the cultured medium/total LDH in the culture dish.

Reverse Transcriptase-Polymerase Chain Reaction. Total RNA was isolated with a High Pure RNA Isolation Kit (Roche, Basel, Switzerland) according to the manufacturer's instructions, and the RNA was treated with RNase-free DNase. Then, 200 ng of RNA was reverse-transcribed with dT adaptor by using a Takara RNA PCR Kit (Takara Inc., Shiga, Japan), according to the manufacturer's instructions. The resulting cDNA was amplified by Ex Taq DNA polymerase (Takara). The PCR conditions were as follows: 96°C for 2 min, 30 cycles of 96°C for 45 s, 55°C for 60 s, and 72°C for 30 s. Claudin-4 and β -actin mRNA were detected by using the following primers: forward primer, 5'-ATGGCGTCTATGGGACTACAGGTCC-

3'; reverse primer, 5'-CCGAGTAGGGCTGTGCTTGCTAC-3'; and forward primer, 5'-TAGATGGGCACAGTGTGGG-3'; reverse primer, 5'-GGCGTGATGGTGGGCATGG-3', respectively.

In Vivo Antitumor Activity. Female BALB/c mice (7–8 weeks old) were obtained from Shimizu Laboratory Supplies Co., Ltd. (Kyoto, Japan). The mice were housed in an environmentally controlled room at 23 ± 1.5°C with a 12-h light/dark cycle. 4T1 cells (2 × 10⁶) in 50 μ l of PBS were injected subcutaneously into the right flank on day 0. Tumor size was determined by measuring two diameters, and the tumor volume was calculated by the following equation: tumor volume = $a \times b \times b/2$, where a represents the maximum tumor diameter and b represents the minimum tumor diameter. After the inoculation of the cells, intratumoral injection of PBS, C-CPE, C-CPE_{Y306A/L315A}-PSIF, or C-CPE-PSIF was performed on days 2, 4, 7, 9, 11, and 14.

Statistical Analysis. Data were analyzed by using two-way analysis of variance followed by the Student's t test. The statistical significance for all comparisons was set at $p < 0.05$.

Results

Preparation of C-CPE-PSIF. Overexpression of claudin-4 is frequently observed in malignant tumors, indicating that the claudins are promising proteins for cancer-targeting therapy (Morin, 2005). We previously developed a novel drug delivery system by using a claudin-4-targeting molecule, the C-terminal of CPE corresponding to 184 to 319 aa (C-CPE₁₈₄₋₃₁₉) (Kondoh et al., 2005; Ebihara et al., 2006). Because of its poor solubility (less than 0.3 mg/ml), it is difficult to use C-CPE₁₈₄₋₃₁₉ in pharmaceutical therapy. Van Itallie et al. (2008) reported that the deletion of the N-terminal 10 aa of C-CPE₁₈₄₋₃₁₉, resulting in C-CPE₁₉₄₋₃₁₉, maintains the claudin binding activity of C-CPE₁₈₄₋₃₁₉ and has high solubility (more than 10 mg/ml) (Van Itallie et al., 2008). We prepared claudin-targeting antitumor agents by genetic fusion of C-CPE₁₉₄₋₃₁₉ and PSIF (C-CPE-PSIF) as shown in Fig. 1A. Here, C-CPE refers to C-CPE₁₉₄₋₃₁₉, PSIF and C-CPE-PSIF were produced by *E. coli* and were purified by affinity chromatography by using a histidine tag. Purification of the proteins was confirmed by SDS-PAGE or immunoblotting (Fig. 1B). The molecular size as determined by SDS-PAGE was identical with the predicted size (C-CPE-PSIF, 60 kDa).

Characterization of C-CPE-PSIF. To evaluate the binding of C-CPE-PSIF to claudins, we used a BV display system. In this system, exogenous proteins are expressed on the surface of the virus particle retaining its function (Sakihama et al., 2008). CPE did not bind to claudin-1 (Fujita et al., 2000). We generated claudin-1-BV and claudin-4-BV and confirmed the claudin expression on BV by Western blotting (data not shown). After the claudin-BV was adsorbed on wells, C-CPE, C-CPE-PSIF, and PSIF proteins were added. We detected the binding of proteins to BV by using anti-histidine-tag antibody. As shown in Fig. 2A, C-CPE specifically interacted with claudin-4-BV but not with wild-type or claudin-1-BV. C-CPE with a mutation on the claudin-4-binding region did not interact with claudin-4-BV (data not shown). These data suggest that fusion with C-CPE provides claudin-4-binding activity to PSIF.

To assess whether C-CPE-PSIF is cytotoxic to claudin-4-expressing cells, we treated claudin-4/L cells with PSIF or C-CPE-PSIF for 24 h. C-CPE-PSIF dose-dependently caused cell death, reaching >90% cell death at 10 ng/ml (Fig. 2B). In contrast, PSIF was not cytotoxic, even at 20 ng/ml. We performed a competition assay to determine whether C-CPE-

PSIF binds to claudin-4 via C-CPE. When the cells were pretreated with BSA or C-CPE before C-CPE-PSIF treatment, C-CPE dose-dependently attenuated the cytotoxic activity of C-CPE-PSIF (Fig. 2C). BSA did not affect the activity even by pretreatment at 10 $\mu\text{g/ml}$, at which C-CPE completely attenuated the cytotoxicity of C-CPE-PSIF. These results suggest that C-CPE-PSIF interacts with claudin-4 via the C-CPE domain.

The claudin family contains 24 members (Furuse and Tsukita, 2006). CPE containing C-CPE as a receptor binding domain bound to claudin-3, -4, -6, and -9 but did not bind to claudin-1, -2, or -5 (Fujita et al., 2000). To confirm the claudin-specificity of C-CPE-PSIF, we investigated the cytotoxic activity of C-CPE-PSIF in L cells expressing claudin-1, -2, -4, or -5 (Fig. 2D). As shown in Fig. 2E, claudin-1/L, -2/L and -5/L cells were less sensitive to C-CPE-PSIF than claudin-4/L cells. Taken together, these results indicate that C-CPE-PSIF may be a claudin-4-targeting cytotoxic agent.

Specificity of C-CPE-PSIF on Cytotoxicity. As mentioned above, C-CPE-PSIF is toxic to exogenous claudin-4-expressing L cells. To examine the specificity of C-CPE-PSIF in the cell lines, we checked the expression of claudin-4 in some human cell lines by immunoblotting and selected two human hepatocarcinoma cell lines: claudin-4-positive HepG2 cells and claudin-4-negative SK-HEP-1 cells (Fig. 3A). Human breast cancer cell line MCF-7 cells were used as a positive control (Ebihara et al., 2006). C-CPE-PSIF was toxic to HepG2 cells and MCF-7 cells, reaching 89 and 53% cell death at 1 and 100 ng/ml, respectively. In contrast, C-CPE-PSIF was not toxic to SK-HEP-1 cells, even at 100 ng/ml (Fig. 3B). Thus, C-CPE-PSIF may have specific toxicity to claudin-4-expressing cells.

Claudin-4 is expressed in various tissues, such as lung,

intestine, liver, and kidney. Most claudins in normal cells would be contained in TJ complexes, whereas the localization of claudins is deregulated in some cancers (Morin, 2005; Kominsky, 2006). C-CPE-PSIF may recognize the deregulated localization of claudin-4, resulting in less toxicity to normal cells. We next examined the effects of C-CPE-PSIF on the human colon carcinoma cell line Caco-2, which expresses claudin-4. Caco-2 cells form a polarized cell monolayer with well developed TJs when they reach confluence, and they are frequently used as a model of polarized cells (Meunier et al., 1995). The claudin-4 protein levels in the confluent culture were greater than the levels in the subconfluent culture (Fig. 3C). As shown in Fig. 3D, C-CPE-PSIF was toxic in the preconfluent cells with fewer TJs (47% cell death at 5 ng/ml). In contrast, the postconfluent cells with well developed TJs were less sensitive to C-CPE-PSIF than the preconfluent cells, and treatment of the cells with C-CPE-PSIF resulted in 40% cell death, even at 200 ng/ml (Fig. 3D).

Early events in epithelial carcinogenesis are the deregulation of cellular polarity and the loss of TJ structures (Wodarz and Näthke, 2007). Next, we examined whether C-CPE-PSIF sensitivity is affected by the cellular polarity in Caco-2 monolayer cell sheets grown on the membrane in Transwell chambers. The Caco-2 monolayer cells exhibit a well differentiated brush border containing TJs on the apical surface, and they are frequently used as an epithelial cell sheet model (Meunier et al., 1995). After the addition of C-CPE-PSIF in the apical or basolateral compartment of the Transwell chamber, we checked the TJ barrier function of the cell sheets by measuring the TER. When C-CPE-PSIF was added to the apical compartment, the TER was not affected for 48 h. In contrast, the addition of C-CPE-PSIF to the basolateral compartment caused a significant reduction in the TER in a dose-dependent manner (Fig. 3E). Furthermore, we found that the addition of C-CPE-PSIF to the basolateral compartment, but not the apical compartment, increased the amount of released LDH, a marker of cytotoxicity (Fig. 3F). Thus, C-CPE-PSIF may have specificity to the cellular polarity.

In Vivo Antitumor Activity of C-CPE-PSIF. We preliminarily investigated the expression of claudin-4 and tumor formation in conventional mice, and we selected a mouse breast cancer cell line, 4T1. Reverse transcription-polymerase chain reaction and immunoblotting analysis revealed the expression of claudin-4 in 4T1 cells (Fig. 4A). To clarify the antitumor activity of a claudin-4-targeting molecule, C-CPE-PSIF, we prepared a fusion protein of PSIF with mutant C-CPE, in which two residues critical for the interaction between C-CPE and claudin-4 (Tyr306 and Leu315) were changed to alanines (Takahashi et al., 2008), resulting in C-CPE_{Y306A/L315A}-PSIF (Fig. 4B). C-CPE_{Y306A/L315A}-PSIF lost the claudin-4-binding activity and was not toxic to CL4/L cells (Fig. 4, C and D). C-CPE-PSIF had dose-dependent cytotoxicity in 4T1 cells, reaching 63% cell death at 100 ng/ml. In contrast, C-CPE_{Y306A/L315A}-PSIF did not show any cytotoxicity, even at 500 ng/ml, indicating that the cytotoxicity of C-CPE-PSIF in 4T1 cells may be mediated by its binding to claudin-4 (Fig. 4E).

Next, to investigate the in vivo antitumor activity of C-CPE-PSIF, 4T1 cells (2×10^6 cells) were inoculated into the right flank of mice on day 0. On day 2, the volume of the tumor exceeded 17 mm³. Vehicle, C-CPE, C-CPE-PSIF, or C-CPE_{Y306A/L315A}-PSIF was intratumorally injected on days




Fig. 1. Preparation of C-CPE-PSIF. A, schematic structure of C-CPE-PSIF. C-CPE-PSIF is a fusion protein of C-CPE and PSIF. C-CPE is the C-terminal fragment of CPE 194 to 319 aa (Van Itallie et al., 2008). The dark area indicates the putative receptor-binding region of C-CPE located in its C terminus (Takahashi et al., 2005). PSIF contains domain II and III of PE. Domain II is critical for the escape of the toxin from the endosome to the cytosol, and domain III is responsible for the inhibition of protein synthesis (Ogata et al., 1990). B, purification of PSIF and C-CPE-PSIF. C-CPE-PSIF was expressed in *E. coli* and isolated by nickel-affinity chromatography. The purification of proteins (5 μg) was confirmed by SDS-PAGE followed by immunoblotting with antibody against the histidine tag. The putative molecular mass of C-CPE-PSIF is approximately 60 kDa.

2, 4, 7, 9, 11, and 14 at 5 $\mu\text{g}/\text{kg}$. Injection of C-CPE-PSIF significantly suppressed tumor growth, resulting in 36% of the tumor volume in the vehicle-treated group on day 16 (Fig. 4F). In contrast, C-CPE and C-CPE_{Y306A/L315A}-PSIF lacking claudin-4-binding activity had no effect on tumor growth, indicating that the antitumor activity of C-CPE-PSIF may be dependent on the claudin-4 targeting. The body weight of the mice did not change after injection of C-CPE-PSIF (data not shown). The appearance of the C-CPE-PSIF-injected mice also indicated no side effects.

Discussion

In the present study, we prepared a claudin-4-targeting ligand C-CPE coupled to PSIF of PE. We found that C-CPE-PSIF was selectively toxic to claudin-4-expressing cells and showed *in vivo* antitumor activity against mouse breast cancer cells, indicating that C-CPE may be a useful tool for claudin-targeted therapy.

Side effects of antitumor agents are a pivotal issue in clinical therapy. To reduce side effects, therapies that specif-




Fig. 2. Characterization of C-CPE-PSIF. A, interaction of C-CPE-PSIF with claudins. Wild-type BV (WT-BV), claudin-1-BV (CL1-BV), or claudin-4-BV (CL4-BV) was adsorbed onto a 96-well immunoplate at 0.5 $\mu\text{g}/\text{well}$. Then, 7.5 μmol C-CPE, C-CPE-PSIF, or PSIF was added to the well, and the protein bound to the BVs was measured by incubation of the anti-histidine-tag Ab, followed by horseradish peroxidase-labeled secondary Ab. The data represent the mean \pm S.D. of three independent experiments. *, significantly different from the WT-BV value ($p < 0.01$). B, cytotoxicity of C-CPE-PSIF in claudin-4/L (CL4/L) cells. After a 24-h treatment of CL4/L cells with PSIF or C-CPE-PSIF at the indicated concentration, the cell viability was measured by WST-8 assay. Viability (percentage) was calculated as a percentage of the vehicle-treated cells. The data represent the mean \pm S.D. of three independent experiments. *, significantly different from the vehicle-treated group ($p < 0.01$). C, competition assay using C-CPE. CL4/L cells were treated with C-CPE or BSA at the indicated concentration for 2 h, and then the cells were treated with C-CPE-PSIF (10 ng/ml) for 24 h. The cell viability was measured by WST-8 assay, as described above. The data represent the mean \pm S.D. of three independent experiments. *, significantly different between BSA and C-CPE-treated groups ($p < 0.01$). D, immunoblot analysis. Lysates of L, CL1/L, CL2/L, CL4/L, or CL5/L cells were subjected to SDS-PAGE, followed by immunoblotting with antibodies against the indicated CL. E, specific cytotoxicity of C-CPE-PSIF. L, CL1/L, CL2/L, CL4/L, and CL5/L cells were treated with C-CPE-PSIF for 24 h at the indicated concentration. The cell viability was assayed by WST-8 assay. The data represent the mean \pm S.D. of three independent experiments. *, significantly different from the L-cells ($p < 0.01$).




Fig. 3. Cytotoxic specificity of C-CPE-PSIF. **A**, expression of claudin-4 in SK-HEP-1 and HepG2 cells. The cell lysates were subjected to SDS-PAGE, followed by Western blotting with anti-claudin-4 Ab. MCF-7 cells were used as a positive control. **B**, cytotoxicity of C-CPE-PSIF in SK-HEP-1 and HepG2 cells. Cells were treated with PSIF or C-CPE-PSIF for 48 h at the indicated concentration. The cell viability was measured by WST-8 assay. The data are representative of at least three independent experiments. Data are the mean \pm S.D. ($n = 3$). *, significantly different from the vehicle-treated cells ($p < 0.01$). **C** and **D**, cytotoxicity of C-CPE-PSIF in TJ-developed or -undeveloped Caco-2 monolayer cells. TJ-developed cells were Caco-2 monolayer cells grown at confluence for 3 days. TJ-undeveloped cells were Caco-2 cells seeded at 10^4 cells/well in 96-well plates. The cell lysates were subjected to SDS-PAGE, followed by Western blotting with anti-claudin-4 Ab (**C**). The cells were treated with the indicated concentrations of C-CPE-PSIF for 48 h, and then the cell viability was measured as above (**D**). The data are representative of at least three independent experiments. Data are the mean \pm S.D. ($n = 3$). *, significantly different from the vehicle-treated cells ($p < 0.05$). **E** and **F**, effect of C-CPE-PSIF on TER and LDH release in Caco-2 monolayer cells. Caco-2 cells were grown on Transwell chambers to form tight junctions. When TER values were constant, C-CPE-PSIF was added to the apical or basolateral side in Transwell chambers at the indicated concentrations. After 0 and 48 h of incubation, TER values were measured (**E**), and the LDH release from the cell was determined (**F**). TER values and LDH release were calculated as the ratio of TER values at 0 h and of the total cellular LDH, respectively. The data are representative of at least three independent experiments. Data are the mean \pm S.D. ($n = 3$). *, significantly different from the vehicle-treated cells ($p < 0.05$).

ically target tumors are needed. Rapid advances in molecular biology and proteomics have allowed the identification of tumor-specific targets for cancer therapy. Therapy that exploits tumor-specific targets has the advantages of high specificity and low systemic toxicity relative to standard chemotherapy (Waldmann, 1991; Allen and Cullis, 2004). Target

molecule selection is critical to the success of targeted therapy. Most targeted therapies have been directed against growth factor receptors, such as ErbB2, which are overexpressed in cancers and are easily accessible because of their cell-surface localization (Zumkeller and Schofield, 1995; Deckert, 2009). A claudin-targeting strategy may have at




Fig. 4. In vivo antitumor activity of C-CPE-PSIF. **A**, expression of claudin-4 in 4T1 cells. Total RNA was extracted from the cells, followed by reverse transcription-polymerase chain reaction analysis (left). Cell lysates were subjected to SDS-PAGE, followed by Western blotting (right). β -Actin is an endogenous control. **B**, schematic structure of C-CPE_{Y306A/L315A}-PSIF. The dark area indicates the putative receptor-binding region of C-CPE in its C terminus (Takahashi et al., 2005). Among 16 amino acids, Tyr306 and Leu315 (indicated by italic letters) play a pivotal role in the binding of C-CPE with claudin-4 (Takahashi et al., 2008). **C**, interaction of C-CPE_{Y306A/L315A}-PSIF with claudin-4. WT-BV, CL1-BV, or CL4-BV was immobilized onto a 96-well immunoplate at 0.5 μ g/well. Then, 7.5 μ mol of C-CPE, C-CPE_{Y306A/L315A}-PSIF, or PSIF was added to the well, and the binding of the proteins to BVs was measured as described in the legend to Fig. 2A. The data represent the mean \pm S.D. of three independent experiments. *, significantly different from the WT-BV value ($p < 0.01$). **D**, cytotoxicity of C-CPE_{Y306A/L315A}-PSIF in CL4/L cells. After a 24-h treatment of CL4/L cells with PSIF, C-CPE-PSIF, or C-CPE_{Y306A/L315A}-PSIF at the indicated concentration, the cell viability was measured as described in the legend to Fig. 2B. The data represent the mean \pm S.D. of three independent experiments. *, significantly different from the vehicle-treated group ($p < 0.01$). **E**, cytotoxicity of C-CPE-PSIFs in 4T1 cells. Cells were treated with PSIF, C-CPE-PSIF, or C-CPE_{Y306A/L315A}-PSIF for 48 h at the indicated concentration. The cell viability was measured as above. The data represent the mean \pm S.D. of three independent experiments. *, significantly different from the vehicle-treated group ($p < 0.05$). **F**, in vivo antitumor activity of C-CPE-PSIF. 4T1 cells (2×10^6 cells) were intradermally inoculated into the right flank of mice on day 0, and each sample (5 μ g/kg) was intratumorally injected on days 2, 4, 7, 9, 11, and 14. Tumor growth was monitored by calculating tumor volume. The data are representative of three independent experiments. Each point is the mean \pm S.D. from five mice. *, significantly different from the vehicle (PBS)-treated group ($p < 0.05$).

least two advantages over the previous targeted methods. First, the claudin-targeting method can recognize different cellular localizations of claudin between normal epithelium and malignant tumors. The expression of claudin is altered in several cancers (Nichols et al., 2004; Morin, 2005; Tsukita et al., 2008). It is noteworthy that claudin-4 is localized to the basolateral cell membrane at sites of cell-cell contact in normal epithelium; in contrast, claudin-4 is localized to the cell membrane even at sites that lack cell-cell contact in poorly differentiated human breast cancers (Kominsky, 2006). The claudin-targeting strategy might have low side effects. Indeed, claudin-targeting therapy using CPE had antitumor activities without side effects in cancers of the breast, ovary, prostate, and pancreas (Long et al., 2001; Michl et al., 2001; Rangel et al., 2003; Kominsky et al., 2004; Santin et al., 2005). Second, a claudin-targeting molecule may be a promising agent for solid tumors. Interstitial pressure is higher in the center of a tumor, and it approaches normal physiological pressure toward the periphery (Jain, 1987, 1989). High-pressure regions usually coincide with regions of poor perfusion and lower vessel surface area; as a result, it is difficult for antitumor agents to be delivered to the intratumor tissue. If an antitumor agent has penetration-enhancing activity, the agent will be effective in therapy for solid tumors. We previously found that C-CPE enhanced intestinal permeability of a drug by modulation of the claudin-claudin interaction. Binding of C-CPE with claudin-4 is critical for modulation of the interaction. Together, C-CPE-PSIF may enhance the intratumoral permeation of antitumor agents by modulation of the claudin-claudin interaction. Combination therapy of a chemical agent with a claudin-4-targeting agent will have a synergistic effect compared with that of a chemical agent alone.

In a previous study, we prepared C-CPE₁₈₄₋₃₁₉-PSIF, in which C-CPE corresponding to 184 to 319 amino acids of CPE was used, and we found that C-CPE₁₈₄₋₃₁₉-PSIF is potently cytotoxic to claudin-4-expressing cells (Ebihara et al., 2006). However, C-CPE₁₈₄₋₃₁₉-PSIF had poor solubility (0.3 mg/ml) and low cytotoxicity ($IC_{50} = 2-3 \mu\text{g/ml}$ in claudin-4/L cells) (Ebihara et al., 2006). In 2008, Van Itallie et al. (2008) found that C-CPE₁₉₄₋₃₁₉ (referred to as C-CPE in the present study), which corresponds to amino acids 194 to 319 of CPE, is a claudin-4 binder with high solubility (10 mg/ml). We prepared a claudin-4-targeting molecule using C-CPE and found that C-CPE-PSIF has high solubility ($\sim 1.0 \text{ mg/ml}$) and high cytotoxicity ($IC_{50} = 1 \text{ ng/ml}$ in claudin-4/L cells). These results indicate that C-CPE is a promising claudin-4-targeting molecule for pharmaceutical therapy.

A targeting ligand coupled to a potent toxin is an attractive antitumor agent, but the application of the targeted ligand to clinical therapy has been limited because of its failure to concentrate at the site of the tumor (Shockley et al., 1992; Kreitman, 1999). The injection of agents directly into the tumor may circumvent these problems because therapy can be concentrated at the site of the tumor, thereby diminishing the risk to nontarget organs. The leakage of the agents from tumors into systemic flow may cause injury to normal tissues. If the claudin-4-targeting molecule leaks from the injected tumor tissue into the systemic flow, the leaked molecule interacts with tissues on the apical side. We found that C-CPE-PSIF is cytotoxic in a cellular polarity-dependent manner (Fig. 3, E and F). Singh et al. (2001) also showed that

Caco-2 cells are more sensitive to CPE when CPE is applied to the basal side rather than the apical side (Singh et al., 2001). These findings indicate that preparation of high affinity and specificity of claudin-4 binder may be critical for clinical application of a claudin-4-targeting ligand such as C-CPE-PSIF.

In summary, we prepared a novel claudin-4-targeting molecule, C-CPE-PSIF, which consists of C-CPE₁₉₄₋₃₁₉ and PSIF, and we found that the cytotoxicity of C-CPE-PSIF is specific to the expression and localization of claudin-4 and that the intratumoral administration of C-CPE-PSIF suppressed tumor growth. This is the first report that a claudin-4-targeting ligand C-CPE is useful for antitumor therapy. Future improvements in the cytotoxicity of the claudin-4-targeting strategy will be useful for its clinical application.

Acknowledgments

We thank the members of our laboratory for useful comments and discussion.

References

- Allen TM and Cullis PR (2004) Drug delivery systems: entering the mainstream. *Science* **303**:1818–1822.
- Anderson JM and Van Itallie CM (1995) Tight junctions and the molecular basis for regulation of paracellular permeability. *Am J Physiol* **269**:G467–G475.
- Deckert PM (2009) Current constructs and targets in clinical development for antibody-based cancer therapy. *Curr Drug Targets* **10**:158–175.
- Dunham W (2007) Report sees 7.6 million global 2007 cancer deaths. 2007 Dec 17, Reuters, New York. Available at <http://www.reuters.com/article/healthNews/idUSN1633064920071217>.
- Ebihara C, Kondoh M, Hasuike N, Harada M, Mizuguchi H, Horiguchi Y, Fujii M, and Watanabe Y (2006) Preparation of a claudin-targeting molecule using a C-terminal fragment of Clostridium perfringens enterotoxin. *J Pharmacol Exp Ther* **316**:255–260.
- Fujita K, Katahira J, Horiguchi Y, Sonoda N, Furuse M, and Tsukita S (2000) Clostridium perfringens enterotoxin binds to the second extracellular loop of claudin-3, a tight junction integral membrane protein. *FEBS Lett* **476**:258–261.
- Furuse M and Tsukita S (2006) Claudins in occluding junctions of humans and flies. *Trends Cell Biol* **16**:181–188.
- Hanna PC, Wiecekowski EU, Mietzner TA, and McClane BA (1992) Mapping of functional regions of Clostridium perfringens type A enterotoxin. *Infect Immun* **60**:2110–2114.
- Jain RK (1987) Transport of molecules in the tumor interstitium: a review. *Cancer Res* **47**:3039–3051.
- Jain RK (1989) Delivery of novel therapeutic agents in tumors: physiological barriers and strategies. *J Natl Cancer Inst* **81**:570–576.
- Jemal A, Siegel R, Ward E, Hao Y, Xu J, Murray T, and Thun MJ (2008) Cancer statistics, 2008. *CA Cancer J Clin* **58**:71–96.
- Katahira J, Inoue N, Horiguchi Y, Matsuda M, and Sugimoto N (1997) Molecular cloning and functional characterization of the receptor for Clostridium perfringens enterotoxin. *J Cell Biol* **136**:1239–1247.
- Kominsky SL (2006) Claudins: emerging targets for cancer therapy. *Expert Rev Mol Med* **8**:1–11.
- Kominsky SL, Vali M, Korz D, Gabig TG, Weitzman SA, Argani P, and Sukumar S (2004) Clostridium perfringens enterotoxin elicits rapid and specific cytolysis of breast carcinoma cells mediated through tight junction proteins claudin 3 and 4. *Am J Pathol* **164**:1627–1633.
- Kondoh M, Masuyama A, Takahashi A, Asano N, Mizuguchi H, Koizumi N, Fujii M, Hayakawa T, Horiguchi Y, and Watanabe Y (2005) A novel strategy for the enhancement of drug absorption using a claudin modulator. *Mol Pharmacol* **67**:749–756.
- Kreitman RJ (1999) Immunotoxins in cancer therapy. *Curr Opin Immunol* **11**:570–578.
- Kreitman RJ and Pastan I (2006) Immunotoxins in the treatment of hematologic malignancies. *Curr Drug Targets* **7**:1301–1311.
- Long H, Crean CD, Lee WH, Cummings OW, and Gabig TG (2001) Expression of Clostridium perfringens enterotoxin receptors claudin-3 and claudin-4 in prostate cancer epithelium. *Cancer Res* **61**:7878–7881.
- McClane BA and Chakrabarti G (2004) New insights into the cytotoxic mechanisms of Clostridium perfringens enterotoxin. *Anaerobe* **10**:107–114.
- Meunier V, Bourrié M, Berger Y, and Fabre G (1995) The human intestinal epithelial cell line Caco-2; pharmacological and pharmacokinetic applications. *Cell Biol Toxicol* **11**:187–194.
- Michl P, Buchholz M, Rolke M, Kunsch S, Löhr M, McClane B, Tsukita S, Leder G, Adler G, and Gress TM (2001) Claudin-4: a new target for pancreatic cancer treatment using Clostridium perfringens enterotoxin. *Gastroenterology* **121**:678–684.
- Michl P and Gress TM (2004) Bacteria and bacterial toxins as therapeutic agents for solid tumors. *Curr Cancer Drug Targets* **4**:689–702.
- Mitic LL and Anderson JM (1998) Molecular architecture of tight junctions. *Annu Rev Physiol* **60**:121–142.

- Morin PJ (2005) Claudin proteins in human cancer: promising new targets for diagnosis and therapy. *Cancer Res* **65**:9603–9606.
- Nichols LS, Ashfaq R, and Iacobuzio-Donahue CA (2004) Claudin 4 protein expression in primary and metastatic pancreatic cancer: support for use as a therapeutic target. *Am J Clin Pathol* **121**:226–230.
- Ogata M, Chaudhary VK, Pastan I, and FitzGerald DJ (1990) Processing of *Pseudomonas* exotoxin by a cellular protease results in the generation of a 37,000-Da toxin fragment that is translocated to the cytosol. *J Biol Chem* **265**:20678–20685.
- Paperna T, Peoples R, Wang YK, Kaplan P, and Francke U (1998) Genes for the CPE receptor (CPETR1) and the human homolog of RVP1 (CPETR2) are localized within the Williams-Beuren syndrome deletion. *Genomics* **54**:453–459.
- Rangel LB, Agarwal R, D'Souza T, Pizer ES, Alò PL, Lancaster WD, Gregoire L, Schwartz DR, Cho KR, and Morin PJ (2003) Tight junction proteins claudin-3 and claudin-4 are frequently overexpressed in ovarian cancer but not in ovarian cystadenomas. *Clin Cancer Res* **9**:2567–2575.
- Sakihama T, Sato T, Iwanari H, Kitamura T, Sakaguchi S, Kodama T, and Hamakubo T (2008) A simple detection method for low-affinity membrane protein interactions by baculoviral display. *PLoS One* **3**:e4024.
- Santin AD, Cané S, Bellone S, Palmieri M, Siegel ER, Thomas M, Roman JJ, Burnett A, Cannon MJ, and Pecorelli S (2005) Treatment of chemotherapy-resistant human ovarian cancer xenografts in C.B-17/SCID mice by intraperitoneal administration of *Clostridium perfringens* enterotoxin. *Cancer Res* **65**:4334–4342.
- Shockley TR, Lin K, Sung C, Nagy JA, Tompkins RG, Dedrick RL, Dvorak HF, and Yarmush ML (1992) A quantitative analysis of tumor specific monoclonal antibody uptake by human melanoma xenografts: effects of antibody immunological properties and tumor antigen expression levels. *Cancer Res* **52**:357–366.
- Singh U, Mitic LL, Wieckowski EU, Anderson JM, and McClane BA (2001) Comparative biochemical and immunocytochemical studies reveal differences in the effects of *Clostridium perfringens* enterotoxin on polarized CaCo-2 cells versus Vero cells. *J Biol Chem* **276**:33402–33412.
- Takahashi A, Komiya E, Kakutani H, Yoshida T, Fujii M, Horiguchi Y, Mizuguchi H, Tsutsumi Y, Tsunoda S, Koizumi N, et al. (2008) Domain mapping of a claudin-4 modulator, the C-terminal region of C-terminal fragment of *Clostridium perfringens* enterotoxin, by site-directed mutagenesis. *Biochem Pharmacol* **75**:1639–1648.
- Takahashi A, Kondoh M, Masuyama A, Fujii M, Mizuguchi H, Horiguchi Y, and Watanabe Y (2005) Role of C-terminal regions of the C-terminal fragment of *Clostridium perfringens* enterotoxin in its interaction with claudin-4. *J Control Release* **108**:56–62.
- Tsukita S and Furuse M (1999) Occludin and claudins in tight-junction strands: leading or supporting players? *Trends Cell Biol* **9**:268–273.
- Tsukita S, Yamazaki Y, Katsuno T, Tamura A, and Tsukita S (2008) Tight junction-based epithelial microenvironment and cell proliferation. *Oncogene* **27**:6930–6938.
- Van Itallie CM, Betts L, Smedley JG 3rd, McClane BA, and Anderson JM (2008) Structure of the claudin-binding domain of *Clostridium perfringens* enterotoxin. *J Biol Chem* **283**:268–274.
- Vermeer PD, Einwalter LA, Moninger TO, Rokhlina T, Kern JA, Zabner J, and Welsh MJ (2003) Segregation of receptor and ligand regulates activation of epithelial growth factor receptor. *Nature* **422**:322–326.
- Waldmann TA (1991) Monoclonal antibodies in diagnosis and therapy. *Science* **252**:1657–1662.
- Wodarz A and Näthke I (2007) Cell polarity in development and cancer. *Nat Cell Biol* **9**:1016–1024.
- Zumkeller W and Schofield PN (1995) Growth factors, cytokines and soluble forms of receptor molecules in cancer patients. *Anticancer Res* **15**:343–348.

Address correspondence to: Dr. Masuo Kondoh, Laboratory of Bio-Functional Molecular Chemistry, Graduate School of Pharmaceutical Sciences, Osaka University, Suita, Osaka 565-0871, Japan. E-mail: masuo@phs.osaka-u.ac.jp

生体バリアを利用した創薬研究

第1回日本 DDS 学会奨励賞(臨床)受賞によせて

Claudin binder as a novel drug delivery system: the 1st Encouragement Award in the 25th Annual Meeting of the Japan Society of Drug Delivery System

Masuo Kondoh · Azusa Takahashi ·
Rie Saeki · Kiyohito Yagi*

Epithelium surrounds the body separating intra and outer body. Passing across epithelium is the first step of drug absorption. Most malignant tumors are derived from epithelium. Moreover, epithelium is the first line of defense against pathological microorganisms. These findings strongly indicate that epithelium is a promising target for drug discovery. Claudin family, which is identified by Dr Tsukita group, is a key molecule for epithelial barrier.

In this review, we discussed about claudin as a potential target for drug development.

進化の過程において、多細胞生物は生体内外および組織内外を隔てる障壁として上皮細胞層を発達させてきた。上皮細胞層の透過過程が薬物吸収の第一ステップであること、悪性腫瘍の90%が上皮由来であること、上皮細胞層は感染性病原微生物の侵入門戸になっていることから、上皮細胞は創薬ターゲットとして有用な性質を有している。1993年以降、上皮細胞バリア構成蛋白質 occludin, claudinなどが同定され、生体バリアの分子基盤が詳らかにされつつある。

本総説では、当研究グループで推進している生体バリアの分子基盤 claudin を利用した創薬研究について、その一端を紹介したい。

60兆個の細胞からなる個体は上皮細胞層により生体内外に区別され、さらに生体内では上皮細胞層および血管内皮細胞層によって脳、血管、腎臓などの

コンパートメントが形成されている。コンパートメント内の恒常性を維持するためには内外の物質移動を制御することが不可欠であり、上皮細胞層や血管内皮細胞層は、生体内外および組織内外を隔てるバリアとして機能することにより、恒常性維持に深く関わっている。これらの細胞層では、隣接する細胞間に tight junction (TJ) が発達しており、TJ によって細胞間隙はシールされ物質の漏れが抑制されている。

70年代には TJ がストランド様の構造をしていることが見いだされていたものの、TJ 分子基盤は長年にわたり不明なままであった¹⁾。93年、京大月田承一郎博士のグループにより TJ 構成蛋白質として occludin が同定され TJ が蛋白質で構成されていることがはじめて明らかにされた²⁾。さらに98年に、同グループにより TJ バリアの機能本体として claudin が同定された³⁾。

Claudin は24種の分子からなる family を形成しており、細胞間隙におけるホモフィリック/ヘテロフィリックな結合により TJ をシールしていると考えられている^{4,5)}。興味深いことにバリア機能には組織特異性が認められ、claudin-1 は皮膚バリア、claudin-5 は血液脳関門バリア、claudin-11 は血液精巣関門バリアを担っており、claudin を利用した新規薬物送達法開発の可能性が強く示唆されている⁶⁻⁸⁾。また、悪性腫瘍の90%を占める上皮がんでは多くのがん種で claudin の発現異常が観察されており、claudin は上皮がんの創薬ターゲットとしても衆目を集めている^{9,10)}。さらに、claudin が粘膜

* Laboratory of Bio-Functional Molecular Chemistry, Graduate School of Pharmaceutical Sciences, Osaka University 大阪大学大学院薬学研究所生体機能分子化学分野

表1 タイトジャンクション研究

年代	研究内容
1973年	TJ ストランドの発見
1982年	脂質ミセル説の提唱
1993年	Occludin の発見
1998年	Claudin の発見
1999年以降	Claudin バリア機能の証明
2005年	Tricellulin の発見

表2 既存の生体バリアを利用した創薬研究

吸収促進剤	生体内標的分子
EDTA	Ca ²⁺ のキレート
オレイン酸	脂質二重膜の流動性亢進
NO 供与剤	不明
カプリン酸ナトリウム	ホスホリパーゼC

免疫組織で高発現していること¹¹⁾、occludin および claudin がC型肝炎ウイルスの感染受容体であることも報告されており^{12,13)}、TJ分子基盤を利用した粘膜ワクチンおよび感染阻害薬創出の可能性も見いだされている。

このように、月田グループによる occludin および claudin の発見に端を発した上皮細胞バリア研究の進展に伴い、従来までの「上皮細胞バリア=物質透過障壁」という概念に加え、「上皮細胞バリア分子基盤=創薬ターゲット」という新たな萌芽が生まれつつある。

本稿では、上述した本邦発の新規創薬ターゲット、上皮細胞バリア制御分子 claudin の可能性について、薬物送達法および抗がん剤開発に焦点を絞り、当研究グループの研究成果を概説する。

上皮細胞バリアの分子基盤

隣接する細胞は10~20 nmの空隙を隔てて向かい合っており、アピカル側では隣接する細胞膜がストランド様の構造体 TJ によって密着し、細胞間の距離がゼロとなるキッキングポイントにより細胞間隙における物質の移動が制御されている¹⁾。電子顕微鏡を用いた解析から、TJ ストランドは網目状にラテラル面に広がり物質の移動を制御していることが1970年代には明らかにされていたものの、このストランドの本体については脂質ミセル説なども提唱され長年にわたり議論が続いていた(表1)¹⁴⁾。

1993年に、京大月田グループにより、はじめての TJ 構成蛋白質として occludin が見いだされ、TJ ストランドが蛋白質で構成されていることが示された²⁾。1998年に新たな TJ 構成蛋白質として claudin が同定され、1999年以降 claudin が TJ バリア能の本体を担うことを示唆する知見が集積しつつある^{3,15,16)}。さらに2005年には、同グループにより、上皮細胞層において三つの細胞が交わるジャンクション(トリセルラージャンクション)の構成蛋白質として tricellulin が同定されている¹⁷⁾。Tricellulin をノックダウンすると上皮細胞バリア機能が低下することから、上皮細胞バリアは2細胞間のシール機構と3細胞間のシール機構により構成されていることが実験的に証明された¹⁷⁾。

上皮細胞バリアを利用した第一世代の創薬研究

前述したように、上皮細胞層は生体内外を隔てる障壁として機能しており、薬物吸収に際しては上皮細胞層の透過が不可欠となっている。すでに30年あまりにわたり上皮細胞バリアを利用した創薬研究として吸収促進剤が研究されてきており、EDTA、オレイン酸、NO 供与剤、カプリン酸ナトリウムなどが吸収促進活性物質として見いだされている(表2)。それぞれの作用点は細胞間隙に存在するカルシウムイオン、細胞膜、phospholipase C などであり、TJ の開口を通じて吸収促進効果を発揮すると考えられている。

これらアプローチが研究開発されていた1980年~98年には TJ 構成蛋白質は未解明であり、吸収促進作用に組織特異性が乏しいこと、TJ の開口に伴い薬物以外の物質の非特異的な流入が生じることなどから、臨床応用されている吸収促進剤はカプリン酸ナトリウムなどにすぎない^{18~20)}。

上皮細胞バリアを利用した第二世代の創薬研究

当研究グループでは、上述した吸収促進剤の有所する課題は TJ を利用した薬物送達法の限界を示しているのではなく、TJ の分子基盤に立脚したア




図1 C-CPEの腸管吸収促進活性

a: 血漿中FD-4濃度の経時変化。b: AUC値。FITCラベルした分子量4000のデキストラン(FD-4)をモデル薬物として用いて、ラット腸管ループ法によりC-CPEの吸収促進活性を解析(Kondoh M et al. 2005²⁵⁾を一部改変)。

ブローチが採られてこなかったことに起因していると考え、上述した吸収促進剤を第一世代のTJ modulatorと定義し、TJの分子基盤に立脚した第二世代のTJ modulatorを利用した薬物送達研究を進めてきた²¹⁾。

1993年、京大月田グループによりTJ構成蛋白質として4回膜貫通蛋白質occludinが見いだされ、TJが蛋白質によって構成されていることがはじめて明らかにされた²⁾。しかしながら、occludinを欠

損させても、機能的、構造的に正常なTJが構成されていたことから、occludinはTJバリア機能分子ではないことが示唆された²²⁾。この報告とほぼ時を同じくして、1998年に新たなTJ構成蛋白質claudinが同定された(表1)³⁾。

Claudinは分子量23 kDaの4回膜貫通蛋白質であり、現在までに24種類の分子が見いだされている。非常に興味深いことに、発現およびバリア機能には組織特異性が認められ、claudin-1欠損マウスでは重層上皮細胞のバリア機能が異常をきたし、分子量600程度の分子が皮膚を透過し、claudin-5欠損マウスでは血液脳関門を分子量800程度の分子が通過する^{5,6,8,15,23)}。さらに、claudin-11は血液精巣関門バリアを担っていることも明らかにされている⁷⁾。これらの報告はclaudinを分子特異的に制御することができれば新たな薬物送達法の開発に繋がることがを意味している。

現在のところ、claudinのバリア機能を阻害する分子としてはウエルシュ菌エンテロトキシン(CPE)のC末断片(C-CPE)のみがclaudin-4のバリア機能を阻害する分子として知られている^{16,24)}。そこでC-CPEをclaudin modulatorのモデル分子として用いて、claudinを利用した薬物送達法創出の可能性について検証を試みた。

まず、C-CPEの薬物送達活性を分子量4000のデキストラン(FD-4)をモデル分子として用いてラット腸管ループ法により解析したところ、C-CPE添加濃度依存的に血漿中FD-4濃度は上昇していた(図1a)。C-CPEの吸収促進活性は臨床応用されている吸収促進剤であるカプリン酸ナトリウムに比して400倍もの値を示しており(図1b)、さらに吸収促進効果には組織特異性も観察されていた²⁵⁾。

CPEはC末30アミノ酸を介して受容体と相互作用することが示唆されている²⁶⁾。そこで、C-CPEの薬物送達活性におけるclaudin-4の関与を検討するために、受容体結合領域をすべて欠損させたC-CPE289およびその一部を欠損させたC-CPE303を作成し、ラット腸管における吸収促進活性を解析したところ、C-CPE289、C-CPE303処理では吸収促進活性が著しく減弱していた(図2a, b)。

以上の結果より、claudinを利用した粘膜吸収促進法の有用性が示唆される。




図2 C-CPEの吸収促進作用における claudin-4の関与

a: C-CPEs. b: 腸管吸収促進効果
C-CPEの claudin-4 結合領域を欠損させた変異体(a)のFD-4吸収促進活性を解析(b). なお、データは Kondoh M et al., 2005²⁵⁾を一部改変.

上皮細胞バリアを利用した第三世代の創薬研究

上述したように、従来の生体バリアを利用した創薬研究は薬物吸収促進に焦点が当てられてきた。2000年以降、悪性腫瘍(年間死者数700万人)の90%を占める上皮由来のがんと claudinとの関連性についても、多方面からの研究が進み、卵巣がん、膵臓がん、膀胱がんなど12種類あまりのがんにおける claudinの高発現が見いだされ、claudinががんターゲットの標的分子としても注目されている^{9,10)}。

さて、正常な上皮細胞層では水平方向に細胞は分裂しコンタクトインヒビションにより細胞増殖は制御されている。一方、上皮細胞ががん化すると細




図3 Claudin-4 指向性分子

a: C-CPE-PSIF. b: Claudin(CL)特異性 *Pseudomonas* exotoxinの蛋白質合成阻害ドメイン(PSIF)とC-CPEとの融合体(a)を作製し、各種 claudin 発現細胞に各濃度24時間処理後にWSTアッセイにより細胞毒性を解析(b). なお、データは Saeki R et al., in press²⁸⁾を一部改変.

胞分裂軸が回転し垂直方向への分裂が始まり、コンタクトインヒビションがかからず腫瘍組織を形成していくと考えられている。このがん化早期イベントである分裂軸の回転を利用したがん治療戦略を構築することができれば、がんの早期診断・早期治療法の開発に繋がるといえる。

当研究グループでは、分裂軸の回転に伴い lateral 面から apical 面に露出する TJ 構成蛋白質 claudin を利用した新規がん治療法の開発を目指し、claudin-4 binder を利用したがん治療戦略の構築を試みている^{27,28)}。

まず、claudin 指向性抗がん剤のモデルとして、claudin-4 結合分子 C-CPE と緑膿菌由来の蛋白質合成阻害因子(PSIF)との融合蛋白質を作製した(図 3a)。C-CPE-PSIF は claudin-1, -2, -5 発現 L 細胞では 10 ng/mL でもまったく細胞障害性を示さないのに対して、claudin-4 発現 L 細胞では 1 ng/mL 処理でも顕著な細胞毒性を発揮していた(図 3b)。さらに、Caco-2 細胞の単層膜培養系に apical 側、もしくは basal 側から C-CPE-PSIF を添加したとこ




図4 生体バリアを利用した創薬研究

る。C-CPE-PSIF を basal 側から加えたときのみ細胞毒性が観察され、C-CPE は claudin の局在性を認識する新しいタイプの抗がん活性分子であると推察された (data not shown)。さらに、マウス乳がん由来細胞株 4T1 細胞を用いて C-CPE-PSIF の抗腫瘍効果を解析したところ、C-CPE-PSIF 処理で *in vitro* および *in vivo* での抗腫瘍活性が認められ体重減少などの副作用は観察されず、claudin-4 結合性を消失させた変異体では *in vitro* および *in vivo* 抗腫瘍活性がともに消失していた²⁸⁾。

これらの結果を踏まえ、現在当研究グループでは、がん化の早期イベント「分裂軸の回転」を標的とした新規がん診断法・治療法の開発を目指し、新規 claudin binder の創出、既存の claudin binder のプローブ化、最適化などを進めている。

まとめ

今回は紙面の都合上割愛させていただいているが、粘膜免疫組織に高発現している claudin を利用したワクチン開発、ウイルスの感染受容体である claudin-1 や occludin を標的とした感染阻害薬

の開発などに関しても研究が進展しつつあり、現在までに claudin を利用した非侵襲性投与技術、claudin 指向性抗がん剤、claudin 指向性粘膜ワクチンの創製に成功し、世界に先駆けて生体バリアの分子基盤を利用した創薬戦略の有用性を提唱してきた^{19, 21, 25, 28-30)}。

周知のように、わが国にはカドヘリンをはじめとした上皮細胞生物学の基礎的知見が集積している。当研究グループでは、本邦固有の上皮細胞生物学の土壌に生まれた創薬研究の萌芽を育み、わが国独自の生体バリアを利用した創薬研究領域を開拓し、本邦発の創薬シーズを一つでも多く“bench side to bed side”へと顕在化できるよう最善を尽くしていきたい(図4)。

本稿で紹介した研究成果は、昭和薬科大学渡邊善照教授、藤井まき子准教授、アスピオファーマ株式会社内田博司博士・花田雄志博士をはじめとした多くの先生方のご指導およびご協力、八木ラボおよび渡辺ラボメンバーの知的好奇心および激的な努力の所産であり、相互作用をいただいたすべての方々へ衷心より御礼申し上げます。

なお、本研究は、文部科学省科学研究費補助金、JST 知的クラスター創成事業、厚生労働省科学研究費補助金、財団法人武田科学振興財団、財団法人薬学奨励財団、財団法人医科学応用財団のグラントサポートにより実施されたものであり、各助成機関に衷心より御礼申し上げます。

文 献

- 1) Staehelin LA : Further observations on the fine structure of freeze-cleaved tight junctions. *J Cell Sci* 13 : 763-786, 1973.
- 2) Furuse M, Hirase T, Itoh M, Nagafuchi A, Yonemura S et al. : Occludin : a novel integral membrane protein localizing at tight junctions. *J Cell Biol* 123 : 1777-1788, 1993.
- 3) Furuse M, Fujita K, Hiiiragi T, Fujimoto F, Tsukita S : Claudin-1 and -2 : novel integral membrane proteins localizing at tight junctions with no sequence similarity to occludin. *J Cell Biol* 141 : 1539-1550, 1998.
- 4) Furuse M, Sasaki H, Tsukita S : Manner of interaction of heterogeneous claudin species within and between tight junction strands. *J Cell Biol* 147 : 891-903, 1999.
- 5) Furuse M, Tsukita S : Claudins in occluding junctions of humans and flies. *Trends Cell Biol* 16 : 181-188, 2006.
- 6) Furuse M, Hata M, Furuse K, Yoshida Y, Haratake A et al. : Claudin-based tight junctions are crucial for the mammalian epidermal barrier : a lesson from claudin-1-deficient mice. *J Cell Biol* 156 : 1099-1111, 2002.
- 7) Gow A, Southwood CM, Li JS, Pariali M, Riordan GP et al. : CNS myelin and sertoli cell tight junction strands are absent in *Osp/claudin-11* null mice. *Cell* 99 : 649-659, 1999.
- 8) Nitta T, Hata M, Gotoh S, Seo Y, Sasaki H et al. : Size-selective loosening of the blood-brain barrier in claudin-5-deficient mice. *J Cell Biol* 161 : 653-660, 2003.
- 9) Kominsky SL : Claudins : emerging targets for cancer therapy. *Expert Rev Mol Med* 8 : 1-11, 2006.
- 10) Morin PJ : Claudin proteins in human cancer : promising new targets for diagnosis and therapy. *Cancer Res* 65 : 9603-9606, 2005.
- 11) Tamagawa H, Takahashi I, Furuse M, Yoshitake-Kitano Y, Tsukita S et al. : Characteristics of claudin expression in follicle-associated epithelium of Peyer's patches : preferential localization of claudin-4 at the apex of the dome region. *Lab Invest* 83 : 1045-1053, 2003.
- 12) Evans MJ, von Hahn T, Tscherner DM, Syder AJ, Panis M et al. : Claudin-1 is a hepatitis C virus co-receptor required for a late step in entry. *Nature* 446 : 801-805, 2007.
- 13) Ploss A, Evans MJ, Gaysinskaya VA, Panis M, You H et al. : Human occludin is a hepatitis C virus entry factor required for infection of mouse cells. *Nature* 457 : 882-886, 2009.
- 14) Kachar B, Reese TS : Evidence for the lipidic nature of tight junction strands. *Nature* 296 : 464-466, 1982.
- 15) Furuse M : Knockout animals and natural mutations as experimental and diagnostic tool for studying tight junction functions in vivo. *Biochim Biophys Acta* 1788 : 813-819, 2009.
- 16) Sonoda N, Furuse M, Sasaki H, Yonemura S, Katahira J et al. : *Clostridium perfringens* enterotoxin fragment removes specific claudins from tight junction strands : Evidence for direct involvement of claudins in tight junction barrier. *J Cell Biol* 147 : 195-204, 1999.
- 17) Ikenouchi J, Furuse M, Furuse K, Sasaki H, Tsukita S et al. : Tricellulin constitutes a novel barrier at tricellular contacts of epithelial cells. *J Cell Biol* 171 : 939-945, 2005.
- 18) Aungst BJ : Intestinal permeation enhancers. *J Pharm Sci* 89 : 429-442, 2000.
- 19) Kondoh M, Yagi K : Progress in absorption enhancers based on tight junction. *Expert Opin Drug Deliv* 4 : 275-286, 2007.
- 20) Utoguchi N, Watanabe Y, Shida T, Matsumoto M : Nitric oxide donors enhance rectal absorption of macromolecules in rabbits. *Pharm Res* 15 : 870-876, 1998.
- 21) Kondoh M, Yoshida T, Kakutani H, Yagi K : Targeting tight junction proteins-significance for drug development. *Drug Discov Today* 13 : 180-186, 2008.
- 22) Saitou M, Fujimoto K, Doi Y, Itoh M, Fujimoto T et al. : Occludin-deficient embryonic stem cells can differentiate into polarized epithelial cells bearing tight junctions. *J Cell Biol* 141 : 397-408, 1998.
- 23) Morita K, Furuse M, Fujimoto K, Tsukita S : Claudin multigene family encoding four-transmembrane domain protein components of tight junction strands. *Proc Natl Acad Sci U S A* 96 : 511-516, 1999.
- 24) Katahira J, Inoue N, Horiguchi Y, Matsuda M, Sugimoto N : Molecular cloning and functional characterization of the receptor for *Clostridium perfringens* enterotoxin. *J Cell Biol* 136 : 1239-1247, 1997.
- 25) Kondoh M, Masuyama A, Takahashi A, Asano N, Mizuguchi H et al. : A novel strategy for the enhancement of drug absorption using a claudin modulator. *Mol Pharmacol* 67 : 749-756, 2005.
- 26) Hanna PC, Mietzner TA, Schoolnik GK, McClane BA : Localization of the receptor-binding region of *Clostridium perfringens* enterotoxin utilizing cloned toxin fragments and synthetic peptides. The 30 C-terminal amino acids define a functional binding region. *J Biol Chem* 266 : 11037-11043, 1991.
- 27) Ebihara C, Kondoh M, Hasuike N, Harada M, Mizuguchi H et al. : Preparation of a claudin-targeting molecule using a C-terminal fragment of *Clostridium perfringens* enterotoxin. *J Pharmacol Exp Ther* 316 : 255-260, 2006.
- 28) Saeki R, Kondoh M, Kakutani H, Tsunoda S, Mochizuki Y et al. : A novel tumor-targeted therapy using a claudin-4-targeting molecule. *Mol Pharmacol*, in press.
- 29) Kondoh M, Yagi K : Tight junction modulators : promising candidates for drug delivery. *Curr Med Chem* 14 : 2482-2488, 2007.
- 30) Matsuhisa K, Kondoh M, Takahashi A, Yagi K : Tight junction modulator and drug delivery. *Expert Opin Drug Deliv* 6 : 509-515, 2009.