

**Figure 1** Correlation between maximal and mean levels of alanine aminotransferase (ALT) (a) and hepatitis B virus (HBV) DNA (b) after discontinuation of nucleos(t)ide analogs (NAs). Open circles indicate patients with detectable hepatitis B e antigen (HBeAg) and closed squares indicate patients without detectable HBeAg.

than 5.7 log copies/mL during the follow-up period after NA discontinuation were not likely to achieve the HBV DNA criterion of a successful discontinuation of below 4.0 log copies/mL. Similarly, it could be inferred that patients reaching ALT levels higher than 79 IU/L would also not likely achieve the ALT criterion of a successful discontinuation of below 30 IU/L.

Based on our findings, we judged that a relapse of hepatitis B occurred when serum ALT exceeded 79 IU/L or when serum HBV DNA exceeded 5.7 log copies/mL

following NA discontinuation. Accordingly, 92 (73%) of the 126 patients enrolled in the present study showed a relapse. We set the follow-up period as discontinuation to relapse for relapse patients and as discontinuation to the last recorded examination for patients without relapse. Whereas re-administration of NAs due to relapse was commenced in 70% of relapse patients in the follow-up period, none was performed in non-relapse patients during that time.

### Elimination of cases likely to show relapse of hepatitis

As it is generally believed that patients who are positive for HBeAg and/or have a higher level of HBV DNA at discontinuation of NAs are likely to relapse, these factors were assessed first. The progression of analyses in the present study and the population structure of each analysis are shown in Figure 2.

The non-relapse rate was compared using the Kaplan–Meier method between 31 patients with HBV DNA equal to or higher than 3.0 log copies/mL and 95 patients with levels lower than 3.0 log copies/mL when NAs were discontinued (Fig. 3). The revised cut-off value of 3.0 log copies/mL was determined by ROC analysis (AUC = 0.709,  $P < 0.001$ ). Thirty (97%) of 31 patients with HBV DNA equal to or higher than 3.0 log copies/mL relapsed within one year of discontinuation. On the other hand, approximately 30% of patients with levels lower than 3.0 log copies/mL showed prolonged non-relapse. Thus, the 31 patients with high HBV DNA at the time of discontinuation were eliminated from the following analyses.

In the remaining 95 patients, the non-relapse rate was compared using the Kaplan–Meier method between 10 patients with detectable HBeAg and 85 patients without HBeAg when NAs were discontinued (Fig. 4). Ninety percent of patients with HBeAg experienced relapse within one year, which was significantly ( $P = 0.005$ ) higher than in cases without HBeAg. In patients without HBeAg, the non-relapse rate decreased rapidly during the first year to approximately 45%, and then decreased relatively slowly over the following 3 years to nearly 30%. It is noteworthy that this subgroup did not relapse afterwards. Since the relapse rate was high among patients with detectable HBeAg, they were excluded from the following analyses as well.

### Factors associated with relapse of hepatitis after discontinuation of NAs

Additional factors associated with relapse of hepatitis were analyzed in the remaining 85 patients who were




Figure 2 The progression of analyses in the present study and population structure of each analysis.

both negative for HBeAg and whose serum HBV DNA was lower than 3.0 log copies/mL at NA cessation. Table 1 shows the comparison of clinical and virological backgrounds between the 53 relapse and 32 non-relapse patients using univariate analysis. Age and gender distributions were similar between the groups. Approximately 75% of the 85 patients had HBV genotype C, but the distribution of genotypes did not differ between the groups. Approximately 90% of patients were being treated with LVD alone at the time of discontinuation, compared with 6% of patients being given ETV. The median duration of NA treatment was about two times longer in patients without relapse. Levels of both HBsAg

and HBcrAg were significantly lower in non-relapse patients than in relapse patients at the time of NA discontinuation. The difference between serum HBsAg was also significant at the initiation of NAs, but not that of HBcrAg. As only patients with HBV DNA lower than 3.0 log copies/mL were analyzed, the majority of these cases showed levels below the 2.6 log copies/mL lower detection limit of the Amplicor assay at NA discontinuation. We therefore also tested HBV DNA with a TaqMan assay, in 43 patients whose serum samples were available. The prevalence of patients having a negative detection signal did not differ between the two groups. The number of




Figure 3 Comparison of non-relapse rates using the Kaplan–Meier method between 31 patients with serum hepatitis B virus (HBV) DNA equal to or higher than 3.0 log copies/mL and 95 patients with serum HBV DNA lower than 3.0 log copies/mL at the time of nucleos(t)ide analog (NA) discontinuation.




Figure 4 Comparison of non-relapse rates using the Kaplan–Meier method between 10 patients with detectable hepatitis B e antigen (HBeAg) and 85 patients without detectable HBeAg at the time of nucleos(t)ide analog (NA) discontinuation.

**Table 1** Comparison of clinical and virological backgrounds between patients with and without relapse of hepatitis at initiation and discontinuation of nucleos(t)ide analogs (NAs)

Background	Non-relapse patients (n = 32)	Relapse patients (n = 53)	P-value
At initiation of NAs			
Age (years)†	47 (17–75)	48 (26–74)	>0.2
Gender (M : F)	23:9	32:21	>0.2
ALT (IU/L)†	183 (9–1182)	187 (20–2052)	>0.2
Genotype (A : B : C : UD)	1:2:21:8	0:3:44:6	0.193
HBeAg (positive)‡	11 (34%)	16 (30%)	>0.2
HBV DNA			
Amplicor assay (log copies/mL)†	6.2 (<2.6–>7.6)	6.5 (<2.6–>7.6)	0.099
HBsAg (log IU/mL)†	2.7 (0.1–4.3)	3.3 (1.6–3.9)	0.018
HBcrAg (log U/mL)†	5.2 (<3.0–>6.8)	5.6 (<3.0–>6.8)	>0.2
At discontinuation of NAs			
Age (years)†	50 (21–78)	49 (26–79)	>0.2
NAs (LVD : LVD+ADV : ETV : ADV)	28:1:3:0	50:0:2:1	>0.2
Duration of NA treatment (months)†	36 (4–129)	17 (4–84)	0.007
Follow-up period after discontinuation of NAs (months)†	45 (6–123)	12 (1–111)	0.002
ALT (IU/L)†	16 (7–38)	20 (9–65)	0.002
HBV DNA			
Amplicor assay (log copies/mL)†	<2.6 (<2.6–2.9)	<2.6 (<2.6–2.9)	>0.2
TaqMan assay (negative signal)‡	5 (23%) (n = 22)	3 (14%) (n = 21)	>0.2
TaqMan assay (negative or positive signal)‡	13 (59%) (n = 22)	13 (62%) (n = 21)	>0.2
HBsAg (log IU/ml)†	2.0 (<–1.5–4.3)	3.1 (0.6–4.0)	0.001
HBcrAg (log IU/mL)†	3.4 (<3.0–4.9)	4.3 (<3.0–>6.8)	0.003

†Data are expressed as the median (range)

‡Data are expressed as a positive number (%)

ADV, adefovir dipivoxil; ALT, alanine aminotransferase; ETV, entecavir; HBcrAg, hepatitis B core-related antigen; HBeAg, hepatitis B e antigen; HBsAg, hepatitis B surface antigen; LVD, lamivudine; UD, undetermined.

patients with a negative detection signal or a positive signal also did not vary significantly. The follow-up period after discontinuation of NAs was significantly shorter in patients with relapse than in those without because formal follow-up ended once patients relapsed. The median period of follow-up was 45 months in patients without relapse.

Multivariate analyses revealed that a shorter duration of NA treatment and higher levels of HBsAg and HBcrAg at discontinuation were significantly associated with the occurrence of hepatitis relapse (Table 2). The cut-off

values that showed the highest significance by ROC analysis were 1.9 log IU/mL for HBsAg (AUC = 0.707,  $P = 0.001$ ), 4.0 log U/mL for HBcrAg (AUC = 0.692,  $P = 0.003$ ), and 16 months (AUC = 0.674,  $P = 0.007$ ) for treatment duration.


### Model for predicting relapse of hepatitis using levels of HBsAg and HBcrAg

The existence of a second cut-off value was suggested by ROC analysis for both of HBsAg (2.9 log IU/mL) and HBcrAg (3.0 log IU/mL) to discriminate between

**Table 2** Multivariate analysis of factors associated with relapse of hepatitis after discontinuation of nucleos(t)ide analogs (NAs)

Factor	Hazard ratio	95%CI	P-value
HBsAg at discontinuation $\geq 1.9$ log IU/mL	5.21	1.87–14.55	0.002
HBcrAg at discontinuation $\geq 4.0$ log U/mL	2.20	1.25–3.87	0.006
Duration of NA treatment $\geq 16$ months	0.54	0.31–0.93	0.027

CI, confidence interval; HBcrAg, hepatitis B core-related antigen; HBsAg, hepatitis B surface antigen.




**Figure 5** Receiver operating characteristic curve (ROC) analysis of hepatitis B surface antigen (HBsAg) and hepatitis B core-related antigen (HBcrAg) to discriminate between patients with and without hepatitis relapse. The existence of two inflection points is suggested for both HBsAg and HBcrAg. Short diagonal lines indicate main inflection points and short broken diagonal lines indicate second inflection points. Vertical lines indicate actual values of antigens that correspond to the main inflection points and vertical broken lines indicate actual values of antigens that correspond to the second inflection points.

patients with and without relapse (Fig. 5). Thus, we set cut-off values as 1.9 and 2.9 log IU/mL for HBsAg and 3.0 and 4.0 log U/mL for HBcrAg in our model for predicting hepatitis relapse.

We tentatively defined three groups using the sum of the scores for HBsAg and HBcrAg levels at the time of NA discontinuation for our model. Conversions were made by assigning a score of 0 for an HBsAg level lower than 1.9 log IU/mL, 1 for a level from 1.9 to 2.8 log IU/mL, and 2 for a level equal to or higher than 2.9 log IU/mL. HBcrAg was scored as 0 for a level lower than 3.0 log U/mL, 1 for a level from 3.0 to 3.9 log U/mL, and 2 for a level equal to or higher than 4.0 log U/mL. Overall, group 1 consisted of patients with a total score of 0, group 2 of patients with a total score of 1 or 2, and group 3 of patients with a total score of 3 or 4.

Patients whose HBV DNA was lower than 3.0 log copies/mL and in whom HBeAg was negative at the time of NA discontinuation were assigned to one of the three groups. Figure 6 shows the comparison of non-relapse rates among the three groups using Kaplan–Meier analysis, which differed significantly. The non-relapse rate was approximately 90% in group 1, as low as 10% in

group 3, and intermediate in group 2. When factors associated with relapse were analyzed in group 3 patients, an age of over 40 years at the time of discontinuation was calculated as a significant factor (hazard



**Figure 6** Comparison of non-relapse rates using the Kaplan–Meier method among three groups classified by the sum of the scores of hepatitis B surface antigen (HBsAg) and hepatitis B core-related antigen (HBcrAg) levels at the time of nucleos(t)ide analog (NA) discontinuation.

ratio = 5.25, range 2.37–11.65,  $P < 0.001$ ). No significant factors were associated with relapse in group 2 patients.

## DISCUSSION

THE EUROPEAN ASSOCIATION for the Study of the Liver recommends continuation of NA treatment until HBsAg is cleared.<sup>25</sup> Liu *et al.* came to a similar conclusion in their study of chronic hepatitis B patients treated with LVD.<sup>14</sup> Indeed, the clearance of HBsAg is a reliable marker for the safe discontinuation of NAs, but the rate of patients who can clear HBsAg is relatively low (1–3%/year).<sup>26–28</sup> Thus, additional factors associated with relapse of hepatitis B after discontinuation of NAs were analyzed in the present study to better identify candidates who could achieve drug-free status. Such studies are relatively few, possibly because patients who discontinue NAs prematurely often experience severe complicating relapse and hepatic failure.<sup>9</sup> Although prospective studies are desirable to obtain accurate results, retrospective studies, such as ours, are also necessary to minimize the risk of adverse complications.

Since HBV cannot be completely eradicated in hosts, the primary goal in treating chronic hepatitis B is to convert symptomatic patients into inactive carriers in whom HBeAg is negative (usually anti-HBe-positive), serum HBV DNA is low, and serum ALT is normal.<sup>1,2,18,29</sup> Thus, we set the clinical conditions of a successful discontinuation of NAs as serum HBV DNA level below 4.0 log copies/mL and ALT below 30 IU/L following NA cessation. Patients who satisfy these conditions are not recommended for treatment by the Japanese guidelines for hepatitis B,<sup>18</sup> and it is also widely accepted that the risk of developing cirrhosis or complicating hepatocellular carcinoma is very low in such patients.<sup>30,31</sup> We used our cohort's mean and maximal values of HBV DNA and ALT for relapse analyses. Mean values were useful for evaluating relapse of hepatitis as a whole since parameter levels often fluctuated after discontinuation, and maximal values were used to evaluate relapse in a real-time fashion during the follow-up period. It is noteworthy that the mean and maximal values correlated very closely for both HBV DNA and ALT. The mean HBV DNA value of 4.0 log copies/mL corresponded to the maximal HBV DNA value of 5.7 by ROC analysis, and similarly the mean ALT value of 30 IU/L corresponded to the maximal ALT value of 79 IU/L. Thus, relapse of hepatitis B was judged to occur when serum ALT became higher than 79 IU/L or when serum HBV DNA surpassed 5.7 log copies/mL after the time of NA discontinuation.

Such criteria may also be useful for physicians to detect relapse at an early phase and avoid the occurrence of severe reactivation or unnecessary discontinuation of NAs.

It is generally understood that patients with a higher level of HBV DNA at the time of NA discontinuation are likely to relapse, but this cut-off value has not been analyzed sufficiently. Our findings using ROC analysis showed that patients with levels lower than 3.0 log copies/mL have a good possibility to achieve successful discontinuation. The presence of HBeAg is also generally accepted as a reliable factor to predict relapse of hepatitis. Our study showed that patients with detectable HBeAg at the time of NA discontinuation were likely to relapse, even if their HBV DNA levels were lower than 3.0 log copies/mL. Therefore, we next analyzed additional factors associated with a relapse of hepatitis after discontinuation of NAs by selecting patients who met both of these criteria.

Nucleos(t)ide analog treatment produces a rapid decrease in serum HBV DNA by suppressing reverse transcription of pregenomic HBV RNA. However, the key intrahepatic HBV replicative intermediate, covalently closed circular DNA (cccDNA), tends to remain and is capable of reinitiating replication once NAs are ceased.<sup>32</sup> Measurement of HBV cccDNA has been reported to be useful for monitoring and predicting responses to antiviral treatments.<sup>33</sup> However, its measurement is difficult in the clinical setting as it requires a liver biopsy. Due to the mechanism of action of NAs mentioned above, serum HBV DNA does not reflect intrahepatic HBV cccDNA in patients undergoing NA treatment.<sup>34</sup> To address this, quantitative measurement of HBV antigens has been reported to be useful for predicting the effect of antiviral treatment in patients with chronic hepatitis B. Although HBsAg is usually used as a serum marker for the diagnosis of HBV infection, several groups have shown that HBsAg levels can also be reflective of the response to peg-interferon in chronic hepatitis B.<sup>28,35,36</sup> The HBcrAg assay measures serum levels of HB core and e antigens simultaneously using monoclonal antibodies that recognize the common epitopes of these two denatured antigens. Since the assay measures all antigens transcribed from the pre-core/core gene, it is regarded as core-related.<sup>37</sup> Serum HBcrAg has been reported to accurately reflect intracellular levels of HBV cccDNA even during NA treatment,<sup>24,34,38</sup> and was found to be useful for identifying patients who were likely to show relapse of hepatitis after the discontinuation of NAs.<sup>39,40</sup> It is possible that levels of HBsAg and HBcrAg have different roles in

monitoring antiviral effects because the transcription of these two antigens are regulated by alternative enhancer-promoter systems in the HBV genome.<sup>3</sup> Therefore, we analyzed both of these antigens to elucidate their ability to predict relapse of hepatitis after discontinuation of NAs.

Multivariate analysis demonstrated that levels of HBsAg and HBcrAg at the time of NA discontinuation were independent factors significantly associated with relapse of hepatitis. Thus, we believe these factors can also be applied for predicting relapse in patients whose HBV DNA is lower than 3.0 log copies/mL and whose HBeAg is negative at NA discontinuation. HBV DNA levels were further analyzed using a highly sensitive assay based on real-time polymerase chain reaction (PCR). However, even the level of a negative signal did not ensure successful discontinuation of NAs. The results obtained here indicate that the combined use of HBV-related antigens are useful makers for monitoring the effect of anti-viral treatment in ways different from HBV DNA. Finally, since prolonged NA administration was also a significant factor associated with safe discontinuation, physicians are advised to continue patient treatment for at least 16 months for the best possible outcome.

From our data, a tentative model for predicting relapse of hepatitis after discontinuation of NAs was constructed using levels of HBsAg and HBcrAg at discontinuation. A negative result for HBeAg and HBV DNA lower than 3.0 log copies/mL at the time of NA discontinuation are the essential conditions in this system. Levels of HBsAg and HBcrAg were each converted into scores from 0 to 2 partly because two cut-off values were needed for each antigen and partly because a scoring system may be more convenient for clinical use. The sum of the two scores, which ranged from 0 to 4, was used to prospect relapse. We found that group 1 patients who had a low score (0) could be recommended to discontinue NAs because nearly 90% of this group achieved successful discontinuation. Further analysis of factors associated with relapse are needed for group 2 patients who had middle range scores (1 or 2), since the odds of achieving successful discontinuation were approximately 50%. Continuation of NA treatment is recommended for group 3 patients having high scores (3 or 4) because nearly 90% of this group relapsed. However, this recommendation may be reconsidered in patients younger than 40 years; such cases tended to have a lower relapse rate in group 3. It is also noteworthy that relapse occurred mainly during the first and second years following NA discontinuation in

all groups, similarly to a report by Liu *et al.*<sup>14</sup> Thus, clinicians should be vigilant in the early phase after discontinuation.

This study has several limitations. The patients who discontinued NAs were recruited retrospectively, and thus the decision to halt NA treatment was made by individual physicians without uniformly established criteria. Based on this, prospective studies are required to confirm our results. Furthermore, as over 90% of the patients we enrolled had genotype C and over 90% of cases were treated with LVD until discontinuation, the results obtained here can not be applied directly to other HBV genotypes or other types of NAs.

In conclusion, the present study showed that maximal levels of serum ALT and HBV DNA were useful for defining relapse patients after discontinuation of NAs. Along with serum HBV DNA of less than 3.0 log copies/mL and negative serum HBeAg, serum levels of HBsAg and HBcrAg at the time of NA discontinuation were able to predict relapse of hepatitis B and should therefore be considered when establishing uniform guidelines regarding the safe withdrawal of NA treatment. To this end, NA administration of more than 16 months is advisable to achieve successful discontinuation.

## ACKNOWLEDGMENTS

THIS RESEARCH WAS supported in part by a research grant from the Ministry of Health, Labor and Welfare of Japan.

We thank Ms. Hiroe Banno for her secretarial assistance and thank Ms. Nozomi Kamiyo and Ms. Etsuko Iigahama for their technical assistance. We also thank Mr Trevor Ralph for his English editorial assistance.

## REFERENCES

- 1 Hoofnagle JH, Doo E, Liang TJ, Fleischer R, Lok AS. Management of hepatitis B: summary of a clinical research workshop. *Hepatology* 2007; 45: 1056–75.
- 2 Lok AS, McMahon BJ. Chronic hepatitis B. *Hepatology* 2007; 45: 507–39.
- 3 Lee WM. Hepatitis B virus infection. *N Engl J Med* 1997; 337: 1733–45.
- 4 Ghany M, Liang TJ. Drug targets and molecular mechanisms of drug resistance in chronic hepatitis B. *Gastroenterology* 2007; 132: 1574–85.
- 5 Liaw YF, Sung JJ, Chow WC *et al.* Lamivudine for patients with chronic hepatitis B and advanced liver disease. *N Engl J Med* 2004; 351: 1521–31.
- 6 Matsumoto A, Tanaka E, Rokuhara A *et al.* Efficacy of lamivudine for preventing hepatocellular carcinoma in chronic

- hepatitis B: a multicenter retrospective study of 2795 patients. *Hepatol Res* 2005; 32: 173–84.
- 7 Suzuki Y, Kumada H, Ikeda K *et al.* Histological changes in liver biopsies after one year of lamivudine treatment in patients with chronic hepatitis B infection. *J Hepatol* 1999; 30: 743–8.
  - 8 Lok AS, Zoulim F, Locarnini S *et al.* Antiviral drug-resistant HBV: standardization of nomenclature and assays and recommendations for management. *Hepatology* 2007; 46: 254–65.
  - 9 Honkoop P, de Man RA, Niesters HG, Zondervan PE, Schalm SW. Acute exacerbation of chronic hepatitis B virus infection after withdrawal of lamivudine therapy. *Hepatology* 2000; 32: 635–9.
  - 10 Honkoop P, de Man RA, Heijtkink RA, Schalm SW. Hepatitis B reactivation after lamivudine. *Lancet* 1995; 346: 1156–7.
  - 11 Chang TT, Gish RG, de Man R *et al.* A comparison of entecavir and lamivudine for HBeAg-positive chronic hepatitis B. *N Engl J Med* 2006; 354: 1001–10.
  - 12 Lai CL, Shouval D, Lok AS *et al.* Entecavir versus lamivudine for patients with HBeAg-negative chronic hepatitis B. *N Engl J Med* 2006; 354: 1011–20.
  - 13 Lai CL, Chien RN, Leung NW *et al.* A one-year trial of lamivudine for chronic hepatitis B. Asia Hepatitis Lamivudine Study Group. *N Engl J Med* 1998; 339: 61–8.
  - 14 Liu F, Wang L, Li XY *et al.* Poor durability of lamivudine effectiveness despite stringent cessation criteria: a prospective clinical study in hepatitis B e antigen-negative chronic hepatitis B patients. *J Gastroenterol Hepatol* 2011; 26: 456–60.
  - 15 Reijnders JG, Perquin MJ, Zhang N, Hansen BE, Janssen HL. Nucleos(t)ide analogues only induce temporary hepatitis B e antigen seroconversion in most patients with chronic hepatitis B. *Gastroenterology* 2010; 139: 491–8.
  - 16 Liaw YF, Leung N, Kao JH *et al.* Asian-Pacific consensus statement on the management of chronic hepatitis B: a 2008 update. *Hepatol Int* 2008; 2: 263–83.
  - 17 Leung N. Recent data on treatment of chronic hepatitis B with nucleos(t)ide analogues. *Hepatol Int* 2008; 2: 163–78.
  - 18 Kumada H, Okanoue T, Onji M *et al.* Guidelines for the treatment of chronic hepatitis and cirrhosis due to hepatitis B virus infection for the fiscal year 2008 in Japan. *Hepatol Res* 2010; 40: 1–7.
  - 19 Schuttler CG, Wend UC, Faupel FM, Lelie PN, Gerlich WH. Antigenic and physicochemical characterization of the 2nd International Standard for hepatitis B virus surface antigen (HBsAg). *J Clin Virol* 2010; 47: 238–42.
  - 20 Dai CY, Yu ML, Chen SC *et al.* Clinical evaluation of the COBAS Amplicor HBV monitor test for measuring serum HBV DNA and comparison with the Quantiplex branched DNA signal amplification assay in Taiwan. *J Clin Pathol* 2004; 57: 141–5.
  - 21 Ronsin C, Pillet A, Bali C, Denoyel GA. Evaluation of the COBAS AmpliPrep-total nucleic acid isolation-COBAS TaqMan hepatitis B virus (HBV) quantitative test and comparison to the VERSANT HBV DNA 3.0 assay. *J Clin Microbiol* 2006; 44: 1390–9.
  - 22 Mizokami M, Nakano T, Orito E *et al.* Hepatitis B virus genotype assignment using restriction fragment length polymorphism patterns. *FEBS Lett* 1999; 450: 66–71.
  - 23 Kimura T, Rokuhara A, Sakamoto Y *et al.* Sensitive enzyme immunoassay for hepatitis B virus core-related antigens and their correlation to virus load. *J Clin Microbiol* 2002; 40: 439–45.
  - 24 Suzuki F, Miyakoshi H, Kobayashi M, Kumada H. Correlation between serum hepatitis B virus core-related antigen and intrahepatic covalently closed circular DNA in chronic hepatitis B patients. *J Med Virol* 2009; 81: 27–33.
  - 25 European Association for the Study of the Liver. EASL Clinical Practice Guidelines: management of chronic hepatitis B. *J Hepatol* 2009; 50: 227–42.
  - 26 Gish RG, Lok AS, Chang TT *et al.* Entecavir therapy for up to 96 weeks in patients with HBeAg-positive chronic hepatitis B. *Gastroenterology* 2007; 133: 1437–44.
  - 27 Marcellin P. Hepatitis B and hepatitis C in 2009. *Liver Int* 2009; 29 (Suppl 1): 1–8.
  - 28 Moucari R, Lada O, Marcellin P. Chronic hepatitis B: back to the future with HBsAg. *Expert Rev Anti Infect Ther* 2009; 7: 633–6.
  - 29 Yokosuka O, Kurosaki M, Imazeki F *et al.* Management of hepatitis B: consensus of the Japan Society of Hepatology 2009. *Hepatol Res* 2011; 41: 1–21.
  - 30 Iloeje UH, Yang HI, Su J, Jen CL, You SL, Chen CJ. Predicting cirrhosis risk based on the level of circulating hepatitis B viral load. *Gastroenterology* 2006; 130: 678–86.
  - 31 Chen CJ, Yang HI, Su J *et al.* Risk of hepatocellular carcinoma across a biological gradient of serum hepatitis B virus DNA level. *JAMA* 2006; 295: 65–73.
  - 32 Werle-Lapostolle B, Bowden S, Locarnini S *et al.* Persistence of cccDNA during the natural history of chronic hepatitis B and decline during adefovir dipivoxil therapy. *Gastroenterology* 2004; 126: 1750–8.
  - 33 Sung JJ, Wong ML, Bowden S *et al.* Intrahepatic hepatitis B virus covalently closed circular DNA can be a predictor of sustained response to therapy. *Gastroenterology* 2005; 128: 1890–7.
  - 34 Wong DK, Tanaka Y, Lai CL, Mizokami M, Fung J, Yuen MF. Hepatitis B virus core-related antigens as markers for monitoring chronic hepatitis B infection. *J Clin Microbiol* 2007; 45: 3942–7.
  - 35 Brunetto MR, Moriconi F, Bonino F *et al.* Hepatitis B virus surface antigen levels: a guide to sustained response to peginterferon alfa-2a in HBeAg-negative chronic hepatitis B. *Hepatology* 2009; 49: 1141–50.
  - 36 Moucari R, Mackiewicz V, Lada O *et al.* Early serum HBsAg drop: a strong predictor of sustained virological response

- to pegylated interferon alfa-2a in HBeAg-negative patients. *Hepatology* 2009; 49: 1151–7.
- 37 Tanaka E, Matsumoto A, Yoshizawa K, Maki N. Hepatitis B core-related antigen assay is useful for monitoring the antiviral effects of nucleoside analogue therapy. *Intervirology* 2008; 51 (Suppl 1): 3–6.
- 38 Hosaka T, Suzuki F, Kobayashi M *et al.* HBcrAg is a predictor of post-treatment recurrence of hepatocellular carcinoma during antiviral therapy. *Liver Int* 2010; 30: 1461–70.
- 39 Matsumoto A, Tanaka E, Minami M *et al.* Low serum level of hepatitis B core-related antigen indicates unlikely reactivation of hepatitis after cessation of lamivudine therapy. *Hepatol Res* 2007; 37: 661–6.
- 40 Shinkai N, Tanaka Y, Orito E *et al.* Measurement of hepatitis B virus core-related antigen as predicting factor for relapse after cessation of lamivudine therapy for chronic hepatitis B virus infection. *Hepatol Res* 2006; 36: 272–6.



