

three patients (30.0%). The moderate adverse events (one each) that developed in five patients were vertigo, peripheral oedema, nasopharyngitis, increase in blood uric acid and in low density lipoprotein, facial palsy and rash, whereas all other adverse events were mild. It is notable that although seven patients discontinued the therapy, none did so owing to adverse events.

#### Antiviral activity

Telaprevir rapidly decreased serum HCV RNA level in all patients enrolled in this study. The median serum HCV RNA level changed from 6.45 log<sub>10</sub> IU/mL (range: 5.1–7.1) just before administration to 4.00 log<sub>10</sub> IU/mL (range: 3.0–4.7) at 16 h after administration and 1.10 log<sub>10</sub> IU/mL (range: 0.5–3.3) on Day 14 (Fig. 1). Telaprevir showed potent antiviral activity: a median log<sub>10</sub> decrease of 2.325 at 16 h and 5.175 on Day 14. During the administration period of 12 weeks, HCV RNA levels decreased to less than the LLOQ of 1.2 log<sub>10</sub> IU/mL in seven patients, and three patients achieved HCV RNA negativity on Day 14 or Day 29. After the decrease in serum HCV RNA, breakthrough occurred in eight patients, and seven of those patients discontinued the trial during the dosing period (from Day 45 to Day 63, Table 1). In addition, one of the remaining three patients who completed the administration of the study drug achieved virus negativity by the end of administration

(Day 86), but relapsed 1 week after completion of drug therapy.

#### Hepatocyte injury markers

As shown in Fig. 2a, the alanine aminotransferase (ALT) and aspartate aminotransferase (AST) levels decreased during and after telaprevir treatment. Median changes from baseline (Day 1) in ALT and AST were –26.5 IU/L (range: –217–5, *N* = 10) and –8.5 IU/L (range: –118–2, *N* = 10) on Day 29, respectively. Fig. 2b shows total bilirubin levels. No clinically significant change in bilirubin was observed in all patients. These data indicate that long-term exposure to telaprevir caused neither damage nor injury in the liver.

#### Sequence analysis of hepatitis C virus NS3

Amino acid substitutions in the NS3 protease domain, which were selected by telaprevir administration, were examined in 39 clones or more for each sample (Table 3). The predominant variants detected during the early time points after administration (on Days 3 and 8) were V36G, T54A and A156V. Subsequently, these variants decreased below the LOD in nine patients, and the predominant variants detected at viral breakthrough after Week 6 of administration (Day 43–86) were single-substituted variants of A156F/T/V and multiple-substituted variants of T54S+A156T and




Fig. 1 Changes in patient hepatitis C virus (HCV) RNA level. For 12 consecutive weeks, all 10 patients received 750 mg telaprevir q8h under feeding conditions. <1.2 LLOQ, below lower limit of quantification of 1.2 log<sub>10</sub> IU/mL; FU, follow-up.




Fig. 2 Change in alanine aminotransferase, aspartate aminotransferase (a) and total bilirubin (b) levels. FU, follow-up.

**Table 3** Means of hepatitis C virus (HCV) RNA levels and representation rates of variants in all subjects during and after telaprevir treatment

	Pre	Day 3	Day 8	Day 14	Day 29	Day 43	Day 50–60	Day 86	FU2W	FU4W	FU8W	FU12W	FU24W
N	10	10	10	10	10	10	9	3	3	3	2	2	2
Mean of HCV RNA level ( $\log_{10}$ IU/mL)	6.35	2.61	1.84	1.45	1.56	2.53	3.22	2.40	2.90	4.13	6.60	6.45	6.45
HCV NS3 variants (%)													
Wild	100.0	40.0	0.2	–	0.2	0.1	–	–	–	18.3	86.4	51.8	97.8
V36A	–	–	–	–	–	–	0.3	–	–	23.8	3.4	38.5	1.1
V36G	–	10.0	0.4	–	2.4	0.8	–	–	–	–	–	–	–
T54A	–	–	9.4	9.5	4.7	0.1	–	–	–	17.9	1.1	5.0	–
A156F	–	–	–	–	–	10.0	25.5	0.8	–	–	–	–	–
A156T	–	–	–	0.5	–	7.5	16.6	31.1	16.3	–	–	1.2	–
A156V	–	–	30.0	–	1.1	15.9	2.3	–	–	–	–	–	–
T54S+A156S	–	–	–	–	–	–	–	–	–	19.2	3.4	1.3	–
T54S+A156T	–	–	–	–	–	9.6	11.4	1.5	16.3	15.0	–	–	–
A156T+V158I	–	–	–	–	–	3.3	10.1	–	–	–	–	–	–

–, not detected; FU, follow-up. Minor substitutions (maximum occupancy in a specimen was less than 10%): T54S, R155G, R155L, A156S, V36A+T54A, V36A+A156S, V36G+A156V, T54A+R155L, T54A+A156S, T54A+A156V, T54S+R155L, T54S+A156V, T54A+V132L, A156S+V132L, T54A+V163I, T54S+A156T+V158I, V36A+T54A+A156S

A156T+V158I; no wild-type virus was detected. In the three patients who completed the administration of telaprevir for 12 weeks, V36A, T54A and T54S+A156S/T were detectable after treatment. In the two patients followed up for 24 weeks, gradual enrichment of the wild-type viruses was observed.

*Pharmacokinetics*


The plasma concentration vs time curves on Days 1, 14 and 85 are shown in Fig. 3a and the  $C_{trough}$  on Days 1, 2, 3, 8, 14, 15, 29, 43, 57 and 85 in Fig. 3b. The pharmacokinetic parameters of telaprevir on Days 1, 14 and 85 are given in Table 4.

As the  $t_{max}$  were similar on Days 1, 14 and 85 with medians of 2.50, 2.49 and 2.72 h, respectively, the repeated

administration under the present conditions was unlikely to cause any change in absorption. The pharmacokinetic parameters of  $C_{max}$ ,  $AUC_{0-8 h}$  and  $C_{trough}$  were lower on Day 1 than those on Days 14 and 85; thus, on Days 1, 14 and 85, the mean values of  $C_{max}$  were respectively 2.24, 3.34 and 3.68  $\mu\text{g/mL}$ , the mean values of  $AUC_{0-8 h}$  were respectively 11.60, 22.31 and 23.98  $\mu\text{g}\cdot\text{h/mL}$ , and the mean values of  $C_{trough}$  at 8 h after the first administration were respectively 1.462, 2.239 and 2.312  $\mu\text{g/mL}$ . The plasma concentration of telaprevir reached steady state on Day 2.

DISCUSSION

During the past decade, the combined use of PEG-IFN and RBV has provided a significant therapeutic advance for



**Fig. 3** Time course of plasma concentration (a) and  $C_{trough}$  (b) of telaprevir. Symbols and error bars indicate mean values and SD, respectively.

Table 4 Pharmacokinetic parameters of plasma telaprevir

	N	$C_{\max}$ ( $\mu\text{g}/\text{mL}$ )	$t_{\max}$ (h) <sup>*</sup>	$\text{AUC}_{0-8\text{ h}}$ ( $\mu\text{g}\cdot\text{h}/\text{mL}$ )	$C_{\text{trough}}$ ( $\mu\text{g}/\text{mL}$ ) <sup>†</sup>	$t_{1/2}$ (h)
Day 1	10	2.24 $\pm$ 0.93	2.50 (2.30–7.92)	11.60 $\pm$ 4.74	1.462 $\pm$ 0.949	5.57 $\pm$ 2.67 <sup>‡,§</sup>
Day 14	10	3.34 $\pm$ 1.11	2.49 (0.98–5.97)	22.31 $\pm$ 8.29	2.239 $\pm$ 0.953	9.64 $\pm$ 6.14 <sup>‡,¶</sup>
Day 85	3	3.68 $\pm$ 1.29	2.72 (2.68–4.00)	23.98 $\pm$ 9.45	2.312 $\pm$ 1.265	18.35 $\pm$ 22.91 <sup>**</sup>

Mean value  $\pm$  SD. <sup>\*</sup>Median (minimum value to maximum value). <sup>†</sup> $C_{\text{trough}}$  at 8 h after the first administration. <sup>‡</sup>Calculated from measured values at 8 h after the first administration. <sup>§</sup> $N = 7$ . <sup>¶</sup> $N = 8$ . <sup>\*\*</sup>Calculated from measured values at 24 h after the first administration.

patients with CHC. Approximately 50% of patients infected with genotype 1 HCV do not, however, achieve SVR with this SOC [3–5]. On the contrary, the treatment with telaprevir-based triple regimen significantly improved SVR rates in patients with genotype 1 HCV. The PROVE 1 and 2 studies of telaprevir use with PEG-IFN and RBV in treatment-naïve patients achieved SVR rates of 61% and 69% (placebo: 41–46%) [11,12]. The Japanese study of the telaprevir-based triple regimen also showed high SVR rates [13–15]. However, the key safety concerns with the telaprevir-based triple regimen were anaemia, rash and IFN-induced systemic symptoms, all of which were most likely caused by the PEG-IFN/RBV treatment. In Japan, there are currently a large number of aged people with genotype 1b HCV and high viral loads, which is one of the most intractable HCV genotypes. As a result of advanced age, many subjects could not tolerate the adverse drug reactions in the telaprevir-based regimen, which was also observed with PEG-IFN/RBV therapies [13,15]. This observation prompted us to re-examine the safety profiles and pharmacokinetics of monotherapy with telaprevir for 12 weeks in Japanese patients.

In this study, 10 treatment-naïve patients with genotype 1b CHC and a high median viral load of 6.45  $\log_{10}$  IU/mL (range: 5.10–7.10) (Table 1) took 750 mg of telaprevir q8h for 12 weeks under feeding conditions. The plasma concentrations of telaprevir reached steady state within 2 days after the initiation of administration in the 750-mg q8h regimen as is shown by the constant  $C_{\text{trough}}$  from Day 2 to Day 85; hence, all the patients enrolled in this study were sufficiently exposed to telaprevir during treatment (Fig. 3b). These results demonstrate that the plasma concentrations of telaprevir were manageable even during the long-term repeated administration. There were no clinically significant events, although the incidence of some events exceeded 20.0% (Table 2). Notably, mild anaemia developed in seven patients (70%) and its occurrence was consistent with the decrease in haemoglobin values, although gradual, during the first 29 days after administration of telaprevir. The incidence of rash, which is reported to develop with a high incidence and high severity in the clinical trials of co-administration of telaprevir with PEG-IFN and RBV [11,12], was also high but its severity was mild in this study. Although exposure to telaprevir was sufficient to eliminate

the virus, neither serious adverse events nor discontinuations because of adverse events occurred during the study period. The results confirmed the high tolerability of telaprevir alone after long-term administration. Although there has been no direct comparison of telaprevir monotherapy and telaprevir-based triple therapy, based on these results, the severe adverse drug reactions reported for telaprevir-based triple therapy including anaemia and rash were likely to be ascribed to the synergistic and/or additive effects of the three drugs, i.e., telaprevir, PEG-IFN, and RBV. The safety information under telaprevir monotherapy described here is very important to understand the aspects of adverse drug reactions, especially anaemia and rash, in telaprevir-based triple therapy. In addition, compared to baseline, the ALT and AST levels were significantly lower during the treatment in all patients, indicating that telaprevir was unlikely to cause direct liver damage or injury even after long-term use.

Although there is a report on HCV RNA mutation after monotherapy with a protease inhibitor for 14 days [16], no information about the selective pressure of such protease inhibitors administered alone for a longer period is available at present. During the treatment period in this study, HCV RNA levels were below the LLOQ in seven patients and undetectable in three patients. Importantly, one patient showed an end-of-treatment response. Viral breakthrough resulting from the selection of Ala<sup>156</sup>-substituted variants with high-level resistance to telaprevir [16] occurred in eight patients. It has been reported that high-level resistance was absent, low-level resistance was minimized, and the majority of the viral population reverted to the wild-type by 3–7 months after telaprevir dosing for 14 days [16]. In the two patients who were studied up to the last visit, enrichment of the wild-type viruses was observed at Week 24 of the follow-up period. It is thus clear that the variants that appeared during prolonged administration of telaprevir for 12 weeks could be replaced by or could revert to the wild-type viruses. This study also provides new knowledge about a selective pathway of the NS3 protease domain of HCV genotype 1b during long-term telaprevir administration (Table 3). It is notable that the wild-type viruses were eliminated promptly by Day 3 of telaprevir monotherapy in all cases, but variants with amino acid substitutions such as V36G, A156V and T54A still remained on Days 3 and 8.

From Day 50 to Day 99, A156T was the predominant variant after viral breakthrough. On Day 43, several substitutions that are rarely reported were found: a single substitution of A156F and multiple substitutions of T54S+A156T and A156T+V158I. In the clonal sequencing analysis in this trial, the observed T54S and V158I substitutions were mostly associated with the A156S/T substitution, and enrichment of multiple-substituted variants was observed under prolonged telaprevir treatment (Fig. S1). A phenotypic enzyme assay suggested that the solo T54S substitution did not change the inhibitory concentration of telaprevir (data not shown). It has also been reported that the T54S and V158I substitutions were also positively selected in the clinical trials of boceprevir, but the solo V158I substitution did not confer telaprevir resistance [17]. Therefore, these two substitutions may be a secondary resistance-associated variant of genotype 1b. Moreover, we could speculate that these variants are susceptible to PEG-IFN and RBV, because the viral variants emerging after the longer selective pressure with telaprevir monotherapy were decreased rapidly by switching the treatment with telaprevir to that with PEG-IFN and RBV [18]. Although it was reported that one patient with low viral load achieved SVR in the treatment regimen in which 750 mg telaprevir was administered q8h for 24 weeks [19], no patients with high viral load achieved SVR in this study. As discussed earlier,

PEG-IFN and RBV-free therapy is an unmet and strong medical need in Japan. Therefore, an oral cocktail therapy for HCV genotype 1b infection using telaprevir and different types of DAAs, for example HCV NS5A or NS5B polymerase inhibitors, would be warranted to improve efficacy and reduce adverse drug reactions of the telaprevir, PEG-IFN and RBV triple therapy.

In conclusion, the results of this study indicate that telaprevir is well tolerated at 750 mg q8h for 12 weeks in Japanese patients with HCV genotype 1b infection. The data obtained in this study on telaprevir monotherapy demonstrate that the severe side effects, rash and anaemia observed in the telaprevir-based triple regimen were likely to be attributable to the additive and/or synergistic effect of telaprevir, PEG-IFN and RBV, and this consideration has encouraged us to evaluate telaprevir in a combination therapy with a different class of DAAs in future.

#### ACKNOWLEDGEMENT

Yamada, Kamiya, Aoki, Sakurai, Kano and Matsui are employees of Mitsubishi Tanabe Pharma Corporation.

#### DISCLOSURE

The others have nothing to declare.

#### REFERENCES

- Liang TJ, Rehermann B, Seeff LB, Hoofnagle JH. Pathogenesis, natural history, treatment and prevention of hepatitis C. *Ann Intern Med* 2000; 132: 296–305.
- Seeff LB. Natural history of chronic hepatitis C. *Hepatology* 2002; 36: S35–S36.
- Manns MP, McHutchison JG, Gordon SC *et al.* Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomised trial. *Lancet* 2001; 358: 958–965.
- Fried MW, Shiffman ML, Reddy KR *et al.* Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 2002; 347: 975–982.
- Akuta N, Suzuki F, Kawamura Y *et al.* Predictive factors of early and sustained responses to peginterferon plus ribavirin combination therapy in Japanese patients infected with hepatitis C virus genotype 1b: amino acid substitutions in the core region and low-density lipoprotein cholesterol levels. *J Hepatol* 2007; 46: 403–410.
- Perni RB, Almquist SJ, Byrn RA *et al.* Preclinical profile of VX-950, a potent, selective, and orally bio-available inhibitor of hepatitis C virus NS3-4A serine protease. *Antimicrob Agents Chemother* 2006; 50: 899–909.
- Reesink HW, Zeuzem S, Weegink CJ *et al.* Rapid decline of viral RNA in hepatitis C patients treated with VX-950: a phase Ib, placebo-controlled, randomized study. *Gastroenterology* 2006; 131: 997–1002.
- Kiffer TL, Sarrazin C, Killer JS *et al.* Telaprevir and pegylated interferon-alpha-2a inhibit wild-type and resistant genotype 1 hepatitis C virus replication in patients. *Hepatology* 2007; 46: 631–639.
- Kumada T, Toyoda H, Honda T *et al.* Treatment of chronic hepatitis C with interferon alone or combined with ribavirin in Japan. *Intervirology* 2006; 49: 112–118.
- Yoshizawa H, Tanaka J, Miyakawa Y. National prevention of hepatocellular carcinoma in Japan based on epidemiology of hepatitis C virus infection in the general population. *Intervirology* 2006; 49: 7–17.
- McHutchison JG, Everson GT, Gordon SC *et al.* Telaprevir with peginterferon and ribavirin for chronic HCV genotype 1 infection. *N Engl J Med* 2009; 360: 1827–1838.
- Hézode C, Forestier N, Dusheiko G *et al.* Telaprevir and peginterferon with or without ribavirin for chronic HCV infection. *N Engl J Med* 2009; 360: 1839–1850.
- Suzuki F, Akuta N, Suzuki Y *et al.* Rapid loss of hepatitis C virus genotype 1b from serum in patients receiving a triple treatment with telaprevir (MP-424), pegylated interferon and ribavirin for 12 weeks. *Hepatol Res* 2009; 39: 1056–1063.
- Akuta N, Suzuki F, Hiraoka M *et al.* Amino acid substitution in hepatitis C virus core region and genetic variation near the interleukin 28B gene predict viral response to telaprevir with peginterferon and ribavirin. *Hepatology* 2010; 52: 421–429.

- 15 Suzuki F, Suzuki Y, Akuta N *et al*. Influence of ITPA polymorphisms on decreases of hemoglobin during treatment with pegylated interferon, ribavirin, and telaprevir. *Hepatology* 2011; 53: 415–421.
- 16 Sarrazin C, Kiffer TL, Bartels D *et al*. Dynamic hepatitis C virus genotypic and phenotypic changes in patients treated with the protease inhibitor telaprevir. *Gastroenterology* 2007; 132: 1767–1777.
- 17 Qiu P, Sanfilippo V, Curry S *et al*. Identification of HCV protease inhibitor resistance mutations by selection pressure-based method. *Nucleic Acids Res* 2009; 37: e74.
- 18 Ozeki I, Akaike J, Karino Y *et al*. Antiviral effects of peginterferon alpha-2b and ribavirin following 24-week monotherapy of telaprevir in Japanese hepatitis C patients. *J Gastroenterol* 2011; 46: 929–937.
- 19 Suzuki F, Suzuki Y, Akuta N *et al*. Sustained virological response in a patient with chronic hepatitis C treated by monotherapy with the NS3-4A protease inhibitor telaprevir. *J Clin Virol* 2010; 47: 76–78.

## SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article:

**Figure S1:** Pie chart of variant occupation at each time point in the

typical two cases. Proposed secondary resistant associated substitutions are underlined.

Please note: Wiley-Blackwell are not responsible for the content or func-

tionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.



sons, those patients who do not achieve SVR need to be identified, so as to free them of unnecessary side effects and reduce costs, preferably before the start of the combination therapy.

Viral- and host-related factors are useful as predictors of treatment efficacy to 48-week IFN/ribavirin combination therapy. With regard to viral factors, amino acid (aa) substitutions at position 70 and/or 91 in the core region of HCV-1b are pretreatment predictors of virological response to combination therapy [1–4], and also affect clinical outcome, including hepatocarcinogenesis [5, 6]. Furthermore, the NS5A region of HCV-1b, including IFN-sensitivity-determining region (ISDR) [7, 8] and IFN/ribavirin resistance-determining region (IRRDR) [9, 10], are also useful as pretreatment predictors of virological response to combination therapy [11, 12]. With regard to host factors, genetic variations near *IL28B* gene (rs8099917, rs12979860) on chromosome 19, which encodes IFN- $\lambda$ -3, are pretreatment predictors of virological response to combination therapy in individuals infected with HCV-1 [13–16], and also affect clinical outcome, including spontaneous clearance of HCV [17]. A recent report identified genetic variation near *IL28B* gene and aa substitution of the core region as predictors of SVR to triple therapy of telaprevir/pegylated (PEG)-IFN/ribavirin in Japanese patients infected with HCV-1b [18]. However, to our knowledge, there are no previous reports of IFN/ribavirin combination therapy based on multivariate analysis to investigate pretreatment predictors, including all of aa substitutions in core aa 70/91 and NS5A-ISDR/IRRDR, and genetic variation near *IL28B* gene.

The aim of the present study was to investigate predictive factors of treatment efficacy, including viral- (aa substitutions in core aa 70/91 and NS5A-ISDR/IRRDR) and host-related factors (genetic variation near *IL28B* gene), to 48-week IFN/ribavirin in Japanese adults infected with HCV-1b.

## Patients and Methods

### Study Population

A total of 1,249 HCV-1b-infected Japanese adult patients were consecutively recruited into the study protocol of combination therapy with IFN (PEG-IFN $\alpha$ -2b or IFN $\alpha$ -2b) plus ribavirin between December 2001 and January 2009 at Toranomon Hospital, Tokyo, Japan. Among these, 490 patients, who could complete a total of 48 weeks of combination therapy, were enrolled in this retrospective study, and fulfilled the following criteria: (1) negativity for hepatitis B surface antigen (HBsAg) in serum; (2) HCV-1b only confirmed by sequence analysis; (3) HCV-RNA levels of  $\geq 5.0$  log IU/ml determined by the COBAS TaqMan HCV test

(Roche Diagnostics, Tokyo, Japan) within the preceding 2 months of enrolment; (4) no hepatocellular carcinoma; (5) body weight  $>40$  kg; (6) lack of coinfection with human immunodeficiency virus; (7) no previous treatment with antiviral or immunosuppressive agents within the preceding 3 months of enrolment; (8) none was an alcoholic; lifetime cumulative alcohol intake was  $<500$  kg; (9) none had other forms of liver diseases, such as hemochromatosis, Wilson disease, primary biliary cirrhosis, alcoholic liver disease, or autoimmune liver disease, and (10) none of the females was pregnant or breastfeeding.

The study protocol was in compliance with the Good Clinical Practice Guidelines and the 1975 Declaration of Helsinki, and was approved by the institutional review board. Each patient gave their informed consent before participating in this trial.

The treatment efficacy was evaluated in terms of HCV-RNA negativity at the end of treatment (end-of-treatment response (ETR)) and 24 weeks after the completion of therapy (SVR), based on the COBAS TaqMan HCV test (Roche Diagnostics). SVR in patients who achieved ETR was defined as SVR after ETR. ETR, SVR, and SVR after ETR could be evaluated in 487 (99%), 448 (91%), and 321 (66%) of 490 patients, respectively.

422 (86%) patients received PEG-IFN $\alpha$ -2b at a median dose of 1.4  $\mu$ g/kg (range 0.7–1.9) subcutaneously each week plus oral ribavirin at a median dose of 11.1 mg/kg (range 3.7–15.1) daily for 48 weeks. The remaining 68 (14%) patients received 6 million units of IFN $\alpha$ -2b intramuscularly each day for 48 weeks (daily for the initial 2 weeks, followed by three times per week for 46 weeks), and oral ribavirin at a median dose of 11.3 mg/kg (range 6.8–13.4) daily for 48 weeks.

Table 1 summarizes the profiles and laboratory data of the 490 patients at the commencement of treatment. They included 310 males and 180 females aged 20–75 years (median 54).

### Measurement of HCV RNA

The antiviral effects of treatment on HCV were assessed by measuring plasma HCV-RNA levels. In this study, HCV-RNA levels were evaluated at least once every month before, during, and after therapy. HCV-RNA concentrations were determined using the COBAS TaqMan HCV test (Roche Diagnostics). The linear dynamic range of the assay was 1.2–7.8 log IU/ml, and the undetectable samples were defined as negative.

### Detection of aa Substitutions in Core, and NS5A Regions of HCV-1b

With the use of HCV-J (accession No. D90208) as a reference [19], the sequence of 1–191 aa in the core protein of HCV-1b was determined and then compared with the consensus sequence constructed on the previous study to detect substitutions at aa 70 of arginine (Arg70) or glutamine/histidine (Gln70/His70) and aa 91 of leucine (Leu91) or methionine (Met91) [1]. The sequence of 2,209–2,248 aa in the NS5A of HCV-1b (ISDR) reported by Enomoto et al. [7, 8] was determined, and the number of aa substitutions in ISDR was defined as wild-type (WT) (0, 1) or non-wild-type (non-WT) ( $\geq 2$ ) in comparison with HCV-J. Furthermore, the sequence of 2,334–2,379 aa in the NS5A of HCV-1b (IRRDR) reported by El-Shamy et al. [9, 10] was determined and then compared with the consensus sequence constructed on the previous study. In the present study, aa substitutions of the core region and NS5A-ISDR/IRRDR of HCV-1b were analyzed by direct sequencing [10, 18].



### Genetic Variation near *IL28B* Gene

Samples for genome-wide association survey were genotyped using the Illumina HumanHap610-Quad Genotyping BeadChip. Genotyping data were subjected to quality control before the data analysis. Genotyping for replication and fine mapping was performed by use of Invader assay, TaqMan assay, or direct sequencing as described previously [20, 21].

In this study, genetic variations near *IL28B* gene (rs8099917), reported as the pretreatment predictors of treatment efficacy in Japanese patients [14, 18], were investigated.

### Statistical Analysis

Non-parametric tests (Mann-Whitney U test,  $\chi^2$  test and Fisher's exact probability test) were used to compare the characteristics of the groups. Correlation analysis was evaluated by the Spearman rank correlation test. Uni- and multivariate logistic regression analyses were used to determine those factors that significantly contributed to ETR, SVR, and SVR after ETR. The odds ratios (OR) and 95% confidence intervals (95% CI) were also calculated. All *p* values <0.05 by the two-tailed test were considered significant. Variables that achieved statistical significance (*p* < 0.05) on univariate analysis were entered into multiple logistic regression analysis to identify significant independent predictive factors. Each variable was transformed into categorical data consisting of two simple ordinal numbers for uni- and multivariate analyses. Potential predictive factors associated with ETR, SVR, and SVR after ETR included the following variables: sex, age, history of blood transfusion, familial history of liver disease, body mass index, aspartate aminotransferase (AST), alanine aminotransferase (ALT), albumin,  $\gamma$ -glutamyl transpeptidase (GGT), leukocyte count, hemoglobin, platelet count, level of viremia,  $\alpha$ -fetoprotein, total cholesterol, high-density lipoprotein cholesterol, low-density lipoprotein cholesterol, triglycerides, uric acid, ribavirin dose/body weight, genetic variation near *IL28B* gene, and aa substitution in the core region, and NS5A-ISDR/IRRDR. Statistical analyses were performed using SPSS software (SPSS Inc., Chicago, Ill., USA).

## Results

### Response to Therapy

ETR was achieved by 372 of 487 (76%) patients, SVR by 244 of 448 (54%), and SVR after ETR by 244 of 321 (76%).

### Number of aa Substitutions in NS5A-ISDR and NS5A-IRRDR

As a whole, 0, 1, and  $\geq 2$  aa substitutions in ISDR were found in 56% (227 of 406), 23% (95 of 406), and 21% (84 of 406) of patients, respectively. Thus, the percentage of patients with  $\leq 1$  aa substitution in ISDR (WT) was 79% (322 of 406). Furthermore,  $\leq 3$ , 4–5, and  $\geq 6$  aa substitutions in IRRDR were found in 36% (73 of 200), 34% (67 of 200), and 30% (60 of 200) of patients, respectively (fig. 1).

**Table 1.** Patient profile and laboratory data at commencement of the 48-week combination therapy of IFN + ribavirin in 490 patients infected with HCV-1b

<i>Demographic data</i>	
Number of patients	490
Male/female	310/180
Age, years	54 (20–75)
History of blood transfusion	169 (34%)
Family history of liver disease	96 (20%)
Body mass index, kg/m <sup>2</sup>	22.6 (15.7–34.7)
<i>Laboratory data</i>	
Level of viremia, log IU/ml	6.4 (2.2–7.7)
Serum AST, IU/l	50 (16–296)
Serum ALT, IU/l	67 (12–836)
Serum albumin, g/dl	3.9 (3.1–4.7)
GGT, IU/l	44 (10–592)
Leukocyte count, n/mm <sup>3</sup>	4,700 (1,200–10,900)
Hemoglobin, g/dl	14.4 (10.6–18.1)
Platelet count, $\times 10^4$ /mm <sup>3</sup>	16.7 (6.4–37.5)
$\alpha$ -Fetoprotein, $\mu$ g/l	5 (1–459)
Total cholesterol, mg/dl	170 (96–284)
High-density lipoprotein cholesterol, mg/dl	46 (13–95)
Low-density lipoprotein cholesterol, mg/dl	100 (32–190)
Triglycerides, mg/dl	90 (33–416)
Uric acid, mg/dl	5.5 (2.3–9.4)
<i>Treatment</i>	
PEG-IFN $\alpha$ -2b/IFN $\alpha$ -2b	422/68
Ribavirin dose, mg/kg	11.2 (3.7–15.1)
<i>aa substitutions in the HCV-1b</i>	
Core aa 70, arginine/glutamine (histidine)	266/151
Core aa 91, leucine/methionine	246/169
ISDR of NS5A, 0/1/ $\geq 2$	227/95/84
IRRDR of NS5A, $\leq 3/4-5/\geq 6$	73/67/60
<i>Genetic variation near IL28B gene</i>	
rs8099917 genotype, TT/TG/GG	150/65/4


Data represent number of patients with percentages in parentheses, or median (range) values.

The correlation between ISDR and IRRDR was analyzed. There was a significant positive correlation between the number of aa substitutions in ISDR and those in IRRDR ( $r = 0.308$ ,  $p < 0.001$ ) (fig. 2).


### aa Substitutions in the Core Region and NS5A-ISDR/IRRDR

Concerning the substitution of core aa 70, the number of aa substitutions in ISDR of 256 patients with Arg70 (median 0) was not significantly different from that of 146 patients with Gln70 (His70) (median 0) (fig. 3a). Fur-





**Fig. 1.** The number of aa substitutions in NS5A-ISDR and NS5A-IRRDR. The percentage of patients with  $\leq 1$  aa substitution in ISDR (WT) was 79%.



**Fig. 2.** Correlation between NS5A-ISDR and NS5A-IRRDR. There was a significant positive correlation between the number of aa substitutions in ISDR and that in IRRDR ( $r = 0.308$ ,  $p < 0.001$ ).

Furthermore, the number of aa substitutions in IRRDR of 123 patients with Arg70 (median 5) was also not significantly different from that of 77 patients with Gln70 (His70) (median 4) (fig. 3b).

Concerning the substitution of core aa 91, the number of aa substitutions in ISDR of 240 patients with Leu91 (median 1) was significantly higher than that of 161 patients with Met91 (median 0) ( $p < 0.001$ ) (fig. 3c). Furthermore, the number of aa substitutions in IRRDR of 111 patients with Leu91 (median 5) was significantly higher than that of 89 patients with Met91 (median 3) ( $p < 0.001$ ) (fig. 3d).

#### *Viremia Level and aa Substitutions in Core Region/ISDR/IRRDR*

Concerning the number of substitutions in ISDR, viremia levels of 321 patients with WT (median 6.5) were significantly higher than those of 84 patients with non-WT (median 5.7) ( $p < 0.001$ ) (fig. 4a).

Concerning the number of substitutions in IRRDR, viremia levels of 140 patients with  $\leq 5$  substitutions (median 6.4) were significantly higher than those of 60 patients with  $\geq 6$  (median 6.1) ( $p = 0.027$ ) (fig. 4b).

Concerning the substitution of core aa 70, viremia levels of 265 patients with Arg70 (median 6.4) were not significantly different from those of 151 patients with Gln70 (His70) (median 6.3) (fig. 4c).

Concerning the substitution of core aa 91, viremia levels of 169 patients with Met91 (median 6.5) were significantly higher than those of 245 patients with Leu91 (median 6.2) ( $p = 0.028$ ) (fig. 4d).

Thus, levels of viremia were influenced by aa substitutions in core aa 91 and ISDR/IRRDR.


#### *Treatment Response according to the Number of aa Substitutions in IRRDR*

Concerning the number of aa substitutions in IRRDR, a significantly higher proportion of patients with  $\geq 4$  aa substitutions (58%) showed SVR compared to patients with  $\leq 3$  (42%) ( $p = 0.039$ ). In contrast, the SVR rate was not significantly different between patients with  $\leq 4$  (49%) and those with  $\geq 5$  (57%) aa substitutions. Likewise, the SVR rate was not significantly different between patients with  $\leq 5$  (51%) and those with  $\geq 6$  (55%) aa substitutions (fig. 5a).

The ETR rate was not significantly different between patients with  $\leq 3$  (74%) and those with  $\geq 4$  (82%) aa substitutions, nor between patients with  $\leq 4$  (76%) and those with  $\geq 5$  (83%). Likewise, the ETR rate was not significantly different between those with  $\leq 5$  (79%) and those with  $\geq 6$  (80%) aa substitutions (fig. 5b).

The SVR rate after ETR was not significantly different between patients with  $\leq 3$  (61%) and those with  $\geq 4$  (74%) aa substitutions, nor between patients with  $\leq 4$  (67%) and those with  $\geq 5$  (72%). Likewise, they were not significantly different between patients with  $\leq 5$  (67%) and those with  $\geq 6$  (75%) aa substitutions (fig. 5c).

Thus, it was useful as predictor of SVR to categorize into two groups of  $\leq 4$  and  $\geq 5$  aa substitutions by univariate analysis. However, the ETR and SVR after ETR rates were not significantly different according to the number of aa substitutions in IRRDR.



**Fig. 3.** aa substitutions in the core region and NS5A-ISDR/IRRDR. **a, b** Concerning the substitution of core aa 70, the number of aa substitutions in ISDR/IRRDR of patients with Arg70 was not significantly different from that of patients with Gln70 (His70). **c, d** Concerning the substitution of core aa 91, the number of aa substitutions in ISDR/IRRDR of patients with Leu91 was significantly higher than that of patients with Met91 ( $p < 0.001$ ).

#### Predictors of SVR as Determined by Uni- and Multivariate Analyses


Univariate analysis identified 15 parameters that correlate with SVR: gender (male sex;  $p < 0.001$ ), age ( $< 55$  years;  $p < 0.001$ ), ribavirin dose ( $\geq 11.0$  mg/kg;  $p = 0.006$ ), AST ( $< 58$  IU/l;  $p = 0.039$ ), leukocyte count ( $\geq 4,500/\text{mm}^3$ ;  $p = 0.043$ ), hemoglobin ( $\geq 14.0$  g/dl;  $p = 0.001$ ), platelet count ( $\geq 15.0 \times 10^4/\text{mm}^3$ ;  $p < 0.001$ ), GGT ( $< 50$  IU/l;  $p = 0.028$ ), uric acid ( $\geq 5.5$  mg/dl;  $p = 0.005$ ), level of viremia ( $< 6.0$  log IU/ml;  $p < 0.001$ ),  $\alpha$ -fetoprotein ( $< 10$   $\mu\text{g/l}$ ;  $p < 0.001$ ), genetic variation in rs8099917 (genotype TT;  $p < 0.001$ ), substitution of aa 70 (Arg70;  $p < 0.001$ ), the number of aa substitutions in ISDR (non-WT;  $p < 0.001$ ) and IRRDR ( $\geq 4$ ;  $p = 0.039$ ). Figure 6 shows the SVR rate according to aa substitution in the core/NS5A region and genetic variation near *IL28B* by univariate analysis.

Multivariate analysis that included the above variables identified 3 parameters that independently influenced

SVR: genetic variation in rs8099917 (genotype TT;  $p < 0.001$ ), gender (male sex;  $p < 0.001$ ), and the number of aa substitutions in ISDR (non-WT;  $p = 0.027$ ) (table 2).


#### Predictors of ETR as Determined by Uni- and Multivariate Analyses

Univariate analysis identified 14 parameters that correlated with ETR: gender (male sex;  $p = 0.001$ ), age ( $< 55$  years;  $p = 0.004$ ), AST ( $< 39$  IU/l;  $p = 0.027$ ), hemoglobin ( $\geq 14.0$  g/dl;  $p = 0.035$ ), platelet count ( $\geq 15.0 \times 10^4/\text{mm}^3$ ;  $p < 0.001$ ), albumin ( $\geq 3.9$  g/dl;  $p = 0.014$ ), GGT ( $< 50$  IU/l;  $p < 0.001$ ), uric acid ( $\geq 5.5$  mg/dl;  $p = 0.003$ ), level of viremia ( $< 6.0$  log IU/ml;  $p = 0.001$ ), low-density lipoprotein cholesterol ( $\geq 85$  mg/dl;  $p = 0.004$ ),  $\alpha$ -fetoprotein ( $< 10$   $\mu\text{g/l}$ ;  $p < 0.001$ ), genetic variation in rs8099917 (genotype TT;  $p < 0.001$ ), substitution of aa 70 (Arg70;  $p < 0.001$ ), and the number of aa substitutions in ISDR (non-WT;  $p = 0.021$ ). Figure 7 shows the ETR rate according to aa



**Fig. 4.** Viremia level and aa substitutions in core region/ISDR/IRRDR. **a** Concerning the number of substitutions in ISDR, viremia levels of patients with WT were significantly higher than those of patients with non-WT ( $p < 0.001$ ). **b** Concerning the number of substitutions in IRRDR, viremia levels of patients with  $\leq 5$  aa substitutions were significantly higher levels than those of patients with  $\geq 6$  ( $p = 0.027$ ). **c** Concerning the substitution of


core aa 70, viremia levels of patients with Arg70 were not significantly different from those of patients with Gln70 (His70). **d** Concerning the substitution of core aa 91, viremia levels of patients with Met91 were significantly higher than those of patients with Leu91 ( $p = 0.028$ ). Thus, levels of viremia might be influenced by aa substitutions in core aa 91 and ISDR/IRRDR.




**Fig. 5.** Treatment response according to the number of aa substitutions in NSSA-IRRDR. **a** A significantly higher proportion of patients with  $\geq 4$  (58%) aa substitutions showed SVR compared to patients with  $\leq 3$  (42%) ( $p = 0.039$ ), and it was useful as predictor

of SVR to categorize into two groups of  $\leq 4$  and  $\geq 5$  aa substitutions by univariate analysis. **b, c** ETR and SVR after ETR rates were not significantly different according to the number of aa substitutions in IRRDR.

**Fig. 6.** SVR rate according to aa substitution in core/NS5A region and genetic variation near *IL28B* by univariate analysis.



**Fig. 7.** ETR rate according to aa substitution in core/NS5A region and genetic variation near *IL28B* by univariate analysis.



**Table 2.** Factors associated with SVR to 48-week IFN + ribavirin combination therapy in patients infected with HCV-1b, identified by multivariate analysis

Factor	Category	OR (95% CI)	p
rs8099917 genotype	1: TG+GG	1	<0.001
	2: TT	16.7 (4.54–61.3)	
Gender	1: Female	1	<0.001
	2: Male	10.5 (3.47–32.3)	
ISDR of NS5A	1: WT	1	0.027
	2: Non-WT	5.68 (1.22–26.3)	


Only variables that achieved statistical significance ( $p < 0.05$ ) on multivariate logistic regression are shown.

**Table 3.** Factors associated with ETR response to 48-week IFN + ribavirin combination therapy in patients infected with HCV-1b, identified by multivariate analysis

Factor	Category	OR (95% CI)	p
rs8099917 genotype	1: TG+GG	1	<0.001
	2: TT	18.2 (6.29–52.6)	
Level of viremia log IU/ml	1: $\geq 6.0$	1	0.001
	2: $< 6.0$	9.20 (2.59–32.6)	
Core aa 70	1: Gln70 (His70)	1	0.004
	2: Arg70	4.68 (1.65–13.3)	
Serum albumin g/dl	1: $< 3.9$	1	0.030
	2: $\geq 3.9$	3.08 (1.11–8.47)	

Only variables that achieved statistical significance ( $p < 0.05$ ) on multivariate logistic regression are shown.

**Fig. 8.** SVR after ETR rate according to aa substitution in core/NS5A region and genetic variation near *IL28B* by univariate analysis.



substitution in the core/NS5A region and genetic variation near *IL28B* by univariate analysis.

Multivariate analysis that included the above variables identified 4 parameters that independently influenced ETR: genetic variation in rs8099917 (genotype TT;  $p < 0.001$ ), level of viremia ( $< 6.0 \log \text{ IU/ml}$ ;  $p = 0.001$ ), substitution of aa 70 (Arg70;  $p = 0.004$ ), and albumin ( $\geq 3.9 \text{ g/dl}$ ;  $p = 0.030$ ) (table 3).

#### Predictors of SVR after ETR as Determined by Uni- and Multivariate Analyses

Univariate analysis identified 11 parameters that influenced SVR after ETR: gender (male sex;  $p < 0.001$ ), age ( $< 55$  years;  $p < 0.001$ ), ribavirin dose ( $\geq 11.0 \text{ mg/kg}$ ;  $p = 0.025$ ), leukocyte count ( $\geq 4,500/\text{mm}^3$ ;  $p = 0.033$ ), hemoglobin ( $\geq 14.0 \text{ g/dl}$ ;  $p = 0.025$ ), platelet count ( $\geq 15.0 \times 10^4/\text{mm}^3$ ;  $p = 0.001$ ), level of viremia ( $< 6.0 \log \text{ IU/ml}$ ;  $p = 0.020$ ), total cholesterol ( $< 170 \text{ mg/dl}$ ;  $p = 0.017$ ),  $\alpha$ -fetoprotein ( $< 10 \mu\text{g/l}$ ;  $p = 0.004$ ), substitution of aa 70 and 91 (Arg70 and/or Leu91;  $p = 0.044$ ), and the number of aa substitutions in ISDR (non-WT;  $p = 0.001$ ). Figure 8 shows the SVR after ETR rate according to aa substitution in the core/NS5A region and genetic variation near *IL28B* by univariate analysis.

Multivariate analysis that included the above variables identified 6 parameters that independently influenced the SVR after ETR: gender (male sex;  $p < 0.001$ ), ribavirin dose ( $\geq 11.0 \text{ mg/kg}$ ;  $p = 0.002$ ), the number of aa substitutions in ISDR (non-WT;  $p = 0.012$ ), substitution of aa 70 and 91 (Arg70 and/or Leu91;  $p = 0.023$ ), platelet count ( $\geq 15.0 \times 10^4/\text{mm}^3$ ;  $p = 0.033$ ), and  $\alpha$ -fetoprotein ( $< 10 \mu\text{g/l}$ ;  $p = 0.042$ ) (table 4).

#### Comparison of Factors Associated with Treatment Efficacy Identified by Multivariate Analysis

Table 5 shows the variables that achieved statistical significance on multivariate logistic regression for each evaluation of treatment efficacy. Rs8099917 genotype was an important predictor of ETR and SVR. With regard to viral factors, core region was an important predictor of ETR, and SVR after ETR. ISDR was an important predictor of SVR, and SVR after ETR. Level of viremia was an important predictor of ETR. Thus, genetic variation near *IL28B* and viral factors (core region, ISDR, and level of viremia) were important predictors of treatment efficacy. Furthermore, gender,  $\alpha$ -fetoprotein, albumin, and platelet count were also identified as other important predictors of treatment efficacy, in addition to genetic variation near *IL28B* and viral factors.

#### Discussion

Using multivariate analysis, the present study identified viral- (aa substitutions in core aa 70/91 and NS5A-ISDR/IRRDR) and host-related factors (genetic variation near *IL28B* gene) that influenced treatment efficacy to 48-week IFN/ribavirin combination therapy, which is in agreement with recent findings [22, 23]. Identification of these viral and host factors before the start of IFN/ribavirin combination therapy should help to select better therapeutic regimens, including triple therapy of telaprevir/PEG-IFN/ribavirin [24–26], for those patients who are less likely to achieve SVR.

According to the number of substitutions in ISDR, a previous report showed that levels of viremia were sig-

**Table 4.** Factors associated with SVR in patients who achieved ETR response to 48-week IFN + ribavirin combination therapy in patients infected with HCV-1b, identified by multivariate analysis

Factor	Category	OR (95% CI)	p
Gender	1: Female	1	<0.001
	2: Male	4.27 (2.15–8.55)	
Ribavirin dose, mg/kg	1: <11.0	1	0.002
	2: ≥11.0	2.95 (1.48–5.86)	
ISDR of NS5A	1: WT	1	0.012
	2: Non-WT	4.00 (1.35–11.8)	
Core aa 70 and 91	1: Gln70 (His70) and Met91	1	0.023
	2: Arg70 and/or Leu91	2.96 (1.16–7.52)	
Platelet count × 10 <sup>4</sup> /mm <sup>3</sup>	1: <15.0	1	0.033
	2: ≥15.0	2.19 (1.07–4.50)	
α-Fetoprotein μg/l	1: ≥10	1	0.042
	2: <10	2.66 (1.04–6.80)	

Only variables that achieved statistical significance (p < 0.05) on multivariate logistic regression are shown.

**Table 5.** Comparison of factors associated with efficacy of 48-week IFN + ribavirin combination therapy in patients infected with HCV-1b, identified by multivariate analysis

Factor	ETR response (at 48 weeks)	SVR after ETR response	SVR
<i>IL28B</i>	rs8099917 p < 0.001, 18.2 (6.29–52.6) <sup>a</sup>		rs8099917 p < 0.001, 16.7 (4.54–61.3) <sup>a</sup>
Virus	Core aa 70 p = 0.004, 4.68 (1.65–13.3) <sup>a</sup>	Core aa 70 and 91 p = 0.023, 2.96 (1.16–7.52) <sup>a</sup>	
	Level of viremia p = 0.001, 9.20 (2.59–32.6) <sup>a</sup>	ISDR p = 0.012, 4.00 (1.35–11.8) <sup>a</sup>	ISDR p = 0.027, 5.68 (1.22–26.3) <sup>a</sup>
Others	Albumin p = 0.030, 3.08 (1.11–8.47) <sup>a</sup>	α-Fetoprotein p = 0.042, 2.66 (1.04–6.80) <sup>a</sup>	
		Platelet count p = 0.033, 2.19 (1.07–4.50) <sup>a</sup>	
		Gender p < 0.001, 4.27 (2.15–8.55) <sup>a</sup>	Gender p < 0.001, 10.5 (3.47–32.3) <sup>a</sup>
		Ribavirin dose p = 0.002, 2.95 (1.48–5.86) <sup>a</sup>	

Only variables that achieved statistical significance (p < 0.05) on multivariate logistic regression are shown.  
<sup>a</sup> OR (95% CI).

nificantly lower in patients with non-WT of ISDR than in those with WT [8]. The present study indicated that substitution of IRRDR and core aa 91, in addition to substitution of ISDR, also significantly influenced levels of viremia. Furthermore, there was a significant positive correlation between the number of aa substitutions in

ISDR and those in IRRDR, and the number of aa substitutions in ISDR/IRRDR of patients with Leu91 was significantly higher than that of patients with Met91. To our knowledge, this is the first report of the relationship between viremia levels and aa substitutions in core region/ISDR/IRRDR. This result might be interpreted to mean



that core aa 91/ISDR/IRRDR might be associated with viremia levels involved in resistance to combination therapy. Further studies that examine the functional impact of aa substitutions to combination therapy should be conducted to confirm the above finding.

The present results showed that  $\alpha$ -fetoprotein, albumin, platelet count, and gender were predictors of virological response to IFN/ribavirin combination therapy. Previous data indicated that absence of advanced liver fibrosis was a positive predictor of SVR to IFN monotherapy and IFN/ribavirin combination therapy [2, 3, 13, 27–29], and that advanced liver fibrosis was usually associated with higher levels of  $\alpha$ -fetoprotein, and lower levels of albumin and platelet count [1, 3, 30–32]. Furthermore, gender is also a predictor of treatment response to IFN/ribavirin combination therapy [2, 3, 14]. In the present study based on a large number of patients, histopathological changes in the liver and gender were identified as independent predictors of virological response, in addition to genetic variation near *IL28B* and viral factors (core region, ISDR, and level of viremia).

In a previous study, multivariate analysis identified core region, gender, and stage of liver fibrosis as parameters that independently influenced the SVR of patients who achieved early virological response, but ISDR was not entered into uni- and multivariate analysis [3]. To our knowledge, the present study based on multivariate analysis is the first report to identify ISDR as pretreatment

predictor of SVR after ETR to combination therapy. Interestingly, ISDR was not a predictor of ETR, but was a significant predictor of SVR to combination therapy. Thus, the underlying mechanisms of failure to develop SVR in those patients who achieve HCV-RNA negativity remain unclear. Further studies that examine the impact of aa substitutions of ISDR to combination therapy should be conducted to confirm the above finding.

One limitation of the present study was that aa substitutions in areas other than the core region and NS5A-ISDR/IRRDR of the HCV genome were not examined. Other limitations were differences in host factors including race [24, 33, 34] and differences in viral factors, such as the distribution of HCV-1a or -1b, and geographic diversities of HCV-1b [35]. Further large-scale prospective studies are necessary to investigate whether the present results relate to the efficacy of 48-week IFN/ribavirin combination therapy, and further understanding of the complex interaction between virus- and host- related factors should facilitate the development of more effective therapeutic regimens.

#### Acknowledgement

This study was supported in part by a Grant-in-Aid from the Ministry of Health, Labor and Welfare, Japan.

#### References

- 1 Akuta N, Suzuki F, Sezaki H, Suzuki Y, Hosaka T, Someya T, Kobayashi M, Saitoh S, Watahiki S, Sato J, Matsuda M, Kobayashi M, Arase Y, Ikeda K, Kumada H: Association of amino acid substitution pattern in core protein of hepatitis C virus genotype 1b high viral load and non-virological response to interferon-ribavirin combination therapy. *Intervirol* 2005;48:372–380.
- 2 Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Arase Y, Ikeda K, Kumada H: Predictive factors of early and sustained responses to peginterferon plus ribavirin combination therapy in Japanese patients infected with hepatitis C virus genotype 1b: amino acid substitutions in the core region and low-density lipoprotein cholesterol levels. *J Hepatol* 2007;46:403–410.
- 3 Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Arase Y, Ikeda K, Kumada H: Predictors of viral kinetics to peginterferon plus ribavirin combination therapy in Japanese patients infected with hepatitis C virus genotype 1b. *J Med Virol* 2007;79:1686–1695.
- 4 Donlin MJ, Cannon NA, Yao E, Li J, Wahed A, Taylor MW, Belle SH, Di Bisceglie AM, Aurora R, Tavis JE: Pretreatment sequence diversity differences in the full-length hepatitis C virus open reading frame correlate with early response to therapy. *J Virol* 2007;81:8211–8224.
- 5 Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Arase Y, Ikeda K, Kumada H: Amino acid substitutions in the hepatitis C virus core region are the important predictor of hepatocarcinogenesis. *Hepatology* 2007;46:1357–1364.
- 6 Fishman SL, Factor SH, Balestrieri C, Fan X, Di Bisceglie AM, Desai SM, Benson G, Branch AD: Mutations in the hepatitis C virus core gene are associated with advanced liver disease and hepatocellular carcinoma. *Clin Cancer Res* 2009;15:3205–3213.
- 7 Enomoto N, Sakuma I, Asahina Y, Kurosaki M, Murakami T, Yamamoto C, Izumi N, Marumo F, Sato C: Comparison of full-length sequences of interferon sensitive and resistant hepatitis C virus 1b. Sensitivity to interferon is conferred by amino acid substitutions in the NS5A region. *J Clin Invest* 1995;96:224–230.
- 8 Enomoto N, Sakuma I, Asahina Y, Kurosaki M, Murakami T, Yamamoto C, Ogura Y, Izumi N, Marumo F, Sato C: Mutations in the nonstructural protein 5A gene and response to interferon in patients with chronic hepatitis C virus 1b infection. *N Engl J Med* 1996;334:77–81.
- 9 El-Shamy A, Sasayama M, Nagano-Fujii M, Sasase N, Imoto S, Kim SR, Hotta H: Prediction of efficient virological response to pegylated interferon/ribavirin combination therapy by NS5A sequences of hepatitis C virus and anti-NS5A antibodies in pre-treatment sera. *Microbiol Immunol* 2007;51:471–482.



- 10 El-Shamy A, Nagano-Fujii M, Sasase N, Imoto S, Kim SR, Hotta H: Sequence variation in hepatitis C virus nonstructural protein 5A predicts clinical outcome of pegylated interferon/ribavirin combination therapy. *Hepatology* 2008;48:38–47.
- 11 Shirakawa H, Matsumoto A, Joshita S, Komatsu M, Tanaka N, Umemura T, Ichijo T, Yoshizawa K, Kiyosawa K, Tanaka E: Nagano Interferon Treatment Research Group: Pre-treatment prediction of virological response to peginterferon plus ribavirin therapy in chronic hepatitis C patients using viral and host factors. *Hepatology* 2008;48:1753–1760.
- 12 Mori N, Imamura M, Kawakami Y, Saneto H, Kawaoka T, Takaki S, Aikata H, Takahashi S, Chayama K: Hiroshima Liver Study Group: Randomized trial of high-dose interferon- $\alpha$ -2b combined with ribavirin in patients with chronic hepatitis C: Correlation between amino acid substitutions in the core/NS5A region and virological response to interferon therapy. *J Med Virol* 2009;81:640–649.
- 13 Ge D, Fellay J, Thompson AJ, Simon JS, Shianna KV, Urban TJ, Heinzen EL, Qiu P, Bertelsen AH, Muir AJ, Sulkowski M, McHutchinson JG, Goldstein DB: Genetic variation in *IL28B* predicts hepatitis C treatment-induced viral clearance. *Nature* 2009;461:399–401.
- 14 Tanaka Y, Nishida N, Sugiyama M, Kurosaki M, Matsuura K, Sakamoto N, Nakagawa M, Korenaga M, Hino K, Hige S, Ito Y, Mita E, Tanaka E, Mochida S, Murawaki Y, Honda M, Sakai A, Hiasa Y, Nishiguchi S, Koike A, Sakaida I, Imamura M, Ito K, Yano K, Masaki N, Sugauchi F, Izumi N, Tokunaga K, Mizokami M: Genome-wide association of *IL28B* with response to pegylated interferon- $\alpha$  and ribavirin therapy for chronic hepatitis C. *Nat Genet* 2009;41:1105–1109.
- 15 Suppiah V, Moldovan M, Ahlenstiel G, Berg T, Weltman M, Abate ML, Bassendine M, Spengler U, Dore GJ, Powell E, Riordan S, Sheridan D, Smedile A, Fragomeli V, Müller T, Bahlo M, Stewart GJ, Booth DR, George J: *IL28B* is associated with response to chronic hepatitis C interferon- $\alpha$  and ribavirin therapy. *Nat Genet* 2009;41:1100–1104.
- 16 Rauch A, Kutalik Z, Descombes P, Cai T, Di Julio J, Mueller T, Bochud M, Battegay M, Bernasconi E, Borovicka J, Colombo S, Cerny A, Dufour JF, Furrer H, Günthard HF, Heim M, Hirschel B, Malinverni R, Moradpour D, Müllhaupt B, Witteck A, Beckmann JS, Berg T, Bergmann S, Negro F, Telenti A, Bochud PY: Swiss Hepatitis C Cohort Study; Swiss HIV Cohort Study: Genetic variation in *IL28B* is associated with chronic hepatitis C and treatment failure: a genome-wide association study. *Gastroenterology* 2010;138:1338–1345.
- 17 Thomas DL, Thio CL, Martin MP, Qi Y, Ge D, O'Huigin C, Kidd J, Kidd K, Khakoo SI, Alexander G, Goedert JJ, Kirk GD, Donfield SM, Rosen HR, Tobler LH, Busch MP, McHutchinson JG, Goldstein DB, Carrington M: Genetic variation in *IL28B* and spontaneous clearance of hepatitis C virus. *Nature* 2009;461:798–801.
- 18 Akuta N, Suzuki F, Hirakawa M, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Saitoh S, Arase Y, Ikeda K, Chayama K, Nakamura Y, Kumada H: Amino acid substitution in HCV core region and genetic variation near the interleukin-28B gene predict viral response to telaprevir with peginterferon and ribavirin. *Hepatology* 2010;52:421–429.
- 19 Kato N, Hijikata M, Ootsuyama Y, Nakagawa M, Ohkoshi S, Sugimura T, Shimotohno K: Molecular cloning of the human hepatitis C virus genome from Japanese patients with non-A, non-B hepatitis. *Proc Natl Acad Sci USA* 1990;87:9524–9528.
- 20 Ohnishi Y, Tanaka T, Ozaki K, Yamada R, Suzuki H, Nakamura Y: A high-throughput SNP typing system for genome-wide association studies. *J Hum Genet* 2001;46:471–477.
- 21 Suzuki A, Yamada R, Chang X, Tokuhiro S, Sawada T, Suzuki M, Nagasaki M, Nakayama-Hamada M, Kawaida R, Ono M, Ohtsuki M, Furukawa H, Yoshino S, Yukioka M, Tohma S, Matsuura K, Wakitani S, Teshima R, Nishioka Y, Sekine A, Iida A, Takahashi A, Tsunoda T, Nakamura Y, Yamamoto K: Functional haplotypes of PADI4, encoding citrullinating enzyme peptidylarginine deiminase 4, are associated with rheumatoid arthritis. *Nat Genet* 2003;34:395–402.
- 22 Kurosaki M, Tanaka Y, Nishida N, Sakamoto N, Enomoto N, Honda M, Sugiyama M, Matsuura K, Sugauchi F, Asahina Y, Nakagawa M, Watanabe M, Sakamoto M, Maekawa S, Sakai A, Kaneko S, Ito K, Masaki N, Tokunaga K, Izumi N, Mizokami M: Pre-treatment prediction of response to pegylated interferon plus ribavirin for chronic hepatitis C using genetic polymorphism in *IL28B* and viral factors. *J Hepatol* 2011;54:439–448.
- 23 Hayes CN, Kobayashi M, Akuta N, Suzuki F, Kumada H, Abe H, Miki D, Imamura M, Ochi H, Kamatani N, Nakamura Y, Chayama K: HCV substitutions and *IL28B* polymorphisms on outcome of peg-interferon plus ribavirin combination therapy. *Gut* 2011;60:261–267.
- 24 McHutchinson JG, Everson GT, Gordon SC, Jacobson IM, Sulkowski M, Kauffman R, McNair L, Alam J, Muir AJ: PROVE1 Study Team: Telaprevir with peginterferon and ribavirin for chronic HCV genotype 1 infection. *N Engl J Med* 2009;360:1827–1838.
- 25 McHutchinson JG, Manns MP, Muir AJ, Terrault NA, Jacobson IM, Afdhal NH, Heathcote EJ, Zeuzem S, Reesink HW, Garg J, Bsharat M, George S, Kauffman RS, Adda N, Di Bisceglie AM: PROVE3 Study Team: Telaprevir for previously treated chronic HCV infection. *N Engl J Med* 2010;362:1292–1303.
- 26 Hézode C, Forestier N, Dusheiko G, Ferenci P, Pol S, Goeser T, Bronowicki JP, Bourlière M, Gharakhanian S, Bengtsson L, McNair L, George S, Kieffer T, Kwong A, Kauffman RS, Alam J, Pawlotsky JM, Zeuzem S: PROVE2 Study Team: Telaprevir and peginterferon with or without ribavirin for chronic HCV infection. *N Engl J Med* 2009;360:1839–1850.
- 27 Jouet P, Roudot-Thoraval F, Dhumeaux D, Metreau JM: Comparative efficacy of interferon alfa in cirrhotic and noncirrhotic patients with non-A, non-B, C hepatitis. *Gastroenterology* 1994;106:686–690.
- 28 Poynard T, McHutchinson J, Goodman Z, Ling MH, Albrecht J: Is an 'a la carte' combination interferon alfa-2b plus ribavirin regimen possible for the first-line treatment in patients with chronic hepatitis C? The AL-GOVIHC Group. *Hepatology* 2000;31:211–218.
- 29 Bruno S, Camma C, Di Marco V, Rumi M, Vinci M, Camozzi M, Rebutti C, Di Bona D, Colombo M, Craxi A, Mondelli MU, Pinzello G: Peginterferon alfa-2b plus ribavirin for naïve patients with genotype 1 chronic hepatitis C: a randomized controlled trial. *J Hepatol* 2004;41:474–481.
- 30 Bayati N, Silverman AL, Gordon SC: Serum  $\alpha$ -fetoprotein levels and liver histology in patients with chronic hepatitis C. *Am J Gastroenterol* 1998;93:2452–2456.
- 31 Chu CW, Hwang SJ, Luo JC, Lai CR, Tsay SH, Li CP, Wu JC, Chang FY, Lee SD: Clinical, virological, and pathologic significance of elevated serum  $\alpha$ -fetoprotein levels in patients with chronic hepatitis C. *J Clin Gastroenterol* 2001;32:240–244.
- 32 Hu KQ, Kyulo NL, Lim N, Elhazin B, Hillebrand DJ, Bock T: Clinical significance of elevated  $\alpha$ -fetoprotein in patients with chronic hepatitis C, but not hepatocellular carcinoma. *Am J Gastroenterol* 2004;99:860–865.
- 33 McHutchinson JG, Poynard T, Pianko S, Gordon SC, Reid AE, Dienstag J, Morgan T, Yao R, Albrecht J: The impact of interferon plus ribavirin on response to therapy in black patients with chronic hepatitis C. The International Hepatitis Interventional Therapy Group. *Gastroenterology* 2000;119:1317–1323.
- 34 Kaplan DE, Sugimoto K, Ikeda F, Stadanlick J, Valiga M, Shetty K, Reddy KR, Chang KM: T-cell response relative to genotype and ethnicity during antiviral therapy for chronic hepatitis C. *Hepatology* 2005;41:1365–1375.
- 35 Nakano I, Fukuda Y, Katanō Y, Nakano S, Kumada T, Hayakawa T: Why is the interferon sensitivity-determining region (ISDR) system useful in Japan? *J Hepatol* 1999;30:1014–1022.

## Amino Acid Substitution in HCV Core Region and Genetic Variation near the *IL28B* Gene Affect Viral Dynamics during Telaprevir, Peginterferon and Ribavirin Treatment

Norio Akuta<sup>a</sup> Fumitaka Suzuki<sup>a</sup> Miharuru Hirakawa<sup>a</sup> Yusuke Kawamura<sup>a</sup>  
Hiromi Yatsuji<sup>a</sup> Hitomi Sezaki<sup>a</sup> Yoshiyuki Suzuki<sup>a</sup> Tetsuya Hosaka<sup>a</sup>  
Masahiro Kobayashi<sup>a</sup> Mariko Kobayashi<sup>b</sup> Satoshi Saitoh<sup>a</sup> Yasuji Arase<sup>a</sup>  
Kenji Ikeda<sup>a</sup> Kazuaki Chayama<sup>d</sup> Yusuke Nakamura<sup>c</sup> Hiromitsu Kumada<sup>a</sup>

<sup>a</sup>Department of Hepatology, and <sup>b</sup>Liver Research Laboratory, Toranomon Hospital, <sup>c</sup>Laboratory of Molecular Medicine, Human Genome Center, Institute of Medical Science, University of Tokyo, Tokyo, and <sup>d</sup>Department of Medical and Molecular Science, Division of Frontier Medical Science, Programs for Biomedical Research, Graduate School of Biomedical Science, Hiroshima University, Hiroshima, Japan

### Key Words

Hepatitis C virus · Core region · *IL28B* · Telaprevir · Peginterferon · Ribavirin · Viral dynamics

### Abstract

**Objectives:** Genetic variation near the *IL28B* gene and substitution of aa 70 and 91 in the core region of HCV-1b are useful as predictors of treatment efficacy to telaprevir/pegylated interferon (PEG-IFN)/ribavirin, but its impact on viral dynamics is not clear. **Methods:** This study investigated predictive factors of viral dynamics during 12- or 24-week regimen of triple therapy in 80 Japanese adults infected with HCV-1b. **Results:** After 24 h of commencement of treatment, the proportion of patients with Arg70 and Leu91 substitutions in the core region who showed  $\geq 3.0$  log drop in HCV RNA level was significantly higher than that of patients with Gln70 (His70) and/or Met91. At 8 and 12 weeks, HCV RNA loss rate of patients with rs8099917 genotype TT near *IL28B* gene was significantly higher than that of patients with non-TT.

Multivariate analysis identified substitution of aa 70 and 91 as a predictor of  $\geq 3.0$  log fall in HCV RNA level at 24 h (Arg70 and Leu91) and SVR (Arg70), and rs8099917 (TT) as a predictor of HCV RNA loss at 12 weeks and SVR. **Conclusions:** This study identified genetic variation near *IL28B* gene and aa substitution of the core region as predictors of viral dynamics during triple therapy.

Copyright © 2011 S. Karger AG, Basel

### Introduction

Hepatitis C virus (HCV) usually causes chronic infection that can result in chronic hepatitis, liver cirrhosis, and hepatocellular carcinoma (HCC) [1, 2]. At present, treatments based on interferon (IFN), in combination with ribavirin, are mainstay for combating HCV infection. In Japan, HCV genotype 1b (HCV-1b) in high viral loads ( $>100$  kIU/ml) accounts for more than 70% of HCV infections, making it difficult to treat patients with chronic hepatitis

KARGER

Fax +41 61 306 12 34  
E-Mail karger@karger.ch  
www.karger.com

© 2011 S. Karger AG, Basel  
0300-5526/11/0000-0000\$38.00/0

Accessible online at:  
www.karger.com/int

Norio Akuta, MD  
Department of Hepatology, Toranomon Hospital  
2-2-2 Toranomon, Minato-ku  
Tokyo 105-0001 (Japan)  
Tel. +81 44 877 5111, E-Mail akuta-gi@umin.ac.jp

C [3]. Such a background calls for efficient treatments of Japanese patients with chronic HCV infection.

Even with pegylated IFN (PEG-IFN) combined with ribavirin, a sustained virological response lasting over 24 weeks after the withdrawal of treatment is achieved in at most 50% of the patients infected with HCV-1b and high viral loads [4, 5]. Recently, a new strategy was introduced in the treatment of chronic HCV infection by means of inhibiting protease in the NS3/NS4 of the HCV polyprotein. Of these, telaprevir (VX-950) was selected as a candidate agent for treatment of chronic HCV infection [6]. Later, it was found that telaprevir, when combined with PEG-IFN and ribavirin, gains a robust antiviral activity [7, 8]. Two previous studies (PROVE1 and PROVE2) showed that the 12- and 24-week regimen of telaprevir/PEG-IFN/ribavirin could achieve sustained virological response rates of 35–60 and 61–69% in patients infected with HCV-1, respectively [9, 10]. Furthermore, a recent study (PROVE3) also showed that the 24- and 48-week regimen of triple therapy could achieve sustained virological response rates of 51 and 53% in HCV-1 infected patients in whom initial PEG-IFN/ribavirin treatment failed, respectively [11].

Amino acid (aa) substitutions at positions 70 and/or 91 in the HCV core region of patients infected with HCV-1b and high viral loads are pretreatment predictors of poor virological response to PEG-IFN plus ribavirin combination therapy [12–14], and also affect clinical outcome, including hepatocarcinogenesis [15, 16]. Furthermore, genetic variations near the *IL28B* gene (rs8099917, rs12979860) on chromosome 19 as host-related factor, which encodes IFN- $\lambda$ -3, are pretreatment predictors of virological response to 48-week PEG-IFN plus ribavirin combination therapy in individuals infected with HCV-1 [17–20], and also affect clinical outcome, including spontaneous clearance of HCV [21]. A recent report identified genetic variation near *IL28B* gene and aa substitution of the core region as predictors of sustained virological response to triple therapy of telaprevir/PEG-IFN/ribavirin in Japanese patients infected with HCV-1b [22]. However, it is not clear at this stage whether genetic variation near the *IL28B* gene and aa substitution of the core region can be used before therapy to predict viral dynamics during triple therapy.

The present study included 80 patients with HCV-1b and high viral loads, who received the triple therapy of telaprevir with PEG-IFN plus ribavirin. The aims of the study were to identify the pretreatment factors that could predict viral dynamics during treatment, including viral- (aa substitutions in the HCV core and NS5A regions) and host-related factors (genetic variation near *IL28B* gene).

## Patients and Methods

### Study Population

Between May 2008 and September 2009, 81 patients infected with HCV were recruited to this study at the Department of Hepatology in Toranomon Hospital in metropolitan Tokyo. The study protocol was in compliance with the Good Clinical Practice Guidelines and the 1975 Declaration of Helsinki, and was approved by the institutional review board. Each patient gave an informed consent before participating in this trial. Patients were divided into two groups: 20 (25%) patients were allocated to a 12-week regimen of triple therapy [telaprevir (MP-424), PEG-IFN and ribavirin] (the T12PR12 group), and 61 patients (75%) were assigned to a 24-week regimen of the same triple therapy for 12 weeks followed by dual therapy of PEG-IFN and ribavirin for 12 weeks (the T12PR24 group).

Eighty of the 81 patients met the following inclusion and exclusion criteria: (1) Diagnosis of chronic hepatitis C. (2) HCV-1b confirmed by sequence analysis. (3) HCV RNA levels of  $\geq 5.0$  log IU/ml determined by the COBAS TaqMan HCV test (Roche Diagnostics, Tokyo, Japan). (4) Japanese (Mongoloid) ethnicity. (5) Age at study entry of 20–65 years. (6) Body weight  $\geq 35$  kg and  $\leq 120$  kg at the time of registration. (7) Lack of decompensated liver cirrhosis. (8) Negativity for hepatitis B surface antigen (HBsAg) in serum. (9) Negative history of HCC. (10) No previous treatment for malignancy. (11) Negative history of autoimmune hepatitis, alcohol liver disease, hemochromatosis, and chronic liver disease other than chronic hepatitis C. (12) Negative history of depression, schizophrenia or suicide attempts, hemoglobinopathies, angina pectoris, cardiac insufficiency, myocardial infarction or severe arrhythmia, uncontrollable hypertension, chronic renal dysfunction or creatinine clearance of  $\leq 50$  ml/min at baseline, diabetes requiring treatment or fasting glucose level of  $\geq 110$  mg/dl, autoimmune disease, cerebrovascular disorders, thyroidal dysfunction uncontrollable by medical treatment, chronic pulmonary disease, allergy to medication or anaphylaxis at baseline. (13) Hemoglobin level of  $\geq 12$  g/dl, neutrophil count  $\geq 1,500/\text{mm}^3$ , and platelet count of  $\geq 100,000/\text{mm}^3$  at baseline. Pregnant or breast-feeding women or those willing to become pregnant during the study and men with a pregnant partner were excluded from the study. In this study, all of the 80 patients were evaluated for the pretreatment predictors for viral dynamics during triple therapy, and 77 of the 80 patients were followed up for at least 24 weeks after the completion of treatment. The treatment efficacy was evaluated by 24 weeks after the completion of therapy (sustained virological response), based on the COBAS TaqMan HCV test (Roche Diagnostics).

Telaprevir (MP-424; Mitsubishi Tanabe Pharma, Osaka, Japan) was administered at 750 or 500 mg three times a day at an 8-hour (q8) interval after the meal. PEG-IFN $\alpha$ -2b (PEG-Intron; Schering Plough, Kenilworth, N.J., USA) was injected subcutaneously at a median dose of 1.5  $\mu\text{g}/\text{kg}$  (range 1.3–2.0  $\mu\text{g}/\text{kg}$ ) once a week. Ribavirin (Rebetol; Schering Plough) was administered at 200–600 mg twice a day after breakfast and dinner (daily dose 600–1,000 mg).

PEG-IFN and ribavirin were discontinued or their doses reduced, as required, upon reduction of hemoglobin level, leukocyte count, neutrophil count or platelet count, or the development of adverse events. Thus, the dose of PEG-IFN was reduced by 50% when the leukocyte count decreased below  $1,500/\text{mm}^3$ , neutro-

**Table 1.** Profile and laboratory data at commencement of telaprevir, peginterferon and ribavirin triple therapy in Japanese patients infected with HCV-1b

<i>Demographic data</i>	
Number of patients	80
Sex, M/F	43/37
Age, years*	55 (23–65)
History of blood transfusion	24 (20.0%)
Family history of liver disease	13 (16.3%)
Body mass index*	22.5 (13.2–32.4)
<i>Laboratory data*</i>	
Level of viremia, log IU/ml	6.8 (5.1–7.6)
Serum aspartate aminotransferase, IU/l	34 (15–118)
Serum alanine aminotransferase, IU/l	42 (12–175)
Serum albumin, g/dl	3.9 (3.3–4.6)
Gamma-glutamyl transpeptidase, IU/l	36 (9–229)
Leukocyte count, per mm <sup>3</sup>	4,800 (2,800–8,100)
Hemoglobin, g/dl	14.3 (11.7–16.8)
Platelet count, × 10 <sup>4</sup> /mm <sup>3</sup>	17.3 (9.5–33.8)
α-Fetoprotein, μg/l	4 (2–39)
Total cholesterol, mg/dl	180 (112–276)
Fasting plasma glucose, mg/dl	92 (64–125)
<i>Treatment</i>	
PEG-IFNα-2b dose, μg/kg*	1.5 (1.3–2.0)
Ribavirin dose, mg/kg*	11.5 (7.2–18.4)
Telaprevir dose, 1,500/2,250 mg/day	10/70
Treatment regimen (T12PR12 group/T12PR24 group)	20/60
<i>Amino acid substitutions in the HCV-1b</i>	
Core aa 70, arginine/glutamine (histidine)	47/33
Core aa 91, leucine/methionine	43/37
ISDR of NS5A, wild-type/non-wild-type	76/4
<i>Genetic variation near IL28B gene</i>	
rs8099917 genotype, TT/TG/GG/ND	46/30/2/2
rs12979860 genotype, CC/CT/TT/ND	43/31/2/4
<i>Past history of IFN therapy</i>	
Treatment naive	27
Relapsers to previous treatment	33
Nonresponders to previous treatment	20
Data are numbers and percentages of patients, except those denoted by *, which represent the median (range) values. ND = Not determined.	

phil count below 750/mm<sup>3</sup> or platelet count below 80,000/mm<sup>3</sup>; PEG-IFN was discontinued when these counts decreased below 1,000/mm<sup>3</sup>, 500/mm<sup>3</sup> or 50,000/mm<sup>3</sup>, respectively. When hemoglobin decreased to <10 g/dl, the daily dose of ribavirin was reduced from 600 to 400, 800 to 600 and 1,000 to 600 mg, depending on the initial dose. Ribavirin was withdrawn when hemoglobin decreased to <8.5 g/dl. However, the dose of telaprevir (MP-424) remained the same, and its administration was stopped when the

discontinuation was appropriate for the development of adverse events. In those patients who discontinued telaprevir, treatment with PEG-IFNα-2b and ribavirin was also terminated.

Table 1 summarizes the profiles and laboratory data of the 80 patients at the commencement of treatment. They included 43 males and 37 females, aged 23–65 years (median 55 years).

#### Measurement of HCV RNA

The antiviral effects of the triple therapy on HCV were assessed by measuring plasma HCV RNA levels. In this study, HCV RNA levels during treatment were evaluated at least once every month before, during, and after therapy. Furthermore, to investigate the pretreatment predictors for viral dynamics, HCV RNA levels during treatment were evaluated at 7 time points; 24 h, 1, 2, 4, 6, 8 and 12 weeks after the commencement of treatment. HCV RNA levels during treatment were evaluated in 80 (100%), 80 (100%), 80 (100%), 79 (98.8%), 75 (93.8%), 74 (92.5%), and 69 (86.3%) of the 80 patients, at the above time intervals, respectively. HCV RNA concentrations were determined using the COBAS TaqMan HCV test (Roche Diagnostics). The linear dynamic range of the assay was 1.2–7.8 log IU/ml, and the undetectable samples were defined as loss of HCV RNA. Especially, falls in HCV RNA levels at 24 h relative to baseline were investigated as very early dynamics.

#### Detection of Amino Acid Substitutions in Core and NS5A Regions of HCV-1b

With the use of HCV-J (accession No. D90208) as a reference [23], the sequence of 1–191 aa in the core protein of HCV-1b was determined and then compared with the consensus sequence constructed on 80 clinical samples to detect substitutions at aa 70 of arginine (Arg70) or glutamine/histidine (Gln70/His70) and aa 91 of leucine (Leu91) or methionine (Met91) [12]. The sequence of 2209–2248 aa in the NS5A of HCV-1b (IFN sensitivity-determining region; ISDR) reported by Enomoto et al. [24] was determined, and the numbers of aa substitutions in ISDR were defined as wild-type (0, 1) or non-wild-type (≥2). In the present study, aa substitutions of the core region and NS5A-ISDR of HCV-1b were analyzed by direct sequencing [22].


#### Genetic Variation near IL28B Gene

Samples for genomewide association survey were genotyped using the Illumina HumanHap610-Quad Genotyping BeadChip. Genotyping data were subjected to quality control before the data analysis. Genotyping for replication and fine mapping was performed by use of the Invader assay, TaqMan assay, or direct sequencing as described previously [25, 26].

In this study, genetic variations near *IL28B* gene (rs8099917, rs12979860), reported as the pretreatment predictors of treatment efficacy and clinical outcome [17–22], were investigated.

#### Statistical Analysis

Nonparametric tests ( $\chi^2$  test and Fisher's exact probability test) were used to compare the characteristics of the groups. Univariate and multivariate logistic regression analyses were used to determine those factors that significantly contributed to viral dynamics and sustained virological response. The ORs and 95%CI were also calculated. All *p* values less than 0.05 by the two-tailed test were considered significant. Variables that achieved statistical significance (*p* < 0.05) on univariate analysis were entered into



**Fig. 1.** **a** Very early dynamics according to amino acid substitutions in core region. After 24 h of commencement of the triple therapy, patients with Arg70 and Leu91 (median 2.9 log IU/ml; range 1.8–4.0 log IU/ml) significantly showed the steeper decline of HCV RNA level than those with Gln70 (His70) and/or Met91 (median 2.7 log IU/ml; range 1.6–3.5 log IU/ml). **b** Very early dynamics according to genetic variation near the *IL28B* gene. After 24 h of commencement of the triple therapy, the decline of HCV RNA level of patients with rs8099917 genotype TT (median 2.8 log IU/ml; range 1.6–4.0 log IU/ml) was not significantly different from that of patients with genotype TG and GG (median 2.7 log IU/ml; range 2.2–3.5 log IU/ml).

multiple logistic regression analysis to identify significant independent predictive factors. Each variable was transformed into categorical data consisting of two simple ordinal numbers for univariate and multivariate analyses. The potential pretreatment factors associated with treatment efficacy included the following variables: sex, age, history of blood transfusion, familial history of liver disease, body mass index, aspartate aminotransferase (AST), alanine aminotransferase (ALT), albumin, gamma-glutamyl transpeptidase ( $\gamma$ GTP), leukocyte count, hemoglobin, platelet count, HCV RNA level,  $\alpha$ -fetoprotein, total cholesterol, fasting blood sugar, PEG-IFN dose/body weight, ribavirin dose/body

weight, telaprevir dose/day, treatment regimen of triple therapy, past history of IFN therapy, genetic variation near the *IL28B* gene, and amino acid substitution in the core region, and NS5A-ISDR. Statistical analyses were performed using the SPSS software (SPSS Inc., Chicago, Ill., USA).

## Results

### *Virological Response to Therapy and Loss of HCV RNA during Treatment*

Sustained virological response was achieved by 63.6% (49 of 77 patients). The disappearance rate of HCV RNA during treatment was 0% (0 of 80), 1.3% (1 of 80), 33.8% (27 of 80), 81.0% (64 of 79), 90.7% (68 of 75), 94.6% (70 of 74), and 89.9% (62 of 69) at 24 hours, 1, 2, 4, 6, 8, and 12 weeks, respectively.

### *Very Early Dynamics according to Amino Acid Substitutions in Core Region and Genetic Variation near the *IL28B* Gene*

After 24 h of commencement of the triple therapy, the proportion of patients with Arg70 and Leu91 substitutions who showed  $\geq 3.0$  log drop in HCV RNA level (45.2%; 14 of 31 patients) was significantly higher than that of patients with Gln70 (His70) and/or Met91 (14.3%; 7 of 49) ( $p = 0.004$ ). Thus, patients with Arg70 and Leu91 (median 2.9 log IU/ml; range 1.8–4.0 log IU/ml) significantly showed the steeper decline of HCV RNA level than those with Gln70 (His70) and/or Met91 (median 2.7 log IU/ml; range 1.6–3.5 log IU/ml) (fig. 1a).

After 24 h of commencement of treatment, the proportion of patients with rs8099917 genotype TT who showed  $\geq 3.0$  log drop in HCV RNA level (30.4%; 14 of 46 patients) was not significantly different from that of patients with genotype TG and GG (21.9%; 7 of 32). Thus, the decline of HCV RNA level of patients with genotype TT (median 2.8 log IU/ml; range 1.6–4.0 log IU/ml) was not significantly different from that of patients with genotype TG and GG (median 2.7 log IU/ml; range 2.2–3.5 log IU/ml) (fig. 1b).

Hence, the fall in HCV RNA level at 24 h was influenced by aa substitution patterns in the core region, but was independent of genetic variation near *IL28B* gene.

### *Rates of Loss of HCV RNA according to Amino Acid Substitutions in Core Region and Genetic Variation near the *IL28B* Gene*

According to the substitution of core aa 70 and 91, the rate of HCV RNA loss of patients with Arg70 and Leu91 was not significantly different from that of patients with