

1  
2  
3 antibiotic that binds to the GC-box in the promoter to block binding of Sp1 or other Sp family proteins  
4 [Blume et al., 1991]. As shown in Fig. 5B, MMA (2.5, 5.0, and 10  $\mu$ M) inhibited complex formation in a  
5 dose-dependent manner. In contrast, the formation of DNA–protein complexes with the MZF-1 probe  
6 was not affected by the addition of MMA, suggesting that the inhibition mediated by MMA was specific  
7 for the GC box–Sp1, and that complex formation with the response element requires the Sp1 binding site.  
8 A supershift assay using nuclear extract from HepG2 cells transfected with a Myc-tagged Sp1 expression  
9 vector and anti-Myc was also performed (Suppl. Fig. 2). The mobility of the HCV response element and  
10 the Sp1 consensus sequence was supershifted partially by addition of anti-Myc (lanes 3 and 6). The  
11 effect of silencing the expression of Sp1 with small interfering RNA (siRNA) was analyzed by EMSA  
12 using nuclear extracts from Sp1-knockdown RzM6-0d and RzM6-LC cells (Fig. 5C). DNA–protein  
13 complexes with the response element or the Sp1 probe were not observed (lanes 2, 4, 6, and 8); however,  
14 formation of DNA–MZF-1 complexes was not influenced by siRNA treatment (lanes 9–12).  
15 Immunoblotting was used to confirm efficient silencing of the Sp1 protein in cells used to generate the  
16 nuclear extracts (Fig. 5D). A significant decrease in the expression of DHCR24 was observed in the  
17 cytosolic fraction from RzM6 cells transfected with siRNA specific for Sp1 (Fig. 5D). Thus, these results  
18 indicate that Sp1, but neither AP-2 $\alpha$  nor MZF-1, bound to the HCV response element, and that Sp1 may  
19 play an important role in the transcriptional regulation of *DHCR24*.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

39  
40 ***Transcriptional regulation of DHCR24 through the HCV response element is mediated by oxidative***  
41 ***stress***  
42

43 DHCR24 functions as a mediator of the cellular response to oxidative stress [Greeve et al., 2000;  
44 Benvenuti et al., 2005; Di Stasi et al., 2005; Luciani et al., 2005; Lu et al., 2008] and is a hydrogen  
45 peroxide scavenger [Lu et al., 2008]. Expression of the *DHCR24* gene is also induced in response to  
46 oxidative stress [Wu et al., 2004; Benvenuti et al., 2006; Kuehnle et al., 2008]. Expression of the HCV  
47 gene elevates the level of reactive oxygen species (ROS) via dysregulation of ER-mediated calcium  
48 homeostasis, which results in oxidative stress [Tardif et al., 2005]. Therefore, the role of oxidative stress  
49 induced by HCV in the regulatory mechanism of the expression of *DHCR24* was examined. HepG2 cells  
50 were treated with hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and transfected with reporter plasmids containing the  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 *DHCR24* promoter deletion mutants. Measurement of promoter activity revealed a significant increase in  
4 transcription in response to oxidative stress (H<sub>2</sub>O<sub>2</sub>) for *DHCR24* promoters containing the HCV response  
5 element (−4976/+113, −2982/+113, −515/+113, and −167/+113) but not for the promoter lacking the  
6 response element (−144/+113; Fig. 6A). Therefore, enhanced transcription in response to oxidative stress  
7 by reporter constructs containing the *DHCR24* promoter may be mediated through the HCV response  
8 element. The formation of complexes containing the response element or Sp1 probe was increased  
9 markedly in the nuclear extracts from the H<sub>2</sub>O<sub>2</sub>-treated HepG2 cells (Fig. 6B) or other hepatic cell lines  
10 (Suppl. Fig. 3), suggesting that oxidative stress enhances the binding affinity of Sp1 to the HCV response  
11 element.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22

#### 23 ***Overexpression of DHCR24 in M6-LC cells is blocked by an ROS scavenger***

24 The increase in the expression of DHCR24 induced by oxidative stress can be blocked by treatment with  
25 an ROS scavenger, *N*-acetylcysteine (NAC) [Wu et al., 2004], which is a precursor of the potent  
26 biological antioxidant glutathione. The H<sub>2</sub>O<sub>2</sub>-induced overexpression of DHCR24 was inhibited by  
27 pre-treatment with NAC and blocked partially by NAC treatment after the induction of oxidative stress  
28 (~50% suppression; Fig. 7A). The enhanced expression of DHCR24 in RzM6-LC cells decreased after  
29 12 or 24 h of treatment with NAC without influencing the level of expression of HCV, suggesting that  
30 overexpression of DHCR24 in cells expressing HCV is mediated through oxidative stress.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41

#### 42 ***Overexpression and enhanced phosphorylation of Sp1 in the cells expressing HCV***

43 Sp1 is a transcription factor that is activated in response to a variety of cellular stressors, including  
44 oxidative stress [Schafer et al., 2003; Chu and Ferro, 2006; Dasari et al., 2006; Qin et al., 2009; Lin et al.,  
45 2011]. Thus, Sp1 may play an important role in linking oxidative stress and augmentation of *DHCR24*  
46 transcription in cells expressing HCV. Sp1 was overexpressed significantly in RzM6-LC cells treated  
47 with H<sub>2</sub>O<sub>2</sub> compared with the control cells (Fig. 8A). Phosphorylation of Sp1 at Ser101 was also  
48 elevated under oxidative stress. Both the basal level and phosphorylation status of nuclear Sp1 were  
49 higher in the presence of HCV (RzM6-LC cells) than in the absence of HCV (RzM6-0d cells; Fig. 8B).  
50  
51  
52  
53  
54  
55  
56  
57

58 Phosphorylation of Sp1 at Ser101 is a target of the DNA damage signaling pathway mediated by ATM  
59  
60

1  
2  
3 (ataxia telangiectasia mutated) and ATR (ATM and Rad3-related) kinases [Olofsson et al., 2007; Iwahori  
4 et al., 2008]. As shown in Fig. 8C, phosphorylation of Sp1 at Ser101 was no longer detectable following  
5 pre-treatment with an ATM kinase inhibitor (KU55933) before exposure to H<sub>2</sub>O<sub>2</sub>. In contrast,  
6 phosphorylation was not affected by other kinase inhibitors (phosphatidylinositol-3 kinase inhibitor,  
7 LY294002 or MEK1 inhibitor, PD98059). Similarly, phosphorylation of Sp1 at Thr453, which is  
8 important for transcriptional activation of Sp1 [Milanini-Mongiat et al., 2002; D'Addario et al., 2006;  
9 Hsu et al., 2006; Lin et al., 2011], was not seen in response to oxidative stress following treatment with  
10 KU55933 (Fig. 8C). The induction of expression of DHCR24 after H<sub>2</sub>O<sub>2</sub> exposure was suppressed  
11 significantly by treatment with KU55933 or NAC, which corresponds with inhibition of Sp1  
12 phosphorylation. In the presence of MMA, the phosphorylation of Sp1 was not inhibited. However, since  
13 MMA blocks the binding of Sp1 [Blume et al., 1991], the induction of expression of DHCR24 by H<sub>2</sub>O<sub>2</sub>  
14 was inhibited. Impairment of DHCR24 induction by H<sub>2</sub>O<sub>2</sub> was also observed after treatment with  
15 siRNAs targeting ATM (Suppl. Fig. 4).  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

29 Studies on the relationship between HCV and ATM have reported that the interaction of NS3/4A with  
30 ATM results in delayed de-phosphorylation of both phosphorylated ATM and phosphorylated histone  
31 H2AX at Ser139 ( $\gamma$ H2AX), which acts as a substrate for ATM in response to DNA damage [Lai et al.,  
32 2008]. In the present study, delayed de-phosphorylation of  $\gamma$ H2AX was also observed in HCV replicon  
33 cells (Suppl. Fig. 5), which corresponded with increased phosphorylation of the H2AX Ser139 residue in  
34 cells expressing HCV (Fig. 8). Similarly, phosphorylation of ATM was sustained in HCV replicon cells  
35 (Suppl. Fig. 6). Therefore, DNA repair may be impaired in cells expressing or replicating HCV, resulting  
36 in sustained DNA damage. As a result, downstream substrates such as Sp1 Ser101 and Thr453 residues  
37 or the H2AX Ser139 residue may be phosphorylated to a greater extent in cells expressing HCV  
38 compared with control cells in the basal state or cells under oxidative stress (Fig. 8A and 8B).  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

50 Taken together, these results indicate that the oxidative stress induced by HCV may produce  
51 quantitative as well as qualitative activation of Sp1, thereby resulting in augmentation of *DHCR24*  
52 transcription.  
53  
54  
55  
56  
57  
58  
59  
60

### Discussion

HCV establishes chronic infection and induces persistent overexpression of DHCR24 in human hepatocytes [Nishimura et al., 2009]. HCV also confers resistance to the apoptosis induced by oxidative stress and suppresses p53 activity by blocking nuclear p53 acetylation and increasing the interaction between p53 and HDM2 (p53-specific E3 ligase) in the cytoplasm, which may be mediated by inhibition of p53 degradation. Thus, the augmentation of DHCR24 by HCV reflects the tumorigenicity of hepatocytes. The present study identified the genomic region of *DHCR24* that is responsive to HCV, and showed that this response is mediated through the activation of Sp1 induced by oxidative stress. In general, expression of the HCV gene elevates the levels of ROS through dysregulation of ER-mediated calcium homeostasis [Tardif et al., 2005]. In healthy cells, ROS usually exist in equilibrium with antioxidants that scavenge ROS and prevent cellular injury. However, this critical balance may be disrupted in the cells infected with HCV, resulting in the accumulation of ROS and the development of constitutive oxidative stress.

Sp1 is a member of the Sp/KLF family of transcription factors that bind to GC elements of promoters [Black et al., 2001; Kaczynski et al., 2003; Chu and Ferro, 2005; Li and Davie, 2010]. Under a variety of endogenous and exogenous stimuli—including oxidative stress and DNA damage—activation of Sp1 may be mediated via induction of expression of Sp1 and post-translational modifications such as acetylation, sumoylation, O-linked glycosylation, and phosphorylation. Sp1 is phosphorylated by several kinases, including DNA-dependent protein kinase, casein kinase II, and cyclin A/cdk2, which exert both positive and negative effects on transcription [Jackson et al., 1990; Armstrong et al., 1997; Fojas de Borja et al., 2001; Ryu et al., 2003]. Sp1 is the only Sp/KLF family member to contain putative consensus SQ/TQ cluster domains within the transactivation domains, which suggests that Sp1 is a substrate of the PI3K-related kinases, e.g., ATM, DNA-dependent protein kinase, and ATR. Indeed, Sp1 is a target of the ATM-dependent DNA damage response pathway [Iwahori et al., 2007; Olofsson et al., 2007; Iwahori et al., 2008]. ATM plays a central role in orchestrating molecular events involved in double-strand break signaling, which is mediated via the phosphorylation of a variety of substrate proteins—including p53 and BRCA1 transcription factors—involved in the DNA damage response. As a result, these phosphorylation events lead to cell cycle checkpoint activation, DNA repair, altered gene

1  
2  
3 expression patterns, and/or apoptosis [Shiloh, 2006].  
4

5 Given the role of Sp1 in oxidative stress [Schafer et al., 2003; Chu and Ferro, 2006; Dasari et al., 2006;  
6 Rojo et al., 2006; Qin et al., 2009; Lin et al., 2010], Sp1 may be regulated by the oxidative stress induced  
7 by HCV and the subsequent phosphorylation, which depends on ATM. However, little is known  
8 regarding the regulation of Sp1 in response to DNA damage. Although the precise role of  
9 phosphorylation of Sp1 at Ser101 in the DNA damage response is unclear, the similar kinetics of Sp1 and  
10  $\gamma$ H2AX phosphorylation [Olofsson et al., 2007] suggest that Sp1 is an early target of the DNA damage  
11 response pathway. Thus, Sp1 may be involved in modulating the cellular response to DNA damage to  
12 prevent cell death [Ryu et al., 2003]. Phosphorylation of Sp1 at Ser101 and histone H2AX, which occurs  
13 in parallel in response to oxidative stress, was enhanced in cells expressing HCV compared with that  
14 observed in control cells (Fig. 8A). Interestingly, augmentation of Sp1 phosphorylation in parallel with  
15 histone H2AX phosphorylation was also detected for cells expressing HCV in the basal state (Fig. 8A  
16 and 8B), which may be primarily due to the increase in endogenous Sp1 protein (Fig. 8A and 8B). In  
17 support of these results, enhanced phosphorylation of Ser101 on Sp1 occurs upon HSV-1 infection, and  
18 is mediated by ATM [Iwahori et al., 2007]. Thus, increased phosphorylation of Sp1 and  $\gamma$ H2AX in cells  
19 expressing HCV is likely to reflect the higher activity of ATM, which may result from the accumulation  
20 and frequency of DNA damage caused by increased generation of endogenous ROS.  
21  
22

23 Oxidative stress is a common mechanism of liver injury [Loguercio and Federico, 2003] and is  
24 mediated by the direct effects of ROS on signal transduction pathways, including extracellular  
25 signal-regulated kinase 1/2 (ERK1/2), c-Jun N-terminal kinase (JNK), and p38 mitogen-activated protein  
26 kinases (MAPKs), which act as downstream kinases in the MAPK cascade to phosphorylate Sp1  
27 Thr453/739 residues [Milanini-Mongiat et al., 2002; D'Addario et al., 2006; Hsu et al., 2006; Chuang et  
28 al., 2008; Lin et al., 2011]. These signal transduction pathways are also stimulated by oxidative stress in  
29 the hepatic cells expressing or replicating HCV, [Qadri et al., 2004; Burdette et al., 2010; Lin et al.,  
30 2010]. Therefore, oxidative stress in response to HCV may induce downstream signaling pathways, such  
31 as ERK1/2, JNK, and p38 MAPK as well as ATM/ATR, to activate Sp1 via post-translational  
32 modifications.  
33  
34

35 Sp1 is a host factor activated by several viral proteins, including HIV-1 Vpr, and HTLV-1 Tax [Peng et  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 al., 2003; Amini et al., 2004; Chang et al., 2005; Zhang et al., 2009]. The HCV core and NS5A proteins  
4 also activate Sp1 [Lee et al., 2001; Xiang et al., 2010]. The HCV core upregulates the DNA-binding  
5 activity and phosphorylation of Sp1 [Lee et al., 2001], and NS5A may also exert a similar effect on Sp1  
6 activity. However, a physical interaction between these proteins and Sp1 has not yet been demonstrated.  
7  
8 Both HCV core and NS5A proteins have a high potential for oxidative stress induction  
9  
10 [Garcia-Mediavilla et al., 2005; Dionisio et al., 2009], which may mediate activation of Sp1. On the  
11  
12 other hand, individual viral proteins were insufficient to increase the expression of DHCR24 (Fig. 1A).  
13  
14 Therefore, in addition to induction of oxidative stress by each viral protein, the persistence of the  
15  
16 signaling pathways induced by oxidative stress, e.g., ATM (Suppl. Fig. 6), may also be required for the  
17  
18 Sp1-mediated increase in the expression of DHCR24.  
19  
20  
21  
22

23 The results of the present study revealed that knockdown of expression of Sp1 almost completely  
24 blocked the enhanced expression of DHCR24. Sp1 is expressed ubiquitously in various mammalian cells  
25 and is involved in regulating the transcriptional activity of genes implicated in many cellular processes  
26 [Black et al., 2001; Kaczynski et al., 2003; Chu and Ferro, 2005; Li and Davie, 2010]. Thus, Sp1 may  
27 represent an essential master regulator among the myriad of transcription factors involved in the direct  
28 regulation of *DHCR24* transcription.  
29  
30  
31  
32  
33  
34

35 In conclusion, HCV was shown to enhance the expression of DHCR24 via the activation of Sp1, which  
36 may shed light on the mechanism of tumorigenesis associated with HCV.  
37  
38  
39  
40  
41

#### 42 Acknowledgements

43 The authors would like to thank Ms. Ryoko Takehara and Yuri Kasama for their technical support, Yuko  
44 Tokunaga for her comments, and Dr. Chieko Kai for her support. This work was supported by grants  
45 from the Ministry of Health and Welfare of Japan; Ministry of Education, Culture, Sports, Science and  
46 Technology of Japan; Program for Promotion of Fundamental Studies in Health Sciences of the National  
47 Institute of Biomedical Innovation; and Cooperative Research Project on Clinical and Epidemiological  
48 Studies of Emerging and Re-emerging Infectious Diseases.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References

- 1  
2  
3  
4  
5 Amini S, Saunders M, Kelley K, Khalili K, Sawaya BE. 2004. Interplay between HIV-1 Vpr and Sp1  
6 modulates p21(WAF1) gene expression in human astrocytes. *J Biol Chem* 279:46046–46056.
- 7 Armstrong SA, Barry DA, Leggett RW, Mueller CR. 1997. Casein kinase II-mediated  
8 phosphorylation of the C terminus of Sp1 decreases its DNA binding activity. *J Biol Chem*  
9 272:13489–13495.
- 10 Benvenuti S, Luciani P, Vannelli GB, Gelmini S, Franceschi E, Serio M, Peri A. 2005. Estrogen and  
11 selective estrogen receptor modulators exert neuroprotective effects and stimulate the  
12 expression of selective Alzheimer's disease indicator-1, a recently discovered antiapoptotic  
13 gene, in human neuroblast long-term cell cultures. *J Clin Endocrinol Metab* 90:1775–1782.
- 14 Benvenuti S, Saccardi R, Luciani P, Urbani S, Deledda C, Cellai I, Francini F, Squecco R, Rosati F,  
15 Danza G, Gelmini S, Greeve I, Rossi M, Maggi R, Serio M, Peri A. 2006. Neuronal  
16 differentiation of human mesenchymal stem cells: changes in the expression of the  
17 Alzheimer's disease-related gene seladin-1. *Exp Cell Res* 312:2592–2604.
- 18 Black AR, Black JD, Azizkhan-Clifford J. 2001. Sp1 and kruppel-like factor family of transcription  
19 factors in cell growth regulation and cancer. *J Cell Physiol* 188:143–160.
- 20 Blume SW, Snyder RC, Ray R, Thomas S, Koller CA, Miller DM. 1991. Mithramycin inhibits SP1  
21 binding and selectively inhibits transcriptional activity of the dihydrofolate reductase gene  
22 in vitro and in vivo. *J Clin Invest* 88:1613–1621.
- 23 Bruchfeld A, Stahle L, Andersson J, Schvarcz R. 2001. Ribavirin treatment in dialysis patients with  
24 chronic hepatitis C virus infection—a pilot study. *J Viral Hepat* 8:287–292.
- 25 Burdette D, Olivarez M, Waris G. 2010. Activation of transcription factor Nrf2 by hepatitis C virus  
26 induces the cell-survival pathway. *J Gen Virol* 91:681–690.
- 27 Chang LK, Chung JY, Hong YR, Ichimura T, Nakao M, Liu ST. 2005. Activation of Sp1-mediated  
28 transcription by Rta of Epstein-Barr virus via an interaction with MCAF1. *Nucleic Acids Res*  
29 33:6528–6539.
- 30 Chu S, Ferro TJ. 2005. Sp1: regulation of gene expression by phosphorylation. *Gene* 348:1–11.
- 31 Chu S, Ferro TJ. 2006. Identification of a hydrogen peroxide-induced PP1-JNK1-Sp1 signaling  
32 pathway for gene regulation. *Am J Physiol Lung Cell Mol Physiol* 291:L983–992.
- 33 Chuang JY, Wang YT, Yeh SH, Liu YW, Chang WC, Hung JJ. 2008. Phosphorylation by c-Jun  
34 NH2-terminal kinase 1 regulates the stability of transcription factor Sp1 during mitosis. *Mol*  
35 *Biol Cell* 19:1139–1151.
- 36 Crameri A, Biondi E, Kuehnle K, Lutjohann D, Thelen KM, Perga S, Dotti CG, Nitsch RM, Ledesma  
37 MD, Mohajeri MH. 2006. The role of seladin-1/DHCR24 in cholesterol biosynthesis, APP  
38 processing and Abeta generation in vivo. *EMBO J* 25:432–443.
- 39 D'Addario M, Arora PD, McCulloch CA. 2006. Role of p38 in stress activation of Sp1. *Gene*  
40 379:51–61.
- 41 Dasari A, Bartholomew JN, Volonte D, Galbiati F. 2006. Oxidative stress induces premature  
42 senescence by stimulating caveolin-1 gene transcription through p38 mitogen-activated  
43 protein kinase/Sp1-mediated activation of two GC-rich promoter elements. *Cancer Res*  
44 66:10805–10814.
- 45 Di Stasi D, Vallacchi V, Campi V, Ranzani T, Daniotti M, Chiodini E, Fiorentini S, Greeve I, Prinetti  
46 A, Rivoltini L, Pierotti MA, Rodolfo M. 2005. DHCR24 gene expression is upregulated in  
47 melanoma metastases and associated to resistance to oxidative stress-induced apoptosis. *Int*  
48 *J Cancer* 115:224–230.
- 49 Dignam JD, Lebovitz RM, Roeder RG. 1983. Accurate transcription initiation by RNA polymerase II  
50 in a soluble extract from isolated mammalian nuclei. *Nucleic Acids Res* 11:1475–1489.
- 51 Dionisio N, Garcia-Mediavilla MV, Sanchez-Campos S, Majano PL, Benedicto I, Rosado JA, Salido  
52 GM, Gonzalez-Gallego J. 2009. Hepatitis C virus NS5A and core proteins induce oxidative  
53 stress-mediated calcium signalling alterations in hepatocytes. *J Hepatol* 50:872–882.
- 54 Fojas de Borja P, Collins NK, Du P, Azizkhan-Clifford J, Mudryj M. 2001. Cyclin A-CDK  
55 phosphorylates Sp1 and enhances Sp1-mediated transcription. *EMBO J* 20:5737–5747.
- 56 García-Mediavilla MV, Sánchez-Campos S, González-Pérez P, Gómez-Gonzalo M, Majano PL,  
57 Lopez-Cabrera M, Clemente G, García-Monzon C, González-Gallego J. 2005. Differential  
58 contribution of hepatitis C virus NS5A and core proteins to the induction of oxidative and  
59 nitrosative stress in human hepatocyte-derived cells. *J Hepatol* 43:606–613.
- 60 Greeve I, Hermans-Borgmeyer I, Brellinger C, Kasper D, Gomez-Isla T, Behl C, Levkau B, Nitsch

- 1  
2  
3 RM. 2000. The human DIMINUTO/DWARF1 homolog seladin-1 confers resistance to  
4 Alzheimer's disease-associated neurodegeneration and oxidative stress. *J Neurosci*  
5 20:7345–7352.
- 6 Hsu MC, Chang HC, Hung WC. 2006. HER-2/neu represses the metastasis suppressor RECK via  
7 ERK and Sp transcription factors to promote cell invasion. *J Biol Chem* 281:4718–4725.
- 8 Iwahori S, Shirata N, Kawaguchi Y, Weller SK, Sato Y, Kudoh A, Nakayama S, Isomura H, Tsurumi  
9 T. 2007. Enhanced phosphorylation of transcription factor sp1 in response to herpes simplex  
10 virus type 1 infection is dependent on the ataxia telangiectasia-mutated protein. *J Virol*  
11 81:9653–9664.
- 12 Iwahori S, Yasui Y, Kudoh A, Sato Y, Nakayama S, Murata T, Isomura H, Tsurumi T. 2008.  
13 Identification of phosphorylation sites on transcription factor Sp1 in response to DNA  
14 damage and its accumulation at damaged sites. *Cell Signal* 20:1795–1803.
- 15 Jackson SP, MacDonald JJ, Lees-Miller S, Tjian R. 1990. GC box binding induces phosphorylation of  
16 Sp1 by a DNA-dependent protein kinase. *Cell* 63:155–165.
- 17 Kaczynski J, Cook T, Urrutia R. 2003. Sp1- and Kruppel-like transcription factors. *Genome Biol*  
18 4:206.
- 19 Kohara M, Tanaka T, Tsukiyama-Kohara K, Tanaka S, Mizokami M, Lau JY, Hattori N. 1995.  
20 Hepatitis C virus genotypes 1 and 2 respond to interferon-alpha with different virologic  
21 kinetics. *J Infect Dis* 172:934–938.
- 22 Koike K. 2007. Hepatitis C virus contributes to hepatocarcinogenesis by modulating metabolic and  
23 intracellular signaling pathways. *J Gastroenterol Hepatol* 22 Suppl 1:S108–111.
- 24 Kuehnle K, Crameri A, Kalin RE, Luciani P, Benvenuti S, Peri A, Ratti F, Rodolfo M, Kulic L,  
25 Heppner FL, Nitsch RM, Mohajeri MH. 2008. Prosurvival effect of DHCR24/Seladin-1 in  
26 acute and chronic responses to oxidative stress. *Mol Cell Biol* 28:539–550.
- 27 Lai CK, Jeng KS, Machida K, Cheng YS, Lai MM. 2008. Hepatitis C virus NS3/4A protein interacts  
28 with ATM, impairs DNA repair and enhances sensitivity to ionizing radiation. *Virology*  
29 370:295–309.
- 30 Lavanchy D. 2009. The global burden of hepatitis C. *Liver Int* 29 Suppl 1:74–81.
- 31 Lee S, Park U, Lee YI. 2001. Hepatitis C virus core protein transactivates insulin-like growth factor  
32 II gene transcription through acting concurrently on Egr1 and Sp1 sites. *Virology*  
33 283:167–177.
- 34 Li L, Davie JR. 2010. The role of Sp1 and Sp3 in normal and cancer cell biology. *Ann Anat*  
35 192:275–283.
- 36 Lin HH, Lai SC, Chau LY. 2011. Heme oxygenase-1/carbon monoxide induces vascular endothelial  
37 growth factor expression via p38 kinase-dependent activation of Sp1. *J Biol Chem*  
38 286:3829–3838.
- 39 Lin W, Tsai WL, Shao RX, Wu G, Peng LF, Barlow LL, Chung WJ, Zhang L, Zhao H, Jang JY, Chung  
40 RT. 2010. Hepatitis C virus regulates transforming growth factor beta1 production through  
41 the generation of reactive oxygen species in a nuclear factor kappaB-dependent manner.  
42 *Gastroenterology* 138:2509–2518.
- 43 Loguercio C, Federico A. 2003. Oxidative stress in viral and alcoholic hepatitis. *Free Radic Biol Med*  
44 34:1–10.
- 45 Lu X, Kambe F, Cao X, Kozaki Y, Kaji T, Ishii T, Seo H. 2008. 3beta-Hydroxysteroid-delta24  
46 reductase is a hydrogen peroxide scavenger, protecting cells from oxidative stress-induced  
47 apoptosis. *Endocrinology* 149:3267–3273.
- 48 Luciani P, Gelmini S, Ferrante E, Lania A, Benvenuti S, Baglioni S, Mantovani G, Cellai I,  
49 Ammannati F, Spada A, Serio M, Peri A. 2005. Expression of the antiapoptotic gene seladin-1  
50 and octreotide-induced apoptosis in growth hormone-secreting and nonfunctioning pituitary  
51 adenomas. *J Clin Endocrinol Metab* 90:6156–6161.
- 52 Milanini-Mongiat J, Pouyssegur J, Pages G. 2002. Identification of two Sp1 phosphorylation sites for  
53 p42/p44 mitogen-activated protein kinases: their implication in vascular endothelial growth  
54 factor gene transcription. *J Biol Chem* 277:20631–20639.
- 55 Nakamura H, Ogawa H, Kuroda T, Yamamoto M, Enomoto H, Kishima Y, Yoshida K, Ito H, Matsuda  
56 M, Noguchi S. 2002. Interferon treatment for patients with chronic hepatitis C infected with  
57 high viral load of genotype 2 virus. *Hepatogastroenterology* 49:1373–1376.
- 58 Nishimura T, Kohara M, Izumi K, Kasama Y, Hirata Y, Huang Y, Shuda M, Mukaidani C, Takano T,  
59 Tokunaga Y, Nuriya H, Satoh M, Saito M, Kai C, Tsukiyama-Kohara K. 2009. Hepatitis C  
60 virus impairs p53 via persistent overexpression of 3beta-hydroxysterol Delta24-reductase. *J*



- 1  
2  
3 Biol Chem 284:36442–36452.
- 4 Olofsson BA, Kelly CM, Kim J, Hornsby SM, Azizkhan-Clifford J. 2007. Phosphorylation of Sp1 in  
5 response to DNA damage by ataxia telangiectasia-mutated kinase. *Mol Cancer Res*  
6 5:1319–1330.
- 7 Peng H, He H, Hay J, Ruyechan WT. 2003. Interaction between the varicella zoster virus IE62 major  
8 transactivator and cellular transcription factor Sp1. *J Biol Chem* 278:38068–38075.
- 9 Qadri I, Iwahashi M, Capasso JM, Hopken MW, Flores S, Schaack J, Simon FR. 2004. Induced  
10 oxidative stress and activated expression of manganese superoxide dismutase during  
11 hepatitis C virus replication: role of JNK, p38 MAPK and AP-1. *Biochem J* 378:919–928.
- 12 Qin K, Zhao L, Ash RD, McDonough WF, Zhao RY. 2009. ATM-mediated transcriptional elevation of  
13 prion in response to copper-induced oxidative stress. *J Biol Chem* 284:4582–4593.
- 14 Rojo AI, Salina M, Salazar M, Takahashi S, Suske G, Calvo V, de Sagarra MR, Cuadrado A. 2006.  
15 Regulation of heme oxygenase-1 gene expression through the phosphatidylinositol  
16 3-kinase/PKC-zeta pathway and Sp1. *Free Radic Biol Med* 41:247–261.
- 17 Ryu H, Lee J, Zaman K, Kubilis J, Ferrante RJ, Ross BD, Neve R, Ratan RR. 2003. Sp1 and Sp3 are  
18 oxidative stress-inducible, antideath transcription factors in cortical neurons. *J Neurosci*  
19 23:3597–3606.
- 20 Schafer G, Cramer T, Suske G, Kemmner W, Wiedenmann B, Hocker M. 2003. Oxidative stress  
21 regulates vascular endothelial growth factor-A gene transcription through Sp1- and  
22 Sp3-dependent activation of two proximal GC-rich promoter elements. *J Biol Chem*  
23 278:8190–8198.
- 24 Shiloh Y. 2006. The ATM-mediated DNA-damage response: taking shape. *Trends in Biochem Sci*  
25 31:402–410.
- 26 Takano T, Kohara M, Kasama Y, Nishimura T, Saito M, Kai C, Tsukiyama-Kohara K. 2011a.  
27 Translocase of outer mitochondrial membrane 70 expression is induced by hepatitis C virus  
28 and is related to the apoptotic response. *J Med Virol* 83:801–809.
- 29 Takano T, Tsukiyama-Kohara K, Hayashi M, Hirata Y, Satoh M, Tokunaga Y, Tateno C, Hayashi Y,  
30 Hishima T, Funata N, Sudo M, Kohara M. 2011b. Augmentation of DHCR24 expression by  
31 hepatitis C virus infection facilitates viral replication in hepatocytes. *J Hepatol* 55:512–521
- 32 Tardif KD, Waris G, Siddiqui A. 2005. Hepatitis C virus, ER stress, and oxidative stress. *Trends*  
33 *Microbiol* 13:159–163.
- 34 Tsukiyama-Kohara K, Tone S, Maruyama I, Inoue K, Katsume A, Nuriya H, Ohmori H, Ohkawa J,  
35 Taira K, Hoshikawa Y, Shibasaki F, Reth M, Minatogawa Y, Kohara M. 2004. Activation of  
36 the CKI-CDK-Rb-E2F pathway in full genome hepatitis C virus-expressing cells. *J Biol*  
37 *Chem* 279:14531–14541.
- 38 Wakita T, Pietschmann T, Kato T, Date T, Miyamoto M, Zhao Z, Murthy K, Habermann A, Krausslich  
39 HG, Mizokami M, Bartenschlager R, Liang TJ. 2005. Production of infectious hepatitis C  
40 virus in tissue culture from a cloned viral genome. *Nat Med* 11:791–796.
- 41 Waterham HR, Koster J, Romeijn GJ, Hennekam RC, Vreken P, Andersson HC, FitzPatrick DR,  
42 Kelley RI, Wanders RJ. 2001. Mutations in the 3beta-hydroxysterol Delta24-reductase gene  
43 cause desmosterolosis, an autosomal recessive disorder of cholesterol biosynthesis. *Am J*  
44 *Hum Genet* 69:685–694.
- 45 Williams T, Admon A, Luscher B, Tjian R. 1988. Cloning and expression of AP-2, a cell-type-specific  
46 transcription factor that activates inducible enhancer elements. *Genes Dev* 2:1557–1569.
- 47 Wu C, Miloslavskaya I, Demontis S, Maestro R, Galaktionov K. 2004. Regulation of cellular response  
48 to oncogenic and oxidative stress by Seladin-1. *Nature* 432:640–645.
- 49 Xiang Z, Qiao L, Zhou Y, Babiuk LA, Liu Q. 2010. Hepatitis C virus nonstructural protein-5A  
50 activates sterol regulatory element-binding protein-1c through transcription factor Sp1.  
51 *Biochem Biophys Res Commun* 402:549–553.
- 52 Zhang L, Zhi H, Liu M, Kuo YL, Giam CZ. 2009. Induction of p21(CIP1/WAF1) expression by human  
53 T-lymphotropic virus type 1 Tax requires transcriptional activation and mRNA stabilization.  
54 *Retrovirology* 6:35.
- 55  
56  
57  
58  
59  
60

1  
2  
3 **Figure legends**  
4

5 **Fig. 1.** *DHCR24* expression is induced in the presence of the complete HCV genome. (A) The amount of  
6 HCV RNA in chimeric mice infected with HCV-R6 (genotype 1b) was quantified by qPCR (left panel).  
7 The amount of *DHCR24* mRNA was measured, and the ratio to the amount of 18S rRNA was calculated  
8 in the tissues (right panel). (B) Western blotting of *DHCR24*, HCV core, and actin protein in RzM6-0d  
9 and LC cells following treatment with the indicated siRNA. (C) Level of *DHCR24* mRNA expression in  
10 cell lines with stable expression of individual HCV proteins, the HCV open reading frame, or the  
11 complete HCV genome. Total RNA from HepG2/Lenti cell lines (emp, core, E1, E2, NS2, NS3/4A,  
12 NS4B, NS5A, and NS5B), CN5 cell lines (CN5-Hep, CN5-Swa, and CN5-Cre), or RzM6 cell lines  
13 (RzM6-0d and RzM6-LC) were prepared, and reverse transcription was performed. Synthesized cDNA  
14 was subjected to quantitative PCR. The level of expression of *DHCR24* mRNA for each sample was  
15 normalized to that of *GAPDH* mRNA and represented as a ratio of HepG2-emp ( $*p < 0.05$ ).  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

29 **Fig. 2.** *DHCR24* promoter activity is augmented by the expression of HCV. (A) The 5'-flanking region of  
30 *DHCR24* was cloned from HepG2 and inserted upstream of the firefly luciferase gene in pGL3  
31 (−4976/+113). A deletion series of the 5'-flanking regions was also constructed (−2982/+113,  
32 −2293/+113, and −515/+113). The black boxes indicate potential binding sites for the ER  
33 stress-responsive transcription factor, XBP-1. TSS, transcription start site (+1). (B) HepG2 cells ( $1 \times 10^4$   
34 cells/well in a 96-well plate) were co-transfected with each *DHCR24* promoter reporter plasmid (0.25  
35 μg/well), a Renilla luciferase expression vector (pRL-TK; 0.025 μg/well), and either an expression  
36 vector containing the HCV full-length genome (pCA-Rz; 0.5 μg/well) or an empty expression vector  
37 (pCAGGS). Luciferase activity at 48 h post-transfection is shown as the ratio of firefly luciferase (FL) to  
38 Renilla luciferase (RL). Data are shown as the mean ± SD from 2 representative experiments performed  
39 in triplicate ( $*p < 0.05$ ).  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53

54 **Fig. 3.** Responsiveness of the *DHCR24* promoter to the expression of HCV. (A) A *DHCR24* promoter  
55 series with sequential 100-bp deletions was constructed and analyzed as described in the legend to Fig.  
56  
57  
58  
59  
60

1  
2  
3 2B ( $*p < 0.05$ ). (B) An additional deletion series (-167/+113, -144/+113, and -120/+113) was  
4  
5 constructed and analyzed as described in (A).  
6  
7  
8

9 **Fig. 4.** The HCV response element mediates the overexpression of *DHCR24* induced by HCV. (A) The  
10 HCV response element (-167/-140) in the 5'-flanking region of *DHCR24* includes sequences with  
11 similarity to the consensus-binding motifs for AP-2 $\alpha$ , Sp1, and MZF-1. (B) Nuclear extracts were  
12 prepared from RzM6-0d and RzM6-LC cells and subjected to an electrophoresis mobility shift assay  
13 (EMSA; 10  $\mu$ g/sample) using the DIG-labeled HCV response element (28-bp), Sp1 (22-bp), or MZF-1  
14 (21-bp) probes. Cold probe indicates unlabeled oligonucleotides. The arrowhead indicates the interaction  
15 between the binding factor(s) and each oligonucleotide. (C) Cured HuH-7/K4 cells and JFH/K4 cells  
16 were co-transfected with each *DHCR24* promoter reporter plasmid (0.5  $\mu$ g/well) and phRL-TK (0.05  
17  $\mu$ g/well) and analyzed as described in Fig. 2B ( $*p < 0.05$ ). (D) Nuclear extracts prepared from cured  
18 HuH-7/K4 cells or JFH/K4 cells were subjected to EMSA (25  $\mu$ g/sample) using the DIG-labeled HCV  
19 response element, Sp1, or MZF-1 probes.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

33 **Fig. 5.** Sp1 binds to the HCV response element. (A) Nuclear extract was prepared from HepG2 cells  
34 transfected with pcDNA6-AP-2 $\alpha$ -myc and subjected to EMSA (10  $\mu$ g/sample) using DIG-labeled HCV  
35 response element or AP-2 $\alpha$  probes (26-bp). For a supershift analysis of myc-tagged AP-2 $\alpha$ , anti-Myc or  
36 control IgG was added to the binding reaction. The closed arrowhead indicates the interaction between  
37 the binding factor(s) and each oligonucleotide, and an additional interaction with antibody is indicated  
38 by an open arrowhead. (B) Nuclear extract from HepG2 cells was pre-incubated at 4°C for 1 h with  
39 different concentrations (2.5, 5, and 10  $\mu$ M) of mithramycin A (MMA) and subjected to EMSA (10  
40  $\mu$ g/sample) using the DIG-labeled HCV response element, Sp1, or MZF-1 probes. (C) Nuclear extracts  
41 were prepared from RzM6 cells transfected with Sp1 siRNA or control siRNA and subjected to EMSA  
42 (10  $\mu$ g/sample) using the DIG-labeled HCV response element, Sp1, or MZF-1 probes. D: Expression of  
43 Sp1, DHCR24, and other proteins was detected in both the nuclear fraction (N), used for the EMSA  
44 shown in Fig. 4C, and in the cytosolic-membrane fraction (C).  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5 **Fig. 6.** Oxidative stress increases the transcription of *DHCR24* through the HCV response element and  
6 Sp1. (A) HepG2 cells ( $1 \times 10^4$  cells/well in a 96-well plate) were co-transfected with individual *DHCR24*  
7 promoter reporter plasmids (0.5  $\mu\text{g}/\text{well}$ ) and phRL-TK (0.05  $\mu\text{g}/\text{well}$ ). Forty-four hours  
8 post-transfection, cells were treated with or without 1 mM  $\text{H}_2\text{O}_2$  for 4 h and analyzed as described in Fig.  
9 2B ( $*p < 0.05$ ). (B) Nuclear extracts prepared from  $\text{H}_2\text{O}_2$ -treated (1 mM, 4 h) or untreated HepG2 cells  
10 were subjected to EMSA (10  $\mu\text{g}/\text{sample}$ ) using the DIG-labeled HCV response element, Sp1, or MZF-1  
11 probes. Densitometric analysis of shifted bands was performed using the Image Quant software. Data are  
12 shown as the mean + SD from triplicate quantifications of two representative experiments ( $* < 0.05$ ).  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22

23 **Fig. 7.** Overexpression of *DHCR24* in the cells expressing HCV is blocked by treatment with an  
24 oxidative stress scavenger. A: HepG2 cells were treated without (lanes 1, 4, and 6) or with (lanes 2 and  
25 5) NAC (10 mM, 8 h). Cells treated with  $\text{H}_2\text{O}_2$  (1 mM, 4 h) were also treated with 10 mM NAC for 8 h  
26 either before (pre; lanes 3 and 4) or after (post; lanes 5 and 6)  $\text{H}_2\text{O}_2$  exposure. Whole-cell lysates (40  
27  $\mu\text{g}/\text{lane}$ ) were analyzed by 10% SDS-PAGE and immunoblotting using a *DHCR24*/Seladin-1 mAb.  
28 Immunoblotting with an actin mAb served as the internal loading control. The ratio of *DHCR24*/actin  
29 was normalized to that of untreated cells (lane 1). B: RzM6-LC cells were treated with NAC (10 mM)  
30 for 12 h (lane 2) or 24 h (lane 4). Whole-cell lysates were analyzed as described in (A). The ratio of  
31 HCV core to actin protein was also calculated. Experiments were performed 3 times, and representative  
32 results are shown.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45

46 **Fig. 8.** Overexpression and elevated phosphorylation of Sp1 in the cells expressing HCV. A: RzM6-0d  
47 and RzM6-LC cells were treated with or without  $\text{H}_2\text{O}_2$  (1 mM, 4 h). Whole-cell lysates (15  $\mu\text{g}/\text{lane}$ ) were  
48 analyzed by 15% SDS-PAGE and immunoblotting using phospho-H2AX (Ser139) ( $\gamma\text{H2AX}$ ) and HCV  
49 core mAbs. An actin mAb served as an internal loading control. Whole-cell lysates (25  $\mu\text{g}/\text{lane}$ ) were  
50 analyzed by 5% SDS-PAGE and immunoblotting using anti-Sp1 (phosphorylated Sp1 and native Sp1, as  
51 indicated) and anti-phospho-Sp1 (Ser101) was performed. B: RzM6-0d and RzM6-LC cells were  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 fractionated to produce nuclear (N) and cytosolic-membrane fractions (C). Fractionated samples (15  
4  $\mu\text{g}/\text{lane}$ ) were analyzed as described in (A). The ratio of phosphorylated Sp1 to Sp1 protein is indicated.  
5  
6 Immunoblotting using anti-HAUSP served as a high-molecular-weight loading control. C: RzM6-0d  
7  
8 cells were pre-treated for 8 h with NAC (10 mM), MMA (10  $\mu\text{M}$ ), KU55933 (KU; 10  $\mu\text{M}$ ), LY294002  
9  
10 (LY; 50  $\mu\text{M}$ ), or PD98059 (PD; 50  $\mu\text{M}$ ) and incubated for 4 h in the absence or presence of  $\text{H}_2\text{O}_2$  (1 mM).  
11  
12 Whole-cell lysates (40  $\mu\text{g}/\text{lane}$ ) were analyzed by 5% SDS-PAGE and immunoblotting using  
13  
14 anti-phospho-Sp1 (Ser101), (Thr453), and polyclonal anti-Sp1 (white arrowhead, phosphorylated Sp1;  
15  
16 black arrowhead, native Sp1). Detection of HAUSP was performed to confirm the quantity of loaded  
17  
18 protein in each lane. Whole-cell lysates (25  $\mu\text{g}/\text{lane}$ ) were analyzed simultaneously by 10% SDS-PAGE  
19  
20 and immunoblotting using anti-DHCR24/seladin-1 mAb, anti-phospho-Akt (Ser473), and  
21  
22 anti-phospho-ERK antibodies.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Figure 1. Saito et al.

EPS  
250x320mm (300 x 300 DPI)




Figure 2. Saito et al.

175x218mm (300 x 300 DPI)




Figure 3. Saito et al.

192x274mm (300 x 300 DPI)






Figure 4.  
Saito et al.

193x230mm (300 x 300 DPI)



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60





Figure 6.  
Saito et al.

TIFF file  
169x248mm (300 x 300 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



**Figure 7.**  
*Saito et al.*

EPS file  
223x358mm (300 x 300 DPI)