

- [5] Bowser DN, Khakh BS. ATP excites interneurons and astrocytes to increase synaptic inhibition in neuronal networks. *J Neurosci* 2004;24:8606–20.
- [6] Bradbury EJ, Burnstock G, McMahon SB. The expression of P2X<sub>3</sub> purinoreceptors in sensory neurons: effects of axotomy and glial-derived neurotrophic factor. *Mol Cell Neurosci* 1998;12:256–68.
- [7] Brusco A, Gomez LA, López EM, Tagliaferro P, Saavedra JP. Relationship between glial organization and the establishment of nerve tracts in rat spinal cord. *Int J Neurosci* 1995;82:25–31.
- [8] Burnstock G. Physiology and pathophysiology of purinergic neurotransmission. *Physiol Rev* 2007;87:659–797.
- [9] Burnstock G. Purines and sensory nerves. *Handb Exp Pharmacol* 2009;194:333–92.
- [10] Chen M, Gu JG. A P2X receptor-mediated nociceptive afferent pathway to lamina I of the spinal cord. *Mol Pain* 2005;1:4.
- [11] Chizh BA, Illes P. P2X receptors and nociception. *Pharmacol Rev* 2001;53:553–68.
- [12] Chu YX, Zhang Y, Zhang YQ, Zhao ZQ. Involvement of microglial P2X<sub>7</sub> receptors and downstream signaling pathways in long-term potentiation of spinal nociceptive responses. *Brain Behav Immun* 2010;24:1176–89.
- [13] Clark AK, Staniland AA, Marchand F, Kaan TK, McMahon SB, Malcangio M. P2X<sub>7</sub>-dependent release of interleukin-1 $\beta$  and nociception in the spinal cord following lipopolysaccharide. *J Neurosci* 2010;30:573–82.
- [14] Collo G, North RA, Kawashima E, Merlo-Pich E, Neidhart S, Surprenant A, Buell G. Cloning of P2X<sub>5</sub> and P2X<sub>6</sub> receptors and the distribution and properties of an extended family of ATP-gated ion channels. *J Neurosci* 1996;16:2495–507.
- [15] D'Mello R, Dickenson AH. Spinal cord mechanisms of pain. *Br J Anaesth* 2008;101:8–16.
- [16] Fam SR, Gallagher CJ, Salter MW. P2Y<sub>1</sub> purinoreceptor-mediated Ca<sup>2+</sup> signaling and Ca<sup>2+</sup> wave propagation in dorsal spinal cord astrocytes. *J Neurosci* 2000;20:2800–8.
- [17] Fellin T, Sul JY, D'Ascenzo M, Takano H, Pascual O, Haydon PG. Bidirectional astrocyte-neuron communication: the many roles of glutamate and ATP. *Novartis Found Symp* 2006;276:208–17.
- [18] Fields RD. Oligodendrocytes changing the rules: action potentials in glia and oligodendrocytes controlling action potentials. *Neuroscientist* 2008;14:540–3.
- [19] Fields RD, Burnstock G. Purinergic signalling in neuron-glia interactions. *Nat Rev Neurosci* 2006;7:423–36.
- [20] Fonnun F, Johnsen A, Hassel B. Use of fluorocitrate and fluoroacetate in the study of brain metabolism. *Glia* 1997;21:106–13.
- [21] Fujiki Y, Yokota S, Okada Y, Oku Y, Miwakeichi F, Tamura Y, Ishiguro M. A standardization method of voltage-imaging data that enables to average and compare spatio-temporal neuronal activities obtained from different samples. *Neurosci Res* 2010;68:e66.
- [22] Fukuda K, Okada Y, Yoshida H, Aoyama R, Nakamura M, Chiba K, Toyama Y. Ischemia-induced disturbance of neural network function in the rat spinal cord analyzed by voltage-imaging. *Neuroscience* 2006;140:1453–65.
- [23] Fyffe RE, Perl ER. Is ATP a central synaptic mediator for certain primary afferent fibers from mammalian skin? *Proc Natl Acad Sci USA* 1984;81:6890–3.
- [24] Gong QJ, Li YY, Xin WJ, Zang Y, Ren WJ, Wei XH, Li YY, Zhang T, Liu XG. ATP induces long-term potentiation of C-fiber-evoked field potentials in spinal dorsal horn: the roles of P2X<sub>4</sub> receptors and p38 MAPK in microglia. *Glia* 2009;57:583–91.
- [25] Gu JG, Bardoni R, Magherini PC, MacDermott AB. Effects of the P2-purinoreceptor antagonists suramin and pyridoxal-phosphate-6-azophenyl-2',4'-disulfonic acid on glutamatergic synaptic transmission in rat dorsal horn neurons of the spinal cord. *Neurosci Lett* 1998;253:167–70.
- [26] Gu JG, MacDermott AB. Activation of ATP P2X receptors elicits glutamate release from sensory neuron synapses. *Nature* 1997;389:749–53.
- [27] Guthrie PB, Knappenberger J, Segal M, Bennett MVL, Charles AC, Kater SB. ATP released from astrocytes mediates glial calcium waves. *J Neurosci* 1999;19:520–8.
- [28] Hamilton N, Vayro S, Wigley R, Butt AM. Axons and astrocytes release ATP and glutamate to evoke calcium signals in NG2-glia. *Glia* 2010;58:66–79.
- [29] Hugel S, Schlichter R. Presynaptic P2X receptors facilitate inhibitory GABAergic transmission between cultured rat spinal cord dorsal horn neurons. *J Neurosci* 2000;20:2121–30.
- [30] Hülsman S, Straub H, Richter DW, Speckmann EJ. Blockade of astrocytes causes stimulation-induced depolarization as revealed by voltage sensitive dyes in mouse brainstem slices. *Exp Brain Res* 2003;150:117–21.
- [31] Ikeda H, Tsuda M, Inoue K, Murase K. Long-term potentiation of neuronal excitation by neuron-glia interactions in the rat spinal dorsal horn. *Eur J Neurosci* 2007;25:1297–306.
- [32] Ikeda R, Takahashi Y, Inoue K, Kato F. NMDA receptor-independent synaptic plasticity in the central amygdala in the rat model of neuropathic pain. *Pain* 2007;127:161–72.
- [33] Inoue K, Kato F, Tsuda M. The modulation of synaptic transmission by the glial purinergic system. *Open Neurosci J* 2010;4:84–92.
- [34] Jahr CE, Jessell TM. ATP excites a subpopulation of rat dorsal horn neurones. *Nature* 1983;304:730–3.
- [35] Jo YH, Schlichter R. Synaptic corelease of ATP and GABA in cultured spinal neurons. *Nat Neurosci* 1999;2:241–5.
- [36] Kanjhan R, Housley GD, Burton LD, Christie DL, Kippenberger A, Thorne PR, Luo L, Ryan AF. Distribution of the P2X<sub>2</sub> receptor subunit of the ATP-gated ion channels in the rat central nervous system. *J Comp Neurol* 1999;407:11–32.
- [37] Kato F, Shigetomi E. Distinct modulation of evoked and spontaneous EPSCs by purinoreceptors in the nucleus tractus solitarius of the rat. *J Physiol* 2001;530:469–86.
- [38] Kawai A, Okada Y, Mckenhoff K, Scheid P. Theophylline and hypoxic ventilatory response in the rat isolated brainstem-spinal cord. *Respir Physiol* 1995;100:25–32.
- [39] Kawamura Y, Manita S, Nakamura T, Inoue M, Kudo Y, Miyakawa H. Glutamate release increases during mossy-CA3 LTP but not during Schaffer-CA1 LTP. *Eur J Neurosci* 2004;19:1591–600.
- [40] Kettenmann H, Backus KH, Schachner M. Aspartate, glutamate and gamma-aminobutyric acid depolarize cultured astrocytes. *Neurosci Lett* 1984;52:25–9.
- [41] Kobayashi K, Fukuoaka T, Yamanaka H, Dai Y, Obata K, Tokunaga A, Noguchi K. Neurons and glial cells differentially express P2Y receptor mRNAs in the rat dorsal root ganglion and spinal cord. *J Comp Neurol* 2006;498:443–54.
- [42] Kojima S, Nakamura T, Nidaira T, Nakamura K, Ooashi N, Ito E, Watase K, Tanaka K, Wada K, Kudo Y, Miyakawa H. Optical detection of synaptically induced glutamate transport in hippocampal slices. *J Neurosci* 1999;19:2580–8.
- [43] Lambrecht G. Agonists and antagonists acting at P2X receptors: selectivity profiles and functional implications. *Naunyn Schmiedeberg Arch Pharmacol* 2000;362:340–50.
- [44] Lê KT, Villeneuve P, Ramjaun AR, McPherson PS, Beaudet A, Séguéla P. Sensory presynaptic and widespread somatodendritic immunolocalization of central ionotropic P2X ATP receptors. *Neuroscience* 1998;83:177–90.
- [45] Li P, Calejesan AA, Zhuo M. ATP P2x receptors and sensory synaptic transmission between primary afferent fibers and spinal dorsal horn neurons in rats. *J Neurophysiol* 1998;80:3356–60.
- [46] Li J, Perl ER. ATP modulation of synaptic transmission in the spinal substantia gelatinosa. *J Neurosci* 1995;15:3357–65.
- [47] Liu YP, Yang CS, Chen MC, Sun SH, Tzeng SF. Ca<sup>2+</sup>-dependent reduction of glutamate aspartate transporter GLAST expression in astrocytes by P2X<sub>7</sub> receptor-mediated phosphoinositide 3-kinase signaling. *J Neurochem* 2010;113:213–27.
- [48] Lozier AP, Kendig JJ. Long-term potentiation in an isolated peripheral nerve-spinal cord preparation. *J Neurophysiol* 1995;74:1001–9.
- [49] Magoski NS, Walz W. Ionic dependence of a P2-purinoreceptor mediated depolarization of cultured astrocytes. *J Neurosci Res* 1992;32:530–8.
- [50] Molander C, Xu Q, Rivero-Melian C, Grant G. Cytoarchitectonic organization of the spinal cord in the rat: II. The cervical and upper thoracic cord. *J Comp Neurol* 1989;289:375–85.
- [51] Nakagawa T, Wakamatsu K, Zhang N, Maeda S, Minami M, Satoh M, Kaneko S. Intrathecal administration of ATP produces long-lasting allodynia in rats: differential mechanisms in the phase of the induction and maintenance. *Neuroscience* 2007;147:445–55.
- [52] Nakamura I, Ohta Y, Kemmotsu O. Characterization of adenosine receptors mediating spinal sensory transmission related to nociceptive information in the rat. *Anesthesiology* 1997;87:577–84.
- [53] Nakatsuka T, Gu JG. ATP P2X receptor-mediated enhancement of glutamate release and evoked EPSCs in dorsal horn neurons of the rat spinal cord. *J Neurosci* 2001;21:6522–31.
- [54] Nakatsuka T, Gu JG. P2X purinoreceptors and sensory transmission. *Pflügers Arch* 2006;452:598–607.
- [55] Nörenberg W, Illes P. Neuronal P2X receptors: localisation and functional properties. *Naunyn Schmiedeberg Arch Pharmacol* 2000;362:324–39.
- [56] Okada Y, Masumiya H, Tamura Y, Oku Y. Respiratory and metabolic acidosis differentially affect the respiratory neuronal network in the ventral medulla of neonatal rats. *Eur J Neurosci* 2007;26:2834–43.
- [57] Oku Y, Masumiya H, Okada Y. Postnatal developmental changes in activation profiles of the respiratory neuronal network in the rat ventral medulla. *J Physiol* 2007;585:175–86.
- [58] Paxinos G, Törk I, Tecott LH, Valentino KL. Atlas of the developing rat brain. San Diego: Academic Press; 1991.
- [59] Rhee JS, Wang ZM, Nabekura J, Inoue K, Akaike N. ATP facilitates spontaneous glycinergic IPSC frequency at dissociated rat dorsal horn interneuron synapses. *J Physiol* 2000;524:471–83.
- [60] Salter MW, De Koninck Y, Henry JL. Physiological roles for adenosine and ATP in synaptic transmission in the spinal dorsal horn. *Prog Neurobiol* 1993;41:125–56.
- [61] Salter MW, Hicks JL. ATP-evoked increases in intracellular calcium in neurons and glia from the dorsal spinal cord. *J Neurosci* 1994;14:1563–75.
- [62] Shigetomi E, Kato F. Action potential-independent release of glutamate by Ca<sup>2+</sup> entry through presynaptic P2X receptors elicits postsynaptic firing in the brainstem autonomic network. *J Neurosci* 2004;24:3125–35.
- [63] Shiokawa H, Nakatsuka T, Furue H, Tsuda M, Katafuchi T, Inoue K, Yoshimura M. Direct excitation of deep dorsal horn neurones in the rat spinal cord by the activation of postsynaptic P2X receptors. *J Physiol* 2006;573:753–63.
- [64] Sokolova E, Nistri A, Giniatullin R. Negative cross talk between anionic GABA<sub>A</sub> and cationic P2X ionotropic receptors of rat dorsal root ganglion neurons. *J Neurosci* 2001;21:4958–68.
- [65] Somjen GG. Evoked sustained focal potentials and membrane potential of neurons and of unresponsive cells of the spinal cord. *J Neurophysiol* 1970;33:562–82.
- [66] Suidicani SO, Brosnan CF, Scemes E. P2X<sub>7</sub> receptors mediate ATP release and amplification of astrocytic intercellular Ca<sup>2+</sup> signaling. *J Neurosci* 2006;26:1378–85.

- [67] Tsuda M, Koizumi S, Kita A, Shigemoto Y, Ueno S, Inoue K. Mechanical allodynia caused by intraplantar injection of P2X receptor agonist in rats: involvement of heteromeric P2X<sub>2/3</sub> receptor signaling in capsaicin-insensitive primary afferent neurons. *J Neurosci* 2000;20:RC90.
- [68] Tsuda M, Shigemoto-Mogami Y, Koizumi S, Mizokoshi A, Kohsaka S, Salter MW, Inoue K. P2X<sub>4</sub> receptors induced in spinal microglia gate tactile allodynia after nerve injury. *Nature* 2003;424:778–83.
- [69] Verkhatsky A, Krishtal OA, Burnstock G. Purinoceptors on neuroglia. *Mol Neurobiol* 2009;39:190–208.
- [70] Vulchanova L, Riedl MS, Shuster SJ, Buell G, Surprenant A, North RA, Elde R. Immunohistochemical study of the P2X<sub>2</sub> and P2X<sub>3</sub> receptor subunits in rat and monkey sensory neurons and their central terminals. *Neuropharmacology* 1997;36:1229–42.
- [71] Vulchanova L, Riedl MS, Shuster SJ, Stone LS, Hargreaves KM, Buell G, Surprenant A, North RA, Elde R. P2X<sub>3</sub> is expressed by DRG neurons that terminate in inner lamina II. *Eur J Neurosci* 1998;10:3470–8.
- [72] Xiang Z, Bo X, Burnstock G. Localization of ATP-gated P2X receptor immunoreactivity in rat sensory and sympathetic ganglia. *Neurosci Lett* 1998;256:105–8.
- [73] Yi DK, Barr GA. Formalin-induced c-fos expression in the spinal cord of fetal rats. *Pain* 1997;73:347–54.
- [74] Yokota S, Tsumori T, Oka T, Nakamura S, Yasui Y. GABAergic neurons in the ventrolateral subnucleus of the nucleus tractus solitarius are in contact with Kölliker-Fuse nucleus neurons projecting to the rostral ventral respiratory group and phrenic nucleus in the rat. *Brain Res* 2008;1228:113–26.
- [75] Yoshida H, Okada Y, Maruiwa H, Fukuda K, Nakamura M, Chiba K, Toyama Y. Synaptic blockade plays a major role in the neural disturbance of experimental spinal cord compression. *J Neurotrauma* 2003;20:1365–76.
- [76] Zeng JW, Liu XH, Zhao YD, Xiao Z, He WJ, Hu ZA, Ruan HZ. Role of P2Y<sub>1</sub> receptor in astroglia-to-neuron signaling at dorsal spinal cord. *J Neurosci Res* 2009;87:2667–76.

