

minimum number of probes of 3 in the region and 0.5 minimum absolute average log ratio for the region. After the first screening described above, the raw data of selected loci were individually scrutinized. As a result, some "aberrations" were removed from the aberration list because they were located within a noisy area, such as the telomeres, and the reliability of the aberration calls in such region was low. Although Genomic Workbench Standard Edition version 5.0.14 adopts hg18 for gene mapping, we referred to Build36.3 in the Map Viewer constructed by NCBI (<http://www.ncbi.nlm.nih.gov/projects/mapview/>) for detailed mapping of genes.

### **V(D)J recombination analysis**

Recombination in the V(D)J region of the immunoglobulin (Ig) heavy chain gene in B-LCLs was analyzed by PCR as described previously with slight modification (Kiyoi et al., 1992; Abe et al., 1994). The primers used for the PCR were 5'-GAG TCG AC(A/T) C(A/G)G C(G/CXG/A)T GTA (T/C)T(T/A) CTG-3' for the V common region and 5'-CCA AGC TTA CCT GAG GAG ACG GTG A-3' for the J common region. PCR was performed using an ExTaq polymerase kit (TaKaRa Bio, Otsu, Shiga, Japan) with a reaction mixture containing 0.4  $\mu$ M of the J common primer and 4  $\mu$ M of the V common primer. An initial incubation at 94°C for 5 min was followed by 40 cycles of

denaturation at 94°C for 10 sec, annealing at 60°C for 1 sec, and extension at 72°C for 15 sec, and finally incubation at 72°C for 5 min. The PCR products were separated on 3% NuSieve GTG agarose gels (Lonza, Basel, Switzerland) and visualized by ethidium bromide staining. Expected product sizes from recombined V(D)J region were in the range 50-200 bp.

## RESULTS

### **The Sonoda-Tajima Cell Collection**

The Sonoda-Tajima cell collection contains more than 3,500 PBMNC samples obtained from ethnic populations across the world (Figure 1). The collection is particularly rich in samples from South America. As stated earlier, any direct use of the PBMNC samples would soon lead to their exhaustion. In comparison to the *in vitro* expansion of the whole genome using PCR, establishing cell lines is a much better method for maintaining the whole genome in a stable manner. With regard to establishing cell lines from PBMNCs, the generation of B cell lines using EBV (B-LCLs) is the most common approach. The genomic DNA of B-LCLs is maintained in a stable fashion. These facts encouraged us to establish B-LCLs using the PBMNCs of the Sonoda-Tajima Cell Collection. In order to include as many ethnic groups as possible from South America

and a sufficient number of individuals in each group, we selected the samples indicated in Table 1 and Figure 2.

When we used either fresh or cryopreserved PBMNCs in good condition to establish the B-LCLs, we were successful in all cases. However, the PBMNC samples in the Sonoda-Tajima Cell Collection have been cryopreserved for over 20 years. In addition, in sampling areas where no electricity was available, the samples were kept at relatively high temperatures prior to cryopreservation. As a consequence of these factors the viability of some of the thawed PBMNC samples was low, i.e., the cells were in poor condition after thawing, and thus it was not easy to establish B-LCLs for these samples. Overall, a success rate of approximately 80% was achieved for the establishment of B-LCLs from the PBMNC samples of the Sonoda-Tajima Cell Collection. Since the success rate was less than 100%, we decided not to use samples for which there were only one or two tubes cryopreserved. Future treatment of such samples remains to be considered. At the moment, we have established 512 B-LCLs from the Sonoda-Tajima Cell Collection. Information on the established B-LCLs is given in Table 1 and Figure 2.

### **Karyotype analysis**

In contrast to other cell lines, such as immortalized cancer cell lines, one of the most significant features of B-LCLs is their chromosome stability. B-LCL cells retain a normal chromosome karyotype even after relatively long-term culture (reviewed by Nilsson, 1992; Lalle et al., 1995; Okubo et al., 2001); thus, they have been widely used for genetic analysis in many research fields. To confirm that the B-LCLs established from the Sonoda-Tajima Cell Collection maintained a normal karyotype, three B-LCL cell lines, WY084, YAN3191 and YAN3268, were karyotyped following G-banding of chromosome preparations. Chromosome numbers were counted in 50 cells (mode-analysis), and then a detailed G-band analysis was performed in 20 of these cells to identify chromosome aberrations. Consistent with previous reports, the vast majority of the cells had a normal karyotype (Figure 3). In the YAN3268 cell line, all of the cells analyzed had a normal chromosome number as well as a normal karyotype (Figure 3A). In the WY084 cell line, 2 of the 50 cells had a reduced chromosome number (Figure 3B). This might have been an artifact of preparation as no structural aberrations were observed in the cells. In the YAN3191 cell line, all 20 karyotyped cells had an elongated pericentromeric region on the long arm of chromosome 1 (Figure 3C). The centric heterochromatin of chromosome 1 is known to be variable and show heteromorphism between individuals ([http://www.rerf.or.jp/dept/genetics/giemsas\\_5\\_e.html](http://www.rerf.or.jp/dept/genetics/giemsas_5_e.html)).

In general, if an irregular karyotype is detected, it is not possible to conclude that it arose during culture because there is a possibility it was already present in the cells of the individual from whom the initial sample was obtained, as in the YAN3191 cell line. In addition, since karyotype analysis has a limited resolution (3-5 Mbp) we cannot detect smaller rearrangements. These limitations prompted us to perform the array CGH analysis on the B-LCLs and their parental PBMNCs.

#### **Array CGH analysis**

To avoid the influence of genetic background, such as copy number variation (CNV), we decided to compare the B-LCL and parental PBMNC from the same person in our experiments. We analyzed ten B-LCLs derived from the Sonoda-Tajima Cell Collection and one B-LCL derived from a healthy Japanese volunteer.

The probe set used in the microarray was uninformative regarding the heterochromatic region of chromosome 1. However, in the YAN3191 cell line we did not detect any variants with respect to the other chromosome 1 probes suggesting that there was no major rearrangement of the chromosome (or of the other chromosomes) and that the cell line essentially retained the innate genomic structure (Figure 4).

A few aberrations were observed in all eleven samples: a deletion in

chromosome 14 (in the variable, diversity or joining regions of the Ig heavy chain); a deletion in chromosome 2 (in the variable or joining regions of the Ig  $\kappa$  light chain); an amplification in chromosome 14 (in the variable or joining regions of the T cell receptor (TCR)  $\alpha$  chain); and, an amplification in chromosome 7 (in the variable, joining or constant regions of the TCR  $\gamma$  chain). Typical examples of chromosome 14 are shown in Figure 5. In addition, a few other aberrations were detected in some but not all cell lines: a deletion at the immunoglobulin  $\lambda$  locus on chromosome 22; and, amplification at the TCR  $\beta$  on chromosome 7. A precise description of each aberration locus and aberration type described above is given in Table 2 and Figure 6. In addition, a few other specific aberrations were observed in some but not all B-LCLs (Figure S1-S4).

In the B-LCLs, it is possible that the deletions detected in the regions associated with the Ig genes were already present in the B lymphocytes prior to EBV infection. Similarly, since the population of B lineage cells in PBMNCs is very small, it is possible that the detected aberrations were already present in the parent B lymphocytes but not in the non-B cells. To examine this latter possibility, we attempted to compare the genomes of B lineage and non-B lineage cells. However, it was impossible to obtain sufficient B lineage cells from the Sonoda-Tajima Collection. Thus, we used a blood sample from one healthy Japanese adult volunteer and two umbilical

cord blood samples obtained from two independent Japanese neonates. CD19<sup>+</sup> and CD19<sup>-</sup> cell populations were collected using the magnetic beads system described earlier (see Materials and methods). After selection, the proportion of CD19<sup>+</sup> cells was approximately 90% (Figure 7B, C).

As expected, most aberrations observed in B-LCL cell lines were also observed in all CD19<sup>+</sup> cells: deletions at the Ig heavy chain on chromosome 14 and at the Ig  $\kappa$  and  $\lambda$  light chains on chromosomes 2 and 22; amplifications at the TCR  $\alpha$  locus on chromosome 14, and at the  $\beta$  and  $\gamma$  loci on chromosome 7. A precise description of each aberration locus and aberration type is also described in Table 2 and Figure 6.

Since a flow cytometry analysis indicated that the major cell population in the PBMNCs was CD3<sup>+</sup> T-lineage cells (Figure 7A), the results of the statistical analysis of array CGH that suggested "amplification" at the T cell receptor loci were highly likely due to deletions at these loci in T-lineage cells rather than amplification in B-lineage cells.

### **V(D)J recombination analysis**

The array CGH analysis clearly demonstrated that B-LCLs possessed rearranged Ig genes. To confirm the rearrangement of the Ig heavy chain, we performed a PCR

analysis on genomic DNAs from the B-LCLs. The variable region of the Ig heavy chain is encoded by 3 gene clusters: approximately 40 functional variable (V) genes, 6 functional joining (J) genes and 25 functional diversity (D) genes. At the later stages of B lymphocyte differentiation, this region undergoes genomic rearrangements such that only one randomly selected gene from each gene cluster is connected to form the mature V(D)J variable gene, and the remaining "left over" genes are removed from the genome (summarized in OMIM #147070, <http://www.ncbi.nlm.nih.gov/omim/>). The PCR analysis identified V(D)J recombination in all B-LCLs (Figure 8). Both the results of the array CGH analysis and the PCR analysis were consistent with a previous report that many B-LCLs secrete Ig into the culture medium (Rosen et al., 1977). Since six of the lines showed a single PCR product, these lines were each presumed to consist of a single B cell clone. However, the other cell lines showed two or more PCR products indicating that they consisted of two or more B cell clones.

## DISCUSSION

Long-term cell culture lines, such as immortalized cancer cell lines, generally possess abnormal karyotypes. However, some cell lines tend to retain a normal karyotype. It is well known that B-LCLs and pluripotent stem cell lines, such as embryonic stem (ES)



cells and induced pluripotent stem (iPS) cells, show this behavior. Due to this characteristic, B-LCLs have been used in many projects, such as the International Histocompatibility Working Group ([http://www.ihwg.org/cellbank/cell\\_lines/blcl.html](http://www.ihwg.org/cellbank/cell_lines/blcl.html)), International HapMap Project (<http://hapmap.ncbi.nlm.nih.gov/>) and Human Genome Diversity Project (<http://www.cephb.fr/en/hgdp/table.php>). High throughput and comprehensive genome analyses have recently been reported using B-LCL collections, e.g., genotyping of blood cell alloantigens (Kroll et al., 2001), detection of copy number variation (CNV) (Redon et al., 2006), and population genetics (Li et al., 2008).

It may be inevitable, however, that long-term cell cultures will accumulate chromosomal aberrations and/or genetic mutations. Additionally, it should be noted that there is still some controversy over whether B-LCLs are truly immortalized cells. Sugimoto et al. (2004) reported that B-LCLs could no longer proliferate following long-term culture, that is, they showed the so-called “crisis” observed in human primary cell cultures. By chance, some B-LCLs can overcome this crisis phase and continue to grow indefinitely in culture, i.e., show immortalization; however, such cells also showed malignant transformation with abnormal karyotypes (Sugimoto et al., 2004). These findings have prompted a vigorous debate on whether the genomes of B-LCLs are really stable.

To address this question, genome-wide evaluations of the genomic stability of B-LCLs have recently been initiated. Whole genome single nucleotide polymorphism (SNP) genotyping indicated that the discrepancies between B-LCLs and their parental PBMNCs were not statistically significant (Simon-Sanchez et al., 2006; Herbeck et al., 2009). In contrast, however, another study has reported different copy numbers of certain genes in B-LCLs and their parental PBMNCs (The Wellcome Trust Case Control Consortium, 2010). In relation to this issue, Bruder et al. analyzed CNVs in monozygotic twins and found somatic mosaicism for CNVs, i.e., CNVs were detected not only between individuals but also within an individual, with a rate of 70-80% of blood cells affected in the most severe cases and of 10-15% in the less affected cases (Bruder et al., 2008). CNVs form continuously in mitosis and meiosis by a diversity of mechanisms, and it is thought that CNVs are one of major driving forces in the rapid evolution of human beings (reviewed by Hastings et al., 2009).

In our array CGH analysis, several aberrations were detected in B-LCLs when they were compared with PBMNCs. However, most of these aberrations were also detected in B lineage cells when compared with non-B lineage cells. Therefore, it is highly likely that most of the aberrations detected in the B-LCLs were present in their parental B cells rather than having occurred de novo during establishment of the cell

cultures. However, with respect to some of the aberrations detected in some B-LCLs derived from the Sonoda-Tajima collection (Figure S1-S4), it was impossible to determine whether they occurred de novo during culture or were present in the parental B cells since insufficient numbers of parental B cells were available for analysis. Of note, a deletion in chromosome 22 (in the variable and joining regions of the Ig  $\lambda$  light chain) was detected in several B-LCLs but not in all lines, suggesting that EBV had infected the B cells at different stages of differentiation, i.e., before and after the rearrangement of the Ig  $\lambda$  light chain.

Interestingly, amplification at 5q35.3 was only detected in one of the two Japanese B-LCLs derived from the same individual (Figure 6). Of course, we cannot formally deny the possibility of de novo amplification during cell culture. However, this observation might indicate somatic mosaicism as described above (Bruder et al., 2008). Even if this is the case, it is not clear at present whether such somatic rearrangements are coupled with RAG transposition (Reddy et al., 2010) or some other mechanism (reviewed by Hastings et al., 2009). Moreover, with respect to aberrations of genes other than of the Ig chains in B-LCLs and primary B cells, the biological implications remain to be determined. When B-LCLs are utilized in genetic research, such aberrations need to be taken into account.

According to our array CGH results, the variation in copy numbers of some genes in B-LCLs compared to their parental PBMNCs might have been present in the parental B cells, and might have been detected if only B lineage cells had been collected and analyzed in a similar fashion to the present study. The possible consequences of any CNV between B cells and non-B cells have yet to be investigated. At present, we cannot formally exclude the possibility that such CNV occurred during the establishment and culture of the B-LCLs.

Overall, our analyses support the contention that B-LCLs provide a valuable source of genomic DNAs for a wide range of genetic studies. However, it is recommended that short-term cell cultures are utilized. Certainly, so-called immortalized B-LCLs with abnormal karyotypes (Okubo et al., 2001; Sugimoto et al., 2004) should not be used for genetic research. In addition, the occurrence of aberrations in B-LCLs when compared to whole mononuclear blood cells should be taken into account when the affected genomic loci are targeted for research.

The Sonoda-Tajima Cell Collection is a very valuable cell collection obtained from various ethnic populations across the world, particularly from South America. Since some of these populations in South America no longer exist, it will be impossible to prepare a similar collection in the future. Here, we describe establishment of B-LCLs

from the Sonoda-Tajima Cell Collection; currently, B-LCLs from more than 500 of the total of about 3,500 samples have been developed. All of these B-LCLs are available from the cell bank held at the Cell Engineering Division of RIKEN BioResource Center in Japan (<http://www.brc.riken.jp/lab/cell/english/>). At the moment, approximately 150 B-LCLs are immediately available, while the others are now under preparation for rapid distribution. In addition, we are willing to establish more B-LCLs on demand from scientists around the world. We believe that our development of this resource will contribute to various fields of science such as human genetics, human evolution, the history of human migrations across continents, the pharmacokinetics of ethnic minority groups, among others.

## **FUNDING**

This work was supported by the Ministry of Education, Culture, Sports, Science, and Technology in Japan.

## **ACKNOWLEDGEMENTS**

We thank Dr. I. Ishiwata at Ishiwata Hospital for providing the umbilical cord blood samples, Dr. Y. Kodama at Radiation Effect Research Foundation for suggestions and

comments on karyotype analysis, Dr. T. Yamamoto at Nagoya University for comments on CNVs, Dr. K. Moriwaki at RIKEN BioResource Center for general discussion and encouragement, H. Sone and J. Niikura for technical assistance, and other members of the Cell Engineering Division of RIKEN BioResource Center for help with various technical matters.

## REFERENCES

- Abe T. et al. 1994. Evidence for defects in V(D)J rearrangements in patients with severe combined immunodeficiency. *J Immunol.* 152: 5504-5513
- Bird AG, McLachlan SM, Britton S. 1981. Cyclosporin A promotes spontaneous outgrowth *in vitro* of Epstein-Barr virus-induced B-cell lines. *Nature.* 289: 300-301
- Bruder CEG. et al. 2008. Phenotypically concordant and discordant monozygotic twins display different DNA Copy-Number-Variation profiles. *Am. J. Human Genet.* 82: 763-771
- Fujiyoshi T. et al. 1999. Characteristic distribution of HTLV type I and HTLV type II carriers among native ethnic groups in South America. *AIDS Res Hum Retroviruses.* 15: 1235-1239
- Hastings PJ. et al. 2009. Mechanisms of change in gene copy number. *Nat Rev Genet.* 10:551-564
- Herbeck JT. et al. 2009. Fidelity of SNP array genotyping using Epstein Barr Virus-transformed B-lymphocyte cell lines: Implications for genome-wide association studies. *PLoS ONE.* 4: e6915
- Ijichi S. et al. 1993. Identification of Human T cell Leukemia virus type IIb infection in the Wayu, an aboriginal population of Colombia. *Jpn J Cancer Res.* 84: 1215-1218

- Kiyoi H, Naoe T, Horibe K, Ohno R. 1992 Characterization of the immunoglobulin heavy chain complementarity determining region (CDR)-III sequences from human B cell precursor acute lymphoblastic leukemia cells. *J Clin Invest.* 89: 739-746
- Komurian-Pradel F, Pelloquin F, Sonoda S, Osame M, de The G. 1992. Geographical subtypes demonstrated by RFLP following PCR in the LTR region of HTLV-1. *AIDS Res Hum Retroviruses.* 8: 429-434
- Kroll H, Carl B, Santoso S, Bux J, Bein G. 2001. Workshop report on the genotyping of blood cell alloantigens. *Transfus Med.* 11: 211-219
- Lalle P. et al. 1995. Genomic stability and wild-type p53 function of lymphoblastoid cells with germ-line *p53* mutation. *Oncogene.* 10: 2447-2454
- Li H-C. et al. 1999. The presence of ancient human T-cell lymphotropic virus type I provirus DNA in an Andean mummy. *Nat Med.* 5: 1428-1432
- Li JZ. et al. 2008. Worldwide human relationships inferred from genome-wide patterns of variation. *Science.* 319: 1100-1104
- Masters JR. et al. 2001. Short tandem repeat profiling provides an international reference standard for human cell lines. *Proc Natl Acad Sci USA.* 98: 8012-8017
- Miura T. et al. 1994. Phylogenetic subtypes of human T-lymphotropic virus type I and their relations to the anthropological background. *Proc Natl Acad Sci USA.* 91:



1124-1127

- Miura T. et al. 1997. Molecular phylogeny of human T-cell leukemia virus type I and II of Amerindians in Colombia and Chile. *J Mol Evol.* 44: S76-S82
- Nilsson K. 1979. The nature of lymphoid cell lines and their relationship to the virus. In: Epstein MA, Achong BG, editor. *The Epstein-Barr Virus*. Berlin: Springer-Verlag. p. 225-266,
- Nilsson K. 1992. Human B-lymphoid cell lines. *Hum Cell.* 5: 25-41
- Okubo M. et al. 2001. Clonal chromosomal aberrations accompanied by strong telomerase activity in immortalization of human B-lymphoblastoid cell lines transformed by Epstein-Barr virus. *Cancer Genet Cytogenet.* 129: 30-34
- Osame M, Usuku K, Ijichi N, Amitani H, Igata A. 1986. HTLV-1 associated myelopathy, a new clinical entity. *Lancet* i:1031-1032
- Reddy YVR, Perkins EJ, Ramsden DA. 2010. Genomic instability due to V(D)J recombination-associated transposition. *Gene Dev.* 20: 1575-1582
- Redon R. et al. 2006. Global variation in copy number in the human genome. *Nature.* 444: 444-454
- Rickinson AB. et al. 1984. T-cell-mediated regression of "spontaneous" and of Epstein-Barr virus-induced B-cell transformation *in vitro*: Studies with Cyclosporin

- A. Cell Immunol. 87: 646-658
- Rosen A, Gergely P, Jodal M, Klein G, Britton S. 1977. Polyclonal Ig production after Epstein-Barr virus infection of human lymphocytes *in vitro*. Nature. 267: 52-54
- Simon-Sanchez J. et al. 2006. Genome-wide SNP assay reveals structural genomic variation, extended homozygosity and cell-line induced alterations in normal individuals. Human Mol Genet. 16:1-14
- Sonoda S. Li HC. Tajima K. 2011. Ethnoepidemiology of HTLV-1 related diseases: Ethnic determinants of HTLV-1 susceptibility and its worldwide dispersal. Cancer Sci. 102: 295-301
- Sugimoto M, Tahara H, Ide T, Furuichi Y. 2004. Steps involved in immortalization and tumorigenesis in human B-lymphoblastoid cell lines transformed by Epstein-Barr virus. Cancer Res. 64: 3361-3364
- Switzer WM. et al. 1995. Phylogenetic relationship and geographic distribution of multiple Human T-cell Lymphotropic Virus type II subtypes. J Virol. 69: 621-632
- The Wellcome Trust Case Control Consortium. 2010. Genome-wide association study of CNVs in 16,000 cases of eight common diseases and 3,000 shared controls. Nature. 464: 713-720
- Uchiyama T, Yodoi J, Sagawa K, Takatsuki K, Uchino H. 1977. Adult T-cell leukemia:

clinical and hematologic features of 16 cases. *Blood*. 50: 481-492

Yamashita M, Ido E, Miura T, Hayami M. 1996. Molecular epidemiology of HTLV-1 in the world. *J Acquir Immune Defic Syndr Hum Retrovirol*. 13: S124-131

Yoshino K. et al. 2006. Essential role for gene profiling analysis in the authentication of human cell lines. *Hum Cell*. 19: 43-38

Yunis JJ, Sawyer JR, Ball DW. 1978. The characterization of high-resolution G-banded chromosomes of man. *Chromosoma*. 67: 293-307

## FIGURE LEGENDS

Fig. 1. Hypothetical pathways of prehistoric human migration.

Red arrows indicate hypothetical prehistoric migration routes of humans after leaving Africa approximately 70 thousand years ago. The numbers in boxes indicate the number of individuals who donated blood samples. The yellow circles indicate the approximate geographic location where the blood samples were collected.

Fig. 2. Geographic locations where blood samples were collected

Red circles indicate approximate locations where the blood samples were collected. The numbers in circles correspond to the location numbers in Table 1.

Fig. 3. Karyotype analysis

The karyotypes of the 3 cell lines YAN3268 (A), WY084 (B), and YAN3191 (C) were analyzed. A representative karyotype of the sample of 20 cells karyotyped and 50 mode-analyzed cells. The chromosome notation system follows the guidelines in ISCN 1991.

Fig. 4. Array CGH of chromosome 1 of the YAN3191 cell line