

Fig. 2. Southern blotting analysis of high molecular DNA obtained from highly purified cell fractions (>99.5%) probed with EBV-terminal repeated (TR) sequence. TCR $\alpha\beta$ ⁺ cells and V δ 2⁺ cells in Patient (Pt)-1, and CD3⁺ cells and CD56⁺ cells in Pt-6 showed >99.9% of purity. EcoRI (E) or BamHI (B)-digested DNA was separated by electrophoresis (2–5 μ g/lane) and hybridized with a ³²P-labeled 5.2-kb BamHI–EcoRI fragment containing the TR repeat sequences. The blotting sensitivity was determined to be at least 0.5% contamination of EBV-infected cells. Arrow, a clonal band. *Non-specific band in each digestion. NC: normal control of EBV-seropositive healthy adult lymphocytes, PC: positive control of Raji cell lines.

CD4⁺T-cells were the major target. A TR band in NK-cells of Patient-2 was the same size as one of biconal bands in CD3⁺T-cells. Patient-3 showed a single TR band in CD4⁺T-cells but not CD8⁺T-cells.

Clonal TR bands were detected in CD3⁺T-cells (>99.5%) of all patients with NK-cell LPD. Blotting analysis of EBV-TR using MNC showed biconal bands in Patient-4. TR(s) in CD3⁺T-cells of Patient-4 (biconal) and Patient-5 (monoclonal) were the same size as those in NK-cells of the same patients. On the other hand, CD3⁺T-cells (>99.9%) from Patient-6 had a larger TR band than NK-cells (>99.9%) (Fig. 2), although the blotting using MNC showed the same sized band as NK-cells.

4.3. EBV load in BM CD34⁺ cells

Virus loads were re-quantified using highly purified subsets from PB and BM cells to further assess the origin of EBV-infection. EBV DNA was not detectable in BM-derived lin⁻CD34⁺ cells of any 5 patients studied, nor in seropositive controls (Fig. 3). EBV was not detected in CD34⁺ cells by the double staining of EBER *in situ* hybridization and immunohistochemistry for BM samples (Fig. 4).

5. Discussion

The current study found that EBV⁺T/NK-cells clonally proliferated in various subsets of patients. EBV clonotype, as determined by EBV TR size, in the minor subsets notably coincided with that in the major subsets of infection. On the other hand, BM CD34⁺ cells carried no EBV. These data suggest that EBV can infect lymphocytes having differentiation ability, that then clonally distribute to T/NK-cells, but not involve infection of CD34⁺ hematopoietic stem cells (HSCs).




Fig. 3. Quantification of EBV DNA in highly purified cell fractions (>99.9%) obtained from patients assessed by real-time PCR. High molecular DNA was extracted from V δ 2, $\alpha\beta$ T, CD4, CD3, and CD56-expressing cells fractionated from peripheral blood, and CD34⁺ cells (hatched bar) obtained from bone marrow cells. Solid bars represent the major infected subsets with high EBV load (black bar), and the minor infected subsets with low EBV load (gray bar). EBV-seropositive healthy carriers or patients with immune thrombocytopenic purpura were used as controls.

It remains controversial in regard to whether CAEBV is an infection or clonal disease. EBER⁺T-cells and NK-cells can be found in the tonsils of acute IM patients.¹⁶ EBV-shedding T/NK-cell lines were established from CAEBV patients.^{17,18} Excessive activation of lytic virus may raise the chance of ectopic infection. While cytotoxic CD8⁺T-cells clonally expand in IM patients,¹⁹ non-CD8⁺T-cells were infected and clonally proliferated in our patients. PB $\gamma\delta$ T-cells




Fig. 4. Immunohistochemical staining of bone marrow (BM) cells obtained from Patient-5 (a) and Patient-6 (b). BM clot samples were stained with anti-CD34 antibody using fuchsin as a chromogen for visualization of alkaline phosphatase activity (positive: red cytosol, *open arrow*), and were then employed for *in situ* hybridization of EBV using diaminobenzidine as a chromogen of peroxidase (positive: brown nucleus, *closed arrow*). There were no double positive cells for CD34 and EBV in BM cells. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of the article.)

consist of polyclonal $\text{V}\delta 2/\text{V}\gamma 9^+$ T-cells in healthy persons. This subset was clonally infected with EBV in Patient-1. Toyabe et al.²⁰ reported that a CAEBV patient with biclonal expansion of $\text{V}\beta 7^+$ and $\text{V}\beta 9^+$ cells in PB $\text{CD}4^+$ T-cells. Both clones were infected with EBV strain that was indistinguishable from each other. Endo et al.²¹ analyzed EBV clonotype in 2 CAEBV patients having biphenotypic expansions of $\text{CD}4^+$ and $\text{CD}8^+$ T-cells. Both subsets carried a TR band with the same size. Germinal center EBV^+ B-cells could evolve LPD.²² Diverse T/NK-cell infection in this line might not represent spreading lytic viruses, but differentiation and expansion of T/NK-cell clones with latent episome. An alternative explanation is integration of the EBV gene into the host genome.²³ EBV latent genes are expressed in heterogeneous and restricted patterns in CAEBV patients.²⁴ Specific integration of EBV gene rarely occurs in transformed cells. Taken together, EBV could infect T/NK-cells at differentiating stages, and these cells proliferate and enter the circulation. Multiple EBV strains affect EBV-associated tumors in a single host.²⁵ Site specific clonality should be investigated to understand EBV^+ T/NK-cell LPD.

The major concern is how clonal EBV^+ T/NK-cells are allowed to proliferate in patients. EBV^+ B-cell LPD arises in X-linked LPD patients. A Caucasian patient with B-cell type CAEBV had defective cytotoxicity due to *PRF1* mutations.²⁶ Although cytotoxic defects are implicated in CAEBV patients,²⁷ EBV-infected T/NK-cells may have altered cytotoxicity. Neither a genetic defect nor familial occurrence has been identified in T/NK-cell type CAEBV or EBV-HLH.^{28,29} The end-to-end length of fused TR itself may drive the clonal selection of EBV-infected cells as an oncogenic component during primary infection.³⁰

This study demonstrated that BM $\text{CD}34^+$ cells were not a niche of EBV infection. Human immunodeficiency virus (HIV)-infected $\text{CD}34^+$ HSCs were detected in HIV-positive patients.³¹ HTLV-1 or HHV8 infects $\text{CD}34^+$ HSCs.³² Primitive HSCs are resistant to HIV infection, although these cells become increasingly susceptible once they begin to differentiate into committed HSCs.³³ EBV might not infect BM, but infect peripheral or thymic $\text{CD}34^+\text{CD}4^-\text{CD}8^+$ -T-cell precursors (DN1/2) prior to the differentiation into $\alpha\beta$ T-cells and $\gamma\delta$ T-cells. Dormant $\text{CD}34^+$ cells may be protected from infection or eliminated after infection. The curative SCT for clonal disease of EBV^+ T/NK-cell progenitors could be explained by the eradication of EBV-infected hematopoietic cells and the reset of virus-specific immunity. Further study on the origin of EBV^+ T/NK-cells may therefore shed some light on the pathogenesis and specific therapy of EBV-associated T/NK-cell LPD/lymphoma.

Conflict of interest

There are no conflicts of interest.

Ethical approval

The genetic study was approved by the ethics committee of Kyushu University, Japan (No. 45).


Acknowledgments

We thank to Drs. Shima T. (Kyushu Univ., Fukuoka, Japan), Wada M. (Otsuka Assay Lab., Tokyo, Japan), and Yamamori S. (Mitsubishi Chem. Med. Corp, Tokyo, Japan) for the quality control of experiments. We thank Dr. Brian Thomas Quinn (Associate Professor, Department of Linguistic Environment, Faculty of Languages and Cultures, Kyushu University) for kindly correcting the manuscript. This work was supported by a Grant-in-Aid for Scientific Research (C2: 60233053) to Ohga S. from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

- Ohga S, Nomura A, Takada H, Hara T. Epstein-Barr virus associated diseases in childhood. Immunological aspects of Epstein-Barr virus infection. *Crit Rev Oncol Hematol* 2002;**44**(3):203–15.
- Kimura H, Hoshino Y, Hara S, Sugaya N, Kawada J, Shibata Y, et al. Differences between T cell-type and natural killer cell-type chronic active Epstein-Barr virus infection. *J Infect Dis* 2005;**191**(4):531–9.
- Ohga S, Sanefuji M, Ishimura M, Nomura A, Torisu H, Kira R, et al. Epstein-Barr virus load in cerebrospinal fluid of patients with chronic active Epstein-Barr virus infection. *Pediatr Infect Dis J* 2008;**27**(11):1027–30.
- Kasahara Y, Yachie A, Takei K, Kanegane C, Okada K, Ohta K, et al. Differential cellular targets of Epstein-Barr virus (EBV) infection between acute EBV-associated hemophagocytic lymphohistiocytosis and chronic active EBV infection. *Blood* 2001;**98**(6):1882–8.
- Ohga S, Nomura A, Takada H, Ihara K, Kawakami K, Yanai F, et al. Epstein-Barr virus (EBV) load and cytokine gene expression in activated T cells of chronic active EBV infection. *J Infect Dis* 2001;**183**(1):1–7.
- Suzuki K, Ohshima K, Karube K, Suzumiya J, Ohga S, Ishihara S, et al. Clinicopathological states of Epstein-Barr virus-associated T/NK-cell lymphoproliferative disorders (severe chronic active EBV infection) of children and young adults. *Int J Oncol* 2004;**24**(5):1165–74.
- Sato E, Ohga S, Kuroda H, Yoshida F, Nishimura M, Nagasawa M, et al. Allogeneic hematopoietic stem cell transplantation for Epstein-Barr virus-associated T/natural killer-cell lymphoproliferative disease in Japan. *Am J Hematol* 2008;**83**(9):721–7.
- Quintanilla-Martinez L, Kimura H, Jaffe ES. EBV-positive T-cell lymphoproliferative disorders of childhood. In: Swerdlow SH, Campo E, Harris NL, et al., editors. *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues*. 4th ed. Lyon, France: International Agency for Research on Cancer; 2008. p. 278–91.

9. Cohen JI, Kimura H, Nakamura S, Ko YH, Jaffe ES. Epstein-Barr virus-associated lymphoproliferative disease in non-immunocompromised hosts: a status report and summary of an international meeting, 8–9 September 2008. *Ann Oncol* 2009;20(9):1472–82.
10. Ohshima K, Kimura H, Yoshino T, Kim CW, Ko YH, Lee SS, et al. Proposed categorization of pathological states of EBV-associated T/natural killer-cell lymphoproliferative disorder (LPD) in children and young adults: overlap with chronic active EBV infection and infantile fulminant EBV T-LPD. *Pathol Int* 2008;58(4):209–17.
11. Elazary AS, Wolf DG, Amir G, Avni B, Rund D, Yehuda DB, et al. Severe Epstein-Barr virus-associated hemophagocytic syndrome in six adult patients. *J Clin Virol* 2007;40(2):156–9.
12. Okano M, Kawa K, Kimura H, Yachie A, Wakiguchi H, Maeda A, et al. Proposed guidelines for diagnosing chronic active Epstein-Barr virus infection. *Am J Hematol* 2005;80(1):64–9.
13. Miyamoto T, Weissman IL, Akashi K. AML1/ETO-expressing nonleukemic stem cells in acute myelogenous leukemia with 8;21 chromosomal translocation. *Proc Natl Acad Sci USA* 2000;97(13):7521–6.
14. Ohga S, Kimura N, Takada H, Nagano M, Ohshima K, Nomura A, et al. Restricted diversification of T-cells in chronic active Epstein-Barr virus infection: potential inclination to T-lymphoproliferative disease. *Am J Hematol* 1999;61(1):26–33.
15. Ohshima K, Karube K, Hamasaki M, Tutiya T, Yamaguchi T, Suefuji H, et al. Differential chemokine, chemokine receptor and cytokine expression in Epstein-Barr virus-associated lymphoproliferative diseases. *Leuk Lymphoma* 2003;44(8):1367–78.
16. Hudnall SD, Ge Y, Wei L, Yang NP, Wang HQ, Chen T. Distribution and phenotype of Epstein-Barr virus-infected cells in human pharyngeal tonsils. *Mod Pathol* 2005;18(4):519–27.
17. Imai S, Sugiura M, Oikawa O, Koizumi S, Hirao M, Kimura H, et al. Epstein-Barr virus (EBV)-carrying and -expressing T-cell lines established from severe chronic active EBV infection. *Blood* 1996;87(4):1446–57.
18. Oyoshi MK, Nagata H, Kimura N, Zhang Y, Demachi A, Hara T, et al. Preferential expansion of V γ 9-J γ P/V δ 2-J δ 3 $\gamma\delta$ T cells in nasal T-cell lymphoma and chronic active Epstein-Barr virus infection. *Am J Pathol* 2003;162(5):1629–38.
19. Maini MK, Gudgeon N, Wedderburn LR, Rickinson AB, Beverley PC. Clonal expansions in acute EBV infection are detectable in the CD8 and not the CD4 subset and persist with a variable CD45 phenotype. *J Immunol* 2000;165(10):5729–37.
20. Toyabe S, Harada W, Uchiyama M. Biclinal expansion of T cells infected with monoclonal Epstein-Barr virus (EBV) in a patient with chronic, active EBV infection. *Clin Exp Immunol* 2003;134(1):92–7.
21. Endo R, Yoshioka M, Ebihara T, Ishiguro N, Kikuta H, Kobayashi K. Clonal expansion of multiphenotypic Epstein-Barr virus-infected lymphocytes in chronic active Epstein-Barr virus infection. *Med Hypotheses* 2004;63(4):582–7.
22. Takakuwa T, Tresnasari K, Rahadiani N, Miwa H, Daibata M, Aozasa K. Cell origin of pyothorax-associated lymphoma: a lymphoma strongly associated with Epstein-Barr virus infection. *Leukemia* 2008;22(3):620–7.
23. Gao J, Luo X, Tang K, Li X, Li G. Epstein-Barr virus integrates frequently into chromosome 4q, 2q, 1q and 7q of Burkitt's lymphoma cell line (Raji). *J Virol Methods* 2006;136(1–2):193–9.
24. Shibata Y, Hoshino Y, Hara S, Yagasaki H, Kojima S, Nishiyama Y, et al. Clonality analysis by sequence variation of the latent membrane protein 1 gene in patients with chronic active Epstein-Barr virus infection. *J Med Virol* 2006;78(6):770–9.
25. Rey J, Xerri L, Bouabdallah R, Keuppens M, Brousset P, Meggetto F. Detection of different clonal EBV strains in Hodgkin lymphoma and nasopharyngeal carcinoma tissues from the same patient. *Br J Haematol* 2008;142(1):79–81.
26. Katano H, Ali MA, Patera AC, Caltafamo M, Jaffe ES, Kimura H, et al. Chronic active Epstein-Barr virus infection associated with mutations in perforin that impair its maturation. *Blood* 2004;103(4):1244–52.
27. Tsuge I, Morishima T, Kimura H, Kuzushima K, Matsuoka H. Impaired cytotoxic T lymphocyte response to Epstein-Barr virus-infected NK cells in patients with severe chronic active EBV infection. *J Med Virol* 2001;64(2):141–8.
28. Ma X, Okamura A, Yosioka M, Ishiguro N, Kikuta H, Kobayashi K. No mutations of SAP/SH2D1A/DSHP and perforin genes in patients with Epstein-Barr virus-associated hemophagocytic syndrome in Japan. *J Med Virol* 2001;65(2):358–61.
29. Lee WI, Chen SH, Hung IJ, Yang CP, Jaing TH, Chen CJ, et al. Clinical aspects, immunologic assessment, and genetic analysis in Taiwanese children with hemophagocytic lymphohistiocytosis. *Pediatr Infect Dis J* 2009;28(1):30–4.
30. Repic AM, Shi M, Scott RS, Sixbey JW. Augmented latent membrane protein 1 expression from Epstein-Barr virus episomes with minimal terminal repeats. *J Virol* 2010;84(5):2236–44.
31. Alexaki A, Liu Y, Wigdahl B. Cellular reservoirs of HIV-1 and their role in viral persistence. *Curr HIV Res* 2008;6(5):388–400.
32. Banerjee P, Sieburg M, Samuelson E, Feuer G. Human T-cell lymphotropic virus type 1 infection of CD34+ hematopoietic progenitor cells induces cell cycle arrest by modulation of p21(cip1/waf1) and survivin. *Stem Cells* 2008;26(12):3047–58.
33. Alexaki A, Wigdahl B. HIV-1 infection of bone marrow hematopoietic progenitor cells and their role in trafficking and viral dissemination. *PLoS Pathog* 2008;4(12):e1000215.



RESEARCH

Open Access

Characterization of Epstein-Barr virus (EBV)-infected cells in EBV-associated hemophagocytic lymphohistiocytosis in two patients with X-linked lymphoproliferative syndrome type 1 and type 2

Xi Yang^{1,2}, Taizo Wada³, Ken-Ichi Imadome⁴, Naonori Nishida¹, Takeo Mukai⁵, Mitsuhiro Fujiwara⁴, Haruka Kawashima⁶, Fumiyo Kato⁶, Shigeyoshi Fujiwara³, Akihiro Yachie³, Xiaodong Zhao², Toshio Miyawaki¹ and Hirokazu Kanegane^{1*}

Abstract

Background: X-linked lymphoproliferative syndrome (XLP) is a rare inherited immunodeficiency by an extreme vulnerability to Epstein-Barr virus (EBV) infection, frequently resulting in hemophagocytic lymphohistiocytosis (HLH). XLP are now divided into type 1 (XLP-1) and type 2 (XLP-2), which are caused by mutations of *SH2D1A/SLAM-associated protein (SAP)* and *X-linked inhibitor of apoptosis protein (XIAP)* genes, respectively. The diagnosis of XLP in individuals with EBV-associated HLH (EBV-HLH) is generally difficult because they show basically similar symptoms to sporadic EBV-HLH. Although EBV-infected cells in sporadic EBV-HLH are known to be mainly in CD8⁺ T cells, the cell-type of EBV-infected cells in EBV-HLH seen in XLP patients remains undetermined.

Methods: EBV-infected cells in two patients (XLP-1 and XLP-2) presenting EBV-HLH were evaluated by in EBV-1 *in situ* hybridization or quantitative PCR methods.

Results: Both XLP patients showed that the dominant population of EBV-infected cells was CD19⁺ B cells, whereas EBV-infected CD8⁺ T cells were very few.

Conclusions: In XLP-related EBV-HLH, EBV-infected cells appear to be predominantly B cells. B cell directed therapy such as rituximab may be a valuable option in the treatment of EBV-HLH in XLP patients.

Keywords: B cells, Epstein Barr virus, Hemophagocytic lymphohistiocytosis, X-linked lymphoproliferative syndrome

Introduction

Hemophagocytic lymphohistiocytosis (HLH) is clinically characterized by prolonged fever, hepatosplenomegaly, hypertriglyceridemia, systemic hypercytokinemia and cytopenia [1]. HLH consists of primary (familial) and secondary (infection, lymphoma or autoimmune disease-associated) types. Approximately half of all infection-associated HLH cases involves the Epstein-Barr virus (EBV) [2]. Most cases of EBV-HLH are sporadic, but a few cases may present the first presentation of X-linked lymphoproliferative syndrome (XLP) [3]. XLP is a rare,

inherited immunodeficiency that is characterized by an extreme vulnerability to EBV infection and shows variable clinical phenotypes, including severe or fatal EBV-HLH (60%), malignant B-cell lymphoma (30%), and progressive dysgammaglobulinemia (30%) [3]. The first gene that is responsible for XLP was identified as the *SH2D1A/SLAM-associated protein (SAP)* gene in 1998 [4-6], and mutations in the *X-linked inhibitor of apoptosis protein (XIAP)* gene can also lead to the clinical phenotype of XLP in 2006 [7]. XLP is now considered to comprise two distinct diseases, namely XLP-1 (SAP deficiency) and XLP-2 (XIAP deficiency).

In addition to B cells, EBV can infect other cell types, including epithelial cells, T cells and natural killer (NK) cells [8]. Studies have shown that activated T cells,

* Correspondence: kanegane@med.u-toyama.ac.jp

¹Department of Pediatrics, Graduate School of Medicine and Pharmaceutical Science, University of Toyama, Toyama, Japan

Full list of author information is available at the end of the article

particularly CD8⁺ T cells, are the primary cellular target of EBV infection in sporadic EBV-HLH [9,10], which reflects the pathogenic role of EBV-infected CD8⁺ T cells in sporadic EBV-HLH. Patients with sporadic EBV-HLH are usually treated with immunochemotherapy, including dexamethasone, cyclosporine A and etoposide, and this therapy can be curable [11]. In contrast, XLP-related EBV-HLH is usually refractory to immunochemotherapy [3]. It is possible that the poor response of XLP-related EBV-HLH to immunochemotherapy can be attributed to the type of EBV-infected cells in this disease, which may differ from the cell type that infected in sporadic EBV-HLH. We investigated the affected cell type in EBV infection of two XLP (XLP-1 and XLP-2) patients with EBV-HLH. Our results demonstrate that the predominant EBV target cells in XLP-related EBV-HLH are CD19⁺ B cells, which appears to be distinct from sporadic EBV-HLH cases.

Patients, materials and methods

Patients

Three patients presented with clinical features of HLH, including persistent fever, hepatosplenomegaly, cytopenia, abnormal liver function, hyperferritinemia and elevated levels of soluble interleukin-2-receptor (Table 1). The clinical features of the patients fulfilled the diagnostic criteria for HLH [1], although hemophagocytosis in

Table 1 Clinical and laboratory finding of the patients in this study

	Patient 1	Patient 2	Patient 3
Family history available	No	Yes	No
Age at the time of the study	4 years	21 months	16 months
Age at onset	3 years	17 months	16 month
Fever	Yes	Yes	Yes
Hepatomegaly	4 cm	5 cm	2.5 cm
Splenomegaly	2 cm	3 cm	1 cm
White blood cells ($\times 10^9/L$)	11.6	6.36	3.03
Neutrophils ($\times 10^9/L$)	1.61	3.915	0.56
Hemoglobin (g/dL)	8.1	9.6	7.5
Platelets ($\times 10^9/L$)	95	56	30
LDH (IU/L)	449	1,693	1,698
AST (IU/L)	88	122	453
ALT (IU/L)	31	25	255
Ferritin ($\mu g/L$)	1,276	26,282	11,129
sIL-2R (U/mL)	3,162	2,880	14,334
IgG (mg/dL)	1,821	806	423
IgA (mg/dL)	302	124	32
IgM (mg/dL)	1,843	40	18
Whole blood EBV-DNA (copies/mL)	140,000	5,700	1,400,000

LDH, lactate dehydrogenase; AST, aspartate amino transferase; ALT, alanine amino transferase, sIL-2R, soluble interleukin-2 receptor; NA: not available.

the bone marrow was not observed in patients 1 and 2. Patient 3 was previously reported as patient HLH3 [10]. The number of EBV-DNA copies in the peripheral blood was increased from the normal level of $\leq 1 \times 10^2$ copies/ml to 1.4×10^5 , 5.7×10^3 and 1.4×10^6 copies/ml in patients 1, 2 and 3, respectively. Blood samples from the patients were obtained using standard ethical procedures with the approval of the Ethics Committee of the University of Toyama, and an analysis of the *SH2D1A* and *XIAP* genes was performed. Patient 1 showed a one-nucleotide insertion (239_240insA) in the *SH2D1A* gene that resulted in a frameshift and a premature stop codon (80KfsX22). Patient 2 carried a two-nucleotides deletion (1021_1022delAA) in the *XIAP* gene that resulted in a frameshift and a premature stop codon (N341YfsX7). Patient 3 had no mutations in the *SH2D1A* or *XIAP* gene.

Cell preparation

Peripheral blood mononuclear cells (PBMCs) were isolated from the patients using Ficoll-Hypaque gradient centrifugation. Lymphocytes were prepared from the PBMCs by depleting the monocytes using anti-CD14 monoclonal antibody (mAb)-coated magnetic beads (Becton Dickinson, San Diego, CA) [10]. The CD19⁺ B cells, CD56⁺ NK cells, CD4⁺ T cells and CD8⁺ T cells were purified by positive selection from the lymphocytes using the respective mAb-coated magnetic beads. The purity of each isolated cell population was assessed by flow cytometry analysis, and each sorted population was found to be higher than 85% pure.

In situ hybridization of EBVRNA

The presence of EBV was estimated by measuring the EBV-encoded small RNA 1 (EBER-1) mRNA using the *in situ* hybridization (ISH) method as described previously [10]. The sorted cells were cytocentrifuged onto silanized slides (Dako, Kyoto, Japan), and the presence of EBER-1 mRNA was determined by ISH using the alkaline phosphatase-conjugated EBER-1 antisense probe (5'-AGCAGAGTCTGGGAAGACAACCACAGACACCGTCCTCACC-3') or a sense probe.

Quantitative PCR for EBV DNA

Quantitative PCR was performed using AmpliTaq Gold and a real-time PCR 7300 system (Applied Biosystems, Foster City, CA) as described previously [12]. The PCR primers for detecting EBV DNA were selected from within the *BALF5* gene, which encodes the viral DNA polymerase. The primers for amplifying the *BALF5* gene sequences were as follows: forward, CGGAAGCCC TCTGGACTTC, and reverse, CCCTGTT TATCCGATGGAATG. The TaqMan probe was FAM-TATACACGCACGAGAAATGCGCC-BFQ. The PCR

conditions were as follows: denaturation at 95°C for 2 minutes, annealing at 58°C for 15 seconds, and extension at 72°C for 15 seconds, and the products were subjected to 45 cycles of PCR amplification. The EBV DNA copy number was considered to be significant when more than 500 copies/ μ g of DNA were observed.

Flow cytometry analysis for the T cell receptor V β repertoire

Flow cytometry analysis of the T cell receptor (TCR) V β repertoire was performed as described previously [10]. In brief, the PBMCs were incubated with the appropriate phycoerythrin-conjugated mAbs with specificity for TCR V β 1-23 (Immunotech, Marseille, France), fluorescein isothiocyanate-conjugated anti-CD8 (Becton Dickinson) and R-PE-Cy5-conjugated anti-CD4 (Dako) mAbs. The stained cells were analyzed using a flow cytometer. TCR V β expression is represented as the percentage of CD4⁺ or CD8⁺ cells for each receptor family.

Results

To determine the localization of EBV infection in the lymphocyte subpopulations of patient 1, CD4⁺ T cells, CD8⁺ T cells, CD19⁺ B cells and CD56⁺ NK cells were sorted using the immunomagnetic bead method and the presence of EBV was evaluated in each lymphocyte subpopulation by EBER-1 ISH (Figure 1A). EBER-1-positive cells were observed in 34.0% of the CD19⁺ B cells, whereas the remaining lymphocyte subpopulations contained fewer than 0.1% EBER-1-positive cells. Therefore, the EBV-infected cells in patient 1 were almost exclusively CD19⁺ B cells. In patient 3, EBER-1-positive cells constituted 75.5% of CD8⁺ T cells, however, they were not detected among CD4⁺ T cells and observed in a few of CD19⁺ B cells and CD56⁺ NK cells (2.8% and 1.7.4%, respectively) (Figure 1B).

The viral loads in the CD4⁺ T cells, CD8⁺ T cells, CD19⁺ B cells and CD56⁺ NK cells in patient 2 were determined by quantitative PCR. The number of EBV DNA genome copies in the CD19⁺ B cells was 1.8×10^4 copies/ μ g, and the copy number in the CD8⁺ T cells was 1.0×10^3 copies/ μ g. The EBV DNA genome could not be detected in either the CD4⁺ T cells or the CD56⁺ NK cells that were isolated from patient 2.


Flow cytometry analysis of the TCR V β repertoire revealed a polyclonal pattern in patients 1 and 2 (Figure 2), which was in contrast to the skewed pattern that is most commonly seen in the CD8⁺ T cells of patients with sporadic EBV-HLH [10]. No clonal dominance in CD8⁺ T cells was demonstrated by mAb in patient 3, but TCR V β 13.3 was predominantly found in the CD8⁺ T cells by complementarity-determining region 3 spectratyping [10].

Discussion

XLP is a severe and rare immunodeficiency disease that is characterized by an extreme vulnerability to EBV infection and frequently results in HLH [3]. XLP was first described as X-linked progressive combined immunodeficiency in 1975 by Purtilo et al. [13]. To better understand and reflect the pathophysiology of this disease, the term "X-linked lymphoproliferative disease or syndrome" has now been used. The first gene to be linked to XLP in 1998 was *SH2D1A* which is located on Xq25 and encodes the SAP [4-6]. Importantly, in 2006, a mutation in the gene that encodes the XIAP was identified as a second XLP-linked gene [7]. Thus, XLP can be divided into XLP-1 (SAP deficiency) and XLP-2 (XIAP deficiency). Most XLP patients present with EBV-HLH. Pachlopnik Schmid et al. [14] reported that the incidence of HLH in XLP-1 and XLP-2 is 55 and 76%, respectively. Currently, hematopoietic stem cell transplantation (HSCT) is the only curative therapy for XLP. Therefore, an early definitive diagnosis and immediate treatment are extremely important for both life-saving intervention and an improved prognosis for XLP patients.

EBV infects the majority of the adult population worldwide and persists in B cells throughout the lifetime of normal individuals, usually without causing disease. EBV is the most common trigger for both the XLP-1 and XLP-2 phenotypes. Prior to being exposed to EBV, most patients with XLP can tolerate infections by other agents, although *in vitro* studies have demonstrated defects of T cell-mediated and humoral immunity. During an acute EBV infection, XLP patients develop normal or high levels of anti-viral capsid antigen IgM antibodies but usually lack heterophile antibodies. Initially, these patients fail to develop EBV-specific cytotoxic T cells, and this results in a massive and overwhelming polyclonal B cell proliferation involving lymphoid and other tissues [8]. SAP binds 2B4, which is a surface molecule involved in activation of NK cell-mediated cytotoxicity. Therefore, SAP-deficient patient shows that NK cell function is impaired, allowing B cell proliferation [15]. SAP has proapoptotic function, and contributes to the maintenance of T cell homeostasis and to the elimination of potentially dangerous DNA-damaged cells. Thus, the loss of this function could be responsible for the uncontrol T cell proliferation in acute EBV infection [16].


B cells are the usual cellular targets of EBV in a primary EBV infection such as infectious mononucleosis and in the sero-positive normal host [8]. After the interaction of the viral surface glycoproteins with the CD21 receptor, EBV entry into B cells is mediated by HLA class II and other co-receptors. However, in cases of



sporadic EBV-HLH, EBV infects primarily T cells and NK cells [9,10,17]. The mechanism of T cell infection by EBV in HLH is still unclear, but one hypothesis is that, in specific situations, CD8⁺ T cells express CD21, which can mediate EBV infection. Although T cells do not express the glycoprotein, they contain mRNA for CD21 [18]. In sporadic EBV-HLH cases, EBV infection into B cells is delayed but occurs during every case of cured EBV-HLH [17]. To the best of our knowledge, this is the first report of EBV infection status in two different types of XLP patients with EBV-HLH. The present study shows that the primary EBV-infected cells in XLP-related EBV-HLH are CD19⁺ B cells and not T cells or NK cells, which are a primary target of EBV infection in sporadic EBV-HLH.

For decades, clinicians and investigators have been puzzled by the differential diagnosis between XLP and sporadic EBV-HLH when they encountered a young boy presenting with EBV-HLH. We believe the differential EBV target cells can provide additional information

to help discriminate between XLP and sporadic EBV-HLH. An evaluation of specific cell type that is infected by EBV should be considered when target therapy is applied. Most patients with sporadic EBV-HLH can achieve remission by immunochemotherapy; however, patients with XLP are usually refractory to this therapy. Recently, B cell-directed therapy using an anti-CD20 mAb (rituximab) was performed in patients with XLP-1 [19]. Two XLP patients who presented with acute EBV infection were successfully treated with rituximab and were free from EBV-HLH and lymphoma for a prolonged period. In addition, rituximab combined with methylprednisolone and intravenous immunoglobulin were administered to an XLP-1 patient with EBV-HLH, and the patient achieved a remission [20]. Patient 1 was also associated with EBV-associated encephalitis and lymphoproliferative disorder. The patient's lymphoproliferative disorder was treated with rituximab, but he died of the disease. Patient 2 was successfully treated with dexamethasone



and immunoglobulin. Our data suggest that B cell target therapy can be a viable therapeutic option for an initial stage of EBV-HLH in both XLP-1 and XLP-2 patients.

Abbreviations

EBER: EBV-encoded small RNA; EBV: Epstein-Barr virus; HLH: Hemophagocytic lymphohistiocytosis; HSCT: Hematopoietic stem cell transplantation; ISH: *In situ* Hybridization; mAb: Monoclonal antibody; NK: Natural killer; PBMC: Peripheral blood mononuclear cells; SAP: SLAM-associated protein; TCR: T cell receptor; XIAP: X-linked inhibitor of apoptosis; XLP: X-linked lymphoproliferative syndrome.

Acknowledgements

The authors would like to thank Ms. Chikako Sakai and Mr. Hitoshi Moriuchi for their excellent technical assistances. This study was supported by a grant-in-aid for scientific research from the Ministry of Education, Culture, Sports, Science and Technology, Japan (TM, HK), and a grant from the Ministry of Health, Labour, and Welfare, Japan (TM).

Author details

¹Department of Pediatrics, Graduate School of Medicine and Pharmaceutical Science, University of Toyama, Toyama, Japan. ²Division of Immunology, Children's Hospital of Chongqing Medical University, Chongqing, China. ³Department of Pediatrics, School of Medicine, Institute of Medical, Pharmaceutical and Health Sciences, Kanazawa University, Kanazawa, Japan. ⁴Department of Infectious Diseases, National Research Institute for Child Health and Development, Tokyo, Japan. ⁵Department of Pediatrics, Kurashiki Central Hospital, Kurashiki, Japan. ⁶Department of Pediatrics, Tokyo Women's Medical University Medical Center East, Tokyo, Japan.

Authors' contributions

XY and HK wrote the manuscript. XY, TW, KI and NN performed the experimental studies. TM, MF, HK and FK managed the patients' care. SF, AY, XDZ and TM revised the manuscript. XY, TW and KI contributed equally to this study. All the authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 1 December 2011 Accepted: 10 February 2012
 Published: 10 February 2012

References

1. Henter JL, Horne A, Aricó M, Egeler RM, Fillipovich AH, Imashuku S, Ladisch S, McClain K, Webb D, Winiarski J, et al: **HLH-2004: Diagnostic and therapeutic guideline for hemophagocytic lymphohistiocytosis.** *Pediatr Blood Cancer* 2007, **48**:124-131.
2. Ishii E, Ohga S, Imashuku S, Yasukawa M, Tsuda H, Miura I, Yamamoto K, Horiuchi H, Takada K, Ohshima K, et al: **Nationwide survey of hemophagocytic lymphohistiocytosis in Japan.** *Int J Hematol* 2007, **86**:58-65.
3. Seemayer TA, Gross TG, Egeler RM, Pirruccello SJ, Davis JR, Kelly CM, Okano M, Lanyi A, Sumegi J: **X-linked lymphoproliferative disease: twenty- five years after the discovery.** *Pediatr Res* 1995, **38**:471-478.
4. Coffey AJ, Brooksbank RA, Brandau O, Oohashi T, Howell GR, Bye JM, Cahn AP, Durham J, Heath P, Wray P, et al: **Host response to EBV infection in X-linked lymphoproliferative disease results from mutations in an SH2-domain encoding gene.** *Nat Genet* 1998, **20**:129-135.
5. Sayos J, Wu C, Morra M, Wang N, Zhang X, Allen D, Van-Schaik S, Notarangelo L, Geha R, Roncarolo MG, et al: **The X-linked lymphoproliferative-disease gene product SAP regulates signals induced through the co-receptor SLAM.** *Nature* 1998, **395**:462-469.
6. Nichols KE, Harkin DP, Levitz S, Krainer M, Kolquist KA, Genovese C, Bernard A, Ferguson M, Zuo L, Snyder E, et al: **Inactivating mutations in an SH2 domain-encoding gene in X-linked lymphoproliferative syndrome.** *Proc Natl Acad Sci USA* 1998, **95**:13765-13770.
7. Rigaud S, Fondanèche MC, Lambert N, Pasquier B, Mateo V, Soulas P, Galicier L, Le-Deist F, Rieux-Laucat F, Revy P, et al: **XIAP deficiency in humans causes an X-linked lymphoproliferative syndrome.** *Nature* 2006, **444**:110-114.
8. Tosato G, Taga K, Angiolillo AL, Sgadari C: **Epstein-Barr virus as an agent of haematological disease.** *Baillière Clin Haematol* 1995, **8**:165-199.
9. Kawaguchi H, Miyashita T, Herbst H, Niedobitek G, Asada M, Tsuchida M, Hanada R, Kinoshita A, Sakurai M, Kobayashi N, et al: **Epstein-Barr virus-infected T lymphocytes in Epstein-Barr virus-associated hemophagocytic syndrome.** *J Clin Invest* 1993, **92**:1444-1450.
10. Toga A, Wada T, Sakakibara Y, Mase S, Araki R, Tone T, Toma T, Kurokawa T, Yanagisawa R, Tamura K, et al: **Clinical significance of cloned expansion and CD5 down-regulation in Epstein-Barr Virus (EBV)-infected CD8+ T lymphocytes in EBV-associated hemophagocytic lymphohistiocytosis.** *J Infect Dis* 2010, **201**:1923-1932.
11. Imashuku S, Hibi S, Ohara T, Iwai A, Sako M, Kato M, Arakawa H, Sotomatsu M, Kataoka S, Asami K, et al: **Effective control of Epstein-Barr virus-related hemophagocytic lymphohistiocytosis with immunochemotherapy.** *Blood* 1999, **93**:1869-1874.
12. Kimura H, Hoshino Y, Kanegane H, Tsuge I, Okamura T, Kawa K, Morishima T: **Clinical and virologic characteristics of chronic active Epstein-Barr virus infection.** *Blood* 2001, **98**:280-286.
13. Purtilo DT, Cassel CK, Yang JP, Harper R: **X-linked recessive progressive combined variable immunodeficiency (Duncan's disease).** *Lancet* 1975, **1**:935-940.
14. Pachlounik Schmid J, Canioni D, Moshous D, Moshous D, Touzot F, Mahlaoui N, Hauck F, Kanegane H, Lopez-Granados E, Mejstrikova E, et al: **Clinical similarities and differences of patients with X-linked lymphoproliferative syndrome type 1 (XLP-1/SAP deficiency) versus type 2 (XLP-2/XIAP deficiency).** *Blood* 2011, **117**:1522-1529.
15. Parolini S, Bottino C, Falco M, Augugliaro R, Gilliani S, Franceschini R, Ochs HD, Wolf H, Bonnefoy JY, Biassoni R, et al: **X-linked lymphoproliferative disease: 2B4 molecules displaying inhibitory rather than activating function are responsible for the inhibit of natural killer cells to kill Epstein-Barr virus-infected cells.** *J Exp Med* 2000, **192**:337-346.
16. Nagy N, Matskova L, Kis LL, Hellmah U, Klein G, Klein E: **The proapoptotic function of SAP provides a clue to the clinical picture of X-linked lymphoproliferative disease.** *Proc Natl Acad Sci USA* 2009, **106**:11966-11971.
17. Kasahara Y, Yachie A: **Cell type specific infection of Epstein-Barr virus (EBV) in EBV-associated hemophagocytic lymphohistiocytosis and chronic active EBV infection.** *Crit Rev Oncol Hematol* 2002, **44**:283-294.
18. Braun M, Melchers I, Peter HH, Illges H: **Human B and T lymphocytes have similar amounts of CD21 mRNA, but differ in surface expression of the CD21 glycoprotein.** *Int Immunol* 1998, **10**:1197-1202.
19. Milone MC, Tsai DE, Hodinka RL, Silverman LB, Malbran A, Wasik MA, Nichols KE: **Treatment of primary Epstein-Barr virus infection in patients**

with X-linked lymphoproliferative disease using B-cell-directed therapy. *Blood* 2005, **105**:994-996.

20. Bond J, Shahdadpuri R, Mc Mahon C, O'Marcaigh A, Cotter M, Smith O: **Successful treatment to acute Epstein-Barr virus infection associated with X-linked lymphoproliferative disorder with rituximab.** *Pediatr Blood Cancer* 2007, **49**:761-762.

doi:10.1186/2042-4280-3-1

Cite this article as: Yang et al.: Characterization of Epstein-Barr virus (EBV)-infected cells in EBV-associated hemophagocytic lymphohistiocytosis in two patients with X-linked lymphoproliferative syndrome type 1 and type 2. *Herpesviridae* 2012 **3**:1.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



Clinical and Genetic Characteristics of XIAP Deficiency in Japan

Xi Yang · Hirokazu Kanegane · Naonori Nishida ·
Toshihiko Imamura · Kazuko Hamamoto ·
Ritsuko Miyashita · Kohsuke Imai ·
Shigeaki Nonoyama · Kazunori Sanayama ·
Akiko Yamaide · Fumiyo Kato · Kozo Nagai ·
Eiichi Ishii · Menno C. van Zelm · Sylvain Latour ·
Xiao-Dong Zhao · Toshio Miyawaki

Received: 26 July 2011 / Accepted: 14 December 2011 / Published online: 8 January 2012
© Springer Science+Business Media, LLC 2012

Abstract Deficiency of X-linked inhibitor of apoptosis (XIAP) caused by *XIAP/BIRC4* gene mutations is an inherited immune defect recognized as X-linked lymphoproliferative syndrome type 2. This disease is mainly observed in patients with hemophagocytic lymphohistiocytosis (HLH) often associated with Epstein–Barr virus infection. We described nine Japanese patients from six unrelated families with XIAP deficiency and studied XIAP protein

expression, *XIAP* gene analysis, invariant natural killer T (iNKT) cell counts, and the cytotoxic activity of CD8⁺ alloantigen-specific cytotoxic T lymphocytes. Of the nine patients, eight patients presented with symptoms in infancy or early childhood. Five patients presented with recurrent HLH, one of whom had severe HLH and died after cord blood transplantation. One patient presented with colitis, as did another patient's maternal uncle, who died of colitis at

X. Yang · H. Kanegane (✉) · N. Nishida · T. Miyawaki
Department of Pediatrics, Graduate School of Medicine,
University of Toyama,
2630 Sugitani,
Toyama 930-0194, Japan
e-mail: kanegane@med.u-toyama.ac.jp

X. Yang · X.-D. Zhao
Division of Immunology,
Children's Hospital of Chongqing Medical University,
Chongqing, China

T. Imamura
Department of Pediatrics, Graduate School of Medical Sciences,
Kyoto Prefectural University of Medicine,
Kyoto, Japan

K. Hamamoto
Department of Pediatrics, Hiroshima Red Cross Hospital,
Hiroshima, Japan

R. Miyashita
Department of Pediatrics, Izumiotsu Municipal Hospital,
Izumiotsu, Japan

K. Imai · S. Nonoyama
Department of Pediatrics, National Defense Medical College,
Tokorozawa, Japan

K. Sanayama
Department of Pediatrics, Japanese Red Cross Narita Hospital,
Narita, Japan

A. Yamaide
Department of Allergy and Rheumatology,
Chiba Children's Hospital,
Chiba, Japan

F. Kato
Department of Pediatrics,
Tokyo Women's Medical University Medical Center East,
Tokyo, Japan

K. Nagai · E. Ishii
Department of Pediatrics,
Ehime University Graduate School of Medicine,
Toon, Japan

M. C. van Zelm
Department of Immunology, Erasmus MC,
Rotterdam, The Netherlands

S. Latour
INSERM U768, Hôpital Necker-Enfants Malades,
Paris, France

4 years of age prior to diagnosis with XIAP deficiency. Interestingly, a 17-year-old patient was asymptomatic, while his younger brother suffered from recurrent HLH and EBV infection. Seven out of eight patients showed decreased XIAP protein expression. iNKT cells from patients with XIAP deficiency were significantly decreased as compared with age-matched healthy controls. These results in our Japanese cohort are compatible with previous studies, confirming the clinical characteristics of XIAP deficiency.

Keywords X-linked lymphoproliferative syndrome · X-linked inhibitor of apoptosis · Epstein–Barr virus · hemophagocytic lymphohistiocytosis · invariant natural killer T cell

Abbreviations

BIR	Baculovirus IAP repeat
CTL	Cytotoxic T lymphocyte
HSCT	Hematopoietic stem cell transplantation
HLH	Hemophagocytic lymphohistiocytosis
IAP	Inhibitor of apoptosis
LCL	Lymphoblastoid cell line
MMC	Mitomycin C
mAb	Monoclonal antibody
MFI	Mean fluorescence intensity
iNKT	Invariant natural killer T
PCR	Polymerase chain reaction
PBMC	Peripheral blood mononuclear cells
TCR	T cell receptor
XIAP	X-linked inhibitor of apoptosis
XLP	X-linked lymphoproliferative syndrome

Introduction

X-linked lymphoproliferative syndrome (XLP) is a rare inherited immunodeficiency estimated to affect approximately one in one million males, although it may be underdiagnosed [1]. XLP is characterized by extreme vulnerability to Epstein–Barr virus (EBV) infection, and the major clinical phenotypes of XLP include fulminant infectious mononucleosis (60%), lymphoproliferative disorder (30%), and dysgammaglobulinemia (30%) [2]. In addition, XLP is associated with a variety of additional clinical phenotypes such as vasculitis, aplastic anemia, and pulmonary lymphoid granulomatosis. Patients with XLP often develop more than one of these phenotypes. The gene responsible for XLP was identified as *SH2D1A*, located on Xq25 and encoding the SLAM-associated protein (SAP) [3–5]. However, gene analysis revealed *SH2D1A* mutations in only 50–60% of presumed XLP patients [6]. Importantly, a mutation in the gene that encodes the X-linked inhibitor of

apoptosis (XIAP) called *XIAP* or *BIRC4* was identified as a second causative gene for XLP [7]. *XIAP* is located close to the *SH2D1A* gene on the X chromosome and consists of six coding exons [8–10]. XIAP produces an anti-apoptotic molecule that belongs to the inhibitor of apoptosis (IAP) family proteins. It contains three baculovirus IAP repeat (BIR) domains that, together with flanking residues, bind to caspases 3, 7, and 9, thereby inhibiting their proteolytic activity [11].

The clinical presentations of XIAP-deficient patients have been frequently reported [7,12,13]. More than 90% of patients with XIAP deficiency develop hemophagocytic lymphohistiocytosis (HLH) which is often recurrent. Therefore, it was recently suggested that the phenotype of XIAP deficiency fits better with the definition of familial HLH than with XLP disease [12]. However, familial HLH is characterized by defects in CD8⁺ T and NK cell cytotoxicity responses, while these responses are normal in XIAP deficiency [7,12]. Other symptoms of XLP, such as splenomegaly, hypogammaglobulinemia, and hemorrhagic colitis, have been reported in patients with XIAP deficiency, but lymphoma has never been noted [7,12–15].

We searched for patients with XIAP deficiency in Japan by detection of *XIAP* gene mutations and flow cytometric assessment of lymphoid XIAP expression. We previously reported the first case of XIAP deficiency in Japan [14]. Thereafter, we identified eight additional cases from five families with XIAP deficiency in our country. In this study, we describe the clinical and laboratory findings from nine patients from six unrelated families with XIAP deficiency, including previous cases, to help further the understanding of the pathogenetic features of this disease.

Materials and Methods

Patient and Family Member Samples

Patients without identified *SH2D1A* mutations but with presumed XLP phenotypes were screened for *XIAP* mutations. Their family members were also screened for the same mutation. Upon identification of *XIAP* mutations, the patients were enrolled in this study. Patient 2.2 passed away before a genetic diagnosis of XIAP deficiency was made, but he was the maternal uncle of patient 2.1 and had presented with a XLP phenotype (Table I). In the end, nine patients from six different families were found to have XIAP deficiencies, three of whom had been reported previously [13,14]. Upon the approval of the Ethics Committee of the University of Toyama and after obtaining informed consent, 5–10 mL heparinized venous blood was collected from the patients, their mothers, and 25 age-matched healthy children (1–13 years of age). All of the samples were

transferred to our laboratory at room temperature within 24 h for analysis.

Mutation Analysis of the *XIAP* Gene

DNA was extracted from peripheral blood using the QuickGene-Mini 80 nucleic acid extraction system (FUJI-FILM Co., Tokyo, Japan). The coding regions and the exon–intron boundaries of the *XIAP* gene were amplified by polymerase chain reaction (PCR) using primers flanking each of the six exons by standard methods. PCR products were sequenced using the BigDye Terminator Cycle Sequencing Kit (Applied Biosystems, Foster City, CA, USA) with the same primers used for PCR amplification. Sequencing analysis was performed on an Applied Biosystems Prism 310 Capillary Sequencer (Applied Biosystems).

Flow Cytometric Analysis of XIAP Protein Expression in Lymphocytes

XIAP protein expression was studied by flow cytometric techniques as previously described [16,17]. Peripheral blood mononuclear cells (PBMC) from patients 1, 2.1, 3.1, 3.2, 4, 5, 6.1, 6.2, and 25 age-matched healthy children were prepared by density gradient centrifugation over Histopaque-1077 (Sigma-Aldrich, Inc., St. Louis, MO, USA). The cells were first fixed in 1% paraformaldehyde in PBS for 30 min at room temperature and then permeabilized in 0.5% saponin in washing buffer. The fixed and permeabilized cells were then incubated with an anti-XIAP monoclonal antibody (mAb) (clone 48 (BD Biosciences, Franklin Lakes, NJ, USA) or clone 2 F1 (Abcam, Cambridge, UK)) for 20 min on ice, washed, and then incubated with a FITC-labeled anti-mouse IgG1 antibody (SouthernBiotech, Birmingham, AL, USA) for 20 min on ice. The stained cells were analyzed on the FC500 flow cytometer (Beckman Coulter, Tokyo, Japan).

Western Blot Analysis of XIAP Protein Expression in Lymphocytes

PBMC from normal controls and patients 3.1, 5, and 6.2 were washed and pelleted. The cells were then lysed in 10 μ L of lysing solution (1% Triton-X 100; 150 mmol/L NaCl; 10 mmol/L Tris–HCl, pH 7.6; 5 mmol/L EDTA–Na; 2 mmol/L phenylmethylsulfonyl fluoride) per 10^6 cells for 30 min on ice. The lysed cells were centrifuged for 10 min at 15,000g to remove nuclei, and the supernatants were diluted in the same volume of Laemmli's sample buffer. Samples were then electrophoresed in sodium dodecyl sulfate–polyacrylamide 10% to 20% gradient gel and blotted on nitrocellulose filters. Blots were blocked in 5% skim milk in PBS for 1 h, treated with anti-XIAP mAb (clone 28 or clone 2F1) for 2 h, and then incubated with peroxidase-conjugated

anti-mouse IgG antibody (Invitrogen, Grand Island, NY, USA) for 1 h. Immunoblots were developed by the ECL Western blotting detection system (GE Healthcare UK Ltd., Buckinghamshire, England).

Flow Cytometric Identification of Invariant Natural Killer T Cells

PBMC from eight patients (1, 2.1, 3.1, 3.2, 4, 5, 6.1, and 6.2) and 25 controls were incubated with fluorochrome-conjugated anti-CD3 (Dako Japan KK, Kyoto, Japan), anti-TCRV α 24, and anti-TCRV β 11 mAbs (Beckman Coulter) to identify invariant natural killer T (iNKT) cells by flow cytometry. After the electronic gating of 100,000 CD3⁺ T cells, iNKT cell populations were defined by the co-expression of TCRV α 24 and TCRV β 11. The iNKT cell counts were evaluated at the diagnosis of XIAP deficiency.

Establishment of Alloantigen-Specific Cytotoxic T Lymphocyte Lines and Analysis of Cytotoxic T Lymphocyte-Mediated Cytotoxicity

Alloantigen-specific CD8⁺ cytotoxic T lymphocyte (CTL) lines were generated as described previously [18,19]. Briefly, PBMC were obtained from patients 1, 2.1, 3.1, and unrelated healthy individuals. These cells were co-cultured with a mitomycin C (MMC)-treated B lymphoblastoid cell line (LCL) established from an HLA-mismatched individual (KI-LCL). Using cell isolation immunomagnetic beads (MACS beads; Miltenyi Biotec, Auburn, CA, USA), CD8⁺ T lymphocytes were isolated from PBMC that had been stimulated with KI-LCL for 6 days. CD8⁺ T lymphocytes were cultured in RPMI 1640 medium supplemented with 10% human serum and 10 IU/mL interleukin-2 (Roche, Mannheim, Germany) and stimulated with MMC-treated KI-LCL three times at 1-week intervals. These lymphocytes were then used as CD8⁺ alloantigen-specific CTL lines. The cytotoxic activity of CTLs was measured by a standard ⁵¹Cr-release assay as described previously [20]. Briefly, alloantigen-specific CTLs were incubated with ⁵¹Cr-labeled allogeneic KI-LCL or TA-LCL, which did not share HLA antigens with KI-LCL, for 5 h at effector/target cell ratios (E/T) of 2.5:1, 5:1, and 10:1. Target cells were also added to a well containing only medium and to a well containing 0.2% Triton X-100 to determine the spontaneous and maximum levels of ⁵¹Cr release, respectively. After 5 h, 0.1 mL of supernatant was collected from each well. The percentage of specific ⁵¹Cr release was calculated as follows: (cpm experimental release – cpm spontaneous release) / (cpm maximal release – cpm spontaneous release) \times 100, where cpm indicates counts per minute.

Table 1 Summary of our data

	Patient 1 [13]	Patient 2.1 [12]	Patient 2.2 [12]	Patient 3.1	Patient 3.2	Patient 4	Patient 5	Patient 6.1	Patient 6.2
Age at initial presentation	20 months	7 months	3 months	2 months	Asymptomatic	2 months	6 months	17 months	15 months
Current age	4 years	Deceased	Died of colitis	12 years	17 years	15 years	2 years	1 year	12 years
Family history	No	Yes	Yes	Yes	Yes	No	No	Yes	Yes
HLH	+	+	-	+	-	-	+	+	+
Recurrent HLH	+	+	-	+	-	-	+	-	+
Fever	+	+	+	+	-	-	+	+	+
Splenomegaly	+	+	ND	-	-	-	-	+	+
Cytopenia	+	+	ND	+	-	-	+	+	+
EBV	+	-	ND	+	-	-	-	+	+
Hypogammaglobulinemia	-	+	ND	-	-	+	-	-	-
Colitis	-	-	+	-	-	-	+	-	-
Treatment	PSL CsA Dex	PSL CsA Dex	ND	PSL CsA	-	IVIG	PSL, Dex CsA, IVIG Infliximab	IVIG, Dex	PSL
Allogeneic HSCT	-	+	-	-	-	-	-	-	-
Mutation	R238X	R381X	ND	W217CfsX27	W217CfsX27	E349del	Del of exons 1-2	N341YfsX7	N341YfsX7
XIAP protein expression	±	-	ND	-	-	+	±	±	±

HLH hemophagocytic lymphohistiocytosis, *ND* no data, *EBV* Epstein–Barr virus, *PSL* prednisolone, *CsA* cyclosporin A, *Dex* dexamethasone, *IVIG* intravenous immunoglobulin, *HSCT* hematopoietic stem cell transplantation, + yes or positive, - no or negative, ± residual expression

Statistical Analysis

Student's *t*-test was used for statistics, with *P*-values <0.05 considered to be statistically significant.

Results

Clinical Manifestations of the Patients

Most of our patients presented with disease symptoms at very early ages; five patients presented in infancy and three patients presented in childhood (Table I). Three of the six families had family history records. Five of the nine patients had recurrent HLH, fever, splenomegaly, and cytopenia. EBV infection and hypogammaglobulinemia were also observed in multiple patients. Most patients with HLH were treated with corticosteroids with or without cyclosporin A to prevent an otherwise rapidly fatal disease course. Patients 2.2 and 5 presented with colitis, whereas patient 2.2 died; patient 5 improved with anti-TNF alpha mAb (infliximab®) treatment. Patient 2.1 underwent cord blood transplantation but died of complications. Patient 4 had a history of recurrent otitis media and pneumonia since 2 months of age, and he was found to have hypogammaglobulinemia. The patient was treated with intravenous immunoglobulin replacement therapy alone, and he is currently doing well. No patient developed lymphoma.

Detection of *XIAP* Mutations

We identified *XIAP* mutations in patients from all six unrelated families (Fig. 1) and analyzed all of the data using the US National Center for Biotechnology Information database




Fig. 1 *XIAP* gene mutations and their consequences for XIAP protein. *XIAP* comprises six exons and encodes the XIAP protein, which consists of 497 amino acids. XIAP contains three BIR domains and one RING domain. Mutations identified in our patients are indicated

(<http://www.ncbi.nlm.nih.gov/SNP>) to check for single-nucleotide polymorphism in the *XIAP* gene. As previously reported, patient 1 possessed a nonsense mutation, 712 C > T, resulting in an early stop codon R238X [14]. Patient 2.1 had a nonsense mutation in exon 5, 1141 C > T, resulting in R381X [13]. Patient 2.2 might have the same mutation as patient 2.1 because patient 2.2 was the maternal uncle of patient 2.1 [13]. Patients 3.1 and 3.2 were siblings and were found to have a one base pair deletion (650delG) in exon 1, resulting in a frameshift and premature stop codon (W217CfsX27). Patient 4 was found to have one amino acid deletion (1045_1047delGAG; E349del) in exon 3. Patient 5 has a large deletion, spanning exons 1 and 2. Patients 6.1 and 6.2 were brothers and had a two-nucleotide deletion (1021_1022delAA), which resulted in a frameshift and premature stop codon (N341YfsX7). All of the mothers of the patients from families 1–5 were heterozygote carriers of the mutations. Interestingly, we could not find any *XIAP* mutation in the mother of patients 6.1 and 6.2. We identified deleterious *XIAP* mutations in nine patients from six unrelated Japanese families that are likely to underlie their XLP phenotypes.

XIAP Expression in Lymphocytes from the Patients and Carriers by Flow Cytometry

XIAP expression levels were analyzed in the lymphocytes of patients from all six families (Fig. 2). The lymphocytes of




Fig. 2 XIAP protein expression in lymphocytes from the patients and their carriers. Flow cytometric detection of intracellular XIAP in lymphocytes from patients and their maternal carriers. The gray and black areas indicate the negative control and anti-XIAP staining, respectively. Anti-XIAP staining was performed using the clones 48 and 2 F1 antibodies where indicated. The number in the box indicates the log scale difference between the mean fluorescence intensity (Δ MFI) stained by the isotype antibody and that by the anti-XIAP antibodies. XIAP expression in 25 normal controls was also analyzed by the clone 48 and 2 F1 antibodies. The data of mean \pm standard deviation of Δ MFI and each representative profile were shown

patients 1, 3.1, 5, 6.1, and 6.2 were examined by two different anti-XIAP mAbs. Using clone 48 antibody, patients 1, 2.1, 3.1, 3.2, 6.1, and 6.2 showed reduced XIAP expression, whereas XIAP was normally expressed in the lymphocytes of patients 4 and 5. In contrast to clone 48, clone 2F1 antibody showed reduced XIAP expression in patient 5. The effects of heterozygous *XIAP* mutations were studied in the lymphocytes of the patients' mothers by anti-XIAP mAb clone 48. The mothers of patients 1, 3.1, and 3.2 showed a bimodal pattern of XIAP protein (Fig. 2). The mothers of patients 2.1, 6.1, and 6.2 did not show a clear mosaic pattern, but all of these patients had reduced XIAP expression levels. Similarly to patients 4 and 5, the mothers of patients 4 and 5 demonstrated a normal XIAP expression pattern.

XIAP Expression in Lymphocytes from the Patients by Western Blot

Western blot analysis was used to evaluate the expression level of XIAP to determine the impact of patient *XIAP* mutations on protein expression and to compare this to the flow cytometric analysis. PBMCs from patients 3.1, 5.1, and 6.2 were available for Western blotting. All of these patients showed a reduction in XIAP protein expression (Fig. 3), fitting with the results obtained by flow cytometric analysis.

iNKT Cell Counts in the Patients

SAP-deficient patients had reduced numbers of NKT cells that expressed an invariantly rearranged T-cell receptor (TCR) consisting of TCRV α 24 and TCRV β 11 chains [21,22]. The rare subset of iNKT cells was originally reported to be reduced in XIAP-deficient patients as well [7] but seemed to be present in normal numbers in a later study involving a larger patient cohort [23]. We analyzed the iNKT cell frequencies in 100,000 CD3⁺ T cells in our XIAP-deficient patients and compared these with healthy controls (Fig. 4). The average frequency of iNKT cells within the CD3⁺ T cell compartment of our XIAP patients was significantly reduced by twofold when compared with healthy




Fig. 3 XIAP expression in lymphocytes from the patients by Western blot. Analysis of XIAP expression in PBMC generated from patients with XIAP deficiency and normal controls using the antibody clone 48 (upper panel), the antibody clone 2 F1 (middle panel), and the β -actin antibody as an internal control (lower panel)




Fig. 4 iNKT cell counts in the patients and healthy controls. **a** Representative flow cytometric analysis of iNKT cells in CD3⁺ lymphocytes from one XIAP-deficient patient and one healthy control. **b** Comparison of the number of iNKT cells in 100,000 CD3⁺ lymphocytes between XIAP-deficient patients and control individuals. Statistical significance between patients and controls was determined with the Student's *t*-test (*p*-value=0.027)

controls (20 vs. 47 per 10⁵ CD3⁺ T cells). Therefore, we concluded that the number of iNKT cells was reduced in our patients with XIAP deficiency.


Functional Analysis of CTL Lines Established from the Patients

To test whether our XIAP-deficient patients have similar defects in CD8⁺ T cell cytotoxicity as described in other subtypes of familial HLH [20,38], we generated CD8⁺ alloantigen-specific CTL from patients 1, 2.1, 3.1, and three healthy controls (Fig. 5). The cytotoxic activity of the CTL of these patients was similar to that of the healthy controls, indicating that XIAP patients clearly differ from other familial HLH patients in this aspect of the disease.

Discussion

XIAP deficiency is a rare but severe and life-threatening inherited immune deficiency [12,13]. Early diagnosis and life-saving treatment such as hematopoietic stem cell transplantation is especially important. The causative gene for

Fig. 5 Cytotoxicity of alloantigen-specific CD8⁺ T cell lines. CD8⁺ T cell lines were generated from PBMC of patients with XIAP deficiency and healthy controls by stimulation with allogeneic LCL (KI-LCL). Their cytotoxicity was determined against allogeneic KI-LCL (circles) and against allogeneic TA-LCL (triangles), which does not share alloantigens with KI-LCL



XIAP deficiency was identified to be *XIAP/BIRC4*, and 25 mutations in the *XIAP* gene have been previously reported [7,12–14]. In the present study, we described four novel mutations (W217CfsX27, E349del, deletion of exons 1 and 2 and N341YfsX7) in the *XIAP* genes as well as previously described patients with R381X and R238X mutations [13,14]. The mother of patients 6.1 and 6.2 had no mutation in the *XIAP* gene. Because this is an X-linked inheritance, the failure to identify the same mutation in the mother suggests that the mother had a germline mosaicism for the mutation. Such mosaicism has not yet been described in XIAP deficiency, but it has been reported in Duchenne muscular dystrophy, X-linked severe combined immunodeficiency, X-linked agammaglobulinemia, and many other inherited diseases [24–26]. HLH is common in XIAP-deficient patients, and it is often recurrent [13,14]. In our study, six patients had HLH and five patients presented with recurrent HLH. Therefore, XIAP deficiency should be suspected in certain boys with HLH, especially in those with family history or recurrent HLH. The reason why XIAP deficiency increases susceptibility to HLH remains unclear. Murine studies have also failed to disclose a mechanism for the development of HLH [27]. Interestingly, *Xiap*-deficient mice possess normal lymphocyte apoptosis induced by a variety of means [28]. Three of our patients presented with EBV-associated HLH. EBV infection has been reported to be a trigger of the first HLH episode in patients with XIAP deficiency [13]. The excess of lymphocyte apoptosis in XIAP deficiency might account for the abnormal immune response to EBV [28]. Splenomegaly is not frequently observed in XLP type 1 or SAP deficiency but might be a common clinical feature in XIAP deficiency [12,13] as four (50%) of eight Japanese patients developed splenomegaly. Pachlopnik Schmid et al. [13] reported that recurrent splenomegaly occurring in the absence of systemic HLH was often

associated with fever and cytopenia. XIAP-deficient patients are at risk for chronic colitis, which is possibly a more frequent cause of mortality than HLH [13]. Our study included two patients who developed colitis, and one of the patients died of colitis at 4 years of age. Although we did not have enough clinical information or samples from that patient because of his early death, his symptoms suggest that he had a XIAP deficiency complicated with colitis because he was the maternal uncle of patient 2.1. The other patient was 2 years old and also suffered from chronic hemorrhagic colitis.

In contrast to SAP deficiency, lymphoma has never been reported in XIAP deficiency, including our patients. Some studies indicate that the XIAP protein is a potential target for the treatment of cancer based on the anti-apoptotic function of XIAP [29]. Therefore, the absence of XIAP may protect patients from cancer, explaining why XIAP-deficient patients do not develop lymphoma. We generated a clinical summary to compare XIAP-deficient patients with the previous reports (Table II). Although our study included a relatively small number of patients, our results appear to be consistent with previous large studies [12,13] and confirm the clinical characteristics of XIAP deficiency.

Flow cytometry can be used for the rapid screening of several primary immunodeficiencies including XLP [30]. XIAP protein has been found to be expressed in various human tissues, including all hematopoietic cells [7,10]. Marsh et al. [16] described that XIAP was readily detectable in normal granulocytes, monocytes, and all lymphocyte subsets. Moreover, patients with *XIAP* mutations had decreased or absent expression of XIAP protein by flow cytometry [14,16]. We investigated XIAP expression in lymphocytes from eight patients by flow cytometry as previously described [16,17]. As demonstrated by Marsh et al. [16], clone 48 antibody provided brighter staining compared

Table II Comparison of patients with XIAP deficiency

	Marsh R et al. [12]	Pachlopnik Schmid J et al. [13]	Our study
Number of patients	10	30	9
HLH	9 (90%)	22/29 (76%)	6/9 (67%)
Recurrent HLH	6 (60%)	11/18 (61%)	5/6 (83%)
EBV-associated HLH	3 (30%)	16/19 (84%)	4/6 (67%)
Splenomegaly	9 (90%)	19/21 (90%)	4/8 (50%)
Hypogammaglobulinemia	2 (20%)	8/24 (33%)	2/8 (25%)
Lymphoma	0	0	0
Colitis	0	5 (17%)	2 (22%)

to clone 2F1 antibody. In patients 5, 6.1, and 6.2, XIAP protein expression was normal when using clone 48 antibody but decreased when using clone 2F1 antibody. Western blot analysis showed XIAP expression in patients 3.1, 5 and 6.2, and using clone 48 antibody, we found a discrepancy between flow cytometry and Western blot. Flow cytometric diagnosis may thus result in false positive results, and the gene sequencing of *XIAP* should be performed even when the patient shows normal XIAP expression levels.

All of the mothers examined in this study except for one were carriers of *XIAP* mutations. Analysis of XIAP expression in the mothers of patients 1, 3.1, and 3.2 revealed a bimodal expression pattern of XIAP in lymphocytes with cellular skewing towards expression of the wild-type XIAP allele as previously demonstrated [16]. However, the mother of patients 2.1, 6.1, and 6.2 demonstrated a normal expression pattern, possibly resulting from an extremely skewed pattern of X chromosome inactivation as shown in XIAP deficiency and other primary immunodeficiencies, and de novo mutations in *XIAP* are also observed [16,31]. The mother of patients 6.1 and 6.2 might have a germline mosaicism for the mutation, resulting in normal XIAP protein expression.

iNKT cells represent a specialized T lymphocyte subpopulation with unique features distinct from conventional T cells [32,33]. Human iNKT cells express an invariant TCR that recognizes self and microbacterial glycosphingolipid antigens presented by the major histocompatibility complex class I-like molecule CD1d [28]. The first series of XIAP-deficient patients showed decreased iNKT cell counts similar to SAP deficiency [7]. However, Xiap-deficient mice have normal numbers of iNKT cells and did not show an abnormal response to apoptotic stimuli [34]. Marsh et al. [23] reported a cohort of XIAP-deficient patients with normal numbers of iNKT cells, indicating that XIAP-deficient patients differ from SAP-deficient patients in this respect. In our cohort, we observed significantly decreased iNKT cell numbers in XIAP-deficient patients compared to healthy controls. However, we could not identify a correlation between the number of iNKT cells and the clinical disease

features. Flow cytometric evaluation of iNKT cell counts can allow for the discrimination of XLP and other primary immunodeficiency diseases because patients may have normal XIAP protein expression in their lymphocytes.

CTLs kill their targets by one of two mechanisms: granule- or receptor-mediated apoptosis [35]. A recent study showed that the main pathway of cytotoxicity mediated by alloantigen-specific human CD4⁺ and CD8⁺ T cells is granule exocytosis and not the FAS/FAS ligand system [18]. Granzyme B is a major effector molecule of granule-mediated killing that rapidly induces cell death after entering the cytoplasm of the target cell [36]. The enzymatic activity of granzyme B is key to its ability to induce cell death. The executioner caspase-3 has been shown to be proteolytically processed and activated by granzyme B [37]. Although XIAP possesses an inhibitory effect for caspases, it is important to study the cytotoxic activities of CTLs in XIAP deficiency. Furthermore, many studies have indicated that some subtypes of patients with familial HLH show a deficiency in their cytotoxic activities [20,38]. To further investigate the function of antigen-specific CTLs, we studied CD8⁺ alloantigen-specific CTL analysis among three XIAP-deficient patients. XIAP-deficient patients showed a normal level of cytotoxic activity, suggesting that XIAP might not play an important role in the cytotoxic responses of CD8⁺ T cells as was previously suggested based on the normal NK cell-mediated cytotoxicity found in XIAP-deficient patients [7,12].

In this study, we have described nine Japanese patients with XIAP deficiency with clinical characteristics similar to those of patients in Europe and USA [12,13].

Acknowledgments This study was supported by Grant-in-Aids for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology (H. Kanegane and T. Miyawaki) and grants from the Ministry of Health, Labour, Welfare of Japan (T. Miyawaki), the XLP Reserch Trust (S. Latour) and Agence Nationale pour la Recherche (ANR-08-MIEN-012-01) and an Erasmus MC Fellowship (M.C. van Zelm). We thank Ms. Chikako Sakai and Mr. Hitoshi Moriuchi for their excellent technical assistance. We are also grateful for the support, cooperation, and trust of the patients and their families.

References

- Sumegi J, Huang D, Lanyi A, Davis JD, Seemayer TA, Maeda A, et al. Correlation of mutations of the SH2D1A gene and Epstein–Barr virus infection with clinical phenotype and outcome in X-linked lymphoproliferative disease. *Blood*. 2000;96:3118–25.
- Seemayer TA, Gross TG, Egeler RM, Pirruccello SJ, Davis JR, Kelly CM, et al. X-linked lymphoproliferative disease: twenty-five years after the discovery. *Pediatr Res*. 1995;38:471–8.
- Sayos J, Wu C, Morra M, Wang N, Zhang X, Allen D, et al. The X-linked lymphoproliferative-disease gene product SAP regulates signals induced through the co-receptor SLAM. *Nature*. 1998;395:462–9.
- Coffey AJ, Brooksbank RA, Brandau O, Oohashi T, Howell GR, Bye JM, et al. Host response to EBV infection in X-linked lymphoproliferative disease results from mutations in an SH2-domain encoding gene. *Nat Genet*. 1998;20:129–35.
- Nichols KE, Harkin DP, Levitz S, Krainer M, Kolquist KA, Genovese C, et al. Inactivating mutations in an SH2 domain-encoding gene in X-linked lymphoproliferative syndrome. *Proc Natl Acad Sci USA*. 1998;95:13765–70.
- Gilmour KC, Cranston T, Jones A, Davies EG, Goldblatt D, Thrasher A, et al. Diagnosis of X-linked lymphoproliferative disease by analysis of SLAM-associated protein expression. *Eur J Immunol*. 2000;30:1691–7.
- Rigaud S, Fondanèche MC, Lambert N, Pasquier B, Mateo V, Soulas P, et al. XIAP deficiency in humans causes an X-linked lymphoproliferative syndrome. *Nature*. 2006;444:110–4.
- Uren AG, Pakusch M, Hawkins CJ, Puls KL, Vaux DL. Cloning and expression of apoptosis inhibitory protein homologs that function to inhibit apoptosis and/or bind tumor necrosis factor receptor-associated factors. *Proc Natl Acad Sci USA*. 1996;93:4974–8.
- Liston P, Roy N, Tamai K, Lefebvre C, Baird S, Cherton-Horvat G, et al. Suppression of apoptosis in mammalian cells by NAIP and a related family of IAP genes. *Nature*. 1996;379:349–53.
- Duckett CS, Nava VE, Gedrich RW, Clem RJ, van Dongen JL, Gilfillan MC, et al. A conserved family of cellular genes related to the baculovirus iap gene and encoding apoptosis inhibitors. *EMBO J*. 1996;15:2685–94.
- Galbán S, Duckett CS. XIAP as a ubiquitin ligase in cellular signaling. *Cell Death Differ*. 2010;17:54–60.
- Marsh RA, Madden L, Kitchen BJ, Mody R, McClimon B, Jordan MB, et al. XIAP deficiency: a unique primary immunodeficiency best classified as X-linked familial hemophagocytic lymphohistiocytosis and not as X-linked lymphoproliferative disease. *Blood*. 2010;7:1079–82.
- Pachlopnik Schmid J, Canioni D, Moshous D, Touzot F, Mahlaoui N, Hauck F, et al. Clinical similarities and differences of patients with X-linked lymphoproliferative syndrome type 1 (XLP-1/SAP-deficiency) versus type 2 (XLP-2/XIAP-deficiency). *Blood*. 2011;117:1522–9.
- Zhao M, Kanegane H, Ouchi K, Imamura T, Latour S, Miyawaki T. A novel XIAP mutation in a Japanese boy with recurrent pancytopenia and splenomegaly. *Haematologica*. 2010;95:688–9.
- Filipovich AH, Zhang K, Snow AL, Marsh RA. X-linked lymphoproliferative syndromes: brothers or distant cousins? *Blood*. 2010;116:3398–408.
- Marsh RA, Villanueva J, Zhang K, Snow AL, Su HC, Madden L, et al. A rapid flow cytometric screening test for X-linked lymphoproliferative disease due to XIAP deficiency. *Cytometry B Clin Cytom*. 2009;76:334–44.
- Marsh RA, Bleesing JJ, Filipovich AH. Using flow cytometry to screen patients for X-linked lymphoproliferative disease due to SAP deficiency and XIAP deficiency. *J Immunol Methods*. 2010;362:1–9.
- Yasukawa M, Ohminami H, Arai J, Kasahara Y, Ishida Y, Fujita S. Granule exocytosis, and not the fas/fas ligand system, is the main pathway of cytotoxicity mediated by alloantigen-specific CD4(+) as well as CD8(+) cytotoxic T lymphocytes in humans. *Blood*. 2000;95:2352–5.
- Yanai F, Ishii E, Kojima K, Hasegawa A, Azuma T, Hirose S, et al. Essential roles of perforin in antigen-specific cytotoxicity mediated by human CD4+ T lymphocytes: analysis using the combination of hereditary perforin-deficient effector cells and Fas-deficient target cells. *J Immunol*. 2003;170:2205–13.
- Ishii E, Ueda I, Shirakawa R, Yamamoto K, Horiuchi H, Ohga S, et al. Genetic subtypes of familial hemophagocytic lymphohistiocytosis: correlations with clinical features and cytotoxic T lymphocyte/natural killer cell functions. *Blood*. 2005;105:3442–8.
- Nichols KE, Hom J, Gong SY, Ganguly A, Ma CS, Cannons JL, et al. Regulation of NKT cell development by SAP, the protein defective in XLP. *Nat Med*. 2005;11:340–5.
- Pasquier B, Yin L, Fondanèche MC, Relouzat F, Bloch-Queyrat C, Lambert N, et al. Defective NKT cell development in mice and humans lacking the adapter SAP, the X-linked lymphoproliferative syndrome gene product. *J Exp Med*. 2005;201:695–701.
- Marsh RA, Villanueva J, Kim MO, Zhang K, Marmer D, Risma KA, et al. Patients with X-linked lymphoproliferative disease due to *BIRC4* mutation have normal invariant natural killer T-cell populations. *Clin Immunol*. 2009;132:116–23.
- Puck JM, Pepper AE, Bedard PM, Laframboise R. Female germ line mosaicism as the origin of a unique IL-2 receptor gamma-chain mutation causing X-linked severe combined immunodeficiency. *J Clin Invest*. 1995;95:895–9.
- O'Marcaigh A, Puck JM, Pepper AE, Santes KD, Cowan MJ. Maternal mosaicism for a novel interleukin-2 receptor gamma-chain mutation causing X-linked severe combined immunodeficiency in a Navajo kindred. *J Clin Immunol*. 1997;17:29–33.
- Sakamoto M, Kanegane H, Fujii H, Tsukada S, Miyawaki T, Shinomiya N. Maternal germinal mosaicism of X-linked agammaglobulinemia. *Am J Med Genet*. 2001;99:234–7.
- Harlin H, Reffey SB, Duckett CS, Lindsten T, Thompson CB. Characterization of XIAP-deficient mice. *Mol Cell Biol*. 2001;21:3604–8.
- Latour S. Natural killer T cells and X-linked lymphoproliferative syndrome. *Curr Opin Allergy Clin Immunol*. 2007;7:510–4.
- Schimmer AD, Dalili S, Batey RA, Riedl SJ. Targeting XIAP for the treatment of malignancy. *Cell Death Differ*. 2006;13:179–88.
- Oliveira JB, Notarangelo LD, Fleisher TA. Applications of flow cytometry for the study of primary immune deficiencies. *Curr Opin Allergy Clin Immunol*. 2008;8:499–509.
- Kanegane H, Futatani T, Wang Y, Nomura K, Shinozaki K, Matsukura H, et al. Clinical and mutational characteristics of X-linked agammaglobulinemia and its carrier identified by flow cytometric assessment combined with genetic analysis. *J Allergy Clin Immunol*. 2001;108:1012–20.
- Godfrey DI, Berzins SP. Control points in NKT-cell development. *Nat Rev Immunol*. 2007;7:505–18.
- Bendelac A, Savage PB, Teyton L. The biology of NKT cells. *Annu Rev Immunol*. 2007;25:297–336.
- Bauler LD, Duckett CS, O'Riordan MX. XIAP regulates cytosol-specific immunity to *Listeria* infection. *PLoS Pathog*. 2008;4:e1000142.
- Hersperger AR, Makedonas G, Betts MR. Flow cytometric detection of perforin upregulation in human CD8 T cells. *Cytometry A*. 2008;73:1050–7.
- Motyka B, Korbitt G, Pinkoski MJ, Heibein JA, Caputo A, Hobman M, et al. Mannose 6-phosphate/insulin-like growth

- factor II receptor is a death receptor for granzyme B during cytotoxic T cell-induced apoptosis. *Cell*. 2000;103:491–500.
37. Martin SJ, Amarante-Mendes GP, Shi L, Chuang TH, Casiano CA, O'Brien GA, et al. The cytotoxic cell protease granzyme B initiates apoptosis in a cell-free system by proteolytic processing and activation of the ICE/CED-3 family protease, CPP32, via a novel two-step mechanism. *EMBO J*. 1996;15:2407–16.
38. zur Stadt U, Rohr J, Seifert W, Koch F, Grieve S, Pagel J, et al. Familial hemophagocytic lymphohistiocytosis type 5 (FHL-5) is caused by mutations in Munc18-2 and impaired binding to syntaxin 11. *Am J Hum Genet*. 2009;85:482–92.

Nationwide Survey of Patients with Primary Immunodeficiency Diseases in Japan

Masataka Ishimura · Hidetoshi Takada · Takehiko Doi · Kousuke Imai ·
Yoji Sasahara · Hirokazu Kanegane · Ryuta Nishikomori · Tomohiro Morio ·
Toshio Heike · Masao Kobayashi · Tadashi Ariga · Shigeru Tsuchiya ·
Shigeaki Nonoyama · Toshio Miyawaki · Toshiro Hara

Received: 7 August 2011 / Accepted: 11 September 2011 / Published online: 29 September 2011
© Springer Science+Business Media, LLC 2011

Abstract To determine the prevalence and clinical characteristics of patients with in Japan, we conducted a nationwide survey of primary immunodeficiency disease (PID) patients for the first time in 30 years. Questionnaires were sent to 1,224 pediatric departments and 1,670 internal medicine departments of Japanese hospitals. A total of 1,240 patients were registered. The estimated number of patients with PID was 2,900 with a prevalence of 2.3 per 100,000 people and homogenous regional distribution in Japan. The male-to-female ratio was 2.3:1 with a median age of 12.8 years. Adolescents or adults constituted 42.8% of the patients. A number of 25 (2.7%) and 78 (8.5%) patients developed malignant disorders and immune-related diseases, respectively, as complications of primary immunodeficiency disease. Close monitoring and appropriate management for these complications in addition to prevention of infectious diseases is important for improving the quality of life of PID patients.

Keywords Primary immunodeficiency disease · epidemiology · nationwide survey · Japan

Abbreviations

APECED	Autoimmune polyendocrinopathy with candidiasis and ectodermal dystrophy
BTK	Bruton's tyrosine kinase
CGD	Chronic granulomatous disease
CID	Combined T and B cell immunodeficiency
CVID	Common variable immunodeficiency disease
FMF	Familial Mediterranean fever
IPEX	Immune dysregulation polyendocrinopathy enteropathy X-linked
NEMO	Nuclear factor kappa B essential modulator
PID	Primary immunodeficiency disease
SIgAD	Selective IgA deficiency
SLE	Systemic lupus erythematosus

M. Ishimura (✉) · H. Takada · T. Doi · T. Hara
Department of Pediatrics, Graduate School of Medical Sciences,
Kyushu University,
3-1-1 Maidashi, Higashi-ku,
Fukuoka 812-8582, Japan
e-mail: ischii@pediatr.med.kyushu-u.ac.jp

K. Imai · S. Nonoyama
Department of Pediatrics, National Defense Medical College,
Tokorozawa, Japan

Y. Sasahara · S. Tsuchiya
Department of Pediatrics, Tohoku University School of Medicine,
Sendai, Japan

H. Kanegane · T. Miyawaki
Department of Pediatrics, Graduate School of Medicine
and Pharmaceutical Science, University of Toyama,
Toyama, Japan

R. Nishikomori · T. Heike
Department of Pediatrics,
Kyoto University Graduate School of Medicine,
Kyoto, Japan

T. Morio
Department of Pediatrics,
Tokyo Medical and Dental University Graduate School,
Bunkyo-ku, Tokyo, Japan

M. Kobayashi
Department of Pediatrics,
Hiroshima University Graduate School of Biomedical Sciences,
Hiroshima, Japan

T. Ariga
Department of Pediatrics, Graduate School of Medicine,
Hokkaido University,
Sapporo, Japan