

CT, with normal urinalysis values and normal renal function. One other patient (under hemodialysis) was diagnosed as having IgG4-related sclerosing cholangitis and end-stage renal disease, with evidence of a mass lesion in the right kidney by nonenhanced CT. Renal biopsy was not performed in these six patients. Another patient showed a right renal mass lesion and a high serum IgG4 level, with normal urinalysis values and mild renal dysfunction. As the tissue obtained by CT-guided renal biopsy of the mass lesion did not include any renal parenchyma, even though marked lymphoplasmacytic infiltration was observed in the adipose tissues, we excluded this patient.

In the 23 patients diagnosed as having renal parenchymal lesions associated with IgG4-related disease, we retrospectively examined the clinical features, data from laboratory and imaging studies, and the clinical response to treatment. Whole-body CT imaging was evaluated in all patients (contrast-enhanced CT in 21 patients). Gallium citrate scintigraphy was performed in 19 of the 23 patients, before therapy in 17 and during therapy in 2. The diagnosis of AIP was made in accordance with the 2006 Japan Pancreas Society revised criteria.⁴² Sialadenitis and dacryadenitis were diagnosed on the basis of physical findings, and the results of imaging studies (CT and gallium citrate scintigraphy) and/or biopsy.

The study was approved by the review board of the Nagaoka Red Cross Hospital and the boards of the various collaborating institutions. All data and samples from patients were collected with their informed consent, and the research was conducted in compliance with the principles of the Declaration of Helsinki.

Renal pathology

Renal pathological examination was conducted at the request of the attending physicians because of urinary abnormalities, renal dysfunction, and/or radiological abnormalities. Renal tissues were obtained by nondirected medical needle biopsy in 21 patients, by open biopsy of the mass lesion in 1 patient (no. 8), and by autopsy in 1 patient (no. 3), who died of lung cancer. All renal tissue specimens were examined by light microscopy. Direct immunofluorescence studies were conducted in 17 patients, and we evaluated 14 specimens because 3 were considered inadequate. For routine light microscopy studies, renal biopsy specimens were fixed in formalin or alcohol-Bouin, embedded in paraffin, and stained with hematoxylin and eosin, periodic acid-Schiff, periodic acid-methenamine silver, and Masson's trichrome. For immunostaining, formalin-fixed, paraffin-embedded biopsy specimens were cut into 3- μ m thick sections, and the sections were immunostained using anti-IgG antibody (Dako, Glostrup, Denmark) and mouse monoclonal antibody against human IgG4 (Zymed Laboratory, San Francisco, CA, USA, or The Binding Site, Birmingham, UK). For direct immunofluorescence studies, tissues were snap frozen, and sections were treated with fluorescein isothiocyanate-conjugated rabbit anti-human IgG, IgA, IgM, C1q, C3c, or fibrinogen (Dako, Carpinteria, CA, USA).

DISCLOSURE

All the authors declared no competing interests.

ACKNOWLEDGMENTS

We thank the members of the IgG4 + MOLPS/Mikulicz's Disease Society of Japan who participated in this study. This work was supported by Health and Labour Sciences Research Grants for the Study of Intractable Diseases from the Ministry of Health, Labor and Welfare, Japan.

REFERENCES

- Kamisawa T, Okamoto A. IgG4-related sclerosing disease. *World J Gastroenterol* 2008; **14**: 3948–3955.
- Yamamoto M, Takahashi H, Ohara M et al. A new conceptualization for Mikulicz's disease as an IgG4-related plasmacytic disease. *Mod Rheumatol* 2006; **16**: 335–340.
- Masaki Y, Dong L, Kurose N et al. Proposal for a new clinical entity, IgG4-positive multi-organ lymphoproliferative syndrome: analysis of 64 cases of IgG4-related disorders. *Ann Rheum Dis* 2009; **68**: 1310–1315.
- Hamano H, Kawa S, Horiuchi A et al. High serum IgG4 concentrations in patients with sclerosing pancreatitis. *N Engl J Med* 2001; **344**: 732–738.
- Zen Y, Fujii T, Harada K et al. Th2 and regulatory immune reaction are increased in immunoglobulin G4-related sclerosing pancreatitis and cholangitis. *Hepatology* 2007; **45**: 1538–1546.
- Kitagawa S, Zen Y, Harada K et al. Abundant IgG4-positive plasma cell infiltration characterizes chronic sclerosing sialadenitis (Kuttner's tumor). *Am J Surg Pathol* 2005; **29**: 783–791.
- Hamano H, Kawa S, Ochi Y et al. Hydronephrosis associated with retroperitoneal fibrosis and sclerosing pancreatitis. *Lancet* 2002; **359**: 1403–1404.
- Zen Y, Inoue D, Kitao A et al. IgG4-related lung and pleural disease: a clinicopathologic study of 21 cases. *Am J Surg Pathol* 2009; **33**: 1886–1893.
- Kasashima S, Zen Y, Kawashima A et al. A new clinicopathological entity of IgG4-related inflammatory abdominal aortic aneurysm. *J Vasc Surg* 2009; **49**: 1264–1271.
- Saida Y, Homma N, Hama H et al. A case of IgG4-related tubulointerstitial nephritis showing the progression of renal dysfunction after a cure for autoimmune pancreatitis. *Jpn J Nephrol* 2010; **52**: 73–79 (in Japanese).
- Takahashi N, Kawashima A, Fletcher JG et al. Renal involvement in patients with autoimmune pancreatitis: CT and MR imaging findings. *Radiology* 2007; **242**: 791–801.
- Uchiyama-Tanaka Y, Mori Y, Kimura T et al. Acute tubulointerstitial nephritis associated with autoimmune-related pancreatitis. *Am J Kidney Dis* 2004; **43**: 18–25.
- Takeda S, Haratake J, Kasai T et al. IgG4-associated idiopathic tubulointerstitial nephritis complicating autoimmune pancreatitis. *Nephrol Dial Transplant* 2004; **19**: 474–476.
- Saeki T, Saito A, Yamazaki H et al. Tubulointerstitial nephritis associated with IgG4-related systemic disease. *Clin Exp Nephrol* 2007; **11**: 168–173.
- Watson SJ, Jenkins DA, Bellamy CO. Nephropathy in IgG4-related systemic disease. *Am J Surg Pathol* 2006; **30**: 1472–1477.
- Rudmik L, Trpkov K, Nash C et al. Autoimmune pancreatitis associated with renal lesion mimicking metastatic tumours. *CMAJ* 2006; **175**: 367–369.
- Murashima M, Tomaszewski J, Glickman JD. Chronic tubulointerstitial nephritis presenting as multiple renal nodules and pancreatic insufficiency. *Am J Kidney Dis* 2007; **49**: E7–E10.
- Yoneda K, Murata K, Katayama K et al. Tubulointerstitial nephritis associated with IgG4-related autoimmune disease. *Am J Kidney Dis* 2007; **50**: 455–462.
- Okazaki K, Chiba T. Autoimmune related pancreatitis. *Gut* 2002; **51**: 1–4.
- Park DH, Kim HM, Chari ST. Recent advances in autoimmune pancreatitis. *Gut* 2009; **58**: 1680–1689.
- Otsuki M, Chung JB, Okazaki K et al. Asian diagnostic criteria for autoimmune pancreatitis: consensus of the Japan-Korea symposium on autoimmune pancreatitis. *J Gastroenterol* 2008; **43**: 403–408.
- Okazaki K, Uchida K, Matsushita M et al. Autoimmune pancreatitis. *Intern Med* 2005; **44**: 1215–1223.
- Naitoh I, Nakazawa T, Ohara H et al. Clinical significance of extrapancreatic lesions in autoimmune pancreatitis. *Pancreas* 2010; **39**: e1–e5.
- Kamisawa T, Anjiki H, Egawa N et al. Allergic manifestations in autoimmune pancreatitis. *Eur J Gastroenterol Hepatol* 2009; **21**: 1136–1139.
- Muraki T, Hamano H, Ochi Y et al. Autoimmune pancreatitis and complement activation system. *Pancreas* 2006; **32**: 16–21.
- Stone JH, Khosroshahi A, Hilgenberg A et al. IgG4-related systemic disease and lymphoplasmacytic aortitis. *Arthritis Rheum* 2009; **60**: 3139–3145.
- Stone JH, Caruso PA, Deshpande V. Case 24–2009: a 26-year-old woman with painful swelling of the neck. *N Engl J Med* 2009; **361**: 511–518.
- Cornell LD, Chicano SL, Deshpande V et al. Pseudotumors due to IgG4 immune-complex tubulointerstitial nephritis associated with autoimmune pancreatocentric disease. *Am J Surg Pathol* 2007; **31**: 1586–1597.

29. Katano K, Hayatsu Y, Matsuda T *et al*. Endocapillary proliferative glomerulonephritis with crescent formation and concurrent tubulointerstitial nephritis complicating retroperitoneal fibrosis with a high serum level of IgG4. *Clin Nephrol* 2007; **68**: 308–314.
30. Morimoto J, Hasegawa Y, Fukushima H *et al*. Membranoproliferative glomerulonephritis-like glomerular disease and concurrent tubulointerstitial nephritis complicating IgG4-related autoimmune pancreatitis. *Intern Med* 2009; **48**: 157–162.
31. Kuroki A, Shibata T, Honda H *et al*. Glomerular and serum IgG subclasses in diffuse proliferative lupus nephritis, membranous lupus nephritis, and idiopathic membranous nephropathy. *Intern Med* 2002; **41**: 936–942.
32. Kuroki A, Iyoda M, Shibata T *et al*. Th2 cytokines increase and stimulate B cells to produce IgG4 in idiopathic membranous nephropathy. *Kidney Intern* 2005; **68**: 302–310.
33. Saeki T, Nishi S, Ito T *et al*. Renal lesions in IgG4-related systemic disease. *Intern Med* 2007; **46**: 1365–1372.
34. Saeki T, Saito A, Hiura T *et al*. Lymphoplasmacytic infiltration of multiple organs with immunoreactivity for IgG4: IgG4-related systemic disease. *Intern Med* 2006; **45**: 163–167.
35. Shimoyama K, Ogawa N, Sawaki T *et al*. A case of Mikulicz's disease complicated with interstitial nephritis successfully treated by high-dose corticosteroid. *Mod Rheumatol* 2006; **16**: 176–182.
36. Nakamura H, Wada H, Origuchi T *et al*. A case of IgG4-related autoimmune disease with multiple organ involvement. *Scand J Rheumatol* 2006; **35**: 69–71.
37. Yamamoto M, Takahashi H, Hasebe K *et al*. The analysis of interleukin-6 in patients with systemic IgG4-related plasmacytic syndrome—expansion of SIPS to the territory of Castleman's disease. *Rheumatology* 2009; **48**: 860–862.
38. Nakada S, Yagi K, Minamizawa K *et al*. A case of Mikulicz's disease with high serum IgG4 concentrations and tubulointerstitial nephritis. *J Chubu Rheum Assoc* 2006; **37**: 102–103. (in Japanese).
39. Saeki T, Imai N, Ito T *et al*. Membranous nephropathy associated with IgG4-related systemic disease and without autoimmune pancreatitis. *Clin Nephrol* 2009; **71**: 173–178.
40. Takamura N, Eguchi K, Migita K *et al*. A male of primary Sjögren's syndrome with interstitial pneumonitis and interstitial tubulo-nephritis in the absence of dry eye and dry mouth: parotid gland MRI is useful diagnostic method for Sjögren's syndrome. *Jpn J Clin Immun* 1995; **18**: 123–132. (in Japanese).
41. Saeki T, Ito T, Yamazaki H *et al*. Hypocomplementemia of unknown etiology: an opportunity to find cases of IgG4-positive multi-organ lymphoproliferative syndrome. *Rheumatol Int*; e-pub ahead of print 18 April 2009.
42. Okazaki K, Kawa S, Kamisawa T *et al*. Clinical diagnostic criteria of autoimmune pancreatitis: revised proposal. *J Gastroenterol* 2006; **41**: 626–631.

IgG4-Related Pleural Disease Diagnosed by a Re-Evaluation of Chronic Bilateral Pleuritis in a Patient Who Experienced Occasional Acute Left Bacterial Pleuritis

Hiroshi Yamamoto¹, Toshiro Suzuki¹, Masanori Yasuo¹, Oriie Kobayashi¹, Kenji Tsushima¹, Michiko Ito¹, Kazuhisa Urushihata¹, Yoshitaka Yamazaki¹, Masayuki Hanaoka¹, Tomonobu Koizumi¹, Takeshi Uehara², Satoshi Kawakami³, Hideaki Hamano⁴, Shige-yuki Kawa⁵ and Keishi Kubo¹

Abstract

A 78-year-old man with cryptogenic chronic bilateral lymphoplasmacytic pleuritis, diagnosed based on left parietal pleural biopsy specimens obtained by pleuroscopy, developed acute left bacterial pleuritis. The left pleural effusion was neutrophil dominant, however, the right pleural effusion showed lymphoplasmacytic infiltration. Laboratory examinations revealed that his serum IgG4 concentration was increased, with a higher level of IgG4 in the right pleural effusion. Re-evaluation of the previous biopsy specimens using an immunostaining method revealed numerous IgG4-positive plasma cell infiltrations with IgG4-positive/IgG-positive plasma cells at 85.4%. Accordingly, the new diagnosis of this patient was considered to be chronic bilateral IgG4-related pleuritis.

Key words: pleuritis, lymphoplasmacytic, IgG4, pleuroscopy

(Intern Med 50: 893-897, 2011)

(DOI: 10.2169/internalmedicine.50.4726)

Introduction

Immunoglobulin (Ig) G4-related lung and plural disease has been receiving increasing attention (1). Recently, Shrestha et al reported 6 patients who were diagnosed with lung involvement in IgG4-related autoimmune pancreatitis (AIP) (2). Interestingly, all 6 patients had fibrinous pleuritis confirmed histopathologically, and one of them showed a right pleural effusion radiologically (2). However, the clinical features of IgG4-related lung and pleural diseases are not fully understood. We herein report a patient who had been diagnosed with chronic bilateral lymphoplasmacytic pleuritis who developed acute left bacterial pleuritis, which was identified to be IgG4-related pleural disease.

Case Report

A 78-year-old man was admitted to our hospital (Shinshu University School of Medicine, Matsumoto, Japan) complaining of general fatigue and fever without pain and dyspnea in July 2008. He had no family history of pancreatic disease, collagen disease, or autoimmune disease. He had a history of gall bladder stones and underwent a cholecystectomy at 68 years of age.

At 74 years of age, he developed a bilateral painless pleural effusion without associated fever. His bilateral pleural effusions were both exudative and mononuclear cell dominant. Fluorodeoxyglucose positron emission tomography (FDG-PET) revealed no abnormal uptake. In 2004, a pleuroscopic

¹The First Department of Internal Medicine, Shinshu University School of Medicine, Japan, ²Department of Pathology, Shinshu University School of Medicine, Japan, ³Department of Radiology, Shinshu University School of Medicine, Japan, ⁴The Second Department of Internal Medicine, Shinshu University School of Medicine, Japan and ⁵Center for Health, Safety and Environmental Management, Shinshu University School of Medicine, Japan

Received for publication October 22, 2010; Accepted for publication January 6, 2011

Correspondence to Dr. Hiroshi Yamamoto, yama5252@shinshu-u.ac.jp




Figure 1. A chest X-ray on admission showed bilateral pleural effusion as described without significant infiltration.



Figure 2. A chest X-ray at 7 days after admission revealed a rapid increase in the left pleural effusion.

parietal pleural biopsy in the left thorax was performed and lymphoplasmacytic pleuritis of unknown etiology was diagnosed. Bacterial culture and polymerase chain reaction (PCR) analysis of the pleural effusion for *Mycobacterium tuberculosis*, *avium*, and *intracellulare* DNA were all negative. Adenosine deaminase (ADA) concentrations in the pleural effusion were measured three times from both sides in 2004. Because the values of ADA in the pleural effusion were 34.1, 36.5, and 46.7 (U/L), we could not conclusively rule out the possibility of tuberculous pleuritis (3). Although the patient had received oral antituberculosis agents for 6 months in 2005, the bilateral pleural effusion did not decrease at all. We therefore believed that his chronic bilateral pleuritis was not tuberculous pleuritis. He was placed under clinical observation with diuretics, but the bilateral pleural effusion still did not show any remarkable change until the time of the present administration.

Laboratory examinations showed the following values (normal range): peripheral white blood cell count, 13,110/ μ L; C-reactive protein, 25.3 mg/dL; total protein (TP), 7.4 g/dL; albumin, 3.3 g/dL; lactate dehydrogenase (LD), 175 IU/L (<220); and IgG, 1,604 mg/dL (<1,700). The serum concentrations of the IgG subclasses were as follows: IgG1, 791 mg/dL (<1,080); IgG2, 777 mg/dL (<931); IgG3, 84




Figure 3. A chest CT at 7 days after admission showed bilateral pleural effusion and mild pleural thickening. The left pleural effusion was more extensive with small air densities (arrows).

mg/dL (<121); and IgG4, 483 mg/dL (<108), and IgG4/IgG 30.1%. The serum autoantibodies were all negative, including antinuclear antibody (ANA), rheumatoid factor (RF), anti-double-stranded DNA antibody, anti-RNP antibody, anti-Sjögren's syndrome (SS)-A antibody, SS-B antibody, and anti-neutrophil cytoplasmic autoantibody (ANCA). The serum level of the soluble interleukin-2 receptor was 907 U/mL (<421) and angiotensin-converting enzyme (ACE) was 12.9 U/L (<25). His electrophoretogram of a blood sample did not show any abnormal bands. His thyroid function was normal and the level of brain natriuretic peptide (BNP) in the blood was normal. Proteinuria was not observed.

A chest X-ray showed bilateral pleural effusion as had been noted prior to the present admission without infiltration in the lung fields (Fig. 1). Cytological analysis was performed for right pleural effusion and the fluid on the day of admission. However, seven days after admission, his chest X-ray revealed a rapid increase of left pleural effusion (Fig. 2). The chest CT showed bilateral pleural effusion and mild pleural thickening without hilar lymph node swelling, and the amount of the left pleural effusion was higher than that in the right side, with small air densities in the effusion (Fig. 3). In order to determine the reason for the rapid change in the effusion in the left thorax, we performed a cytological examination of the left pleural effusion percutaneously. Both left and right pleural effusions were exudative, however, the cell populations were entirely different. The cell fraction of the right pleural effusion was mononuclear cell (mainly lymphocyte and plasma cell) dominant (Fig. 4a) and the left pleural effusion was polymorphonuclear cell (mainly neutrophil) dominant (Fig. 4b). The concentrations of several markers in the right and left pleural effusion were as follows: pH, 8.0 and 7.5; TP, 6.1, 4.9 (g/dL); LDH, 191, 2,297 (IU/L); glucose, 116, 2 (mg/dL); and IgG, 2,515, 1,491 (mg/dL), respectively. Although the cultures of bacteria and acid-fast bacilli in both the left and right pleural effusions were negative, we diagnosed acute bacterial pleuritis for the left thorax. In addition to administration of antiimi-




Figure 4. Cytology of the right pleural effusion showed lymphoplasmacytic infiltration (a) and left pleural effusion revealed marked neutrophil infiltration (b). (Giemsa stain $\times 100$)




Figure 5. The left parietal pleural biopsy specimens showed lymphoplasmacytic infiltration with mild fibrosis in Hematoxylin and Eosin staining (a). An immunohistochemical examination of the specimen revealed dense infiltration of IgG4-positive plasma cells (b). ($\times 400$)

crobial agents, pleural lavage was performed for 5 days via a catheter. As a result, the left pleural effusion was controlled and the patient was discharged 27 days after admission. At the time of discharge, his bilateral pleural effusion was the same as that observed on admission.

In order to determine the etiology of his chronic bilateral pleuritis, we re-evaluated the previously obtained parietal pleural biopsy specimens from the left thorax using an immunostaining method. The numbers of both IgG4-positive and IgG-positive cells were counted in regions of the highest density and averaged in the three most cellular high-power fields (HPF) (2). The specimens of the left parietal pleural tissue showed lymphoplasmacytic infiltration with mild fibrosis and no granulomas were seen in Hematoxylin

and Eosin (HE) staining (Fig. 5a). Interestingly, immunohistochemical examinations showed infiltration by numerous IgG4-positive plasma cells in the specimens (Fig. 5b). The number of IgG4-positive plasma cells per HPF was 17.6, scored as moderate (10-30/HPF) according to Kamisawa et al. (4), and the percentage of IgG4-positive to IgG-positive plasma cells (IgG4+/IgG+) was 85.4%. There was a mixture of kappa-positive and lambda-positive cells in the specimens. Furthermore, the concentrations of IgG and IgG4 in the right pleural effusion were higher than those in the serum (the IgG subtypes in the right pleural effusion were as follows: IgG1, 758 mg/dL; IgG2, 651 mg/dL; IgG3, 69 mg/dL; and IgG4, 590 mg/dL). Therefore, the chronic lymphoplasmacytic pleuritis in this patient was diagnosed as




Figure 6. A chest X-ray in 2010 of recent data showing that there was not significant increase in the bilateral pleural effusion.

IgG4-related pleural disease.

One year later, he developed colon cancer, and underwent a curative resection in 2009. The FDG-PET before the operation showed an abnormal accumulation of FDG in the colon tumor and an ill-defined nodule of the left lung, which was suspected to be an inflammatory lesion. However, no other abnormal uptake was observed, even in the pleura. Furthermore, the thoraco-abdominal CT showed no remarkable abnormality except for bilateral pleural effusion and colon cancer.

Two years after the discharge, the patient treated with diuretics had no respiratory symptoms and his bilateral pleural effusion had not changed significantly (Fig. 6). His serum concentrations of the IgG subclasses and interleukin-6 (IL-6) in 2010 were as follows: IgG1 1280 mg/dL; IgG2 1000 mg/dL; IgG3 136 mg/dL; and IgG4 754 mg/dL (IgG4/IgG 34.1%), and 1.76 pg/mL (<2.41).

Discussion

We herein described an elderly man with bilateral chronic lymphoplasmacytic pleuritis who developed acute left bacterial pleuritis and finally was identified as having IgG4-related disease 4 years after the appearance of the bilateral pleural effusion. A re-evaluation of the previously obtained biopsy specimens using an immunostaining method revealed numerous IgG4-positive plasma cells in the parietal pleura. Because IgG4-related disease was seldom reported in 2004, we did not measure the serum IgG4 concentration at that time, and thus overlooked the diagnosis of bilateral lymphoplasmacytic pleuritis with IgG4-related pleural disease in this patient.

Rigorous diagnostic criteria for IgG4-related lung and pleural diseases have not been established, however, a high serum concentration of IgG4 and marked IgG4-positive plasma cell infiltration in the biopsy specimens are accepted as the characteristic features of this disease (1, 2). The lung and pleural involvement of AIP and Mikulicz's disease were thought to meet the criteria for IgG4-related lung and pleu-

ral diseases. Mikulicz's disease is an IgG4-related systemic disease that is identified by an enlargement of the lachrymal and salivary glands, which differs substantially from Sjögren's syndrome (5). Fujinaga et al reported that the lung lesions were detected in 25 (54%) out of 46 patients with AIP who had undergone thin-slice chest CT scanning (6). Various extra-pancreatic lesions had been reported in AIP and a new clinicopathological entity of IgG4-related autoimmune disease was submitted (4). Meanwhile, Masaki et al reported that 6 (9.4%) of 64 patients with Mikulicz's disease showed interstitial lung disease and 11 (17.2%) of 64 patients had AIP (7). They proposed a new clinical entity, IgG4-positive multiorgan lymphoproliferative syndrome (IgG4+MOLPS), a syndrome characterized by hyper-IgG4 gammaglobulinemia (>135 mg/dL) and IgG4-positive plasma cell infiltration in the tissues (IgG4+/IgG+ >50%) (7). However, there are important limitations of these reports because a large proportion of the lung lesions were diagnosed by radiological findings rather than by the pathological findings of the lungs (4, 6, 7). On the other hand, Zen et al reported 21 patients with IgG4-related lung and pleural disease diagnosed on the basis of the histopathological findings of the lung and pleura (1). These patients had characteristic HE findings such as diffuse lymphoplasmacytic infiltration, irregular fibrosis, and occasional eosinophil infiltration. The immunostaining showed diffuse IgG4-positive plasma cell infiltration and a ratio of IgG4+/IgG+ greater than 30%. In addition, 9 (43%) out of the 21 patients had IgG4-related disease in other organs, such as the pancreas and submandibular glands (1).

Miyake et al reported a male patient with left pleural effusion and swelling of the submandibular glands diagnosed with Mikulicz's disease (5). Their patient had a high concentration of serum IgG4 and his biopsy specimens of the submandibular gland showed abundant IgG4-bearing mononuclear cell infiltrations. Although he had not undergone a lung or pleural biopsy, pleurocentesis was performed on the left thorax, and numerous mononuclear cells (lymphocytes and plasma cells) were found to be present in his pleural effusion. Additionally, his IgG4 concentration in the pleural effusion was higher than that in the serum (5). Similarly, the present patient showed a high serum IgG4 concentration, and his effusion of non-bacterial pleuritis was mononuclear cell (mainly lymphocyte and plasma cell) dominant, and the IgG4 concentration in the effusion was higher than that in the serum. Furthermore, the biopsy specimens of the left parietal pleura showed infiltration of numerous IgG4-positive plasma cells, in agreement with the criteria for IgG4-related lung and pleural diseases proposed by Zen et al (1) and IgG4+MOLPS (5). In the present patient, although the serum concentrations of IgG and IgG4 were high, the electrophoretogram of his blood did not show a monoclonal band. His serum autoantibodies were all negative and the serum concentrations of ACE, ANCA, and IL-6 were normal. Because the other differential diagnoses such as multiple myeloma, collagen vascular disease, sarcoidosis (8),

Wegener's granulomatosis, Castleman disease (7, 9), and other malignant diseases were not in agreement with the chronic bilateral pleuritis of the present patient pathologically, we considered the final diagnosis of the patient to be chronic bilateral IgG4-related pleuritis. IgG4-related lesions and lymphomatoid granulomatosis grade 1 (LYG-G1) lesions have been reported to be morphologically indistinguishable in the lung (1, 10). We also think that the lack of any atypical cells and the presence of IgG4-positive plasma cells may therefore be more indicative of IgG4-related disease over LYG-G1 (10).

To the best of our knowledge, there have been only three other published cases with IgG4-related lung and pleural diseases, diagnosed on the basis of pathological findings of the lung and pleura or pleural effusion, which revealed pleural effusion radiologically (2, 5, 10). The patient with Mikulicz's disease was treated with 30 mg/day prednisolone (PSL) for 14 days, followed by a tapering of the PSL dosage. As a result, the pleural effusion showed a drastic reduction (5). One patient with right pleural effusion was successfully treated with diuretics after the thoroscopic biopsy (10). Although the present patient did not receive oral corticosteroid therapy after the diagnosis, his bilateral pleural effusion did not increase and IgG4-related disease in other organs was not found by FDG-PET and CT. It was reported that FDG-PET is a sensitive tool for detecting lesions related to IgG4-related disease, and that the levels of FDG uptake revealed the activity of the lesions (11). However, there have been no reports of patients with IgG4-related pleural disease who were examined by FDG-PET. Because no abnormal accumulation of FDG in the pleura or pleural effusion was observed in the present case, it seems that the activity of IgG4-related pleural disease is relatively low. Furthermore, IgG4-related pleural disease might thus have a relatively favorable prognosis.

In conclusion, there have been some reports of IgG4-related diseases, however, the present case is rare due to the fact that this patient was diagnosed to have IgG4-related localized pleural disease based on the findings of a pleural biopsy; in addition, the patient was followed for a substantial length of time and also demonstrated a relatively favorable prognosis.

The authors state that they have no Conflict of Interest (COI).

This report was supported by Intractable Disease, the Health and Labour Sciences Research Grants from Ministry of Health, Labor and Welfare.

References

1. Zen Y, Inoue D, Kitao A, et al. IgG4-related lung and pleural disease: a clinicopathologic study of 21 cases. *Am J Surg Pathol* **33**: 1886-1893, 2009.
2. Shrestha B, Sekiguchi H, Colby TV, et al. Distinctive pulmonary histopathology with increased IgG4-positive plasma cells in patients with autoimmune pancreatitis: report of 6 and 12 cases with similar histopathology. *Am J Surg Pathol* **33**: 1450-1462, 2009.
3. Gopi A, Madhavan SM, Sharma SK, Sahn SA. Diagnosis and treatment of tuberculous pleural effusion in 2006. *Chest* **131**: 880-889, 2007.
4. Kamisawa T, Funata N, Hayashi Y, et al. A new clinicopathological entity of IgG4-related autoimmune disease. *J Gastroenterol* **38**: 982-984, 2003.
5. Miyake K, Moriyama M, Aizawa K, et al. Peripheral CD4+ T cells showing a Th2 phenotype in a patient with Mikulicz's disease associated with lymphadenopathy and pleural effusion. *Mod Rheumatol* **18**: 86-90, 2008.
6. Fujinaga Y, Kadoya M, Kawa S, et al. Characteristic findings in images of extra-pancreatic lesions associated with autoimmune pancreatitis. *Eur J Radiol* **76**: 228-238, 2010.
7. Masaki Y, Dong L, Kurose N, et al. Proposal for a new clinical entity, IgG4-positive multiorgan lymphoproliferative syndrome: analysis of 64 cases of IgG4-related disorders. *Ann Rheum Dis* **68**: 1310-1315, 2009.
8. Watarai M, Yazawa M, Yamanda K, Yamamoto H, Yamazaki Y. Pulmonary sarcoidosis with associated bloody pleurisy. *Intern Med* **41**: 1021-1023, 2002.
9. Ko SF, Ng SH, Hsieh MJ, et al. Castleman disease of the pleura: experience with eight surgically proven cases. *Ann Thorac Surg* **76**: 219-224, 2003.
10. Yamashita K, Haga H, Kobashi Y, Miyagawa-Hayashino A, Yoshizawa A, Manabe T. Lung involvement in IgG4-related lymphoplasmacytic vasculitis and interstitial fibrosis: report of 3 cases and review of the literature. *Am J Surg Pathol* **32**: 1620-1626, 2008.
11. Tanabe T, Tsushima K, Yasuo M, et al. IgG4-associated multifocal systemic fibrosis complicating sclerosing sialadenitis, hypophysitis, and retroperitoneal fibrosis, but lacking pancreatic involvement. *Intern Med* **45**: 1243-1247, 2006.

