

「新規疾患, IgG4関連多臓器リンパ増殖性疾患(IgG4+MOLPS)の確立のための研究」班

区 分	氏 名	所 属 等	職 名
研 究 代 表 者	梅原 久範	金沢医科大学大学院医科学研究科血液免疫内科学	教 授
研 究 分 担 者	三森 経世	京都大学大学院医学研究科臨床免疫学	教 授
	住田 孝之	筑波大学大学院人間総合科学研究科臨床免疫学	教 授
	坪田 一男	慶應義塾大学医学部眼科学教室	教 授
	吉野 正	岡山大学大学院病態制御学腫瘍制御学病理学	教 授
	岡崎 和一	関西医科大学内科学第三講座	教 授
	川 茂幸	信州大学健康安全センター	教 授
	竹上 勉	金沢医科大学総合医学研究所生命科学研究領域	教 授
	友杉 直久	金沢医科大学総合医学研究所先端医療研究領域	教 授
	田中 良哉	産業医科大学第一内科学講座	教 授
	正木 康史	金沢医科大学大学院医科学研究科血液免疫内科学	准 教 授
	川野 充弘	金沢大学附属病院リウマチ・膠原病内科	科 長
研 究 協 力 者	北川 和子	金沢医科大学大学院医科学研究科眼科学	教 授
	中村 栄男	名古屋大学医学部・大学院医学系研究科病理組織医学	教 授
	折口 智樹	長崎大学大学院 医歯薬学総合研究科 リハビリテーション科学講座	教 授
	横山 仁	金沢医科大学腎臓内科学	教 授
	久保 惠嗣	信州大学医学部呼吸器内科	教 授
	伊藤 邦彦	静岡県立大学薬学部臨床薬効解析学分野	教 授
	利波 久雄	金沢医科大学放射線科	教 授
	高橋 裕樹	札幌医科大学医学部第一内科	准 教 授
	山本 元久	札幌医科大学医学部第一内科	助 教
	松井 祥子	富山大学保健管理センター	准 教 授
	佐伯 敬子	長岡赤十字病院内科	部 長
	小島 勝	獨協医科大学病理学形態	教 授
	神澤 輝実	東京都立駒込病院消化器内科	部 長
	浜野 英明	信州大学医学部消化器内科	助 教
	西山 進	倉敷成人病センターリウマチ科	部 長
	廣川 満良	医療法人神甲会限病院病理細胞診断部	部 長
	鈴木 律朗	名古屋大学医学部造血細胞移植情報管理部・生体統計学	准 教 授
	安積 淳	神戸海星病院眼科	部 長
	全 陽	金沢大学医学部病理部	准 教 授
	黒瀬 望	金沢医科大学病態診断医学	助 教
	石垣 靖人	金沢医科大学総合医学研究所生命科学研究領域	准 教 授
	佐藤 康晴	岡山大学大学院病態制御学腫瘍制御学病理学	講 師
	瀬戸 加大	愛知がんセンター研究所 (遺伝子医療研究部)	副 所 長
	塚本 憲史	群馬大学医学部附属病院腫瘍センター	センター長
	松本 守生	独立行政法人国立病院機構西群馬病院血液内科	医 長
	松本 洋典	京都府立医科大学血液・腫瘍内科	講 師
	坂井 晃	広島大学病院血液内科	講 師
	尾山 徳秀	新潟大学歯学総合病院眼科眼腫瘍・眼形成	医 員
	今村 好章	福井大学医学部附属病院病理部	部 長
	高比良雅之	金沢大学付属病院 (眼腫瘍・眼窩疾患)	講 師
	井上 大	富山県立中央病院放射線科	副 医 長
	藤川 敬太	健康保険諫早総合病院リウマチ科	医 長
	村山佳子	群馬県立がんセンター	部 長
	葉師神芳洋	愛媛大学医学部附属病院腫瘍センター	准 教 授
	菅井 進	久藤総合病院	院 長
	源 誠二郎	大阪府立呼吸器アレルギー医療センター アレルギー内科	部 長
	川端 大介	京都大学大学院医学研究科臨床免疫学	助 教
	小川 葉子	慶應義塾大学医学部眼科学教室	特任准教授
	早稲田優子	金沢大学付属病院呼吸器内科	医 員
	伊藤 直子	金沢大学大学院臓器機能制御学 (第二内科)	助 教
	八木 邦公	金沢大学大学院臓器機能制御学 (第二内科)	講 師
	坪井 洋人	筑波大学大学院人間総合科学研究科臨床免疫学	講 師
	山本 洋	信州大学医学部内科学第一講座	講 師
宮下賜一郎	独立行政法人国立病院機構 長崎医療センター	医 長	
平田信太郎	産業医科大学第一内科学講座	講 師	
山田 和徳	金沢大学附属病院リウマチ・膠原病内科	医 員	
森本 尚孝	公立三豊総合病院 内科	医 長	
田中 真生	金沢医科大学血液免疫内科学	准 教 授	
藤田 義正	金沢医科大学血液免疫内科学	講 師	
瀧木 俊興	金沢医科大学血液免疫内科学	助 教	
三木美由貴	金沢医科大学血液免疫内科学	助 教	
岩男 悠	金沢医科大学血液免疫内科学	助 教	
中島 章夫	金沢医科大学血液免疫内科学	助 教	
中村 拓路	金沢医科大学血液免疫内科学	大学院生	
事 務 局	良永 幸恵	金沢医科大学血液免疫内科学 〒920-0293 石川県河北郡内灘町大学1-1 T E L 0 7 6 - 2 8 6 - 2 2 1 1 (内線3539) F A X 0 7 6 - 2 8 6 - 9 2 9 0 e-mail yuki@kanazawa-med.ac.jp	
経理事務担当者	中川 邦子	金沢医科大学 研究推進課 T E L 0 7 6 - 2 8 6 - 2 2 1 1 (内線3052) F A X 0 7 6 - 2 8 6 - 2 3 4 6 e-mail hrc-jimu@kanazawa-med.ac.jp	

平成 23 年度第 2 回総会出席者名簿

平成 23 年 2 月 18 日 (土)

参加者 53 名 (敬称略)

班 長	岡崎 和一	関西医科大学内科学第三講座
分担研究者	梅原 久範	金沢医科大学血液免疫制御学
	神澤 輝実	都立駒込病院 消化器内科
	川 茂幸	信州大学総合健康安全センター
	坪内 博仁	鹿児島大学大学院医歯学総合研究科
	千葉 勉	京都大学大学院医学研究科消化器内科学講座
	日比 紀文	慶應義塾大学医学部消化器内科
	松田 文房	京都大学ゲノム医学センター
研究協力者	大原 弘隆	名古屋市立大学大学院医学研究科
	中沢 貴宏	名古屋市立大学大学院医学研究科
	太田 正穂	信州大学法医学教室
	乾 和郎	藤田保健衛生大学坂文種報徳會病院消化器内科
	洪 繁	名古屋大学医学部附属病院消化器内科
	中村 誠司	九州大学大学院研究院口腔顎顔面病態学講座
	能登原 憲司	倉敷中央病院 病理検査科
	西野 隆義	東京女子医科大学八千代医療センター消化器内科
	平野 賢二	東京大学医学部附属病院 消化器内科
	水野 伸匡	愛知県がんセンター中央病院消化器内科部
	吉田 仁	昭和大学医学部内科学講座消化器内科学部門
	濱野 英明	信州大学医学部附属病院医療情報部
	長谷部 修	長野市民病院 消化器内科
参加協力者	森山 雅文	前原 隆 田中 昭彦 九州大学顎顔面口腔外科
	渡邊 智裕	塩川 雅広 孟 凡斌 宇座 徳之 児玉 裕三
	栗田 亮	田辺 渉 京都大学消化器内科
	寺尾知可史	京都大学ゲノム医学センター
	半田 知宏	京都大学呼吸器内科
	佐伯 恵太	慶應義塾大学
	五十嵐久人	藤森 尚 九州大学病態制御内科
	菅野 敦	東北大学大学院医学系研究科
	宮部 勝之	内藤 格 名古屋市立大学
	寄山 敏男	鹿児島大学消化器内科
	栗山 勝利	大津赤十字病院
	小泉 理美	都立駒込病院
	中山 新士	岸本 真房 栗島 亜希子 中島 淳 楠田 武生
	山科 雅央	福井 由理 富山 尚 岡崎 敬 住本 貴美 高橋 悠
	山口 隆志	西尾 彰功 関西医科大学
事務局	内田 一茂	長谷川 也真 関西医科大学内科学第三講座 (消化器肝臓内科)

X. 研究成果の刊行物・別刷

Comprehensive diagnostic criteria for IgG4-related disease (IgG4-RD), 2011

Hisanori Umehara · Kazuichi Okazaki · Yasufumi Masaki · Mitsuhiro Kawano · Motohisa Yamamoto · Takako Saeki · Shoko Matsui · Tadashi Yoshino · Shigeo Nakamura · Shigeyuki Kawa · Hideaki Hamano · Terumi Kamisawa · Toru Shimosegawa · Akira Shimatsu · Seiji Nakamura · Tetsuhide Ito · Kenji Notohara · Takayuki Sumida · Yoshiya Tanaka · Tsuneyo Mimori · Tsutomu Chiba · Michiaki Mishima · Toshifumi Hibi · Hirohito Tsubouchi · Kazuo Inui · Hirotaka Ohara

Received: 14 October 2011 / Accepted: 19 November 2011 / Published online: 5 January 2012
© Japan College of Rheumatology 2011

Abstract

Background IgG4-related disease (IgG4-RD) is a novel clinical disease entity characterized by elevated serum IgG4 concentration and tumefaction or tissue infiltration by IgG4+ plasma cells. Although IgG4-RD is not rare and is clinically important, its clinical diagnostic criteria have not been established. Comprehensive diagnostic criteria for

IgG4-RD, including the involvement of various organs, are intended for the practical use of general physicians and nonspecialists.

Methods Two IgG4-RD study groups, the Umehara and Okazaki teams, were organized by the Ministry of Health, Labor and Welfare Japan. As IgG4-RD comprises a wide variety of diseases, these groups consist of physicians and researchers in various disciplines, including rheumatology, hematology, gastroenterology, nephrology, pulmonology, ophthalmology, odontology, pathology, statistics, and basic and molecular immunology throughout Japan, with 66 and 56 members of the Umehara and Okazaki teams, respectively. Collaborations of the two study groups involved

For the All Japan IgG4 team.

Professional collaborators of the All Japan G4 team are given in the Appendix.

H. Umehara and K. Okazaki equally contributed to this work.

H. Umehara · Y. Masaki
Department of Hematology and Immunology,
Kanazawa Medical University, Kanazawa, Ishikawa, Japan

T. Saeki
Department of Internal Medicine, Nagaoka Red Cross Hospital,
Nagaoka, Niigata, Japan

H. Umehara (✉)
Division of Hematology and Immunology,
Department of Internal Medicine, Kanazawa Medical University,
1-1 Daigaku, Uchinada-machi, Kahoku-gun,
Kanazawa, Ishikawa 920-0293, Japan
e-mail: umehara@kanazawa-med.ac.jp

S. Matsui
Health Administration Center, Sugitani Campus,
University of Toyama, Toyama, Japan

K. Okazaki (✉)
Division of Gastroenterology and Hepatology,
The Third Department of Internal Medicine,
Kansai Medical University, Hirakata, Osaka 573-1191, Japan

T. Yoshino
Department of Pathology, Okayama University Graduate School
of Medicine, Dentistry and Pharmaceutical Sciences,
Okayama, Japan

M. Kawano
Division of Rheumatology, Department of Internal Medicine,
Kanazawa University Graduate School of Medical Science,
Kanazawa, Ishikawa, Japan

S. Nakamura
Department of Pathology and Laboratory Medicine,
Nagoya University Hospital, Nagoya, Japan

M. Yamamoto
The First Department of Internal Medicine,
Sapporo Medical University, Sapporo, Hokkaido, Japan

S. Kawa
Center for Health, Safety and Environmental Management,
Shinshu University, Matsumoto, Japan

H. Hamano
Department of Gastroenterology, Shinshu University School
of Medicine, Matsumoto, Japan

detailed analyses of clinical symptoms, laboratory results, and biopsy specimens of patients with IgG4-RD, resulting in the establishment of comprehensive diagnostic criteria for IgG4-RD.

Results Although many patients with IgG4-RD have lesions in several organs, either synchronously or meta-chronously, and the pathological features of each organ differ, consensus has been reached on two diagnostic criteria for IgG4RD: (1) serum IgG4 concentration >135 mg/dl, and (2) >40% of IgG+ plasma cells being IgG4+ and >10 cells/high powered field of biopsy sample. Although the comprehensive diagnostic criteria are not sufficiently sensitive for the diagnosis of type 1 IgG4-related autoimmune pancreatitis (IgG4-related AIP), they are adequately sensitive for IgG4-related Mikulicz's disease (MD) and kidney disease (KD). In addition, the comprehensive diagnostic criteria, combined with organ-specific diagnostic criteria, have increased the sensitivity of diagnosis to 100% for IgG4-related MD, KD, and AIP.

Conclusion Our comprehensive diagnostic criteria for IgG4-RD are practically useful for general physicians and nonspecialists.

Keywords IgG4-related disease · Criteria · Mikulicz's disease · Autoimmune pancreatitis · Interstitial nephritis

T. Kamisawa
Department of Internal Medicine, Tokyo Metropolitan
Komagome Hospital, Tokyo, Japan

T. Shimosegawa
Division of Gastroenterology, Tohoku University Graduate
School of Medicine, Sendai, Miyagi, Japan

A. Shimatsu
Clinical Research Institute, National Hospital Organization
Kyoto Medical Center, Kyoto, Japan

S. Nakamura
Section of Oral and Maxillofacial Oncology,
Division of Maxillofacial Diagnostic and Surgical Sciences,
Faculty of Dental Science, Kyushu University, Fukuoka, Japan

T. Ito
Department of Medicine and Bioregulatory Science,
Graduate School of Medical Sciences, Kyushu University,
Fukuoka, Japan

K. Notohara
Department of Anatomic Pathology, Kurashiki Central Hospital,
Kurashiki, Japan

K. Notohara
Department of Pathology and Laboratory Medicine,
Kanazawa Medical University, Kanazawa, Ishikawa, Japan

Abbreviations

IgG4-RD	IgG4-related disease
MD	Mikulicz's disease
AIP	Autoimmune pancreatitis
KD	Kidney disease
TIN	Tubulointerstitial nephritis
SS	Sjögren's syndrome
MHLW	Japan Ministry of Health, Labor and Welfare Japan; familial multifocal fibrosclerosis
RPF	Retroperitoneal fibrosis
TIN	Tubulointerstitial nephritis
MOLPS	Multiorgan lymphoproliferative syndrome
SIPS	Systemic IgG4 plasmacytic syndrome

Introduction

IgG4-related disease (IgG4-RD) is a new emerging disease entity of unknown etiology with multiorgan involvement [1]. IgG4-RD has been found to affect the pancreas, bile duct, lacrimal glands, salivary glands, central nervous system, thyroid, lungs, liver, gastrointestinal tract, kidney, prostate, retroperitoneum, arteries, lymph nodes, skin, and breast. Therefore, IgG4-RD includes a wide variety of diseases, including Mikulicz's disease (MD) [2, 3], autoimmune pancreatitis (AIP) [4], hypophysitis, Riedel

T. Sumida
Doctoral Programs in Clinical Science, Department of Clinical
Immunology, Graduate School of Comprehensive Human
Science, University of Tsukuba, Tsukuba, Ibaraki, Japan

Y. Tanaka
First Department of Internal Medicine, School of Medicine,
University of Occupational and Environmental Health, Fukuoka,
Japan

T. Mimori
Department of Rheumatology and Clinical Immunology,
Graduate School of Medicine, Kyoto University, Kyoto, Japan

T. Chiba
Department of Gastroenterology and Hepatology,
Graduate School of Medicine, Kyoto University,
Kyoto, Japan

M. Mishima
Department of Respiratory Medicine, Graduate School
of Medicine, Kyoto University, Kyoto, Japan

T. Hibi
Division of Gastroenterology and Hepatology,
Department of Internal Medicine,
Keio University School of Medicine,
Tokyo, Japan

thyroiditis [5], interstitial pneumonitis [6, 7], interstitial nephritis [8, 9], prostatitis, lymphadenopathy [10, 11], retroperitoneal fibrosis [12, 13], inflammatory aortic aneurysm [14], and inflammatory pseudotumor. Although IgG4-RD is not rare and is clinically important, its clinical diagnostic criteria have not yet been established. Two study groups were thus organized by the Ministry of Health, Labor and Welfare (MHLW) Japan. One group, the Umehara team, chaired by Professor Umehara of Kanazawa Medical University, is seeking to establish diagnostic criteria for IgG4-RD; the second group, the Okazaki team, chaired by Professor Okazaki of Kansai Medical University, is seeking to understand the etiology and pathogenesis of IgG4-RD. These groups consist of physicians and researchers in various fields, including rheumatology, hematology, gastroenterology, nephrology, pulmonology, ophthalmology, odontology, pathology, statistics, and basic and molecular immunology from all over Japan, with 66 and 56 members of the Umehara and Okazaki teams, respectively.

Background for establishing diagnostic criteria for IgG4-RD

General concepts of IgG4-RD

Although the two groups independently analyzed the clinical features and conditions of IgG4-RD, they collaborated closely, which resulted in the following consensus: (1) IgG4-RD can occur in various organs, including the central nervous system, salivary glands, thyroid gland, lungs, pancreas, biliary duct, liver, gastrointestinal tract, kidneys, prostate gland, retroperitoneum, and lymph nodes, with clinical symptoms depending on lesion location. (2) IgG4-RD mainly affects middle-aged to elderly men; its clinical symptoms are relatively mild, and the condition usually comes to clinical attention due to organ swelling or damage. (3) Many patients with IgG4-RD can be treated effectively by steroid therapy. (4) Although the infiltration of IgG4+

cells and increased serum concentrations of IgG4 are characteristic of IgG4-RD, the severity of fibrosis is dependent on the individual organs involved. The common characteristics of these conditions include elevated serum IgG4 concentrations and tissue infiltration by IgG4+ plasma cells, accompanied by tissue fibrosis and sclerosis [1].

Naming of IgG4-related disease

Many terms have been used to describe IgG4-RD, including IgG4-related sclerosing disease [15], IgG4-related autoimmune disease [16], systemic IgG4 plasmacytic syndrome (SIPS) [17], and IgG4-related multiorgan lymphoproliferative syndrome (IgG4-MOLPS) [3]. The members of the Umehara and Okazaki teams carefully examined reports using these different nomenclatures and concluded that they referred to the same condition, and the two teams finally agreed to use a uniform nomenclature—IgG4-related disease (IgG4-RD)—for several reasons: (1) Although infiltration of IgG4+ cells and increased serum concentrations of IgG4 are characteristic of IgG4-RD, the severity of fibrosis is dependent on the individual organs involved. For example, storiform fibrosis is characteristic of IgG4-related autoimmune pancreatitis (IgG4-related AIP), IgG4-related retroperitoneal fibrosis (IgG4-related RPF), and IgG4-related tubulointerstitial nephritis (IgG4-related TIN), but is very seldom found in patients with IgG4-related MD and IgG4-related lymphadenopathy. (2) Although many patients with this IgG4-RD have lesions in several organs, either synchronously or metachronously, other patients show involvement of a single organ. (3) As there have been several reports describing patients with IgG4-associated conditions concomitant with malignant tumors, such as pancreatic and salivary carcinomas [18–21] and ocular adnexal lymphoma [22, 23], using the term systemic may lead to an incorrect diagnosis of an IgG4-related condition in a patient with malignant tumors in other organs [24].

Prevalence of IgG4-RD

It is difficult to ascertain the number of patients with IgG4-RD because the awareness of this disease is low and its diagnostic criteria have not yet been established. The Umehara team attempted to estimate the number of individuals with IgG4-RD throughout Japan by using as an example Ishikawa Prefecture, which contains 1.16 million people with little population inflow/outflow. The incidence of this disease throughout Japan was estimated to be 0.28–1.08/100,000, with 336–1,300 patients newly diagnosed per year and approximately 6,700–26,000 patients who developed IgG4-RD over the past 20 years [1]. In contrast, the Okazaki team attempted to estimate the

H. Tsubouchi
Division of Digestive and Lifestyle Disease,
Department of Human and Environmental Sciences,
Kagoshima University Graduate School of Medical and Dental
Sciences, Kagoshima, Japan

K. Inui
Department of Internal Medicine, Second Teaching Hospital,
Fujita Health University, Toyoake, Aichi, Japan

H. Ohara
Department of Internal Medicine and Bioregulation,
Nagoya City University Graduate School of Medical Sciences,
Nagoya, Aichi, Japan

incidence of IgG4-RD through a network of Japanese researchers in an AIP study; they reported that 8,000 patients throughout Japan had IgG4-RD.

Proposal of comprehensive diagnostic criteria for IgG4-RD

Concept of comprehensive diagnostic criteria for IgG4-RD

IgG4-RD may occur, either synchronously or metachronously, in a variety of organs throughout the body, including the pancreas, bile duct, lacrimal gland, salivary gland, thyroid, lung, liver, gastrointestinal tract, kidney, and retroperitoneum [1]. As clinical symptoms and pathological features depend on lesion location, it is probably impossible to establish criteria that include all patients with IgG4-RD. Detailed diagnostic criteria are needed for the involvement of each organ, including clinical symptoms, serological and histological findings, and radiological images. To date, diagnostic criteria for IgG4-related MD [25] (Table 1), IgG4-related AIP type 1 [26] (Table 2), and IgG4-related kidney disease (KD) [27] (Table 3) have been established. However, these organ-specific criteria are not suitable for the diagnosis of patients with involvement of other organs. In addition, organ-specific criteria may not be

familiar to general clinicians and specialists in diseases of those organs, although all clinicians should become aware of this new disease entity and its diagnosis. Therefore, comprehensive diagnostic criteria are necessary for practical use and to differentiate among malignancies.

Comprehensive diagnostic criteria for IgG4-RD

The comprehensive diagnostic criteria we have proposed for IgG4-RD (Table 4) consist of three parts: concept, diagnostic criteria, and explanatory notes. The concept clarifies the features characteristic of IgG4-RD, such as lesion location, symptoms, and prognosis. Diagnostic criteria are based on two major characteristics of IgG4-RD: increased serum concentrations of IgG4 and infiltration of IgG4+ cells. The cutoff value for serum IgG4 concentration, 135 mg/dl, was based on receiver operating characteristic (ROC) curves, and its validity was confirmed in patients with AIP [26, 28]. Although tissue biopsies are difficult to obtain from some organs, including the pancreas, retroperitoneum, and ocular cavity, histopathological examination is important. Because IgG4+ plasma cell infiltration has been reported in various diseases and clinical conditions, such as rheumatoid synovitis, inflammatory oral and skin lesions, and carcinomas with a peritumoral inflammatory response [29], pathological criteria should be rigorous. Histopathological findings of marked IgG4+ cell

Table 1 Diagnostic criteria for IgG4+ Mikulicz's disease [25] (approved by the Japanese Society for Sjögren's Syndrome, 2008)

1. Symmetrical swelling of at least two pairs of lacrimal, parotid, and submandibular glands continuing for more than 3 months; and
 2. Elevated serum IgG4 (>135 mg/dl);
- or
3. Histopathological features including lymphocyte and IgG4+ plasma-cell infiltration (IgG4+ plasma cells/IgG+ plasma cells >50%) with typical tissue fibrosis or sclerosis

Differential diagnosis is necessary from other disorders, including sarcoidosis, Castleman's disease, Wegener's granulomatosis, lymphoma, and cancer. Although the diagnostic criteria for Sjögren's syndrome (SS) may also include some patients with IgG4+ Mikulicz's disease, the clinicopathological conditions of patients with typical SS and IgG4+ Mikulicz's disease are different

Table 2 Clinical diagnostic criteria for autoimmune pancreatitis in Japan (2006) [26]

1. Diffuse or segmental narrowing of the main pancreatic duct with irregular walls and diffuse or localized enlargement of the pancreas on imaging modalities, including abdominal ultrasound, computed tomography, and magnetic resonance imaging
2. High-serum F-globulin, IgG, or IgG4 concentration or the presence of autoantibodies, such as antinuclear antibodies and rheumatoid factor
3. Marked interlobular fibrosis and prominent infiltration of lymphocytes and plasma cells to the periductal area, occasionally accompanied by lymphoid follicles in the pancreas

For diagnosis, criterion 1 must be present, together with criterion 2 and/or 3

However, it is necessary to exclude malignant diseases such as pancreatic and biliary cancers

Table 3 Diagnostic criteria for IgG4-related kidney disease [27]

1. Presence of some kidney damage, as manifested by abnormal urinalysis or urine marker(s) or decreased kidney function with either elevated serum IgG or IgE or hypocomplementemia	
2. Abnormal renal radiologic findings:	
a. Multiple low-density lesions on enhanced computed tomography	
b. Diffuse kidney enlargement	
c. Hypovascular solitary mass in the kidney	
d. Hypertrophic lesion of the renal pelvic wall without irregularities of the renal pelvic surface	
3. Elevated serum IgG4 level (>135 mg/dl)	
4. Histological findings in the kidney:	
a. Dense lymphoplasmacytic infiltration by >10 IgG4+ plasma cells/high power field (HPF) and/or IgG4+/IgG+ plasma cells >40%	
b. Characteristic (sclero-) fibrosis surrounding nests of lymphocytes and/or plasma cells	
5. Histological findings in extra-renal organ(s):	
Dense lymphoplasmacytic infiltration by >10 IgG4+ plasma cells/HPF and/or IgG4+/IgG+ plasma cells >40%	
Definite:	1 + 3 + 4 a, b
	2 + 3 + 4 a, b
	2 + 3 + 5
	1 + 3 + 4 a + 5
Probable:	1 + 4 a, b
	2 + 4 a, b
	2 + 5
	3 + 4 a, (b)
Possible:	1 + 3
	2 + 3
	1 + 4 a
	2 + 4 a
Appendix:	
1. Clinically and histologically, the following diseases should be excluded: Wegener's granulomatosis, Churg–Strauss syndrome, extramedullary plasmacytoma	
2. Radiologically, the following diseases should be excluded: malignant lymphoma, urinary tract carcinomas, renal infarction, and pyelonephritis (rarely, Wegener's granulomatosis, sarcoidosis and metastatic carcinoma)	

infiltration [>10 cells/high power field (HPF)] and an IgG4+/IgG+ cell ratio $>40\%$ are diagnostic of IgG4-RD. Explanatory notes describe clinical characteristics of IgG4-RD specific to each organ, as well as blood tests and pathologic findings, responses to steroids, and differential diagnoses.

Algorithm for diagnosing IgG4-RD

A diagnostic algorithm for IgG4-RD, using comprehensive diagnostic criteria combined with organ-specific criteria, is shown in Fig. 1. A diagnosis of IgG4-RD is definitive in patients with: (1) organ enlargement, mass or nodular lesions, or organ dysfunction; (2) a serum IgG4 concentration >135 mg/dl; and (3) histopathological findings of >10 IgG4 cells/HPF and an IgG4+/IgG+ cell ratio $>40\%$ (category 1). A diagnosis of IgG4-RD is possible in patients who fulfill criteria (1) and (2), but with negative results on histopathology or without histopathologic

examination (category 2 and 3), whereas a diagnosis of IgG4-RD is probable in patients with organ involvement (1) and fulfilled histopathologic criteria, but without increased serum IgG4 concentration (2) (category 4). Patients with organ symptoms without satisfying serologic or histopathologic criteria are considered unlikely to have IgG4-RD (category 5 and 6). For patients in categories 2–5, organ-specific criteria for IgG4-RD could be applied, such as those for AIP [26], MD [25], and KD [27] associated with IgG4. Patients who fulfill the organ-specific criteria for IgG4-RD have a definite diagnosis of this disease (category 7).

Validation of comprehensive diagnostic criteria in previous reports of patients with IgG4-RD

To validate the comprehensive diagnostic criteria, we applied them to patients described in two studies of IgG4-related MD [3, 30], two of IgG4-related KD [9, 27], and

Table 4 Comprehensive diagnostic criteria for IgG4-related disease, 2011**I. Concept**

IgG4-related disease (IgG4-RD) shows organ enlargement or nodular/hyperplastic lesions in various organs concurrently or metachronously, due to marked infiltration of lymphocytes and IgG4+ plasma cells, as well as fibrosis of unknown etiology. IgG4-RD affects various organs, including the pancreas, bile duct, lacrimal gland, salivary gland, central nervous system, thyroid, lung, liver, gastrointestinal tract, kidney, prostate, retroperitoneum, arteries, lymph nodes, skin, and breast. Although many patients with IgG4-RD have lesions in several organs, either synchronously or metachronously, others show involvement of a single organ. Clinical symptoms vary depending on the affected organ, and some patients may experience serious complications, such as obstruction or compression symptoms due to organomegaly or hypertrophy, and organ dysfunction caused by cellular infiltration or fibrosis. Steroid therapy is often effective

II. [Comprehensive clinical diagnostic criteria for IgG4-RD]

1. Clinical examination showing characteristic diffuse/localized swelling or masses in single or multiple organs

2. Hematological examination shows elevated serum IgG4 concentrations (≥ 135 mg/dl)

3. Histopathologic examination shows:

(1) Marked lymphocyte and plasmacyte infiltration and fibrosis.

(2) Infiltration of IgG4+ plasma cells: ratio of IgG4+/IgG+ cells > 40% and >10 IgG4+ plasma cells/HPF

Definite: 1) + 2) + 3)

Probable: 1) + 3)

Possible: 1) + 2)

However, it is important to differentiate IgG4-RD from malignant tumors of each organ (e.g. cancer, lymphoma) and similar diseases (e.g. Sjögren's syndrome, primary sclerosing cholangitis, Castleman's disease, secondary retroperitoneal fibrosis, Wegener's granulomatosis, sarcoidosis, Churg–Strauss syndrome) by additional histopathological examination

Even when patients cannot be diagnosed using the CCD criteria, they may be diagnosed using organ-specific diagnostic criteria for IgG4RD

III. Explanatory notes

1. The comprehensive diagnostic criteria are the minimal consensus to aid general practitioners and other nonspecialist physicians in the clinical diagnosis of IgG4-RD. For each affected organ, organ-specific diagnostic criteria established for IgG4-related Mikulicz's disease, IgG4-related autoimmune pancreatitis, and IgG4-related kidney disease, should be used concurrently

2. Concept:

The difference from multifocal fibrosclerosis is unclear although these diseases may be IgG4-RD. Many patients show multiple organ involvement and are characterized as having systemic disease, whereas other patients show involvement of a single organ

(a) Autoimmune pancreatitis, type 1 (IgG4-related autoimmune pancreatitis): This disease is synonymous with IgG4-related sclerosing pancreatitis/lymphoplasmacytic sclerosing pancreatitis (LPSP). It can be diagnosed using the clinical diagnostic criteria for autoimmune pancreatitis established by the Ministry of Health, Labor and Welfare, Japan Pancreas Society, in 2006 [26]

(b) IgG4-related sclerosing cholangitis: This disease is characterized by sclerotic changes with diffuse or localized stenosis in the intrahepatic/extrahepatic bile duct and gallbladder. Circumferential wall thickening is observed at the site of stenosis, with similar changes in areas without stenosis. Obstructive jaundice often develops, making it important to differentiate this condition from tumors, such as cholangiocarcinoma and pancreatic cancer, and from primary sclerosing cholangitis. It is also necessary to exclude secondary sclerosing cholangitis as an apparent cause

(c) IgG4-related lacrimal, orbital, and salivary-gland lesions: This condition includes IgG4-related Mikulicz's disease characterized by symmetrical (sometimes unilateral) swelling of any of the lacrimal, parotid, submandibular, sublingual glands, and some minor salivary glands. Nodular/infiltrative lesions may also occur in orbital tissue other than the lacrimal glands. IgG4-related Mikulicz's disease can be diagnosed by the organ-specific diagnostic criteria for IgG4-related Mikulicz's disease established by the Sjögren's Syndrome Study Group of Japan in 2008 [25]

(d) IgG4-related central nervous system lesions: These lesions include infundibular hypophysitis, hypertrophic pachymeningitis, and intracerebral inflammatory pseudotumor

Table 4 continued

(e) IgG4-related respiratory lesions: These lesions occur primarily in the interstitium, such as bronchovascular bundles, interlobular septum, alveolar septum, and pleura. They are frequently accompanied by mediastinal and hilar lymphadenopathy, along with X-ray evidence of a mass or infiltration of the lung. Some patients have asthma-like symptoms. It is important to differentiate these lesions from malignant tumors, sarcoidosis, collagen diseases of the lung, and infection

(f) IgG4-related renal lesions: Abnormal imaging findings include diffuse renal enlargement, multifocal contrast defects of the renal parenchyma, renal mass lesions, and pelvic wall thickening. Renal histology shows mainly interstitial nephritis, but glomerular lesions (e.g., membranous nephropathy), may also be present. IgG4-related tubulointerstitial nephritis can be diagnosed using the organ-specific diagnostic criteria for IgG4-related kidney disease [27]

(g) IgG4-related retroperitoneal fibrosis/periarterial lesions: This disease is characterized by thickening of the abdominal aortic adventitia and periurethral soft tissue, often accompanied by hydronephrosis or mass lesions. Periarteritis may occur around the aorta or relatively large branches and is evident as arterial wall thickening on radiological imaging. Magnetic resonance imaging (MRI) and positron emission tomography (PET) have been shown to be helpful for diagnosing retroperitoneal fibrosis in addition to X-ray, which may include CT scan. Biopsy is often inconclusive, making it difficult to differentiate this condition from secondary retroperitoneal fibrosis due to malignant tumors or infectious diseases

(h) Other tumefactive lesions: Proliferation of IgG4+ plasma cells and lymphocytes may accompany fibrosis. Including some conventional inflammatory pseudotumors, these lesions have been reported in the brain, orbit, lung, breast, liver, pancreas, retroperitoneum, kidney, and lymph nodes

IV. Blood test findings

1. Polyclonal serum γ -globulin, IgG, and IgE are often elevated, and hypocomplementemia may occur
2. Elevated IgG4 can also be seen in other diseases (e.g., atopic dermatitis, pemphigus, asthma, and multicentric Castleman's disease) and is therefore not specific to IgG4-RD
3. On rare occasions, serum IgG4 concentration may be elevated in patients with malignant tumors. However, patients with >270 mg/dl IgG4 are unlikely to have pancreatic cancer
4. In patients with single-organ involvement and serum IgG4 concentration <135 mg/dl, the IgG4+/IgG+ ratio may be helpful in making a diagnosis
5. At present, the significance of elevated IgG4 in the pathogenesis/pathophysiology of IgG4-RD is unknown

V. Histopathological findings

1. Storiform or swirling fibrosis or obliterative phlebitis are characteristic of IgG4-RD and may be important in its diagnosis
2. Eosinophilic infiltration often occurs, along with infiltration of IgG4+ cells
3. Reactive infiltration of IgG4+ cells and fibrosis may also occur, such as at the periphery of pancreatic cancers

VI. Imaging studies

IgG4-RD may occur, either synchronously or metachronously, in a variety of organs throughout the body, including the pancreas, bile duct, lacrimal gland, salivary gland, thyroid, lung, liver, gastrointestinal tract, kidney, and retroperitoneum. MRI and fluorodeoxyglucose (FDG)-PET have been shown to be helpful for detecting multiorgan involvements

VII. Steroids

1. Patients with malignant lymphoma or paraneoplastic lesions can sometimes be improved by steroid administration. Therefore, steroid trials should be strictly avoided
2. Efforts should be made to collect tissue samples for diagnosis. However, patients having disease in organs difficult to biopsy, such as pancreas, retroperitoneum, and pituitary, and who respond to steroids may possibly have IgG4-RD
3. In accordance with the guidelines for treatment of autoimmune pancreatitis, patients should be started on 0.5–0.6 mg/kg per day of prednisolone. If patients do not respond to the initial steroid therapy, the diagnosis should be reviewed

VIII. Diseases to be excluded or differentiated

1. To exclude malignancies (e.g., cancer, lymphoma) in involved organs, it is essential to determine histopathologically whether malignant cells are present
2. Similar diseases (e.g., Sjögren's syndrome, primary sclerosing cholangitis, multicentric Castleman's disease, idiopathic retroperitoneal fibrosis, Wegener's granulomatosis, sarcoidosis, Churg–Strauss syndrome) are diagnosed using the diagnostic criteria for each disease
3. Multicentric Castleman's disease is a hyper-interleukin (IL)-6 syndrome and is not included among the IgG4-RDs, even if the diagnostic criteria for IgG4-RD are fulfilled

three of IgG4-related AIP [31] (Table 5). The sensitivity of these criteria were comparatively good for diagnosing IgG4-related MD (83 and 70%) and KD (87 and 85%). In contrast, patients with IgG4-related AIP could not be diagnosed by the comprehensive diagnostic criteria (0% for

definite, nearly 70% for possible, and 10–30% for unlikely) because biopsies could not be obtained from most of these patients. Application of organ-specific criteria to undiagnosed patients increased the sensitivity of diagnosis to 100%, even for patients with IgG4-related AIP (Table 5).




Fig. 1 Diagnostic algorithm performance for comprehensive diagnostic criteria for IgG4-related disease (IgG4-RD) using comprehensive diagnostic criteria combined with organ-specific criteria. A diagnosis of IgG4-RD is definitive in patients with (1) organ enlargement, mass or nodular lesions, or organ dysfunction, (2) a serum IgG4 concentration >135 mg/dl, and (3) histopathological findings of >10 IgG4+ cells/HPF and an IgG4+/IgG+ cell ratio >40% (C1). A diagnosis of IgG4-RD is possible in patients who fulfill criteria (1) and (2), but with negative results on histopathology or without histopathologic examination (C2, C3), whereas a diagnosis of IgG4-RD is probable in patients with (1) organ involvement and (2) fulfilled histopathologic criteria, but without increased serum IgG4 concentration (C4). Patients with organ symptoms without satisfying the serologic or histopathologic criteria are considered unlikely to have IgG4-RD (C5, C6). For patients in C2–C6, organ-specific criteria for IgG4-related autoimmune pancreatitis (AIP), IgG4-related Mikulicz's disease (MD), and IgG4-related kidney disease (KD). Patients who fulfill the organ-specific criteria have a definite diagnosis of IgG4-RD (C7)

Discussion

Although there is increased interest in IgG4-RD, awareness of it remains low and diagnostic criteria have not yet been published. Therefore, IgG4-RD has often been misdiagnosed as a malignant tumor, lymphoma, Sjögren's syndrome, or other diseases despite the effectiveness of steroid therapy. As IgG4-RD affects various organs, its clinical symptoms vary, and each patient with IgG4-RD may visit specialists addressing organ-specific lesions. Organ-specific

criteria have been established for IgG4-related AIP [26], MD [25] and KD [27], but these criteria are not suitable for diagnosing patients with other involved organs, and they are not familiar to general clinicians and nonspecialists. Comprehensive diagnostic criteria are therefore needed for practical use by such physicians. Although it is difficult to obtain tissue biopsy samples from some organs, including the pancreas, retroperitoneum, and brain, histopathologic examination is highly important to exclude malignancies [18–21] and other types of disease [29]. Indeed, most patients with IgG4-related AIP could be diagnosed without biopsy. The comprehensive diagnostic criteria for IgG4-RD have relatively low sensitivity in patients with IgG4-related AIP because of a lack of biopsy samples but were sufficiently sensitive for IgG4-related MD and KD. Patients who could not be diagnosed by the comprehensive diagnostic criteria could be diagnosed by organ-specific criteria, indicating the complementarity of comprehensive diagnostic criteria and organ-specific criteria for IgG4-RD.

Acknowledgments This work was supported by Intractable Diseases, the Health and Labor Sciences Research Grants from Ministry of Health, Labor and Welfare, Japan. We sincerely thank the many contributing researchers and collaborators who participated in the All Japan IgG4 team.

Conflict of interest None.

Appendix

The authors thank the many patients who participated in this registry. In addition to the listed authors, other professional collaborators of the All Japan G4 team in the Research Program for Intractable Disease by Ministry of Health, Labor and Welfare (MHLW) Japan, include: Atsushi Azumi (Kobe Kaisei Hospital); Keiji Kubo, and Hiroshi Yamamoto (Shinshu University); Daisuke Kawabata (Kyoto University); Seiji Minamoto (Osaka Respiratory and Allergy Center); Susumu Nishiyama (Kurashiki Hospital); Kazuo Tsubota and Yoko Ogawa (Keio University); Shintaro Hirata (University of Occupational and Environmental Health); Tomoki Origuchi (Nagasaki University); Yasuharu Sato (Okayama University); Susumu Sugai (Kudou Hospital); Hiroki Takahashi (Sapporo Medical University); Hiroto Tsuboi (Tsukuba University); Dai Inoue, Masayuki Takahira and Yuko Waseda (Kanazawa University); Masaru Kojima (Dokkyo University School of Medicine); Norifumi Tsukamoto (Gunma University); Morio Matsumoto (Nishigunma National Hospital); Kayoko Murayama (Gunma Prefectural Cancer Center); Ritsuro Suzuki and Shigeru Ko (Nagoya University); Takahiro Nakazawa and Osamu Hasebe (Nagoya City University);

Table 5 Sensitivity for diagnosis by comprehensive diagnostic criteria for IgG4-RD

Main organ	Definite	Probable	Possible	Denial	References
Mikulicz (64)	53 (83%)	4 (6%)	7 (11%)	0 (0%)	Masaki et al. [3]
+OS criteria		4/4	7/7		
Total	64 (100%)				
Mikulicz (40)	28 (70%)	0 (0%)	12 (30%)	0 (0%)	Yamamoto et al. [30]
+OS criteria			12/12		
Total	40/40 (100%)				
Kidney (23) ^a	20 (87%)	0 (0%)	0 (0%)	3 (13%)	Saeki et al. [9]
+OS criteria				3/3	
Total	23 (100%)				
Kidney (41) ^a	35 (85%)	0 (0%)	3 (7%)	3 (7%)	Kawano et al. [27]
+OS criteria			3/3	3/3	
Total	41/41 (100%)				
AIP (60)	0 (0%)	0 (0%)	41 (68%)	19 (32%)	Takuma et al. [32]
+OS criteria			41/41	19/19	
Total	60 (100%)				
AIP (54)	0 (0%)	0 (0%)	42 (78%)	12 (22%)	Okazaki et al. [31]
+OS criteria			42/42	12/12	
Total	54 (100%)				
<i>OS criteria organ-specific criteria</i>					
AIP (90)	0 (0%)	3 (3%)	70 (78%)	9 (10%)	Fujiwara et al. [33]
+OS criteria		3/3	70/70	9/9	
Total	90 (100%)				

^a 10 patients were included both in Refs. [9, 27]

Mitsuyoshi Hirokawa (Kuma Hospital); Masao Seto and Nobumasa Mizuno (Aichi Cancer Center); Yosuke Matsumoto (Kyoto Prefecture University); Tokuhide Oyama (Niigata University); Akira Sakai (Hiroshima University); Yoshiaki Imamura (Fukui University); Keita Fujikawa (Isahaya hospital); Yoshihiro Yakushijin (Ehime University); Akira Sakai (Hiroshima University); Takayoshi Nishino (Tokyo Womens Medical University), Kenji Hirano (Tokyo University), Hitoshi Yoshida (Showa University), Tatsuo Kinashi, Kazushige Uchida (Kansai Medical University), Kunihiro Itoh (University of Shizuoka) and Kazuko Kitagawa, Tsutomu Takegami, Naohisa Tomosugi, Hisanori Tonami, Nozomu Kurose, Yasuhito Ishigaki, Takayuki Nojima, Hitoshi Yokoyama, Toshihiro Fukushima, Masao Tanaka, Yoshimasa Fujita, Toshioki Sawaki, Takafumi Kawanami, Miyuki Miki, Haruka Iwao, Akio Nakajima, and Takuji Nakamura (Kanazawa Medical University).

References

- Umehara H, Okazaki K, Masaki Y, Kawano M, Yamamoto M, Saeki T, et al. A novel clinical entity, IgG4-related disease (IgG4RD): general concept and details. *Mod Rheumatol* 2011; [Epub ahead of print]
- Yamamoto M, Ohara M, Suzuki C, Naishiro Y, Yamamoto H, Takahashi H, et al. Elevated IgG4 concentrations in serum of patients with Mikulicz's disease. *Scand J Rheumatol*. 2004;33:432–3.
- Masaki Y, Dong L, Kurose N, Kitagawa K, Morikawa Y, Yamamoto M, et al. Proposal for a new clinical entity, IgG4-positive multi-organ lymphoproliferative syndrome: Analysis of 64 cases of IgG4-related disorders. *Ann Rheum Dis*. 2009;63:1310–5.
- Okazaki K, Uchida K, Koyabu M, Miyoshi H, Takaoka M. Recent advances in the concept and diagnosis of autoimmune pancreatitis and IgG4-related disease. *J Gastroenterol*. 2011;46(3):277–88.
- Dahlgren M, Khosroshahi A, Nielsen GP, Deshpande V, Stone JH. Riedel's thyroiditis and multifocal fibrosclerosis are part of the IgG4-related systemic disease spectrum. *Arthritis Care Res (Hoboken)*. 2010;62(9):1312–8.
- Zen Y, Inoue D, Kitao A, Onodera M, Abo H, Miyayama S, et al. IgG4-related lung and pleural disease: a clinicopathologic study of 21 cases. *Am J Surg Pathol*. 2009;33(12):1886–93.
- Inoue D, Zen Y, Abo H, Gabata T, Demachi H, Kobayashi T, et al. Immunoglobulin G4-related lung disease: CT findings with pathologic correlations. *Radiology*. 2009;251(1):260–70.
- Saeki T, Saito A, Yamazaki H, Emura I, Imai N, Ueno M, et al. Tubulointerstitial nephritis associated with IgG4-related systemic disease. *Clin Exp Nephrol*. 2007;11(2):168–73.
- Saeki T, Nishi S, Imai N, Ito T, Yamazaki H, Kawano M, et al. Clinicopathological characteristics of patients with IgG4-related tubulointerstitial nephritis. *Kidney Int*. 2010;78(10):1016–23.
- Sato Y, Kojima M, Takata K, Morito T, Asaoku H, Takeuchi T, et al. Systemic IgG4-related lymphadenopathy: a clinical and pathologic comparison to multicentric Castlemann's disease. *Mod Pathol*. 2009;22(4):589–99.
- Sato Y, Notohara K, Kojima M, Takata K, Masaki Y, Yoshino T. IgG4-related disease: historical overview and pathology of hematological disorders. *Pathol Int*. 2010;60(4):247–58.
- Hamano H, Kawa S, Ochi Y, Unno H, Shiba N, Wajiki M, et al. Hydronephrosis associated with retroperitoneal fibrosis and sclerosing pancreatitis. *Lancet*. 2002;359(9315):1403–4.

13. Zen Y, Onodera M, Inoue D, Kitao A, Matsui O, Nohara T, et al. Retroperitoneal fibrosis: a clinicopathologic study with respect to immunoglobulin G4. *Am J Surg Pathol*. 2009;33(12):1833–9.
14. Stone JH, Khosroshahi A, Hilgenberg A, Spooner A, Isselbacher EM, Stone JR. IgG4-related systemic disease and lymphoplasmacytic aortitis. *Arthritis Rheum*. 2009;60(10):3139–45.
15. Kamisawa T, Nakajima H, Egawa N, et al. IgG4-related sclerosing disease incorporating sclerosing pancreatitis, cholangitis, sialadenitis and retroperitoneal fibrosis with lymphadenopathy. *Pancreatology* 2006;6:132–7.
16. Kamisawa T, Funata N, Hayashi Y, Eishi Y, Koike M, Tsuruta K, et al. A new clinicopathological entity of IgG4-related autoimmune disease. *J Gastroenterol*. 2003;38(10):982–4.
17. Yamamoto M, Takahashi H, Hasebe K, Suzuki C, Naishiro Y, Hayashi T, et al. The analysis of interleukin-6 in patients with systemic IgG4-related plasmacytic syndrome-expansion of SIPS to the territory of Castleman's disease. *Rheumatology (Oxford)*. 2009;48:860–2.
18. Kamisawa T, Chen PY, Tu Y, Nakajima H, Egawa N, Tsuruta K, et al. Pancreatic cancer with a high serum IgG4 concentration. *World J Gastroenterol*. 2006;12(38):6225–8.
19. Joo M, Chang SH, Kim H, Gardner JM, Ro JY. Primary gastrointestinal clear cell sarcoma: report of 2 cases, one case associated with IgG4-related sclerosing disease, and review of literature. *Ann Diagn Pathol*. 2009;13(1):30–5.
20. Gill J, Angelo N, Yeong ML, McIvor N. Salivary duct carcinoma arising in IgG4-related autoimmune disease of the parotid gland. *Hum Pathol*. 2009;40(6):881–6.
21. Straub BK, Esposito I, Gotthardt D, Radeleff B, Antolovic D, Flechtenmacher C, et al. IgG4-associated cholangitis with cholangiocarcinoma. *Virchows Arch*. 2011;458(6):761–5.
22. Cheuk W, Yuen HK, Chan AC, Shih LY, Kuo TT, Ma MW, et al. Ocular adnexal lymphoma associated with IgG4+ chronic sclerosing dacryoadenitis: a previously undescribed complication of IgG4-related sclerosing disease. *Am J Surg Pathol*. 2008;32(8):1159–67.
23. Kubota T, Moritani S, Yoshino T, Nagai H, Terasaki H. Ocular adnexal marginal zone B cell lymphoma infiltrated by IgG4-positive plasma cells. *J Clin Pathol*. 2010;63(12):1059–65.
24. Yamamoto M, Takahashi H, Tabeya T, Suzuki C, Naishiro Y, Ishigami K, et al. Risk of malignancies in IgG4-related disease. *Mod Rheumatol* 2011; [Epub ahead of print]
25. Masaki Y, Sugai S, Umehara H. IgG4-related diseases including Mikulicz's disease and sclerosing pancreatitis: diagnostic insights. *J Rheum*. 2010;37:1380–5.
26. Okazaki K, Kawa S, Kamisawa T, Naruse S, Tanaka S, Nishimori I, et al. Clinical diagnostic criteria of autoimmune pancreatitis: revised proposal. *J Gastroenterol*. 2006;41(7):626–31.
27. Kawano M, Saeki T, Nakashima H, Nishi S, Yamaguchi Y, Hisano S, et al. Proposal for diagnostic criteria for IgG4-related kidney disease. *Clin Exp Nephrol* 2011; 15(5):615–26.
28. Hamano H, Kawa S, Horiuchi A, Unno H, Furuya N, Akamatsu T, et al. High serum IgG4 concentrations in patients with sclerosing pancreatitis. *N Engl J Med*. 2001;344:732–8.
29. Strehl JD, Hartmann A, Agaimy A. Numerous IgG4-positive plasma cells are ubiquitous in diverse localised non-specific chronic inflammatory conditions and need to be distinguished from IgG4-related systemic disorders. *J Clin Pathol*. 2011;64(3):237–43.
30. Yamamoto M, Takahashi H, Naishiro Y, Isshiki H, Ohara M, Suzuki C, et al. Mikulicz's disease and systemic IgG4-related plasmacytic syndrome (SIPS). *Nihon Rinsho Meneki Gakkai Kaishi*. 2008;31(1):1–8.
31. Okazaki K, Uchida K, Fukui T. Recent advances in autoimmune pancreatitis: concept, diagnosis, and pathogenesis. *J Gastroenterol*. 2008;43(6):409–18.
32. Takuma K, Kamisawa T, Igarashi Y. Autoimmune pancreatitis and IgG4-related sclerosing cholangitis. *Curr Opin Rheumatol*. 2011;23(1):80–7.
33. Fujinaga Y, Kadoya M, Kawa S, Hamano H, Ueda K, Momose M, et al. Characteristic findings in images of extra-pancreatic lesions associated with autoimmune pancreatitis. *Eur J Radiol*. 2010;76(2):228–38.

A novel clinical entity, IgG4-related disease (IgG4RD): general concept and details

Hisanori Umehara · Kazuichi Okazaki · Yasufumi Masaki · Mitsuhiro Kawano · Motohisa Yamamoto · Takako Saeki · Shoko Matsui · Takayuki Sumida · Tsuneyo Mimori · Yoshiya Tanaka · Kazuo Tsubota · Tadashi Yoshino · Shigeyuki Kawa · Ritsuro Suzuki · Tsutomu Takegami · Naohisa Tomosugi · Nozomu Kurose · Yasuhito Ishigaki · Atsushi Azumi · Masaru Kojima · Shigeo Nakamura · Dai Inoue · The Research Program for Intractable Disease by Ministry of Health, Labor and Welfare (MHLW) Japan G4 team

Received: 17 May 2011 / Accepted: 28 July 2011 / Published online: 1 September 2011
© Japan College of Rheumatology 2011

Abstract IgG4-related disease (IgG4RD) is a novel clinical disease entity characterized by elevated serum IgG4 concentration and tumefaction or tissue infiltration by IgG4-positive plasma cells. IgG4RD may be present in a certain proportion of patients with a wide variety of diseases, including Mikulicz's disease, autoimmune pancreatitis, hypophysitis, Riedel thyroiditis, interstitial pneumonitis, interstitial nephritis,

prostatitis, lymphadenopathy, retroperitoneal fibrosis, inflammatory aortic aneurysm, and inflammatory pseudotumor. Although IgG4RD forms a distinct, clinically independent disease category and is attracting strong attention as a new clinical entity, many questions and problems still remain to be elucidated, including its pathogenesis, the establishment of diagnostic criteria, and the role of IgG4. Here we describe the concept of IgG4RD and up-to-date information on this emerging disease entity.

H. Umehara and K. Okazaki contributed equally to this work.

Names of the collaborators of The Research Program for Intractable Disease by Ministry of Health, Labor and Welfare (MHLW) Japan G4 team are listed in Appendix.

H. Umehara (✉) · Y. Masaki
Division of Hematology and Immunology,
Department of Internal Medicine, Kanazawa
Medical University, 1-1 Daigaku, Uchinada-machi,
Kahoku-gun, 920-0293 Ishikawa, Japan
e-mail: umehara@kanazawa-med.ac.jp

K. Okazaki
Division of Gastroenterology and Hepatology,
The Third Department of Internal Medicine,
Kansai Medical University, Osaka, Japan

M. Kawano
Division of Rheumatology,
Department of Internal Medicine,
Kanazawa University Graduate School
of Medical Science, Ishikawa, Japan

M. Yamamoto
The First Department of Internal Medicine,
Sapporo Medical University, Sapporo,
Hokkaido, Japan

T. Saeki
Department of Internal Medicine,
Nagaoka Red Cross Hospital, Niigata, Japan

S. Matsui
Health Administration Center, Sugitani Campus,
University of Toyama, Toyama, Japan

T. Sumida
Department of Clinical Immunology, Doctoral Programs in
Clinical Science, Graduate School of Comprehensive Human
Science, University of Tsukuba, Ibaraki, Japan

T. Mimori
Department of Rheumatology and Clinical Immunology,
Kyoto University Graduate School of Medicine,
Kyoto, Japan

Y. Tanaka
First Department of Internal Medicine,
School of Medicine, University of Occupational and
Environmental Health,
Fukuoka, Japan

K. Tsubota
Department of Ophthalmology,
Keio University School of Medicine, Tokyo, Japan

Keywords IgG4-related diseases · Mikulicz's disease · Sjögren's syndrome · Autoimmune pancreatitis · Castleman's disease

Abbreviations

IgG4RD	IgG4-related disease
MD	Mikulicz's disease
SS	Sjögren's syndrome
MHLW Japan	Ministry of Health, Labor and Welfare Japan
LPSP	Lymphoplasmacytic sclerosing pancreatitis
AIP	Autoimmune pancreatitis
FMF	Familial multifocal fibrosclerosis
ANA	Anti-nuclear antibody

Introduction

In 1892, Dr. Johann von Mikulicz, also known as Jan Mikulicz-Radecki, published a paper describing a patient with symmetrical swelling of the lachrymal, parotid, and submandibular glands, with massive infiltration of these glands by mononuclear cells [1]. Following reports describing similar patients, this condition was called Mikulicz's disease (MD). In contrast, patients with similar symptoms, but with diseases such as leukemia, malignant lymphoma, and sarcoidosis, were reported to have

Mikulicz's syndrome [2]. In 1930, Dr. Henrik Sjögren, an ophthalmologist, published a paper describing a woman with rheumatoid arthritis accompanied by keratoconjunctivitis sicca and severe swelling of the parotid glands, a condition that has been recognized as Sjögren's syndrome (SS) [3]. In 1953, Morgan and Castleman examined 18 patients with MD and concluded that this condition is one manifestation of SS [4]. Since then, MD has attracted very little interest in western countries. In Japan, however, there have been many patients with MD, such that differences between MD and SS have been clarified [5–7]. For example, their gender distribution is quite different, in that MD occurs in both men and women, whereas SS occurs mainly in women. Second, patients with MD have relatively mild xerostomia and xerophthalmia, despite significant enlargement of their lachrymal and salivary glands. Further, MD is accompanied by more complications, such as autoimmune pancreatitis (AIP). Patients with MD show a better response to glucocorticoid therapy than patients with SS. Finally, it has become clear that MD is related to elevated serum IgG4 concentrations and infiltration of IgG4-positive cells [5–9].

Following the description of a patient with chronic pancreatitis due to an autoimmune mechanism [10], lymphoplasmacytic sclerosing pancreatitis (LPSP) was found to be a characteristic histopathological finding in patients with AIP [11]. These findings led to the concept of AIP, which has characteristics similar to those of other autoimmune diseases, such as hypergammaglobulinemia, the

T. Yoshino
Department of Pathology,
Okayama University Graduate School of Medicine,
Dentistry and Pharmaceutical Sciences,
Okayama, Japan

S. Kawa
Center for Health, Safety and Environmental Management,
Shinshu University, Nagano, Japan

R. Suzuki
Department of HSCT Data Management and Biostatistics,
Graduate School of Medicine,
Nagoya University, Nagoya, Japan

T. Takegami
Division of Molecular Oncology and Virology,
Medical Research Institute, Kanazawa Medical University,
Ishikawa, Japan

N. Tomosugi
Department of Advanced Medicine, Medical Research Institute,
Kanazawa Medical University, Ishikawa, Japan

N. Kurose
Department of Pathology and Laboratory Medicine,
Kanazawa Medical University, Ishikawa, Japan

Y. Ishigaki
Division of Core Facility, Medical Research Institute,
Kanazawa Medical University, Ishikawa, Japan

A. Azumi
Ophthalmology Department, Kobe Kaisei Hospital,
Hyogo, Japan

M. Kojima
Department of Anatomic and Diagnostic Pathology,
Dokkyo University School of Medicine, Mibu, Japan

S. Nakamura
Department of Pathology and Laboratory Medicine,
Nagoya University Hospital, Nagoya, Japan

D. Inoue
Department of Radiology,
Kanazawa University Graduate
School of Medical Science,
Ishikawa, Japan




Fig. 1 IgG4-related conditions. Many diseases have been reported to be IgG4-related

presence of various autoantibodies, lymphocytic infiltration into pancreatic tissue, and good responsiveness to steroids [12]. Following a report showing elevated serum IgG4 concentrations in patients with AIP [13], the pancreatic research team of the Ministry of Health, Labor and Welfare Japan (MHLW Japan) showed that AIP was related to IgG4 [14].

IgG4-positive plasma cell infiltration has also been observed in patients with other conditions, including retroperitoneal and mediastinal fibrosis [15, 16], inflammatory pseudotumor of the lung and liver [17], Küttner tumor [18], and interstitial nephritis [19], indicating that these diseases and conditions collectively constitute a new disease concept, IgG4-related disease (Fig. 1). These findings have led to the organization of two study groups by MHLW Japan to analyze the condition of IgG4-related disease. These groups consist of doctors and researchers in various fields, including rheumatology, hematology, gastroenterology, nephrology, pulmonology, ophthalmology, odontology, pathology, statistics, and basic and molecular immunology, from all over Japan. One of these groups, chaired by Professor Umehara of Kanazawa Medical University, is seeking to establish diagnostic criteria for IgG4-related multi-organ lymphoproliferative syndrome (IgG4-MOLPS), whereas the second group, chaired by Professor Okazaki of Kasai Medical University, is seeking to understand the etiology and pathogenesis of IgG4-related systemic disease.

Unification of different nomenclatures for IgG4-related disease (IgG4RD)

The concept of IgG4RD arose when elevated serum IgG4 concentrations were first reported in patients with sclerosing pancreatitis [13]. Autoimmune pancreatitis (AIP) is also

Table 1 Nomenclatures of IgG-related conditions

IgG4-related autoimmune disease	Kamisawa [21]
IgG4-associated multifocal systemic fibrosis	van der Vliet [76]
IgG4-related systemic disease	Kamisawa [20]
IgG4-related sclerosing disease	Kamisawa [15]
Hyper-IgG4 disease	Neild [59]
IgG4-related disease (IgG4-RD)	Zen [77]
Systemic IgG4 plasmacytic syndrome (SIPS)	Yamamoto [22]
IgG4-related multi-organ lymphoproliferative syndrome (IgG4-MOLPS)	Masaki [29]
IgG4-associated disease	Geyer [78]

associated with a variety of extrapancreatic lesions, including sclerosing cholangitis, sclerosing sialadenitis, and dacryoadenitis, resulting in the concept of IgG4-related systemic disease [20], also called IgG4-related autoimmune disease [21] or IgG4-related sclerosing disease [15]. The finding of elevated serum IgG4 and IgG4-positive plasma cell infiltration in MD suggested that MD was a systemic disease, which was called systemic IgG4 plasmacytic syndrome (SIPS) [22]. Further, a comparison of patients with MD and those with typical SS resulted in the formulation of a new clinical entity, IgG4+MOLPS [23]. Although many reports from Japan and other countries have described IgG4-related conditions under different names (Table 1), these may refer to the same condition, familial multifocal fibrosclerosis (FMF). Indeed, retroperitoneal fibrosis (RPF), mediastinal fibrosis, sclerosing cholangitis, Riedel's thyroiditis, and pseudotumor of the orbit may all be different manifestations of a single disease [24].

The name "IgG4-related sclerosing disease" is mainly based on the swelling of fibrous organs, such as the pancreas and retroperitoneum, whereas "SIPS" and "IgG4+MOLPS" are based on lymphoplasmacytic proliferation in glands and swollen lymph nodes without fibrosis. Although many patients with this condition (i.e., IgG4-related sclerosing disease, etc.) have lesions in several organs, either synchronously or metachronously, other patients show involvement of only a single organ. At this point, it is unclear whether the pathogenetic mechanism of this disease is systemic or whether it consists of manifestations in individual organs. In addition, several reports have described patients with IgG4-associated conditions concomitant with malignant tumors such as pancreatic [25, 26] and salivary [27] carcinomas, and ocular adnexal lymphoma [28]. Therefore, using the term 'systemic' may lead to an incorrect diagnosis of an IgG4-related condition in a patient with malignant




Fig. 2 Histopathology of IgG4-related disease (IgG4RD). IgG4RD is characterized histopathologically by the infiltration of IgG4-positive plasma cells and fibrosis. However, the severity of fibrosis is dependent

on the individual organs involved. For example, storiform fibrosis and obliterative phlebitis are characteristic of retroperitoneal lesions, but are very seldom observed in salivary glands ($\times 40$)


tumors in other organs. Based on these reasons, the members of the two MHLW Japan research teams agreed, at their second meeting in Kanazawa on February 11, 2010, to use the term “IgG4-related disease (IgG4RD)”.

General concept of IgG4RD

After the unification of the disease name as IgG4RD, both MHLW Japan research teams have sought to determine its pathogenesis and to formulate diagnostic criteria. The two teams reached a consensus that IgG4RD can occur in various organs, including the central nervous system, salivary glands, thyroid gland, lungs, pancreas, biliary duct, liver, gastrointestinal tract, kidneys, prostate gland, retroperitoneum, and lymph nodes, but that clinical symptoms depend on the location of the lesion. IgG4RD mainly affects middle-aged to elderly men. Its clinical symptoms are relatively mild, and the condition usually comes to clinical attention due to organ swelling or damage. Many patients with IgG4RD are treated effectively by steroid therapy. Although the infiltration of IgG4-positive cells and increased serum concentrations of IgG4 are characteristic of IgG4RD, the severity of fibrosis is dependent on the individual organs involved. For example, storiform fibrosis and obliterative phlebitis are characteristic of pancreatic, biliary tract, and retroperitoneal lesions, but are very seldom found in salivary glands or lymph nodes (Fig. 2).

Prevalence of IgG4RD

It is difficult to ascertain the number of patients with IgG4RD because its diagnostic criteria have not yet been established, the awareness of this disease is low, and its symptoms vary. An attempt was made to estimate the number of individuals with IgG4RD throughout Japan by



	KMU	KUH	total
2003	2	2	4
2004	0	1	1
2005	1	3	4
2006	1	3	4
2007	1	4	5
2008	1	3	4
2009	1	6	7
	7	22	29

Fig. 3 Prevalence of patients with IgG4RD. An attempt was made to estimate the number of individuals with IgG4RD throughout Japan by using as an example Ishikawa Prefecture (population 1.14 million people) with little population inflow/outflow. If all new patients with IgG4RD visit Kanazawa Medical University Hospital (KMU) or Kanazawa University Hospital (KUH), the incidence of this disease throughout Japan would be 0.28–1.08/100,000 population, with 336–1,300 patients newly diagnosed per year. If life expectancy after diagnosis is 20 years, then approximately 6,700–26,000 patients in Japan would have developed IgG4RD over the past 20 years. The numbers in the table represent the numbers of patients who visited KMU or KUH each year

using as an example Ishikawa Prefecture, which has a population of 1.14 million people with little population inflow/outflow (Fig. 3). In Ishikawa Prefecture, there are

two University Hospitals, Kanazawa Medical University Hospital (KMU) and Kanazawa University Hospital (KUH). Assuming that new patients with IgG4RD would visit one of these two hospitals, it was estimated that the incidence of this disease throughout Japan would be 0.28–1.08/100,000 population, with 336–1,300 patients newly diagnosed per year. Because the median age of onset of IgG4RD is 58 years and the clinical symptoms are relatively mild, with slow progression and good response to steroid therapy, life expectancy after diagnosis was estimated at 20 years. Thus, an estimated 6,700–26,000 individuals in Japan would have developed IgG4RD over the past 20 years.

Clinicopathological features of IgG4RD

Differences between IgG4-related MD and Sjögren's syndrome

Since elevated serum IgG4 was first reported in patients with MD [6], the members of the Japanese Society of Sjögren's Syndrome have assessed the clinical symptoms, laboratory findings, and detailed histopathology in patients with MD (characterized by symmetrical swelling of the lachrymal, submandibular, and parotid glands), nationwide, since 2004. Some patients did not show typical symptoms of MD such as swelling of the lachrymal, parotid, or submandibular glands but showed elevated serum IgG4 and other indices indicative of MD according to the criteria for the diagnosis of IgG4-related MD shown in Table 2 [8]. Sixty-four patients with MD or elevated serum IgG4 (>135 mg/dl) and characteristic histological findings were initially diagnosed with IgG4RD (formerly called IgG4+MOLPS) based on proposed guidelines for the diagnosis of IgG4RD (Table 3). A comparison of patients with IgG4RD and those with typical SS showed: (1) compared with SS patients, fewer patients with IgG4RD had symptoms of xerophthalmia, xerostomia, or arthralgia, whereas many had coexisting AIP, interstitial nephritis, allergic rhinitis, and/or bronchial asthma (Fig. 4a); (2) most patients with IgG4RD were negative for anti-SS-A and anti-SS-B antibodies, as well as for rheumatoid factor (RF) and anti-nuclear antibody (ANA) (Fig. 4b); (3) serum IgG4 and IgE concentrations were significantly higher in IgG4RD than in SS patients (Fig. 4c); and (4) steroid therapy was extremely effective in patients with IgG4RD but had limited effect in patients with SS [29].

The histopathological features of IgG4RD are unique, though both IgG4RD and SS show marked lymphocytic infiltration. IgG4RD is characterized by the formation of lymphoid follicles but lower levels of lymphocytic infiltration into the salivary ducts, such that their structure remains intact (Fig. 5a). Therefore, the absence of


Table 2 Diagnostic criteria of IgG4+ Mikulicz's disease [8] (approved by the Japanese Society for Sjögren's Syndrome 2008)

1. Symmetrical swelling of at least 2 pairs of lachrymal, parotid, or submandibular glands for at least 3 months
AND
2. Elevated serum IgG4 (>135 mg/dl)
OR
3. Histopathological features including lymphocyte and IgG4+ plasma cell infiltration (IgG4+ plasma cells/IgG+ plasma cells >50%) with typical tissue fibrosis or sclerosis
Differential diagnosis is necessary to distinguish IgG4+ Mikulicz's disease from other distinct disorders, including sarcoidosis, Castleman's disease, Wegener's granulomatosis, lymphoma, and cancer. The diagnostic criteria for Sjögren's syndrome (SS) may also include some patients with IgG4+ Mikulicz's disease; however, the clinicopathological conditions of patients with typical SS and IgG4+ Mikulicz's disease are different

Table 3 Guidelines for diagnosis of IgG4RD (proposed by the Research Program for Intractable Disease Ministry of Health, Labor and Welfare Japan, G4 team)

Clinical features highly suggestive of IgG4RD
1. Symmetrical swelling of lachrymal, parotid, or submandibular glands
2. Autoimmune pancreatitis
3. Inflammatory pseudotumor
4. Retroperitoneal fibrosis
5. Suspicion of Castleman's disease
Laboratory data highly suggestive of IgG4RD
1. Serum IgG4 >135 mg/dl
2. IgG4+ cells/IgG+ cells >40% in biopsy
Clinical features suggestive of IgG4RD
1. Unilateral swelling of at least one lachrymal, parotid, or submandibular gland
2. Orbital pseudotumor
3. Sclerosing cholangitis
4. Prostatitis
5. Hypertrophic pachymeningitis
6. Interstitial pneumonitis
7. Interstitial nephritis
8. Thyroiditis/hypo-function of thyroid
9. Hypophysitis
10. Inflammatory aneurysm
Laboratory data suggestive of IgG4RD
1. Hypergammaglobulinemia of unknown origin
2. Hypocomplementemia or existence of immune complex
3. Increase of IgE or eosinophils
4. Tumefactive lesions or lymph node swelling detected by gallium scan or fluoro-D-glucose positron emission tomography (FDG-PET)

Fig. 4 Comparison of clinical symptoms and laboratory findings in IgG4RD and typical Sjögren's syndrome (SS) [29]. **a** Clinical symptoms, **b** immunological findings, and **c** subclasses of immunoglobulins and IgG observed in patients with IgG4RD ($n = 61$) and typical SS ($n = 31$). Data are expressed as percentages. P values are for comparisons of IgG4RD with typical SS. Patients with typical SS fulfilled both Japanese and European SS criteria and were positive for both anti-SSA/Ro and anti-SSB/La antibodies



lymphoepithelial lesions in patients with IgG4RD, in contrast to SS, may explain the lower rate of dryness in the former, despite the marked swelling of lachrymal and


salivary glands. The most important difference between IgG4RD and SS is that the former is characterized by marked infiltration of IgG4-positive plasma cells, with a

Fig. 5 Histopathological findings of minor labial salivary gland biopsies in patients with IgG4RD and typical SS.

a Massive infiltration of lymphocytes and plasma cells was observed in patients with IgG4RD and those with typical SS ($\times 200$). IgG4RD, however, was characterized by lymphoid follicle formation but ducts were intact without lymphocytic infiltration. H&E staining.

b IgG4RD showed scattered IgG4+ plasma cells in the periphery of the follicles ($\times 200$), whereas typical SS showed few or no IgG4+ cells. IgG4 immunostaining.

c Staining for immunoglobulin κ - and λ -chains ($\times 200$)



ratio of IgG4-positive to IgG-positive cells of $>40\%$, a finding almost never seen in patients with SS (Fig. 5b). Moreover, most patients with IgG4RD show polyclonal B-cell proliferation, with equal staining for immunoglobulin κ - and λ -chains (Fig. 5c). Thus, despite their similarities in organ involvement, IgG4-MD and SS are quite different conditions, with distinct clinical and pathological characteristics [7–9, 22, 29–31].

IgG4-related Küttner tumor

Küttner tumor, a unilateral sclerosing sialadenitis, is an IgG4RD [18]. A common feature of MD and Küttner tumor is that both manifest sialadenitis, as in IgG4RD. Histologically, Küttner tumors are very severe fibrous sclerotic lesions containing IgG4-positive plasma cells [32]. In contrast, fibrosis tends to be less severe in MD, although

fibrosis in MD is frequently not examined extensively, because MD is generally diagnosed by the biopsy of minor labial salivary glands. Therefore, at present, it is difficult to set a strict boundary between MD and Küttner tumor.

IgG4-related autoimmune pancreatitis (IgG4-related AIP)

Recent studies have suggested that AIP manifests as two distinct subtypes, called types 1 and 2 (Table 4) [33]. Clinically, type 1 AIP seems to be the pancreatic manifestation of IgG4RD, characterized by: (1) mild abdominal symptoms, usually without acute attacks of pancreatitis; (2) occasional occurrence of obstructive jaundice; (3) increased serum gammaglobulin, IgG, and/or IgG4 concentrations; (4) presence of autoantibodies; (5) diffuse enlargement of the pancreas with a capsule-like low-density rim; (6)