

II. 研究成果の刊行に関する一覧表

研究成果の刊行に関する一覧表

雑誌

| 発表者氏名 | 論文タイトル名 | 発表誌名 | 巻号 | 頁 | 出版年 |
|--|--|---------------------------------------|----|-----------|------|
| Moriyuki Nakamura, Taisuke Fujibayashi, Akira Tominaga, Norifumi Satoh, Taketo Kawarai, Osamu Shinozuka, Haruo Watanabe, Tsuneyoshi Yamazaki, and <u>Hidenobu Sempuku.</u> | Hinokitiol inhibits <i>Candida albicans</i> adherence to oral epithelial cells. | Journal of Oral Biosciences | 52 | 42-50 | 2010 |
| Tadayoshi Arakawa, Takeshi Fujimaru, Tsutomu Ishizaki, Hiroaki Takeuchi, Masato Kageyama, Takuji Ikemi, Nobuhiro Hanada, Haruo Watanabe, and <u>Hidenobu Sempuku.</u> | Unique functions of hydroxyapatite to adherence of mutans streptococci. | Quintessence International. | 41 | e11-9 | 2010 |
| Yoshihide Nishiyama, Erika Inaba, Hiroshi Uematsu and <u>Hidenobu Sempuku.</u> | Effects of mucosal care on oral pathogens in professional oral hygiene to the elderly. | Archives of Gerontology & Geriatrics. | 51 | e139-e143 | 2010 |
| <u>Hidenobu Sempuku,</u> Hideo Miyazaki, Saori Yoneda, Akihiro Yoshihara, and Akio Tada. | A quick statistically accurate diagnosis for caries risk in the elderly. | Clinical Laboratory | 56 | 505-512 | 2010 |

| | | | | | |
|---|---|-----------------------------|-----|---------|------|
| 泉福英信 | 歯科医療機関における院内感染対策の導入について | 日本歯科評論 | 774 | 135-140 | 2007 |
| Kentaro Okuda, Nobuhiro Hanada, Yoshie Usui, Hiroaki Takeuchi, Hidehiko Koba, Ryoma Nakao, Haruo Watanabe, and <u>Hidenobu Senpuku</u> | Inhibition of <i>Streptococcus mutans</i> adherence and biofilm formation using analogues of the SspB peptide. | Archives of Oral Biology | 55 | 754-762 | 2010 |
| 泉福英信 | 多剤性耐性菌アシネトバクター &NDM-1 の歯科医療における感染防止対策 | Dental Diamond | 11 | 192-196 | 2010 |
| <u>狩山玲子、公文裕巳</u> | バイオフィルム形成とクオラムセンシング 臨床医学編 2. 泌尿器感染症とバイオフィルム. | 化学療法の領域 | 26 | 71-78 | 2010 |
| Sugi N, Naruishi K, Kudo C, Hisaeda-Kako A, Kono T, Maeda H, <u>Takashiba S.</u> | Prognosis of periodontitis recurrence after intensive periodontal treatment using examination of serum IgG antibody titer against periodontal bacteria. | J Clin Lab Anal | 25 | 25-32 | 2011 |
| Koide Y, Maeda H, Yamabe K, Naruishi K, Yamamoto T, Kokeguchi S, <u>Takashiba S.</u> | Rapid detection of <i>mecA</i> and <i>spa</i> by the loop-mediated isothermal amplification (LAMP) method. | Lett Appl Microbiol | 50 | 386-392 | 2010 |
| Soga Y, Sugiura Y, Takahashi K, Nishimoto H, Maeda Y, Tanimoto M, <u>Takashiba S.</u> | Progress of oral care and reduction of oral mucositis—a pilot study in a hematopoietic stem cell transplantation ward. | Support Care Cancer | 19 | 303-307 | 2011 |

| | | | | | |
|---|--|--------------------------|------|----------|------|
| Soga Y, Maeda Y, Ishimaru F, Tanimoto M, Maeda H, Nishimura F, Takashiba S | Bacterial substitution of coagulase-negative staphylococci for streptococci on the oral mucosa after hematopoietic cell transplantation. | Support Care Cancer | 19 | 995-1000 | 2011 |
| 金中章江, 岩田宏隆, 大谷久美, 森本祥代, 前田知子, 井本有香, 塩見千尋, 長島義之, 高柴正悟 | 介護高齢者に対するのチームアプローチ 口腔機能の向上から栄養状態の改善を目指して | 感染防止 | 20 | 14-22 | 2010 |
| Yamabe K, Maeda H, Kokeguchi S , Soga Y, Meguro M, Naruishi K, Asakawa S, Takashiba S | Antigenic group II chaperonin in <i>Methanobrevibacter oralis</i> may cross-react with human chaperonin CCT. | Mol Oral Microbiol. | 25 | 112-122 | 2010 |
| 佐藤法仁, 渡辺朱理, 苔口 進 | 医療従事者の身だしなみに関する研究-医師・歯科医師の身だしなみに対する不快感調査 | 日本医事新報 | 4498 | 95-98 | 2010 |
| 山部こころ, 苔口 進 , 前田博史 | メタン生成古細菌は歯周病の病原因子? | 化学と生物 | 48 | 463-470 | 2010 |
| 苔口 進 , 山部こころ, 前田博史 | メタン産生古細菌 <i>Methanobrevibacter</i> の口腔疾患への関わり | 日本臨床腸内微生物学会誌 | 12 | 17-25 | 2010 |
| 渡辺朱理, 佐藤法仁, 苔口 進 | 歯科衛生士学校生における市中感染型メチシリン耐性ブドウ球菌の保菌調査を通しての感染予防対策教育の向上 | 日本歯科衛生学会雑誌 | 5 | 69-76 | 2011 |
| Akio Tada, Hidenobu Senpuku | Attitudes towards HIV-infected patients, knowledge related to HIV/Universal precautions, and infection control practices of Japanese dentists. | Journal of Dental Health | 61 | 273-281 | 2011 |

| | | | | | |
|--|---|-------------------------|-----|-----------|------|
| Naoki Narisawa, Taketo Kawarai, Naomi Suzuki, Yutaka Sato, Kuniyasu Ochiai, Makoto Ohnishi, Haruo Watanabe, and <u>Hidenobu Senpuku</u> | Competence-dependent endogenous DNA rearrangement and uptake of extracellular DNA gives a natural variant of <i>Streptococcus mutans</i> without biofilm formation. | Journal of Bacteriology | 193 | 5147-5154 | 2011 |
| Ryoma Nakao, Hideki Hasegawa, Kuniyasu Ochiai, Shogo Takashiba, Akira Ainai, Makoto Ohnishi, Haruo Watanabe, and <u>Hidenobu Senpuku</u> | Outer membrane vesicles of <i>Porphyromonas gingivalis</i> elicit a mucosal immune response | PLoS ONE | 6 | e26163 | 2011 |
| Tatsuro Ito, Takahide Maeda, and <u>Hidenobu Senpuku</u> | Roles of salivary components in <i>Streptococcus mutans</i> colonization in a new animal model using NOD/SCID. <i>e2f1</i> ^{-/-} mice. | PLoS ONE | 7 | e32063 | 2011 |
| 泉福英信 | 歯科領域と口腔感染症について | バムサジャーナル | 23 | 3-6 | 2011 |
| 小澤寿子, 中野雅子, 木村泰子, 新井 高 | 歯科用ユニット給水管路の新クリーンシステムの評価 | 緑膿菌感染症研究会講演記録 | 54 | 193-200 | 2011 |
| 狩山玲子、金原和秀、高野和潔、妹尾典久、大森啓士、光畑律子、桐田泰三、 <u>公文裕巳</u> | 緑膿菌性バイオフィルム形成阻害剤のスクリーニングにおける新規マイクロデバイスの有用性について | 日本歯科衛生学会雑誌 | 45 | 84-88 | 2011 |
| <u>佐藤法仁</u> | シリーズ院内感染対策塾 第3回 滅菌・消毒・洗浄の基礎(1) | ZOOM UP | 133 | 32-33 | 2011 |

| | | | | | |
|------|------------------------------------|----------|-----|-------|------|
| 佐藤法仁 | シリーズ院内感染対策塾 第4回 滅菌・消毒・洗浄の基礎 (2) | ZOOM UP | 134 | 34-35 | 2012 |
| 佐藤法仁 | 感染制御学ノート vol.1 感染制御と微生物 | DH style | 6 | 14-15 | 2012 |
| 佐藤法仁 | 感染制御学ノート vol.2 細菌 | DH style | 67 | 14-15 | 2012 |
| 佐藤法仁 | 感染制御学ノート vol.3 ウイルス | DH style | 68 | 14-16 | 2012 |

III. 研究成果の刊行物・別刷

ORIGINAL

Hinokitiol Inhibits *Candida albicans* Adherence to Oral Epithelial Cells

Moriyuki Nakamura^{1,4)}, Taisuke Fujibayashi^{1,4)}, Akira Tominaga^{2,4)}, Norifumi Satoh^{3,4)},
Taketo Kawarai⁴⁾, Osamu Shinozuka¹⁾, Haruo Watanabe⁴⁾, Tsuneyoshi Yamazaki¹⁾
and Hidenobu Senpuku⁴⁾ §

¹⁾ Dentistry for Persons with Disabilities, Tokyo Medical & Dental University

²⁾ Department of Oral Surgery, Tokyo Medical University

³⁾ EN Otsuka Pharmaceutical Co., Ltd.

⁴⁾ Department of Bacteriology, National Institute of Infectious Diseases

¹⁾ 1-5-45 Yushima, Bunkyo-ku Tokyo 113-8549, Japan

²⁾ 6-7-1 Nishishinjuku, Shinjuku-ku, Tokyo 160-0023, Japan

³⁾ 2-1-20, Kandasurugadai, Chiyoda-ku, Tokyo 101-0062, Japan

⁴⁾ 1-23-1, Toyama, Shinjuku-ku, Tokyo 162-8640, Japan

[Received on July 27, 2009 ; Accepted on November 25, 2009]

Key words : *Candida albicans* / epithelial cell / hinokitiol / cell adherence

Abstract : *Candida* spp. are an opportunistic pathogen causing serious local and systemic infections, especially in immuno-compromised hosts such as the elderly and HIV-positive patients. Hinokitiol C₁₀H₁₂O₂ (β -thujaplicin) is a component of the essential oils isolated from Cupressaceae and shows antibacterial activities for various bacteria. The aim of this study was to demonstrate the preventive effects of hinokitiol on the adherence of seven species of *Candida* to oral epithelial cells and to establish a safe and useful method for oral hygiene. A short-time treatment (30 min) with 0.25 mM hinokitiol showed 30–70% inhibition of adherence of *Candida* spp. to oral epithelial cells, inhibited about 11% biofilm formation, and did not inhibit the cell growth of *Candida* spp. Furthermore, short treatment with 0.25 mM hinokitiol was a safe method for oral hygiene against *Candida* infection because it did not inhibit the cell growth of commensal bacteria, such as oral streptococci existing in normal flora, or damage the epithelial cells. However, long-time treatment and a high concentration of hinokitiol demonstrated both the adherence inhibition of *Candida* and damage to commensal bacteria and epithelial cells. Our data suggest an appropriate procedure to apply hinokitiol that may be beneficial for the prevention of opportunistic pathogens such as *Candida* spp. in the oral cavity. The clinical and daily use of hinokitiol under an appropriate procedure may be a preventive and realistic therapy for *Candida* infection in immune-compromised hosts.

Introduction

Candida spp. are human commensals and can par-

ticipate in biofilm formation on mucosal and cutaneous surfaces as well as on device surfaces, *e. g.*, dentures and catheters^{1–7)}. Many manifestations of candidiasis are associated with the formation of biofilms on host tissues (*e. g.*, oral thrush) and on indwelling medical devices (central venous catheter-associated candidemia)⁸⁾. Biofilms are a community of microor-

§ Corresponding author

E-mail : hsenpuku@nih.go.jp

ganisms attached to a surface and surrounded by an extracellular matrix. A clinically significant characteristic of microbial biofilms is their enhanced resistance to antimicrobial agents^{9,10}. Anti-fungal agents include ketoconazole, fluconazole, itraconazole, ravuconazole, miconazole, amphotericin B, nystatin, chlorhexidine, hydrogen peroxide and povidone-iodine. Long-term treatment of oropharyngeal candidiasis with anti-fungal therapy, such as fluconazole, itraconazole, and ketoconazole, sometimes leads to the emergence of drug-resistant *Candida albicans*¹¹. Chlorhexidine, hydrogen peroxide and povidone-iodine are potent anti-microbial and chemical agents that reduce various pathogens. Their clinical application is also limited because they induce various adverse reactions, such as anaphylactic shock^{12,13}, damage to human tissues, and destruction of normal beneficial flora including the oral streptococci that exist in high concentrations in the oral cavity.

C. albicans, one of the *Candida* spp., is the primary causative agent of candidiasis, among the most important nosocomial infections of humans. The microorganism is an opportunistic pathogen that can cause serious local as well as systemic infections, especially in immuno-compromised hosts such as the elderly and HIV-positive patients¹⁴. *C. albicans* is a pleomorphic fungus having the ability to transition between yeast cell and filamentous forms that is essential for virulence; and therefore constitutes a central component in *C. albicans* pathogenesis. Various models have been established to study candidiasis: *Candida*-epithelial cell interactions using immortalized human epithelial cell lines in tissue culture¹⁵⁻¹⁷ where *C. albicans* adheres to the epithelial cells of the oral mucosa and also invades these cells. Invasion into the epithelial cell limit of the oral mucosa is characteristic of both human and experimental animal models of oropharyngeal candidiasis¹⁸⁻²¹.

Hinokitiol C₁₀H₁₂O₂ (β -thujaplicin) is a component of the essential oils isolated from Cupressaceae and shows antibacterial activities against various bacteria: MRSA, Legionella, colon bacillus and oral bacteria²²⁻²⁷. It is reported as being used in Japan as a preservative for flowers, mushrooms and vegetables,

a plant growth stimulator (inhibit bacteria), and anti-bacterial or fungicidal additive in food, cosmetics, toothpaste, hair tonics and tick repellents²⁸. However, the clinical application of hinokitiol may be limited because it has anti-fungal and broad spectrum anti-bacterial activities, and killing activities of human tissues. Conditioning studies for the preventive effects of hinokitiol on *Candida* infection and biofilm formation are needed and the aim here was to demonstrate the preventive and safe effects of hinokitiol on adherence in seven species of *Candida* on oral epithelial cells. Our data indicate the beneficial effects of hinokitiol in preventing *Candida* from colonizing human tissues. The clinical use of hinokitiol may be considered for use in preventive therapy for *Candida* infections.

Materials and Methods

1. Strains and culture conditions

The strains used were *C. albicans* SC5314, *Candida tropicalis* IFO0618, *Candida dubliniensis* CD36, *Candida dubliniensis* CD57, *Candida parapsilosis* ATCC 22019, *Candida parapsilosis* FRCP-0201, *Candida glabrata* 850821, *Streptococcus mutans* MT6229, *S. mutans* OMZ175, *Streptococcus sobrinus* 6715, *Streptococcus salivarius* TCM5707, *Streptococcus sanguinis* ATCC 10556 and *Streptococcus mitis* ATCC 6249. *Candida* spp. were provided by Dr. Masakazu Niimi of the National Institute of Infectious Diseases. *Candida* spp. were incubated for 24 h at 37°C in Yeast Peptone Dextrose (YPD; 2% Bacto peptone, 2% dextrose and 1% yeast extract) broth. Streptococci were grown in Brain Heart Infusion broth (BHI; Difco Laboratory, Detroit, MI, USA) in an aerobic atmosphere of 5% CO₂, 75% N₂ and 20% O₂ (GasPack CO₂; Mitsubishi, Tokyo) at 37°C for 24 h before beginning each experiment.

2. Effects of hinokitiol on growth

Cell suspensions 1 × 10⁸ CFU/mL of *Candida* strains and various concentrations (0.0625, 0.125, 0.25 and 0.5 mM) of hinokitiol (Wako Pure Chemical Industries Ltd., Osaka, Japan) were mixed and incubated in YPD for 24 h at 37°C using aerobic conditions. Cell

growth was measured as absorbance at 660 nm at 0, 4, 6, 8, 10 and 24 h during incubation. Treatment assays were performed independently three times.

3. Effects on adherence and cytotoxic activity on epithelial cells of hinokitiol

C. albicans SC5314 1×10^8 CFU were treated using 0.25 mM hinokitiol for 0, 1, 2, 5, 10, 20, 30 and 60 min at 37°C in 1 mL sterile phosphate-buffered saline (PBS), and washed using sterile PBS two times. Pellets after the last centrifugation were suspended in 1 mL sterile PBS, added to a monolayer of Ca9-22 epithelium cells in 24-well polystyrene plates that contained 70–90% confluent, and incubated for 24 h at 37°C under aerobic conditions. Cells attached to *Candida* were washed using sterile PBS three times and detached using 0.05% trypsin-EDTA. The detached cell suspensions were collected in 0.05% trypsin-EDTA using pipetting, and plated on YPD using an EDDY JET spiral plating system (IUL, S. A., Barcelona, Spain).

Short-treatment effect of hinokitiol on *C. albicans* viability was also tested. After treatment with 0.25 mM hinokitiol for 0, 1, 2, 5, 10, 20, 30 and 60 min and washing with sterile PBS two times, they were then poured onto a YPD agar plate using an EDDY JET spiral plating system and incubated for 24 h at 37°C under aerobic conditions.

Other *Candida* strains at 1×10^8 CFU were also treated using various concentrations of hinokitiol : 0, 0.0625, 0.125, 0.25, 0.5, 1.0 and 5.0 mM for 30 min in PBS at 37°C, and added to a monolayer of Ca9-22 epithelium cells in 24-well polystyrene plates, washed with sterile PBS two times, and incubated for 30 min at 37°C. The effects of hinokitiol on *Candida* species viabilities were also observed in a previously described method. Moreover, it was tested whether hinokitiol has effects on the viability of commensal bacteria, such as oral streptococci, *S. mutans* MT6229, *S. mutans* OMZ175, *S. sobrinus* 6715, *S. salivarius* TCM5707, *S. sanguinis* ATCC 10556 and *S. mitis* ATCC 6249. They were treated with 0.25 mM or 1.0 mM for 30 min and poured onto a Mitis-Salivarius agar plate (MS, Becton/Dickinson, Sparks, MD, USA).

After incubation for 24 h at 37°C under aerobic or 5% CO₂ aerobic conditions (Gaspack CO₂; Becton/Deckinson), the colonies of *Candida* or streptococci in all experiments were counted on the YPD or MS agar plate and the colony number rates of tested organisms were calculated in comparison with the control : non-treated organisms (100%) or time zero.

4. Biofilm formation

C. albicans SC5314 incubated for 24 h at 37°C in YPD broth was adjusted to OD=0.5 at 660 nm ; harvested using centrifugation ; and washed in sterile PBS two times. *C. albicans* suspensions were diluted using RPMI1640, treated using various concentrations (0, 0.125, 0.25, 0.5, 1.0 and 5.0 mM) of hinokitiol, and added to 96-well microtiter plates. Chemically defined RPMI1640 medium containing minimal (0.2%) glucose was used as a nutrient-poor condition for the biofilm formation assay. After incubation for 24 h at 37°C, biofilms formed in wells were washed using sterile PBS two times. Biofilm formation was tested using the XTT assay at 492 nm. XTT reduction has been widely used to measure biofilm activity and allows the detection of small differences in metabolic activity between strains^{29–31}.

5. Epithelial cell monolayer damage assay

Epithelial cell (monolayer cells of Ca9-22) damage caused by hinokitiol was determined by the release of lactate dehydrogenase (LDH) into the medium using a Cytotoxicity Detection Kit-LDH (Roche Diagnostics K. K., Tokyo). The assays were performed according to the manufacturer's instructions and the measurements were performed in triplicate.

6. Statistical analysis

All data were analyzed using the Statistical Package for SPSS for Windows (version 100 ; Chicago, IL, USA). Student's *t*-test with the Bonferroni method was used to compare control data and the data of *Candida* spp. or streptococci after incubation. *p*-Values less than 0.05 were considered to be significant.




Fig. 1 Long-term treatment using hinokitiol on *Candida* growth

Various *Candida* strains were incubated in YPD broth containing each concentration (0, 0.0625, 0.125, 0.25 or 0.5 mM) of hinokitiol. Absorbance at 660 nm was measured at 0, 4, 6, 8, 10, and 24 h incubation with *C. albicans* SC5314, *C. tropicalis* IFO0618, *C. dubliniensis* CD36 and CD57 suspensions. Results are the mean \pm standard deviation of three independent experiments, each performed using triplicate assays.

Results

1. Effects of hinokitiol on *Candida* growth and adherence

At first, to clarify the effects of hinokitiol on *Candida* growth, various concentrations of hinokitiol were added for 24 h in the growth of *Candida* species. The 0 and 0.0625 mM hinokitiol did not inhibit the growth of *C. albicans* SC5314, *C. tropicalis* IFO0618 and *C. dubliniensis* CD36 and CD57, whereas 0.125 mM hinokitiol inhibited growth intermediately, and 0.25 and 0.5 mM hinokitiol completely inhibited fungal cell growth of all *Candida* strains at all measurement times (Fig. 1). Similar inhibition data were also observed for other *Candida* species (*C. parapsilosis* ATCC 22019, *C. parapsilosis* FRCP-0201 and *C. glabrata* 850821) (data not shown). If hinokitiol is used in the oral cavity, long-time application may damage oral flora and epithelial cells. Short-term treatment of hinokitiol is better for application in the oral cavity than long-time treatment. Therefore, short-term (1, 2, 5, 10, 20, 30 and 60 min) treatment with 0.25 mM hinokitiol was performed in the *C. albicans* viability experiment. Inhibition effects were not

observed at all treatment times (Fig. 2). Further, it was tested whether short-term treatment inhibited the adherence of *C. albicans* to a monolayer of Ca9-22 epithelial cells. Inhibition effects were significantly observed at 10, 20, 30 and 60 min using 0.25 mM hinokitiol; further, the inhibition levels increased in a treatment time-dependent manner (Fig. 2). It was also observed whether short-time treatment damaged epithelial cells. Cytotoxic activities (LDH activities) against epithelial cells were less than 2% with treatment of 0.25 mM hinokitiol for less than 30 min and increased slightly with the treatment time (60 min) in comparison with other treatment times (Fig. 3). Taken together, treatment for 30 min using 0.25 mM hinokitiol is appropriate for not only inhibition of the adherence of *C. albicans* but also safeguarding epithelial cells in the oral cavity.

Short-term treatment effects of hinokitiol were also tested with the adherence of other *Candida* species on monolayers of oral epithelial cells. To check the viability of *Candida* after treatment, the treated cells were immediately poured onto a YPD agar plate, and colony counts were performed after overnight incubation at 37°C. Short-term treatment (30 min) of 0.125–0.25 mM hinokitiol inhibited adherence of




Fig. 2 Short-term treatment of hinokitiol on *Candida* growth and attachment

C. albicans SC9314 were pretreated using 0.25 mM hinokitiol for 1, 2, 5, 10, 20, 30 and 60 min. Pretreated *Candida* were poured onto a YPD agar plate and applied on Ca9-22 monolayer cells. The *C. albicans* attached to Ca9-22 cells were also poured onto a YPD agar plate. The colony-forming units (CFU) were counted and the percent of hinokitiol-treated *C. albicans* CFU was recalculated as 100% to control data (non-treatment). Results are the mean \pm standard deviation of three independent experiments, each performed using triplicate assays; and compared to the control (0 mM hinokitiol) (*: $p < 0.05$, **: $p < 0.01$).

Candida spp. to Ca9-22 cells by 30–70%, but showed little effect on the cell numbers of *Candida* (Fig. 4). More than 1 mM hinokitiol markedly inhibited both the adherence and cell number of *Candida* spp. Treatment using 0.5 mM hinokitiol inhibited adherence by more than 60% and the cell number of *Candida* spp. by about 20%, except for *C. grabrata* 850821. Short-treatment using 1 mM hinokitiol did not show a different appearance of the cell surface from non-treated *C. albicans* using electron microscope observation (data not shown).

2. Effects of hinokitiol on *C. albicans* biofilm formation and streptococci viability

To determine the effects of hinokitiol on *Candida* biofilm formation, various concentrations of hinokitiol




Fig. 3 Short-term treatment of hinokitiol on YPD monolayer cells

YPD monolayer cells were treated using 0.25 mM hinokitiol for 1, 2, 5, 10, 20, 30 and 60 min. Cell damage caused by hinokitiol was determined by the release of lactate dehydrogenase (LDH) into the medium. The assays were performed according to the manufacturer's instructions and the measurements were performed in triplicate. To measure epithelial cell damage, the following calculation was used: $(E_{Ca} - C1 - C2) / (100L - C1) \times 100 = \text{relative cytotoxicity (\%)}$. Absorbance measured at OD 490–600 directly correlates with LDH activity. E_{Ca}, epithelial cells treated with hinokitiol; C1, control 1, non-treated epithelial cells; C2, control 2, only *C. albicans*; 100L, 100% lysis (0.2% Triton-X 100). Controls 1, 2 and 100% lysis were determined individually for each treatment.

were applied during biofilm formation with *C. albicans*. Hinokitiol inhibited biofilm formation in a dose-dependent manner (Fig. 5); however, treatment with 0.25 mM hinokitiol, inhibiting the adherence of *Candida* species on oral epithelial cells by 30–70%, inhibited biofilm formation by about 11%. Treatment with 0.5 mM and 1.0 mM hinokitiol inhibited biofilm formation by about 45% and 75%. Therefore, 0.25 mM hinokitiol has significant ability to inhibit both adherence and biofilm formation; but no bactericidal effects against *Candida* species were observed. The effect of using 0.25 mM hinokitiol is a mild treatment for *Candida* infection. To determine the effects on commensal bacteria of hinokitiol in the oral cavity, 0.25 mM and 1.0 mM hinokitiol were applied to vari-




Fig. 4 Short-term treatment of hinokitiol on *Candida* adherence

Various *Candida* strains were pretreated using each concentration (0, 0.0625, 0.125, 0.25, 0.5, 1.0 and 5.0 mM) of hinokitiol for 30 min. Pretreated *Candida* strains were applied to YPD or Ca9-22 monolayer cells. Ca9-22 cells attached to a *Candida* strain were collected in 0.05% trypsin-EDTA and poured onto the YPD agar plate. Colony formation on YPD was counted 24 h after incubation with *C. albicans* SC5314, *C. tropicalis* IFO0618, *C. dubliniensis* CD57, *C. parapsilosis* ATCC22019, *C. glabrata* 850821, *C. parapsilosis* FRCP-0201 suspensions. Percent of hinokitiol-treated *Candida* CFU was calculated in comparison to non-treated *Candida* (100%). Results are the mean \pm standard deviation of three independent experiments each performed using triplicate assays; and compared to the control (0 mM hinokitiol) (*: $p < 0.05$, **: $p < 0.01$).

ous oral streptococci. Treatment of 1.0 mM inhibited all streptococci, whereas treatment with 0.25 mM did not significantly inhibit the viability of streptococci (Fig. 6).

Discussion

Here we show that *Candida* adhesion to oral epithelial cells was reduced with lower concentrations (0.0625–0.25 mM) of hinokitiol. Short 30-min treatment with 0.25 mM hinokitiol is safe, and may be useful for the prevention of *Candida* infections in human tissues. Further, 0.25 mM hinokitiol may not destroy normal fungal and bacterial flora or damage epithelial cells. In contrast, at higher concentrations (>0.5 mM), hinokitiol prevented adhesion and biofilm for-

mation and also prevented the cell viability of *Candida* spp. and oral commensal streptococci. Higher concentrations of hinokitiol have anti-fungal and -microbial broad spectrum activities and may have side effects allowing damage to human tissues and microbial substitution of the normal microflora, which causes infection by opportunistic pathogens^{32,33}. Therefore, daily oral care with 0.25 mM hinokitiol is safe and may be effective for continuing prevention of *Candida* infections on human tissues.

Several studies concerning *C. albicans* biofilms have examined the activities of CHX, ethanol, hydrogen peroxide, betadine, and povidone-iodine; however, the findings are not consistent^{10,34–36}. CHX was uniformly effective against strains of common borne microorganisms, but was disqualified for clinical use




Fig. 5 Effects of hinokitiol on *Candida* biofilm formation

C. albicans SC5314 was incubated in chemically defined RPMI1640 medium containing each concentration (0, 0.125, 0.25, 0.5, 1 and 5 mM) of hinokitiol for 24 h. Biofilm formation was tested using the XTT assay. Absorbance at 492 nm was measured 24 h after incubation with *C. albicans*. Results are the mean \pm standard deviation of three independent experiments each performed using triplicate assays; and are compared to the control (non-treatment) (* : $p < 0.05$, ** : $p < 0.01$).

because it was highly cytotoxic³⁷). Sanchez *et al.* investigated the effectiveness of using 30-min exposure to dilutions of CHX in PBS on canine embryonic fibroblasts and on cultures of *S. aureus*. They found that the antiseptics tested were lethal to canine fibroblasts at bactericidal concentrations *in vitro*³⁸). Hydrogen peroxide at concentrations of 3% and 5% was the most effective to reduce the biofilm density of *C. albicans* and *Staphylococcus epidermidis* in the elimination of biofilms and killing fungi and bacteria^{39,40}); however, a drawback of hydrogen peroxide is its toxicity and irritation to the skin or tissue when used for longer contact periods⁴¹). Thus, several reports have proposed the cytotoxic effects of anti-fungal and microbiological agents on human cells.

Hinokitiol has cytotoxic effects on human cell lines. Yamano *et al.* evaluated the toxic effect of hinokitiol on Hela 229 cells⁴²). Although no decrease in activity was observed at hinokitiol concentrations of less than 1 $\mu\text{g}/\text{mL}$, activity decreased to 50% of the control value with long exposure (over 72 h) to 32 $\mu\text{g}/\text{mL}$ hinokitiol; no morphological changes were apparent




Fig. 6 Short-term treatment of hinokitiol on streptococci growth

S. mutans MT6229, *S. mutans* OMZ175, *S. sobrinus* 6715, *S. salivarius* TCM5707, *S. sanguinis* ATCC 10556 and *S. mitis* ATCC 6249 were treated with each concentration (0, 0.25 or 1.0 mM) of hinokitiol. The treated strains were then poured onto an MS agar plate. After incubation for 24 h, the colonies of streptococci were counted. Percent of hinokitiol-treated streptococci CFU was calculated in comparison with non-treated streptococci (100%). Results are the mean \pm standard deviation of three independent experiments each performed using triplicate assays, and compared to the control (0 mM hinokitiol) (* : $p < 0.05$, ** : $p < 0.01$).

by Giemsa Staining at 32 $\mu\text{g}/\text{mL}$. The agent must be effective against bacteria within a short exposure time without harming host cells or tissues to use in the oral cavity³³). Here we show that short treatment (5–30 min) at 0.25 mM (42.05 $\mu\text{g}/\text{mL}$) hinokitiol, a higher concentration than 32 $\mu\text{g}/\text{mL}$, significantly inhibited *Candida* adherence, and poor cytotoxicity to the monolayer Ca9-22 cells (Fig. 2). In particular, 30-min treatment was a convenient or appropriate time for treatment using 0.25 mM hinokitiol (Figs. 2, 3).

Taken together, our data show that it is important for anti-fungal and anti-bacterial agents to be used at the appropriate concentrations in practice, at the correct concentrations, and for adequate contact times⁴³). In this study, short treatment using 0.25 mM hinokitiol showed a preventive effect on *Candida* adherence to oral epithelial cells rather than an anti-fungal effect, and this concentration prevented *Candida* biofilm formation. The clinical and daily use of hinokitiol under the appropriate procedure may be a preventive and realistic therapy for *Candida* infection in

immune-compromised hosts. However, a definitive conclusion regarding the mechanism of the inhibition effect using short treatment of 0.25 mM on the adherence of *Candida* spp. to the monolayer of epithelial cells requires further investigation, changing interactions between the surface adhesins of *Candida* and receptors on epithelial cells.


Acknowledgements

The authors thank Naoki Narisawa and Saori Yoneda for their technical support, helpful discussions, and advice. This work was supported in part by a grant-in-aid for the Development of Scientific Research (15390571, 19659559 and 18592011) from the Ministry of Education, Science, and Culture of Japan ; and by a grant from the Ministry of Health, Labor and Welfare (H16-Medical Services-014 and H19-Medical Services-007).

References

- 1) Douglass, L. J. : *Candida* biofilms and their role in infection. Trends Microbiol. **11** : 30—36, 2003.
- 2) Douglass, L. J. : Medical importance of biofilms in *Candida* infections. Rev. Iberoam. Micol. **19** : 139—143, 2002.
- 3) Kumamoto, C. A. : *Candida* biofilms. Curr. Opin. Microbiol. **5** : 608—611, 2002.
- 4) Kawamura-Sato, K., Wachino, J., Kondo, T., Ito, H. and Arakawa, Y. : Reduction of disinfectant bactericidal activities in clinically isolated *Actinobacter* species in the presence of organic material. J. Antimicrob. Chemother. **61** : 568—576, 2008.
- 5) Lopez-Ribot, J. L., McAtee, R. K., Perea, S., Kirkpatrick, W. P., Rinalci, M. G. and Petterson, T. F. : Multiple resistance phenotypes of *Candida albicans* coexist during episodes of oropharyngeal candidiasis in human immunodeficiency virus-infected patients. Antimicrob. Agents Chemother. **43** : 1621—1630, 1999.
- 6) Mukherjee, P. K. and Clandra, J. : *Candida* biofilm resistance. Drug Resist. Updat. **7** : 301—309, 2004.
- 7) Ramage, G., Wickes, B. L. and Lopez-Ribot, J. L. : Biofilms of *Candida albicans* and their associated resistance to antifungal agents. Am. Clin. Lab. **20** : 42—44, 2001.
- 8) Bailli, G. S. and Douglas, L. J. : Matrix polymers of *Candida* biofilms and their possible role in biofilm resistance to antifungal agents. J. Antimicrob. Chemother. **46** : 397—403, 2000.
- 9) Donlan, R. M. and Costerton, J. W. : Biofilms : survival mechanisms of clinically relevant microorganisms. Clin. Microbiol. Rev. **15** : 167—193, 2002.
- 10) Kumamoto, C. A. and Vines, M. D. : Alternative *Candida albicans* lifestyles : growth on surfaces. Annu. Rev. Microbiol. **59** : 113—133, 2005.
- 11) Lopez-Ribot, J. L. : *Candida albicans* biofilms : more than filamentation. Curr. Biol. **15** : R453—R455, 2005.
- 12) Moghadam, B. K., Disko, C. L. and Gier, R. E. : Chlorhexidine mouth wash-induced fixed eruption. Case report and review of the literature. Oral Surg. Oral Med. Oral Pathol. **71** : 431—434, 1991.
- 13) Waren, K. D. and Munsick, R. A. : Anaphylaxis from povidone-iodine. Lancet **345** : 1506, 1995.
- 14) Reichart, P. A., Philipsen, H. P., Schmidt-Westhausen, A. and Samaranayake, L. P. : Pseudomembranous oral candidiasis in HIV infection : ultrastructural findings. J. Oral Pathol. Med. **24** : 268—281, 1995.
- 15) Fu, Y., Rieg, G., Fonzi, W. A., Belanger, P. H., Edwards, J. E. and Filler, S. G. : Expression of the *Candida albicans* gene ALS1 in *Saccharomyces cerevisiae* induces adherence to endothelial cells. Infect. Immun. **66** : 1783—1786, 1998.
- 16) McDonnell, G. E. : Antisepsis, disinfection, and sterilization ; types, action, and resistance. AZM Press, Washington, DC, 2007.
- 17) Sandovsky-Losica, H., Chauhan, N., Calderone, R. and Segal, E. : Gene transcription studies of *Candida albicans* following infection of HEp2 epithelial cells. Med. Mycol. **44** : 329—334, 2006.
- 18) Cawson, R. A. and Rajasingham, K. C. : Ultrastructural features of the invasive phase of *Candida albicans*. Br. J. Dermatol. **87** : 435—443, 1972.
- 19) Eversole, L. R., Reichart, P. A., Ficarra, G., Schmidt-Westhausen, A., Romagnoli, P. and Pimpinelli, N. : Oral keratinocyte immune responses in HIV-associated candidiasis. Oral Surg. Oral Med. Oral Pathol. Oral Radiol. Endodont. **84** : 372—380, 1997.
- 20) Kamai, Y., Kobota, M., Hosokawa, T., Fukuoka, T. and Filler, S. G. : New model of oropharyngeal candidiasis in mice. Antimicrob. Agents Chemother. **45** : 3195—3197, 2001.
- 21) Montes, L. F. and Wilbom, W. H. : Ultrastructural features of host-parasite relationship in oral candidiasis. J. Bacteriol. **96** : 1349—1356, 1986.

- 22) Arima, Y., Nakai, Y., Hayakawa, R. and Nishino, T. : Antibacterial effect of beta-thujaplicin on staphylococci isolated from atopic dermatitis : relationship between changes in the number of viable bacterial cells and clinical improvement in an eczematous lesion of atopic dermatitis. *J. Antimicrob. Chemother.* **51** : 113—122, 2003.
- 23) Chandra, J., Mukherjee, P.K., Leidich, S. D., Faddoul, F. F., Hoyer, L. L., Douglas, L. J. and Ghannoum, M. A. : Antifungal resistance of candidal biofilms formed on denture acrylic in vitro. *J. Dent. Res.* **80** : 903—908, 2001.
- 24) Eedtman, Y. and Gripenberg, J. : Antibiotic substances from the heart wood of *Thuja plicata* Don. *Nature* **161** : 719, 1948.
- 25) Inamori, Y., Shinohara, S., Tsujibo, H., Okabe, T., Morita, Y., Sakagami, Y., Kameda, Y. and Ishida, N. : Antimicrobial activity and metalloprotease inhibition of hinokitiol-related compounds of *Thujopsis dolabrata* S. and *Z. hondai* MAK. *Biol. Pharm. Bull.* **22** : 990—993, 1999.
- 26) Inamori, Y., Muro, C., Sajima, E., Katagiri, M., Okamoto, Y., Tanaka, H., Sakagami, Y. and Tsujibo, H. : Biological activity of purpurogalia. *Biosci. Biotechnol. Biochem.* **61** : 890—892, 1997.
- 27) Saeki, Y., Ito, Y., Shibata, M., Sato, Y., Okuda, K. and Takazoe, I. : Antimicrobial action of natural substances on oral bacteria. *Bull. Tokyo Dent. Coll.* **30** : 129—135, 1989.
- 28) Bentley, R. : A fresh look at natural tropolonoids. *Nat. Prod. Rep.* **25** : 118—138, 2008.
- 29) Hawse, S. P. and Douglas, L. J. : Biofilm formation by *Candida* species on the surface of catheter materials in vitro. *Infect. Immun.* **62** : 915—921, 1994.
- 30) Kuhn, D. M. and Ghannoum, M. A. : *Candida* biofilms : antifungal resistance and emerging therapeutic options. *Curr. Opin. Invest. Drugs* **5** : 186—197, 2004.
- 31) Mukherjee, P. K., Zhou, G., Munton, R. and Ghannoum, M. A. : *Candida* biofilm : a well-designed protected environment. *Med. Mycol.* **43** : 191—208, 2005.
- 32) Senpuku, H., Tada, A., Nakao, R., Yonezawa, H., Yoneda, S., Yoshihara, A. and Miyazaki, H. : Relationships of anti-PAc (361-386) peptide salivary IgA antibody, eosinophils and basophils with periodontal status in the elderly. *FEMS Immunol. Med. Microbiol.* **49** : 84—90, 2007.
- 33) Tada, A., Senpuku, H., Motozawa, Y., Yoshihara, A., Hanada, N. and Tanzawa, H. : Association between commensal bacteria and opportunistic pathogens in the dental plaque of elderly individuals. *Clin. Microbiol. Infect.* **12** : 776—781, 2006.
- 34) Chisty, M. M., Nargis, M., Inaba, T., Ishida, K., Osanai, A. and Kamiya, H. : Transmission electron microscopy of *Sclustosoma mansoni* cercariae treated with hinokitiol (beta-thujaplicin), a compound for potential skin application against cercarial penetration. *Tohoku J. Exp. Med.* **202** : 63—67, 2004.
- 35) Chambers, S. T., Peddie, B. and Pithie, A. : Ethanol disinfection of plastic-adherent micro-organisms. *J. Hosp. Infect.* **63** : 193—196, 2006.
- 36) Suci, P. A. and Tyler, B. J. : Action of chlorhexidine digluconate against yeast and filamentous forms in an early-stage *Candida albicans* biofilm. *Antimicrob. Agents Chemother.* **46** : 3522—3531, 2002.
- 37) Müller, G. and Kramer, A. : Biocompatibility index of antiseptic agents by parallel assessment of antimicrobial activity and cellular cytotoxicity. *J. Antimicrob. Chemother.* **61** : 1281—1287, 2008.
- 38) Sanchez, I. R., Nusbaum, K. E., Swaim, S. F., Hale, A. S., Henderson, R. A. and McGuire, J. A. : Chlorhexidine diacetate and povidone-iodine cytotoxicity to canine embryonic fibroblasts and *Staphylococcus aureus*. *Vet. Surg.* **17** : 182—185, 1988.
- 39) Malic, S., Hill, K. E., Ralphs, J. R., Hayes, A., Thomas, D. W., Potts, A. J. and Williams, D. W. : Characterization of *Candida albicans* infection of an *in vitro* oral epithelial model using confocal laser scanning microscopy. *Oral Microbiol. Immunol.* **22** : 188—194, 2007.
- 40) Presterl, E., Suchomel, M., Eder, M., Reichmann, S., Lassnigg, A., Graninger, W. and Rotter, M. : Effects of alcohols, povidone-iodine and hydrogen peroxide on biofilms of *Staphylococcus epidermidis*. *J. Antimicrob. Chemother.* **60** : 417—420, 2007.
- 41) Howell, J. M. and Chisholm, C. D. : Outpatient wound preparation and care : a national survey. *Ann. Emerg. Med.* **21** : 976—981, 1992.
- 42) Yamano, H., Yamazaki, T., Sato, K., Shiga, S., Hagiwara, T., Ouchi, K. and Kishimoto, T. : In vitro inhibitory effects of hinokitiol on proliferation of *Chlamydia trachomatis*. *Antimicrob. Agents Chemother.* **49** : 2519—2521, 2005.
- 43) Kuhn, D. M., George, T., Chandra, J., Mukherjee, P. K. and Ghannoum, M. A. : Antifungal susceptibility of *Candida* biofilms : Unique efficacy of amphotericin B lipid formulations and echinocandims. *Antimicrob. Agents Chemother.* **46** : 1773—1780, 2002.



Contents lists available at ScienceDirect

Archives of Gerontology and Geriatrics

journal homepage: www.elsevier.com/locate/archger

Effects of mucosal care on oral pathogens in professional oral hygiene to the elderly

Yoshihide Nishiyama^a, Erika Inaba^{b,c}, Hiroshi Uematsu^b, Hidenobu Senpuku^{c,*}

^a Dental Support Co., Ltd., Chiba, 261-8501, Japan

^b Department of Gerodontology, Tokyo Medical and Dental University, Tokyo, 113-8510, Japan

^c Department of Bacteriology, National Institute of Infectious Diseases, Toyama 1-23-1, Shinjuku-ku, Tokyo, 162-8640, Japan

ARTICLE INFO

Article history:

Received 3 June 2009
Received in revised form 12 January 2010
Accepted 9 April 2010
Available online 21 May 2010

Keywords:

Mucosal care
Oral professional care
Oral hygiene
Mutans streptococci
Candida sp.
Elderly

ABSTRACT

To elucidate the effects of mucosal care on colonization of mutans streptococci and infection of *Candida* species, we randomized 50 functionally dependent nursing home residents, mean age 76.1 (± 7.8 years), 76% females, to receive oral professional care with or without mucosal care. During 12 months of follow-up, significant reduction in the numbers of mutans streptococci in saliva and dental plaque samples were observed in both treatment groups. However, there was no reduction in the numbers of mutans streptococci in tongue samples in the non-mucosal group ($p = 0.150$). In the comparison between the treatment with and without mucosal care, the treatment with mucosal care showed a significant inhibition effects on numbers of mutans streptococci on the tongue surface at 6 months after the treatment as compared with the treatment without mucosal care ($p = 0.043$). Mucosal care was not associated with any significant reduction in the rate of opportunistic infection with *Candida* species in any of the three sites. However, more subjects without mucosal care had *Candida* infection in dental plaque during the follow-up ($p = 0.046$). Professional dental care with (versus without) mucosal care had no effect on mutans streptococci colonization in saliva or plaque. Lack of mucosal care seemed to be associated with higher rates opportunistic infections with *Candida* species.

© 2010 Elsevier Ireland Ltd. All rights reserved.

1. Introduction

Institutionalized elderly individuals who need systemic care have poorer oral health than those who live independently at home (Simons et al., 1999; Yoneyama et al., 1999; Senpuku et al., 2003). Diminished oral health, in turn, may affect their quality of life (Karuza et al., 1992). Poor oral health increases the risk of systemic disease such as aspiration pneumonia and septicemia (Scannapieco, 1999; Shay, 2002) as well as for dental diseases such as caries (Tada et al., 2004).

Although various species of bacteria have been found in oral cavity, mutans streptococci (mutans streptococci, *Streptococcus mutans* and *Streptococcus sobrinus*) are shown to be the principal causative pathogens related to dental caries in humans (Köhler and Persson, 1991; Kolenbrander and London, 1993; Loesche et al., 1995). The fungus *Candida albicans* and other *Candida* species constitute a part of the normal microbial flora in the oral cavity. However, some carriers with a high density of *Candida* species may become infected (oral candidosis) and has a higher prevalence in the elderly; and may prompt the dentist or clinician to look out for underlying pathology changes in the seemingly healthy individuals.

Therefore, oral *Candida* may be an indicator of pneumonia because of fungal aspiration from the oral cavity.

Oral professional care is an important practice to maintain the oral health of the elderly (Yoneyama et al., 1999; Abe et al., 2001). However, little is known about how mucosal oral care controls mutans streptococci colonization and infection of *Candida* in the oral cavity. To elucidate the effects of mucosal care on colonization of mutans streptococci and infection of *Candida* to maintain oral health of elderly, we studied two groups of elderly patients treated using oral professional care with or without mucosal care.

2. Materials and methods

2.1. Human subjects

Fifty elderly residents from an Itabashi-ward Tokyo nursing home who are dependently living in long-term nursing care participated in this study. The study was conducted from June 2006 to June 2007. The subjects were randomly selected from the residents using a random-numbers table and their identity was held blind from the investigators. Prior to the survey, the aim and details of the study were explained and consent was obtained from all subjects before registration. The study was approved by the Ethics Committee of the Tokyo Medical and Dental University and performed according to the rules of the Helsinki Declaration. One

* Corresponding author. Tel.: +81 3 5285 1111x2223; fax: +81 3 5285 1163.
E-mail address: hsempuku@nih.go.jp (H. Senpuku).

subject dropped out at 2 months, one at 6 months and four at 12 months after beginning the study where they moved to other institutions. Forty subjects (75.9 ± 7.5 years old; 29 females, 11 males) with more than 10 teeth were selected to compare the infections with streptococci and lactobacilli after oral professional care and with or without mucosal care, because the numbers of teeth influence the amount of colonization of streptococci and lactobacilli in the oral cavity. To study *Candida* species infections, all 50 subjects (76.1 ± 7.8 years old; 38 females, 12 males) were sampled before and after professional care. Dental examinations to determine the presence of dental caries, periodontal inflammation, denture usage, oral hygiene level, coated tongue and other typical oral conditions were performed under artificial white light by trained dentists before the study; and then routine oral professional care began. Oral examinations were performed according to WHO oral examination procedures (WHO, 1986). Coronal caries status indicators were: the number of coronal and root decayed teeth, the number of coronal and root filled teeth, the number of coronal and root surface decayed teeth and the number of coronal and root surface filled teeth. The general oral status of the subjects before the investigation is shown in Table 1. The subjects were randomly divided into two groups; Groups treated with professional oral care with and without mucosal care. The care givers and other staff at the institution remained unchanged during the study period.

2.2. Oral care

During the initial examination, all patients had a routine dental and medical check-up. Oral care techniques by dental hygienists were standardized before beginning oral care. For daily oral care, all subjects who were able used the sink facilities in their room performed daily care 2–3 times a day after each meal. Further, dental hygienists provided professional care for 20 min once per week, i.e. dental brushing of teeth surfaces, mucosal cleaning with a sponge brush (NIPPON ZETTOC, Tokyo, Japan), tongue cleaning with tongue brush (Ebis Co., Ltd, Nara, Japan), denture cleaning, and oral washing with tap water in addition to the daily oral care. After professional care, mucosal cleaning was performed by sponge brush soaked in tap water for 1 min in oral professional care group with mucosal care. The routine oral care including professional care was performed for 12 months. The subjects received no standard or professional care prior to entering the institution and the first sampling. No antibiotic therapy was administered during the 2 weeks prior to this study and during the 12-month study period; and none of these subjects suffered from severe infections. There was no significant difference in systemic disease incidence between the two groups with and without mucosal care prior to beginning the study (Table 1). There was no information concerning antibiotic therapy of the subjects before entering the institution.

2.3. Sampling of saliva, plaque and tongue

Subjects were asked not to brush their teeth 2 h prior to sampling. All sampling was performed before oral professional care and at 0, 1, 2, 3, 6 and 12 months after beginning the study. Whole saliva was stimulated by chewing paraffin gum for 5 min. For bacterial counts, a sterile cotton swab (Seedswab No. 1, Eiken Chemical Co. Ltd., Tokyo) was immersed in the saliva for 10 s and placed in transport fluid (0.4% agar, 0.15% thioglycolate/phosphate). Supragingival plaque samples were collected by swabbing five times from the posteroanterior buccal surface of the upper right second premolar and first molar using a cotton swab. Subjects having partial dentures ($n = 29$) and not having the above teeth were sampled from the opposite side or other remaining teeth. Tongue samples were collected by swabbing five times from the

Table 1
Baseline characteristics in subjects.

| | Total | With mucosal care | Without mucosal care |
|--------------------------|------------|-------------------|----------------------|
| Age | 76.1 ± 7.8 | 75.6 ± 8.4 | 76.8 ± 6.1 |
| Gender (female:male) | 38:12 | 21:6 | 18:6 |
| Tooth brush/day | 2.5 ± 2.0 | 2.5 ± 1.0 | 2.5 ± 1.0 |
| Existing tooth | 17.0 ± 7.7 | 17.8 ± 7.6 | 16.3 ± 7.9 |
| Decayed tooth | 0.9 ± 2.0 | 0.8 ± 1.2 | 1.0 ± 2.5 |
| Missing tooth | 10.9 ± 7.8 | 10.2 ± 7.7 | 11.8 ± 7.9 |
| Filled tooth | 9.5 ± 5.3 | 9.4 ± 4.2 | 9.6 ± 6.4 |
| Partial denture usage | | | |
| Do | 29 | 14 | 15 |
| Do not | 21 | 12 | 9 |
| Status of oral hygiene | | | |
| Good | 9 | 5 | 4 |
| Moderate | 35 | 17 | 18 |
| Bad | 6 | 14 | 2 |
| Inflammation of gingival | | | |
| No | 0 | 0 | 0 |
| Mild | 17 | 9 | 8 |
| Moderate | 27 | 15 | 12 |
| Heavy | 6 | 2 | 4 |
| Coating of tongue | | | |
| –1/3 | 11 | 6 | 5 |
| 1/3–2/3 | 23 | 10 | 13 |
| 2/3– | 12 | 6 | 6 |
| Oral dryness | | | |
| Normal | 27 | 15 | 15 |
| Mild | 2 | 2 | 2 |
| Moderate | 10 | 6 | 6 |
| Severe | 11 | 3 | 3 |
| Medical problem | | | |
| Sinus problem | 4 | 1 | 3 |
| Pneumonia | 9 | 4 | 5 |
| GI disorder | 7 | 4 | 3 |
| Diabetes mellitus | 7 | 5 | 2 |
| Liver complaint | 7 | 5 | 2 |
| Blood disorder | 2 | 2 | 2 |
| Auto immune disease | 2 | 1 | 1 |
| Cancer | 2 | 0 | 2 |
| Cardiac disease | 10 | 6 | 4 |
| High-blood pressure | 20 | 9 | 11 |
| Kidney diseases | 2 | 2 | 0 |
| Smoking habit | 5 | 4 | 1 |

Values indicated mean (\pm standard deviation) of age, tooth brush/day, existing tooth, decayed tooth, missing tooth and filled tooth. Other values indicated numbers of person.

center of the tongue. Totally two swabs were acquired to analyze the bacteria and fungi in each sampling. After placing in transport fluid, all samples were immediately taken to the Biomedical Laboratory (BML, Tokyo, Japan) for analysis to detect mutans streptococci, total Streptococci (TS), Lactobacilli (LB) and *Candida* species.

2.4. Counting of bacteria and identification of fungi

Each sample was poured onto Mitis-Salivarius agar (Nippon Becton Dickinson Co. Ltd, Tokyo, Japan) to count TS, modified Mitis-Salivarius agar containing 0.2 U/m bacitracin to count mutans streptococci, and Rogosa selective lactobacillus agar to count LB using an EDDY JET spiral plating system (IUL, S.A., Barcelona, Spain); incubated at 37 °C using anaerobic conditions for 48 h (Tsuha et al., 2004); and the numbers of colony forming units (CFU) were counted. Colonies of mutans streptococci were identified by their characteristic appearance.

For *Candida* species identification, each sample was poured directly onto sabouroud dextrose agar plate and incubated for 48 h

at 35 °C to identify *Candida* species colonies, which were identified by their characteristic morphological appearance and color, and using *Candida* check (Iatron Laboratories Inc., Tokyo, Japan).

2.5. Statistical procedures

All data were analyzed using the Statistical Package for SPSS for Windows (version 100, Chicago, IL, USA). The repeated data for bacterial number (Log10) and *Candida* species identification were compared using the one way ANOVA and Friedman test in Table 2. *T*-test with the Bonferroni method was used to compare primary data with data after oral professional care at each month or the two groups of oral professional care with and without oral mucosal care. In comparison between the treatment with and without oral mucosal care, the data subtracted data after the treatment from primary data in the numbers of mutans streptococci were used for analyses in the *T*-test with the Bonferroni method. The data of parameters excepting for mean ± SD in Table 1 and all data in Table 3 were compared between two groups using a χ^2 test for equal and unequal variations. *p*-Values less than 0.05 were considered to be significant.

3. Results

3.1. General and oral status

Comparing systemic diseases such as sinus problems, pneumonia, cardiac disorders, diabetes mellitus and others showed there were no significant differences between oral mucosal care-treated and -non-treated groups (Table 1). The dental status of the groups with and without oral mucosal treatment is shown in Table 1

where there were no significant differences between the two groups in all parameters.

3.2. Effects of oral mucosal treatment on oral bacteria colonization

The roles of oral mucosal treatment for the oral pathogenic bacteria, mutans streptococci, lactobacilli and total streptococci were studied for the tongue, plaque and saliva samples from both groups at various time points after oral professional care. Significant differences ($\dagger p < 0.05$ and $\dagger\dagger p < 0.01$) in the decrease in the repeated data of mutans streptococci numbers were shown using the one way ANOVA in all samples from both groups except for the tongue sample from elderly patients without oral mucosal care (Table 2). Further, in the two data comparisons using the *T*-test with the Bonferroni method, the numbers of mutans streptococci decreased significantly in the plaque samples at 6 and 12 months after the treatment of elderly patients without oral mucosal care in comparison to the primary data (at 0 month, $*p < 0.05$, Table 3); whereas, a significant decrease in the saliva sample was observed throughout the study from 1 to 12 months ($*p < 0.05$ and $**p < 0.01$, Table 2). However, there were no significant differences in tongue samples (Table 2). Further, a significant decrease in mutans streptococci numbers was also observed throughout the study from 1 to 12 months in all samples from the elderly patients with oral mucosal care ($*p < 0.05$ and $**p < 0.01$, Table 2). In the comparison between the treatment with and without oral mucosal care in Table 2, a significant difference was only recognized at 6 months in tongue sample (*T*-test with the Bonferroni method, $p = 0.043$) but not at other months and in other samples. The treatment with mucosal care showed a significant inhibition effects on numbers of mutans streptococci on the tongue

Table 2
Numbers of mutans streptococci in elderly subjects with and without oral mucosal care.

| Subject group Sample | After start of oral professional care (month) | | | | | | One way ANOVA (<i>p</i>) |
|------------------------------------|---|-------------|-------------|-------------|-------------|-------------|-------------------------------|
| | 0 | 1 | 2 | 3 | 6 | 12 | |
| With mucosal care (N=21) | | | | | | | |
| Saliva | 5.0 ± 2.1 | 3.5 ± 2.2* | 3.5 ± 2.0** | 3.4 ± 2.1** | 3.0 ± 1.9** | 3.0 ± 1.9** | 0.000 ^{††} |
| Plaque | 5.3 ± 2.0 | 4.1 ± 2.5* | 4.2 ± 2.3* | 3.7 ± 2.5* | 3.5 ± 2.3** | 3.7 ± 2.0* | 0.000 ^{††} |
| Tongue | 5.1 ± 2.0 | 3.5 ± 2.2** | 3.8 ± 2.2* | 3.5 ± 2.2** | 2.9 ± 1.9** | 3.2 ± 2.2** | 0.003 ^{††} |
| Without mucosal care (N=19) | | | | | | | |
| Saliva | 5.1 ± 1.8 | 4.1 ± 1.4** | 4.0 ± 1.7* | 4.1 ± 1.4* | 3.4 ± 1.5** | 3.5 ± 1.7** | 0.000 ^{††} |
| Plaque | 5.2 ± 2.1 | 4.9 ± 1.3 | 4.8 ± 2.0 | 4.6 ± 2.0 | 4.0 ± 2.0* | 4.0 ± 2.2** | 0.019 [†] |
| Tongue | 4.5 ± 2.4 | 4.0 ± 1.6 | 4.1 ± 1.5 | 4.0 ± 1.7 | 3.6 ± 1.5 | 3.7 ± 1.5 | 0.150 |

The results are expressed as mean (±standard deviation) of number (Log10/ml) of mutans streptococci. Asterisks indicate significant differences between 0 month and other months in *T*-test with Bonferroni method ($*p < 0.05$; $**p < 0.01$) and among 0–12 months in one way ANOVA ($\dagger p < 0.05$; $\dagger\dagger p < 0.01$). *N*: numbers of subjects with mucosal care or without mucosal care.

Table 3
Numbers of subjects which detected *Candida* species.

| Subject group Sample | After start of oral professional care (month) | | | | | | Friedman test (<i>p</i>) |
|------------------------------------|---|----------|---------|----------|----------|-----------|-------------------------------|
| | 0 | 1 | 2 | 3 | 6 | 12 | |
| With mucosal care (N=26) | | | | | | | |
| Saliva | 1 (3.8) | 3 (11.5) | 0 (0.0) | 1 (3.8) | 2 (7.7) | 1 (3.8)* | 0.487 |
| Plaque | 2 (7.7) | 3 (11.5) | 0 (0.0) | 0 (0)** | 1 (3.8) | 0 (0)** | 0.127 |
| Tongue | 2 (7.7) | 3 (11.5) | 0 (0.0) | 2 (7.7) | 0 (0) | 2 (7.7) | 0.298 |
| Total | 2 (7.7) | 4 (15.4) | 0 (0.0) | 2 (7.7)* | 3 (11.5) | 3 (11.5)* | 0.437 |
| Without mucosal care (N=24) | | | | | | | |
| Saliva | 2 (8.3) | 2 (8.3) | 2 (8.3) | 4 (16.7) | 4 (16.7) | 6 (25.0) | 0.299 |
| Plaque | 3 (12.5) | 4 (16.7) | 1 (4.2) | 5 (20.8) | 5 (20.8) | 8 (33.3) | 0.046 [†] |
| Tongue | 2 (8.3) | 4 (16.7) | 0 (0) | 4 (16.7) | 2 (8.3) | 5 (20.8) | 0.133 |
| Total | 3 (12.5) | 4 (16.7) | 2 (8.3) | 8 (33.3) | 5 (20.8) | 9 (37.5) | 0.006 ^{††} |

The results are expressed as numbers (% to total number) of subjects which detected *Candida* species. Total was counted subject numbers which detected *Candida* species in saliva, plaque or tongue. Asterisks indicate significant differences between professional treatment with mucosal care and without mucosal care in χ^2 -test ($*p < 0.05$; $**p < 0.01$) in each sample and month, and among 0–12 months in Friedman test ($\dagger p < 0.05$; $\dagger\dagger p < 0.01$). *N*: numbers of subjects with mucosal care or without mucosal care.

surface at 6 months after the treatment as compared with the treatment without mucosal care. Significant differences for the repeated data of lactobacilli were not observed in all samples from both groups (data not shown). Mucosal care had no significant association with reduction in lactobacilli count. Significant differences in the decrease in the repeated data of total streptococci numbers were indicated in all samples from both groups (data not shown). Dental care with or without mucosal care had significant association with reduction in total streptococci count.

3.3. Effects of oral mucosal treatment on *Candida* species infection

The roles of oral mucosal treatment for the *Candida* species were also studied for the same samples from both groups at various time points after oral professional care. The infection of *Candida* species was qualitatively assessed in each sample. Significant differences ($^1p < 0.05$ and $^{11}p < 0.01$) in the increase in the repeated data of isolation frequency of *Candida* species were shown using the Friedman test in plaque samples and total of samples from elderly patients without oral mucosal care (Table 3). However, in elderly patients with oral mucosal care, there were no significant differences. Significant differences between elderly with and without oral mucosal care were also observed in isolation frequency of *Candida* species from total of saliva, plaque and tongue at 3 and 12 months after treatment with professional oral hygiene ($^*p < 0.05$, Table 3). In particular, the differences were observed in saliva at 12 months ($^*p < 0.05$) and plaque at 3 and 12 months ($^{**}p < 0.01$). However, in other months and tongue, there were no significant differences.

Microbiology studies previously suggested the colonized denture acted as a possible reservoir of *Candida* infection (Budtz-Jørgensen, 1974). However, we found no significant difference in denture usage between groups treated by professional care with or without mucosal care.

4. Discussion

Long-term decreasing effects of mucosal care on mutans streptococci number were recognized on the tongue surface in which physical attachment of bacteria is due to existence of many papillae. On the other hand, the decreasing effects of mucosal care appeared shortly after start of the professional treatment, but were lost during long-term treatment on the tooth surface in which bacterial attachment is supported by salivary binding receptors absorbed on the enamel surface to oral streptococci. The difference of efficacies might be dependent on physical and biochemical interactions between the tissue surfaces and bacteria. The mucosal care might have more benefit to reduce mutans streptococci numbers in the physical attachment to tongue surface than in the biochemical attachment to tooth surface during long-term treatment. We suggest at least oral mucosal care reduces infection directly and reduces the attachment of mutans streptococci on the tongue in the oral cavity; and this was further reduced in the biofilm using professional treatment. However, the effects of mucosal care on the planktonic cell numbers of mutans streptococci in saliva samples were not observed in comparison with data before start of oral professional care. Oral hygiene procedures with mucosal care are used to remove the biofilm and micro-colonies formed by oral microorganisms on teeth and tongue surfaces, and mucosal epithelial cells attached to oral microorganisms on mucosal surfaces. Further, a healthy and fresh environment is created in the oral cavity. Oral mucosal care may be more effective in reducing reattachment of remaining oral microorganisms including pathogens such as mutans streptococci attached to tongue after professional oral treatment, and supports

the effects of professional treatment in the oral cavity without the biofilm after professional treatment than by using professional care without mucosal care that does not completely remove microorganisms on oral mucosal tissues.

In contrast, the oral mucosal care may not affect the decrease or increase of lactobacilli numbers during treatment using oral professional care. The mucosal care may loosen adhesive bacteria such as mutans streptococci and some of the oral streptococci on the tissues but not non-adhesive bacteria such as lactobacilli or some of the planktonic cells in the oral cavity.

The oral streptococci are the first to attach in early dental plaque formation and have a specific temporal and spatial distribution that is essential for the development of the oral biofilm (Rickard et al., 2003). Reports show early colonizers of oral streptococci species (mitis group streptococci) can suppress the growth of *S. mutans* by producing hydrogen peroxide (Kreth et al., 2005). The streptococci compete for adhesion-binding sites on the saliva-coated tooth surface (Nobbs et al., 2007). Recently, *Streptococcus oligofermentans* was found as a new oral commensal species and has an inverse correlation with *S. mutans* within human dental plaque (Tong et al., 2007). In combination cultures, *S. salivarius* strongly inhibits biofilm formation when cultured with *S. mutans* (Tamura et al., 2009). This inhibition occurs in the early phase of biofilm formation and was dependent on inactivation of the competence stimulating peptide of *S. mutans*. Therefore, some oral streptococci may function as competitive strains to inhibit *S. mutans* colonization. When the oral streptococci are refreshed by professional treatment with oral mucosal care, this treatment may create a potential for a protective environment to inhibit infection by mutans streptococci.

It is generally considered the mechanisms employed when a healthy microflora is present may interfere with the adhesion of invading pathogens and includes competitive exclusion (Reid and Tieszer, 1993), displacement (Millsap et al., 1994), production of antibacterial compounds (Klaenhammer, 1988), and release of bio-surfactants (Frandsen et al., 1991). Oral streptococci have an antagonistic effect toward many bacterial species (Holmberg and Hallander, 1973; Willcox and Drucker, 1988; Bormard and Stinson, 1999) and have an inverse correlation with the detection of opportunistic pathogens (Tada et al., 2007). Therefore, we consider oral professional care with mucosal care cleanses the tongue and mucosal surface where the treatment allows re-establishment and growth by fresh commensal bacteria such as total streptococci; and disturbs the infection by opportunistic pathogens such as *Candida* in the oral cavity of the elderly. In this study, the significant effects of treatment with mucosal care on *Candida* species were observed in saliva and plaque samples but not in tongue sample. The results of inverse influence on samples were shown in the effects of mucosal care on mutans streptococci. It was also considered that the difference of efficacies on *Candida* species and mutans streptococci infections might be dependent on physical and biochemical interactions between the tissue surfaces and pathogens. However, to attain these potential effects treatment may be required for long periods to disturb increase of *Candida* species infections because the differences of isolate frequency of *Candida* between treatment with and without oral mucosal care were observed after 3 months after professional care treatment began, and the increases of repeated data were also observed during 12 months in the treatment without mucosal care. The treatment with mucosal care might disturb the increase of *Candida* species infection in the oral cavity. In other studies, long duration of professional care treatment was suggested to be effective in decreasing the risk of oral infection and inflammation in institutionalized elderly (Adachi et al., 2002; Yoneyama et al., 2002; Kokubu et al., 2008). Dental care of the institutionalized elderly is often limited to emergency care and is not aimed at