

研究成果の刊行に関する一覧表

書籍

著者氏名	論文タイトル名	書籍全体の 編集者名	書 籍 名	出版社名	出版地	出版年	ページ

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Doi K, <u>Wei M</u> , Kitano M, Uematsu N, Inoue M, <u>Wanibuchi H.</u>	Enhancement of preneoplastic lesion yield by chios mastic Ggm in a rat liver medium-term carcinogenesis bioassay.	Toxicol.Appl.Pharmacol.	234	135-142	2009
Kushida M, <u>Wanibuchi H</u> , <u>Wei M</u> , Kakehashi A, Ozaki K, Sukata T, Miyata K, Ogata K, Uwagawa S, Fukushima S.	Ethanol dose not promote MeIQx-initiated rat colon carcinogenesis based on evidence from analysis of a colon cancer surrogate marker.	J Toxicol.Pathol.	22	65-70	2009
Tanaka-Okamoto M, Hori K, Ishizaki H, Hosoi A, Itoh Y, <u>Wei M</u> , <u>Wanibuchi H</u> , Mizogchi A, Nakamura H, Miyoshi J.	Increased susceptibility to spontaneous lung cancer in mice lacking LIM-domain only 7.	Cancer Sci.	100	608-616	2009
Fukushima S, Kakehashi A, <u>Wei M</u> , <u>Wanibuchi H.</u>	Existence of a threshold for the genotoxic carcinogens: Evidence from mechanism-based carcinogenicity studies.	Genes and Environment	31	33-36	2009
Kakehashi A, Inoue M, <u>Wei M</u> , Fukushima S, <u>Wanibuchi H.</u>	Cytokeratin 8/18 overexpression and complex formation as an indicator of GST-P positive foci transformation into hepatocellular carcinomas.	Toxicol.Appl.Pharmacol.	238	71-79	2009
Romanenko A,	Urinary bladder carcinogenesis induced by	Carcinogenesis	30	1821-18	2009

Kakehashi A, Morimura K, <u>Wanibuchi H</u> , <u>Wei M</u> , Voizianov A, Fukushima S.	chronic exposure to persistent low-dose ionizing radiation after Chernobyl accident.			31	
Kakehashi A, Kato A, Inoue M, Ishii N, Okazaki E, <u>Wei M</u> , Tachibana T, <u>Wanibuchi H</u> .	Cytokeratin 8/18 as a new marker of mouse liver preneoplastic lesions.	Toxicol.Appl.Pharmacol.	242	47-55	2009
<u>Wei M</u> , Hamoud AS, Yamaguchi T, Kakehashi A, Morimura K, Doi K, Kushida M, Kitano M, <u>Wanibuchi H</u> , Fukushima S.	Patassium bromated enhances N-Ethyl-N-Hydroxyethylnitrosamine- Induced kidney carcinogenesis only at high dose in wistar rats: Indication of the existence of an enhancement threshold.	Toxicol Pathol.	37	983-991	2009
Yokohira M, Hashimoto N, Yamakawa K, Suzuki S, Saoo K, Kuno T, <u>Imaida K</u> .	Lack of modifying effects of intratracheal instillation of quartz or dextran sulfate sodium (DSS) in drinking water on lung tumor development initiated with 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) in female A/J mice.	J. Toxicol. Pathol.	22	179-85	2009
Hashimoto N, Yachida S, Okano K, Wakabayashi H, <u>Imaida K</u> , Kurokohchi K, Masaki T, Kinoshita H, Tominaga M, Ajiki T, Ku Y, Okabayashi T, Hanazaki K, Hiroi M, Izumi S, Mano S, Okada S, Karasawa V, Maeba V, Suzuki Y.	Immunohisto- chemically detected expression of p27kip1 and skp2 predicts survival in patients with intrahepatic cholangiocarcinomas.	Ann. Surg.Oncol.	16	395-403	2009
Matsuda Y, Takeuchi H, Yokohira M, Saoo K, Hosokawa K, Yamakawa K, Zeng Y, Totsuka Y, Wakabayashi K, <u>Imaida K</u> .	Enhancing effects of a high fat diet on 2-amino-3,8-dimethylimidazo [4,5-f]quinoxaline (MeIQx) induced lung tumorigenesis in female A/J mice.	Molecular Medicine Reports	2	701-706	2009
Takeuchi H, Saoo K, Matsuda Y, Yokohira	8-Methoxy-psoralen, a potent human CYP2A6 inhibitor, inhibits lung adenocarcinoma	Molecular Medicine Reports	2	585-588	2009

M, Yamakawa K, Zeng Y, Kuno T, Kamataki T, <u>Imaida K.</u>	development induced by 4-(methyl-nitrosamino)-1-(3-pyridyl)-1-butanone (NNK) in female A/J mice.				
Yokohira M, Hashimoto N, Yamakawa K, Suzuki S, Saoo K, Kuno T, <u>Imaida K.</u>	A lung carcinogenic bioassay of CuO and TiO <sub>2</sub> nanoparticles with intratracheal instillation using F344 male rats.	J. Toxicol. Pathol.	22	71-78	2009
Yokohira M, Mastuda Y, Suzuki S, Hosokawa K, Yamakawa K, Hashimoto N, Saoo K, Nabae K, Doi Y, Kuno T, <u>Imaida K.</u>	Equivocal colonic carcinogenicity of Aloe arborescens Miller var. natalensis Berger at high dose level in a Wistar Hannover rat 2-year Study.	J. Food Sci.	77	T24-30	2009
Yokohira M, Kuno T, Yamakawa K, Hashimoto N, Ninomiya F, Suzuki S, Saoo K, <u>Imaida K.</u>	An intratracheal instillation bioassay system for detection of lung toxicity due to fine particles in F344 rats.	J. Toxicol. Pathol.	22	1-10	2009
Doi K, Sakai K, Tanaka R, Toma K, Yamaguchi T, <u>Wei M</u> , Fukushima S, <u>Wanibuchi H.</u>	Chemopreventive effects of 13alpha, 14alpha-epoxy-3beta-methoxyserratane-21beta-ol (PJJ-34), a serratane-type triterpenoid, in a rat multi-organ carcinogenesis bioassay.	Cancer Lett.	289	161-169	2010
Suzuki S, Arnold LL, Pennington KL, Kakiuchi-Kiyota S, <u>Wei M</u> , <u>Wanibuchi H.</u> , Cohen SM.	Effects of pioglitazone, a peroxisome proliferator-activated receptor gamma agonist, on the urine and urothelium of the rat.	Toxicol Sci.	113	349-357	2010
Takahashi Y, Hara Y, Imanaka M, <u>Wanibuchi H.</u> , Tanaka K, Ishikawa T, Mori S, Fukusato T.	No inhibitory effects of (–)-epigallocatechin gallate and lycopene on spontaneous hepatotumorigenesis in C3H/HeN mice.	Fukushima J. Med. Sci.	56	17-27	2010
Fukushima S, <u>Wei M</u> , Kakehashi A, <u>Wanibuchi H.</u>	Thresholds for genotoxic carcinogens: Evidence from mechanism-based carcinogenicity studies.	Cancer Risk Assessment	8	207-221	2010
Tago Y, <u>Wei M</u> , Ishii N, Kakehashi A, <u>Wanibuchi H.</u>	Evaluation of the subchronic toxicity of dietary administered <i>Equisetum arvense</i> in F344 rats.	J Toxicol. Pathol.	23	245-251	2010
Yachida S, <u>Imaida K.</u> , Yokohira M, Hashimoto N, Suzuki	Jun activation domain binding protein 1 is overexpressed from the very early stages of hepatocarcinogenesis.	Surgical Oncol.	17	3386-3393	2010

S, Okano K, Wakabayashi H, Maeta H, Suzuki Y.					
Kohno M, Haramoto M, Nakajima O, Yang L, Hinotsu S, Yokohira M, <u>Imaida K</u> , Kawakami K.	Antedrug budesonide by intrapulmonary treatment attenuates bleomycin-induced lung injury in rats with minimal systemic adverse effects.	Biol. Pharm. Bull.	33	1206-1211	2010
<u>Imaida K</u> , Yokohira M, Hashimoto N, Kuno T.	Risk analysis of environmental chemicals on lung carcinogenesis.	Asian Pacific J Cancer Prev	11	9-12	2010
Yokohira M, Hashimoto N, Yamakawa K, Suzuki S, Saoo K, Kuno T, <u>Imaida K</u> .	Potassium octatitanate fibers (TISMO) induce pleural mesothelial cell reactions with iron accumulation in female A/J mice.	Oncol. Lett.	1	273-278	2010
Tsuda H, Futakuchi M, Fukamachi K, Shirai T, <u>Imaida K</u> , Fukushima S, Tatematsu M, Furukawa F, Tamano S, Ito N.	A medium-term, rapid rat bioassay model for the detection of carcinogenic potential of chemicals.	Toxicol Pathol.	38	182-187	2010
Kuno T, Hirose Y, Yamada Y, <u>Imaida K</u> , Tatematsu K, Mori Y, Mori H.	Chemoprevention of 1,2-dimethyl-hydrazine-induced colonic preneoplastic lesions in Fischer rats by 6-methylsulfinylhexyl isothiocyanate, a wasabi derivative.	Oncology Letters	1	273-278	2010
Yamakawa K, Kuno T, Hashimoto N, Yokohira M, Suzuki S, Nakano Y, Saoo K, <u>Imaida K</u> .	Molecular analysis of carcinogen induced rodent lung tumors-Involvement of microRNA expression and Kras or EGFR mutation.	Molecular Medicine Reports	3	141-147	2010
Takeuchi H, Saoo K, Yamakawa K, Matsuda Y, Yokohira M, Zeng Y, Kuno T, Totsuka Y, Takahashi M, Wakabayashi K and <u>Imaida K</u> .	Tumorigenesis of 2-Amino-3,8-dimethylimidazo [4,5-f]quinoxaline (MeIQx), but not enhancing effects of concomitant high fat diet, on lung carcinogenesis in female A/J mice.	Oncol. Lett.	1	137-142	2010
Suzuki S, Yokohira M, Hashimoto N, Saoo K, Matsuda Y, Yamakawa K, Nakano Y, Kuno T, <u>Imaida K</u> .	Different threshold levels for 2-amino-3,8-dimethylimidazo-[4,5-f]quinoxaline (MeIQx) initiation of lung and colon carcinogenesis and the effects of an additional initiation by 4-(methylnitrosamino)-1-(3-pyridyl) -1-butanone (NNK) in A/J mice.	Molecular Medicine Reports	3	301-307	2010

Tsujino M, Fujii M, Okabe K, Mori T, Fukushima N, <u>Tsujiuchi T.</u>	Differential expressions and DNA methylation patterns of lysophosphatidic acid receptor genes in human colon cancer cells.	Virchows Arch	457	669-676	2010
Okabe K, Hayashi M, Fujii M, Honoki K, Mori T, Fukushima N, <u>Tsujiuchi T.</u>	Mutations of lysophosphatidic acid receptor genes in human osteosarcoma cells.	Pathobiology	77	278-282	2010
Okabe K, Hayashi M, Wakabayashi N, Yamawaki Y, Teranishi M, Fukushima N, <u>Tsujiuchi T.</u>	Differential expressions and DNA methylation patterns of lysophosphatidic acid receptor genes in mouse tumor cells.	Pathobiology	77	309-314	2010
Ishii N, <u>Wei M</u> , Kakehashi A, Doi K, Yamano S, Inaba M, <u>Wanibuchi H.</u>	Enhanced urinary bladder, liver and colon carcinogenesis in Zucker diabetic fatty rats in a multi-organ carcinogenesis: Evidence for mechanisms involving activation of PI3K signaling and impairment of p53 on urinary bladder carcinogenesis.	J Toxicol Pathol.	24	1-12	2011
<u>Wei M</u> , <u>Wanibuchi H</u> , Nakae D, Tsuda H, Takahashi H, Hirose M, Totsuka M, Tatematsu M, Fukushima S.	Low-dose carcinogenicity of 2-amino-3-methylimidazo[4,5-f]quinoline in rats: Evidence for the existence of no-effect levels and a mechanism involving p21Cip/WAF1.	Cancer Sci.	102	88-94	2011
Kakehashi A, Ishii N, Shibata T, <u>Wei M</u> , Okazaki E, Tachibana T, Fukushima S, <u>Wanibuchi H.</u>	Mitochondrial prohibitins and septin 9 are implicated in the onset of rat hepatocarcinogenesis.	Toxicol Sci.	119	61-72	2011
Chusiri Y, Wongpoomchai R, Kakehashi A, <u>Wei M</u> , <u>Wanibuchi H</u> , Vinitketkumnuan U, Fukushima S.	Non-genotoxic mode of action and possible threshold for hepatocarcinogenicity of Kojic acid in F344 rats.	Food Chem Toxicol.	49	471-176	2011
Hoshi H, Sawada T, Uchida M, Saito H, Iijima H, Toda-Agetsuma M, Wada T, Yamazoe S, Tanaka H, Kimura K, Kakehashi A, <u>Wei</u>	Tumor-associated MUC5AC stimulates in vivo tumorigenicity of human pancreatic cancer.	International journal of oncology.	38	619-627	2011

<u>M, Hirakawa K, Wanibuchi H.</u>					
Kohata Y, Fujiwara Y, Machida H, Okazaki H, Yamagami H, Tanigawa T, Watanabe K, Watanabe T, Tominaga K, <u>Wei M, Wanibuchi H, Arakawa T.</u>	Role of Th-2 cytokines in the development of Barrett's esophagus in rats.	J Gastroenterol.	46	883-893	2011
Chung K, Nishiyama N, Yamano S, Komatsu H, Hanada S, <u>Wei M, Wanibuchi H, Suehiro S, Kakehashi A.</u>	Serum AGR2 as an early diagnostic and postoperative prognostic biomarker of human lung adenocarcinoma.	Cancer Biomark.	10	101-107	2011
Yokohira M, Hashimoto N, Yamakawa K, Saoo K, Kuno T, <u>Imaida K.</u>	Lack of promoting effects from physical pulmonary collapse in a female A/J mouse lung tumor initiated with 4-(methyl- nitrosamino)-1-(3-pyridyl)-1- butanone (NNK) with remarkable mesothelial cell reactions in the thoracic cavity by the polymer.	Exp. Toxicol. Pathol.	63	181-185	2011
Okabe K, Hayashi M, Yoshida I, Nishimura K, Fukushima N, <u>Tsujiuchi T.</u>	Distinct DNA methylation patterns of lysophosphatidic acid receptor genes during rat hepatocarcinogenesis induced by a choline deficient L-amino acid defined diet.	Arch Toxicol.	85	1303-1310	2011
Hayashi M, Okabe K, Yamawaki Y, Teranishi M, Honoki K, Mori T, Fukushima N, <u>Tsujiuchi T.</u>	Loss of lysophosphatidic acid receptor-3 enhances cell migration in rat lung tumor cells.	Bioch. Bioph. Res. Com.	405	450-454	2011
Okabe K, Hayashi M, Yamawaki Y, Teranishi M, Honoki K, Mori T, Fukushima N, <u>Tsujiuchi T.</u>	Possible involvement of lysophosphatidic acid receptor-5 gene in the aquisition of growth advantage of rat tumor cells.	Mol Carcinog	50	635-642	2011
Okabe K, Kato K, Teranishi M, Okumura M, Fukui R, Mori T,	Induction of lysophosphatidic acid receptor-3 by 12-O-tertadecanolphorbol-13 -acetate stimulates cell migration of rat liver cells.	Cancer Lett	309	236-242	2011

Fukushima N, Tsujuchi T.					
Nakatani S, <u>Wei M</u> , Ishimura E, Kakehashi A, Mori K, Nishizawa Y, Inaba M, <u>Wanibuchi</u> H.	Proteome analysis of laser microdissected glomeruli from formalin-fixed paraffin-embedded kidneys of autopsies of diabetic patients: nephronectin is associated with the development of diabetic glomerulosclerosis.	Nephrol Dial Transpl.	27	1889-1897	2012
Hayashi M, Okabe K, Kato K, Okumura M, Fukui R, Fukushima N, <u>Tsujuchi T.</u>	Different function of lysophosphatidic acid receptors in cell proliferation and migration of neuroblastoma cells.	Cancer Lett	316	91-96	2012
Okabe K, Hayashi M, Kato K, Okumura M, Fukui R, Honoki K, Fukushima N, <u>Tsujuchi T.</u>	Lysophosphatidic acid receptor-3 increases tumorigenicity and aggressiveness of rat hepatoma RH7777 cells.	Mol. Carcinog		DOI: 10.1002/ mc.2185 1.	2012
Kato K, Fukui R, Okabe K, Tanabe E, Kitayoshi M, Fukushima N, <u>Tsujuchi T.</u>	Constitutively active lysophosphatidic acid receptor-1 enhances the induction of matrix metalloproteinase-2..	Biochem. Biophys. Res. Commun.	417	790-793	2012
Nakatani S, Kakehashi A, Ishimura E, Yamano S, Mori K, <u>Wei M</u> , Inaba M, <u>Wanibuchi H.</u>	Targeted proteomics of isolated glomeruli from the kidneys of diabetic rats: Sorbin and SH3 domain containing 2 is a novel protein associated with diabetic nephropathy.	Exp Diabetes Res.		doi:10.1 155/201 1/97935 4	2012
Xie XL, <u>Wei M</u> , Kakehashi A, Yamano S, Tajiri M, <u>Wanibuchi H.</u>	2-amino-3-methylimidazo [4,5-f] quinoline (IQ) promotes mouse hepatocarcinogenesis by activating transforming growth factor- $\beta$ and Wnt/ $\beta$ -catenin signaling pathways.	Toxicol Sci.	125	392-400	2012

