

TABLE I (Continued)

No	Patient	Year of birth	Sex	Exchange blood transfusion during newborn period	Thrombocytopenia during childhood	ADAMTS13:AC (%)	Disease-causing ADAMTS13 gene mutations	Age of TTP diagnosis	Prophylactic FFP infusion	Remarks	Ref.
									From when		
16	M3	1969	F	—	—	< 0.5	C-Hetero p.R193W/ p.R349C	33 y	— —		[50]
17	M4	1971	F	—	—	< 0.5	C-Hetero p.R193W/ p.R349C	30 y	— —		[50]
18	N6	1986	F	+	+	< 0.5	C-Hetero p.H234R/ c.3220delTACC	4 y	+ 4 y		[17]
19	O4	1958	F	—	—	< 0.5	C-Hetero p.I178T/ p.Q929X	26 y	+ 26 y		[50]
20	P3	1971	M	—	+	< 0.5	C-Hetero p.C908Y/ p.C322G, p.T323R, p.F324L	3 y	+ 21 y		[42]
21	Q1	1983	M	+	+	< 0.5–0.7	C-Hetero p.G227R/ p.C908Y	6 y	+ 11 y		[56]
22	Q2	1988	M	+	+	< 0.5	C-Hetero p.G227R/ p.C908Y	2 y	+ 7 y		[56]
23	R5	1982	F	—	+	< 0.5	C-Hetero p.R193W/ p.A606P	23 y	+ 23 y		[50]
24	S3	1982	F	—	+	0.9	Not determined	4 y	+ *		
25	T4	1981	F	+	+	< 0.5	Homo c.3220delTACC	1 m	+ *		[56]
26	U3	1990	F	+	+	< 0.5	Homo c.2259delA	4 m	+ *		[56]
27	V3	1983	F	+	+	< 0.5	C-Hetero p.W1081X/ p.R193W	6 y	+ 6 y		[56]
28	W4	1990	F	—	+	< 0.5	C-Hetero p.G550R/Not determined	15 y	+ 15 y		[56]
29	X5	1963	F	—	—	< 0.5	Not determined	40 y	— —		
30	Y3	1960	F	—	+	< 0.5	C-Hetero p.G385E/ p.R1206X	45 y	+ 45 y		[56]
31	Z3	1971	F	—	+	< 0.5	Homo p.R193W	25 y	— —		[50]
32	AA3	1987	F	—	—	< 0.5	Not determined	19 y	— —		
33	BB3	1947	M	—	—	< 0.5	Homo p.R193W	55 y	— —		[56]
34	CC5	2004	M	+	+	< 0.5	C-Hetero p.Q723K/ p.R398C	2 y	+ 2 y		[56]

H. Yagi, M. Matsumoto, Y. Fujimura

Please cite this article in press as: Hideo Yagi et al., Paradigm shift of childhood thrombotic thrombocytopenic purpura with severe ADAMTS13 deficiency, Presse Med (2012), doi: 10.1016/j.pmed.2011.10.027.

TABLE 1 (Continued)

No	Patient	Year of birth	Sex	Exchange blood transfusion during newborn period	Thrombocytopenia during childhood	ADAMTS13:AC (%)	Disease-causing ADAMTS13 gene mutations	Age of TTP diagnosis	Prophylactic FFP infusion	Remarks	Ref.
35	DD5	2007	F	-	+	< 0.5	C-Hetero p.R268P/p.Y304C	1 m	-		[56]
36	EE4	2003	M	+	+	< 0.5	Homo c.2259delA	4 y	-		[56]
37	FF3	1991	F	-	+	< 0.5	Homo p.Q449X	6 y	+ 6 y		[56]
38	GG2	1931	M	-	-	2.4-3.4	Homo p.C1024R	63 y	+ 63 y	Dead (stroke at the age of 79)	[56]
39	HH4	2003	F	+	+	< 0.5	C-Hetero p.Q449X/c.4119delG	1 y	-		[56]
40	II3	1977	F	+	+	< 0.5	Not determined	9 m	+ 10 y		
41	JJ3	1980	M	-	+	< 0.5	C-Hetero c.1885delT/p.C908Y	12 y	+ 25 y	Hemodialysis	[56]

C-Hetero: compound heterozygotes; Homo: homozygotes.

There have been two fatal USS cases, one is the above-mentioned USS-C3 and the other is patient USS-I4 (male, born in 1972), whose natural history was previously described in detail [51]. Briefly, patient USS-I4 was diagnosed with late-onset USS when he was 50 years old. The next year he received a cholecystectomy and then experienced a bout of TTP, which led to renal insufficiency. He received extensive treatment, including PE and hemodialysis but did not improve, and he died of renal insufficiency at the age of 51 years.

#### ADAMTS13 gene analysis

The parents of USS patients are usually asymptomatic carriers, and a major population of patients from unrelated parents is a compound heterozygote, while a minor population of patients from related parents is a homozygote [42,43,51-56].


We performed ADAMTS13 gene analyses in 38 out of 41 USS patients and disease-causing mutations were identified in 37 patients: nine with homozygous and 28 with compound heterozygous ADAMTS13 gene mutations. Furthermore, five of these 37 patients were siblings. Therefore, within 64 [2 × (37 - 5)] allelic numbers (*n*) for ADAMTS13 gene mutations, the three most frequently found mutations were in the following order: p.R193W (*n* = 8), p.Q449X (*n* = 5), and p.C908Y (*n* = 4). All these mutations were unique to Japanese individuals, perhaps to East-Asians, and were totally different from Europeans and white and black Americans. In addition, to date, we have not found an apparent association between specific ADAMTS13 mutations and clinical phenotypes. However, Camilleri et al. [57] reported that some single nucleotide polymorphisms in the ADAMTS13 gene could modulate ADAMTS13:AC and its secretion, indicating that further investigations are required.

Patient USS-GG2 (male, born in 1931) suddenly developed a bout of TTP when he was 63 years old. After this incident, he had repeated TTP bouts and required prophylactic FFP infusions under a clinical diagnosis of CR-TTP. Even under these circumstances, he developed a cerebellar infarction at 76 years of age. During the infusion intervals, ADAMTS13:AC was often measured and determined to be 2.4-3.4% of the normal levels but ADAMTS13:INH was not detected. Most recently, the patient was diagnosed with USS with the homozygous missense mutation C1024R based on an ADAMTS13 gene analysis (unpublished).

#### Treatment

Except for exchange blood transfusions to treat jaundice in newborns, USS patients usually respond well to small FFP infusions. Therefore, the question arises; what is the best marker for deciding this indication? As suggested above, mild thrombocytopenia seems to occasionally occur in USS patients during childhood, but this condition might be overlooked

**Family  
USS-L**



**FIGURE 3**  
**Family pedigree and ADAMTS13 analyses in family USS-L**

The propositus is L2, and L3 is her younger sister. Both siblings had an abortion along with thrombocytopenia of an unknown etiology. When propositus L2 became pregnant the second time, she had mild thrombocytopenia and her physician recommended low-dose aspirin, which enabled her to maintain the pregnancy until 30 weeks of gestation. However, it was uncertain why and how aspirin worked in this occasion. She delivered a live but premature baby by caesarean section. Then, she successfully bore three more children with the same treatment. All of the babies were premature and alive. During childhood, L2 had no episodes of thrombocytopenia, but L3 was diagnosed with ITP at 3 years of age. At 25 years of age, propositus L2 was diagnosed with USS based on an analysis of ADAMTS13:AC and ADAMTS13:INH. In addition, an ADAMTS13 gene analysis provided solid evidence that the two siblings are compound heterozygotes for ADAMTS13 gene mutations (p.R125fsx6/p.Q1302X). Squares and circles indicate males and females, respectively, and an arrow with P indicates the proposita. Closed circles and closed squares represent USS patients. The half-closed circles and squares represent asymptomatic carriers. The cross indicates deceased (cited from [50] with a slight modification).

because of the paucity of clinical signs. Thus, mild thrombocytopenia alone may not be a good marker. However, in clinical practice, some USS patients receive prophylactic FFP infusions (5–10 mL/kg BW) every 2–3 weeks because the half-life of ADAMTS13:AC in the plasma is thought to be 2–3 days, while other patients receive FFP infusions only when acute TTP bouts develop. In our registry, 26 of 41 (63%) USS patients received prophylactic FFP infusions. Currently, USS patients receive FFP infusions based on the physician's observations and the frequency of TTP bouts. However, the efficacy of prophylactic FFP infusions needs to be more precisely evaluated over a long observation period because our two patients who developed renal insufficiency had been

receiving FFP infusions since they were clinically diagnosed with CR-TTP.

One serious adverse effect of repeated plasma infusions is that nine out of 41 (22%) USS patients were infected with hepatitis C virus. In this regard, virus-free rADAMTS13 preparations would be a promising product for USS patients.

**Acquired thrombotic thrombocytopenic purpura/ADAMTS13:AC deficiency**

Figure 4 shows age and gender distribution of acquired TTP. Of 195 patients with acquired TTP and a severe ADAMTS13:AC deficiency due to ADAMTS13:INH, 17 (6%) were childhood patients, including 14 with acquired idiopathic TTP (ai-TTP)




FIGURE 1

**Age and gender distribution of patients with acquired idiopathic (ai)-thrombotic thrombocytopenic purpura (TTP) and connective tissue disease (CTD)-associated TTP**

The upper panel shows the age and gender distribution of 195 ai-TTP patients who were registered at Nara Medical University during 1998–2008 and were determined to have a severe ADAMTS13:AC deficiency and be positive for ADAMTS13:INH. The largest population peak is found at approximately 60 years old, and we identified 14 patients (14/195, 7.2%) aged less than 15 years. Of note, five were very young children below 2 years of age.

The lower panel shows the age and gender distribution of 46 CTD-TTP patients who were registered at our institution during the same time period and were determined to have a severe ADAMTS13:AC deficiency and be positive for ADAMTS13:INH. This CTD-TTP patient population is widely distributed across ages, but is more common in patients between 30–60 years old. There is an apparent female predominance (41/46, 89%) in CTD-TTP patients in this registry, and three childhood patients were identified.

and three with connective tissue disease (CTD)-associated TTP. Of note, the former group included five patients who were less than 2 years of age and were initially misdiagnosed with other thrombocytopenic disorders, such as ITP, HUS, and hemophagocytic syndrome (HPS). It is important to examine and compare the detailed clinical features of these TTP patients to adulthood patients in order for physicians to ascertain that TTP is not a rare childhood disease. Therefore, we herein describe the clinico-laboratory features of these five young infants with ai-TTP and three childhood patients with CTD-associated TTP.

**Idiopathic (ai)- thrombotic thrombocytopenic purpura**

Table II summarizes the features of the 14 childhood patients with ai-TTP, including five infantile patients. Case 1 was previously reported [58] and case 5 was more recently described [59]. Interestingly, in contrast to the adulthood ai-TTP

patients with a severe ADAMTS13:AC deficiency, the childhood patients had a slightly male predominance (female:male = 5:9). Five patients (5/14, 33%) had apparent prodromal illnesses, such as an upper respiratory tract infection (n = 3), Rotavirus infection (n = 1), or urinary tract infection (n = 1). Ten patients (10/14, 67%) had neurological findings, including headache (n = 5), altered mental status (n = 4), hemiparesis (n = 1), seizures (n = 1), and vision disturbance (n = 1). These patients exclusively presented with renal involvement (11/14, 73%) and fever (13/14, 93%). All the patients had hemolytic anemia (Hb, 4.5–11.3 g/dL) and thrombocytopenia (platelet, 7–38 × 10<sup>9</sup>/L), but their serum creatinine levels (Cr, 0.19–1.0 mg/dL) remained within the normal range. Most of the childhood patients had five clinical signs that are characteristic of classic TTP ('pentad'), but six patients, including five young infants aged below 2 years, were initially misdiagnosed with other thrombocytopenic disorders, such as ITP (n = 2), HUS (n = 2), HPS (n = 1), and paroxysmal

H Yagi, M Matsumoto, Y Fujimura

TABLE II

Clinical features in childhood patients with acquired TTP with severe ADAMTS13 deficiency

Case	Age	Sex	Prodromal illness	Initial diagnosis	Clinical findings on admission					
					Neurological symptom	Renal involvement	Cr (mg/dL)	Fever	Hb (g/dL)	Platelet ( $\times 10^9/L$ )
1	9 m	F	Rotavirus	HUS	Altered mental status	Yes	0.3	Yes	4.5	2
2	19 m	F	URI	IITP	No	Yes	0.19	Yes	11.3	38
3	19 m	M	URI	HUS	Altered mental status	Yes	1	Yes	5.4	9
4	12 m	M	No	HPS	Hemiparesis	Yes	0.31	Yes	8.5	38
5	8 m	M	No	IITP	No	No	0.19	Yes	8.7	25
6	7 y	M	No	TTP	Altered mental status	No	0.32	Yes	5.4	7
7	10 y	M	UTI	TTP	No	Yes	0.4	No	10.5	19
8	11 y	M	No	TTP	Headache	Yes	0.7	Yes	8.8	3
9	11 y	F	URI	TTP	No	Yes	0.6	Yes	5.5	6
10	13 y	F	No	TTP	Headache	No	0.58	Yes	5.7	12
11	14 y	F	No	TTP	Altered mental status, convulsion	No	0.4	Yes	7.8	6
12	15 y	M	No	PNH	Headache	Yes	1	Yes	10.5	17
13	15 y	M	No	TTP	Headache	Yes	0.91	Yes	8.1	11
14	14 y	M	No	TTP	Headache, visual disturbance	Yes	1.05	Yes	7	28
15	8 y	F	SLE	DIC	Altered mental status, headache	Yes	0.44	Yes	8.8	38
16	11 y	F	MCTD	TTP	Headache	Yes	0.6	Yes	11.9	43
17	12 y	F	SLE	TTP	No	No	0.61	Yes	5.6	1

Case	ADAMTS13: AC (%)	ADAMTS13: INH (BU/ml)	Treatments				Outcome		Clinical course remarks	Ref.
			PE (times)	FFP infusions	Immunosuppressive agents	Platelet transfusion	Relapse	Prognosis		
1	< 0.5	> 100	19	No	SP	Yes	No	Alive	Cerebral infarction	[58]
2	< 0.5	4.3	3	Yes	SP, PSL	Yes	Yes	Dead		
3	< 0.5	2.3	3	Yes	PSL	Yes	No	Alive		
4	< 0.5	1.7	2	No	PSL	Yes	No	Alive	Cerebral infarction	
5	< 0.5	4.8	6	No	SP, PSL	Yes	No	Alive		[59]
6	< 0.5	2.1	3	No	Rituximab	No	No	Alive		
7	< 0.5	0.5	No	No	SP, PSL, MZR	Yes	No	Alive		
8	< 0.5	1.3	No	Yes	SP	No	No	Alive		
9	< 0.5	5.6	5	No	PSL	Yes	No	Alive		
10	< 0.5	2	9	No	SP, PSL	No	No	Alive		

TABLE II (Continued)

Case	ADAMTS13: AC (%)	ADAMTS13: INH (BU/ml)	Treatments				Outcome		Clinical course remarks	Ref.
			PE (times)	FFP infusions	Immunosuppressive agents	Platelet transfusion	Relapse	Prognosis		
11	< 0.5	3.2	17	No	SP, PSL, VCR	No	No	Alive		
12	< 0.5	34	39	No	PSL, VCR, CSA	No	Yes	Alive		
13	< 0.5	2.3	3	No	SP, PSL, AZI	No	No	Alive		
14	< 0.5	6.8	30	Yes	PSL	No	No	Alive		
15	< 0.5	1.2	5	Yes	SP, PSL, CY	No	No	Alive		
16	< 0.5	1.8	3	No	SP, PSL	No	No	Alive		
17	< 0.5	0.7	9	Yes	SP, PSL	No	No	Alive		

Cr: creatinine; Hb: hemoglobin; ADAMTS13:AC: ADAMTS13 activity; ADAMTS13:INH: ADAMTS13 inhibitor; PE: plasma exchange; FFP: fresh frozen plasma; URI: upper respiratory infection; UTI: urinary tract infection; SLE: systemic lupus erythematosus; MCTD: mixed connective tissue disease; TTP: thrombotic thrombocytopenic purpura; ITP: idiopathic thrombocytopenic purpura; HUS: hemolytic uremic syndrome; HPS: hemophagocytic syndrome; PNH: paroxysmal nocturnal hemoglobinuria; SP: steroid pulse; PSL: prednisolone; MZR: mizoribine; VCR: vincristine; CSA: cyclosporine A; AZI: azathioprine; Cy: cyclophosphamide.

nocturnal hemoglobinuria (PNH,  $n = 1$ ). After analyzing ADAMTS13, they were all correctly diagnosed with ai-TTP. Of these 14 childhood patients with ai-TTP, 13 received plasma exchange (PE, 2–39 times, median 5 times), including four patients who subsequently received a FFP infusion. They also received immunosuppressive therapy, including steroid pulse ( $n = 7$ ), prednisolone ( $n = 12$ ), vincristine ( $n = 2$ ), cyclosporin ( $n = 1$ ), azathioprine ( $n = 1$ ), mizoribine ( $n = 1$ ), and rituximab ( $n = 1$ ). As an adjunctive therapy, the patients were given intravenous immune globulin ( $n = 3$ ) or an antiplatelet agent ( $n = 2$ ). Of note, seven patients received platelet transfusions before or after they were diagnosed with acquired TTP. In five of the seven patients who received platelet transfusions, there were no apparent serious complications. However, case 1 developed general convulsions soon after the platelet transfusion, and case 2 died from bleeding without appreciable hemostatic effects from the platelet transfusion. As consequence, 13 out of 14 childhood patients with ai-TTP achieved one clinical remission, but two patients relapsed, including one who died. We think that clinicians should be aware of the existence of ai-TTP during very early childhood, and herein we present a short summary for each of these five infants with ai-TTP.

### Case 1

In March 2000, a 9-month-old girl presented with a fever. She subsequently showed loss of appetite, a drop in physical activity, a pale complexion, and vomiting followed by diarrhea related to a Rotavirus infection. On the following day, these symptoms continued and generalized petechiae appeared. She was taken to a family doctor who determined that she had

severe anemia and thrombocytopenia. As a result, she was admitted to a nearby hospital for treatment. Upon admission, she was drowsy and her laboratory findings showed severe anemia (Hb, 4.5 g/dL), thrombocytopenia (platelet,  $2.0 \times 10^9/L$ ), hyperbilirubinemia (total bilirubin, 2.6 mg/dL), and an elevated LDH level (2,925 IU/L). Both direct and indirect Coombs' tests were negative, and fragmented RBCs were detected in the blood film. The hemostatic tests showed the following: prothrombin time (PT, 14.0 sec), activated PTT (35.9 sec), fibrinogen (268 mg/dL), thrombin-antithrombin complex (TAT, 31.7  $\mu\text{g/L}$ ), D-dimer (7.14  $\mu\text{g/mL}$ ), and fibrin degradation product (FDP, 82.3  $\mu\text{g/mL}$ ). The levels of blood urea nitrogen (BUN) and creatinine were within the normal ranges (25 mg/dL and 0.3 mg/dL, respectively). Neither Shiga-like toxin nor *E. coli* O157:H7 was detected in her stool. However, she had proteinuria, hematuria, and marked petechiae on her body due to thrombocytopenia. She was tentatively diagnosed with HUS, and she received five units of platelet transfusion. Soon after completing the platelet transfusion, she developed generalized convulsions followed by right hemiplegia, and therefore, PE therapy was immediately instituted. On the following day, both CT and MRI examinations of her brain revealed a diffuse hemorrhagic infarction in the left posterior region. The PE therapy was continued for the next 37 days on a total of 19 occasions, along with steroid pulse therapy and high-dose intravenous immunoglobulin (IVIg) infusions until clinical improvements were noted. The ADAMTS13:AC and ADAMTS13:INH titers measured by the VWFM assay were less than 3% and greater than 100 BU/mL, respectively (later, both values were re-evaluated by the chromogenic act-ELISA using deep-frozen plasmas, and the respective data were

H Yagi, M Matsumoto, Y Fujimura

less than 0.5% and greater than 100 BU/mL [58]). Now, almost 10 years have passed, and the patient is apparently healthy with a minimal sequela of the right hemiplegia.

### Case 2

In July 2000, a 19-month-old girl presented with a fever and cough. The next day, she was taken to a family doctor and then transferred to a nearby hospital because of thrombocytopenia (platelet,  $21 \times 10^9/L$ ). Upon admission, her laboratory data revealed slight anemia (Hb, 11.3 g/dL), thrombocytopenia (platelet,  $38 \times 10^9/L$ ), and an elevated LDH level (994 IU/L). An analysis of a bone marrow aspiration showed no abnormalities, and therefore she was suspected to have ITP. She was administered high-dose IVIG with steroid therapy but her platelet count did not increase. Her platelet count slightly increased soon after the platelet transfusions, while the number of schistocytes in the blood films gradually increased. This patient never had renal dysfunction or neurological signs. Thus, her physician suspected that the patient had USS but did not measure ADAMTS13:AC. The patient was given an infusion of 80 mL of FFP, but her platelet count did not increase. During this period, she was alert and no clinical deterioration was noted. Three months after admission, plasma samples from this patient were sent to our laboratory for ADAMTS13:AC and ADAMTS13:INH testing. Based on the results of the VWFM assays, the patient was diagnosed with a severe ADAMTS13:AC deficiency (< 3%) with ADAMTS13:INH (4.0 BU/mL). However, in those days we were unable to clearly determine whether this patient had USS and developed alloantibodies or acquired TTP with autoantibodies to ADAMTS13. The patient was not given PE therapy because she did not show any clinical deterioration during the subsequent 3 months. Therefore, she was discharged and then carefully observed at the outpatient clinic. However, 1 month after discharge, she was re-admitted to the hospital and received PE therapy because of exacerbated anemia and thrombocytopenia. However, her clinical signs did not improve, even after whole blood exchange therapy. Thus, she was treated with RBC and platelet transfusions, but 2 weeks later she fell into coma and died of tracheal bleeding, which was 8 months after her first hospital admission (plasma ADAMTS13:AC and ADAMTS13:INH in this patient were measured only on one occasion. In recent years, both values were re-evaluated by the chromogenic act-ELISA using deep-frozen plasma samples, and the data were less than 0.5% and 4.3 BU/mL, respectively).

### Case 3

In July 2002, a 19-month-old boy developed a low-grade fever and cough followed by petechiae. He was taken to a family doctor because his nasal bleeding did not stop. The doctor noted thrombocytopenia and anemia and suspected HUS, and the patient was transferred to a local hospital. He had mild thrombocytopenia (platelet,  $70 \times 10^9/L$ ) soon after birth,

which spontaneously improved. Upon admission, he was drowsy and his laboratory data showed anemia (Hb, 5.4 g/dL), thrombocytopenia (platelet,  $9 \times 10^9/L$ ), elevated LDH (1991 IU/L), and proteinuria. He was administered FFP infusions, steroid therapy, and IVIG. A platelet transfusion was performed but his platelet counts did not significantly increase. Since he was negative for DIC markers, the patient was clinically diagnosed with TTP and then administered PE therapy. After three consecutive PE therapies, he became alert, recovered, and his laboratory markers returned to normal levels. The ADAMTS13:AC and ADAMTS13:INH titers were measured by classic VWFM assays using frozen plasma that was obtained before the PE therapy was administered, and the results were less than 3% and 2.3 BU/mL, respectively (later, chromogenic act-ELISA gave values of less than 0.5% and 2.3 BU/mL, respectively). His plasma ADAMTS13:AC deficiency with ADAMTS13:INH continued for more than 6 months but without appreciable clinical manifestations. After 4 years, ADAMTS13:AC (77%) had normalized and ADAMTS13:INH (< 0.5 BU/ml) was absent.

### Case 4

In June 2002, a 13-month-old boy developed a fever followed by dark urine and diarrhea. He was taken to a nearby clinic, where he was determined to have leukocytosis (WBC, 12,000/ $\mu$ L), anemia (Hb, 7.2 g/dL), and thrombocytopenia (platelet,  $46 \times 10^9/L$ ). In addition, a peripheral blood film showed phagocytosis and therefore the patient was diagnosed with suspected HPS. He was transferred to a university hospital where platelet transfusions were performed twice for two consecutive days, but his platelet counts only transiently increased. Soon after the second platelet transfusion, a bone marrow aspiration was performed, but the HPS diagnosis was not confirmed. On the other hand, there was a transient increase in his platelet count (platelet,  $40 \times 10^9/L$ ) after he was infused with a small amount of FFP, and therefore the physician suspected a diagnosis of USS. Therefore, the patient received a daily plasma infusion therapy for the next 5 days. However, hematuria developed followed by right hemiplegia. An MRI revealed a hemorrhagic infarction (3  $\times$  4 cm) in the left parieto-occipital region. Based on the clinical course, he was eventually diagnosed with ai-TTP. After he received PE therapy for two consecutive days with orally administered prednisolone, his clinical conditions rapidly improved and his laboratory findings recovered. After his recovery, the ADAMTS13:AC and ADAMTS13:INH levels were tested using the classic VWFM assay and deep-frozen plasma samples that were obtained before the PE therapy, and the results were less than 3% and 1.9 BU/mL, respectively (later, these values were re-examined with the chromogenic act-ELISA, and the results were less than 0.5% and 1.9 BU/mL, respectively). He subsequently improved and was discharged. Three years later, his plasma

TABLE III

Comparison of clinical features and outcomes between childhood and adulthood patients with acquired idiopathic (ai)-TTP

Ai-TTP (n = 195)	Childhood patients (n = 14)	Adulthood patients (n = 181)
Age at the onset of TTP bouts (years old), Median (25, 75 percentile)	11 (1.6, 14)	57 (41, 65)
Female (%)	35.7	55.2
‘Pentad’		
(1) Platelet count ( $\times 10^9/L$ ), Median (25, 75 percentile)	15 (8, 24)	10 (7, 17)
(2) Hemoglobin (g/dL), Median (25, 75 percentile)	8.0 (5.6, 8.8)	7.5 (6.3, 8.8)
(3) Renal involvement (%)	71.4	75.7
Serum creatinine (mg/dL), Median (25, 75 percentile)	0.5 (0.3, 0.9)	1.0 (0.7, 1.3)
(4) Central nervous system involvements (%)	71.4	79.6
(5) Fever ( $\geq 37.0^\circ C$ ) (%)	92.9	69.6
Times of plasma exchange	5.5 (3.0-17.5)	ND
Mortality in the current episode of TTP bouts (%)	7.1	15.5

ND: not determined.

ADAMTS13:AC had normalized. At present, he has fully recovered and has no residual right hemiplegia.

### Case 5

In January 2005, a 9-month-old boy with generalized petechiae and a fever was referred to a local hospital, where he was determined to have thrombocytopenia (platelet,  $9 \times 10^9/L$ ). He was admitted to a university hospital and diagnosed with acquired TTP based on ADAMTS13:AC ( $< 3\%$ ) and ADAMTS13:INH titers (2.8 BU/mL) that were determined using the classic VWFM assays. (Later, these values were re-examined by the chromogenic act-ELISA, and the results were less than 0.5% and 4.8 BU/mL, respectively). After he was diagnosed with ai-TTP, he was administered PE therapy for six consecutive days at a different hospital. His clinical symptoms rapidly improved, but the increase in platelet counts was only transient and his platelet count was consistently lower than  $10 \times 10^9/L$ . To prevent serious bleeding complications, the physician administered oral prednisolone, together with continuous low-dose platelet transfusions. Two months later, he was discharged, despite having an ADAMTS13:AC deficiency with ADAMTS13:INH that lasted for at least 8 months. Two years later, we were able to examine the plasma ADAMTS13:AC and ADAMTS13:INH in this patient, and determine that ADAMTS13 had fully normalized [59].

Table III compares the clinical features and outcomes of the childhood patients (n = 14) and adulthood patients (n = 181) with ai-TTP in our registry [13].

### Connective tissue disease-associated thrombotic thrombocytopenic purpura

In 1939, Gitlow and Goldmark [60] first reported a close relationship between ‘classic’ TTP and systemic lupus erythematosus (SLE). In 1999, Brenner et al. [61] described five patients with childhood-onset ‘classic’ TTP and reviewed 30 other patients who were previously described in the literature. According to their analysis, nine (9/35, 26%) fulfilled four or more ACR criteria for SLE and eight (8/35, 23%) had incipient SLE. Interestingly, of the five patients who were initially diagnosed with ‘classic’ idiopathic TTP in their laboratory, three were diagnosed with SLE within 3 years, and the remaining two patients fulfilled the ACR classification criteria for SLE within 4 years of disease onset. However, at that time, ADAMTS13:AC assays were not generally available, and therefore no data on ADAMTS13 was provided in their report.

In our registry of 919 patients with TMAs, 221 had CTD-associated TMA, of which 92 had SLE-associated TMA [13]. For the 221 CTD-TMA and/or 92 SLE-TMA patients, the number of patients with a severe ADAMTS13:AC deficiency with ADAMTS13:INH was 46 and 24, respectively. Furthermore, within the 221 patients with CTD-TMA, 11 developed the disease in childhood (less than 15 years of age), including eight patients with SLE, 1 with RA (rheumatoid arthritis), and two with MCTD (mixed connective tissue disease), in whom three had a severe ADAMTS13:AC deficiency. These three patients included two SLE- and one MCTD-associated TTP patients, and they uniformly had relatively low titers of ADAMTS13:INH (0.7–1.8 BU/mL) at



H Yagi, M Matsumoto, Y Fujimura

the onset of TTP, which slightly differed from those with ai-TTP (table II). Here we briefly describe these three childhood patients with CTD-associated TTP (cases 15–17) due to their relevance in clinical practice.

#### Case 15

In April 2005, a 7-year-old girl was determined to have proteinuria and occult blood in her urine based on a school health examination. The patient was admitted to a nearby university hospital for further examination in June 2005, where she was diagnosed with SLE (Lupus nephritis) based on her clinical manifestations and the following laboratory findings: proteinuria, positive for anti-nuclear antibodies and anti-double stranded DNA antibodies, and low complementemia. She was treated with prednisolone, mizoribine, and azathioprine. Because her clinical signs significantly improved with these treatments, she was discharged in September 2005.

In April 2006 (8 years of age), this patient noticed proteinuria and hematuria based on a self-examination at home. The next day, she was admitted to the same hospital. Her laboratory data at the second admission were as follows: Hb (8.8 g/dL), LDH (1608 IU/L), platelet ( $2 \times 10^9/L$ ), PT (11.0 sec), PTT (31.1 sec), fibrinogen (355 mg/dL), antithrombin (140%), TAT (9.06  $\mu\text{g/L}$ ), D-dimer (5.25 ng/mL), FDP (7.9  $\mu\text{g/mL}$ ), and schistocytes in a peripheral blood smear. She had a DIC score of 6 according to the DIC diagnostic criteria from the International Society of Thrombosis and Haemostasis [15]. As a result, she was initially treated with nafamostat, but there were no clinical improvements. ADAMTS13:AC and ADAMTS13:INH assays were performed for a differential diagnosis of TTP, and the results were less than 0.5% and 1.2 BU/mL, respectively. Thus, she was diagnosed with SLE-associated TTP and treated with PE (five times), steroid pulse therapy, and cyclophosphamide. These treatments saved her life, and to date, she has not had a relapse of TTP.

#### Case 16

In July 2006, an 11-year-old girl developed general fatigue, headache, and vomiting. Two days later, she was admitted to a local hospital where laboratory tests indicated the following: Hb (11.9 g/dL), LDH (1636 IU/L), total bilirubin (9.3 mg/dL), platelet ( $33 \times 10^9/L$ ), PT (12.7 sec), PTT (34.0 sec), fibrinogen (290 mg/dL), antithrombin (> 75%), D-dimer (< 2 ng/mL), FDP (< 5  $\mu\text{g/mL}$ ), BUN (18.0 mg/dL), and schistocytes in a peripheral blood smear. For a differential diagnosis, her plasma ADAMTS13:AC and ADAMTS13:INH levels were determined to be less than 0.5% and 1.8 BU/mL, respectively. In addition, upon admission she simultaneously had Raynaud's phenomenon and was positive for anti-nuclear antibodies and anti-RNP antibodies. Thus, she was diagnosed with MCTD-associated TTP, and was administered PE with steroid pulse therapy starting on the third day of hospitalization. During the third PE, she had anaphylactic shock, perhaps related to the infused plasma.

Thus, she stopped the PE therapy and continued the steroid pulse therapy alone. As a result of these treatments, she recovered and on the hospital day 14 her ADAMTS13:AC increased to 67% of normal and ADAMTS13:INH became negative. To date, she has had no episodes of TTP.

#### Case 17

In October 2007, a 12-year-old girl suddenly developed jaundice with a fever. She was admitted to a nearby university hospital, and her routine laboratory data provided the following: Hb (5.6 g/dL), platelet ( $1 \times 10^9/L$ ), PT (14.0 sec), PTT (40.1 sec), fibrinogen (333 mg/dL), FDP (20.5  $\mu\text{g/mL}$ ), total bilirubin (5.5 mg/dL), and schistocytes in a peripheral blood smear. Upon admission, she had low levels of complement, and was positive for anti-nuclear antibodies, anti-double stranded DNA antibodies, and anti-SS-A antibodies. Plasma ADAMTS13:AC and ADAMTS13:INH were simultaneously measured by the act-ELISA and were less than 0.5% and 0.7 BU/mL, respectively. Based on these results, she was diagnosed with SLE-associated TTP and PE therapy was initiated. After three consecutive PE treatments with steroid pulse therapy, her platelet count increased. However, on hospital day 8, her platelet count decreased again, and her ADAMTS13:INH titer increased to 2.2 BU/mL. The PE therapy was re-initiated with steroid pulse therapy. A total of nine rounds of PE therapy and two courses of steroid pulse therapy resulted in remission on hospital day 23. At this time, ADAMTS13:AC and ADAMTS13:INH were 86% and less than 0.5 BU/mL, respectively. To date, she has had no TTP relapses.

#### Treatment of acquired thrombotic thrombocytopenic purpura

Plasma exchange (PE) is the first line therapy that was demonstrated to be effective in randomized clinical trials for acquired TTP [62]. PE removes ADAMTS13:INH, UL-VWFm, and hazardous cytokines from the circulation in TTP patients, and replenishes ADAMTS13 without circulatory overload. Corticosteroids are often used as an adjunctive treatment. In relapsing or refractory cases, other immunosuppressive drugs such as cyclosporine, cyclophosphamide, vincristine, and rituximab are empirically used. PE therapy should be initiated as soon as possible after TTP is diagnosed, but the onset of therapy tends to be delayed in childhood patients with acquired TTP because of difficult differential diagnoses, especially with HUS, unless ADAMTS13:AC is measured.

In regard to platelet transfusions in TTP patients with a severe ADAMTS13:AC deficiency, these transfusions have consistently been viewed as a contraindication because they may enhance thrombotic complications due to platelet aggregation and thrombus formation under high shear stress generated in the microvasculature. Our experience also partially supports this concept, but such adverse effects happened only on very few occasions. We believe that prophylactic platelet

transfusions should be avoided in TTP patients with a severe ADAMTS13:AC deficiency, but that platelet transfusions must be done if patients experience overt bleeding.

In our 17 childhood patients with acquired TTP, 15 patients were promptly treated with PE and corticosteroid therapy, and 16 children (94%) achieved a first remission. Recently, McDonald et al. [63] reported that the number of PE courses to first remission was higher in children (median, 22.5; range, 10–30) than in adults (median, 15.5; range, 3–93) [64], suggesting that childhood TTP may be more resistant to treatment. By contrast, our results indicated that patients with acquired TTP and a severe ADAMTS13:AC deficiency responded well to PE (median number of PE courses, 5.5; range, 2–39), but two patients (2/17, 11.8%) relapsed and one (1/17, 5.9%) died. Furthermore, in this study, we observed that the children with a high ADAMTS13:INH titer (> 5 BU) tended to require more frequent PE courses to achieve remission.

Fakhouri et al. [65] recently reported that adulthood TTP patients with high-titer ADAMTS13:INH could be successfully treated with a combination of PE and rituximab, a chimeric monoclonal antibody to CD20. The efficacy of rituximab in such patients is apparently due to a reduction in anti-ADAMTS13 IgG antibodies by depleting the patient's B-lymphocytes [65,66]. Recently, there have been many successful cases [67–69], and to date, no significant adverse effects have been reported. In our registry, only one childhood TTP patient (7 years old) with acquired TTP with ADAMTS13:INH was successfully treated with PE followed by rituximab, as shown in table II. However, the best choice or combination in regard to immunosuppressants

for treating children with acquired TTP and a severe ADAMTS13:AC deficiency needs to be carefully determined in future studies.

## Conclusion

The discovery of ADAMTS13 provided a breakthrough in our understanding of the mechanism of platelet thrombus formation under high shear stress and directly linked this enzyme to TTP pathogenesis in humans. Subsequently, the recent development of rapid and sensitive ADAMTS13 assays and their utilization in clinical practice have shown that the early- and late-onset phenotypes of USS are not different diseases and are likely affected by both acquired endogenous and exogenous circumstances. Furthermore, we have presented a novel category of ai-TTP that occurs during very early childhood (less than 2 years of age), which was perhaps totally overlooked or misdiagnosed before 2002 [39]. Thus, TTP should be recognized as a life-threatening generalized disease that not only occurs in adulthood, but also in childhood, causing a paradigm shift in our clinical understanding of TTP since the first discovery by Moschowitz in 1924.

Disclosure of interest: Yoshihiro Fujimura is a clinical advisory board for Baxter Bioscience.

Acknowledgments: This work was supported in part by research grants from the Ministry of Education, Culture, Sports, Science and Technology of Japan and from the Ministry of Health, Labor, and Welfare of Japan.

## References

- [1] Moake JL. Thrombotic microangiopathies. *N Engl J Med* 2002;347:589-600.
- [2] George JN. How I treat patients with thrombotic thrombocytopenic purpura. *Blood* 2010;116:4060-9.
- [3] Coppo P, Veyradier A. Thrombotic microangiopathies: towards a pathophysiology-based classification. *Cardiovasc Hematol Disord Drug Targets* 2009;9:36-50.
- [4] Coppo P, Schwarzinger M, Buffet M, Wynckel A, Clabault K, Presne C *et al*. Predictive features of severe acquired ADAMTS13 deficiency in idiopathic thrombotic microangiopathies: the French TMA reference center experience. *PLoS One* 2010;5:e10208.
- [5] Sadler JE. Von Willebrand factor, ADAMTS13, and thrombotic thrombocytopenic purpura. *Blood* 2008;112:11-8.
- [6] Furlan M, Robles R, Galbusera M, Remuzzi G, Kyrle PA, Brenner B *et al*. von Willebrand factor-cleaving protease in thrombotic thrombocytopenic purpura and the hemolytic-uremic syndrome. *N Engl J Med* 1998;339:1578-84.
- [7] Tsai HM, Lian EC. Antibodies to von Willebrand factor-cleaving protease in acute thrombotic thrombocytopenic purpura. *N Engl J Med* 1998;339:1585-94.
- [8] Moschowitz E. Hyaline thrombosis of the terminal arterioles and capillaries: a hitherto undescribed disease. *Proc N Y Pathol Soc* 1924;24:21-4.
- [9] Amorosi EL, Ullmann JE. Thrombotic thrombocytopenic purpura: report of 16 cases and review of the literature. *Medicine* 1966;45:139-59.
- [10] Torok TJ, Holman RC, Chorba TL. Increasing mortality from thrombotic thrombocytopenic purpura in the United States-analysis of national mortality data, 1968–1991. *Am J Hematol* 1995;50:84-90.
- [11] Gasser C, Gautier E, Steck A, Siebenmann RE, Dechslin R. Hemolytic-uremic syndrome: bilateral necrosis of the renal cortex in acute acquired hemolytic anemia. *Schweiz Med Wochenschr* 1955;85:905-9.
- [12] Karmali MA, Petric M, Lim C, Fleming PC, Arbus GS, Lior H. The association between idiopathic hemolytic uremic syndrome and infection by verotoxin-producing *Escherichia coli*. *J Infect Dis* 1985;151:775-82.
- [13] Fujimura Y, Matsumoto M. Registry of 919 patients with thrombotic microangiopathies across Japan: database of Nara Medical University during 1998–2008. *Intern Med* 2010;49:7-15.
- [14] Wada H, Wakita Y, Nakase T, Shimura M, Hiyoyama K, Nagaya S *et al*. Increased plasma-soluble fibrin monomer levels in patients with disseminated intravascular coagulation. *Am J Hematol* 1996;51:255-60.
- [15] Taylor FBJr, Toh CH, Hoots WK, Wada H, Levi M. Towards definition, clinical and laboratory criteria, and a scoring system for disseminated intravascular coagulation. *Thromb Haemost* 2001;86:1327-30.

H Yagi, M Matsumoto, Y Fujimura

- [16] Furlan M, Robles R, Lämmle B. Partial purification and characterization of a protease from human plasma cleaving von Willebrand factor to fragments produced by *in vivo* proteolysis. *Blood* 1996;87:4223-34.
- [17] Kinoshita S, Yoshioaka A, Park YD, Ishizashi H, Konno M, Funato M *et al.* Upshaw-Schulman syndrome revisited: a concept of congenital thrombotic thrombocytopenic purpura. *Int J Hematol* 2001;74:101-8.
- [18] Kato S, Matsumoto M, Matsuyama T, Isonishi A, Hiura H, Fujimura Y. Novel monoclonal antibody-based enzyme immunoassay for determining plasma levels of ADAMTS13 activity. *Transfusion* 2006;46:1444-52.
- [19] Kasper CK, Aledort L, Aronson D, Counts R, Edson JR, van Eys J *et al.* Proceedings: a more uniform measurement of factor VIII inhibitors. *Thromb Diath Haemorrh* 1975;34:612.
- [20] Zheng X, Chung D, Takayama TK, Majerus EM, Sadler JE, Fujikawa K. Structure of von Willebrand factor-cleaving protease (ADAMTS13), a metalloprotease involved in thrombotic thrombocytopenic purpura. *J Biol Chem* 2001;276:41059-63.
- [21] Uemura M, Tatsumi K, Matsumoto M, Fujimoto M, Matsuyama T, Ishikawa M *et al.* Localization of ADAMTS13 to the stellate cells of human liver. *Blood* 2005;106:922-4.
- [22] Suzuki M, Murata M, Matsubara Y, Uchida T, Ishihara H, Shibano T *et al.* Detection of von Willebrand factor-cleaving protease (ADAMTS-13) in human platelets. *Biochem Biophys Res Commun* 2004;313:212-6.
- [23] Turner N, Nolasco L, Tao Z, Dong JF, Moake J. Human endothelial cells synthesize and release ADAMTS-13. *J Thromb Haemost* 2006;4:1396-404.
- [24] Manea M, Kristoffersson A, Schneppenheim R, Solelem MA, Mathieson PW, Morgelin M *et al.* Podocytes express ADAMTS13 in normal renal cortex and in patients with thrombotic thrombocytopenic purpura. *Br J Haematol* 2007;138:651-62.
- [25] Matsumoto M, Chisuwa H, Nakazawa Y, Ikegami T, Hashikura Y, Kawasaki S *et al.* Liver transplantation rescues a deficient state of von Willebrand factor-cleaving protease activity in patients with liver cirrhosis due to congenital biliary atresia. *Blood* 2000;96:636a ([abstract]).
- [26] Uemura M, Fujimura Y, Matsumoto M, Ishizashi H, Kato S, Matsuyama T *et al.* Comprehensive analysis of ADAMTS13 in patients with liver cirrhosis. *Thromb Haemost* 2008;99:1019-29.
- [27] Fujimura Y, Matsumoto M, Yagi H. *Thrombotic microangiopathy*. Tokyo: Springer; 2008. (p. 625-39).
- [28] Padilla A, Moake JL, Bernardo A, Ball C, Wang Y, Arya M *et al.* P-selectin anchors newly released ultralarge von Willebrand factor multimers to the endothelial cell surface. *Blood* 2004;103:2150-6.
- [29] Akiyama M, Takeda S, Kokame K, Takagi J, Miyata T. Crystal structures of the noncatalytic domains of ADAMTS13 reveal multiple discontinuous exosites for von Willebrand factor. *Proc Natl Acad Sci U S A* 2009;106:19274-9.
- [30] Tandon NN, Rock G, Jameson GA. Anti-CD36 antibodies in thrombotic thrombocytopenic purpura. *Br J Haematol* 1994;88:816-25.
- [31] Davis AK, Makar RS, Stowell CP, Kuter DJ, Dzik WH. ADAMTS13 binds to CD36: a potential mechanism for platelet and endothelial localization of ADAMTS13. *Transfusion* 2009;49:206-13.
- [32] Yagi H, Konno M, Kinoshita S, Matsumoto M, Ishizashi H, Matsui T *et al.* Plasma of patients with Upshaw-Schulman syndrome, a congenital deficiency of von Willebrand factor-cleaving protease activity, enhances the aggregation of normal platelets under high shear stress. *Br J Haematol* 2001;115:991-7.
- [33] Zhang Q, Zhou YF, Zhang CZ, Zhang X, Lu C, Springer TA. Structural specializations of A2, a force-sensing domain in the ultralarge vascular protein von Willebrand factor. *Proc Natl Acad Sci U S A* 2009;106:9226-31.
- [34] Zanardelli S, Chion AC, Groot E, Lenting PJ, McKinnon TA, Laffan MA *et al.* A novel binding site for ADAMTS13 constitutively exposed on the surface of globular VWF. *Blood* 2009;114:2819-28.
- [35] Soejima K, Matsumoto M, Kokame K, Yagi H, Ishizashi H, Maeda H *et al.* ADAMTS-13 cysteine-rich/spacer domains are functionally essential for von Willebrand factor cleavage. *Blood* 2003;102:3232-7.
- [36] Klaus C, Plaimauer B, Studt JD, Dörner F, Lämmle B, Mannucci PM *et al.* Epitope mapping of ADAMTS13 autoantibodies in acquired thrombotic thrombocytopenic purpura. *Blood* 2004;103:4514-9.
- [37] Luken BM, Turenhout EA, Kaijen PH, Greuter MJ, Pos W, van Mourik JA *et al.* Amino acid regions 572-579 and 657-666 of the spacer domain of ADAMTS13 provide a common antigenic core required for binding of antibodies in patients with acquired TTP. *Thromb Haemost* 2006;96:295-301.
- [38] Pos W, Crawley JT, Fijnheer R, Voorberg J, Lane DA, Luken BM. An autoantibody epitope comprising residues R660, Y661, and Y665 in the ADAMTS13 spacer domain identifies a binding site for the A2 domain of VWF. *Blood* 2010;115:1640-9.
- [39] Fujimura Y, Matsumoto M, Yagi H, Yoshioaka A, Matsui T, Titani K. Von Willebrand factor-cleaving protease and Upshaw-Schulman syndrome. *Int J Hematol* 2002;75:25-34.
- [40] Moake JL, Rudy CK, Troll JH, Weinstein MJ, Colaninno NM, Azocar J *et al.* Unusually large plasma factor VIII: von Willebrand factor multimers in chronic relapsing thrombotic thrombocytopenic purpura. *N Engl J Med* 1982;307:1432-5.
- [41] Furlan M, Robles R, Solenthaler M, Wassmer M, Sandoz P, Lämmle B. Deficient activity of von Willebrand factor-cleaving protease in chronic relapsing thrombotic thrombocytopenic purpura. *Blood* 1997;89:3097-103.
- [42] Kokame K, Aoyama Y, Matsumoto M, Fujimura Y, Miyata T. Inherited and *de novo* mutations of ADAMTS13 in a patient with Upshaw-Schulman syndrome. *J Thromb Haemost* 2008;6:213-5.
- [43] Levy GG, Nichols WC, Lian EC, Foroud T, McClintick JN, McGee BM *et al.* Mutations in a member of the ADAMTS gene family cause thrombotic thrombocytopenic purpura. *Nature* 2001;413:488-94.
- [44] Tsai HM. Pathophysiology of thrombotic thrombocytopenic purpura. *Int J Hematol* 2010;91:1-19.
- [45] Bernardo A, Ball C, Nolasco L, Moake JF, Dong JF. Effects of inflammatory cytokines on the release and cleavage of the endothelial cell-derived ultralarge von Willebrand factor multimers under flow. *Blood* 2004;104:100-6.
- [46] Motto DG, Chauhan AK, Zhu G, Homeister J, Lamb CB, Desch KC *et al.* Shiga toxin triggers thrombotic thrombocytopenic purpura in genetically susceptible ADAMTS13-deficient mice. *J Clin Invest* 2005;115:2752-61.
- [47] Banno F, Kokame K, Okuda T, Honda S, Miyata S, Kato H *et al.* Complete deficiency in ADAMTS13 is prothrombotic, but it alone is not sufficient to cause thrombotic thrombocytopenic purpura. *Blood* 2006;107:3161-6.
- [48] Hara T, Kitano A, Kajiwara T, Kondo T, Sakai K, Hamasaki Y. Factor VIII concentrate-responsive thrombocytopenia, hemolytic anemia, and nephropathy. Evidence that factor VIII: von Willebrand factor is involved in its pathogenesis. *Am J Pediatr Hematol Oncol* 1986;8:324-8.
- [49] Veyradier A, Meyer D, Loirat C. Desmopressin, an unexpected link between nocturnal enuresis and inherited thrombotic thrombocytopenic purpura (Upshaw-Schulman syndrome). *J Thromb Haemost* 2006;4:700-1.
- [50] Fujimura Y, Matsumoto M, Kokame K, Isonishi A, Soejima K, Akiyama N *et al.* Pregnancy-induced thrombocytopenia and TTP, and the risk of fetal death, in Upshaw-Schulman syndrome: a series of 15 pregnancies in 9 genotyped patients. *Br J Haematol* 2009;144:742-54.
- [51] Uchida T, Wada H, Mizutani M, Iwashita M, Ishihara H, Shibano T *et al.* Identification of novel mutations in ADAMTS13 in an adult patient with congenital thrombotic thrombocytopenic purpura. *Blood* 2004;104:2081-3.
- [52] Schneppenheim R, Budde U, Oyen F, Angerhaus D, Aumann V, Drewke E *et al.* von Willebrand factor cleaving protease and

- ADAMTS13 mutations in childhood TTP. *Blood* 2003;101:1845-50.
- [53] Kokame K, Matsumoto M, Soejima K, Yagi H, Ishizashi H, Funato M *et al*. Mutations and common polymorphisms in ADAMTS13 gene responsible for von Willebrand factor-cleaving protease activity. *Proc Natl Acad Sci U S A* 2002;99:11902-7.
- [54] Matsumoto M, Kokame K, Soejima K, Miura M, Hayashi S, Fujii Y *et al*. Molecular characterization of ADAMTS13 gene mutations in Japanese patients with Upshaw-Schulman syndrome. *Blood* 2004;103:1305-10.
- [55] Shibagaki Y, Matsumoto M, Kokame K, Ohba S, Miyata T, Fujimura Y *et al*. Novel compound heterozygote mutations (H234Q/R1206X) of the ADAMTS13 gene in an adult patient with Upshaw-Schulman syndrome showing predominant episodes of repeated acute renal failure. *Nephrol Dial Transplant* 2006;21:1289-92.
- [56] Fujimura Y, Matsumoto M, Isonishi A *et al*. Natural history of Upshaw-Schulman syndrome based on ADAMTS13 gene analysis in Japan. *J Thromb Haemost* 2011;9(Suppl. 1): 283-301.
- [57] Camilleri RS, Cohen H, Mackie IJ, Scully M, Starke RD, Crawley JT *et al*. Prevalence of the ADAMTS-13 missense mutation R1060W in late onset adult thrombotic thrombocytopenic purpura. *J Thromb Haemost* 2008;6:331-8.
- [58] Ashida A, Nakamura H, Yoden A, Tamai H, Ishizashi H, Yagi H *et al*. Successful treatment of a young infant who developed high-titer inhibitors against VWF-cleaving protease (ADAMTS-13): Important discrimination from upshaw-Schulman syndrome. *Am J Hematol* 2002;71:318-22.
- [59] Sato A, Hoshi Y, Onuma M, Sato R, Tsunematsu Y, Isonishi A *et al*. A 9-month-old infant with acquired idiopathic thrombotic thrombocytopenic purpura caused by inhibitory IgG-autoantibody to ADAMTS13. *Pediatr Hematol Oncol* 2010;27:53-8.
- [60] Gitlow S, Goldmark C. Generalized capillary and arteriolar thrombosis. Report of two cases with a discussion of the literature. *Ann Intern Med* 1939;13:1046-67.
- [61] Brunner HJ, Freedman M, Silverman ED. Close relationship between systemic lupus erythematosus and thrombotic thrombocytopenic purpura in childhood. *Arthritis Rheum* 1999;42:2346-55.
- [62] Rock GA, Shumak KH, Buskard NA, Blanchette VS, Kelton JG, Nair RC *et al*. Comparison of plasma exchange with plasma infusion in the treatment of thrombotic thrombocytopenic purpura. Canadian Apheresis Study Group. *N Engl J Med* 1991;325:393-7.
- [63] McDonald V, Liesner R, Grainger J, Gattens M, Machin SJ, Scully M. Acquired, noncongenital thrombotic thrombocytopenic purpura in children and adolescents: clinical management and the use of ADAMTS 13 assays. *Blood Coagul Fibrinolysis* 2010;21:245-50.
- [64] Scully M, Yarranton H, Liesner R, Cavenagh J, Hunt B, Benjamin S *et al*. TTP registry: correlation with laboratory ADAMTS 13 analysis and clinical features. *Br J Haematol* 2008;142:819-26.
- [65] Fakhouri F, Vernant JP, Veyradier A, Wolf M, Kaplanski G, Binast R *et al*. Efficiency of curative and prophylactic treatment with rituximab in ADAMTS13-deficient thrombotic thrombocytopenic purpura: a study of 11 cases. *Blood* 2005;106:1932-7.
- [66] Scully M, Cohen H, Cavenagh J, Benjamin S, Starke R, Killick S *et al*. Remission in acute refractory and relapsing thrombotic thrombocytopenic purpura following rituximab is associated with a reduction in IgG antibodies to ADAMTS-13. *Br J Haematol* 2007;136:451-61.
- [67] Albaranki JH, Teo J, Alexander SI. Rituximab therapy in two children with autoimmune thrombotic thrombocytopenic purpura. *Pediatr Nephrol* 2009;24:1749-52.
- [68] Harambat J, Lamireau D, Delmas Y, Ryman A, Llanas B, Brissaud O. Successful treatment with rituximab for acute refractory thrombotic thrombocytopenic purpura related to acquired ADAMTS13 deficiency: a pediatric report and literature review. *Pediatr Crit Care Med* 2011;12:e90-3.
- [69] Jayabose S, Dunbar J, Nowicki TS, Tugal O, Ozkaynak MF, Sandoval C. Rituximab therapy to prevent relapse in chronic relapsing thrombotic thrombocytopenic purpura (TTP) in a child. *Pediatr Hematol Oncol* 2011;28(2):167-72.

