

表3 各臓器の絶対・相対重量及び最終体重 (ホルムアルデヒド:22時間/日×28日間)

肝

Exp. Group	Liver (g/100g)	Liver (g)	Body weight (g)
Control	5.304 ± 0.498	1.459 ± 0.150	27.5 ± 1.0
0.1 ppm	5.363 ± 0.642	1.458 ± 0.180	27.2 ± 0.5
0.3 ppm	5.478 ± 0.153	1.477 ± 0.085	27.0 ± 1.1
1.0 ppm	5.431 ± 0.161	1.438 ± 0.042	26.5 ± 0.7

腎

Exp. Group	Kidney (g/100g)	Kidney (g)	Body weight (g)
Control	1.087 ± 0.091	0.299 ± 0.030	27.5 ± 1.0
0.1 ppm	1.111 ± 0.065	0.302 ± 0.014	27.2 ± 0.5
0.3 ppm	1.109 ± 0.052	0.299 ± 0.017	27.0 ± 1.1
1.0 ppm	1.102 ± 0.063	0.292 ± 0.021	26.5 ± 0.7

脾

Exp. Group	Spleen (g/100g)	Spleen (g)	Body weight (g)
Control	0.266 ± 0.029	0.073 ± 0.009	27.5 ± 1.0
0.1 ppm	0.278 ± 0.032	0.076 ± 0.008	27.2 ± 0.5
0.3 ppm	0.239 ± 0.020	0.065 ± 0.008	27.0 ± 1.1
1.0 ppm	0.239 ± 0.023	0.063 ± 0.006	26.5 ± 0.7

心

Exp. Group	Heart (g/100g)	Heart (g)	Body weight (g)
Control	0.509 ± 0.045	0.140 ± 0.013	27.5 ± 1.0
0.1 ppm	0.504 ± 0.054	0.137 ± 0.013	27.2 ± 0.5
0.3 ppm	0.497 ± 0.015	0.134 ± 0.003	27.0 ± 1.1
1.0 ppm	0.490 ± 0.015	0.130 ± 0.005	26.5 ± 0.7

脳

Exp. Group	Brain (g/100g)	Brain(g)	Body weight (g)
Control	1.665 ± 0.093	0.457 ± 0.015	27.5 ± 1.0
0.1 ppm	1.693 ± 0.044	0.460 ± 0.006	27.2 ± 0.5
0.3 ppm	1.701 ± 0.075	0.458 ± 0.008	27.0 ± 1.1
1.0 ppm	1.682 ± 0.051	0.446 ± 0.016	26.5 ± 0.7

平均値±標準偏差 (* P<0.05, ** P<0.01)

表4 病理組織学的検索結果

臓器名	所見	対照群	0.1 ppm群	0.3 ppm群	1.0 ppm群
		検索数	6	6	6
脳		著変なし	著変なし	著変なし	著変なし
肝臓	炎症性細胞集簇	1+:2	1+:5, 2+:1	1+:6	1+:3
腎臓	好塩基性尿管		1+:1		1+:1
	乳頭部鈣質沈着			1+:1	
脾臓		著変なし	著変なし	著変なし	著変なし
胸腺		著変なし	著変なし	著変なし	著変なし
心臓		著変なし	著変なし	著変なし	著変なし

1+:軽度、2+:中等度

III. 研究成果の刊行に関する一覧表

書籍

著者氏名	論文タイトル名	書籍全体の 編集者名	書籍名	出版社名	出版地	出版年	ページ
Aisaki K and Kanno J	15 STANDARDIZATION OF GENE- EXPRESSION INFORMATION FOR THE SAFETY EVALUATION: ACTIVITIES IN JAPAN	Darrell R. Boverhof, B. Bhaskar Gollapudi	Applications of Toxicogeno mics in Safety Evaluation and Risk Assessment	John Wiley & Sons, Inc.	New York, USA	2011	323-329
Kanno J, Aisaki K, Igarashi K, Nakatsu N, Kodama Y, Sekita K, Takagi A and Kitajima S	Chapter11 Application of Percellome Toxicogenomics to Food Safety	Ingemar Pongratz, Linda Bergander	Hormone- Disruptive Chemical Contaminan ts in Food	Royal Society of Chemist ry	London, England	2011	184-198
菅野 純	Percellome トキシコ ゲノミクスの進捗		医学のあゆ み	医 歯 薬 出 版 株 式 会 社	Tokyo, Japan	2011	1125- 1126


雑誌

発表者氏名	論文タイトル名	発表誌名	巻名	ページ	出版年
Igarashi K, Kitajima S, Aisaki K, Tanemura K, Taquahashi Y, Moriyama N, Ikeno E, Matsuda N, Saga Y, Blumberg B, and Kanno J	Development of Humanized Steroid and Xenobiotic Receptor Mouse by homologous knock-in of the human Steroid and Xenobiotic Receptor Ligand Binding Domain sequence.	J Toxicol Sci	37	373- 380	2012

Hang NT, Lien LT, Kobayashi N, Shimbo T, Sakurada S, Thuong PH, Hong LT, Tam DB, Hijikata M, Matsushita I, Hung NV, Higuchi K, Harada N, Keicho N.	Analysis of factors lowering sensitivity of interferon-gamma release assay for tuberculosis.	PLoS One	6	e23806	2011
Tanaka T, Sakurada S, Kano K, Takahashi E, Yasuda K, Hirano H, Kaburagi Y, Kobayashi N, Hang N, Lien L, Matsushita I, Hijikata M, Uchida T, Keicho N.	Identification of tuberculosis-associated proteins in whole blood supernatant.	BMC Infect Dis	11	71	2011
Fujiki R, Hashiba W, Sekine H, Yokoyama A, Chikanishi T, Ito S, Imai Y, Kim J, He HH, Igarashi K, Kanno J, Ohtake F, Kitagawa H, Roeder RG, Brown M, Kato S.	GlcNAcylation of histone H2B facilitates its monoubiquitination.	Nature	480	557-560	2011
Fujimoto N, Kitamura S, Kanno J.	Androgen dependent transcription of a mouse prostatic protein gene, PSP94: Involvement of estrogen receptors.	J Steroid Biochem Mol Biol	127	301-306	2011
Nagano, K., Gotoh, K., Kasai, T., Aiso, S., Nishizawa, T., Ohnishi, M., Ikawa, N., Eitaki, Y., Yamada, K., Arito, H. and Fukushima, S.	Two- and 13-week Inhalation Toxicities of Indium-Tin Oxide and Indium Oxide in Rats.	J Occup Health	53	51-63	2011
Nagano, K., Nishizawa, T., Eitaki, Y., Ohnishi, M., Noguchi, T., Arito, H. and Fukushima, S.	Pulmonary Toxicity in Mice by 2- and 13-week Inhalation Exposures to Indium-tin Oxide and Indium Oxide Aerosols.	J Occup Health	53,	234-239	2011

IV. 研究成果の刊行物・別刷り

Applications of Toxicogenomics in Safety Evaluation and Risk Assessment



Edited by

Darrell R. Boverhof and B. Bhaskar Gollapudi

15

STANDARDIZATION OF GENE-EXPRESSION INFORMATION FOR THE SAFETY EVALUATION: ACTIVITIES IN JAPAN

Ken-ichi Aisaki and Jun Kanno

*Division of Cellular and Molecular Toxicology, Biological Safety Research
Center, National Institute of Health Sciences, Tokyo, Japan*

15.1	Introduction	324
15.2	Technical Information Essential for RNA Standardization	324
15.3	Information on Domestic and International Activities	324
15.3.1	Domestic Activities	325
15.3.1.1	<i>The Toxicogenomics Project of NIH</i>	325
15.3.1.2	<i>The Percellome Project</i>	325
15.3.1.3	<i>The Japan Microarray Consortium</i>	327
15.3.1.4	<i>The New Energy Development Organization</i>	327
15.3.1.5	<i>Other Domestic Initiatives</i>	327
15.3.2	International Activities	328
15.4	Closing Summary	328
	Acknowledgment	329
	References	329

Applications of Toxicogenomics in Safety Evaluation and Risk Assessment, First Edition.

Edited by Darrell R. Boverhof and B. Bhaskar Gollapudi.

© 2011 John Wiley & Sons, Inc. Published 2011 by John Wiley & Sons, Inc.

15.1 INTRODUCTION

Recent developments in cDNA microarray and quantitative PCR methodologies have demonstrated their applicabilities to medicine, food (including food additives and supplements), and industrial chemicals for the evaluation of their efficacies and safety. In the United States, the Food and Drug Administration (FDA) has started to accept voluntarily submitted gene-expression data for the drug approval processes. The European Medicines Agency (EMA) is also considering a guideline for such processes. In Japan, the regulatory agencies involved in the approval of medicines have not been officially taking action or developing guidelines for the use of genomics data. An exception is a Ministry of Health, Labor, and Welfare (MHLW) funded research group that has been preparing to educate corporations, government, and consumers on the urgency of this matter. The general thought among Japanese chemical corporations is that the safety evaluation of toxicogenomics is second only to the safety evaluation of medicine. Here we report the study results by the above-mentioned research group, which was organized by scientists who specialize in mRNA measurements.

15.2 TECHNICAL INFORMATION ESSENTIAL FOR RNA STANDARDIZATION

For the mRNA to be utilized for the safety evaluation of new chemical substances, the standardization techniques should meet the following conditions:

1. Be commonly applicable to quantitative PCR, any kind of microarray, and new techniques such as comprehensive high-throughput sequencing.
2. Be based on a universal scale independent of the time and place of measurement
3. Be highly reproducible
4. Be easy to conduct

Among various methodologies proposed, currently available, and most fitting to these conditions would be the Percellome method, which measures mRNA in a copy number per one cell basis (Kanno et al., 2006). Conditions 1 through 3 have been satisfied, and for condition 4, a manual on the use of robotics is available. In addition a database with over 100 chemicals for rodent liver has been prepared and will be publically accessible in the near future.

15.3 INFORMATION ON DOMESTIC AND INTERNATIONAL ACTIVITIES

There exists more than one activity domestically and internationally for mRNA standardization.

15.3.1 Domestic Activities

15.3.1.1 *The Toxicogenomics Project of NIHS*

The Toxicogenomics Project (TGP 1; 2002–2006, TGP 2; 2007–2012) is a joint research project of the government partnering fifty-fifty with the private sector. It is run by the National Institute of Health Sciences (NIHS) and an alliance of 17 Japanese pharmaceutical companies. The mission is to streamline the safety evaluation process for pharmaceuticals. Rat liver was chosen as a primary research target because the member companies already had their own reference data. The project adopted the Percellome method, which was championed by a group of scientists of the NIHS (led by Jun Kanno). Up to 150 chemicals were monitored for rat liver transcriptome responses after acute dosing (3, 6, 9, and 24 h after single oral administration) and repeated dosing (3, 7, 14, and 28 days of repeated oral dosage), with four dose levels. The project included a parallel examination of responses using in vitro assays of rat and human hepatocyte culture systems (Affymetrix GeneChip, Rat Expression 230A, Rat Genome 230 2.0 Array and Human Genome U133 Plus 2.0 Array). In 2005 the entire project was moved from the NIHS location to the newly established National Institute of Biomedical Innovation (NIBIO) in the Osaka region. Because of that point the member pharmaceutical companies had their own cumulative data (which were non-Percellome data), to save resources, it was decided to move away from the Percellome quality control and analysis, and instead adopt a common normalization such as the 75 percentile global normalization with superficial quality control. However, the sample processing protocol was not changed, including essential processes of the Percellome method, and retaining Percellome spike data. A repeating of arrays due to bad data quality was discovered by Percellome quality control and is no longer performed (later validation of the data revealed that most of the data were of adequate Percellome quality). For the in vitro studies, on the other hand, the TGP confirmed that the Percellome method is appropriate for normalizing in vitro data, and thus this methodology has remained. In 2007, the second five-year-project was started, again as a government-private sector joint research project by NIBIO, an alliance of 13 Japanese pharmaceutical companies and 2 groups of scientists, including Kanno's group, charged with the development of informatics using the database established in the first five-year project (<http://www.tgp.nibio.go.jp/>). Because the project is of a proprietary nature due to the membership contracts with the investing companies, the data will gradually become available to the public.

15.3.1.2 *The Percellome Project*

The Percellome Project to develop a mouse transcriptome began in 2003, one year after to the TGP project was initiated. This Percellome Project, which is supported by grants from the MHLW and other public institutions, has as its mission facilitating the chemical safety processes. The project is run by the Division of Cellular and Molecular Toxicology, Biological Safety Research Center of the NIHS (headed by Jun Kanno). The first study (designated as TTG1) ran from 2003 through

2005 and involved the development of a database of 90 chemicals on the mouse liver acute response (2, 4, 8, and 24 h after single oral exposure). From 2005 through 2007, an Inhalation Toxicogenomics (ITG) study was added to collect lung transcriptome data at very low exposure levels, near the levels associated with the “sick building” syndrome. From 2006 through 2008, a second stage of the Percellome Project (TTG2) was launched for repeated exposure studies and multi-organ responses studies. Studies of fetus (development) and brain behavior (designated as NTG) were also launched (Affymetrix GeneChip, Mouse Expression 430A, Mouse Genome 430 2.0 Array).

In the TTG1 studies we monitored the time- and dose-dependent gene expressions induced by a single oral administration of a test chemical in mouse liver at four time points (2, 4, 8, and 24 h) and at four dose levels (0, 1×, 3×, 10×), in triplicate (48 per chemical; Figure 15.1). Currently our database contains data for more than 100 chemicals. The data for each 100 chemicals is expressed as a 3-D graph (time × dose × mRNA in copies per cell) made of about 45,000 surfaces corresponding to the probe sets of the Affymetrix MOE430 2.0 GeneChip (see <http://www.toxicomics.nih.gov/db/>). Two kinds of unsupervised gene clustering methods are developed on the 3-D surface shape. (Aisaki et al., 2007, Matsumoto et al., 2005).




Figure 15.1 Structure of Percellome data showing the surface plot for a single transcript. In mouse liver transcriptome analysis, the mRNA copy numbers per cell of a probe set are plotted on a 4 × 4 (4 time points and 4 dosage levels) matrix to comprise a 3-D graph. One data set of a test chemical consists of about 45,000 layers of 3-D surfaces. Further analysis, such as unsupervised clustering and extraction of significant genes are performed on this 3-D surface data. (See insert for color representation of the figure.)

In the TTG2 study, three types of data were generated; the repeated dosing up to 14 days, multiorgan assessments (liver, kidney, lung, heart, testis, brain) after single dosing, and fetus/developmental toxicogenomics (whole embryo, embryonic stem cells, embryoid body). The ITG studies (lung and liver) repeated the dosages for up to 7 days. A series of in-house developed software programs were used and were mostly specified for the 3-D surface data sets to generate time-dependent triggered gene lists and common gene clusters among chemicals. A putative target gene of the knockout phenotypes turned out to completely agree with the estimated effects of the test chemical.

In 2009 a third Percellome Project was launched to comprehensively highlight the toxicity networks from the transcriptomic data collected by the preceding Percellome methods. The additional systems biology approach was incorporated for this purpose. The intention was to keep the transcriptomic data from being biased by the known toxicological information. Without phenotypic anchoring, we attempted to interpret the transcriptomic data by referring to our own transcriptomic database, and genome-based knowledge.

As of 2010 we have compiled a database of more than 100 chemicals, and this database will be available to the public on a “one-chemical one-report” basis. With each report we will include a comprehensive list of genes showing the biologically significant alteration. The home-page for the Percellome database should be available soon.

15.3.1.3 The Japan Microarray Consortium

The Japan Microarray Consortium (JMAC: <http://www.jmaq.org/>) has been operating since 2007 and consists of up to seventy of the biochip producing and related companies under the control of the Ministry of Economy, Trade and Industry. This consortium regularly holds working group meetings for the activity of industrial promotion and market creation along with an interest in standardization.

15.3.1.4 The New Energy Development Organization

The New Energy Development Organization (NEDO) had funded a carcinogenesis toxicogenomics project headed by Prof. T. Shirai of Nagoya City University. For up to 28 days, a total of 85 known hepato-carcinogenic, nonhepato-carcinogenic, or noncarcinogenic compounds were given to rats and livers were sampled at days 1, 3, 7, 14, and 28. The gene expression was monitored by in-house microarray (NEDO-ToxArray) and a GeneChip system (RGU34A, Affymetrix). (<http://www.nedo.go.jp/iinkai/kenkyuu/bunkakai/18h/jigo/49/1/5-2.pdf>, in Japanese).

15.3.1.5 Other Domestic Initiatives

In the domains of clinical medicine, drug development, and chemical/industrial manufacturing, there is broad agreement that comprehensive toxicogenomics data analysis is necessary for the more precise information on the toxicity of ingested chemicals. Yet each sector has been reluctant to make an effort toward the development of a standardization scheme. There are comments that suggest a

desire for concerted action by administrative authorities. The Pharmaceuticals and Medical Devices Agency (PMDA; <http://www.pmda.go.jp/english/index.html>), which approves new drug applications as commissioned by MHLW, has been holding informal meetings on pharmacogenomics with pharmaceutical companies. In July and October of 2007, PMDA joined the joint FDA/EMEA biomarker validation workshops as an observer. PMDA also asked the Japanese Society of Clinical Pharmacology and Therapeutics to prepare a report on the current status and challenges for pharmacogenomics in drug evaluation (<http://www.pmda.go.jp/topics/file/h191113kohyo.pdf>, in Japanese). Overall, pharmacogenomics is slowly but steadily moving toward some form of standardization.

15.3.2 International Activities

Major international activities include the MicroArray Quality Control Project (MAQC, US), the External RNA Controls Consortium (ERCC, US), and the European Medicines Agency (EMA). The MAQC and ERCC bring together scientists from microarray manufacturers, reagent manufacturers, pharmaceutical companies, academia, and government administrative offices. They hold periodic meetings and are rigorously incorporating information from related activities and research. In 2007 they held hearings on the Percellome Project, and the JMAC. MAQC-III has evolved into SEQC (Sequencing Quality Control), and so is charged with the quality control of next-generation high-speed sequencing platforms (<http://www.fda.gov/ScienceResearch/BioinformaticsTools/MicroarrayQualityControlProject/default.htm#MAQC-IIIalsoknownasSEQC>). At the E15 of the International Conference on Harmonisation of Technical Requirements for Registration of Pharmaceuticals for Human Use (ICH), a *Guidance for Industry; E15 Definitions for Genomic Biomarkers, Pharmacogenomics, Pharmacogenetics, Genomic Data and Sample Coding Categories* was published. This document has set the stage for future contributions to the standardization of terminology in pharmacogenomics. OECD and WHO are also jointly holding toxicogenomics workshops to discuss the future standardization and utilization of toxicogenomics in chemicals safety assessments (http://www.oecd.org/document/29/0,3343,en_2649_34377_34704669_1_1_1_1,00.html and <http://www.who.int/ipcs/methods/toxicogenomics/en/>).

15.4 CLOSING SUMMARY

The regulatory activities for the standardization of toxicogenomics are progressing rather slowly in Japan. The first area of interest would be the development of toxicopharmacogenomics data for drug approval agencies; it is expected that industrial chemicals would then follow.

The Percellome projects have made all reagents, protocols, and basic analytical tools for quality control available to their collaborators (kanno@nihs.go.jp). The Percellome database will be released to the public sometime in the near future.

ACKNOWLEDGMENT

The authors thank all the members of the Division of Cellular and Molecular Toxicology, NIHS for the support of Percellome Projects and especially Nae Matsuda, Tomoko Ando, Noriko Moriyama, Yuko Kondo, Maki Abe, Koichi Morita, Hisako Aihara and Chiyuri Aoyagi for technical support. This study was supported in part by MHLW Health Sciences Research Grants H19-Iyaku-Ippan-001, H18-Kagaku-Ippan-001, H15-Kagaku-002, and H14-Toxico-001.

REFERENCES

- Aisaki, K., Aizawa, S., Fujii, H., Kanno, J., and Kanno, H. (2007). Glycolytic inhibition by mutation of pyruvate kinase gene increases oxidative stress and causes apoptosis of a pyruvate kinase deficient cell line. *Exp Hematol* **35**, 1190–1200.
- Kanno, J., Aisaki, K., Igarashi, K., Nakatsu, N., Ono, A., Kodama, Y., and Nagao, T. (2006). “Per cell” normalization method for mRNA measurement by quantitative PCR and microarrays. *BMC Genomics* **7**, 64.
- Matsumoto, S., Aisaki, K., and Kanno, J. (2005). Mass distributed clustering: a new algorithm for repeated measurements in gene expression data. *Genome Info* **16**, 183–194.

CHAPTER 11

Application of Percellome Toxicogenomics to Food Safety

J. KANNO,^a K. AISAKI,^a K. IGARASHI,^a N. NAKATSU,^{a,b}
Y. KODAMA,^a K. SEKITA,^a A. TAKAGI^a AND
S. KITAJIMA^a

^aNational Institute of Health Sciences, Tokyo 158-8501, Japan; ^bNational Institute of Biomedical Innovation, Osaka 567-0085, Japan

11.1 Introduction

The approaches employed to assess chemical toxicity are seldom applied to food safety assessment for a number of reasons. In the first place, routine toxicity assessment involves administration of a pure compound to experimental animals at different doses up to the maximal tolerated dose (MTD) to determine the no observed (adverse) effect level [NO(A)EL], while food is a complex mixture of chemicals. Secondly, the sensitivity of routine toxicity tests is inadequate for foods. The maximal amount of the test substance applied is limited by guidelines to 5% of the diet or, in the case of oral gavage, $10 \text{ mL}\cdot\text{kg}^{-1}$ (approximately $10 \text{ g}\cdot\text{kg}^{-1}$ for liquids), in order not to jeopardize reliability (especially with respect to chronic toxicity and carcinogenesis) by inducing malnutrition. For example, low body weight alone can result in tumors at low incidences.¹⁻³ When an item of food is found to be non-toxic at the dose of $10 \text{ g}\cdot\text{kg}^{-1}$, this value is taken as NOEL and the tolerable daily intake (TDI) calculated, using a safety factor of 1/100, to be $100 \text{ mg}\cdot\text{kg}^{-1}$. Thus the grams or even tens of grams per kg body weight that we consume daily are not allowed in routine toxicity testing and, moreover, according to such standard approaches, if a food showed any toxicity in routine testing, such food would be highly toxic.

Even though flavorings are added to food and other consumed products at relatively low concentrations, these substances are often not subject to toxicity testing at the MTD from other reasons. One practical consideration is that the odor of many flavorings at MTD is too strong to be tolerated by the test animal or the staff that care for them. Moreover, in some cases, the amount of a flavoring required to conduct an MTD test is more than its annual production and/or prohibitively costly. At the practical dose levels, flavoring does not exert any detectable histopathological effect on experimental animals. However, such doses do stimulate, at least, olfactory (and perhaps other) receptors. Thus, more sensitive approaches to assessing the toxicity of food and flavors are highly desirable. One approach is to use particularly sensitive animal models, such as transgenic mice to evaluate certain biological phenomena, including carcinogenic potential. Another strategy is to detect early molecular changes that occur prior to morphological alterations and/or other symptoms. Toxicogenomics is one example of the latter.

Our Percellome Toxicogenomics Project is designed to identify dynamic and extensive networks of genes whose time- and dose-dependent patterns of expression in response to a chemical, or a mixture of chemicals such as food, allows its toxic effects to be predicted. Here, we describe this project briefly, employing a flavoring compound, estragole, as a peroxisome proliferator-activated receptor alpha (PPAR-alpha) agonist whose effects are similar to those of clofibrate. Among the more than 100 chemicals tested, 45 are food related (Table 11.1), including food additives, components of food, functional health foods, agrochemicals, and other food contaminants (*e.g.* that might leach out of packaging materials).

11.2 Materials and Methods

Forty-eight male C57BL/6 Cr Slc or C57BL/6J mice (SLC, Hamamatsu, Charles River Japan, Atsugi, Japan) were maintained for two weeks in an enclosed environment (diet: CRF-1, Oriental Yeast Co.) with a 12-h photoperiod (starting at 8:00 a.m.) and then, at 12 weeks of age, a single oral administration of the test chemical at 10:00~10:20 a.m. and at the doses indicated in Table 11.1. At 2, 4, 8, and 24 hours later, the livers of three animals at each dose were collected. To minimize circadian oscillations in the transcriptome, 12 animals were processed at each time-point at a rate of 2.5 min from initiation ether anesthesia to immersion of the liver and other organ samples into RNAlater solution (Ambion, TX, USA), thereby keeping the total elapsed to within 30 min. Prior to analysis, the blocks of liver tissue were soaked in RNAlater overnight at 4 °C. This solution was then replaced with RLT buffer (Qiagen, Germany), the tissue homogenized, and a small aliquot of each homogenate treated with DNase-free RNase A, followed by proteinase K; the DNA concentration was then determined with PicoGreen fluorescent dye (Molecular Probes, USA). Next, on the basis of its DNA content, each sample was spiked with the same relative amount of GSC cocktail (containing

Table 11.1 Food-related chemicals in Percellome Database.

<i>Food-related Chemicals (including agrochemicals)</i>	<i>CAS No.</i>	<i>Dose (mg/kg)</i>	<i>Vehicle</i>	<i>Percellome study No. TTG</i>
Acephate	30560-19-1	0, 0.7, 2, 7, 20, 70 mg/kg	MC 0.5%	67, 109
Alpha lipoic acid	1077-28-7	0, 10, 30, 100 mg/kg	MC 0.5%	128
Aluminum ammonium sulfate	7784-26-1	0, 30, 100, 300 mg/kg	MC 0.5%(WAKO)	155
Aluminum lactate	18917-91-4	0, 30, 100, 300 mg/kg	MC 0.5%(WAKO)	163
Aluminum sulfate	10043-01-3	0, 100, 300, 1000 mg/kg	MC 0.5%(WAKO)	178
Bisphenol A	1980-5-7	0, 30, 100, 300 mg/kg	DMSO 0.1% + MC 0.5%	47
Caffeine	1958-8-2	0, 3, 10, 30 mg/kg	MC 0.5%	59
Carbaryl	63-25-2	0, 2, 7, 10, 20, 30, 100 mg/kg	MC 0.5%	68, 166
Chlorpyrifos	2921-88-2	0, 3, 10, 30 mg/kg	MC 0.5%(WAKO)	165
Citric acid-calcium salt	813-94-5	0, 100, 300, 1000 mg/kg	MC 0.5%	39
Coenzyme Q10	303-98-0	0, 30, 100, 300 mg/kg	corn oil	86
Curcumin	458-37-7	0, 70, 200, 700 mg/kg	corn oil	132
DEHP (di(2-ethylhexyl)phthalate)	117-81-7	0, 200, 700, 2000 mg/kg (<i>cf.</i> MEHP)	corn oil	98
Dexamethasone	1950-2-2	0, 10, 30, 100 ug/kg	DMSO 0.1% + MC 0.5%	62
Estragole	140-67-0	0, 10, 30, 100 mg/kg	corn oil	147
Ethanol	64-17-5	0, 150, 500, 1500 mg/kg (<i>d</i> = 0.79)	MC 0.5%	78
Food Red No.104	18472-87-2	0, 10, 30, 100 mg/kg	MC 0.5%(WAKO)	156
Forskolin	66575-29-9	0, 1, 3, 10 mg/kg	MC 0.5%	60
Genistein	446-72-0	0, 10, 30, 100 mg/kg	DMSO 0.1% + MC 0.5%	48
Glycyrrhizin2K	1405-86-3	0, 3, 10, 30 mg/kg	DMSO 0.1% + MC 0.5%	110
Hydroxycitric Acid	27750-10-3	0, 100, 300, 1000 mg/kg	MC 0.5%	38

Indigo	482-89-3	0, 30, 100, 300 mg/kg	corn oil	57
Maltol	118-71-8	0, 10, 30, 100 mg/kg	corn oil	133
MEHP (mono(2-ethylhexyl)phthalate)	4376-20-9	0, 70, 200, 700 mg/kg (<i>cf.</i> MEHP)	corn oil	104
Methanol	67-56-1	0, 30, 100, 300 mg/kg (<i>cf.</i> Formalin, EtOH)	MC 0.5%	79
Methoprene	40596-69-8	0, 0.3, 1, 3 mg/kg	DMSO 0.1% + corn oil	66
Methyl dihydro jasmonate	24851-98-7	0, 10, 30, 100 mg/kg	corn oil	135
Nerolidol	cis: [3790-78-1], trans:[40716-66-3]	0, 5.9, 17.7, 59.1 mg/kg	corn oil	134
Paraquat dichloride	1910-42-5	0, 1, 3, 10 mg/kg	MC 0.5%	61
Pentachlorophenol	87-86-5	0, 10, 30, 100 mg/kg	MC 0.5%	16
Permethrin	52645-53-1	0, 10, 30, 100 mg/kg	corn oil	97
Phytol	7541-49-3	0, 10, 30, 100 mg/kg	corn oil	136
Pyriproxyfen	95737-68-1	0, 0.3, 1, 3 mg/kg	DMSO 0.1% + corn oil	87
Sodium Dehydroacetate	4418-26-2	0, 30, 100, 300 mg/kg	MC 0.5%(WAKO)	154
Tebufenozide	112410-23-8	0, 0.3, 1, 3 mg/kg	DMSO 0.1% + corn oil	88
Tributyltin	56573-85-4	0, 30,100, 300 ug/kg	DMSO 0.5% + MC 0.5%	85
Tributyltin chloride	1461-22-9	0, 3,10, 30 ug/kg	DMSO 0.1% + MC 0.5%	49
Verbenone	18308-32-5	0, 10, 30, 100 mg/kg	corn oil	148
Warfarin (sodium salt)	81-81-2	0, 10, 30, 100 mg/kg	MC 0.5%	81
FD&C Blue No.1 (brilliant blue)	3844-45-9	0, 30, 100, 300 mg/kg	MC 0.5%(WAKO)	152
FD&C Blue No.2 (Indigo carmine)	860-22-0	0, 30, 100, 300 mg/kg	MC 0.5%(WAKO)	139
Food Red No.102	2611-82-7	0, 100, 300, 1000 mg/kg	MC 0.5%(WAKO)	177
FD&C Red No.40	25956-17-6	0, 30, 100, 300 mg/kg	MC 0.5%(WAKO)	140
FD&C Green No.8 (pyranine conc)	6358-69-6	0, 30, 100, 300 mg/kg	MC 0.5%(WAKO)	176
FD&C Yellow No.5 (Food Yellow No.4 Japan)	1934-21-0	0, 100, 300, 1000 mg/kg	MC 0.5%(WAKO)	161

MC: methylcellulose

five *Bacillus subtilis* RNA sequences corresponding to the Affymetrix GeneChip probe sets AFFX-ThrX-3_at, AFFX-LysX-3_at, AFFX-PheX-3_at, AFFX-DapX-3_at, and AFFX-TrpX-3_at), and subsequently processed in accordance with the Affymetrix Standard protocol for application to Mouse 430 2.0 GeneChips. After conversion of the raw read-outs to Percellome data,⁴ the values obtained (with four doses and five time-points, using 2-h vehicle value as 0-h) were plotted on the three-dimensional graphs (*cf.* <http://toxicomics.nih.gov/db/>) (Figure 11.1).

The highest dose that, 24 hours after a single oral gavage, did not influence body weight or cause macroscopic changes in internal organs, microscopic alterations in liver observed with H&E staining, and/or abnormal behavior was employed as the maximal dose. In general, the maximal doses in the Percellome studies are 10- to 100-fold lower than the reported MTDs.

The Percellome data set was first screened with an RSort program,⁵ which identifies the number of peaks on a surface, distinguishes between the surfaces for 45,101 probe sets simply on the basis of their shapes, includes which of the peaks is significantly different ($p < 0.05$) from the corresponding control (vehicle) values, as evaluated by Student's *t*-test, and excludes probe sets with less than one copy per liver cell. Thereafter, taking into consideration the *t*-test values and ± 1 standard deviation, the surface shape was checked manually by




Figure 11.1 A three-dimensional representation of Percellome data. The dose- and time-dependent alterations in the expression of a probe set of genes is expressed as the absolute number of mRNA molecule per cell on the vertical axis. The mean of data from three different animals at each time and dose is represented, along with the SD (colored more lightly). Asterisks indicate $p < 0.05$ from concurrent control by student's *t*-test; small crosses represents data for each animal.




Figure 11.2 The chemical structure of estragole.

S.K. and/or J.K. Another original program referred to as Percellome Explorer⁵ was utilized to compare pairwise the RSort results of all evaluations in the Percellome project with lists of shared probe sets.

The flavor compound estragole (1-allyl-4-methoxybenzene; CAS no. 140-67-0) (Figure 11.2) is a major component of tarragon, as well as of oils obtained from basil and other plants. We found that pure estragole (98%) from Sigma-Aldrich (catalog no. A29208-25G, lot no. 05202AH) at a maximal dose of 100 mg kg⁻¹ caused no observable effects after 24 hours, so the Percellome investigation was conducted at doses of 0, 10, 30, and 100 mg kg⁻¹ dissolved in corn oil (Sigma-Aldrich, C8267) and delivered by gavage.

11.3 Results

First, the Percellome data obtained were filtered automatically with the RSort program to identify surfaces with less than four peaks in the 4 x 4 matrix, a *p*-value of <0.05 at any time-point when comparing the control value to the corresponding maximal or middle dose, and a maximal peak of more than three copies of mRNA per cell. In this manner, 1214 up-regulated and 244 down-regulated sets were automatically selected out of the 45,101 probe sets in the mouse GeneChip, and subsequently 167 of those were selected manually as biologically feasible. The surfaces of several of the probe sets selected are depicted in Figure 11.3. Among those up-regulated, Ingenuity Pathway Analysis (IPA) highlighted a certain number genes encoding proteins associated with the PPAR-alpha. Of the metabolic enzymes up-regulated, Cyp4a10, Cyp4a14, and Cyp4a31, the last has been reported to be regulated by PPAR-alpha. Expression of the up-regulated transporters Slc16a5, Slc25a20, and Slc27a1 is also considered to be regulated by PPAR-alpha. A limited number of genes related to oxidative stress, apoptosis, and cell proliferation were induced (Table 11.2). The genes down-regulated could not be assigned to any known pathway by IPA or by searching the literature. Therefore, we conclude that administration of a single gavage of estragole up-regulates the expression of genes controlled by PPAR-alpha.

11.4 Discussion

Among others, PPAR-alpha regulates a set of genes whose products participate in the β -oxidation of fatty acids, as well as, in rodents at least, peroxisome




Figure 11.3 Representative three-dimensional surfaces of probe sets induced by estragole; Acot3, Acs11, Crat, Ehhadh, Pex11a, Cyp4a14, Slc16a5, Slc25a20, and Slc27a1 are considered to be regulated by PPAR-alpha; the findings with respect to genes related to oxidative stress, apoptosis, and cell proliferation are also shown (*cf.* Table 11.2).