

Table 3
Point mutations of the five reference Beijing strains detected by comparative genomic sequencing (CGS).

Strain	A05N056	ID381	4558	4994	4991/M
Sublineage	G1/2	G3	G4	G5/6	G7
No. of point mutations*					
Total	1071 (815)	1077 (843)	1127 (881)	1119 (900)	1148 (1009)
Common to at least two strains	851 (813)	886 (843)	928 (878)	950 (896)	949 (901)
Specific to each strain	220 (2)	191 (0)	199 (3)	169 (4)	199 (108)

* Numbers in parentheses denote the number of point mutations common to K-backbone (Niemann et al., 2009).

genotypes from G4; 24 of 106 genotypes from G5/6; 15 of 54 genotypes from G7.

They were selected from a total population of isolates of pulmonary tuberculosis patients in Kobe City during 2002 and 2007, which had already been reported (Iwamoto et al., 2009). A minimum spanning tree (MST) based on their VNTR types (Fig. S1) was constructed using software (Bionumerics ver. 4.2; Applied Maths, Sint-Martens-Latem, Belgium) according to descriptions in previous reports (Wada and Iwamoto, 2009; Wada et al., 2009a).

3. Results

3.1. Comparative genomics of referential strains

To ascertain the genetic diversity of the five sublineages of Beijing family, CGS analysis was conducted for the five reference strains. In total, ROIs common to at least two strains (1133 positions) and specific to each strain (A05N056: 273 positions, ID381: 263 positions, 4558: 266 positions, 4994: 329 positions, and 4991/M: 220 positions) were verified using resequencing arrays to determine their exact substitutions. As results of two steps of hybridization, approximately 1100 positions (76–85% of ROIs) were determined as point mutations in each strain (Table 3).

To evaluate the reliability of detection, these positions were compared with existing mutation data of Beijing strains based on a next-generation sequencer (NGS) (Niemann et al., 2009). Two Beijing strains isolated in Kazakhstan, K-1 and K-2, whose genotypes were identical, were analysed in their study. Their common SNPs (K-backbone) comprised 1209 point mutations. Our microarray data revealed that the number of common mutations (each targeted strain and K-backbone) were correspondent to the phylogenetic divergence (G1/2–G7, 815 [67.4%]–1009 [83.5%] positions). Approximately 200 mutations were specific to respective strains (Table 3). Most mutations that were specific to four of five strains (except for 4991/M) were not identical to the K-backbone. They were estimated as peculiar mutations that occurred after their phylogenetic divergence in the evolutionary process to the modern Beijing subfamily. However, 108 of 199 point mutations specific to 4991/M were common to the K-backbone. All positions designated as SNPs by the CGS are listed in Tables S1 and S2.

3.2. Subdivision of clinical strains population

In CGS analysis, 129 positions were estimated as intergenic SNPs that were specific to each strain. When they were verified using a conventional sequencer, 123 (96.1%) positions were amplified by PCR, and 111 positions (including two deletion mutations) were confirmed as mutations that were exclusive to each strain (Table S3). These specific mutations were used to characterise 103 clinical strains belonging to the sublineages studied to ascertain the inner phylogenetic divergence of each sublineage. Results show that all strains possessed common mutations (G1/2, 6 of 20 positions; G3, 8 of 25; G4, 6 of 22; G5/6 1 of 18; and G7, 12 of 26) in the respective sublineages, which means that these

Table 4
Phylogenetic subdivision of *M. tuberculosis* Beijing family in Japan based on intergenic point mutations identified in this study.

Sublineage	Subdivision [†]	No. of subdivided strains*	No. of intergenic mutations of respective subdivisions
Total	–	–	–
G1/2	G1/2-0 [†]	1	20/20
	G1/2-1	4	19/20
	G1/2-2	1	14/20
	G1/2-3	2	13/20
	G1/2-4	2	7/20
	G1/2-5	6	6/20
G3	G3-1	2	10/25
	G3-2	21	9/25
	G3-3	1	8/25
G4	G4-1	4	17/22
	G4-2	19	15/22
	G4-3	1	6/22
G5/6	G5/6-1	3	11/18
	G5/6-2	3	10/18
	G5/6-3	1	3/18
	G5/6-4	17	1/18
G7	G7-1	1	25/26
	G7-2	9	14/26
	G7-3	2	13/26
	G7-4	3	12/26

* Genotypes of strains were defined by 15 loci of variable number of tandem repeats (Supply et al., 2006).

[†] Correspondent to divergence position illustrated in Fig. 1.

[‡] Identical to the reference strain A05N056.

sublineages have been monophyletic with respective bottlenecks in their evolution (Table 4, Fig. 1). The phylogenetic divergences and VNTR types of the strains used for this setting are listed in Table S4.

Each sublineage was subdivided into 3–6 sub-sublineages (Fig. 1). They were regarded as having been diverted at respective phylogenetic points. In our setting, four sublineages (except for G1/2) were observed to include the phylogenetic divergence of major components of the analysed strains (G3-2, 21 of 24 strains; G4-2, 19 of 24; G5/6-4, 17 of 24; and G7-2, 9 of 15).

4. Discussion

The ancient (atypical) Beijing subfamily has predominated in Japan despite dissemination of the modern Beijing subfamily throughout the world. The genetic diversity of Beijing lineage in Japan has been regarded as broader than in other areas. That outstanding difference encouraged us to scrutinise the genetic characteristics and varieties of Beijing strains in Japan, which might yield various clues suggesting not only the evolutionary process of the lineage but also the fine adaptation of the pathogen to endemic areas.

In this study, we verified that each of the five sublineages of Beijing family in Japan was monophyletic with a bottleneck

(Research on Emerging and Re-emerging Infectious Diseases, Health Sciences Research Grants), and the Kurozumi Medical Foundation (Epidemiology).

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.meegid.2012.04.029>.

References

- Abadia, E., Zhang, J., dos Vultos, T., Ritacco, V., Kremer, K., Aktas, E., Matsumoto, T., Refregier, G., van Soolingen, D., Gicquel, B., Sola, C., 2010. Resolving lineage assignment on *Mycobacterium tuberculosis* clinical isolates classified by spoligotyping with a new high-throughput 3R SNPs based method. *Infect. Genet. Evol.* 10, 1066–1074.
- Albert, T.J., Dailidiene, D., Dailide, G., Norton, J.E., Kalia, A., Richmond, T.A., Molla, M., Singh, J., Green, R.D., Berg, D.E., 2005. Mutation discovery in bacterial genomes: metronidazole resistance in *Helicobacter pylori*. *Nat. Methods* 2, 951–953.
- Cole, S.T., Brosch, R., Parkhill, J., Garnier, T., Churcher, C., Harris, D., Gordon, S.V., Eiglmeier, K., Gas, S., Barry IIIrd, C.E., Teakaia, F., Badcock, K., Basham, D., Brown, D., Chillingworth, T., Connor, R., Davies, R., Devlin, K., Feltwell, T., Gentles, S., Hamlin, N., Holroyd, S., Hornsby, T., Jagels, K., Krogh, A., McLean, J., Moule, S., Murphy, L., Oliver, K., Osborne, J., Quail, M.A., Rajandream, M.A., Rogers, J., Rutter, S., Seeger, K., Skelton, J., Squares, R., Squares, S., Sulston, J.E., Taylor, K., Whitehead, S., Barrell, B.G., 1998. Deciphering the biology of *Mycobacterium tuberculosis* from the complete genome sequence. *Nature* 393, 537–544.
- Comas, I., Homolka, S., Niemann, S., Gagneux, S., 2009. Genotyping of genetically monomorphic bacteria: DNA sequencing in *Mycobacterium tuberculosis* highlights the limitations of current methodologies. *PLoS One* 4, e7815.
- Faksri, K., Drobniowski, F., Nikolayevskiy, V., Brown, T., Prammananan, T., Palittapongpim, P., Prayoonwiwat, N., Chairasert, A., 2011. Genetic diversity of the *Mycobacterium tuberculosis* Beijing family based on IS6110, SNP, LSP and VNTR profiles from Thailand. *Infect. Genet. Evol.* 11, 1142–1149.
- Filliol, I., Motiwala, A.S., Cavatore, M., Qi, W., Hazbon, M.H., Bobadilla del Valle, M., Fyfe, J., Garcia-Garcia, L., Rastogi, N., Sola, C., Zozio, T., Guerrero, M.I., Leon, C.I., Crabtree, J., Angiuoli, S., Eisenach, K.D., Durmaz, R., Joloba, M.L., Rendon, A., Sifuentes-Osorio, J., Ponce de Leon, A., Cave, M.D., Fleischmann, R., Whittam, T.S., Alland, D., 2006. Global phylogeny of *Mycobacterium tuberculosis* based on single nucleotide polymorphism (SNP) analysis: insights into tuberculosis evolution, phylogenetic accuracy of other DNA fingerprinting systems, and recommendations for a minimal standard SNP set. *J. Bacteriol.* 188, 759–772.
- Gagneux, S., DeRiemer, K., Van, T., Kato-Maeda, M., de Jong, B.C., Narayanan, S., Nicol, M., Niemann, S., Kremer, K., Gutierrez, M.C., Hilty, M., Hopewell, P.C., Small, P.M., 2006. Variable host-pathogen compatibility in *Mycobacterium tuberculosis*. *Proc. Natl. Acad. Sci. USA* 103, 2869–2873.
- Gagneux, S., Small, P.M., 2007. Global phylogeography of *Mycobacterium tuberculosis* and implications for tuberculosis product development. *Lancet Infect. Dis.* 7, 328–337.
- Glynn, J.R., Whiteley, J., Bifani, P.J., Kremer, K., van Soolingen, D., 2002. Worldwide occurrence of Beijing/W strains of *Mycobacterium tuberculosis*: a systematic review. *Emerg. Infect. Dis.* 8, 843–849.
- Hanekom, M., van der Spuy, G.D., Streicher, E., Ndabambi, S.L., McEvoy, C.R., Kidd, M., Beyers, N., Victor, T.C., van Helden, P.D., Warren, R.M., 2007. A recently evolved sublineage of the *Mycobacterium tuberculosis* Beijing strain family is associated with an increased ability to spread and cause disease. *J. Clin. Microbiol.* 45, 1483–1490.
- Herring, C.D., Palsson, B.O., 2007. An evaluation of Comparative Genome Sequencing (CGS) by comparing two previously-sequenced bacterial genomes. *BMC Genomics* 8, 274.
- Iwamoto, T., Fujiyama, R., Yoshida, S., Wada, T., Shirai, C., Kawakami, Y., 2009. Population structure dynamics of *Mycobacterium tuberculosis* Beijing strains during past decades in Japan. *J. Clin. Microbiol.* 47, 3340–3343.
- Iwamoto, T., Yoshida, S., Suzuki, K., Wada, T., 2008. Population structure analysis of the *Mycobacterium tuberculosis* Beijing family indicates an association between certain sublineages and multidrug resistance. *Antimicrob. Agents Chemother.* 52, 3805–3809.
- Luo, T., Yang, C., Gagneux, S., Gicquel, B., Jian, M., Gao, Q., 2012. Combination of single nucleotide polymorphism and variable number tandem repeats for genotyping homogenous population of *Mycobacterium tuberculosis* Beijing strains in China. *J. Clin. Microbiol.* 50, 633–639.
- Maeda, S., Wada, T., Iwamoto, T., Murase, Y., Mitarai, S., Sugawara, I., Kato, S., 2010. Beijing family *Mycobacterium tuberculosis* isolated from throughout Japan: phylogeny and genetic features. *Int. J. Tuberc. Lung. Dis.* 14, 1201–1204.
- Millet, J., Miyagi-Shiohira, C., Yamane, N., Mokrousov, I., Rastogi, N., 2012. High-resolution MIRU-VNTRs typing reveals the unique nature of *Mycobacterium tuberculosis* Beijing genotype in Okinawa. *Japan Infect. Genet. Evol.* 12, 637–641.
- Mokrousov, I., Ly, H.M., Otten, T., Lan, N.N., Vyshnevskiy, B., Hoffner, S., Narvskaya, O., 2005. Origin and primary dispersal of the *Mycobacterium tuberculosis* Beijing genotype: clues from human phylogeography. *Genome Res.* 15, 1357–1364.
- Mokrousov, I., Narvskaya, O., Otten, T., Vyazovaya, A., Limeschenko, E., Steklova, L., Vyshnevskiy, B., 2002. Phylogenetic reconstruction within *Mycobacterium tuberculosis* Beijing genotype in northwestern Russia. *Res. Microbiol.* 153, 629–637.
- Niemann, S., Koser, C.U., Gagneux, S., Plinke, C., Homolka, S., Bignell, H., Carter, R.J., Cheetham, R.K., Cox, A., Gormley, N.A., Kokko-Gonzales, P., Murray, L.J., Rigatti, R., Smith, V.P., Arends, F.P., Cox, H.S., Smith, G., Archer, J.A., 2009. Genomic diversity among drug sensitive and multidrug resistant isolates of *Mycobacterium tuberculosis* with identical DNA fingerprints. *PLoS One* 4, e7407.
- Okamoto, S., Tamaru, A., Nakajima, C., Nishimura, K., Tanaka, Y., Tokuyama, S., Suzuki, Y., Ochi, K., 2007. Loss of a conserved 7-methylguanosine modification in 16S rRNA confers low-level streptomycin resistance in bacteria. *Mol. Microbiol.* 63, 1096–1106.
- Parish, T., Stoker, N.G., 1998. *Mycobacteria* protocols. In: Walker, J.M. (Ed.), *Methods in Molecular Biology*. Humana Press, pp. 31–44.
- Schurch, A.C., Kremer, K., Hendriks, A.C., Freyee, B., McEvoy, C.R., van Crevel, R., Boeree, M.J., van Helden, P., Warren, R.M., Siezen, R.J., van Soolingen, D., 2011. SNP/RD typing of *Mycobacterium tuberculosis* Beijing strains reveals local and worldwide disseminated clonal complexes. *PLoS One* 6, e28365.
- Supply, P., Allix, C., Lesjean, S., Cardoso-Oelemann, M., Rusch-Gerdes, S., Willery, E., Savine, E., de Haas, P., van Deutekom, H., Roring, S., Bifani, P., Kurepina, N., Kreiswirth, B., Sola, C., Rastogi, N., Vatin, V., Gutierrez, M.C., Fauville, M., Niemann, S., Skuce, R., Kremer, K., Locht, C., van Soolingen, D., 2006. Proposal for standardization of optimized mycobacterial interspersed repetitive unit-variable-number tandem repeat typing of *Mycobacterium tuberculosis*. *J. Clin. Microbiol.* 44, 4498–4510.
- Tsolaki, A.G., Gagneux, S., Pym, A.S., Goguet de la Salmoniere, Y.O., Kreiswirth, B.N., Van Soolingen, D., Small, P.M., 2005. Genomic deletions classify the Beijing/W strains as a distinct genetic lineage of *Mycobacterium tuberculosis*. *J. Clin. Microbiol.* 43, 3185–3191.
- van Soolingen, D., Qian, L., de Haas, P.E., Douglas, J.T., Traore, H., Portaels, F., Qing, H.Z., Enkhsaikan, D., Nymadawa, P., van Embden, J.D., 1995. Predominance of a single genotype of *Mycobacterium tuberculosis* in countries of East Asia. *J. Clin. Microbiol.* 33, 3234–3238.
- Wada, T., Fujihara, S., Shimouchi, A., Harada, M., Ogura, H., Matsumoto, S., Hase, A., 2009a. High transmissibility of the modern Beijing *Mycobacterium tuberculosis* in homeless patients of Japan. *Tuberculosis (Edinb)* 89, 252–255.
- Wada, T., Iwamoto, T., 2009. Allelic diversity of variable number of tandem repeats provides phylogenetic clues regarding the *Mycobacterium tuberculosis* Beijing family. *Infect. Genet. Evol.* 9, 921–926.
- Wada, T., Iwamoto, T., Maeda, S., 2009b. Genetic diversity of the *Mycobacterium tuberculosis* Beijing family in East Asia revealed through refined population structure analysis. *FEMS Microbiol. Lett.* 291, 35–43.
- Yokoyama, E., Hachisu, Y., Hashimoto, R., Kishida, K., 2012. Population genetic analysis of *Mycobacterium tuberculosis* Beijing subgroup strains. *Infect. Genet. Evol.* 12, 630–636.
- Zhang, Y., Chen, C., Liu, J., Deng, H., Pan, A., Zhang, L., Zhao, X., Huang, M., Lu, B., Dong, H., Du, P., Chen, W., Wan, K., 2011. Complete genome sequences of *Mycobacterium tuberculosis* strains CCDC5079 and CCDC5080, which belong to the Beijing family. *J. Bacteriol.* 193, 5591–5592.

Association of *IFNGR2* gene polymorphisms with pulmonary tuberculosis among the Vietnamese

Minako Hijikata · Junko Shojima · Ikumi Matsushita · Katsushi Tokunaga · Jun Ohashi · Nguyen T. L. Hang · Toru Horie · Shinsaku Sakurada · Nguyen P. Hoang · Pham H. Thuong · Luu T. Lien · Naoto Keicho

Received: 16 August 2011 / Accepted: 24 October 2011 / Published online: 6 November 2011
© The Author(s) 2011. This article is published with open access at Springerlink.com

Abstract Interferon- γ (IFN- γ) is a key molecule of T helper 1 (Th1)-immune response against tuberculosis (TB), and rare genetic defects of IFN- γ receptors cause disseminated mycobacterial infection. The aim of the present study was to investigate whether genetic polymorphisms found in the Th1-immune response genes play a role in TB. In our study, DNA samples were collected from two series of cases including 832 patients with new smear-positive TB and 506 unrelated individuals with no history of TB in the general

population of Hanoi, Vietnam. Alleles of eight microsatellite markers located around Th1-immune response-related genes and single nucleotide polymorphisms near the promising microsatellites were genotyped. A set of polymorphisms within the interferon gamma receptor 2 gene (*IFNGR2*) showed a significant association with protection against TB ($P = 0.00054$). Resistant alleles tend to be less frequently found in younger age at diagnosis ($P = 0.011$). Luciferase assays revealed high transcriptional activity of the promoter segment in linkage disequilibrium with resistant alleles. We conclude that the polymorphisms of *IFNGR2* may confer resistance to the TB development of newly infected individuals. Contribution of the genetic factors to TB appeared to be different depending on age at diagnosis.

Electronic supplementary material The online version of this article (doi:10.1007/s00439-011-1112-8) contains supplementary material, which is available to authorized users.

M. Hijikata · J. Shojima · I. Matsushita · S. Sakurada · N. Keicho (✉)
Department of Respiratory Diseases, Research Institute, National Center for Global Health and Medicine, 1-21-1 Toyama, Shinjuku-ku, Tokyo 162-8655, Japan
e-mail: nkeicho-ky@umin.ac.jp

K. Tokunaga
Department of Human Genetics, Graduate School of Medicine, The University of Tokyo, Tokyo 113-0033, Japan

J. Ohashi
Molecular and Genetic Epidemiology, Department of Public Health Medicine, Doctoral Program in Life System Medical Sciences, Graduate School of Comprehensive Human Sciences, University of Tsukuba, Ibaraki 305-8077, Japan

N. T. L. Hang
NCGM-BMH Medical Collaboration Center, Hanoi, Vietnam

T. Horie
JICA-Guatemala Chagas Disease Project, Guatemala City, Guatemala

N. P. Hoang · P. H. Thuong · L. T. Lien
Hanoi Lung Hospital, Hanoi, Vietnam

Introduction

Tuberculosis (TB) remains one of the major health problems worldwide (Lopez et al. 2006): According to an estimate, approximately one-third of the world's population is infected with *Mycobacterium tuberculosis* (*M. tuberculosis*), and more than 9 million people develop active TB disease every year. Of these patients, 80% are from 22 high-burden countries including Vietnam (World Health Organization 2010).

Development of TB has been considered to be a two-stage process, infection with *M. tuberculosis* and progression to disease. In total, 5–10% of immunocompetent individuals initially infected with *M. tuberculosis* develop active TB during their lifetime (Frieden et al. 2003). When young vulnerable individuals fail to inhibit growth of the pathogen, they often develop the disease within 2 years of infection. In the remaining individuals, containment of *M. tuberculosis* is successful, though the agent is not

eliminated completely, which leads to life-long latent infection (Russell 2007). When immune levels are impaired after years of infection, reactivation of dormant bacteria leads to disease manifestation, which contributes to the development of elderly TB, though new TB patients affected by re-infection also have to be taken into account (Tufariello et al. 2003). Protective immunity to control the initial infection, orchestrated by immune cells including T cells and macrophages, is influenced by a variety of factors including genetic predisposition (Möller et al. 2010a).

T helper 1 (Th1)-type immune system is crucial to protection against mycobacterial diseases, in which interferon- γ (IFN- γ) has a key role (Lin and Flynn 2010). Although T cell response to mycobacterial infection in human beings is difficult to be addressed experimentally (Cooper 2009), it is known that genetic defects of Th1 molecules can be found in genes such as interferon gamma receptor 1 (*IFNGR1*), interferon gamma receptor 2 (*IFNGR2*), signal transducer and activator of transcription 1, 91 kDa (*STAT1*), interleukin 12B (*IL12B*) and interleukin 12 receptor, beta 1 (*IL12RB1*) cause severe mycobacterial diseases (Zhang et al. 2008). These observations have highlighted IFN- γ /interleukin-12 (IL-12) axis and their polymorphisms have been investigated in mycobacterial infection: association of promoter polymorphism in *IFNGR1* with TB was reported in African populations in independent studies, whereas association of *IFNGR2* with TB has not been published in the literature (Cooke et al. 2006; Stein et al. 2007). Associations with *IL12B* and *IL12RB1* were not consistently shown (Möller et al. 2010b). In the present study, we analyzed genetic polymorphisms of major Th1 cytokine receptors (*IFNGR1*, *IFNGR2*, *IL12RB1* and *IL12RB2*) and signal transduction molecules (*STAT1* and *STAT4*) in Hanoi-Vietnamese and reported a disease association and functional significance of polymorphisms in *IFNGR2*.

Materials and methods

Study population

The patients and control subjects were recruited in Hanoi, Vietnam (Horie et al. 2007). In total, 832 smear-positive pulmonary TB patients without previous episodes of TB (age 41 ± 14.4 , males 77.6%) and 506 healthy volunteers without previous and present history of TB (age 37 ± 10.3 , males 50.0%) participated in this study. All of them were unrelated Hanoi, Vietnamese. TB patients were all recruited immediately after the diagnosis was made. The TB panel A ($n = 277$, age 41 ± 13.5 , males 73.3%) was collected in 2003–2004, and the second TB panel B ($n = 555$, age 41 ± 14.8 , males 79.8%) was collected in

2007–2009. Pulmonary physicians diagnosed all the patients as new active pulmonary TB and treated them with anti-TB drugs based on the guidelines of the national TB program. Informed consent was obtained from all participants. The study protocol was approved by the ethics committees of the Ministry of Health, Vietnam and the National Center for Global Health and Medicine, Japan. Since 4 patients in panel A were human immunodeficiency virus (HIV) positive by previously described PCR assay (Panteleeff et al. 1999) with minor modifications and 49 patients in panel B were HIV seropositive (Hang et al. 2011), they were excluded from further analysis.

Microsatellite markers

We used eight microsatellite markers (*IFNGR1*-MS1, *IFNGR2*-MS1, *IFNGR2*-MS2, *IL12RB1*-MS1, *IL12RB2*-MS1, *IL12RB2*-MS2, *STAT1*-MS1 and *STAT4*-MS1) located in the major Th1-immune response genes (Tanaka et al. 2005) for screening of genetic polymorphisms associated with active TB. A part of the samples, 98 TB patients from the TB panel A and 200 controls were analyzed as described under (Tanaka et al. 2005).

Single nucleotide polymorphisms (SNP) screening in *IFNGR2* of Vietnamese samples

Forty-eight control samples were subjected to PCR amplifications of promoter and seven exon regions of *IFNGR2* and their sequences were analyzed for polymorphisms. GC content of genomic sequence upstream of the translation initiation codon was high (78.2% of nucleotides -1 to -500), and PCR condition was optimized for GC-rich template. The genomic DNA was extracted from anticoagulated blood with QIAamp DNA midi kit (QIAGEN, Hamburg, Germany). PCR was performed using TaKaRa LA Taq with GC buffer I (TaKaRa, Shiga, Japan) with primers 5'-CTCC CAACAGGCGTCAAACGACATGGTG-3' and 5'-TGGTC CCTGCTCCACCGCTGCTACTACAAA-3'. PCR cycling condition was 40 cycles of 95°C for 30 s, 67°C for 30 s and 72°C for 2 min. Amplified products (1,607 bp) were purified and sequenced with BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Foster City, CA, USA) using 3100 Genetic Analyzer (Applied Biosystems), with primers 5'-AGCTTAATATGTACTTTGGGG-3' and 5'-CACCCACTCTGAGCACCCGG-3'. This method was also used for the typing of three promoter SNPs, rs8134145, rs8126756 and rs17882748. Sequencing primers that have the allelic variant at their 3'-end 5'-GGAGGGGTGGGGGC TCCAGGAAA-3', 5'-GCAGGGCCCGCTCTTCCCGA GCA-3' and 5'-GGGCTCCAGGAAAGCCCGGGGT-3' were also designed, and allele-specific sequencing was

performed to directly determine the haplotypes of the three promoter SNPs.

Selection of representative SNPs around *IFNGR2* and genotyping

Representative SNPs around *IFNGR2* were selected from HapMap database (The International HapMap Consortium 2005). SNP genotype data of Han Chinese in Beijing (CHB) encompassing 350 kb from *IL10RB* to *CRYZL1* were analyzed by Haploview 4.2 (Barrett et al. 2005), and 27 representative SNPs were chosen based on the method of block-by-block tags in linkage disequilibrium (LD) blocks determined by confidence interval method (Gabriel et al. 2002). The *IFNGR2* SNPs identified as mentioned above and selected SNPs were genotyped in 273 TB patients of panel A and 506 controls. Genotyping was performed by the Digitag2 assay that has previously been described in another study (Nishida et al. 2007).

Rapid amplification of cDNA end (5'-RACE) of *IFNGR2*

The exact 5' end of exon 1 was confirmed with FirstChoice RLM-RACE Kit (Ambion, Austin, TX, USA) using total RNA of THP-1 cells (ATCC TIB-202) stimulated with 10 ng/ml of phorbol myristate acetate (Schwende et al. 1996), U937 cells (ATCC CRL-1593.2) and Jurkat cells (ATCC TIB-152).

Luciferase assay

Promoter region of *IFNGR2* (Rhee et al. 1996) consisting of 1,167 bp (position -1,172 to -6 of initiation codon) was amplified by PCR and inserted into *Xho* I and *Bgl* II sites of pGL4.10 vector (Promega, Madison, WI, USA). Three plasmids of the observed haplotypes (CCC, ATC, ATT of rs8134145, rs8126756 and rs17882748) were constructed, and their sequences were confirmed to be devoid of any additional nucleotide difference. Reporter plasmids were mixed with pRL-TK (Promega) and transfected to Jurkat human T-cell leukemia cells with Lipofectamin LTX (Invitrogen, Carlsbad, CA, USA) in triplicate. Cells were harvested after 24 h and luciferase activity was measured using Dual-Luciferase Reporter Assay System (Promega). The transfection experiments were repeated twice with three independent subclones of each plasmid.

Statistical analysis

Disease associations with markers were assessed by Chi-square test or Fisher's exact test, and *P* values less than 0.05 were considered significant. Statistical analysis was

performed using Stata version 10 (StataCorp, College Station, TX, USA). When necessary, *P* values were subjected to Bonferroni's correction for multiple comparisons. To determine whether genotype frequencies in the populations are compatible with Hardy–Weinberg equilibrium, Hardy–Weinberg exact tests were carried out using the program Arlequin version 3.11 (Excoffier et al. 2007). To assess pairwise LD between polymorphisms, we calculated Lewontin's *D'* and *r* square (*r*²) for polymorphisms by Haploview version 4.2 (Barrett et al. 2005). TB disease associated with genetic variations was assessed by odds ratios unadjusted or adjusted for sex, age at recruitment and its interaction using logistic models. Tendency of having resistant alleles in the order of age at diagnosis was also tested using a similar logistic model within the TB group. Difference in luciferase activity between the haplotype under consideration and the other haplotypes was assessed by Wilcoxon rank sum/Mann–Whitney *U* test.

Results

Microsatellite markers

Microsatellite marker *IFNGR2*-MS1 located in 5'-upstream region of *IFNGR2* showed significant association with TB even after Bonferroni's correction (Table 1) and the frequency of *IFNGR2*-MS1-325 allele was significantly lower in TB patients than in controls (Supplementary table 1). *IFNGR2*-MS2, the other microsatellite was located in intron 2 of *IFNGR2* and the frequency of *IFNGR2*-MS2-252 allele was also lower in TB patients than in controls (uncorrected *P* = 0.0024), but not significant after Bonferroni's correction. *IFNGR2*-MS1-325 allele and *IFNGR2*-MS2-252 allele were in LD (*D'* = 0.91, *r*² = 0.64).

Screening of genetic polymorphisms in *IFNGR2*

Forty-eight control samples were subjected to PCR amplifications of promoter and seven exons of *IFNGR2* and their sequences were analyzed for possible polymorphisms. In the exonic sequences of *IFNGR2*, a non-synonymous SNP, rs9808753 was found in exon 2, and another SNP, rs1059293 was shown in 3'-untranslated region (UTR) of exon 7, while there were no SNPs in exon–intron boundaries. In the 5' region up to -850 bp of the translation initiation codon, three SNPs, rs8134145, rs8126756 and rs17882748 were also identified.

Genotyping of selected SNPs around *IFNGR2*

Association of microsatellite markers of *IFNGR2* with TB prompted us to identify relevant SNPs that may show

Table 1 Association results for microsatellite markers

Marker	Locus	No. of alleles (total)	No. of alleles (<5% grouped ^a)	<i>P</i> value ^b (2 × <i>m</i>)	Minimum <i>P</i> value ^b (2 × 2)	<i>P</i> value ^c Corrected	<i>P</i> value HWE
<i>IFNGR1</i> -MS1	6q23.3	14	7	0.419	0.0549	NS	0.4858
<i>IFNGR2</i> -MS1	21q22.11	8	5	0.016	0.0009	0.036	0.2762
<i>IFNGR2</i> -MS2		6	4	0.013	0.0024	NS	0.0326
<i>IL12RB1</i> -MS1	19p13.1	4	4	0.366	0.1600	NS	0.1606
<i>IL12RB2</i> -MS1	1p31.3-p31.2	12	6	0.155	0.0267	NS	0.7499
<i>IL12RB2</i> -MS2		6	4	0.540	0.2228	NS	0.7289
<i>STAT1</i> -MS1	2q32.2	13	5	0.563	0.3010	NS	0.0254
<i>STAT4</i> -MS1	2q32.2-q32.3	11	5	0.232	0.1046	NS	0.5243


NS not significant, HWE Hardy–Weinberg equilibrium

^a Alleles with frequencies less than 5% were grouped

^b Fisher's exact test

^c Bonferroni's correction. 2 × 2 minimum *P* value was multiplied by the numbers of total alleles (40)

Fig. 1 Association *P* values and pairwise LD of genotyped polymorphisms around *IFNGR2* region. **a** Association results for 32 SNPs in 273 patients (panel A) and 506 controls are shown. *P* values of microsatellite markers in 98 patients and 200 controls are also included. Positions of genes are shown on the top. **b** Pairwise LD (r^2) between 32 SNPs and 2 microsatellite markers determined by the Haploview program is shown. In the calculation of pairwise LD, microsatellite alleles except for one showing the smallest *P* value were grouped, and the microsatellite locus was regarded as having only two alleles



further association. We genotyped 27 SNPs selected around *IFNGR2*, 3 SNPs in the 5' upstream region, a non-synonymous SNP in exon 2, and a 3'UTR SNP in exon 7 of *IFNGR2* in 273 TB patients (panel A) and 506 controls (Fig. 1, Supplementary table 2). The rs2834213 SNP in intron 2 and the rs1059293 SNP in 3'UTR were associated with TB ($P = 0.0073$, OR 0.69 95% confidence interval [CI] 0.52–0.91; $P = 0.0088$, OR 0.70 95% CI 0.54–0.92). These SNPs were in Hardy–Weinberg equilibrium in the control group. We confirmed that SNPs in other nearby genes were not associated with the disease (Supplementary table 2). As expected, the resistant G allele of rs2834213

and C allele of rs1059293 were both in LD with *IFNGR2*-MS1-325 allele and *IFNGR2*-MS2-252 allele (Supplementary Fig. 1). Particularly, the resistant G allele of rs2834213 in intron 2 was in high LD with *IFNGR2*-MS1-325 allele ($D' = 0.94$, $r^2 = 0.83$), that is located 1.9 kb upstream of the translation initiation codon.

Directly determined haplotypes consisting of three SNPs in the 5' GC-rich region of *IFNGR2*

In addition to single SNPs associated with the disease, we also characterized a set of SNPs in the 5' GC-rich region of

the gene, rs8134145, rs8126756 and rs17882748, since these three SNPs are closely located within 300 bp upstream of the transcription start site as discussed below, which may influence *IFNGR2* expression. When we directly determined haplotypes of three 5' SNPs by allele-specific sequencing in 273 patients and in 506 controls, three common haplotypes (CCC, ATC and ATT) accounted for 99.7% of chromosomes. The haplotype ATC was in high LD with the intron 2 SNP rs2834213 ($D' = 0.97$, $r^2 = 0.82$), and frequencies of the ATC haplotype were significantly lower in patients than in controls ($P = 0.036$, OR 0.76 95% CI 0.58–0.99). Haplotypes carrying SNPs in the entire *IFNGR2* region and their frequencies were estimated in 273 patients and in 506 controls. Consequently, the G allele of the intron 2 SNP rs2834213, the C allele of the 3'UTR SNP rs1059293 and the directly determined haplotype ATC, are uniquely contained in the same haplotype as shown in Supplementary table 3.

Transcription start site (TSS) of *IFNGR2*

In the public database, the aforementioned 5' SNPs, rs8134145, rs8126756, and rs17882748 are regarded as variants in 5' UTR, since TSS of the reference cDNA sequence (NM_005534.3) is located at position –648 of the translation initiation codon. However, multiple TSS were actually reported in *IFNGR2*, the positions of which were distributed from the initiation codon to almost 990 bp upstream, presumably due to cell type differences (Rhee et al. 1996). For this reason, we determined the 5' ends by 5' RACE in our study. As a result, TSS obtained from all immune cell lines tested were 121 bp upstream of the initiation codon. Thus, the positions of the three SNPs were calculated as –295, –285 and –8 from the TSS, indicating that they are promoter variants in these cell types.

Association results of TB panel B

We selected the intron 2 SNP, rs2834213 as a representative SNP for the disease-resistant polymorphisms and genotyped 503 patients in TB panel B, which were compared with the original control subjects ($N = 506$) in Table 2. The G allele of rs2834213 was significantly associated with TB in panel B ($P = 0.0025$, OR 0.71 95% CI 0.57–0.89). In a logistic model to assess possible confounders, adjusted odds ratios was compared with non-adjusted odds ratios for the G allele, which were hardly affected by sex, age at recruitment and its interaction term, indicating that the *IFNGR2* SNP remained significantly associated with TB in dominant and recessive models respectively ($P = 0.016$ and $P = 0.004$; table not shown).

Furthermore, we set up another logistic model to examine the relationship between having the TB-resistant

Table 2 Association results of rs2834213 A/G SNP

Sample	Allele (frequency)		Genotype (%)			P value			OR (95% CI)		
	A	G	A/A	A/G	G/G	Allele	Genotype		Allele	Genotype	
							Dominant	Recessive		Dominant	Recessive
TB panel A	452 (0.837)	88 (0.163)	186 (68.9)	80 (29.6)	4 (1.5)	0.0073	0.047	0.0050	0.69 (0.52–0.91)	0.73 (0.57–0.92)	0.25 (0.08–0.72)
TB panel B	838 (0.833)	168 (0.167)	347 (67.0)	144 (28.6)	12 (2.4)	0.0025	0.015	0.0068	0.71 (0.57–0.89)	0.72 (0.56–0.95)	0.40 (0.20–0.80)
TB combined	1290 (0.834)	256 (0.166)	533 (69.0)	224 (29.0)	16 (2.1)	0.00054	0.0075	0.00048	0.70 (0.57–0.86)	0.73 (0.57–0.92)	0.35 (0.18–0.65)
Controls	786 (0.780)	222 (0.220)	311 (61.7)	164 (32.5)	29 (5.8)						

TB tuberculosis, OR odds ratio, CI confidence interval

Table 3 Tendency of having G allele (rs2834213) in the order of age strata at the time of diagnosis ($N = 757$)

Age at diagnosis (year)	GA or GG genotype (n/N)	(%)	Odds ratio per 10-year change* (95% CI)
16–25	35/124	28.2	0.88 (0.79–0.98)
26–35	43/171	25.1	
36–45	53/165	32.1	
46–55	54/171	31.6	
56–65	35/87	40.2	
65–	15/39	38.5	

* In a logistic model, the trend of having the G allele was calculated as odds ratio when the patients are 10-years younger at the time of diagnosis ($P = 0.019$)

G alleles (as binary outcome) and age at diagnosis (as a continuous variable). In patients from panel A and B ($n = 757$), the TB-resistant G allele was less frequently found, as the age at diagnosis was younger ($P = 0.011$). Similarly, in the age-stratified analysis, when the patients are 10 years younger at the time of diagnosis, the odds ratio (OR) for having the G allele was 0.88 (95% CI, 0.79–0.98) and this trend remained significant ($P = 0.019$) (Table 3).

Luciferase assay

We constructed plasmids containing 5' fragments in which only nucleotide sequences of the three promoter SNPs rs8134145, rs8126756, and rs17882748 are different and measured transcriptional activity of the three promoter segments (CCC, ATC, and ATT) in Jurkat human T-cell leukemia cells. Consequently, the resistant ATC haplotype had significantly higher transcriptional activity than CCC haplotype and ATT haplotype ($P = 0.037$ respectively) by Mann–Whitney U test (Fig. 2).

Discussion

IFN- γ plays a crucial role in host defense against intracellular pathogens mainly through activation of macrophages and regulation of Th1 cell response (Boehm et al. 1997). IL-12 released from dendritic cells and macrophages drives production of IFN- γ via IL-12 receptors, IL12RB1 and IL12RB2, on Th1 cells and subsequent activation of STAT4. In turn, IFN- γ binds to IFN- γ receptors composed of IFNGR1 and IFNGR2 subunits and transduces STAT1 signals to target cells (Bach et al. 1997).

In this study, we first screened eight microsatellite markers within the genes encoding these Th1 cytokine receptors and signal transducers, and demonstrated that the *IFNGR2* marker alleles showed significant association with




Fig. 2 Dual luciferase reporter assays. The ratios of Firefly luciferase activity (signal S) to Renilla luciferase activity (control C) are displayed using box and whisker plots. Three subcloned plasmids were prepared, and each subcloned plasmid was tested in triplicate and S/C values were averaged. The experiments were carried out twice independently. As a result, six independent S/C values were obtained for each haplotype. ATC haplotype showed significantly higher transcriptional activity than CCC haplotype and ATT haplotype (*, ** $P = 0.037$, respectively) by Mann–Whitney U test. No significant difference was observed between CCC and ATT haplotypes (data not shown)

active TB. SNPs around the gene were analyzed and a strong disease association with the intron 2 SNP rs2834213, the 3'UTR SNP rs1059293 and the 5' promoter segment characterized by three SNPs was thus demonstrated. Possible influence of population substructure was kept to a minimum, since their ethnicity was Hanoi Vietnamese in which more than 99% were the Kinh people (Hoa et al. 2008).

To our knowledge, this is the first report of *IFNGR2* polymorphisms associated with TB. Intron 2 SNP, rs2834213 was most robustly associated with TB, but its biological importance is currently unclear. Indeed, it was not located near the splice sites (5,582 nucleotides downstream of splice donor site and 877 nucleotides upstream of splice acceptor site). A SNP in 3'UTR of exon 7 rs1059293 was in strong LD with the rs2834213 and also associated with TB, but it was 99 nucleotides upstream of polyadenylation signal. *IFNGR2* did not carry any non-synonymous SNP in high LD with rs2834213.

In an attempt to search functional polymorphism(s) in strong LD with the intron SNP (rs2834213) further, we identified the 300 bp promoter segment containing three SNPs. HapMap database does not have data of the three promoter SNPs, presumably due to high GC content that hinders high throughput genotyping method. The direct haplotyping revealed that it was also associated with the disease as well as the intron 2 SNP. Although we demonstrated that the promoter ATC haplotype showing an inverse disease association has high transcriptional activity in vitro and may confer resistance to TB, we could not

conclude which polymorphism around *IFNGR2* is primarily responsible for the disease until the functional roles of other SNPs showing more robust association are fully studied.

Among previous TB association studies with Th1-related genes, CC genotype at the -56 C/T SNP (rs2234711) of *IFNGR1* was repeatedly associated with TB in African populations (Cooke et al. 2006; Stein et al. 2007). In our study, *IFNGR1*-MS1-158 allele was in strong LD with -56 SNP (Tanaka et al. 2005), but this *IFNGR1* marker allele was not associated with TB. The lack of association is presumably because of insufficient power to detect weak genetic effects. Otherwise, it could be due to population-specific LD, when the true causative variant was not -56 SNP itself.

Experimental data have shown that *IFNGR2* is a key regulator for IFN- γ -STAT1 signaling in T cells (Schroder et al. 2004; Regis et al. 2006). During the development of Th1 cells, *IFNGR2* transcription is reduced in the IFN- γ rich condition and this reduction alleviates a potentially harmful anti-proliferative action of IFN- γ -STAT1 signaling. However, *IFNGR2* expression is not completely suppressed, because temporary activation of STAT1 is still necessary for Th1 system. *IFNGR2* transcription is thus fine-tuned during the Th1 differentiation process. In the promoter region, the transcriptional activity of the resistant haplotype ATC was higher than the other two common haplotypes in the Jurkat T cell line at baseline levels. Although physiological modulation of *IFNGR2* expression is not easily simulated in a single cell-type model, this segment may have a potential to influence Th1 function through *IFNGR2* regulation.

In this study, another interesting finding is that the resistant allele tend to be less frequently observed in younger patients at the time of diagnosis, a surrogate for age at onset in new patients. This effect was moderate but significant. The allele frequency in older age at diagnosis nearly reached the level of the control population. It is likely that the elderly kept latent infection of *M. tuberculosis* for long years, and the age-associated decline in immune response caused development of active TB, while the younger patients developed active TB soon after initial infection (Tufariello et al. 2003). In intermediate or low burden countries, there are more elderly patients and the effect of the resistant allele of *IFNGR2* may be smaller.

Moreover, in African countries with high rates of TB and HIV co-infection, HIV is the strongest risk factor for TB development (Reid et al. 2006). By contrast, the proportion of HIV-positive TB patients is only 8.8% in the Vietnamese TB panel B and 1.4% in TB panel A, therefore possible effect of the resistant allele on HIV infection could not be determined in this study. In the

previous reports, other polymorphisms of *IFNGR2* were associated with liver fibrosis of chronic hepatitis C virus infection and with viremia of hepatitis B virus infection (Nalpas et al. 2010; Huang et al. 2011). Because IFN- γ is a key cytokine for the control of infectious diseases, association of *IFNGR2* polymorphisms with HIV infection needs be clarified.

One limitation in our study is a single control panel of the Vietnamese population. Results of the first case-control set were only partially confirmed because of incomplete independence of the two study sets, though sample size itself was not small. Another limitation is that our control panel may include asymptomatic individuals with latent TB infection, because performing tuberculin testing is not common in Vietnam. Considering two-stage process of infection with the pathogen and progression to disease, we cannot directly specify which stage of TB was more affected by *IFNGR2* in our study population. Future use of interferon gamma release assays to detect latent infection of *M. tuberculosis* in this field might be helpful to arrive at a solution (Pai et al. 2008). Because of the complexity of LD structure and the age-dependent effect as regards these variations, carefully conducted studies should be undertaken to reproduce our results in other populations. Validation studies by re-sequencing are also warranted. In non-Asian populations, however, the LD of rs2834213 does not appear to reach the promoter region of *IFNGR2* (data not shown), indicating that the functional promoter haplotype may not be easily found in disease marker association studies by the conventional tag SNP-based approach in other populations.

We conclude that the polymorphisms of *IFNGR2* may confer resistance to TB in Vietnam. It appeared to be different depending on age at diagnosis. Further functional studies are needed to elucidate the genetic susceptibility to TB, fully considering complicated immune process regarding early or late onset of the disease.

Ethical standards We declare that these experiments comply with the current laws of Japan and Vietnam.

Acknowledgments We thank Izumi Matsumoto, Masako Okochi and Keiko Wakabayashi for their technical assistance. The authors thank Kazuko Tanabe D.V.M. and Mr. John Crosskey for their critical reading of this manuscript. This work was partly supported by a grant from the Program of Japan Initiative for Global Research Network on Infectious Diseases (J-GRID), Ministry of Education, Culture, Sports, Science and Technology, Japan.

Conflict of interest The authors declare that they have no conflict of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Bach EA, Aguet M, Schreiber RD (1997) The IFN gamma receptor: a paradigm for cytokine receptor signaling. *Annu Rev Immunol* 15:563–591
- Barrett JC, Fry B, Maller J, Daly MJ (2005) Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics* 21:263–265
- Boehm U, Klamp T, Groot M, Howard JC (1997) Cellular responses to interferon-gamma. *Annu Rev Immunol* 15:749–795
- Cooke GS, Campbell SJ, Sillah J, Gustafson P, Bah B, Sirugo G, Bennett S, McAdam KP, Sow O, Lienhardt C, Hill AV (2006) Polymorphism within the interferon-gamma/receptor complex is associated with pulmonary tuberculosis. *Am J Respir Crit Care Med* 174:339–343
- Cooper AM (2009) T cells in mycobacterial infection and disease. *Curr Opin Immunol* 21:378–384
- Excoffier L, Laval G, Schneider S (2007) Arlequin (version 3.0): an integrated software package for population genetics data analysis. *Evol Bioinform Online* 1:47–50
- Frieden TR, Sterling TR, Munsiff SS, Watt CJ, Dye C (2003) Tuberculosis. *Lancet* 362:887–899
- Gabriel SB, Schaffner SF, Nguyen H, Moore JM, Roy J, Blumenstiel B, Higgins J, DeFelice M, Lochner A, Faggart M, Liu-Cordero SN, Rotimi C, Adeyemo A, Cooper R, Ward R, Lander ES, Daly MJ, Altshuler D (2002) The structure of haplotype blocks in the human genome. *Science* 296:2225–2229
- Hang NT, Lien LT, Kobayashi N, Shimbo T, Sakurada S, Thuong PH, le Hong T, Tam do B, Hijikata M, Matsushita I, Hung NV, Higuchi K, Harada N, Keicho N (2011) Analysis of factors lowering sensitivity of interferon- γ release assay for tuberculosis. *PLoS One* 6:e23806
- Hoa BK, Hang NT, Kashiwase K, Ohashi J, Lien LT, Horie T, Shojima J, Hijikata M, Sakurada S, Satake M, Tokunaga K, Sasazuki T, Keicho N (2008) HLA-A, -B, -C, -DRB1 and -DQB1 alleles and haplotypes in the Kinh population in Vietnam. *Tissue Antigens* 71:127–134
- Horie T, Lien LT, Tuan LA, Tuan PL, Sakurada S, Yanai H, Keicho N, Nakata K (2007) A survey of tuberculosis prevalence in Hanoi, Vietnam. *Int J Tuberc Lung Dis* 11:562–566
- Huang HH, Shih WL, Li YH, Wu CF, Chen PJ, Lin CL, Liu CJ, Liaw YF, Lin SM, Lee SD, Yu MW (2011) Hepatitis B viraemia: its heritability and association with common genetic variation in the interferon gamma signalling pathway. *Gut* 60:99–107
- International HapMap Consortium (2005) A haplotype map of the human genome. *Nature* 437:1299–1320
- Lin PL, Flynn JL (2010) Understanding latent tuberculosis: a moving target. *J Immunol* 185:15–22
- Lopez AD, Mathers CD, Ezzati M, Jamison DT, Murray CJ (2006) Global and regional burden of disease and risk factors, 2001: systematic analysis of population health data. *Lancet* 367:1747–1757
- Möller M, de Wit E, Hoal EG (2010a) Past, present and future directions in human genetic susceptibility to tuberculosis. *FEMS Immunol Med Microbiol* 58:3–26
- Möller M, Nebel A, van Helden PD, Schreiber S, Hoal EG (2010b) Analysis of eight genes modulating interferon gamma and human genetic susceptibility to tuberculosis: a case-control association study. *BMC Infect Dis* 10:154
- Nalpas B, Lavielle-Meziani R, Plancoulaine S, Jouanguy E, Nalpas A, Munteanu M, Charlotte F, Ranque B, Patin E, Heath S, Fontaine H, Vallet-Pichard A, Pontoire D, Bourlière M, Casanova JL, Lathrop M, Bréchet C, Poynard T, Matsuda F, Pol S, Abel L (2010) Interferon gamma receptor 2 gene variants are associated with liver fibrosis in patients with chronic hepatitis C infection. *Gut* 59:1120–1126
- Nishida N, Tanabe T, Takasu M, Suyama A, Tokunaga K (2007) Further development of multiplex single nucleotide polymorphism typing method, the DigiTag2 assay. *Anal Biochem* 364:78–85
- Pai M, Zwerling A, Menzies D (2008) Systematic review: T-cell-based assays for the diagnosis of latent tuberculosis infection: an update. *Ann Intern Med* 149:177–184
- Panteleeff DD, John G, Nduati R, Mbori-Ngacha D, Richardson B, Kreiss J, Overbaugh J (1999) Rapid method for screening dried blood samples on filter paper for human immunodeficiency virus type 1 DNA. *J Clin Microbiol* 37:350–353
- Regis G, Conti L, Boselli D, Novelli F (2006) IFN γ STAT1 trafficking tunes IFN γ -STAT1 signaling in T lymphocytes. *Trends Immunol* 27:96–101
- Reid A, Scano F, Getahun H, Williams B, Dye C, Nunn P, De Cock KM, Hankins C, Miller B, Castro KG, Ravigliione MC (2006) Towards universal access to HIV prevention, treatment, care, and support: the role of tuberculosis/HIV collaboration. *Lancet Infect Dis* 6:483–495
- Rhee S, Ebensperger C, Dembic Z, Pestka S (1996) The structure of the gene for the second chain of the human interferon-gamma receptor. *J Biol Chem* 271:28947–28952
- Russell DG (2007) Who puts the tubercle in tuberculosis? *Nat Rev Microbiol* 5:39–47
- Schroder K, Hertzog PJ, Ravasi T, Hume DA (2004) Interferon-gamma: an overview of signals, mechanisms and functions. *J Leukoc Biol* 75:163–189
- Schwende H, Fitzke E, Amb S, Dieter P (1996) Differences in the state of differentiation of THP-1 cells induced by phorbol ester and 1, 25-dihydroxyvitamin D₃. *J Leukoc Biol* 59:555–561
- Stein CM, Zalwango S, Chiunda AB, Millard C, Leontiev DV, Horvath AL, Cartier KC, Chervenak K, Boom WH, Elston RC, Mugerwa RD, Whalen CC, Iyengar SK (2007) Linkage and association analysis of candidate genes for TB and TNF α cytokine expression: evidence for association with IFNGR1, IL-10, and TNF receptor 1 genes. *Hum Genet* 121:663–673
- Tanaka G, Matsushita I, Ohashi J, Tsuchiya N, Ikushima S, Oritsu M, Hijikata M, Nagata T, Yamamoto K, Tokunaga K, Keicho N (2005) Evaluation of microsatellite markers in association studies: a search for an immune-related susceptibility gene in sarcoidosis. *Immunogenetics* 56:861–870
- Tufariello JM, Chan J, Flynn JL (2003) Latent tuberculosis: mechanisms of host and bacillus that contribute to persistent infection. *Lancet Infect Dis* 3:578–590
- World Health Organization (2010) Global tuberculosis control 2010. WHO website http://www.who.int/tb/publications/global_report/2010/
- Zhang SY, Boisson-Dupuis S, Chappier A, Yang K, Bustamante J, Puel A, Picard C, Abel L, Jouanguy E, Casanova JL (2008) Inborn errors of interferon (IFN)-mediated immunity in humans: insights into the respective roles of IFN- α /beta, IFN- γ , and IFN- λ in host defense. *Immunol Rev* 226:29–40

Circulating Levels of Adiponectin, Leptin, Fetuin-A and Retinol-Binding Protein in Patients with Tuberculosis: Markers of Metabolism and Inflammation

Naoto Keicho^{1*}, Ikumi Matsushita¹, Takahiro Tanaka¹, Takuro Shimbo², Nguyen Thi Le Hang³, Shinsaku Sakurada¹, Nobuyuki Kobayashi⁴, Minako Hijikata¹, Pham Huu Thuong⁵, Luu Thi Lien⁵

1 Department of Respiratory Diseases, Research Institute, National Center for Global Health and Medicine, Tokyo, Japan, **2** Department of Clinical Research and Informatics, International Clinical Research Center, National Center for Global Health and Medicine, Tokyo, Japan, **3** National Center for Global Health and Medicine—Bach Mai Hospital Medical Collaboration Center, Hanoi, Viet Nam, **4** Department of Respiratory Medicine, National Center for Global Health and Medicine, Tokyo, Japan, **5** Hanoi Lung Hospital, Hanoi, Viet Nam

Abstract

Background: Wasting is known as a prominent feature of tuberculosis (TB). To monitor the disease state, markers of metabolism and inflammation are potentially useful. We thus analyzed two major adipokines, adiponectin and leptin, and two other metabolic markers, fetuin-A and retinol-binding protein 4 (RBP4).

Methods: The plasma levels of these markers were measured using enzyme-linked immunosorbent assays in 84 apparently healthy individuals (=no-symptom group) and 46 patients with active pulmonary TB around the time of treatment, including at the midpoint evaluation (=active-disease group) and compared them with body mass index (BMI), C-reactive protein (CRP), chest radiographs and TB-antigen specific response by interferon- γ release assay (IGRA).

Results: In the no-symptom group, adiponectin and leptin showed negative and positive correlation with BMI respectively. In the active-disease group, at the time of diagnosis, leptin, fetuin-A and RBP4 levels were lower than in the no-symptom group [adjusted means 2.01 versus 4.50 ng/ml, $P < 0.0001$; 185.58 versus 252.27 $\mu\text{g/ml}$, $P < 0.0001$; 23.88 versus 43.79 $\mu\text{g/ml}$, $P < 0.0001$, respectively]. High adiponectin and low leptin levels were associated with large infiltrates on chest radiographs even after adjustment for BMI and other covariates ($P = 0.0033$ and $P = 0.0020$). During treatment, adiponectin levels increased further and then decreased. Leptin levels remained low. Initial low levels of fetuin-A and RBP4 almost returned to the normal reference range in concert with reduced CRP.

Conclusions: Our data and recent literature suggest that low fat store and underlying inflammation may regulate these metabolic markers in TB in a different way. Decreased leptin, increased adiponectin, or this ratio may be a promising marker for severity of the disease independent of BMI. We should further investigate pathological roles of the balance between these adipokines.

Citation: Keicho N, Matsushita I, Tanaka T, Shimbo T, Hang NTL, et al. (2012) Circulating Levels of Adiponectin, Leptin, Fetuin-A and Retinol-Binding Protein in Patients with Tuberculosis: Markers of Metabolism and Inflammation. PLoS ONE 7(6): e38703. doi:10.1371/journal.pone.0038703

Editor: Pere-Joan Cardona, Fundació Institut d'Investigació en Ciències de la Salut Germans Trias i Pujol, Universitat Autònoma de Barcelona, CIBERES, Spain

Received: January 19, 2012; **Accepted:** May 9, 2012; **Published:** June 7, 2012

Copyright: © 2012 Keicho et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by a grant from the Program of Japan Initiative for Global Research Network on Infectious Diseases, MEXT, Japan. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: nkeicho-ky@umin.ac.jp

Introduction

Tuberculosis (TB) is a major infectious cause of death around the world, with most of the 1.5 million deaths per year attributable to the disease occurring in developing countries. Negative energy balance in chronic inflammation has been recognized as a prominent feature of TB and one of the major obstacles to manage the patients [1,2]. Recent emergence of drug resistant TB is assumed to be driven by poorly implemented drug regimens, but malnutrition as well as HIV co-infection might worsen the condition: Inflammatory responses evoked by infection increase the demand for anabolic energy, leading to a synergistic vicious circle and further deterioration of the clinical condition [3].

It is generally believed that undernourishment diminishes protective immunity against *Mycobacterium tuberculosis*. [4]. A series of animal experiments, particularly aerosol-infected guinea pig models have demonstrated that chronic protein-energy malnutrition reduces secretion of T-helper 1 (Th1) cytokines [5]. It is rapidly reversed with alimentary supplement, indicating a pivotal role of nutrition, although it remains unclear what the optimal nutritional interventions are for improving the human disease in an effective manner [4].

On the other hand, in many countries today, rapid industrialization and urbanization are accompanied by changing patterns of diet and physical activity and this results in over-nutrition [6]. Consequently, a combination of these two unfavor-

Table 1. Characteristics of study population.

	no-symptom group (N=84)	active-disease group (N=46)	P values
Male/Female (n)	41/43	42/4	<0.0001
Age (year)*	40.0 (28.1–48.6)	47.2 (34.7–55.0)	0.0064
BMI (kg/m ²)*	21.8 (20.0–23.7)	18.3 (17.1–19.5)	<0.0001
BCG history (yes/no/unknown)	33/28/23	10/3/33	<0.0001
positive/negative results of IGRA (n)	55/29	41/4**	0.0015

*Median and 25-to-75 percentiles in parenthesis are shown.

**One indeterminate case is not shown here.

doi:10.1371/journal.pone.0038703.t001

able conditions, a slow decline of infectious diseases associated with undernutrition and a rapid increase in obesity and diabetes are a serious double burden to public health and clinical medicine in resource limited settings [7].

Mainly in studies carried out in industrialized countries, fat-cell-derived hormones/cytokines designated as adipokines and relevant mediators have been investigated extensively and proposed as markers of obesity and diabetes [8]. Of these adipokines, adiponectin is a unique insulin sensitizer with atheroprotective role [9,10]. Plasma levels of adiponectin are inversely correlated with body weight and visceral fat mass [11,12]. Leptin is another major adipokine in proportion to fat stores [13,14] and one of the key mediators of energy metabolism [2]. Even mild weight loss induced by dietary restriction is known to reduce leptin levels [11]. These markers supposedly shift towards the opposite in lean patients with wasting diseases. However, the significance of these metabolic markers in chronic infectious diseases like TB has not been fully understood [2].

We have recently conducted a proteomic research and demonstrated that plasma levels of fetuin-A and retinol-binding protein 4 (RBP4), also closely linked to the metabolic and inflammatory state, were significantly lower in patients with active pulmonary TB than in control subjects [15]. Fetuin-A, also known as α 2-*Heremans-Schmid* glycoprotein, is an abundant plasma

component of hepatic origin [16] and a negative regulator of insulin signaling [17,18]. Elevation of plasma fetuin-A is strongly associated with atherogenic lipid profile as well as fatty liver in obese patients [18]. Lipid components in the liver presumably upregulate fetuin-A expression, which may in turn repress adiponectin and impair adipocyte function [19,20]. Fetuin-A is also downregulated in acute inflammation as a negative acute-phase protein [21]. RBP4, synthesized in the liver and adipose tissue, has recently been identified as another adipokine involved in the development of insulin resistance [22]. In humans, similar to leptin, circulating RBP4 levels are high in obesity and decreased after calorie-restriction induced weight loss [11,23]. RBP4 is also known as a specific transporter protein for retinol (vitamin A) and can be used to assess the short-term fluctuation of nutritional states as a rapid turnover protein [24].

Alteration of the circulating levels of these markers should be investigated in TB, since they are expected to provide a basis of a critical link among nutritional status, metabolism and immunity of the disease, and hopefully to consider efficient nutritional interventions. In the present study, we thus measured circulating adiponectin and leptin in addition to fetuin-A and RBP4 levels in patients with active pulmonary TB versus apparently healthy individuals and compared the levels with body mass index (BMI), a simple estimate of adiposity [25] and C-reactive protein (CRP),

Table 2. Correlation of tested marker levels with BMI, CRP and IGRA values in each of the no-symptom and active-disease groups.

Variable	no-symptom group (N=84)			active-disease group (N=46)		
	Pearson's <i>r</i> (P values) ^a					
	by BMI (kg/m ²)	by CRP (μ g/ml)	by IFN- γ (IU/ml) ^b	by BMI (kg/m ²)	by CRP (μ g/ml)	by IFN- γ (IU/ml) ^b
Adiponectin (μ g/ml)	-0.4530 (<0.0001)*	-0.2892 (0.0076)	-0.2254 (0.0393)	-0.4421 (0.0021)	0.1477 (0.3274)	-0.1092 (0.4700)
Leptin (ng/ml)	0.4518 (<0.0001)*	0.1694 (0.1234)	0.1179 (0.2855)	0.2771 (0.0623)	-0.0918 (0.5442)	0.3568 (0.0149)
Leptin/adiponectin ratio	0.5820 (<0.0001)*	0.2793 (0.0101)	0.2067 (0.0592)	0.4901 (0.0005)*	-0.1633 (0.2783)	0.2804 (0.0591)
Fetuin-A (μ g/ml)	0.0309 (0.7805)	0.0415 (0.7079)	0.0322 (0.7714)	0.1243 (0.4105)	-0.1833 (0.2226)	0.2402 (0.1078)
RBP4 (μ g/ml)	0.1605 (0.1447)	-0.0213 (0.8475)	0.0716 (0.5173)	0.1535 (0.3085)	-0.3018 (0.0415)	-0.0916 (0.5448)

^aPearson's correlation coefficients with P values were calculated. Plasma concentrations were analyzed after logarithmic transformation.

^bTB-antigen stimulated IFN- γ response

*Statistically significant when the significance level is set as $P < 0.002$ based on the Bonferroni correction.

doi:10.1371/journal.pone.0038703.t002

Table 3. BMI, CRP and tested marker levels in IGRA-positive and -negative subgroups in the no-symptom group.

marker	IGRA-negative (N = 29)		IGRA-positive (N = 55)		P values (ANCOVA)
	adjusted mean ^a	(95%CI)	adjusted mean ^a	(95%CI)	
BMI (kg/m ²)	21.52	(20.58–22.46)	21.48	(20.74–22.22)	0.9392
CRP (μg/ml)	1.12	(0.60–2.08)	1.30	(0.80–2.12)	0.6663
Adiponectin (μg/ml)	7.19	(5.67–9.11)	6.39	(5.30–7.70)	0.3792
Leptin (ng/ml)	4.50	(3.34–6.05)	4.38	(3.47–5.54)	0.8783
Leptin/adiponectin ratio	0.63	(0.40–0.97)	0.69	(0.49–0.97)	0.7080
Fetuin-A (μg/ml)	234.22	(212.40–258.29)	263.88	(244.26–285.06)	0.0333
RBP4 (μg/ml)	39.64	(32.28–48.69)	42.88	(36.45–50.43)	0.4997

^aEstimated means of plasma concentrations were compared after logarithmic transformation, being adjusted for gender and age as covariates. The data shown are transformed back to the original unit.

No P values were statistically significant when the significance level is set as $P < 0.007$ based on the Bonferroni correction.

doi:10.1371/journal.pone.0038703.t003

a representative positive acute phase protein [26]. We further characterized their relationship with disease severity and alterations during the course of treatment.

Methods

Study design

We randomly selected and used plasma samples and demographic information in 46 patients with active pulmonary TB (= active-disease group) without treatment history as a biomarker sub-study of a large cohort study [27]. All patients entered the study from July 2007 to March 2009. Diagnosis of active pulmonary TB was made clinically and radiologically and confirmed bacteriologically in Hanoi Lung Hospital. A sputum smear test showed positive results in all of the patients in the active disease group and all of them completed anti-TB treatment following the national standard regimen, 2 months of streptomycin, isoniazid, rifampicin, and pyrazinamide followed by 6 months of isoniazid and ethambutol (2SHRZ/6HE).

Chest radiographs were taken at the time of diagnosis and interpreted by two readers independently in a blind manner. The presence of cavitory lesions and the number of lung zones (zero to six corresponding to the upper, middle, and lower fields on the

right and left sides of the lung) affected by infiltrates were recorded [28]. HIV status was examined before starting anti-TB treatment. The proportion of HIV co-infection is less than 10% in this study area and those with HIV positive were excluded from the drawing up of this sub-study.

As a reference, we also measured plasma samples derived from 84 apparently healthy men and women who may have chances of direct or indirect contacts with TB patients as health care staff (= no-symptom group). All participants were tested for TB-antigen specific interferon- γ response by the commercially available enzyme-linked immunosorbent assay (ELISA)-based interferon- γ release assay (IGRA), QuantiFERON-TB Gold In-Tube™ (Cellestis, Victoria, Australia). In the no-symptom group, IGRA-positive individuals suspected of latent TB infection were recommended to take chest radiography and to confirm there were no active pulmonary lesions. Subsequently a chance of receiving isoniazid prophylactic therapy was given. The protocol was approved by ethical committees of the Ministry of Health, Viet Nam and National Center for Global Health and Medicine, Japan respectively and written informed consent was obtained from each participant.

Table 4. BMI, CRP and tested marker levels in the no-symptom and active-disease groups after adjustment for gender and age.

marker	no-symptom group (N = 84)		active-disease group (N = 46)		P values (ANCOVA)
	adjusted mean ^a	(95%CI)	adjusted mean ^a	(95%CI)	
BMI (kg/m ²)	21.68	(21.06–22.30)	17.65	(16.66–18.65)	<0.0001*
CRP (μg/ml)	1.22	(0.86–1.74)	36.88	(20.94–64.94)	<0.0001*
Adiponectin (μg/ml)	6.82	(5.73–8.12)	9.29	(7.02–12.30)	0.0136
Leptin (ng/ml)	4.50	(3.78–5.35)	2.01	(1.52–2.66)	<0.0001*
Leptin/adiponectin ratio	0.66	(0.50–0.88)	0.22	(0.14–0.34)	<0.0001*
Fetuin-A (μg/ml)	252.27	(234.55–271.33)	185.58	(165.07–208.64)	<0.0001*
RBP4 (μg/ml)	43.79	(38.09–50.34)	23.88	(19.08–29.88)	<0.0001*

^aEstimated means of plasma concentrations were compared after logarithmic transformation, being adjusted for gender and age as covariates. The data shown are transformed back to the original unit.

*Statistically significant when the significance level is set as $P < 0.007$ based on the Bonferroni correction.

doi:10.1371/journal.pone.0038703.t004

Table 5. CRP and tested marker levels in the no-symptom and active-disease groups after adjustment for gender, age and BMI.

marker	no-symptom group (N = 84)		active-disease group (N = 46)		P values (ANCOVA)
	adjusted mean ^a	(95%CI)	adjusted mean ^a	(95%CI)	
CRP (μg/ml)	1.11	(0.77–1.60)	47.80	(25.36–90.09)	<0.0001*
Adiponectin (μg/ml)	7.80	(6.63–9.19)	6.39	(4.81–8.49)	0.1671
Leptin (ng/ml)	3.77	(3.26–4.37)	3.28	(2.54–4.24)	0.2790
Leptin/adiponectin ratio	0.48	(0.38–0.61)	0.51	(0.35–0.76)	0.7704
Fetuin-A (μg/ml)	248.04	(229.95–267.57)	194.46	(170.48–221.80)	0.0004*
RBP4 (μg/ml)	42.90	(37.08–49.63)	25.27	(19.62–32.55)	0.0001*

^aEstimated means of plasma concentrations were compared after logarithmic transformation, being adjusted for gender, age and BMI as covariates. The data shown are transformed back to the original unit.

*Statistically significant when the significance level is set as $P < 0.008$ based on the Bonferroni correction.

doi:10.1371/journal.pone.0038703.t005

Measurements of markers of metabolism and inflammation

Immediately after making the diagnosis of active TB disease, heparinized blood samples were drawn for IGRA before starting anti-TB treatment (0 month) and the remaining plasma without mixing any stimulants was reserved in a -80°C freezer until measurement. Samples were collected twice again, after the initial phase of treatment (2 months) and at the end of treatment (7 months) in the active disease group. This study was originally intended to identify a variety of biomarkers associated with TB phenotypes [15] and the participants were not obliged to keep fasting. The blood was collected in the daytime between 8 am and 4 pm at the outpatient clinic to avoid interference in dosing schedule of anti-TB drugs.

The AssayMax Human C-Reactive Protein ELISA kit was used for detection of human c-reactive protein (CRP) in plasma (Assaypro LLC, St. Charles, MO, USA). The minimum detectable dose was less than 0.25 ng/ml. The Quantikine[®] Human Total Adiponectin/Acrp30 Immunoassay kit was used to detect total (low, middle and high molecular weight) human adiponectin in plasma (R&D Systems, Inc.; Minneapolis, MN, USA). The mean

minimum detectable dose was 0.246 ng/ml. The Quantikine[®] Human Leptin Immunoassay kit was used to detect human leptin in plasma (R&D Systems, Inc.). The mean minimum detectable dose was 7.8 pg/ml. The AHSB ELISA kit was used to detect fetuin-A in plasma (BioVender Laboratory Medicine Inc.; Modrice, Czech Republic). The detection limit was 0.35 ng/ml. A competitive ELISA for quantitative determination of RBP4 in human plasma was also applied (AdipoGen Inc.; Seoul, Korea) and the detection limit was 1 ng/ml. All were performed according to the manufacturer's instructions. Differences in measured concentrations between EDTA plasma samples as reference and these heparin samples were within a range of variation generally accepted in ELISA (coefficient of variance <15%) (data not shown).

Statistical analysis

Plasma protein levels were served for subsequent statistical analysis after logarithmic transformation of the measurements to minimize distortion of the data distribution. Means of demographic data between two groups were compared by analysis of variance (ANOVA) after testing for equal variances and

Table 6. BMI, CRP and tested marker levels in patients with small and large infiltrates on chest radiographs after adjustment for gender and age.

marker	small infiltrates ^a (N = 22)		large infiltrates ^a (N = 23)		P values (ANCOVA)
	adjusted mean ^b	(95%CI)	adjusted mean ^b	(95%CI)	
BMI (kg/m ²)	18.73	(16.74–20.71)	18.11	(15.95–20.27)	0.3065
CRP (μg/ml)	26.14	(12.63–54.10)	35.92	(16.29–79.21)	0.1520
Adiponectin (μg/ml)	10.28	(5.38–19.66)	18.83	(9.31–38.11)	0.0033*
Leptin (ng/ml)	2.42	(1.64–3.57)	1.65	(1.08–2.52)	0.0020*
Leptin/adiponectin ratio	0.24	(0.11–0.52)	0.09	(0.04–0.21)	0.0002*
Fetuin-A (μg/ml)	201.97	(149.87–272.18)	184.68	(133.52–255.46)	0.3222
RBP4 (μg/ml)	36.14	(21.76–60.03)	31.56	(18.17–54.79)	0.3770
IFN-γ (IU/ml) ^c	11.04	(2.13–57.16)	5.80	(0.97–34.82)	0.2039

^aSmall infiltrates = less than 3 of 6 zones in the lung affected, large infiltrates = 3 or more than 3 of 6 zones affected

^bEstimated means of plasma concentrations were compared after logarithmic transformation, being adjusted for gender and age as covariates. The data shown are transformed back to the original unit.

^cTB-antigen stimulated IFN-γ response

*Statistically significant when the significance level is set as $P < 0.006$ based on the Bonferroni correction.

doi:10.1371/journal.pone.0038703.t006

Table 7. CRP and tested marker levels in patients with small and large infiltrates on chest radiographs after adjustment for gender, age and BMI.

marker	small infiltrates ^a (N=22)		large infiltrates ^a (N=23)		P values (ANCOVA)
	adjusted mean ^b	(95%CI)	adjusted mean ^b	(95%CI)	
CRP ($\mu\text{g/ml}$)	26.59	(12.78–55.28)	35.50	(16.02–78.63)	0.1991
Adiponectin ($\mu\text{g/ml}$)	10.84	(6.01–19.53)	18.15	(9.57–34.40)	0.0061*
Leptin (ng/ml)	2.37	(1.63–3.47)	1.67	(1.11–2.52)	0.0040*
Leptin/adiponectin ratio	0.22	(0.11–0.44)	0.09	(0.04–0.20)	0.0002*
Fetuin-A ($\mu\text{g/ml}$)	200.77	(148.59–271.28)	185.46	(133.74–257.18)	0.3886
RBP4 ($\mu\text{g/ml}$)	35.69	(21.43–59.46)	31.83	(18.29–55.42)	0.4626
IFN- γ (IU/ml) ^c	11.41	(2.17–59.90)	5.68	(0.94–34.53)	0.1760

^aSmall infiltrates=less than 3 of 6 zones in the lung affected, large infiltrates=3 or more than 3 of 6 zones affected

^bEstimated means of plasma concentrations were compared after logarithmic transformation, being adjusted for gender, age and BMI as covariates. The data shown are transformed back to the original unit.

^cTB-antigen stimulated IFN- γ response

*Statistically significant when the significance level is set as $P < 0.007$ based on the Bonferroni correction.

doi:10.1371/journal.pone.0038703.t007

proportions between two groups were compared by the chi-squared test. Since it is well known that levels of adipokines such as leptin are influenced by gender and age, measurements of protein markers in any two groups were compared by analysis of covariance (ANCOVA) to allow for the covariates. The relationship between markers and other parameters were assessed by Pearson's correlation coefficients. Overall alterations of the measurements at three time points were initially analyzed by repeated-measures ANOVA and only when statistically significant, post-hoc comparisons were proceeded to: Difference of values between two time points was assessed by the paired-T test, under

normal approximation based on the central limit theorem. P values < 0.05 were considered to be statistically significant in general. When the Bonferroni correction was applied, however, a level of statistical significance was set as $0.05/n$ (n = the number of comparisons). Statistical analysis was performed using Stata version 11 (StataCorp, College Station, TX, USA).




Figure 1. CRP and tested marker levels in patients with active TB before (0 month), during (2 months) and at the end (7 months) of anti-TB treatment (N=46). Vertical bars with diamonds on the left side (M and F) indicate reference values, means \pm SEM of the values in men (N=41) and women (N=43) of the no-symptom group. A horizontal bar indicates the grand mean of the values in each condition. * indicates $P < 0.05$ by paired comparison between 0 month and 2 months. When significant, 2 months and 7 months were also compared. doi:10.1371/journal.pone.0038703.g001

Results

Characteristics of study population

The no-symptom group consisted of 84 apparently healthy individuals, whose blood samples were used to obtain the standard values of markers in the study population. This group includes an approximately equal number of men and women with median age of 40, and more than half of the individuals had latent TB infection diagnosed by the IGRA method (Table 1). The active-disease group members were 46 patients with smear-positive active pulmonary TB. The majority of the patients were male with low body mass index ($BMI < 18.5 \text{ kg/m}^2$) and the median age was 47, slightly older than in the non-symptom group.

Correlation of adiponectin, leptin, fetuin-A and RBP4 levels with BMI, CRP and IGRA values in the no-symptom and active-disease groups

Correlation coefficients (r) were calculated in the no-symptom and active-disease groups respectively (Table 2). Adiponectin and leptin showed negative and positive correlations with BMI respectively in the no-symptom group ($r = -0.4530$, $P < 0.0001$; $r = 0.4518$, $P < 0.0001$). Leptin/adiponectin ratio showed a positive correlation with BMI in the active-disease group ($r = 0.4901$, $P = 0.0005$) as well as in the no-symptom group ($r = 0.5820$, $P < 0.0001$). These correlations were statistically significant even after Bonferroni correction for multiple comparisons. The other possible correlations including a pair of leptin and TB-antigen stimulated IFN- γ response did not reach significant levels in this study, when Bonferroni correction was applied.

Pairwise correlations between four tested markers

Pairwise correlation coefficients (r) between four tested metabolic markers were further calculated in the no-symptom and active-disease groups respectively (Table S1). A significant correlation was found only between fetuin-A and RBP4 levels ($r = 0.4007$, $P = 0.0058$) in the active disease group.

Adiponectin, leptin, fetuin-A and RBP4 levels with IGRA-positive and -negative subgroups in the no-symptom group

IGRA-positive values higher than the cutoff value, 0.35 IU/ml are regarded as latent TB infection after active disease is ruled out. We thus categorized the no-symptom group into IGRA-positive and -negative subgroups and compared plasma concentrations of the above markers. However, none of the marker levels including fetuin-A were significantly different between IGRA-positive and -negative subgroups after adjustment for gender and age, when considering the number of comparisons (Table 3).

Adiponectin, leptin, fetuin-A and RBP4 levels in the no-symptom and active-disease groups

The active-disease group had significantly low BMI and very high CRP levels at the time of diagnosis, when assessed by using ANCOVA with adjusted means (Table 4). In the disease group, leptin, leptin/adiponectin ratio, fetuin-A and RBP4 levels were remarkably lower than in the no-symptom group ($P < 0.0001$ respectively) after adjustment for gender and age and these differences were statistically significant even after Bonferroni correction (Table 4).

Since BMI was strongly correlated with some of the adipokine values as shown in Table 2, we further analyzed levels of the four markers after adjustment for BMI as well as gender and age. Consequently, adiponectin and leptin levels were not significantly

different between the two groups any more, whereas fetuin-A and RBP4 levels remained significant ($P = 0.0004$ and $P = 0.0001$) (Table 5)

Adiponectin, leptin, fetuin-A and RBP4 levels in patients with mild and severe disease

At the time of diagnosis, severity of the disease was assessed by spread of infiltrates on chest radiographs (Table 6). Small infiltrates affecting less than 3 of the 6 lung zones and large ones affecting more, categorized the patients into two subgroups (= mild and severe disease) half-and-half.

After adjustment for gender and age, adiponectin levels were higher and leptin levels were lower in patients with large infiltrates than in those with small infiltrates ($P = 0.0033$ and $P = 0.0020$). Interestingly, differences in the levels of these two adipokines between small and large infiltrates were significant respectively ($P = 0.0061$ and $P = 0.0040$), even after adjustment for BMI as well as gender and age (Table 7). Leptin/adiponectin ratio was lower, or adiponectin/leptin ratio was higher, in patients with large infiltrates than in those with small infiltrates independent of BMI ($P = 0.0002$). None of the markers were associated with the presence of cavity on the chest radiographs (data not shown).

Adiponectin, leptin, fetuin-A and RBP4 levels in patients with active TB before, during and at the end of anti-TB treatment

Figure 1 shows plasma values at the time points before (0 month), during (2 months) and at the end (7 months) of anti-TB treatment. Mean values in men ($N = 41$) and women ($N = 43$) of the no-symptom group are shown as a reference, in which gender difference was observed in leptin levels and leptin/adiponectin ratio ($P < 0.0001$).

Overall differences of the measurements during anti-TB treatment in all of these four markers were statistically significant by repeated-measures ANOVA ($P < 0.01$). Post-hoc analysis showed that adiponectin levels increased transiently ($P = 0.0004$; 0 month vs. 2 months) and then decreased close to the reference range by the end of treatment ($P < 0.0001$; 2 months vs. 7 months). Leptin levels remained low throughout the treatment course, though gradually elevated ($P = 0.0226$; 0 month vs. 2 months). Initial low levels of fetuin-A and RBP4 significantly improved during treatment ($P = 0.0001$ and $P = 0.0016$; 0 month vs. 2 months), almost reaching the reference range by the end in concert with reduced CRP levels.

Discussion

We assessed the clinical significance of four metabolic markers, adiponectin, leptin, fetuin-A and RBP4 in patients with active TB, analyzing them in relation to classical nutritional and inflammatory parameters, BMI and CRP, severity of disease and treatment course. BMI is known to be lower in patients with active TB than in control subjects [1,2]. After effective treatment, weight often increases but patients may remain underweight [11].

Plasma levels of adiponectin were inversely correlated with BMI in concordance with previous results [11,12]. The adiponectin levels tended to be elevated in the active-disease group characterized by low BMI, though it did not reach significant levels, which was also shown by others [29]. Interestingly in our study, adiponectin levels were significantly higher in severe disease with extensive pulmonary lesions than in mild disease, even after adjustment for BMI. Adiponectin as a modulator of inflammation in a variety of diseases has recently been highlighted [30]. For instance, in critically ill patients, adiponectin levels appear to be

transiently suppressed at the initial phase and then gradually elevated at the recovery phase [31,32]. The plasma concentrations in patients with active TB were further increased after starting treatment and then decreased close to the reference range by the end of treatment. Elevated adiponectin levels in chronic inflammatory diseases may be explained by compensatory response to the underlying disease as well as concomitant low body fat mass, which is postulated by others [33,34]. A study designed to measure alteration of adiponectin and BMI simultaneously throughout the treatment period would be able to characterize it further.

In most recent reports, leptin levels are low in TB [29,35–38], though other earlier or smaller studies have shown conflicting results [39–42]. In the present study, using a commercial ELISA, significantly lower levels of leptin were demonstrated in patients with active TB, which could be mostly explained by marked undernutrition in our disease population. Within the active-disease group, however, correlation between leptin and BMI was less clear. BMI-independent regulation of plasma leptin concentrations should also be taken into consideration in TB at least in part [13,37]. This idea is also supported by an *ex vivo* study by others demonstrating that continuous exposure of IL-1 or TNF- α provides a signal to downregulate leptin in human adipose tissue [43], though acute inflammation such as sepsis may rather upregulate circulating leptin levels transiently [44–46]. In addition to relatively high levels of adiponectin, low levels of leptin were observed in patients with large infiltrates, even after adjustment for BMI. This is concordant with a recent study showing that leptin levels were low in severe TB disease [29]. We have further demonstrated that low leptin/adiponectin ratio, or high adiponectin/leptin ratio is characteristic to severe TB disease in this study. This ratio was originally proposed as an atherogenic index indicating a balance between the two markers bearing apparently opposite functions in inflammation [47]. Our findings support the idea that suppressed production of leptin may be detrimental to host defense against TB by virtue of impairment of Th1 cell-mediated immunity [13,29,48]. After starting treatment, leptin levels were slightly elevated, but remained low during the treatment period. This is also compatible with reports made by others [37,38], although the mechanism remains unknown. Long-lasting low levels of leptin may be attributed to individual predisposition to TB or delayed recovery from wasting disease.

In our study, fetuin-A levels were considerably low in TB even after adjustment for BMI. Soon after starting treatment, the levels were increased in inverse proportion to the decrease in CRP. In TB, fetuin-A may be downregulated by at least dual mechanisms, strongly mediated by underlying inflammation [21] and partly controlled by depleted liver fat due to wasting or malnutrition [18]. Low fetuin-A levels may also result in impairment of macrophage function to kill the pathogen and ectopic calcification possibly in TB lesions [49,50].

RBP4 levels were also low in TB even after adjustment for BMI. Throughout the treatment course, the levels were gradually elevated close to the reference range inversely with the decrease in CRP. These findings are supported by a recent report demon-

strating that RBP4 rapidly decreases during acute inflammation, possibly acting as a negative acute phase reactant, similar to fetuin-A, albumin and prealbumin [21,51,52]. This may partly explain a close positive correlation with fetuin-A demonstrated in the active-disease group. In addition to dual regulation of RBP4 by underlying inflammation and low body fat mass, reduced renal function is also known to cause retention of the circulating levels, such that further caution is needed to interpret RBP4 measurement in disease state [53].

Our study has several limitations. Firstly, many types of nutrients including micronutrients are essential to the human body but the potential interplay between each component of nutrients was not within our scope at that time. Secondly, since change of BMI was not measured during treatment, direct comparison of improved BMI with the corresponding marker levels was not possible. Thirdly, blood was collected during the daytime without enforced fasting. Although, of course, this increases the variance of measurements, it can be inferred that daytime variations on circulating adipokines and leptin [54] are not as large as to seriously affect conclusive results of comparisons within and between groups in this study. Finally, computer tomography, which has advantages over chest radiography as an imaging tool, was not available in our setting.

Overall, our data and recent literature would suggest that all of the four markers tested are controlled partly by low fat store and partly by inflammation in TB but their regulatory mechanisms are more or less different and interactions with other relevant factors including insulin sensitivity and cellular immunity are worth further investigation. In particular, leptin, adiponectin and their ratio may be promising markers for severity of the wasting disease. Since nutritional intervention has a potential to improve prognosis of intractable TB such as HIV co-infection and MDR-TB, large-scale prospective studies using selected biomarkers to investigate metabolic contributors to disease phenotype are desired. The more fully we understand the mechanisms linking diet, health, and disease, the more effective will be our ability to design optimal interventions.

Supporting Information

Table S1 Pairwise correlations between four tested markers. (DOC)

Acknowledgments

We thank Tokie Totsu and Fumi Toshioka for their technical assistance. The authors thank Kazuko Tanabe D.V.M. and Mr. John Crosskey for their critical reading of this manuscript.

Author Contributions

Conceived and designed the experiments: N. Keicho IM TT N. Kobayashi SS. Performed the experiments: IM. Analyzed the data: N. Keicho IM NTLH TS. Contributed reagents/materials/analysis tools: IM TT NTLH SS MH PHT LTL. Wrote the paper: N. Keicho.

References

1. Tverdal A (1986) Body mass index and incidence of tuberculosis. *Eur J Respir Dis* 69: 355–362.
2. Schaible UE, Kaufmann SH (2007) Malnutrition and infection: complex mechanisms and global impacts. *PLoS Med* 4: e115.
3. Prentice AM, Gershwin ME, Schaible UE, Keusch GT, Victora CG, et al. (2008) New challenges in studying nutrition-disease interactions in the developing world. *J Clin Invest* 118: 1322–1329.
4. Cegielski JP, McMurray DN (2004) The relationship between malnutrition and tuberculosis: evidence from studies in humans and experimental animals. *Int J Tuberc Lung Dis* 8: 286–298.
5. Dai G, McMurray DN (1998) Altered cytokine production and impaired antimycobacterial immunity in protein-malnourished guinea pigs. *Infect Immun* 66: 3562–3568.
6. Leung CC, Lam TH, Chan WM, Yew WW, Ho KS, et al. (2007) Lower risk of tuberculosis in obesity. *Arch Intern Med* 167: 1297–1304.
7. Dooley KE, Chaisson RE (2009) Tuberculosis and diabetes mellitus: convergence of two epidemics. *Lancet Infect Dis* 9: 737–746.
8. Msaad S, Haynes EN (2007) Biomarkers of obesity and subsequent cardiovascular events. *Epidemiol Rev* 29: 98–114.

9. Yang WS, Lee WJ, Funahashi T, Tanaka S, Matsuzawa Y, et al. (2001) Weight reduction increases plasma levels of an adipose-derived anti-inflammatory protein, adiponectin. *J Clin Endocrinol Metab* 86: 3815–3819.
10. Pitombo C, Araújo EP, De Souza CT, Pareja JC, Geloneze B, et al. (2006) Amelioration of diet-induced diabetes mellitus by removal of visceral fat. *J Endocrinol* 191: 699–706.
11. Klempel MC, Varady KA (2011) Reliability of leptin, but not adiponectin, as a biomarker for diet-induced weight loss in humans. *Nutr Rev* 69: 145–154.
12. Kuo SM, Halpern MM (2011) Lack of association between body mass index and plasma adiponectin levels in healthy adults. *Int J Obes (Lond)* 35: 1487–1494.
13. Malli F, Papaioannou AI, Gourgoulis KI, Daniil Z (2010) The role of leptin in the respiratory system: an overview. *Respir Res* 11: 152.
14. Klimcakova E, Kovacicova M, Stich V, Langin D (2010) Adipokines and dietary interventions in human obesity. *Obes Rev* 11: 446–456.
15. Tamaka T, Sakurada S, Kano K, Takahashi E, Yasuda K, et al. (2011) Identification of tuberculosis-associated proteins in whole blood supernatant. *BMC Infect Dis* 11: 71.
16. Denecke B, Gräber S, Schäfer C, Heiss A, Wölje M, et al. (2003) Tissue distribution and activity testing suggest a similar but not identical function of fetuin-B and fetuin-A. *Biochem J* 376: 135–145.
17. Mori K, Emoto M, Yokoyama H, Araki T, Teramura M, et al. (2006) Association of serum fetuin-A with insulin resistance in type 2 diabetic and nondiabetic subjects. *Diabetes Care* 29: 468.
18. Stefan N, Hennige AM, Staiger H, Machann J, Schick F, et al. (2006) Alpha2-Heremans-Schmid glycoprotein/fetuin-A is associated with insulin resistance and fat accumulation in the liver in humans. *Diabetes Care* 29: 853–857.
19. Hennige AM, Staiger H, Wicke C, Machicao F, Fritsche A, et al. (2008) Fetuin-A induces cytokine expression and suppresses adiponectin production. *PLoS One* 3: e1765.
20. Dasgupta S, Bhattacharya S, Biswas A, Majumdar SS, Mukhopadhyay S, et al. (2010) NF-kappaB mediates lipid-induced fetuin-A expression in hepatocytes that impairs adipocyte function effecting insulin resistance. *Biochem J* 429: 451–462.
21. Gangneux C, Daveau M, Hiron M, Derambure C, Papaconstantinou J, et al. (2003) The inflammation-induced down-regulation of plasma Fetuin-A (alpha2HS-Glycoprotein) in liver results from the loss of interaction between long C/EBP isoforms at two neighbouring binding sites. *Nucleic Acids Res* 31: 5957–5970.
22. Yang Q, Graham TE, Mody N, Preitner F, Peroni OD, et al. (2005) Serum retinol binding protein 4 contributes to insulin resistance in obesity and type 2 diabetes. *Nature* 436: 356–362.
23. Vitkova M, Klimcakova E, Kovacicova M, Valle C, Moro C, et al. (2007) Plasma levels and adipose tissue messenger ribonucleic acid expression of retinol-binding protein 4 are reduced during calorie restriction in obese subjects but are not related to diet-induced changes in insulin sensitivity. *J Clin Endocrinol Metab* 92: 2330–2335.
24. Marshall WJ (2008) Nutritional assessment: its role in the provision of nutritional support. *J Clin Pathol* 61: 1083–1088.
25. van Lettow M, Fawzi WW, Semba RD (2003) Triple trouble: the role of malnutrition in tuberculosis and human immunodeficiency virus co-infection. *Nutr Rev* 61: 81–90.
26. Schultz DR, Arnold PI (1990) Properties of four acute phase proteins: C-reactive protein, serum amyloid A protein, alpha 1-acid glycoprotein, and fibrinogen. *Semin Arthritis Rheum* 20: 129–147.
27. Hang NT, Lien LT, Kobayashi N, Shimbo T, Sakurada S, et al. (2011) Analysis of factors lowering sensitivity of interferon- γ release assay for tuberculosis. *PLoS One* 6: e23806.
28. Sakurada S, Hang NT, Ishizuka N, Toyota E, Hung LD, et al. (2012) Inter-rater agreement in the assessment of abnormal chest X-ray findings for tuberculosis between two Asian countries. *BMC Infect Dis* 12: 31.
29. Santucci N, D'Attilio L, Kovalevski L, Bozza V, Besedovsky H, et al. (2011) A multifaceted analysis of immune-endocrine-metabolic alterations in patients with pulmonary tuberculosis. *PLoS One* 6: e26363.
30. Robinson K, Prins J, Venkatesh B (2011) Clinical review: adiponectin biology and its role in inflammation and critical illness. *Crit Care* 15: 221.
31. Langouche L, Vander Perre S, Frystyk J, Flyvbjerg A, Hansen TK, et al. (2009) Adiponectin, retinol-binding protein 4, and leptin in protracted critical illness of pulmonary origin. *Crit Care* 13: R112.
32. Walkey AJ, Rice TW, Konter J, Ouchi N, Shibata R, et al. (2010) Plasma adiponectin and mortality in critically ill subjects with acute respiratory failure. *Crit Care Med* 38: 2329–2334.
33. Moriconi N, Kraenzlin M, Müller B, Keller U, Nusbaumer CP, et al. (2006) Body composition and adiponectin serum concentrations in adult patients with cystic fibrosis. *J Clin Endocrinol Metab* 91: 1506–1509.
34. Sood A (2010) Obesity, adipokines, and lung disease. *J Appl Physiol* 108: 744–753.
35. Schwenk A, Hodgson L, Rayner CF, Griffin GE, Macallan DC (2003) Leptin and energy metabolism in pulmonary tuberculosis. *Am J Clin Nutr* 77: 392–398.
36. van Lettow M, van der Meer JW, West CE, van Crevel R, Semba RD (2005) Interleukin-6 and human immunodeficiency virus load, but not plasma leptin concentration, predict anorexia and wasting in adults with pulmonary tuberculosis in Malawi. *J Clin Endocrinol Metab* 90: 4771–4776.
37. van Crevel R, Karyadi E, Netea MG, Verhoef H, Nelwan RH, et al. (2002) Decreased plasma leptin concentrations in tuberculosis patients are associated with wasting and inflammation. *J Clin Endocrinol Metab* 87: 758–763.
38. Buyukoglan H, Gulmez I, Kelesimur F, Kart L, Oymak FS, et al. (2007) Leptin levels in various manifestations of pulmonary tuberculosis. *Mediators Inflamm* 2007: 64859.
39. Yuksel I, Sencan M, Dokmetas HS, Dokmetas I, Ataseven H, et al. (2003) The relation between serum leptin levels and body fat mass in patients with active lung tuberculosis. *Endocr Res* 29: 257–264.
40. Cakir B, Yonem A, Guler S, Odabasi E, Demirbas B, et al. (1999) Relation of leptin and tumour necrosis factor alpha to body weight changes in patients with pulmonary tuberculosis. *Horm Res* 52: 279–283.
41. Bornstein SR, Preas HL, Chrousos GP, Sufredini AF (1998) Circulating leptin levels during acute experimental endotoxemia anti-inflammatory therapy in humans. *J Infect Dis* 178: 887–890.
42. Kim JH, Lee CT, Yoon HI, Song J, Shin WG, et al. (2010) Relation of ghrelin, leptin and inflammatory markers to nutritional status in active pulmonary tuberculosis. *Clin Nutr* 29: 512–8.
43. Bruun JM, Pedersen SB, Kristensen K, Richelsen B (2002) Effects of pro-inflammatory cytokines and chemokines on leptin production in human adipose tissue in vitro. *Mol Cell Endocrinol* 190: 91–99.
44. Arnalich F, Lopez J, Codoceo R, Jim M, Madero R, et al. (1999) Relationship of plasma leptin to plasma cytokines and human survival in sepsis and septic shock. *J Infect Dis* 180: 908–911.
45. Guadillo O, Eiras S, Lago F, Diéguez C, Casanueva FF (2000) Elevated serum leptin concentrations induced by experimental acute inflammation. *Life Sci* 67: 2433–2441.
46. Wallace AM, Sattar N, Mcmillan DC (2000) The co-ordinated cytokine/hormone response to acute injury incorporates leptin. *Cytokine* 12: 1042–1045.
47. Satoh N, Naruse M, Usui T, Tagami T, Suganami T, et al. (2004) Leptin-to-adiponectin ratio as a potential atherogenic index in obese type 2 diabetic patients. *Diabetes Care* 27: 2488–2490.
48. Wieland CW, Florquin S, Chan ED, Leemans JC, Weijer S, et al. (2005) Pulmonary Mycobacterium tuberculosis infection in leptin-deficient ob/ob mice. *Int Immunol* 17: 1399–1408.
49. Jahnen-Dechent W, Schäfer C, Ketteler M, McKee MD (2008) Mineral chaperones: a role for fetuin-A and osteopontin in the inhibition and regression of pathologic calcification. *J Mol Med (Berl)* 86: 379–389.
50. Jersmann HP, Dransfield I, Hart SP (2003) Fetuin/alpha2-HS glycoprotein enhances phagocytosis of apoptotic cells and macrophage apoptosis by human macrophages. *Clin Sci (Lond)* 105: 273–278.
51. Koch A, Weiskirchen R, Sanson E, Zimmermann HW, Voigt S, et al. (2010) Circulating retinol binding protein 4 in critically ill patients before specific treatment: prognostic impact and correlation with organ function, metabolism and inflammation. *Crit Care* 14: R179.
52. Fuhrman MP, Charney P, Mueller CM (2004) Hepatic proteins and nutrition assessment. *J Am Diet Assoc* 104: 1258–1264.
53. Ingenbleek Y, Van Den Schrieck HG, De Nayer P, De Visscher M (1975) Albumin, transferrin and the thyroxine-binding prealbumin/retinol-binding protein (TBPA-RBP) complex in assessment of malnutrition. *Clin Chim Acta* 63: 61–67.
54. Gavriła A, Peng CK, Chan JL, Mietus JE, Goldberger AL, et al. (2003) Diurnal and ultradian dynamics of serum adiponectin in healthy men: comparison with leptin, circulating soluble leptin receptor, and cortisol patterns. *J Clin Endocrinol Metab* 88: 2838–2843.

RESEARCH ARTICLE

Open Access

Inter-rater agreement in the assessment of abnormal chest X-ray findings for tuberculosis between two Asian countries

Shinsaku Sakurada^{1†}, Nguyen TL Hang^{2†}, Naoki Ishizuka¹, Emiko Toyota³, Le D Hung⁴, Pham T Chuc⁴, Luu T Lien⁴, Pham H Thuong⁴, Pham TN Bich², Naoto Keicho^{1*} and Nobuyuki Kobayashi¹

Abstract

Background: Inter-rater agreement in the interpretation of chest X-ray (CXR) films is crucial for clinical and epidemiological studies of tuberculosis. We compared the readings of CXR films used for a survey of tuberculosis between raters from two Asian countries.

Methods: Of the 11,624 people enrolled in a prevalence survey in Hanoi, Viet Nam, in 2003, we studied 258 individuals whose CXR films did not exclude the possibility of active tuberculosis. Follow-up films obtained from accessible individuals in 2006 were also analyzed. Two Japanese and two Vietnamese raters read the CXR films based on a coding system proposed by Den Boon et al. and another system newly developed in this study. Inter-rater agreement was evaluated by kappa statistics. Marginal homogeneity was evaluated by the generalized estimating equation (GEE).

Results: CXR findings suspected of tuberculosis differed between the four raters. The frequencies of infiltrates and fibrosis/scarring detected on the films significantly differed between the raters from the two countries ($P < 0.0001$ and $P = 0.0082$, respectively, by GEE). The definition of findings such as primary cavity, used in the coding systems also affected the degree of agreement.

Conclusions: CXR findings were inconsistent between the raters with different backgrounds. High inter-rater agreement is a component necessary for an optimal CXR coding system, particularly in international studies. An analysis of reading results and a thorough discussion to achieve a consensus would be necessary to achieve further consistency and high quality of reading.

Background

Despite its several disadvantages, chest radiography remains an important supporting tool in tuberculosis (TB) surveys and clinical management of active disease [1-3]. Chest X-ray (CXR) findings should be carefully assessed because of its potential problems such as low specificity and insufficient reproducibility [4].

In this context, reading methods that are less influenced by raters are required and several CXR coding systems have been proposed [5-7]. In general, complex interpretation codes hamper intra- and inter-rater

agreement and simple codes are preferred [6,7], because reproducible and validated coding system may be useful in monitoring disease in clinical and epidemiological studies [8,9].

Previous studies suggest that variability in CXR interpretation among raters is attributed to subjective reading accompanied by insufficient experience or different professional background of the raters [7,10-12]. However, the relationship between agreement levels and relevant factors that may cause disagreement, particularly influence of medical background including different national origins has not been characterized.

In the present study, Vietnamese and Japanese raters studied the readings of suspected TB lesions on CXR films taken during a survey of TB prevalence in Hanoi,

* Correspondence: nkeicho-ty@umin.ac.jp

† Contributed equally

¹National Center for Global Health and Medicine, Tokyo, Japan
Full list of author information is available at the end of the article

Viet Nam [3]. The follow-up films were also compared with the initial films. As analytical tools, two different types of coding systems were used: One was previously reported by another group [5] and the other was newly developed in this study. The aim of the study was to highlight inter-rater agreement between raters with different medical backgrounds. We also attempted to characterize the optimal codes or coding systems used in international studies for a simple and objective evaluation of CXR findings suspected of TB.

Methods

Ethics approval

This study was approved by the ethics committees of the Ministry of Health, Viet Nam and the National Center for Global Health and Medicine (formerly, International Medical Center of Japan). Written informed consent was obtained from each participant prior to the investigations, including the prevalence survey and the follow-up study.

Study population

A population-based TB prevalence survey of 11,624 people aged 15 and over was conducted in Hanoi in 2003 as reported previously [3]. Briefly, subjects suspected of having active TB based on CXR or on symptoms underwent sputum smear microscopy and/or mycobacterial culture. Details of HIV status were not obtained from the study subjects. According to the report of World Health Organization during this period, estimated prevalence of HIV co-infection in new TB patients aged 15-49 was relatively low (2.8%) in Viet Nam [13].

Barring 317 individuals, active TB was radiographically excluded for the rest. Of these 317 individuals, 22 (6.9%) were diagnosed by bacteriological methods, including sputum culture [3]. In 2004, individuals who presented with radiographic findings during the initial survey were advised to undergo sputum smear and culture tests following the World Health Organization recommendation [14,15]. In the 2006 follow-up, in which the same group of individuals was recalled for plain chest radiographic examination (AGFA X-ray film, Beijing, China; Shimadzu UD 150L-30V, Kyoto, Japan) and sputum test, including direct smear and culture. Using a questionnaire, we collected information regarding individual history, additional examinations performed, and treatment for TB undergone after the initial survey. Demographic information (including addresses) collected during the prevalence survey was used to trace the target group in the follow-up period.

The CXR films analyzed in this study were those in which active TB had not been radiographically excluded during the prevalence survey and were those taken during the follow-up in 2006. In total, 258 of the 317 films in the

prevalence survey and 93 follow-up films were available at the time of analysis in this study. The rest of TB-suspected films in the prevalence survey were missing.

CXR coding systems and reading of films

Two coding systems were used to classify the CXR findings. The chest radiograph reading and recording system (CRRS) was developed in 2005 to detect TB and other forms of lung disease [5]. Profusion score and details of abnormalities unrelated to TB were omitted. All the other coding items of this system were retained. A Japan-Vietnam CXR coding system (JVCS) (Figure 1) consisting of rather simple codes was also used: We newly developed this system, considering a registration form used in a public payment system for TB treatment expenses in Japan and reading practice in Viet Nam. CRRS classifies parenchymal abnormalities as primary or secondary lesions depending on the significance of the lesion. In contrast, JVCS does not consider the significance of the lesion, though it records pleural effusion and thickening separately. Additionally, CRRS classifies nodules based on their size and calcification, whereas JVCS separately records nodules and calcification.

Two Japanese pulmonary physicians (E.T. and N.K.) and two Vietnamese radiologists (L.D.H. and P.T.C.) read the CXR films. These readers were different from those who read the CXR films during the initial survey. All CXR films were first read using CRRS. After the completion of readings by CRRS, CXR films were read using JVCS without the results of CRRS being made known to the readers. Each reader was also blinded to the others' readings and clinical information. Instruction and training regarding the two coding systems were given prior to the actual reading. The four raters were asked to reach a consensus while assessing 10 standard films from Japan and another 10 films from Viet Nam.

Statistical analysis

We adopted a double entry system of data entry. JMP version 7.0.1 (SAS Institute Inc., Cary, NC, USA) and SAS version 9.1 (SAS Institute Inc.) were used for analysis. Kappa statistics were used to investigate inter-rater agreement on the presence or absence of lesions of interest. We adopted the following guidelines for interpretation of kappa coefficients: < 0, poor agreement; 0-0.20, slight; 0.21-0.40, fair; 0.41-0.60, moderate; 0.61-0.80, good; and 0.81-1.00, very good [16-18]. Weighted kappa was used to assess inter-rater agreement on variables with more than two categories. McNemar's test or its extension, Bowker's test of symmetry, was used to investigate the symmetry of disagreement between two raters, which tests whether the frequency of an abnormality detected by one rater is significantly different from that by another rater. The generalized estimation equation (GEE) was also used to test the