

Fig. 4 – Protein spots with differing intensities between tumors with vascular invasion and those without. The ratio of means was calculated by dividing the average intensities of the samples of tumor tissues with vascular invasion with those with no vascular invasion. The localization of these protein spots is shown in Supplementary Fig. 2, and the detailed data for protein identification are shown in Supplementary Tables 5 and 6.

3.6. Functional significance of CapG overexpression in tumor cells

To examine the functional significance of the upregulation of CapG in the tumor cells, we examined the effects of CapG silencing on the malignant potential of liver cancer cells. Western blotting showed that CapG silencing was achieved by siRNA transfection (Fig. 6A). The invasion potential was remarkably inhibited by CapG silencing (Fig. 6B and C). In contrast, cell proliferation was not affected by CapG silencing by siRNA transfection (Fig. 6D).

4. Discussion

Prediction of early recurrence is critically important for risk-stratification therapy in HCC. The patients who experience recurrence within 1 year following curative resection have poor prognosis compared to that of others [1]. Various anti-cancer drugs are used to prevent recurrence after surgical resection of HCC [4–9]. The therapeutic strategy is improved by selecting patients who are most likely to have early recurrence after surgery. Alternatively, it may be possible to




Fig. 5 – Validation of CapG expression by western blotting. CapG expression was examined in the samples used for the 2D-DIGE experiments by using specific antibodies against CapG. Rabbit and chicken antibodies against CapG were used in A and B, respectively. The upper panels show the western blot image, and the lower panels show the results of the quantitative analysis of the western blot data. Note that the expression level of CapG was equivalent between non-tumor and tumor tissues and was significantly different between the tumor tissues from the patients with and without vascular invasion.

avoid seriously invasive treatment in patients whose prognosis is expected to be poor even after extensive treatment. Vascular invasion reflects the invasiveness of HCC and has been considered as an independent risk factor for early recurrence and poor prognosis in HCC [2,22–24]. Therefore, understanding the molecular mechanisms underlying vascular invasion will lead to the development of novel therapies and will eventually benefit patients with HCC.

We identified the association of a higher expression level of CapG with vascular invasion in surgically resected tissues by using a proteomic approach. CapG was originally identified as the protein that binds to the morphologically defined barbed end of actin filaments [25]. Overexpression of CapG is highly associated with the malignant features of tumor cells in lung cancer [26,27], cholangiocarcinoma [15], colorectal cancer [28], breast cancer [21,29,30], ovarian cancer [31], oral cancer [20], and pancreatic cancer [17]. The presence of higher CapG expression in the tumor stroma than in the normal counterpart tissue has also been observed in nasopharyngeal carcinoma [32]. Interestingly, the higher expression of CapG is unique to the tumor margin in the case of breast cancer, suggesting that CapG plays an especially important role in metastasis [29]. In contrast, another study reported that CapG might act as a tumor suppressor gene in stomach cancer, lung cancer, and melanoma [33]. Therefore, the roles of CapG in tumor cells may depend on the type of malignancy. As there have been no reports concerning the expression of CapG in HCC to date, our study will provide further understanding of the molecular mechanisms underlying HCC.

Previous studies have reported the mechanisms underlying the regulation of CapG expression. The activation of tyrosine kinases [34], hypoxia-inducible factor 1 (HIF1) [35], integrin [36], and AP-1 [37] causes upregulation of CapG. The aberrant expression of these proteins has been reported in HCC and is correlated with a poor clinical outcome [38–41]. These proteins may regulate CapG expression in a coordinated manner and contribute to the progression of the HCC. It is worth investigating whether the overexpression of CapG directly promotes the vascular invasion or functions as one of the components of protein network for malignant features.

We found that CapG silencing resulted in reduced invasion but not growth inhibition of the cells. These observations were similar to the results observed in a previous study, wherein Tonack et al. reported that the overexpression of CapG induced enhanced motility but did not induce any changes in cellular proliferation in pancreatic cancer cells [42]. CapG may contribute to venous invasion only by promoting the invasiveness of tumor cells, and there may be the other proteins which upregulate cell proliferation and promote invasion in a coordinate manner with CapG.

Higher expression levels of CapG have been found to be related to malignant features of tumor cells, and the clinical usefulness of CapG suggests that it should form part of the approach for lung cancer [27,43], cholangiocarcinoma [15], colorectal cancer [28,43], ovarian cancer [31], and pancreatic cancer treatments [17]. In this study, we were unable to reproduce the correlation between CapG expression and cancer progression by immunohistochemical analysis using




Fig. 6 – Functional assessment of CapG in liver cancer cells in vitro. **A:** Western blotting showed that the expression of CapG was markedly diminished by transfection with siRNAs (siRNA1 and 2) for CapG. **B:** Cell growth was not affected by transfection with siRNAs (siRNA1 and 2) for CapG. **C:** The appearance of liver cancer cells after transfection with the control, siRNA1, and siRNA2. The number of invasive cells decreased after transfection with siRNAs (siRNA1 and 2). **D:** Quantitative results of the invasion assay. Note that the silencing of CapG by siRNA resulted in a significant reduction of the invasive potential of liver cancer cells.

the antibodies for western blotting (Fig. 5) (data not shown), and the clinical usefulness of CapG in HCC remains to be challenged. The antibody suitable for immunochemical validation will be key to the prognostic and predictive applications of CapG. It is worth considering the information in the

antibody databases such as the Human Protein Atlas (www.proteinatlas.org) and the Antibodypedia (www.antibodypedia.com) for the further immunochemical validation studies.

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.jprot.2012.10.004>.

Acknowledgments

This study was supported by the National Cancer Center Research Core Facility and by the National Cancer Center Research and Development Fund (23-A-8 and 23-A-10).

REFERENCES

- [1] Thorgeirsson SS, Grisham JW. Molecular pathogenesis of human hepatocellular carcinoma. *Nat Genet* 2002;31:339-46.
- [2] Poon RT, Fan ST, Ng IO, Lo CM, Liu CL, Wong J. Different risk factors and prognosis for early and late intrahepatic recurrence after resection of hepatocellular carcinoma. *Cancer* 2000;89:500-7.
- [3] el-Assal ON, Yamanoi A, Soda Y, Yamaguchi M, Yu L, Nagasue N. Proposal of invasiveness score to predict recurrence and survival after curative hepatic resection for hepatocellular carcinoma. *Surgery* 1997;122:571-7.
- [4] Monaco AP. The role of mTOR inhibitors in the management of posttransplant malignancy. *Transplantation* 2009;87:157-63.
- [5] Feng YX, Wang T, Deng YZ, Yang P, Li JJ, Guan DX, et al. Sorafenib suppresses postsurgical recurrence and metastasis of hepatocellular carcinoma in an orthotopic mouse model. *Hepatology* 2011;53:483-92.
- [6] Mazzaferro V, Romito R, Schiavo M, Mariani L, Camerini T, Bhoori S, et al. Prevention of hepatocellular carcinoma recurrence with alpha-interferon after liver resection in HCV cirrhosis. *Hepatology* 2006;44:1543-54.
- [7] Qu LS, Jin F, Huang XW, Shen XZ. Interferon-alpha therapy after curative resection prevents early recurrence and improves survival in patients with hepatitis B virus-related hepatocellular carcinoma. *J Surg Oncol* 2010;102:796-801.
- [8] Shimizu M, Takai K, Moriwaki H. Strategy and mechanism for the prevention of hepatocellular carcinoma: phosphorylated retinoid X receptor alpha is a critical target for hepatocellular carcinoma chemoprevention. *Cancer Sci* 2009;100:369-74.
- [9] Wang SM, Zhu J, Pan LF, Liu YK. Inhibitory effect of dimeric beta peptide on the recurrence and metastasis of hepatocellular carcinoma in vitro and in mice. *World J Gastroenterol* 2008;14:3054-8.
- [10] Yokoo H, Kondo T, Okano T, Nakanishi K, Sakamoto M, Kosuge T, et al. Protein expression associated with early intrahepatic recurrence of hepatocellular carcinoma after curative surgery. *Cancer Sci* 2007;98:665-73.
- [11] Orimo T, Ojima H, Hiraoka N, Saito S, Kosuge T, Kakisaka T, et al. Proteomic profiling reveals the prognostic value of adenomatous polyposis coli-end-binding protein 1 in hepatocellular carcinoma. *Hepatology* 2008;48:1851-63.
- [12] Budhu A, Forgues M, Ye QH, Jia HL, He P, Zanetti KA, et al. Prediction of venous metastases, recurrence, and prognosis in hepatocellular carcinoma based on a unique immune response signature of the liver microenvironment. *Cancer Cell* 2006;10:99-111.
- [13] Wong CM, Wong CC, Lee JM, Fan DN, Au SL, Ng IO. Sequential alterations of miRNA expression in hepatocellular carcinoma development and venous metastasis. *Hepatology* 2012;55:1453-61.
- [14] Kondo T, Hirohashi S. Application of highly sensitive fluorescent dyes (CyDye DIGE Fluor saturation dyes) to laser microdissection and two-dimensional difference gel electrophoresis (2D-DIGE) for cancer proteomics. *Nat Protoc* 2007;1:2940-56.
- [15] Morofuji N, Ojima H, Onaya H, Okusaka T, Shimada K, Sakamoto Y, et al. Macrophage-capping protein as a tissue biomarker for prediction of response to gemcitabine treatment and prognosis in cholangiocarcinoma. *J Proteomics* 2012;75:1577-89.
- [16] Dahl E, Sadr-Nabavi A, Klopocki E, Betz B, Grube S, Kreutzfeld R, et al. Systematic identification and molecular characterization of genes differentially expressed in breast and ovarian cancer. *J Pathol* 2005;205:21-8.
- [17] Thompson CC, Ashcroft FJ, Patel S, Saraga G, Vimalachandran D, Prime W, et al. Pancreatic cancer cells overexpress gelsolin family-capping proteins, which contribute to their cell motility. *Gut* 2007;56:95-106.
- [18] Lal A, Lash AE, Altschul SF, Velculescu V, Zhang L, McLendon RE, et al. A public database for gene expression in human cancers. *Cancer Res* 1999;59:5403-7.
- [19] Van Ginkel PR, Gee RL, Walker TM, Hu DN, Heizmann CW, Polans AS. The identification and differential expression of calcium-binding proteins associated with ocular melanoma. *Biochim Biophys Acta* 1998;1448:290-7.
- [20] Nomura H, Uzawa K, Ishigami T, Kouzu Y, Koike H, Ogawara K, et al. Clinical significance of gelsolin-like actin-capping protein expression in oral carcinogenesis: an immunohistochemical study of premalignant and malignant lesions of the oral cavity. *BMC Cancer* 2008;8:39.
- [21] Renz M, Betz B, Niederacher D, Bender HG, Langowski J. Invasive breast cancer cells exhibit increased mobility of the actin-binding protein CapG. *Int J Cancer* 2008;122:1476-82.
- [22] Utsunomiya T, Shimada M, Taguchi KI, Hasegawa H, Yamashita Y, Hamatsu T, et al. Clinicopathologic features and postoperative prognosis of multicentric small hepatocellular carcinoma. *J Am Coll Surg* 2000;190:331-5.
- [23] Paquet KJ, Gad HA, Lazar A, Koussouris P, Mercado MA, Heine WD, et al. Analysis of factors affecting outcome after hepatectomy of patients with liver cirrhosis and small hepatocellular carcinoma. *Eur J Surg* 1998;164:513-9.
- [24] Schoniger-Hekele M, Muller C, Kutilek M, Oesterreicher C, Ferenci P, Gangl A. Hepatocellular carcinoma in Central Europe: prognostic features and survival. *Gut* 2001;48:103-9.
- [25] Casella JF, Maack DJ, Lin S. Purification and initial characterization of a protein from skeletal muscle that caps the barbed ends of actin filaments. *J Biol Chem* 1986;261:10915-21.
- [26] Shao F, Zhang R, Don L, Ying K. Overexpression of gelsolin-like actin-capping protein is associated with progression of lung adenocarcinoma. *Tohoku J Exp Med* 2011;225:95-101.
- [27] Zhu WY, Hunag YY, Liu XG, He JY, Chen DD, Zeng F, et al. Prognostic evaluation of CapG, gelsolin, P-gp, GSTP1, and topo-II proteins in non-small cell lung cancer. *Anat Rec (Hoboken)* 2012;295:208-14.
- [28] Wu JH, Tian XY, Hao CY. The significance of a group of molecular markers and clinicopathological factors in identifying colorectal liver metastasis. *Hepatogastroenterology* 2011;58:1182-8.
- [29] Kang S, Kim MJ, An H, Kim BG, Choi YP, Kang KS, et al. Proteomic molecular portrait of interface zone in breast cancer. *J Proteome Res* 2010;9:5638-45.
- [30] Xu SG, Yan PJ, Shao ZM. Differential proteomic analysis of a highly metastatic variant of human breast cancer cells using two-dimensional differential gel electrophoresis. *J Cancer Res Clin Oncol* 2010;136:1545-56.
- [31] Partheen K, Levan K, Osterberg L, Claesson I, Fallenius G, Sundfeldt K, et al. Four potential biomarkers as prognostic factors in stage III serous ovarian adenocarcinomas. *Int J Cancer* 2008;123:2130-7.
- [32] Li MX, Xiao ZQ, Chen YH, Peng F, Li C, Zhang PF, et al. Proteomic analysis of the stroma-related proteins in nasopharyngeal carcinoma and normal nasopharyngeal epithelial tissues. *Med Oncol* 2010;27:134-44.
- [33] Watari A, Takaki K, Higashiyama S, Li Y, Satomi Y, Takao T, et al. Suppression of tumorigenicity, but not anchorage

- independence, of human cancer cells by new candidate tumor suppressor gene CapG. *Oncogene* 2006;25:7373-80.
- [34] Pierce A, Unwin RD, Evans CA, Griffiths S, Carney L, Zhang L, et al. Eight-channel iTRAQ enables comparison of the activity of six leukemogenic tyrosine kinases. *Mol Cell Proteomics* 2008;7:853-63.
- [35] Liao SH, Zhao XY, Han YH, Zhang J, Wang LS, Xia L, et al. Proteomics-based identification of two novel direct targets of hypoxia-inducible factor-1 and their potential roles in migration/invasion of cancer cells. *Proteomics* 2009;9:3901-12.
- [36] Chen M, Sinha M, Luxon BA, Bresnick AR, O'Connor KL. Integrin alpha6beta4 controls the expression of genes associated with cell motility, invasion, and metastasis, including S100A4/metastasin. *J Biol Chem* 2009;284:1484-94.
- [37] Bahassi el M, Karyala S, Tomlinson CR, Sartor MA, Medvedovic M, Hennigan RF. Critical regulation of genes for tumor cell migration by AP-1. *Clin Exp Metastasis* 2004;21:293-304.
- [38] Huynh H, Ong RW, Li PY, Lee SS, Yang S, Chong LW, et al. Targeting receptor tyrosine kinase pathways in hepatocellular carcinoma. *Anticancer Agents Med Chem* 2011;11:560-75.
- [39] Kong J, Pan B, Ke S, Dong S, Li X, Zhou A, et al. Insufficient radiofrequency ablation promotes angiogenesis of residual hepatocellular carcinoma via HIF-1alpha/VEGFA. *PLoS One* 2012;7:e37266.
- [40] Bergamini C, Sgarra C, Trerotoli P, Lupo L, Azzariti A, Antonaci S, et al. Laminin-5 stimulates hepatocellular carcinoma growth through a different function of alpha6beta4 and alpha3beta1 integrins. *Hepatology* 2007;46:1801-9.
- [41] Guo LL, Xiao S, Guo Y. Activation of transcription factors NF-kappaB and AP-1 and their relations with apoptosis associated-proteins in hepatocellular carcinoma. *World J Gastroenterol* 2005;11:3860-5.
- [42] Tonack S, Patel S, Jalali M, Nedjadi T, Jenkins RE, Goldring C, et al. Tetracycline-inducible protein expression in pancreatic cancer cells: effects of CapG overexpression. *World J Gastroenterol* 2011;17:1947-60.
- [43] Zheng Z, Li J, He X, Chen X, Yu B, Ji J, et al. Involvement of RhoGDI2 in the resistance of colon cancer cells to 5-fluorouracil. *Hepatogastroenterology* 2010;57:1106-12.



Fbxw7 regulates lipid metabolism and cell fate decisions in the mouse liver

Ichiro Onoyama,^{1,2} Atsushi Suzuki,^{3,4} Akinobu Matsumoto,^{1,2} Kengo Tomita,⁵ Hideki Katagiri,⁶ Yuichi Oike,^{4,7} Keiko Nakayama,^{2,8} and Keiichi I. Nakayama^{1,2}

¹Department of Molecular and Cellular Biology, Medical Institute of Bioregulation, Kyushu University, Higashi-ku, Fukuoka, Fukuoka, Japan.

²CREST, Japan Science and Technology Agency, Kawaguchi, Saitama, Japan. ³Division of Organogenesis and Regeneration, Medical Institute of Bioregulation, Kyushu University, Higashi-ku, Fukuoka, Fukuoka, Japan. ⁴PRESTO, Japan Science and Technology Agency, Kawaguchi, Saitama, Japan.

⁵Division of Gastroenterology and Hepatology, Department of Internal Medicine, National Defense Medical College, Tokorozawa, Saitama, Japan.

⁶Division of Advanced Therapeutics for Metabolic Diseases, Center for Translational and Advanced Animal Research, Tohoku University Graduate School of Medicine, Sendai, Japan. ⁷Department of Molecular Genetics, Graduate School of Medical Sciences, Kumamoto University, Kumamoto, Japan. ⁸Division of Developmental Genetics, Center for Translational and Advanced Animal Research, Tohoku University Graduate School of Medicine, Sendai, Japan.

E3 ubiquitin ligase complexes of the SCF type consist of ring-box 1 (Rbx1), cullin 1 (Cul1), S-phase kinase-associated protein 1 (Skp1), and a member of the F-box family of proteins. The identity of the F-box protein determines the substrate specificity of the complex. The F-box family member F-box- and WD repeat domain-containing 7 (Fbxw7; also known as Fbw7, SEL-10, hCdc4, and hAgo) targets for degradation proteins with wide-ranging functions, and uncovering its *in vivo* role has been difficult, because *Fbxw7*^{-/-} embryos die in utero. Using two different Cre-loxP systems (*Mx1-Cre* and *Alb-Cre*), we generated mice with liver-specific null mutations of *Fbxw7*. Hepatic ablation of *Fbxw7* resulted in hepatomegaly and steatohepatitis, with massive deposition of triglyceride, a phenotype similar to that observed in humans with nonalcoholic steatohepatitis. Both cell proliferation and the abundance of *Fbxw7* substrates were increased in the *Fbxw7*-deficient liver. Long-term *Fbxw7* deficiency resulted in marked proliferation of the biliary system and the development of hamartomas. *Fbxw7* deficiency also skewed the differentiation of liver stem cells toward the cholangiocyte lineage rather than the hepatocyte lineage *in vitro*. This bias was corrected by additional loss of the Notch cofactor RBP-J, suggesting that Notch accumulation triggered the abnormal proliferation of the biliary system. Together, our results suggest that *Fbxw7* plays key roles, regulating lipogenesis and cell proliferation and differentiation in the liver.

Introduction

The abundance of cellular proteins is regulated in a coordinated manner at the levels of their synthesis and degradation. In particular, intracellular proteolysis is thought to be subject to highly specific regulation. The ubiquitin-proteasome system is responsible for such specific degradation of proteins, with ubiquitylation playing the regulatory role in this process. Ubiquitylation of target proteins is mediated by the sequential action of 3 enzymes: a ubiquitin-activating enzyme (E1), a ubiquitin-conjugating enzyme (E2), and a ubiquitin ligase (E3). The ubiquitylated substrates are then selectively recognized and degraded by the 26S proteasome (1). Uncontrolled proteolysis is implicated in dysregulation of cell proliferation and aberrant cell differentiation and is thought to underlie many human malignancies (2).

F-box proteins determine the substrate specificity of the SCF-type E3 complex, which consists of the RING-finger protein ring-box 1 (Rbx1; also known as Roc1 and Hrt1), the scaffold protein cullin 1 (Cul1), and the adaptor protein S-phase kinase-associated protein 1 (Skp1) in addition to an F-box protein (2–4). F-box- and WD repeat domain-containing 7 (Fbxw7; also known as Fbw7, SEL-10, hCdc4, and hAgo) is a member of the F-box protein family that was initially identified as a negative regulator of LIN-12-mediated (Notch-mediated) signaling in *Caenorhabditis elegans* by genetic analysis (5, 6). Fbxw7 also interacts with Notch family proteins and promotes their ubiquitin-dependent turnover in mammalian cells (5, 7, 8).

Furthermore, it targets for degradation various mammalian proteins that control cell cycle progression (2, 4), including cyclin E (9–11), c-Myc (12, 13), and c-Jun (14, 15), as well as other proteins that do not contribute directly to cell cycle control, such as SREBPs (16–18), mammalian target of rapamycin (mTOR) (19), and PPAR- γ coactivator-1 α (PGC-1 α) (20).

Given its ability to promote degradation of cyclin E, c-Myc, c-Jun, and Notch, all of which are products of proto-oncogenes, Fbxw7 was expected to function as an oncosuppressor protein. Indeed, mutations in the *Fbxw7* gene have been detected in many types of human malignancy, including cholangiocarcinoma and T cell acute lymphoblastic leukemia as well as pancreatic, gastric, colorectal, prostate, and endometrial cancer (21–31). The study of Fbxw7 is thus important not only from the point of view of basic biology but also from the medical standpoint.

To analyze the functions of Fbxw7 *in vivo*, we and others have generated Fbxw7-deficient mice. However, *Fbxw7*^{-/-} embryos were found to die in utero at E10.5, manifesting marked abnormalities in vascular development as a result of dysregulation of Notch signaling (32, 33). To avoid this early embryonic mortality, we have established mice in which *Fbxw7* is conditionally disrupted in T cells (34) or in hematopoietic stem cells (35), and we have also examined the effects of *Fbxw7* ablation in mouse embryonic fibroblasts (36). The loss of Fbxw7 in immature T cells results in the failure of these cells to exit the cell cycle, leading to thymic hyperplasia and the subsequent development of lymphoma. Among known targets of Fbxw7, only c-Myc and Notch accumulated in the Fbxw7-deficient thymocytes, and c-Myc accumulation was found

Conflict of interest: The authors have declared that no conflict of interest exists.

Citation for this article: *J Clin Invest.* 2010;121(1):342–354. doi:10.1172/JCI40725. **123**



to be primarily responsible for the hyperproliferation phenotype. In contrast to that in immature T cells, the accumulation of c-Myc apparent in Fbxw7-null mature T cells induced expression of p53, which in turn led to cell cycle arrest and apoptosis. Furthermore, we found that Fbxw7 contributes to the long-term maintenance of hematopoietic stem cells. Most of the phenotypes of Fbxw7 deficiency in these conditional mouse mutants are related to cell proliferation or death and appear to be attributable to deregulation of c-Myc and Notch. Although Fbxw7 targets many substrates that do not participate directly in cell cycle control for degradation, the physiological roles of Fbxw7-mediated degradation of such targets have been largely unclear.

We have now examined the consequences of Fbxw7 deficiency in the liver. Unexpectedly, the major phenotypes associated with such deficiency were abnormalities in lipid metabolism and cell differentiation, which differ markedly from those in hematopoietic cell lineages and fibroblasts, in which Fbxw7 contributes primarily to the control of cell proliferation and apoptosis. We thus propose that Fbxw7 targets different groups of proteins for ubiquitin-dependent degradation and thereby contributes to distinct biological functions in a tissue-specific manner.

Results

Conditional inactivation of Fbxw7 in the liver by 2 Cre-loxP systems. We generated mice harboring floxed Fbxw7 alleles (referred to herein as Fbxw7^{F/F} mice) in which exon 5 (which encodes the F-box domain) is flanked by loxP sites (34). To ablate Fbxw7 in the liver, we crossed these Fbxw7^{F/F} mice with mice harboring a Cre transgene under the control of the promoter for the myxovirus resistance 1 (Mx1) or albumin (Alb) genes (Mx1-Cre or Alb-Cre mice). We confirmed that almost all floxed alleles were inactivated by Cre recombinase in the livers of Alb-Cre/Fbxw7^{F/F} mice as well as in those of Mx1-Cre/Fbxw7^{F/F} mice at 3 weeks after the last of 3 i.p. injections of poly(I)-poly(C) (pIpC) to activate the Mx1 gene promoter (Figure 1A). For subsequent experiments, we examined the effects of short- or long-term Fbxw7 deficiency in Mx1-Cre/Fbxw7^{F/F} mice and those of long-term a priori deficiency in Alb-Cre/Fbxw7^{F/F} mice.

Massive lipid deposition and nonalcoholic steatohepatitis-like lesions in the Fbxw7-deficient liver. Mx1-Cre/Fbxw7^{F/F} mice at 8 weeks of age were subjected to i.p. injection of pIpC every other day for 3 days to activate the Mx1 gene promoter. At 3 weeks after the last injection of pIpC, the livers of these mice were enlarged and lighter in color compared with those of control animals (Figure 1B). The liver-to-body weight ratio of these Mx1-Cre/Fbxw7^{F/F} mice was increased by approximately 30% relative to that of control mice (Figure 1C). Histological examination revealed that the nuclei of cells in the enlarged liver remained centrally located, whereas the corresponding cytoplasm was only weakly eosinophilic and contained numerous microvesicular vacuoles (Figure 1, D and E). Staining with Oil red O (Figure 1, F-I) also revealed massive lipid deposition, predominantly in the area around central veins (Figure 1G). Similar lipid deposition was also observed in the livers of Alb-Cre/Fbxw7^{F/F} mice at as early as 12 weeks of age (Figure 1, J and K). The mechanism underlying such an uneven localization of lipid deposition is unclear and awaits further investigation.

Lobular infiltration of inflammatory cells such as lymphocytes and neutrophils (Figure 1, L and M; arrowhead), as well as the presence of many ballooned hepatocytes (occasionally containing Mallory body-like eosinophilic inclusions) (Figure 1, N and O; arrow), were observed in the livers of older mutant mice at approximately

50 weeks of age. Sinusoidal fibrogenic changes in the liver as revealed by Masson's trichrome staining were also evident (Figure 1, P and Q), and serum levels of aspartate aminotransferase (AST) and alanine aminotransferase (ALT) were significantly increased (Figure 1R) in Alb-Cre/Fbxw7^{F/F} mice at 50 weeks of age. The serum level of bilirubin tended to be higher in the mutant animals than in age-matched controls, suggestive of the destruction of liver tissue in the mutant mice, although this difference was not statistically significant (Supplemental Figure 1; supplemental material available online with this article; doi:10.1172/JCI40725DS1). The onset of inflammatory changes occurred later than that of steatosis, but feeding Mx1-Cre/Fbxw7^{F/F} mice a methionine- and choline-deficient (MCD) diet resulted in acceleration of inflammation (Figure 2A). The extents of steatosis and hepatitis were less pronounced in Alb-Cre/Fbxw7^{F/F} mice than in Mx1-Cre/Fbxw7^{F/F} mice subjected to acute ablation of Fbxw7, probably as a result of compensatory mechanisms operative during development in the former animals. However, massive steatosis and inflammation were also apparent in Alb-Cre/Fbxw7^{F/F} mice fed the MCD diet, whereas control animals did not show such marked changes (Figure 2B). These results suggested that Alb-Cre/Fbxw7^{F/F} mice are also more sensitive to steatohepatitis than are controls. The histological findings in both types of Fbxw7-deficient mice are highly similar to those associated with nonalcoholic steatohepatitis (NASH) in humans (37).

Expression of adipogenic and lipogenic genes in the Fbxw7-deficient liver. We next determined lipid concentrations in liver extracts. Triglyceride levels were significantly increased in the livers of Mx1-Cre/Fbxw7^{F/F} mice compared with those in control animals at 3 weeks after the final pIpC injection, whereas the concentration of total cholesterol was not affected in the mutant livers (Figure 3A). Given that triglyceride synthesis is regulated predominantly by transcriptional activators, such as SREBPs, carbohydrate response element-binding protein (ChREBP), and PPAR- γ , we examined the expression of these proteins and their downstream targets in the liver. Immunoblot analysis revealed that the abundance of nuclear SREBP1, which is the major SREBP in the liver and a target of Fbxw7-mediated proteolysis (16, 17), was increased both in pIpC-injected Mx1-Cre/Fbxw7^{F/F} mice and in Alb-Cre/Fbxw7^{F/F} mice (Figure 3B and Supplemental Figure 2). The intensity of the more slowly migrating band, likely corresponding to the phosphorylated form of SREBP1, was especially increased, consistent with the previous observation that the phosphorylated forms of SREBPs are targeted by Fbxw7 (16-18), as is generally the case for Fbxw7 substrates (11, 34). In contrast, the amounts of ChREBP and PPAR- γ were decreased in the mutant mice compared with those in control animals, suggestive of the operation of a negative feedback loop triggered by triglyceride accumulation. Consistent with this notion, the abundance of Pparg mRNA in the liver was increased in SREBP cleavage-activating protein-deficient mice, in which the SREBP pathway is inactivated (38). The levels of PGC-1 α and mTOR (total or phosphorylated forms) were unaffected by hepatic deletion of Fbxw7.

RT and real-time PCR analysis revealed that the abundance of mRNAs for the adipogenic and lipogenic transcriptional activators SREBP1c, ChREBP, and Pparg was decreased in the Fbxw7-deficient liver (Figure 3C), suggesting that the transcription of these genes is suppressed by a negative feedback loop triggered by the high level of triglyceride. At the protein level, the precursor form of SREBP1 was reduced, probably as a result of the decrease in the abundance of its mRNA, whereas the mature cleaved form was increased (Supplemental Figure 3). Among the downstream targets of SREBPs,





Figure 1

Development of NASH-like liver disease as a result of *Fbxw7* deletion. (A) Genomic PCR analysis from the mouse liver of the indicated genotypes. The positions of amplified fragments corresponding to WT, floxed, and exon 5–deleted ($\Delta E5$) alleles are indicated. (B) Gross appearance of the livers of indicated genotypes treated as in A. Scale bar: 10 mm. (C) Liver/body weight ratio of mice treated as in A. Data are mean \pm SD from 5 animals of each genotype. $***P < 0.005$. (D and E) H&E staining of liver sections from *Mx1-Cre/Fbxw7^{+F/F}* (+/ Δ) and *Mx1-Cre/Fbxw7^{F/F}* (Δ/Δ) mice, respectively, treated as in A. (F and G) Oil red O staining of liver sections treated as in A. (H and I) Higher-magnification views of images in F and G, respectively. (J and K) Oil red O staining of liver sections from *Fbxw7^{F/F}* (F/F) and *Alb-Cre/Fbxw7^{F/F}* (Δ/Δ) mice, respectively, at 12 weeks of age. (L–O) H&E staining of liver sections from *Mx1-Cre/Fbxw7^{+F/F}* (+/ Δ) (L and N) and *Mx1-Cre/Fbxw7^{F/F}* (Δ/Δ) (M and O) mice at 50 weeks after the final injection of plpC. Lobular infiltration of inflammatory cells is indicated by the arrowhead, and Mallory body–like eosinophilic inclusion is indicated by the arrow. (P and Q) Masson’s trichrome staining of liver sections from *Fbxw7^{F/F}* (F/F) and *Alb-Cre/Fbxw7^{F/F}* (Δ/Δ) mice, respectively, at 50 weeks of age. Scale bar: 50 μ m (D, E, H, I, and L–O); 100 μ m (F, G, J, K, P, and Q). (R) Serum AST and ALT activities in *Fbxw7^{F/F}* ($n = 6$) and *Alb-Cre/Fbxw7^{F/F}* ($n = 10$) mice at 50 weeks of age. Data are mean \pm SD. $*P < 0.05$.

**Figure 2**

Increased susceptibility to a NASH-like condition conferred by *Fbxw7* ablation in the liver. (A) *Mx1-Cre/Fbxw7^{+/F}* (control) and *Mx1-Cre/Fbxw7^{F/F}* mice were injected with pIpC at 8 weeks of age and then fed an MCD diet for 2 weeks. Liver sections were then subjected to H&E staining. Lower- and higher-magnification views are shown (top and bottom panels, respectively). In addition to fatty degeneration, many foci of lobular infiltration of inflammatory cells (arrowheads) were apparent in the livers of *Mx1-Cre/Fbxw7^{F/F}* mice. Scale bar: 50 μm . (B) *Alb-Cre/Fbxw7^{+/F}* (control) and *Alb-Cre/Fbxw7^{F/F}* mice at 12 weeks of age were fed an MCD diet for 4 weeks and then analyzed as in A. Lower- and higher-magnification views are shown (top and bottom panels, respectively). Control mice developed a small extent of fatty degeneration, whereas *Alb-Cre/Fbxw7^{F/F}* mice showed massive accumulation of lipid droplets and many foci of lobular infiltration of inflammatory cells (arrowheads) similar to those apparent in *Mx1-Cre/Fbxw7^{F/F}* mice. Scale bar: 50 μm .

the amounts of mRNAs for fatty acid synthase (*Fas*) and stearoyl-CoA desaturase-1 (*Scd1*) were increased, whereas those for the LDL receptor (*Ldlr*) and HMG-CoA synthase (*Hmgcs1*) were decreased, in the mutant liver (Figure 3C). Immunostaining also showed that SREBP1 accumulated in the region around the central veins (Figure 3D), corresponding to the area of lipid deposition, even though deletion of *Fbxw7* appears to occur throughout almost the entire liver. The expression of SCD-1 was also increased in the region around the central veins in which SREBP1 was upregulated (Figure 3E). Collectively, these results suggested that the accumulation of SREBP proteins as a result of *Fbxw7* ablation results in triglyceride deposition in the liver, which in turn affects the expression of other adipogenic and lipogenic genes as well as their downstream targets via a negative feedback loop.

Increased proliferation of *Fbxw7*-deficient hepatocytes. We compared the abundance of cyclin E and c-Myc between the livers of *Mx1-Cre/Fbxw7^{+/F}* and *Mx1-Cre/Fbxw7^{F/F}* mice at 3 or 50 weeks after the final pIpC injection, beginning at 8 weeks of age. Immunoblot analysis revealed that the amount of cyclin E in the livers of *Mx1-Cre/Fbxw7^{F/F}* mice was increased compared with that in *Mx1-Cre/Fbxw7^{+/F}* mice at 3 weeks after pIpC injection but not at 50 weeks (Figure 4A). The abundance of c-Myc was not affected by the loss of *Fbxw7* in the liver, at either 3 or 50 weeks after injection. To measure the rate of hepatocyte proliferation, we subjected *Mx1-Cre/Fbxw7^{+/F}* and *Mx1-Cre/Fbxw7^{F/F}* mice to i.p. injection with BrdU for 3 consecutive days, beginning at 3 or 50 weeks after the final pIpC injection. Immunostaining of the liver with antibodies

to BrdU at 1 day after the last BrdU injection revealed that the rate of BrdU incorporation was markedly increased in *Fbxw7*-deficient liver cells compared with that in control cells (Figure 4B). Most of the BrdU-positive cells were also reactive with antibodies to albumin but not with those to cytokeratin 19 (CK19) at 3 weeks after pIpC injection (Figure 4C), suggesting that the proliferating cells are predominantly hepatocytes. In contrast, at 50 weeks after pIpC injection, both hepatocytes and cholangiocytes in *Mx1-Cre/Fbxw7^{F/F}* mice incorporated BrdU to a greater extent than did those in control mice. The TUNEL assay revealed that the frequency of apoptosis was also increased in the *Fbxw7*-deficient liver at 3 weeks after pIpC injection (Figure 4, D and E), suggesting that the loss of *Fbxw7* transiently promotes cell cycle progression but eventually results in apoptosis in the liver, as it does in T lymphocytes (34).

Development of hamartomas with hyperproliferation of the biliary system in the *Fbxw7*-deficient liver. We next examined in more detail the long-term effects of *Fbxw7* loss in the liver. Macroscopic examination of *Mx1-Cre/Fbxw7^{F/F}* mice at 50 weeks after pIpC injection at 8 weeks of age revealed that the mutant liver was enlarged and darker in color compared with the control liver and possessed a rough surface as a result of the presence of several nodules (Figure 5A). We confirmed that the *Fbxw7* gene was deleted in such nodules (Figure 5B), which were grossly demarcated and readily excised from the liver. Histological examination revealed structural abnormalities characterized by marked dilation of intrahepatic bile ducts as well as apparent proliferation of the biliary system




Figure 3

Deposition of triglyceride and accumulation of SREBP1 in the *Fbxw7*-deficient liver. **(A)** Triglyceride and total cholesterol concentrations in the livers of *Mx1-Cre/Fbxw7^{F/F}* (control) and *Mx1-Cre/Fbxw7^{F/F}* mice at 3 weeks after the final injection of plpC, beginning at 8 weeks of age. Data are mean ± SD from 3 mice of each genotype. **P* < 0.05. **(B)** Protein extracts of the livers of *Mx1-Cre/Fbxw7^{F/F}* (control) and *Mx1-Cre/Fbxw7^{F/F}* mice at 3 weeks after the final injection of plpC, beginning at 8 weeks of age, were subjected to IB analysis with antibodies to the indicated proteins (left panel). Liver extracts of *Fbxw7^{F/F}* (control) and *Alb-Cre/Fbxw7^{F/F}* mice at 12 weeks of age were similarly analyzed (right panel). Three animals were examined for each genotype. Hsp90 was analyzed as a loading control. p-mTOR, phosphorylated mTOR. **(C)** RT and real-time PCR analysis of the indicated mRNAs in the livers of *Mx1-Cre/Fbxw7^{F/F}* (control) and *Mx1-Cre/Fbxw7^{F/F}* mice treated as in **A**. Normalized data are expressed relative to the corresponding value for control mice and are mean ± SD from 3 independent experiments. **P* < 0.05. **(D and E)** Liver sections of *Mx1-Cre/Fbxw7^{F/F}* (+Δ) and *Mx1-Cre/Fbxw7^{F/F}* (Δ/Δ) mice, treated as in **A**, were subjected to immunofluorescence staining with antibodies **(D)** to SREBP1 and **(E)** to SCD-1. CV, central vein. Scale bar: 100 µm.

(Figure 5, C–F); these abnormalities were pathologically diagnosed as hamartomas. Such lesions were also observed, albeit to a lesser extent, in *Alb-Cre/Fbxw7^{F/F}* mice at as early as 12 weeks of age (Figure 5, G and H). Hamartomas, which are reactive with antibodies to CK19 (Figure 5, I and J), developed in all mutant mice of both genotypes examined (*n* = 14). These results suggested that the loss of *Fbxw7* may promote proliferation of the biliary system and shift the development of hepatic stem cells toward the cholangiocyte lineage rather than the hepatocyte lineage.

We examined the abundance of mRNAs for *Alb* (Figure 5K) and *CK19* (Figure 5L) as markers of hepatocyte and cholangiocyte lineages, respectively. The amount of *CK19* mRNA in the liver was increased

as early as 2 weeks after pIpC injection in *Mx1-Cre/Fbxw7^{F/F}* mice and showed a more than 40-fold increase at 50 weeks after *Fbxw7* deletion. In contrast, the abundance of *Alb* mRNA in the mutant liver at 50 weeks after pIpC injection was decreased by 40% compared with that in control liver. These changes in differentiation markers were thus consistent with a marked proliferation of the biliary system in the *Fbxw7*-deficient liver.

Skewed hepatic differentiation induced by Notch1 accumulation in the Fbxw7-deficient liver. The hepatic cell fate decision is thought to be largely dependent on Notch signaling (39–44). We therefore examined the expression of Notch, a target of *Fbxw7*, in the *Fbxw7*-deficient liver. Although immunoblot analysis did not reveal an




Figure 4

Increased proliferation and apoptosis of Fbxw7-deficient hepatocytes. (A) IB analysis of cyclin E, c-Myc, and Hsp90 (loading control) in liver extracts from *Mx1-Cre/Fbxw7^{+/F}* and *Mx1-Cre/Fbxw7^{F/F}* mice at 3 or 50 weeks after *Fbxw7* deletion by plpC injection, beginning at 8 weeks of age. (B) Representative immunostaining for BrdU in liver sections from *Mx1-Cre/Fbxw7^{+/F}* (+/Δ) and *Mx1-Cre/Fbxw7^{F/F}* (Δ/Δ) mice injected with BrdU on 3 consecutive days, beginning 3 weeks after the final plpC injection as in A. Arrowheads indicate BrdU-positive nuclei. Scale bar: 50 μm. (C) The proportion of BrdU-positive hepatocytes or cholangiocytes was determined from immunostaining for BrdU in combination with that for albumin or CK19 in the livers of *Mx1-Cre/Fbxw7^{+/F}* (control) or *Mx1-Cre/Fbxw7^{F/F}* mice at 3 or 50 weeks after deletion of *Fbxw7* as in A. Data are mean ± SD from 10 fields from 3 mice of each genotype. **P* < 0.05. (D) Representative TUNEL staining for liver sections of *Mx1-Cre/Fbxw7^{+/F}* (+/Δ) and *Mx1-Cre/Fbxw7^{F/F}* (Δ/Δ) mice 3 weeks after the final plpC injection as in A. Arrowheads indicate TUNEL-positive cells. Scale bar: 50 μm. (E) The proportion of TUNEL-positive liver cells was determined from images similar to those in D. Data are mean ± SD from 3 animals of each genotype. ****P* < 0.005.

increase in the abundance of any of the isoforms of Notch in the liver of *Mx1-Cre/Fbxw7^{F/F}* mice at either 3 or 50 weeks after *Fbxw7* deletion (data not shown), confocal microscopic analysis revealed that the intracellular domain of Notch1 was highly concentrated in both the cytoplasm and the nucleus of Fbxw7-deficient liver at 3 weeks after pIpC injection (Figure 6A). Consistent with the observed upregulation of Notch1, the abundance of Notch1 target genes, including those for *Hes1* and *Hey1*, was simultaneously

increased in the livers of *Mx1-Cre/Fbxw7^{F/F}* mice (Figure 6B and Supplemental Figure 5A, respectively). At 15 weeks after plpC injection, Notch1 accumulated in the hepatocyte-like cells residing around the portal area, and these cells were reactive to antibodies to CK7 (Figure 6C), another marker of cholangiocytes, suggesting that such cells might be in the process of transdifferentiation to the cholangiocytes by Notch activation. At 50 weeks after plpC injection, Notch1 was observed in the form of aggregates in the




Figure 5

Hamartoma development as a result of long-term ablation of *Fbxw7* in the liver. (A) Gross appearance of the livers of *Mx1-Cre/Fbxw7^{F/F}* and *Mx1-Cre/Fbxw7^{F/F}* mice at 50 weeks after the final injection of plpC, beginning at 8 weeks of age. Scale bar: 10 mm. (B) PCR analysis of genomic DNA from the dilated bile ducts excised from hamartomas in the livers of 2 *Mx1-Cre/Fbxw7^{F/F}* mice. Genomic DNA from control mice was also analyzed. (C–F) H&E staining of liver sections from a *Mx1-Cre/Fbxw7^{F/F}* (+/Δ) mouse (C) and from a *Mx1-Cre/Fbxw7^{F/F}* (Δ/Δ) mouse (D–F) that developed hamartoma after *Fbxw7* deletion as in A. (G and H) H&E staining of liver sections from *Fbxw7^{F/F}* (F/F) and *Alb-Cre/Fbxw7^{F/F}* (Δ/Δ) mice, respectively, at 12 weeks of age. Arrowheads indicate malformation of the ductal plate. Scale bar: 50 μm (F); 100 μm (E, G, and H); 200 μm (C and D). (I and J) Immunofluorescence staining for CK19 in the livers of *Mx1-Cre/Fbxw7^{F/F}* (+/Δ) and *Mx1-Cre/Fbxw7^{F/F}* (Δ/Δ) mice at 50 weeks after the final injection of plpC, beginning at 8 weeks of age. The dashed line indicates the outer boundary of portal vein. PV, portal vein; BD, bile duct. Scale bar: 25 μm. (K and L) RT and real-time PCR analysis of *Alb* and *CK19* mRNAs, respectively, in the livers of *Mx1-Cre/Fbxw7^{F/F}* (control) and *Mx1-Cre/Fbxw7^{F/F}* mice at 1, 2, or 50 weeks after deletion of *Fbxw7* as in A. Normalized data are expressed relative to the corresponding value for control mice. Data are mean ± SEM from 3 independent experiments. **P* < 0.05.



cytoplasm or the nucleus (Supplemental Figure 4A). The increase in *Hes1* or *Hey1* was not detected by immunostaining analysis at this period (Supplemental Figure 4B and Supplemental Figure 5B), but the abundance of mRNAs for *Hes1*, *Hey1*, and *Hey2* was increased in the livers of *Mx1-Cre/Fbxw7^{F/F}* mice at 50 weeks after pIpC injection (Figure 6D). Neither *Notch2*, mutations in the gene in which mutations result in Alagille disease, nor *Notch3* or *Notch4*, the expression of both of which is increased in hepatocellular carcinoma, were detected by immunofluorescence analysis in the livers of either control or *Mx1-Cre/Fbxw7^{F/F}* mice (data not shown). However, neither the expression of Notch ligands, such as *Dll-1* and *Jagged-1*, nor that of the Notch cofactor RBP-J in the liver appeared to be affected by the loss of *Fbxw7* (Supplemental Figure 6). We also examined the expression of *TSC1* and *TSC2*, given that the loss of function of either *TSC1* or *TSC2* is known to result in the development of hamartoma in humans. However, no difference in expression of *TSC1* or *TSC2* was found between *Fbxw7*-deficient and control mice (Supplemental Figure 6).

To investigate whether the skewed developmental orientation toward the cholangiocyte lineage apparent in the *Fbxw7*-deficient liver is dependent on *Notch1* accumulation, we examined the differentiation of hepatic stem cells in culture (45). A fraction containing hepatic stem cells was prepared from the livers of *Fbxw7^{+/+}* and *Fbxw7^{F/F}* embryos and was then infected with a retrovirus encoding Cre recombinase or with the empty virus alone to generate *Fbxw7^{+/+}*, *Fbxw7^{+Δ}*, *Fbxw7^{F/F}*, and *Fbxw7^{ΔΔ}* cells in the presence of HGF and EGF. Immunofluorescence analysis revealed that most of the *Fbxw7^{+/+}*, *Fbxw7^{+Δ}*, and *Fbxw7^{F/F}* cells differentiated into the hepatocyte lineage, characterized by albumin expression, with only a small subset of cells differentiating into the cholangiocyte lineage (Figure 7A). In contrast, the percentage of *Fbxw7^{ΔΔ}* cells that differentiated into the cholangiocyte lineage, characterized by expression of *CK7*, was markedly increased compared with that for cells of the control genotypes. To confirm these results in a quantitative manner, we performed RT and real-time PCR analysis of *Alb* and *CK19* mRNAs. Consistent with the immunofluorescence data, the amount of *CK19* mRNA was significantly increased in *Fbxw7^{ΔΔ}* cells compared with that in *Fbxw7^{+Δ}* cells, whereas the abundance of *Alb* mRNA did not differ between the 2 genotypes (Figure 7B).

Notch signaling is implicated in the differentiation of liver stem cells into the cholangiocyte lineage. Indeed, immunofluorescence analysis revealed that *Notch1* accumulated in *Fbxw7^{ΔΔ}* cells to a greater extent than in *Fbxw7^{+Δ}* cells (Supplemental Figure 7). We therefore examined whether additional ablation of the *Notch* cofactor RBP-J might correct the abnormal development of *Fbxw7*-deficient liver stem cells. We generated *Fbxw7*-deficient hepatic stem cells with additional deletion of either *Rbpj* or *Myc* genes and examined the level of *CK19* mRNA. The abundance of *CK19* mRNA was increased in *Fbxw7^{ΔΔ}Myc^{ΔΔ}* cells but not in *Fbxw7^{ΔΔ}Rbpj^{ΔΔ}* cells (Figure 7B). These results indicate that the skewed developmental orientation of hepatic stem cells to the cholangiocyte lineage is dependent on *Notch1* accumulation induced by the loss of *Fbxw7*.

Discussion

Given that the substrates of *Fbxw7* include key proteins that contribute to diverse biological processes, including the cell cycle, cell differentiation, and apoptosis, and that the binding of *Fbxw7* to its substrates depends on their phosphorylation, the function

of this protein is likely complex. Although much attention has focused on the relation between the accumulation of cyclin E due to loss of *Fbxw7* function and tumorigenesis, *Notch* degradation by *Fbxw7* is critical during embryogenesis, suggesting that *Fbxw7* functions in development- and tissue-dependent manners. To provide insight into the physiological and pathological relevance of *Fbxw7*, we have induced conditional inactivation of *Fbxw7* in several mouse tissues. Our previous studies have shown that ablation of *Fbxw7* in hematopoietic cells or fibroblasts results in abnormalities that are mainly related to the cell cycle and apoptosis. We now show that liver-specific ablation of *Fbxw7* induced fatty liver and abnormal cell differentiation, likely as a result of the accumulation of SREBPs and *Notch1*, respectively, as well as promoted cell proliferation (Figure 8).

We generated 2 types of mice with liver-specific deficiency of *Fbxw7* with the use of the *Mx1* or *Alb* gene promoters to drive Cre expression. The phenotypes of *Alb-Cre/Fbxw7^{F/F}* mice are milder than those induced by acute ablation of *Fbxw7* in *Mx1-Cre/Fbxw7^{F/F}* mice, probably because of the operation of compensatory mechanisms during development in the former animals. In *Mx1-Cre/Fbxw7^{F/F}* mice, it would be expected for *Fbxw7* to be deleted in cells and tissues other than the liver, such as hematopoietic cells. To exclude the possibility that ablation of *Fbxw7* in hematopoietic cell lineages might be responsible for steatohepatitis, we have generated *Lck-Cre/Fbxw7^{F/F}* and *CD4-Cre/Fbxw7^{F/F}* mice (in both of which *Fbxw7* deletion occurs in T cells), *CD19-Cre/Fbxw7^{F/F}* mice (*Fbxw7* deletion occurs in B cells), and *LysM-Cre/Fbxw7^{F/F}* mice (*Fbxw7* deletion occurs in myeloid cells). None of these animals showed either fatty liver or hepatic inflammation (data not shown). Furthermore, *Alb-Cre/Fbxw7^{F/F}* mice manifested pronounced hepatic infiltration of inflammatory cells when they were fed an MCD diet, confirming that the steatohepatitis induced by *Fbxw7* deletion is attributable to an effect that is intrinsic to the liver.

Nonalcoholic fatty liver disease (NAFLD) is a growing health concern, due to its rapidly increasing prevalence worldwide. NASH is a progressive form of NAFLD that has the potential to develop into hepatocellular carcinoma. We now show that mice with liver-specific ablation of *Fbxw7* developed clinicopathologic features similar to those of NAFLD or NASH in humans, including triglyceride deposition around central veins, pericellular fibrosis, infiltration of inflammatory mononuclear cells, and the appearance of Mallory bodies in the liver as well as increases in the serum levels of ALT and AST. However, these animals were not found to develop hepatocellular carcinoma. Genetic mouse models for human NASH have been established by functional deletion of leptin (46) or its receptor (47), phosphatase and tensin homolog (PTEN) (48), NEMO (also known as IKK- γ) (49), interleukin-1 receptor α (50), galectin-3 (51), or retinoic acid receptor α (52). Mice transgenic for SREBP1c also manifest pronounced NASH (53). SREBP1c is degraded in an *Fbxw7*-dependent manner (16), and we have now shown that it accumulated in the *Fbxw7*-deficient liver. These findings thus suggest that an *Fbxw7*-SREBP1 axis plays a key physiological role in the regulation of lipid metabolism in the liver as well as a pathological role in the development of NASH.

Whereas steatosis develops in the acute phase of liver-specific *Fbxw7* deficiency, hamartoma develops in the chronic phase. *Fbxw7* targets mTOR for degradation (19). The TSC complex, consisting of *TSC1* (hamartin) and *TSC2* (tuberin), is the major negative regulator of mTOR, and its genetic loss results in mTOR






Figure 6

Accumulation of Notch1 and activation of its target genes in the Fbxw7-deficient liver. (**A** and **B**) Representative immunostaining for the intracellular domain of (**A**) Notch1 and for (**B**) Hes1 in liver sections from *Mx1-Cre/Fbxw7^{+F}* (*+/\Delta*) and *Mx1-Cre/Fbxw7^{F/F}* (Δ/Δ) mice at 3 weeks after *Fbxw7* deletion by plpC injection, beginning at 8 weeks of age. Arrowheads indicate accumulating (**A**) Notch1 intracellular domain and (**B**) Hes1 in the nucleus. (**C**) Immunofluorescence staining for the intracellular domain of Notch1 and for CK7 in the livers of *Mx1-Cre/Fbxw7^{+F}* (*+/\Delta*) and *Mx1-Cre/Fbxw7^{F/F}* (Δ/Δ) mice at 15 weeks after the final injection of plpC, beginning at 8 weeks of age. Intense Notch1 staining was detected in the Fbxw7-deficient liver, and most of the Notch1-positive cells express CK7 (arrowheads). Scale bar: 10 μ m (**A**); 50 μ m (**B** and **C**). (**D**) RT and real-time PCR analysis of Notch target genes in the livers of *Mx1-Cre/Fbxw7^{+F}* (control) and *Mx1-Cre/Fbxw7^{F/F}* mice at 50 weeks after *Fbxw7* deletion. Normalized data for *Hes1*, *Hey1*, and *Hey2* mRNAs are expressed relative to the corresponding value for control mice and are mean \pm SD from 3 independent experiments. **P* < 0.05.

**Figure 7**

Skewed differentiation of hepatic stem cells from *Fbxw7*-deficient mice. (A) Primary cultured liver cells of the indicated genotypes were subjected to immunofluorescence staining with antibodies to albumin or to CK7. Clusters of cells expressing CK7 were apparent only with *Fbxw7*^{Δ/Δ} cells. Scale bar: 50 μm. (B) RT and real-time PCR analysis of *Alb* and *CK19* mRNAs in primary cultured liver cells of the indicated genotypes. Normalized data for *Alb* and *CK19* mRNAs are expressed relative to the corresponding value for *Fbxw7*^{+/Δ} cells. Data are mean ± SD from 3 independent experiments. ****P* < 0.005.

activation and development of hamartoma in humans (54, 55). However, the abundance of mTOR or TSC1/2 was not altered in the *Fbxw7*-deficient livers of mice, suggesting that the accumulation of mTOR or the loss of TSC1/2 is not responsible for hamartoma development in these animals. Microscopic examination revealed over proliferation of the biliary system in the hamartomas, suggesting that deregulated differentiation of liver stem cells into the cholangiocyte lineage might be largely responsible for hamartoma development. Liver stem cells are able to differentiate into either the hepatocyte or cholangiocyte lineages, with the Notch signaling pathway having been implicated in regulation of the cell fate decision by skewing differentiation toward the cholangiocyte lineage (41). We have now shown that both Notch1 and its target genes were overexpressed in the *Fbxw7*-deficient livers of mice and that the abnormal cell differentiation induced by *Fbxw7* loss was corrected by the additional loss of the Notch cofactor RBP-J. These results suggest that Notch1 accumulation as a result of *Fbxw7* loss is primarily responsible for the abnormal cell differentiation in the *Fbxw7*-deficient mouse liver. Although the origin of hamartomas as well as the mechanism of their development in the *Fbxw7*-deficient liver are currently unclear, transient activation of Notch proteins as a result of *Fbxw7* loss may lead to a shift in cell differentiation from hepatocytes to cholangiocytes, and the generation of such abnormally differentiated cells might confer a predisposition to hamartoma development that is realized if the cells undergo an additional gene mutation. Mice lacking both *Foxa1* and *Foxa2* were recently shown to display a similar liver phenotype (hyperplasia of the biliary tree) (56). However, neither

differentiation of hepatocytes nor Notch signaling were affected in *Foxa1/2*-deficient mice, whereas hyperactivation of Notch signaling seems to be attributable to the bile duct hamartoma in *Fbxw7*-deficient mice. Furthermore, proliferation of relatively small and uniform bile ducts is prominent in *Foxa1/2*-deficient mice, whereas the abnormal bile ducts in *Fbxw7*-deficient mice are large and heterogeneous in size. We therefore concluded that the mechanism underlying the development of proliferative bile ducts is likely different between these mutant mice.

Although *Fbxw7* had been thought to function primarily in cell cycle control by regulating cyclin E, c-Myc, Notch, and c-Jun, the recent identification of additional substrates has suggested new cell cycle-independent roles for *Fbxw7*. We now provide genetic evidence that the major substrates of *Fbxw7* in the liver are SREBP1 and are responsible for liver steatosis and hamartoma development, respectively. These results contrast with our previous observations that deletion of *Fbxw7* in the hematopoietic system and fibroblasts results primarily in deregulation of the cell cycle or of apoptosis due to activation of the p53-dependent checkpoint (34–36). Why does the function of *Fbxw7* differ in different tissues? We propose that the biological relevance of *Fbxw7* is determined by 3 factors: (a) the expression of *Fbxw7*; (b) the expression and activation of protein kinases that phosphorylate the Cdc4 phosphodegron, an amino acid sequence that is recognized by *Fbxw7*; and (c) the expression of substrate molecules. The combination of these 3 factors may define the role of *Fbxw7* in a tissue-specific manner, with the different phenotypes associated with *Fbxw7* deficiency




Figure 8

A model for Fbxw7 functions in vivo. Fbxw7 mediates ubiquitin-dependent degradation of substrates in different functional categories. For example, Fbxw7 controls cell proliferation by targeting c-Myc, cyclin E, and c-Jun for degradation. However, it also regulates lipid metabolism and cell differentiation by targeting SREBP and Notch proteins, respectively. Major phenotypes associated with Fbxw7 deficiency in different tissues are shown in red. Ub, ubiquitin.

being attributable to different expression patterns of Fbxw7, its substrates, and kinases that phosphorylate each substrate.

Methods

Generation of conditional knockout mice. Mice homozygous for the floxed *Fbxw7* allele (*Fbxw7^{FL/FL}* mice) (34) were crossed with *Mx1-Cre* transgenic mice (57) provided by K. Rajewsky (Harvard Medical School, Boston, Massachusetts, USA) or *Alb-Cre* transgenic mice (58) purchased from The Jackson Laboratory. Expression of Cre recombinase in the resulting offspring of the former cross was induced by i.p. injection of 500 µg pIpC (GE Healthcare Biosciences) on 3 alternate days. Deletion of exon 5 of the floxed *Fbxw7* allele was confirmed by PCR analysis of genomic DNA as previously described (34). *Fbxw7^{FL/FL}* mice were also crossed with *Rbpj^{FL/FL}* mice (59) provided by T. Honjo (Kyoto University, Kyoto, Japan) or *Myc^{FL/FL}* mice (60) provided by I.M. de Alborán (National Center for Biotechnology, Madrid, Spain). The experimental protocols were approved by the Institutional Animal Care and Use Committee of Kyushu University.

Histological and biochemical analysis. Liver tissue was fixed with 4% paraformaldehyde in PBS, embedded in paraffin, and stained with H&E or Masson’s trichrome solution. Some sections were stained with Oil red O (Nakalai Tesque) according to standard procedures, in order to examine the extent of lipid accumulation in hepatocytes. Serum levels of AST and ALT were measured with a standard clinical autoanalyzer.

Dietary model of NASH. Mice were fed with an MCD diet (Funabashi Farm) for the indicated periods (see the legend for Figure 2) and analyzed.

Measurement of triglyceride and total cholesterol levels in the liver. Frozen liver tissue was homogenized, and triglyceride and total cholesterol were extracted from the homogenate with chloroform/methanol (2:1, vol/vol) 3362.

dried, and resuspended in 2-propanol. The amounts of triglyceride and total cholesterol in the extract were measured with the use of Lipidos liquid and Cholesterol liquid kits (Toyobo), respectively.

Immunoblot analysis. Total protein extracts were prepared from liver with RIPA buffer. The extracts (30 µg) were subjected to immunoblot analysis as described previously (61) with antibodies to cyclin E (M-20), to c-Myc (N-262), to ChREBP (P-13), or to PPAR-γ (E-8), all of which were obtained from Santa Cruz Biotechnology Inc.; with antibodies to Ser²⁴⁴⁸-phosphorylated or total (7C10) forms of mTOR (Cell Signaling Technology); with antibodies to SREBP1 (2A4, NeoMarkers); or with antibodies to PGC-1α (Chemicon). As a control, each membrane was stripped and then probed with antibodies to Hsp90 (BD Transduction Laboratories).

RT and real-time PCR analysis. Total RNA was extracted from liver using the guanidinium thiocyanate-phenol-chloroform method, purified, and subjected (1 µg) to RT with random hexanucleotide primers (ReverTra Ace α, Toyobo). The resulting cDNA was subjected to real-time PCR in a reaction mixture that contained 1× SYBR Green PCR Master Mix (Applied Biosystems) and 200 nM of gene-specific primers. Assays were performed in triplicate with an ABI Prism 7700 Sequence Detector (Applied Biosystems). The PCR protocol comprised 40 cycles of incubation at 60°C for 30 seconds and 95°C for 5 seconds. The sequences of the PCR primers (sense and antisense, respectively) were 5'-TGCTCCCAGCTGCAGGC-3' and 5'-GCCCGGTAGCTCTGGGTGTA-3' for *Fas*, 5'-TGGGTTGGCTGCTTGTG-3' and 5'-GCGTGGCAGGATGAAG-3' for *Scd1*, 5'-CTGCCGACCTGATGAATTCC-3' and 5'-TAGGGCCATCACACTGTGTC-3' for *Ldlr*, 5'-GCTCTCCATACAGTGCTACC-3' and 5'-GAGTGAAA-GATCATGAAGCC-3' for *Hmgs1*, 5'-AGAGATGCCATCTCCAGCCTC-3' and 5'-CTTGGTCTTAGGGTCTTCAGG-3' for *ChREBP*, 5'-CTGTGAAGTTCAAT-GCACTGGAA-3' and 5'-CCTCGATGGGCTTCACGTT-3' for *Pparg*, 5'-CAT-GGATTGCACATTTGAAG-3' and 5'-CCTGTGTCCCCTGTCTCA-3' for *SREBP1c*, 5'-TCCTGTGCTGCAGCCTTTCTCA-3' and 5'-CCAGGTTCCCA-CAAAGGCATCA-3' for fatty acid-binding protein 4, 5'-GTCCTACA-GATTGACAATGC-3' and 5'-CACGCTCTGGATCTGTGACAG-3' for *CK19*, 5'-CATGACACCATGCCTGTGAT-3' and 5'-CTCTGATCTTCAG-GAAGTGATAC-3' for *Alb*, 5'-CATTCCAAGCTAGAGAAGCAAG-3' and 5'-TATTTCCCCAACACGCTCG-3' for *Hes1*, 5'-AAAATGCTGCACACTG-CAGG-3' and 5'-CGAGTCCTTCAATGATGCTCAG-3' for *Hey1*, 5'-AAAC-GACCTCCGAAAGCGA-3' and 5'-CGGTGAATTGGACCTCATCACT-3' for *Hey2*, and 5'-GGAACATAGCCGTAAACTGC-3' and 5'-TCACTGTGCCT-GAACTTACC-3' for β-tubulin. Reactions for β-tubulin mRNA were performed concurrently on the same plate as those for the test mRNAs, and results were normalized by the corresponding amount of β-tubulin mRNA.

BrdU incorporation in vivo. Mice were injected with BrdU (1 mg, i.p.) on 3 consecutive days. The liver was removed 24 hours after the third injection of BrdU, and BrdU incorporation was examined with an In Situ BrdU Detection Kit (BD Biosciences). BrdU-positive cells were counted in 10 different fields at high (×400) magnification, and the percentage of BrdU-positive cells was calculated.

Immunofluorescence microscopy. Liver tissue was fixed with 4% paraformaldehyde in PBS and sectioned at a thickness of 40 µm with a vibratome. Sections were then immunostained with antibodies to the intracellular domain of Notch1 or to SCD-1 (both from Cell Signaling Technology), to Hes1 (AB5702, Millipore), to SREBP1 (2A4, NeoMarkers), to albumin (Biogenesis), to CK19 (45), or to CK7 (MAB3226, Chemicon). Immune complexes were detected with Alexa Fluor 488- or Alexa Fluor 546-conjugated goat antibodies to mouse or rabbit IgG (Invitrogen). Cultured liver cells were also subjected to immunostaining, as described previously (45), with the antibodies to albumin and to CK7. For confocal microscopic analysis, we used Zeiss LSM 510 META Confocal Microscope (Carl Zeiss MicroImaging).

TUNEL assay. The TUNEL assay was performed as described previously (62). In brief, paraffin-embedded sections of liver were treated with H₂O₂,



permeabilized for 15 minutes at 37°C with proteinase K (20 µg/ml, Sigma-Aldrich), and then incubated for 1 hour at 37°C with a reaction mixture containing terminal deoxynucleotidyl transferase (Invitrogen) and biotinylated dUTP (Boehringer Ingelheim). Labeled DNA was visualized with an ABC Kit (Vector Laboratories) and diaminobenzidine.

Primary culture of fetal hepatocytes. For the preparation of a single-cell suspension, the livers of mice at E13.5 were dissociated in culture medium (DMEM supplemented with 10% FBS, γ -insulin [1 µg/ml, Wako], 0.1 µM dexamethasone [Sigma-Aldrich], 10 mM nicotinamide [Sigma-Aldrich], 2 mM L-glutamine [Gibco BRL], 50 µM β -mercaptoethanol [Sigma-Aldrich], 5 mM HEPES [Wako], and penicillin-streptomycin [Gibco BRL]) by repeated passage of the tissue through the mouth of a pipette. Human recombinant HGF (50 ng/ml, Sigma-Aldrich) and EGF (20 ng/ml, Sigma-Aldrich) were added to the cells at 24 hours after culture initiation. Cells were seeded at a density of 1×10^6 cells per well in 6-well plates for infection with retroviruses as described below (45).

Gene deletion in cultured cells by retroviral infection. cDNA encoding Cre recombinase was subcloned into the retroviral vector pMX-puro provided by T. Kitamura (University of Tokyo, Tokyo, Japan), and the resulting construct was introduced into Plat E packaging cells (63) with the use of the FuGENE6 reagent (Roche). The resulting culture supernatants containing the recombinant ecotropic retrovirus were harvested and incubated for 24 hours in the presence of Polybrene (2 µg/ml; Sigma-Aldrich) with proliferating liver cells harboring floxed alleles of *Fbxw7*, *Rbpj*, or *Myc*. The cells were cultured for an additional 24 hours in virus-free med-

ium, subjected to selection in medium containing puromycin (3 µg/ml), cultured for 96 hours in puromycin-free medium, and then harvested.

Statistics. Data are presented as mean \pm SD and were analyzed using 2-tailed Student's *t* test. A *P* value of less than 0.05 was considered statistically significant.

Acknowledgments

We thank T. Honjo for *Rbpj* floxed mice; I.M. de Alborán for *Myc* floxed mice; K. Rajewsky for *Mx1*-Cre transgenic mice; T. Kitamura for pMX-puro; S. Aishima, Y. Nishihara, M. Sakamoto, and R. Irie for discussion; N. Kitajima, Y. Yamada, and K. Takeda for technical assistance; members of our laboratories for comments on the manuscript; and A. Ohta and M. Kimura for help in the preparation of the manuscript. This work was supported in part by a grant from the Ministry of Education, Culture, Sports, Science, and Technology of Japan and by a research grant from the Takeda Science Foundation.

Received for publication August 6, 2009, and accepted in revised form September 29, 2010.

Address correspondence to: Keiichi I. Nakayama, Department of Molecular and Cellular Biology, Medical Institute of Bioregulation, Kyushu University, 3-1-1 Maidashi, Higashi-ku, Fukuoka, Fukuoka 812-8582, Japan. Phone: 81.92.642.6815; Fax: 81.92.642.6819; E-mail: nakayak1@bioreg.kyushu-u.ac.jp.

- Hershko A, Ciechanover A. The ubiquitin system. *Annu Rev Biochem.* 1998;67:425–479.
- Nakayama KI, Nakayama K. Ubiquitin ligases: cell-cycle control and cancer. *Nat Rev Cancer.* 2006; 6(5):369–381.
- Frescas D, Pagano M. Deregulated proteolysis by the F-box proteins SKP2 and beta-TrCP: tipping the scales of cancer. *Nat Rev Cancer.* 2008;8(6):438–449.
- Welcker M, Clurman BE. FBW7 ubiquitin ligase: a tumour suppressor at the crossroads of cell division, growth and differentiation. *Nat Rev Cancer.* 2008;8(2):83–93.
- Hubbard EJ, Wu G, Kitajewski J, Greenwald I. *sel-10*, a negative regulator of *lin-12* activity in *Caenorhabditis elegans*, encodes a member of the CDC4 family of proteins. *Genes Dev.* 1997;11(23):3182–3193.
- Sundaram M, Greenwald I. Suppressors of a *lin-12* hypomorph define genes that interact with both *lin-12* and *glp-1* in *Caenorhabditis elegans*. *Genetics.* 1993;135(3):765–783.
- Gupta-Rossi N, et al. Functional interaction between SEL-10, an F-box protein, and the nuclear form of activated Notch1 receptor. *J Biol Chem.* 2001;276(37):34371–34378.
- Oberg C, Li J, Pauley A, Wolf E, Gurney M, Lendahl U. The Notch intracellular domain is ubiquitinated and negatively regulated by the mammalian Sel-10 homolog. *J Biol Chem.* 2001;276(38):35847–35853.
- Koepf DM, et al. Phosphorylation-dependent ubiquitination of cyclin E by the SCF^{FBW7} ubiquitin ligase. *Science.* 2001;294(5540):173–177.
- Moberg KH, Bell DW, Wahrer DC, Haber DA, Hariharan IK. Archipelago regulates Cyclin E levels in *Drosophila* and is mutated in human cancer cell lines. *Nature.* 2001;413(6853):311–316.
- Strohmaier H, Spruck CH, Kaiser P, Won KA, Sangfelt O, Reed SI. Human F-box protein hCdc4 targets cyclin E for proteolysis and is mutated in a breast cancer cell line. *Nature.* 2001;413(6853):316–322.
- Yada M, et al. Phosphorylation-dependent degradation of c-Myc is mediated by the F-box protein Fbw7. *EMBO J.* 2004;23(10):2116–2125.
- Welcker M, et al. The Fbw7 tumor suppressor regulates glycogen synthase kinase 3 phosphorylation-dependent c-Myc protein degradation. *Proc Natl Acad Sci U S A.* 2004;101(24):9085–9090.
- Nateri AS, Riera-Sans L, Da Costa C, Behrens A. The ubiquitin ligase SCF^{FBW7} antagonizes apoptotic JNK signaling. *Science.* 2004;303(5662):1374–1378.
- Wei W, Jin J, Schlisio S, Harper JW, Kaelin WG Jr. The v-Jun point mutation allows c-Jun to escape GSK3-dependent recognition and destruction by the Fbw7 ubiquitin ligase. *Cancer Cell.* 2005;8(1):25–33.
- Sundqvist A, et al. Control of lipid metabolism by phosphorylation-dependent degradation of the SREBP family of transcription factors by SCF^{FBW7}. *Cell Metab.* 2005;1(6):379–391.
- Punga T, Bengoechea-Alonso MT, Ericsson J. Phosphorylation and ubiquitination of the transcription factor sterol regulatory element-binding protein-1 in response to DNA binding. *J Biol Chem.* 2006;281(35):25278–25286.
- Bengoechea-Alonso MT, Ericsson J. A phosphorylation cascade controls the degradation of active SREBP1. *J Biol Chem.* 2009;284(9):5885–5895.
- Mao JH, et al. FBXW7 targets mTOR for degradation and cooperates with PTEN in tumor suppression. *Science.* 2008;321(5895):1499–1502.
- Olson BL, et al. SCF^{Cdc4} acts antagonistically to the PGC-1alpha transcriptional coactivator by targeting it for ubiquitin-mediated proteolysis. *Genes Dev.* 2008;22(2):252–264.
- Maser RS, et al. Chromosomally unstable mouse tumours have genomic alterations similar to diverse human cancers. *Nature.* 2007;447(7147):966–971.
- Lee JW, et al. Mutational analysis of the hCDC4 gene in gastric carcinomas. *Eur J Cancer.* 2006; 42(14):2369–2373.
- Kemp Z, et al. CDC4 mutations occur in a subset of colorectal cancers but are not predicted to cause loss of function and are not associated with chromosomal instability. *Cancer Res.* 2005; 65(24):11361–11366.
- Hubalek MM, et al. Cyclin E dysregulation and chromosomal instability in endometrial cancer. *Oncogene.* 2004;23(23):4187–4192.
- Koh MS, Ittmann M, Kadmon D, Thompson TC, Leach FS. CDC4 gene expression as potential biomarker for targeted therapy in prostate cancer. *Cancer Biol Ther.* 2006;5(4):384–383.
- Calhoun ES, et al. BRAF and FBXW7 (CDC4, FBW7, AGO, SEL10) mutations in distinct subsets of pancreatic cancer: potential therapeutic targets. *Am J Pathol.* 2003;163(4):1255–1260.
- Akhoodi S, et al. FBXW7/hCDC4 is a general tumor suppressor in human cancer. *Cancer Res.* 2007;67(19):9006–9012.
- Song JH, Schnitke N, Zaat A, Walsh CS, Miller CW. FBXW7 mutation in adult T-cell and B-cell acute lymphocytic leukemias. *Leuk Res.* 2008; 32(11):1751–1755.
- Thompson BJ, et al. The SCF^{FBW7} ubiquitin ligase complex as a tumor suppressor in T cell leukemia. *J Exp Med.* 2007;204(8):1825–1835.
- O'Neil J, et al. FBW7 mutations in leukemic cells mediate NOTCH pathway activation and resistance to gamma-secretase inhibitors. *J Exp Med.* 2007;204(8):1813–1824.
- Malyukova A, et al. The tumor suppressor gene hCDC4 is frequently mutated in human T-cell acute lymphoblastic leukemia with functional consequences for Notch signaling. *Cancer Res.* 2007;67(12):5611–5616.
- Tsunematsu R, et al. Mouse Fbw7/Sel-10/Cdc4 is required for notch degradation during vascular development. *J Biol Chem.* 2004;279(10):9417–9423.
- Tetzlaff MT, et al. Defective cardiovascular development and elevated cyclin E and Notch proteins in mice lacking the Fbw7 F-box protein. *Proc Natl Acad Sci U S A.* 2004;101(10):3338–3345.
- Onoyama I, et al. Conditional inactivation of Fbxw7 impairs cell-cycle exit during T cell differentiation and results in lymphomagenesis. *J Exp Med.* 2007;204(12):2875–2888.
- Matsuoka S, et al. Fbxw7 acts as a critical fail-safe against premature loss of hematopoietic stem cells and development of T-ALL. *Genes Dev.* 2008;22(8):986–991.
- Ishikawa Y, Onoyama I, Nakayama KI, Nakayama K. Notch-dependent cell cycle arrest and apoptosis in mouse embryonic fibroblasts lacking Fbxw7. *Oncogene.* 2008;27(47):6164–6174.
- Matteoni CA, Younossi ZM, Gramlich T, Boparai N, Liu YC, McCullough AJ. Nonalcoholic fatty liver disease: a spectrum of clinical and pathological



- severity. *Gastroenterology*. 1999;116(6):1413-1419.
38. Horton JD, et al. Combined analysis of oligonucleotide microarray data from transgenic and knockout mice identifies direct SREBP target genes. *Proc Natl Acad Sci U S A*. 2003;100(21):12027-12032.
 39. Kim BJ, Fulton AB. The genetics and ocular findings of Alagille syndrome. *Semin Ophthalmol*. 2007;22(4):205-210.
 40. Loomes KM, et al. Bile duct proliferation in liver-specific Jag1 conditional knockout mice: effects of gene dosage. *Hepatology*. 2007;45(2):323-330.
 41. Nishikawa Y, et al. Transdifferentiation of mature rat hepatocytes into bile duct-like cells in vitro. *Am J Pathol*. 2005;166(4):1077-1088.
 42. McCright B, Lozier J, Gridley T. A mouse model of Alagille syndrome: Notch2 as a genetic modifier of Jag1 haploinsufficiency. *Development*. 2002;129(4):1075-1082.
 43. McDaniel R, et al. NOTCH2 mutations cause Alagille syndrome, a heterogeneous disorder of the notch signaling pathway. *Am J Hum Genet*. 2006;79(1):169-173.
 44. Kodama Y, Hijikata M, Kageyama R, Shimotohno K, Chiba T. The role of notch signaling in the development of intrahepatic bile ducts. *Gastroenterology*. 2004;127(6):1775-1786.
 45. Suzuki A, et al. Flow-cytometric separation and enrichment of hepatic progenitor cells in the developing mouse liver. *Hepatology*. 2000;32(6):1230-1239.
 46. Leclercq IA, Farrell GC, Schriemer R, Robertson GR. Leptin is essential for the hepatic fibrogenic response to chronic liver injury. *J Hepatol*. 2002;37(2):206-213.
 47. Sahai A, et al. Obese and diabetic *db/db* mice develop marked liver fibrosis in a model of nonalcoholic steatohepatitis: role of short-form leptin receptors and osteopontin. *Am J Physiol Gastrointest Liver Physiol*. 2004;287(5):G1035-G1043.
 48. Horie Y, et al. Hepatocyte-specific Pten deficiency results in steatohepatitis and hepatocellular carcinomas. *J Clin Invest*. 2004;113(12):1774-1783.
 49. Luedde T, et al. Deletion of NEMO/IKKgamma in liver parenchymal cells causes steatohepatitis and hepatocellular carcinoma. *Cancer Cell*. 2007;11(2):119-132.
 50. Isoda K, et al. Deficiency of interleukin-1 receptor antagonist deteriorates fatty liver and cholesterol metabolism in hypercholesterolemic mice. *J Biol Chem*. 2005;280(8):7002-7009.
 51. Nakanishi Y, et al. Nonalcoholic steatohepatitis and hepatocellular carcinoma in galectin-3 knockout mice. *Hepatology Res*. 2008;38(12):1241-1251.
 52. Yanagitani A, et al. Retinoic acid receptor alpha dominant negative form causes steatohepatitis and liver tumors in transgenic mice. *Hepatology*. 2004;40(2):366-375.
 53. Nakayama H, et al. Transgenic mice expressing nuclear sterol regulatory element-binding protein 1c in adipose tissue exhibit liver histology similar to nonalcoholic steatohepatitis. *Metabolism*. 2007;56(4):470-475.
 54. Consortium ECTS. Identification and characterization of the tuberous sclerosis gene on chromosome 16. *Cell*. 1993;75(7):1305-1315.
 55. van Slegtenhorst M, et al. Identification of the tuberous sclerosis gene TSC1 on chromosome 9q34. *Science*. 1997;277(5327):805-808.
 56. Li Z, White P, Tuteja G, Rubins N, Sackett S, Kaestner KH. Foxa1 and Foxa2 regulate bile duct development in mice. *J Clin Invest*. 2009;119(6):1537-1545.
 57. Kuhn R, Schwenk F, Aguet M, Rajewsky K. Inducible gene targeting in mice. *Science*. 1995;269(5229):1427-1429.
 58. Postic C, Magnuson MA. DNA excision in liver by an albumin-Cre transgene occurs progressively with age. *Genesis*. 2000;26(2):149-150.
 59. Tanigaki K, et al. Regulation of alphabeta/gammadelta T cell lineage commitment and peripheral T cell responses by Notch/RBP-J signaling. *Immunity*. 2004;20(5):611-622.
 60. de Alboran IM, et al. Analysis of C-MYC function in normal cells via conditional gene-targeted mutation. *Immunity*. 2001;14(1):45-55.
 61. Kamura T, et al. Cytoplasmic ubiquitin ligase KPC regulates proteolysis of p27^{Kip1} at G1 phase. *Nat Cell Biol*. 2004;6(12):1229-1235.
 62. Nishiyama M, Nakayama K, Tsunematsu R, Tsukiyama T, Kikuchi A, Nakayama KI. Early embryonic death in mice lacking the beta-catenin-binding protein Duplin. *Mol Cell Biol*. 2004;24(19):8386-8394.
 63. Morita S, Kojima T, Kitamura T. Plat-E: an efficient and stable system for transient packaging of retroviruses. *Gene Ther*. 2000;7(12):1063-1066.

