

| | | | | | |
|--------------------|---|----------------------------|-----|---------|------|
| 仲 哲治 (Naka, T.) | Plasma membrane proteomics identifies bone marrow stromal antigen 2 as a potential therapeutic target in endometrial cancer. | Int. J. Cancer | 132 | 472-84 | 2013 |
| 同 上 | Periostin facilitates Skin sclerosis via PI3K/Akt dependent mechanism in a mouse model of scleroderma. | PLoS One | 7 | E41994 | 2012 |
| 同 上 | Serum HE4 as a diagnostic and prognostic marker for lung cancer. 33(4), (2012). | Tumour Biol. | 33 | 1141-9 | 2012 |
| 同 上 | Serum leucine-rich alpha-2 glycoprotein is a disease activity biomarker in ulcerative colitis. 潰瘍性大腸炎の疾患活動性マーカーとしての血清ロイシンリッチアルファ2グリコプロテイン | Intestine | 17 | 107-9 | 2013 |
| 同 上 | Therapeutic strategies in epithelial ovarian cancer. | J. Exp. Clin. Cancer Res. | 31 | 14 | 2012 |
| 同 上 | Dysregulation of melanocyte function by Th17-related cytokines: significance of Th17 cell infiltration in autoimmune Vitiligo Vulgaris. | Pigment Cell Melanoma Res. | 25 | 219-30 | 2012 |
| 同 上 | Periostin, a matrix-cellular protein, accelerates cutaneous wound repair by activating dermal fibroblasts. | Exp. Dermatol. | 21 | 331-6 | 2012 |
| 同 上 | Serum leucine-rich alpha-2 glycoprotein is a disease activity biomarker in ulcerative colitis. | Inflamm. Bowel Dis. | 17 | 491-502 | 2011 |

| | | | | | |
|--------------------------|---|----------------------------|----------|---------|------|
| 仲 哲治 (Naka, T.) | Targeting annexin A4 to counteract chemoresistance in clear cell carcinoma of the ovary. | Expert Opin. Ther. Targets | 14 | 963-71 | 2010 |
| 同 上 | iTRAQ-based proteomic identification of leucine rich alpha 2 glycoprotein (LRG) as a novel inflammatory biomarker in autoimmune diseases. | Ann. Rheum. Dis. | 69 | 770-4 | 2010 |
| 同 上 | Enhanced expression of Annexin A4 in clear cell carcinoma of the ovary and its association with chemoresistance to carboplatin. | Int. J. Cancer | 125 | 2316-22 | 2009 |
| 中山敬一 (Nakayama, K.I.) | Regulation of c-Myc ubiquitination controls chronic myelogenous leukemia initiation and progression. | Cancer Cell | 23 | 362-75 | 2013 |
| 同 上 | Ablation of fbwx7 eliminates leukemia-initiating cells by preventing quiescence. Cancer Cell | Cancer Cell | 23 | 347-61 | 2013 |
| 同 上 | FBXL21 regulates oscillation of the circadian clock through ubiquitination and stabilization of cryptochromes. | Cell | 152 | 1106-18 | 2013 |
| 同 上 | Selective escape of proteins from the mitochondria during mitophagy. | Nature Commun. | 4 | 1410 | 2013 |
| 同 上 | Comprehensive identification of substrates for F-box proteins by differential proteomics analysis. | J. Proteome Res. | In press | | 2013 |

| | | | | | |
|--------------------------|---|------------------|-----|----------|------|
| 中山敬一 (Nakayama, K.I.) | Skp2-SCF E3 ligase regulates Akt ubiquitination, glycolysis, herceptin sensitivity, and tumorigenesis. | Cell | 149 | 1098-111 | 2012 |
| 同 上 | Proteome-wide identification of ubiquitylation sites by conjugation of engineered lysine-less ubiquitin. | J. Proteome Res. | 11 | 796-807 | 2012 |
| 同 上 | The FBXL5- IRP2 axis is integral to control of iron metabolism in vivo. | Cell Metab. | 14 | 339-51 | 2011 |
| 同 上 | p57 is required for quiescence and maintenance of adult hematopoietic stem cells. | Cell Stem Cell | 9 | 262-71 | 2011 |
| 同 上 | p57 ^{Kip2} and p27 ^{Kip1} cooperate to maintain hematopoietic stem cell quiescence through interactions with Hsc70. | Cell Stem Cell | 9 | 247-61 | 2011 |
| 同 上 | SCF ^{FBW7} regulates cellular apoptosis by targeting MCL1 for ubiquitylation and destruction. | Nature | 471 | 104-9 | 2011 |
| 同 上 | UBE4B promotes Hdm2-mediated degradation of the tumor suppressor p53. | Nature Med. | 17 | 347-355 | 2011 |
| 同 上 | Fbxw7 regulates lipid metabolism and cell fate decisions in the mouse liver. | J. Clin. Invest. | 121 | 342-354 | 2011 |

| | | | | | |
|--------------------------|---|-------------------|-----|---------|------|
| 中山敬一 (Nakayama, K.I.) | Deficiency of Rap1-binding protein RAPL causes lymphoproliferative disorders through mislocalization of p27 ^{kip1} . | Immunity | 34 | 24-38 | 2011 |
| 同 上 | Deciphering the transcriptional complex critical for RhoA gene expression and cancer metastasis. | Nature Cell Biol. | 12 | 457-67 | 2010 |
| 同 上 | Skp2 targeting suppresses tumorigenesis by Arf-p53- independent cellular senescence. | Nature | 464 | 374-9 | 2010 |
| 同 上 | KDM7 is a dual demethylase for histone H3 Lys 9 and Lys 27 and functions in brain development. | Genes Dev. | 24 | 432-7 | 2010 |
| 同 上 | Skp2 is required for survival of aberrantly proliferating Rb1-deficient cells and for tumorigenesis in Rb1 ^{+/-} mice. | Nature Genet. | 42 | 83-8 | 2010 |
| 同 上 | Large-scale proteomic analysis of tyrosine-phosphorylation induced by T-cell receptor or B-cell receptor activation reveals new signaling pathways. | Proteomics | 9 | 3549-63 | 2009 |
| 同 上 | Phosphorylation-dependent regulation of cytosolic localization and oncogenic function of Skp2 by Akt/PKB. | Nature Cell Biol. | 11 | 420-32 | 2009 |

| | | | | | |
|--------------------------|--|----------------------------|----------|----------|------|
| 中山敬一 (Nakayama, K.I.) | Common and specific roles of the related CDK inhibitors p27 and p57 revealed by a knock-in mouse model. | Proc. Natl. Acad. Sci. USA | 106 | 5192-97 | 2009 |
| 同上 | CHD8 suppresses p53-mediated apoptosis through histone H1 recruitment during early embryogenesis. Nature Cell Biol. | Nature Cell Biol. | 11 | 172-82 | 2009 |
| 平野 久 (Hirano, H.) | Differential proteome analysis of serum proteins associated with the development of type 2 diabetes mellitus in the KK- <i>Ay</i> mouse model using the iTRAQ technique. | J. Proteomics | In press | | 2013 |
| 同上 | Identification of phosphorylated proteins involved in the oncogenesis of prostate cancer via Pin1-proteomic analysis. | Prostate | 72 | 626-37 | 2012 |
| 同上 | N-Myristoylation of the Rpt2 subunit regulates intracellular localization of the yeast 26S proteasome. | Biochemistry | 51 | 8856-66 | 2012 |
| 同上 | Effects of growth hormone on the salmon pituitary proteome. | J. Proteomics | 75 | 1718-31 | 2012 |
| 同上 | Structural basis for inhibition of xyloglucan-specific endo- β -1,4-glucanase (XEG) by XEG-protein inhibitor. | J. Biol. Chem. | 287 | 18710-16 | 2012 |

| | | | | | |
|---------------------|--|---------------|-----|---------|------|
| 平野 久 | 卵巣明細胞腺がん関連タンパク質の発現調節. | 生物物理化学 | 55 | 5-8 | 2011 |
| 同 上 (Hirano, H.) | N ^α -Acetylation of yeast ribosomal proteins: identification by 2D-DIGE MS/MS and analysis of the effect on proteins synthesis. | J. Proteomics | 74 | 431-41 | 2011 |
| 同 上 | Cell polarity in <i>Saccharomyces cerevisiae</i> depends on proper localization of the Bud9 landmark protein by the EKC/KEOPS complex. | Genetics | 188 | 871-882 | 2011 |
| 同 上 | Wild-type p53 enhances annexin IV gene expression in ovarian clear cell adenocarcinoma. | FEBS J. | 278 | 1470-83 | 2011 |
| 同 上 | AAA+ proteins RUVBL1 and RUVBL2 coordinate PIKK family and function in nonsense-mediated mRNA decay. 3, 1-13 (2010). | Sci. Signal | 3 | 1-13 | 2010 |
| 同 上 | Co- and post-translational modifications of the 26S proteasome in yeast. | Proteomics | 10 | 2769-79 | 2010 |
| 同 上 | Characterization of multiple alternative forms of heterogeneous nuclear ribonucleoprotein K by phosphate-affinity electrophoresis. | Proteomics | 10 | 3884-95 | 2010 |

| | | | | | |
|------------------|--|----------------------|------|--------------------|------|
| 尾野雅哉 (Ono, M) | Quantitative targeted absolute proteomics-based large-scale quantification of proline-hydroxylated alpha-fibrinogen in plasma for pancreatic cancer diagnosis. | J. Proteome Res | 12 | 753-62 | 2013 |
| 同 上 | Proteomic Approaches to the Discovery of Cancer Biomarkers for Early Detection and Personalized Medicine. | Jpn. J. Clin. Oncol. | 43 | 103-9 | 2013 |
| 同 上 | ADP-ribosylation of guanosine by SCO5461 protein secreted from Streptomyces coelicolor. | Toxicon. | 63 | 55-63 | 2013 |
| 同 上 | プロテオーム解析によるがん診断と治療への道 | 分子消化器病 | 9 | 71-6 | 2012 |
| 同 上 | Carbonic anhydrase I as a new plasma biomarker for prostate cancer. | ISRN Oncol. | 2012 | Article ID: 768190 | 2012 |
| 同 上 | DDX31 regulates the p53-HDM2 pathway and rRNA gene transcription through its interaction with NPM1 in renal cell carcinomas. | Cancer Res. | 72 | 5867-77 | 2012 |
| 同 上 | プロテオミクス解析システム 2DICAL を用いたがんバイオマーカー開発. | 細胞 | 44 | 46-49 | 2012 |

| | | | | | |
|-------------------|--|-----------------------------|------|-------------------|------|
| 尾野雅哉 (Ono, M.) | Biomarker Discovery of Pancreatic and Gastrointestinal Cancer by 2DICAL: 2-Dimensional Image-Converted Analysis of Liquid Chromatography and Mass Spectrometry. | Int. J. Proteomics | 2012 | Article ID 897412 | 2012 |
| 同 上 | Identification of 14-3-3gamma as a Miep-interacting protein and its role in mitochondrial quality control. | Sci, Rep. | 2 | 379 | 2012 |
| 同 上 | Use of quantitative shotgun proteomics to identify fibronectin 1 as a potential plasma biomarker for clear cell carcinoma of the kidney. | Cancer Biomark | 10 | 175-83 | 2012 |
| 同 上 | Identification of adipophilin as a potential plasma biomarker for colorectal cancer using label-free quantitative mass spectrometry and protein microarray. Cancer | Epidemiol. Biomarkers Prev. | 20 | 2195-203 | 2011 |
| 同 上 | Combined functional genome survey of therapeutic targets for clear cell carcinoma of the kidney. | Jpn. J. Clin. Oncol. | 41 | 847-853 | 2011 |

| | | | | | |
|-------------------|---|------------------------------------|-----|----------|------|
| 尾野雅哉 (Ono, M.) | Plasma biomarker discovery and validation for colorectal cancer by quantitative shotgun mass spectrometry and protein microarray. | Cancer Sci. | 102 | 630-638 | 2011 |
| 同 上 | Possible Existence of Lysosome-Like Organella within Mitochondria and Its Role in Mitochondrial Quality Control. | PLoS ONE | 6 | e16054 | 2011 |
| 同 上 | Reduced Plasma Level of CXC Chemokine Ligand 7 in Patients with Pancreatic Cancer. | Cancer Epidemiol. Biomarkers Prev. | 20 | 160-171 | 2011 |
| 同 上 | Traf2- and Nck-interacting kinase is essential for Wnt signaling and colorectal cancer growth. | Cancer Res. | 70 | 5024-33 | 2010 |
| 同 上 | Traf2- and Nck-interacting kinase is essential for canonical Wnt signaling in Xenopus axis formation. | J. Biol. Chem. | 285 | 26289-94 | 2010 |
| 同 上 | Survival prediction for pancreatic cancer patients receiving gemcitabine treatment. | Mol. Cell. proteomics | 9 | 695-704 | 2010 |

| | | | | | |
|------------------------|---|-----------------------|----|---------|------|
| 南野直人 (Minamino, N.) | Direct immunochemi-luminescent assay for proBNP and total BNP in human plasma: proBNP and total BNP levels in healthy individuals and heart failure patients. | Plos One | 8 | e53233 | 2013 |
| 同 上 | Large-scale identification of endogenous secretory peptides using electron transfer dissociation mass spectrometry. | Mol. Cell. Proteomics | 12 | 700-9 | 2013 |
| 同 上 | Peptidomics-based discovery of an antimicrobial peptide derived from insulin-like growth factor-binding protein 5. | J. Proteome Res. | 10 | 1870-80 | 2011 |
| 同 上 | A peptidomics strategy for discovering endogenous bioactive peptides. | J. Proteome Res. | 9 | 5047-52 | 2010 |
| 同 上 | Snapshot peptidomics of the regulated secretory pathway. | Mol. Cell. Proteomics | 8 | 1638-47 | 2009 |
| 高坂新一 (Kohsaka, S) | Robo1 regulates the migration and laminar distribution of upper-layer pyramidal neurons of the cerebral cortex. | Cerebral Cortex | 23 | 1495-50 | 2013 |

| | | | | | |
|----------------------|--|------------------|-----|---------|------|
| 高坂新一 (Kohsaka, S) | Accumulating microglia phagocytose injured neurons in hippocampal slice cultures: involvement of p38 MAP kinase. | PLoS One | 7 | 1-11 | 2012 |
| 同 上 | Long-term changes of spine dynamics and microglia after transient peripheral immune response triggered by LPS in vivo. | Mol. Brain | 17 | 4-27 | 2011 |
| 同 上 | NMDA receptor regulates migration of newly generated neurons in the adult hippocampus via Disrupted-In-Schizophrenia 1(DISC1). | J. Neurochem. | 118 | 34-44 | 2011 |
| 同 上 | Appearance of phagocytic microglia adjacent to motoneurons in spinal cord tissue from a presymptomatic transgenic rat model of amyotrophic lateral sclerosis. | J Neurosci. Res. | 88 | 2736-46 | 2010 |
| 同 上 | Inhibition of N-methyl-d-aspartate receptor activity resulted in aberrant neuronal migration caused by delayed morphological development in the mouse neocortex. | Neuroscience | 169 | 609-18 | 2010 |

| | | | | | |
|------------------------|--|-----------------|-----|----------|------|
| 高坂新一 (Kohsaka, S.) | Pigment epithelium-derived factor up-regulation induced by memantine, an N-methyl- d- aspartate receptor antagonist, is involved in increased proliferation of hippocampal progenitor cells. | Neuroscience | 167 | 372-83 | 2010 |
| 同 上 | P2Y12 receptor-mediated integrin-β1 activation regulates microglial process extension induced by ATP. | Glia | 58 | 790-801 | 2010 |
| 同 上 | The Alzheimer's disease drug memantine increases the number of radial glia-like progenitor cells in adult hippocampus. | Glia | 57 | 1082-90 | 2009 |
| 同 上 | Neuroprotective effect of Sonic hedgehog up-regulated in Schwann cells following sciatic nerve injury. | J. Neurochem. | 107 | 918-27 | 2008 |
| 宮本泰傑 (Miyamoto, Y.) | Androgen-regulated transcriptional control of sialyltransferases in prostate cancer cells. | PLoS One | 7 | e31234 | 2012 |
| 同 上 | Analysis of polarized secretion of fucosylated alpha-fetoprotein in HepG2 cells. | J. Proteome Res | 11 | 2798-806 | 2012 |
| 同 上 | 13-Cis retinoic acid can enhance the antitumor activity of non-replicating Sendai virus particle against neuroblastoma. | Cancer Sci. | 104 | 238-44 | 2013 |

| | | | | | |
|------------------------|--|-----------------------|----------|---------|------|
| 宮本泰傑 (Miyamoto, Y.) | Occurrence of free deaminoneuraminic acid (KDN)-containing complex-type N-glycans in human prostate cancers. | Glycobiology | In press | | 2013 |
| 同 上 | Accumulation of free Neu5Ac- containing complex-type N-glycans in human pancreatic cancers. | Glycoconj. J. | 30 | 247-56 | 2013 |
| 同 上 | Expression of gangliosides, GD1a and sialyl paragloboside, is regulated by NF-kappaB-dependent transcriptional control of alpha2,3-sialyltransferase I, II and VI in human castration-resistant prostate cancer cells. | Int. J. Cancer | 129 | 1838-47 | 2011 |
| 同 上 | N-Glycosylation profiling of recombinant mouse extracellular superoxide dismutase produced in Chinese hamster ovary cells. | Glycoconj. J. | 28 | 183-96 | 2011 |
| 加藤菊也 (Kato, K.) | Dynamics of cancer cell subpopulations in primary and metastatic colorectal tumors. | Clin. Exp. Metastasis | 28 | 427-35 | 2011 |
| 同 上 | Quantitative detection of EGFR mutations in circulating tumor DNA derived from lung adenocarcinomas. | Clin. Cancer Res. | 17 | 7808-15 | 2011 |

| | | | | | |
|------------------------|---|----------------|------|----------|------|
| 宮本泰豪 (Miyamoto, Y.) | Identification of various types of α 2-HS glycoprotein in sera of patients with pancreatic cancer: Possible implication in resistance to protease treatment. | Mol. Med. Rep. | 3 | 651-6 | 2010 |
| 同 上 | Involvement of ST6Gal I in the biosynthesis of a unique human colon cancer biomarker candidate, α 2,6-sialylated blood group type 2H (ST2H) antigen. | J. Biochem. | 148 | 359-70 | 2010 |
| 同 上 | Novel ganglioside found in adenocarcinoma cells of Lewis-negative patients. | Glycobiology | 20 | 1594-606 | 2010 |
| 加藤菊也 (Kato, K.) | Transcriptional induction and translational inhibition of Arc and Cugbp2 in mice hippocampus after transient global ischemia under normothermic condition. | Brain Res. | 1287 | 136-45 | 2009 |
| 同 上 | Algorithm for in vitro diagnostic multivariate index assay. | Breast Cancer. | 16 | 248-51 | 2009 |

| | | | | | |
|------------------------|--|-------------------------------|-----|-----------|------|
| 加藤菊也 (Kato, K.) | Impact of the next generation DNA sequencers. | Int. J. Clin. Exp. Med. | 2 | 193-202 | 2009 |
| 同 上 | Putative precursor cancer cells in human colorectal cancer tissue. | Int. J. Clin. Exp. Pathol. | 2 | 154-62 | 2009 |
| 同 上 | Optimal aggregation of binary classifiers for multiclass cancer diagnosis using gene expression profiles. IEEE/ACM Trans Comput. Biol. Bioinform. | Trans Comput. Biol. Bioinform | 6 | 333-43 | 2009 |
| 同 上 | Using gene expression profiling to identify a prognostic molecular spectrum in gliomas. | Cancer Sci. | 100 | 165-72 | 2009 |
| 宮本泰豪 (Miyamoto, Y.) | Comprehensive clinico-glycomic study of 16 colorectal cancer specimens: elucidation of aberrant glycosylation and its mechanistic causes in colorectal cancer cells. | J. Proteome Res. | 8 | 2990-3005 | 2009 |
| 野村文夫 (Nomura, F.) | Proteomic analysis of gingival crevicular fluid for discovery of novel periodontal disease markers. | Proteomics | 12 | 2190-202 | 2012 |

| | | | | | |
|----------------------|---|-------------------------|----------|---------|------|
| 野村文夫 (Nomura, F.) | Increased serum levels of pigment epithelium-derived factor by excessive alcohol consumption-detection and identification by a three-steps serum proteome analysis. | Alcohol Clin. Exp. Res. | 35 | 211-7 | 2011 |
| 同 上 | Serum level of fibrinogen A α chain fragment increases in chronic thromboembolic pulmonary hypertension. 75 , (2011). | Circ. J. | 75 | 2675-82 | 2011 |
| 荒木令江 (Araki, N.) | Integrated proteomics identified a novel activation signaling of dynein IC2-GR-COX-1 in NF1 disease model cells. | Mol. Cell. Proteomics | 12 | 1377-94 | 2013 |
| 同 上 | Glioma initiating cells form a differentiation niche via the induction of extracellular matrices and integrin alpha V. | PLOS ONE | In press | | 2013 |
| 同 上 | Epiplakin modifies the motility of the HeLa cells and accumulates at the outer surfaces of three-dimensional cell clusters. | J. Dermatol. | 40 | 249-58 | 2013 |
| 同 上 | Enhancement of human cancer cell motility and invasiveness by anaphylatoxin C5a via aberrantly expressed C5a-receptor (CD88). | Clin. Cancer Res. | 19 | 1-10 | 2013 |

| | | | | | |
|---------------------|--|----------------------------|-------------|----------|------|
| 荒木令江 (Araki, N.) | Phosphorylation by PKD2 at Ser171 of SET reduced its inhibitory effect on PP2A phosphatase in T cells. | PLOS ONE | 7 | e51242 | 2012 |
| 同 上 | Expression and purification of human FROUNT, a common cytosolic regulator of CCR2 and CCR5. | Protein Expr. Purif | 77 | 86-91 | 2011 |
| 同 上 | A novel serum carbohydrate marker on MUC5AC: values for diagnostic and prognostic indicators for cholangiocarcinoma. | Cancer | 117 | 3393-403 | 2011 |
| 同 上 | Possible involvement of cyclophilin a processing in fumagillin- induced suppression of cholangiocarcinoma cell proliferation. | Asian Pac. J. Cancer Prev. | 13 | 137-41 | 2012 |
| 同 上 | Serial analysis of gene expression reveals promising therapeutic targets for liver fluke-associated cholangiocarcinoma. | Asian Pac. J. Cancer Prev. | 13 Suppl | 89-93 | 2012 |
| 同 上 | Contribution of RIZ1 to regulation of proliferation and migration of a liver fluke-related cholangiocarcinoma cell. | Asian Pac. J. Cancer Prev. | 13 | 4007-11 | 2012 |
| 同 上 | The contribution of BCR-ABL-independent activation of ERK1/2 to acquired imatinib resistance in K562 chronic myeloid leukemia cells. | Cancer Sci. | 101 | 137-42 | 2010 |

| | | | | | |
|------------------------|--|-----------------------|-----|---------|------|
| 荒木令江 (Araki, N.) | An integrated approach of differential Mass Spectrometry and gene ontology analysis identified novel proteins regulating neuronal differentiation and survival. | Mol. Cell. Proteomics | 8 | 2350-67 | 2009 |
| 同 上 | Suppression of galectin-3 expression enhances apoptosis and chemosensitivity in liver fluke associated cholangio-carcinoma. | Cancer Sci. | 38 | 249-61 | 2009 |
| 同 上 | Silver ion unusually stabilizes the structure of parallel-motif DNA triplex. | J. Am. Chem. Soc. | 131 | 3826-7 | 2009 |
| 同 上 | Involvement of PI3K-Akt-Bad pathway in apoptosis induced by 2,6-di-O-methyl- β -cyclo-dextrin, not 2,6-di-O-methyl- α -cyclodextrin, through cholesterol depletion from lipid rafts on plasma membranes in cells. | Eur. J. Pharm. Sci. | 38 | 249-61 | 2009 |
| 中村和行 (Nakamura, K.) | Proteomic analysis showed down-regulation of nucleophosmin in progressive tumor cells compared to regressive tumor cells. | Anticancer Res. | 33 | 153-160 | 2012 |
| 同 上 | Proteomic differential display identifies upregulated vinculin as a possible biomarker of pancreatic cancer. | Oncol. Rep. | 28 | 1845-50 | 2012 |

| | | | | | |
|------------------------|--|------------------|----|---------|------|
| 中村和行 (Nakamura, K.) | Glyoxalase I (GLO1) is up-regulated in pancreatic cancerous tissues compared with related non-cancerous tissues. | Anticancer. Res. | 32 | 3219-22 | 2012 |
| 同上 | Heat-shock protein 27 plays the key role in gemcitabine-resistance of pancreatic cancer cells. 32, 2 (2012). | Anticancer Res. | 32 | 295-99 | 2012 |
| 同上 | Identification of four isoforms of aldolase B down-regulated in hepatocellular carcinoma tissues by means of two-dimensional Western blotting. | In Vivo | 25 | 881-6 | 2011 |
| | Up-regulation of 42 kDa tubulin alpha-6 chain fragment in well-differentiated hepatocellular carcinoma tissues from patients infected with hepatitis C virus. | Anticancer Res. | 31 | 3331-6 | 2011 |
| 同上 | Differential expression of up-regulated cofilin-1 and down-regulated cofilin-2 characteristic of pancreatic cancer tissues. | Oncol. Rep. | 26 | 1595-9 | 2011 |
| 同上 | Proteomic differential display analysis for TS-1-resistant and -sensitive pancreatic cancer cells using two-dimensional gel electrophoresis and mass spectrometry. | Anticancer Res. | 31 | 2103-8 | 2011 |

| | | | | | |
|------------------------|---|-----------------|----|---------|------|
| 中村和行 (Nakamura, K.) | Identification of differentially expressed proteins in tumor necrosis factor-alpha-resistant and -sensitive rat hepatoma cells. | Anticancer Res. | 31 | 2059-63 | 2011 |
| 同 上 | Screening for serological biomarkers of pancreatic cancer by two-dimensional electrophoresis and liquid chromatography-tandem mass spectrometry., | Oncol. Rep. | 26 | 287-92 | 2011 |
| 同 上 | Two-dimensional gel electrophoresis using immobilized pH gradient strips and Flamingo™ fluorescent gel stain identified non-nuclear proteins possibly related to malignant tumor progression. | Anticancer Res. | 31 | 1259-63 | 2011 |
| 同 上 | KNK437 down regulates heat shock protein 27 of pancreatic cancer cells and enhances the cytotoxic effect of gemcitabine. | Chemotherapy | 57 | 12-6 | 2011 |
| 同 上 | Proteomic differential display analysis shows up-regulation of 14-3-3 protein sigma in human scirrhous-type gastric carcinoma cells. | Anticancer Res. | 30 | 4459-65 | 2010 |
| 同 上 | Staining with highly sensitive Coomassie Brilliant Blue SeePico™ stain after Flamingo™ fluorescent gel stain is useful for cancer proteomic analysis by means of | Anticancer Res. | 30 | 4001-5 | 2010 |