

- [11] Vaux, D.L. and Silke, J. (2005) IAPs, RINGs and ubiquitylation. *Nat. Rev. Mol. Cell Biol.* 6, 287–297.
- [12] Gyrd-Hansen, M. and Meier, P. (2010) IAPs: from caspase inhibitors to modulators of NF-kappaB, inflammation and cancer. *Nat. Rev. Cancer* 10, 561–574.
- [13] Sekine, K., Takubo, K., Kikuchi, R., Nishimoto, M., Kitagawa, M., Abe, F., Nishikawa, K., Tsuruo, T. and Naito, M. (2008) Small molecules destabilize cIAP1 by activating auto-ubiquitylation. *J. Biol. Chem.* 283, 8961–8968.
- [14] Kim, S., Ohoka, N., Okuhira, K., Sai, K., Nishimaki-Mogami, T. and Naito, M. (2010) Modulation of RIP1 ubiquitylation and distribution by MeBS to sensitize cancer cells to tumor necrosis factor alpha-induced apoptosis. *Cancer Sci.* 101, 2425–2429.
- [15] Sato, S., Aoyama, H., Miyachi, H., Naito, M. and Hashimoto, Y. (2008) Demonstration of direct binding of cIAP1 degradation-promoting bestatin analogs to BIR3 domain: synthesis and application of fluorescent bestatin ester analogs. *Bioorg. Med. Chem. Lett.* 18, 3354–3358.
- [16] Sato, S., Tetsuhashi, M., Sekine, K., Miyachi, H., Naito, M., Hashimoto, Y. and Aoyama, H. (2008) Degradation-promoters of cellular inhibitor of apoptosis protein 1 based on bestatin and actinonin. *Bioorg. Med. Chem.* 16, 4685–4698.
- [17] Itoh, Y., Ishikawa, M., Naito, M. and Hashimoto, Y. (2010) Protein knockdown using methyl bestatin-ligand hybrid molecules: design and synthesis of inducers of ubiquitination-mediated degradation of cellular retinoic acid-binding proteins. *J. Am. Chem. Soc.* 132, 5820–5826.
- [18] Uren, A.G., Pakusch, M., Hawkins, C.J., Puls, K.L. and Vaux, D.L. (1996) Cloning and expression of apoptosis inhibitory protein homologs that function to inhibit apoptosis and/or bind tumor necrosis factor receptor-associated factors. *Proc. Natl. Acad. Sci. USA* 93, 4974–4978.

