

compression myelopathy. One month after G-CSF administration, mean recovery rate of JOA score was 29.1%. In contrast, it was 1.1% in the control group at 1 month after initial treatment. In addition, we observed that both motor power and pain sensation scores significantly increased in the G-CSF group compared with the control group at 1 month after treatment. No surgical treatment was performed in patients of either group during the month after G-CSF administration or initial treatment, and they were equally provided conservative treatment such as bed rest. Thus, the present results strongly suggest that G-CSF administration exhibited a neuroprotective effect for the injured spinal cord in patients with worsening symptoms of thoracic myelopathy and improved the myelopathy.

To the best of our knowledge, there has been no other medical treatment that has provided reliable evidence for improvement of thoracic myelopathy. This study provides evidence that G-CSF neuroprotective therapy may be useful as a medical treatment of patients with worsening symptoms of thoracic compression myelopathy. The G-CSF therapy may be especially useful for patients in whom the treatment of complications other than myelopathy needs to be given priority and thus requires a long waiting period before surgery.

In our present trial, no severe side effects occurred. Thus, we suggest that the dose (10 $\mu\text{g}/\text{kg}$ per d), duration (5 consecutive days), and route (intravenous administration) of G-CSF administration used in this study are principally safe for the treatment of patients with thoracic myelopathy.

The biggest limitation of this study was that the trial was performed as an open-labeled study and the selection of patients to the G-CSF group and the control group was not randomized. We cannot deny the possibility that a placebo effect of injection may participate in the improvement of neurological symptoms. To increase the level of evidence, in the next stage the study design should be a randomized, double-blind placebo-controlled study. By conducting a phase IIb clinical trial in a large number of patients with the study design described earlier, we will be able to reach a better conclusion regarding the effectiveness of G-CSF neuroprotective therapy for patients with worsening symptoms of thoracic compression myelopathy.

➤ Key Points

- ❑ A multicenter prospective controlled clinical trial was performed to confirm the feasibility of G-CSF administration for patients with worsening symptoms of thoracic myelopathy.
- ❑ For 10 patients with progressive myelopathy, G-CSF (10 $\mu\text{g}/\text{kg}$ per day) was intravenously administered for 5 consecutive days.
- ❑ The administration of G-CSF caused neurological recovery in the patients.

References

1. Roberts AW. G-CSF: a key regulator of neutrophil production, but that's not all! *Growth Factors* 2005;23:33–41.
2. Harada M, Qin Y, Takano H, et al. G-CSF prevents cardiac remodeling after myocardial infarction by activating the Jak–Stat pathway in cardiomyocytes. *Nat Med* 2005;11:305–11.
3. Schäbitz WR, Kollmar R, Schwaninger M, et al. Neuroprotective effect of granulocyte colony-stimulating factor after focal cerebral ischemia. *Stroke* 2003;34:745–51.
4. Koda M, Nishio Y, Kamada T, et al. Granulocyte colony-stimulating factor (G-CSF) mobilizes bone marrow-derived cells into injured spinal cord and promotes functional recovery after compression-induced spinal cord injury in mice. *Brain Res* 2007;1149:223–31.
5. Nishio Y, Koda M, Kamada T, et al. Granulocyte colony-stimulating factor attenuates neuronal death and promotes functional recovery after spinal cord injury in mice. *J Neuropathol Exp Neurol* 2007;66:724–31.
6. Kawabe J, Koda M, Hashimoto M, et al. Granulocyte colony-stimulating factor (G-CSF) exerts neuroprotective effects via promoting angiogenesis after spinal cord injury in rats. *J Neurosurg Spine* 2011;15:414–21.
7. Sakuma T, Yamazaki M, Okawa A, et al. Neuroprotective therapy using granulocyte-colony stimulating factor for patients with worsening symptoms of compression myelopathy, part 1: a phase I and IIa clinical trial [published online ahead of print September 21, 2011]. *Eur Spine J* 2012;21:482–9.
8. Zhang Y, Wang L, Fu Y, et al. Preliminary investigation of effect of granulocyte colony stimulating factor on amyotrophic lateral sclerosis. *Amyotroph Lateral Scler* 2009;10:430–1.
9. Nefussy B, Artamonov I, Deutsch V, et al. Recombinant human granulocyte-colony stimulating factor administration for treating amyotrophic lateral sclerosis: a pilot study. *Amyotroph Lateral Scler* 2010;11:187–93.
10. Shyu WC, Lin SZ, Lee CC, et al. Granulocyte colony-stimulating factor for acute ischemic stroke: a randomized controlled trial. *CMAJ* 2006;174:927–33.

Accumulation of Activated Invariant Natural Killer T Cells in the Tumor Microenvironment after α -Galactosylceramide-Pulsed Antigen Presenting Cells

Kaoru Nagato · Shinichiro Motohashi ·
Fumihiko Ishibashi · Kohsuke Okita ·
Kazuki Yamasaki · Yasumitsu Moriya ·
Hidehisa Hoshino · Shigetoshi Yoshida ·
Hideki Hanaoka · Shin-ichiro Fujii ·
Masaru Taniguchi · Ichiro Yoshino ·
Toshinori Nakayama

Received: 31 January 2012 / Accepted: 10 April 2012 / Published online: 26 April 2012
© Springer Science+Business Media, LLC 2012

Abstract

Purpose The intravenous administration of α -Galactosylceramide (α -GalCer)-pulsed antigen presenting cells (APCs) is well tolerated and the increased IFN- γ producing cells in the peripheral blood after the treatment appeared to be associated with prolonged survival. An exploratory study protocol was designed with the preoperative administration of α -GalCer-pulsed APCs to clarify the mechanisms of these findings, while especially focusing on the precise tumor site.

Methods Patients with operable advanced lung cancer received an intravenous injection of α -GalCer-pulsed APCs before surgery. The resected lung and tumor infiltrating lymphocytes (TILs) as well as peripheral blood mononuclear cells

were collected and the invariant NKT (iNKT) cell-specific immune responses were analyzed.

Results Four patients completed the study protocol. We observed a significant increase in iNKT cell numbers in the TILs and augmented IFN- γ production by the α -GalCer-stimulated TILs.

Conclusion The administration of α -GalCer-pulsed APCs successfully induced the dramatic infiltration and activation of iNKT cells in the tumor microenvironment.

Keywords Invariant NKT cell · antigen presenting cell · immunotherapy · tumor infiltrating lymphocyte · non-small cell lung cancer

K. Nagato · S. Motohashi · F. Ishibashi · K. Okita · K. Yamasaki ·
T. Nakayama
Department of Immunology, Graduate School of Medicine, Chiba
University, Clinical Research Center, Chiba University Hospital,
1-8-1 Inohana, Chuo-ku,
Chiba 260-8670, Japan

H. Hanaoka
Graduate School of Medicine, Chiba University,
Clinical Research Center, Chiba University Hospital,
1-8-1 Inohana, Chuo-ku,
Chiba 260-8670, Japan

S.-i. Fujii
Research Unit for Cellular Immunotherapy,
RIKEN Research Center for Allergy and Immunology,
1-7-22, Suehiro-cho, Tsurumi-ku,
Yokohama, Japan

K. Nagato · S. Motohashi (✉) · F. Ishibashi · Y. Moriya ·
H. Hoshino · S. Yoshida · I. Yoshino
General Thoracic Surgery, Graduate School of Medicine, Chiba
University, Clinical Research Center, Chiba University Hospital,
1-8-1 Inohana, Chuo-ku,
Chiba 260-8670, Japan
e-mail: motohashi@faculty.chiba-u.jp

M. Taniguchi
Laboratory for Immune Regulation,
RIKEN Research Center for Allergy and Immunology,
1-7-22, Suehiro-cho, Tsurumi-ku,
Yokohama, Japan

Introduction

V α 24 invariant natural killer T (V α 24 iNKT) cells are a unique innate lymphocyte subpopulation characterized by the expression of a canonical invariant T cell receptor with a specific α -chain gene rearrangement (V α 24-J α 18) and pairing mostly with a V β 11 β -chain in human. Synthetic glycolipid, α -Galactosylceramide (α -GalCer) is a mouse and human iNKT cell ligand, presented by a monomorphic class I-like antigen presenting molecule CD1d [1–3]. Ligand activated iNKT cells exhibit both direct and indirect potent anti-tumor activity.

Patients with malignant diseases show either a decreased number or functionally impaired V α 24 iNKT cells in human peripheral blood mononuclear cells (PBMCs) [4–9]. Head and neck cancer patients with poor circulating iNKT cell number show significantly worse clinical outcomes, suggesting an important contribution of iNKT cells to anti-tumor responses [10]. In addition, the ability to produce IFN- γ from circulating iNKT cells in cancer patients is preserved even though the absolute number of iNKT cells decreases, and thus, residual iNKT cells might still have a good competence to exert anti-tumor responses. Therefore, the expansion and activation of these cells *in vivo* may be therapeutically meaningful in patients with severely decreased or functionally deficient V α 24 iNKT cells. Clinical studies of α -GalCer-pulsed antigen presenting cells (APCs) have been conducted to recover a functionally sufficient number of V α 24 iNKT cells [11–14]. A phase I/II study of α -GalCer-pulsed APCs in patients with advanced or recurrent non-small cell lung cancer (NSCLC) found that the treatment elicits V α 24 iNKT cell-dependent immune responses, which are correlated with prolonged overall survival time [13]. The mechanisms that underlie this positive clinical outcome are still unclear.

The current clinical trial focused on the iNKT cell-specific immunological responses in the tumor microenvironments to investigate further anti-tumor mechanisms of V α 24 iNKT cells after α -GalCer-pulsed APC treatment. Therefore, in this exploratory study, the preoperative administration of α -GalCer-pulsed APCs was performed to clarify the iNKT cell specific immune responses at the tumor site more precisely. The results indicated that α -GalCer-pulsed APCs successfully induced the activation of tumor infiltrating V α 24 iNKT cells in the lung.

Material and Methods

Patient Eligibility Criteria

The study included patients between 20 and 80 years of age, with a diagnosis of clinical stage IIB or IIIA NSCLC that was to be treated surgically. Further inclusion criteria were a

performance status of 0, 1, or 2; normal or near normal renal, hepatic and hematopoietic function; and no chemotherapy or radiotherapy received for at least 4 weeks before enrollment. V α 24⁺V β 11⁺ iNKT cells were detected by flow cytometry in the enrolled patients at a level of >10 cells in 1 ml peripheral blood. The exclusion criteria were a positive response to HIV, hepatitis C virus, or human T-cell lymphotropic virus antibodies; positive for hepatitis B antigen; the presence of active inflammatory disease or active autoimmune disease; a history of hepatitis; pregnancy or lactation; concurrent corticosteroid therapy and evidence for another active malignant neoplasm. The α -GalCer-pulsed APC non-treatment cases were investigated as a control group to elucidate the effects of α -GalCer-pulsed APC treatment. The inclusion and exclusion criteria of the control group were the same as for the treatment group. The histological type, tumor-node-metastasis classification and the anti-tumor effect of treatment were classified according to the general rules for the clinical and pathologic recording of lung cancer as described by the Japan Lung Cancer Society.

Clinical Protocol and Study Design

The study was carried out in the Department of Chest Surgery, Chiba University Hospital, Japan, according to the standards of Good Clinical Practice for Trials on Medicinal Products in Japan. The protocol was approved by the Institutional Ethics Committee (No. 1972). In addition, this trial underwent ad hoc reviews by the Chiba University Quality Assurance Committee on Cell Therapy.

The study design is illustrated in Fig. 1. Written informed consent was obtained from all of the patients before undergoing a screening evaluation to determine eligibility. Clinical and laboratory assessments were conducted once a week, including of a complete physical examination and standard laboratory values. Any adverse events and changes in laboratory values were graded according to the National Cancer Institute Common Toxicity Criteria version 4.0.

Preparation of APCs from Peripheral Blood

All procedures were carried out according to the Good Manufacturing Practice standards. Eligible patients underwent peripheral blood leukapheresis (COBE Spectra, Gambro BCT,




Fig. 1 Study design of α -GalCer-pulsed APC administration. The patients received α -GalCer-pulsed APCs. The timing for both apheresis and α -GalCer-pulsed APC administration are shown. APC, α -GalCer-pulsed APC administration

Inc., Lakewood, CO) and PBMCs were collected and further separated by density gradient centrifugation (OptiPrep, Nycomed Amersham, Oslo, Norway). Thereafter, whole PBMCs were cultured with GM-CSF and IL-2, as previously described [11, 15]. Briefly, PBMCs were washed three times and resuspended in AIM-V (Invitrogen Corp., Carlsbad, CA) with 800 units/ml of human granulocyte macrophage colony-stimulating factor (GeneTech Co., Ltd., China) and 100 Japanese reference units per milliliter of recombinant human IL-2 (Imunace, Shionogi, Osaka, Japan). The cultured cells were pulsed with 100 ng/ml of specific ligand, α -GalCer (KRN7000; Kirin Brewery, Gunma, Japan) on the day before administration. Whole cells were harvested after 7 days of cultivation, washed 3 times and resuspended in 100 ml of 2.5 % albumin in saline. The patients received an intravenous injection of the cultured cells once (Fig. 1). The criteria for α -GalCer-pulsed APC administration included a negative bacterial culture 48 h before APC injection, cell viability >70 % and an endotoxin test 48 h before APC injection with a result <0.7 Ehrlich units/ml. The patients were injected with 1×10^9 cells/m²/injection of APCs.

Phenotype Evaluation of APCs

The phenotypes of α -GalCer-pulsed APCs were determined using a FACSCalibur flow cytometer (BD biosciences). The monoclonal antibodies (mAb) used were FITC-labeled anti-HLA-DR, CD83, CD14; phycoerythrin-labeled anti-CD86, CD1d; and allophycocyanin-labeled anti-CD11c, CD40 (Becton Dickinson, San Diego, CA). Isotype-matched control mAbs were used as negative controls.

Preparation of Tumor Infiltrating Lymphocytes, Tumor Cells, Normal Lung Mononuclear Cells and Lymph Nodes Mononuclear Cells

Fresh tumor tissue specimens were obtained from the surgical specimens and the tissue was cut into small pieces with scissors. The tissue specimen was placed in a flask with a mixture of 0.1 mg/ml DNase type I, 1 mg/ml collagenase type IV and 0.5 mg/ml hyaluronidase type V (all from Sigma, St. Louis, MO) in RPMI 1640 and stirred at room temperature for 1 h. The resultant cell suspension was washed in HBSS and subjected to two-layered (75 and 100 %) Ficoll-Hypaque discontinuous density gradient centrifugation at 1200 g for 20 min. The cells from the 100 % interface and 75 % interface were used as tumor infiltrating lymphocytes (TILs) and tumor cells, respectively. Normal lung tissue and lymph nodes were excised from the surgical specimen, cut with scissors in RPMI 1640 containing enzymes and passed through a gauze filter. The resultant cell suspension was washed in HBSS and subjected to Ficoll-Hypaque gradient centrifugation. The

interface was collected and used as either a normal lung or lymph node.

Immunological Monitoring

PBMC samples were obtained at least twice before APC administration and 1 week after APC injection.

Flow Cytometric Analysis of V α 24⁺V β 11⁺ Inkt Cells in the Peripheral Blood and TILs

The cell concentrations of V α 24⁺V β 11⁺ iNKT cells in PBMCs, TILs and mononuclear cells from normal lung tissue or lymph node were assessed by flow cytometry. Mononuclear cells were three-color stained with FITC-conjugated anti-T-cell receptor (TCR) V α 24 mAb (C15; Immunotech, Marseilles, France), phycoerythrin-conjugated anti-TCR V β 11 mAb (C21, Immunotech) and APC-conjugated anti-CD3 mAb (UCTH1; BD Bioscience). The stained cells were subjected to flow cytometry and the percentages of V α 24⁺V β 11⁺CD3⁺ cells among mononuclear cells were calculated. Thereafter, the number of iNKT cells (counts/ml) was estimated based on the PBMC counts.

Single-Cell Enzyme-Linked Immunospot Assay

PBMCs, TILs and cells from normal lung tissue or lymph nodes were washed 3 times with PBS and then were stored in liquid nitrogen until use. IFN- γ -secreting cells were assayed in 96-well filtration plates (Millipore, Bedford, MA) coated with mouse anti-human IFN- γ (10 μ g/ml; Mabtech, Nacka Strand, Sweden). The cells (5×10^5 per well) were incubated for 16 h with or without α -GalCer (100 ng/ml) in 10%FCS containing RPMI. Phorbol 12-myristate 13-acetate (10 μ g/ml) plus ionomycin (10 nmol/l) was used as a positive control. After culture, the plates were washed and incubated with biotinylated anti-IFN- γ (1 μ g/ml; Mabtech). Spot-forming cells were quantified by microscopy.

Quantitative Real Time PCR of V α 24 Invariant TCR and CD1d Expression

Total RNA was extracted from the tumors, normal lung tissue and lymph nodes using TRIzol Reagent (Sigma Aldrich) and reverse transcribed using Superscript II RT (Invitrogen Life Technologies) and oligo (dT12–18) primers (Invitrogen Life Technologies). The primers specific for the constant region of TCR α chain (C α) (sense, CGCCTTCAA CAACAGCATTA; antisense, ACCAGCTTGACATCA CAGGA), TCR V α 24 (sense, GCAAAGCTCTCT GCACATCA; antisense, CCAGGGTTGAGCCTCTGTC), CD1d (sense,gtcaggaagtgcgaactga; antisense, atcctgagacatggcacacc) were used with 5 μ g of sample cDNA and

amplified with *Taq* polymerase (Promega). Quantitative real-time PCR was performed using real-time Taq-Man technology and an ABI PRISM 7000 sequence detector (Applied Biosystems, Foster City, CA). The expression was normalized using the $C\alpha$ signal for $V\alpha 24$ and GAPDH for CD1d.

Statistical Methods

Statistical analyses were performed using Student's *t*-test.

Results

Patient Characteristics

A total of 4 patients met the inclusion criteria and were enrolled in the study. The patient characteristics are summarized in Table I. The study included one patient with adenocarcinoma and three patients with squamous cell carcinoma. Two patients were stage IIb and two were stage IIIa primary lung cancer. No patients had received any previous treatments.

In addition, a total of 6 patients who had not received α -GalCer-pulsed APC injection were enrolled as the control group. Fresh tumor tissue, normal lung tissue and lymph nodes were excised from the surgical specimens. The patient characteristics of the control group are also listed in Table I.

Phenotypes of α GalCer-Pulsed APCs

The phenotypes of α GalCer-pulsed APCs prepared for administration were analyzed by flow cytometry. All profiles for each patient are shown in Fig. 2. The percentages of HLA-DR⁺, CD11c⁺, CD86⁺, CD40⁺, CD83⁺ and CD1d⁺ cells were determined by the overtone subtraction test using the population comparison platform in the FlowJo software package. More than 50 % of the cultured cells were HLA-DR⁺ cells, 10 % to 50 % were CD11c⁺ cells and 50 % to

80 % were CD86⁺ cells. Interestingly, the majority of the cultured cells were CD3⁺ T cells or CD56⁺CD3⁻ NK cells, indicating the expression of HLA-DR⁺, CD11c⁺ or CD86⁺ on human T cells or NK cells (data not shown). Some variations were observed in the expression of CD83 (23.9–55.4 %), CD40 (8.5–17.1 %) and CD1d (23.1–69.9 %; Fig. 2).

Adverse Events

No serous (grade >2) toxicity or severe side effects were observed in any patients.

Immunological Monitoring of PBMCs and Resected Specimens

Immunological assays were conducted for all patients. The frequency of peripheral blood $V\alpha 24$ iNKT cells in all patients was measured by FACS. Figure 3 shows that two patients (cases 002 and 004) showed an increased number of circulating $V\alpha 24$ iNKT cells after the α -GalCer-pulsed APC administration. No clear relationship was found between the number of circulating $V\alpha 24$ iNKT cells and the α -GalCer-pulsed APC administration in the remaining two patients (cases 001 and 003).


Immunological assays were also performed for TILs and mononuclear cells (MNC)s from normal lung and lymph node tissues. TILs from all 4 cases in the α -GalCer-pulsed APC administration group contained a high percentage of $V\alpha 24$ iNKT cells in comparison to the normal lung MNCs (TILs; 1.86 %, 0.32 %, 0.15 % and 0.39 % vs. lung MNCs; 0.031 %, 0.013 %, 0.003 % and 0.01 %, Fig. 4a). The frequency of $V\alpha 24$ iNKT cells in the TILs in case 001 was 60 times higher than the normal lung MNCs. Though the content of $V\alpha 24$ iNKT in the normal lung MNCs was extremely low in case 003, the $V\alpha 24$ iNKT cells were found to have accumulated in the TILs. The average percentage of $V\alpha 24$ iNKT cells in the TILs was 50 times higher than that

Table I Patient characteristics of α -GalCer-pulsed APC group and control group

Case	Treat ^a	Age/Sex	Histology	c-stage	Operation method
001	APC ^b	75/M	Ad ^d	T2N1M0 (stage IIB)	Lobectomy+LND ^g
002	APC	76/M	Sq ^e	T2N1M0 (stage IIB)	Lobectomy+LND
003	APC	74/M	Sq	T1N2M0 (stage IIIA)	Lobectomy+LND
004	APC	68/M	Sq	T3N1M0 (stage IIIA)	Pneumonectomy+LND
c-01	cont ^c	71 M	Sq	T2N1M0 (stage IIB)	Lobectomy+LND
c-02	cont	55 M	large ^f	T2N1M0 (stage IIB)	Lobectomy+LND
c-03	cont	70 M	Sq	T3N0M0 (stage IIB)	Lobectomy+LND
c-04	cont	72 M	Sq	T3N1M0 (stage IIIA)	Bilobectomy+LND
c-05	cont	56/M	Sq	T2N1M0 (stage IIB)	Lobectomy+LND
c-06	cont	63/M	Ad	T2N2M0 (stage IIIA)	Lobectomy+LND

^aTreat, Treatment; ^bAPC, α -GalCer-pulsed APC administration; ^c cont, control; ^dAd, Adenocarcinoma; ^e Sq, Squamous cell carcinoma; ^f large, large cell carcinoma; ^g LND, Lymph Node dissection

Fig. 2 Flow cytometric analysis of α -GalCer-pulsed APCs. The expression levels of HLA-DR, CD11c, CD86, CD83, CD1d and CD40 were assessed by flow cytometry. Shaded areas: background staining with an iso-type control. Solid lines: staining profiles of the indicated molecules. Values represent the percentages of positive cells




in normal lung MNCs. The $V\alpha 24$ iNKT cell frequency in the draining lymph nodes of each case was almost the same as that in the normal lung MNCs (Fig. 4a).

The proportion of $V\alpha 24$ iNKT cells in the control group showed a relatively high percentage of TILs in comparison to the normal lung MNCs (TILs; 0.031 %, 0.058 %, 0.13 %, 0.47 %, 0.18 % and 0.12 % vs. lung MNCs; 0.034 %, 0.011 %, 0.004 %, 0.039 %, 0.014 % and 0.02 %,

Fig. 4b). The average percentage of $V\alpha 24$ iNKT cells in the TILs was only 8 times higher than that in the normal lung MNCs.

Normal lung MNCs in the control group demonstrated a trend toward a higher $V\alpha 24$ iNKT cell rate in comparison to the treatment group (Fig. 4c). On the other hand, the proportion of $V\alpha 24$ iNKT cells in TILs tended to increase in the α -GalCer-pulsed APC injected group in comparison to the

Fig. 3 Immunological monitoring of PBMCs of patients with α -GalCer-pulsed APC administration. The absolute number of peripheral blood iNKT cells ($V\alpha 24^+V\beta 11^+$ cells) and NK cells ($CD56^+CD3^-$ cells). Flow cytometric analysis and automated full blood counts (Chiba University Hospital) indicated the absolute number of $V\alpha 24$ iNKT cells and NK cells. APC, α -GalCer-pulsed APC administration; ope., operation






Fig. 4 The frequencies of Vα24 iNKT cells in TILs and mononuclear cells from normal lung and lymph node tissues. **a**, The proportion of Vα24 iNKT cells (Vα24⁺Vβ11⁺ cells) in PBMCs on day 14, TILs, normal lung MNCs and lymph node MNCs in the α-GalCer-pulsed APC administration group were assessed by flow cytometry. The lymph node MNCs column depicts one representative MNC profile in the draining lymph nodes including hilar (#10, 11 and 12) and mediastinal (#1, 3, 4 and 7) nodes. **b**, The proportion of Vα24 iNKT cells in PBMCs, TILs and normal lung MNCs in the control group were assessed by flow cytometry. **c-d**, The comparison between the Vα24 iNKT cell contents in normal lung MNCs **c** and TILs **d** in the α-GalCer-pulsed APC treatment group and the control group. **e** The TIL/Normal Lung MNC ratio of Vα24 iNKT cell proportion. control, control group; Tx, α-GalCer-pulsed APC administration group; **p*=0.0008

control group (mean percentage, 0.68 % and 0.165 %, Fig. 4d). The Vα24 iNKT cell ratio of TILs/normal lung in the α-GalCer-pulsed APC administration group was significantly higher than that of the control group (*p*=0.0008, Fig. 4e).

The number of IFN-γ-producing cells after restimulation with α-GalCer in vitro was concurrently monitored in PBMCs, TILs, normal lung MNCs and lymph node MNCs using an ELISPOT assay. An analysis of the PBMCs and resected specimen showed the highest value of α-GalCer-responsive IFN-γ-producing cell number in the TILs of the α-GalCer-pulsed APC treated group (Fig. 5a). The absolute number of α-GalCer-responsive IFN-γ-producing cells in the TILs was apparently high in cases 001 and 002, whereas a relatively low value was seen in cases 003 and 004. This observation was not detected with the use of control group specimens (Fig. 5b). Together with the results in Fig. 4, the administration of α-GalCer-pulsed APCs induced the mobilization of endogenous Vα24 iNKT cells into the primary site of the lung cancer and augmented the IFN-γ-producing ability of tumor infiltrating Vα24 iNKT cells.

In addition, the number of IFN-γ-producing cells in PBMCs was determined after restimulation with α-GalCer




Fig. 5 Detection of α-GalCer-reactive IFN-γ-producing cells by enzyme-linked immunospot assay. **a** Cryopreserved PBMCs, TILs, normal lung MNCs and lymph node MNCs of the α-GalCer-pulsed APC treated group were thawed and cultured overnight with either α-GalCer or vehicle. The presence of IFN-γ-producing cells was quantified by an enzyme-linked immunospot assay. The resected draining

lymph nodes including hilar (#10, 11 and 12) and mediastinal (#1, 3, 4 and 7) nodes are shown. Spot number of IFN-γ with standard deviation for triplicate culture of 4 cases are shown. **b** Spot-forming cell number in the control group. SFC, Spot Forming Cell; pre, pretreatment; d14, day 14; Nor, normal lung MNCs; n.d., not done; #11 s, lymph node #11 superior; #11i, lymph node #11 inferior




Fig. 5 (continued)

in vitro. In Fig. 5a, the number of IFN- γ producing cells in PMBC increased 14 days after treatment in case 002 and 004, indicating that global NKT cell activation in these patients.

The mRNA Expression Level of V α 24 Inkt Cell Receptor and CD1d


Quantitative RT-PCR was performed to further confirm the increase in the number of V α 24 iNKT cells. The mRNA from the primary tumor, normal lung tissue and lymph node samples was obtained from the α -GalCer-pulsed APC treated patients and the relative expression level of V α 24 invariant TCR and the constant region of TCR (C α) mRNA were evaluated. The mRNA for V α 24 TCR was highly expressed in the tumor (Fig. 6a). The V α 24-J α 18 invariant TCR mRNA could not be detected in the normal lung tissue since the lung parenchyma included mainly alveolar epithelial cells and only a low copy number of C α mRNA was detected. The relative gene expression of CD1d was also evaluated by quantitative RT-PCR. CD1d expression was ascertained both in normal lung tissue and tumor tissue in all

4 cases and the expression level appeared to be higher in normal lung tissue (Fig. 6b).

Discussion

The major aim of this study was to investigate the V α 24 iNKT cell-specific immune responses in the primary tumor site after the intravenous injection of α -GalCer-pulsed APCs in patients with advanced NSCLC. The development of these immunotherapeutic approaches with the administration of α -GalCer-pulsed APCs requires a thorough knowledge of local immune responses. V α 24 iNKT cells accumulate in lung cancer lesions [5], as observed in this report (Fig. 4e). A significant increase in the tumor infiltrating V α 24 iNKT cell population was detected after the administration of α -GalCer-pulsed APCs in comparison to the non-injected control group. This observation is quite reasonable for the iNKT cell-targeted therapy aimed at activation of V α 24 iNKT cells in the tumor located site in vivo. The activation status of CTLs, rather than just the existence

Fig. 6 The relative mRNA expression of the V α 24⁺ TCR in tumor, normal lungs and lymph nodes. Cancer tissue, non-cancerous lung tissue and lymph nodes were obtained from α -GalCer-pulsed APC treated patients. **a** The expression level of V α 24 TCR mRNA in each sample was analyzed by quantitative RT-PCR. Each mRNA was quantified by the standard curve method and copy numbers of V α 24 TCR were normalized by the copy number of the constant region of the TCR α chain (C α mRNA. **b** The expression level of CD1d mRNA in each sample was analyzed by quantitative RT-PCR. Each mRNA was quantified by the standard curve method and copy numbers of CD1d were normalized by the copy number of the GAPDH mRNA. LN, lymph node; Normal, normal lung tissue; Tumor, tumor tissue; n.d., not detected; #11 s, lymph node #11 superior; #11i, lymph node #11 inferior



of CTLs, has great prognostic significance [16–18]. Therefore, IFN- γ -producing cells were also monitored in PBMCs, TILs and MNCs from normal lung tissue and lymph nodes using an ELISPOT assay. After starting the protocol, standard preoperative chemo-radiotherapy was introduced to treat locally advanced NSCLC, and such a change of treatment strategy hindered the entry of patients into this protocol. In spite of the limited number of patients analyzed, the results obtained by ELISPOT assay indicated that tumor infiltrating mononuclear cells had augmented IFN- γ

producing capacity, which may have a positive impact on the tumor microenvironment.

Tumor infiltrating lymphocytes are found in a variety of cancers and they are thought to be a result of a host immune response directed against tumor cells. Several reports have shown that the presence of large numbers of tumor-infiltrating CD8⁺ T cells are associated with a favorable prognosis in esophageal carcinoma [18, 19], colorectal cancer [20, 21], ovarian cancer [22], and pancreatic carcinoma [23], while the infiltration of CD4⁺ T cells that possess regulatory function,

such as Foxp3⁺ regulatory T cells (Treg) are associated with the poor prognosis of ovarian cancer [24, 25]. The balance between CD8⁺ CTLs and Tregs in tumors is critical for disease progression and survival [24, 26, 27]. These diverse results indicate that the functional roles of TILs are complicated and uncertain and the effects of TILs might vary with the type and stage of cancers. In addition to CD8⁺ cytotoxic T cells, tumor infiltrating V α 24 iNKT cells have been reported to be a positive prognostic factor for colorectal carcinoma [28]. The current results indicated that the injection of α -GalCer-pulsed APCs could induce the accumulation of V α 24 iNKT cells in TILs, which would therefore lead to a good prognosis after a complete surgical resection.

Although a complete surgical resection is regarded as the optimal treatment for NSCLC, only around 25 % of NSCLC are suitable for potentially curative resection. Despite optimal surgical management, the 5-year survival rate of resected NSCLC ranges between 85.9 % for pathological stage Ia and 41 % for pathological stage IIIa [29]. Approximately 50 % or more of patients with NSCLC who undergo surgery experience relapse due to the existence of microscopic lesions that could not be detected by preoperative screening. Recently, adjuvant chemotherapy given after surgery has been shown to improve survival [30–32]. A meta-analysis suggested that cisplatin-based adjuvant chemotherapy could yield an absolute overall survival advantage of 5 % at 5 years [33]. At the same time, chemotherapeutic agents often show severe toxic effects and it was reported that in patients with early-stage disease have deleterious effects on long-term survival. This emphasizes the importance of development of less-invasive preoperative or postoperative therapy to suppress the growth of micrometastases. Therefore, immune cells for tumor surveillance, such as NK and iNKT cells, which possess anti-tumor activity, should be beneficial and post-surgical adjuvant immunotherapy by the use of these cells may be favorable since the residual tumor is quite small after a complete resection.

Conclusions

α -GalCer-pulsed APC administration successfully induced the dramatic infiltration and activation of V α 24 iNKT cells in the tumor lesion. This report is the first clinical trial of V α 24 iNKT cell targeted immunotherapy that shows a functional V α 24 iNKT cell accumulation in the tumor microenvironment. These results encourage the further development of immunotherapy aimed at the activation of endogenous V α 24 iNKT cells in the lung.

Acknowledgements We thank Kyowa Hakko Kirin Co. Ltd. for providing clinical grade α -GalCer (KRN7000) for these studies. We also thank all the nurses and staff surgeons in the Department of the Thoracic Surgery, Chiba University Hospital, Chiba, Japan, for their excellent help with patient care and continuous support.

This work was supported by Global COE Program (Global Center for Education and Research in Immune System Regulation and Treatment), City Area Program (Kazusa/Chiba Area) MEXT (Japan), and by grants from the Ministry of Education, Culture, Sports, Science and Technology (Japan) (Grants-in-Aid: for Scientific Research on Priority Areas #17016010; Scientific Research [B] #21390147, and Scientific Research [C] #21591808, and Cancer Translational Research Project), the Ministry of Health, Labor and Welfare (Japan), Uehara Memorial Foundation, Mochida Foundation, Chiba Foundation for Health Promotion and Disease Prevention and Mitsui Life Social Welfare Foundation.

Conflict of interest The authors declare that they have no conflict of interest.

References

- Kawano T, Cui J, Koezuka Y, Toura I, Kaneko Y, Motoki K, et al. CD1d-restricted and TCR-mediated activation of V α 14 NKT cells by glycosylceramides. *Science*. 1997;278:1626–9.
- Brossay L, Chioda M, Burdin N, Koezuka Y, Casorati G, Dellabona P, et al. CD1d-mediated recognition of an α -galactosylceramide by natural killer T cells is highly conserved through mammalian evolution. *J Exp Med*. 1998;188:1521–8.
- Spada FM, Koezuka Y, Porcelli SA. CD1d-restricted recognition of synthetic glycolipid antigens by human natural killer T cells. *J Exp Med*. 1998;188:1529–34.
- Tahir SM, Cheng O, Shaulov A, Koezuka Y, Bublej GJ, Wilson SB, et al. Loss of IFN- γ production by invariant NK T cells in advanced cancer. *J Immunol*. 2001;167:4046–50.
- Motohashi S, Kobayashi S, Ito T, Magara KK, Mikuni O, Kamada N, et al. Preserved IFN- γ production of circulating V α 24 NKT cells in primary lung cancer patients. *Int J Cancer*. 2002;102:159–65.
- Molling JW, Kolgen W, van der Vliet HJ, Boomsma MF, Kruizenga H, Smorenburg CH, et al. Peripheral blood IFN- γ -secreting V α 24⁺V β 11⁺ NKT cell numbers are decreased in cancer patients independent of tumor type or tumor load. *Int J Cancer*. 2005;116:87–93.
- van der Vliet HJ, Nishi N, Koezuka Y, von Blomberg BM, van den Eertwegh AJ, Porcelli SA, et al. Potent expansion of human natural killer T cells using α -galactosylceramide (KRN7000)-loaded monocyte-derived dendritic cells, cultured in the presence of IL-7 and IL-15. *J Immunol Methods*. 2001;247:61–72.
- Yanagisawa K, Seino K, Ishikawa Y, Nozue M, Todoroki T, Fukao K. Impaired proliferative response of V α 24 NKT cells from cancer patients against α -galactosylceramide. *J Immunol*. 2002;168:6494–9.
- Dhodapkar MV, Geller MD, Chang DH, Shimizu K, Fujii S, Dhodapkar KM, et al. A reversible defect in natural killer T cell function characterizes the progression of premalignant to malignant multiple myeloma. *J Exp Med*. 2003;197:1667–76.
- Molling JW, Langius JA, Langendijk JA, Leemans CR, Bontkes HJ, van der Vliet HJ, et al. Low levels of circulating invariant natural killer T cells predict poor clinical outcome in patients with head and neck squamous cell carcinoma. *J Clin Oncol*. 2007;25:862–8.
- Ishikawa A, Motohashi S, Ishikawa E, Fuchida H, Higashino K, Otsuji M, et al. A phase I study of α -galactosylceramide (KRN7000)-pulsed dendritic cells in patients with advanced and recurrent non-small cell lung cancer. *Clin Cancer Res*. 2005;11:1910–7.
- Motohashi S, Ishikawa A, Ishikawa E, Otsuji M, Iizasa T, Hanaoka H, et al. A phase I study of in vitro expanded natural killer T cells

- in patients with advanced and recurrent non-small cell lung cancer. *Clin Cancer Res.* 2006;12:6079–86.
13. Motohashi S, Nagato K, Kunii N, Yamamoto H, Yamasaki K, Okita K, et al. A phase I-II study of α -galactosylceramide-pulsed IL-2/GM-CSF-cultured peripheral blood mononuclear cells in patients with advanced and recurrent non-small cell lung cancer. *J Immunol.* 2009;182:2492–501.
 14. Motohashi S, Nakayama T. Invariant natural killer T cell-based immunotherapy for cancer. *Immunotherapy.* 2009;1:73–82.
 15. Ishikawa E, Motohashi S, Ishikawa A, Ito T, Uchida T, Kaneko T, et al. Dendritic cell maturation by CD11c⁻ T cells and V α 24⁺ natural killer T-cell activation by α -galactosylceramide. *Int J Cancer.* 2005;117:265–73.
 16. Grabenbauer GG, Lahmer G, Distel L, Niedobitek G. Tumor-infiltrating cytotoxic T cells but not regulatory T cells predict outcome in anal squamous cell carcinoma. *Clin Cancer Res.* 2006;12:3355–60.
 17. van Beek J, zur Hausen A, Snel SN, Berkhof J, Kranenbarg EK, van de Velde CJ, et al. Morphological evidence of an activated cytotoxic T-cell infiltrate in EBV-positive gastric carcinoma preventing lymph node metastases. *Am J Surg Pathol.* 2006;30:59–65.
 18. Schumacher K, Haensch W, Roefzaad C, Schlag PM. Prognostic significance of activated CD8(+) T cell infiltrations within esophageal carcinomas. *Cancer Res.* 2001;61:3932–6.
 19. Cho Y, Miyamoto M, Kato K, Fukunaga A, Shichinohe T, Kawarada Y, et al. CD4⁺ and CD8⁺ T cells cooperate to improve prognosis of patients with esophageal squamous cell carcinoma. *Cancer Res.* 2003;63:1555–9.
 20. Menon AG, Janssen-van Rhijn CM, Morreau H, Putter H, Tollenaar RA, van de Velde CJ, et al. Immune system and prognosis in colorectal cancer: a detailed immunohistochemical analysis. *Lab Invest.* 2004;84:493–501.
 21. Prall F, Duhrkop T, Weirich V, Ostwald C, Lenz P, Nizze H, et al. Prognostic role of CD8⁺ tumor-infiltrating lymphocytes in stage III colorectal cancer with and without microsatellite instability. *Hum Pathol.* 2004;35:808–16.
 22. Zhang L, Conejo-Garcia JR, Katsaros D, Gimotty PA, Massobrio M, Regnani G, et al. Intratumoral T cells, recurrence, and survival in epithelial ovarian cancer. *N Engl J Med.* 2003;348:203–13.
 23. Ryschich E, Notzel T, Hinz U, Autschbach F, Ferguson J, Simon I, et al. Control of T-cell-mediated immune response by HLA class I in human pancreatic carcinoma. *Clin Cancer Res.* 2005;11:498–504.
 24. Sato E, Olson SH, Ahn J, Bundy B, Nishikawa H, Qian F, et al. Intraepithelial CD8⁺ tumor-infiltrating lymphocytes and a high CD8⁺/regulatory T cell ratio are associated with favorable prognosis in ovarian cancer. *Proc Natl Acad Sci U S A.* 2005;102:18538–43.
 25. Curiel TJ, Coukos G, Zou L, Alvarez X, Cheng P, Mottram P, et al. Specific recruitment of regulatory T cells in ovarian carcinoma fosters immune privilege and predicts reduced survival. *Nat Med.* 2004;10:942–9.
 26. Alvaro T, Lejeune M, Salvado MT, Bosch R, Garcia JF, Jaen J, et al. Outcome in Hodgkin's lymphoma can be predicted from the presence of accompanying cytotoxic and regulatory T cells. *Clin Cancer Res.* 2005;11:1467–73.
 27. Gao Q, Qiu SJ, Fan J, Zhou J, Wang XY, Xiao YS, et al. Intratumoral balance of regulatory and cytotoxic T cells is associated with prognosis of hepatocellular carcinoma after resection. *J Clin Oncol.* 2007;25:2586–93.
 28. Tachibana T, Onodera H, Tsuruyama T, Mori A, Nagayama S, Hiai H, et al. Increased intratumor V α 24-positive natural killer T cells: a prognostic factor for primary colorectal carcinomas. *Clin Cancer Res.* 2005;11:7322–7.
 29. Sawabata N, Miyaoka E, Asamura H, Nakanishi Y, Eguchi K, Mori M, et al. Japanese lung cancer registry study of 11,663 surgical cases in 2004: demographic and prognosis changes over decade. *J Thorac Oncol.* 2011;6:1229–35.
 30. Arriagada R, Bergman B, Dunant A, Le Chevalier T, Pignon JP, Vansteenkiste J. Cisplatin-based adjuvant chemotherapy in patients with completely resected non-small-cell lung cancer. *N Engl J Med.* 2004;350:351–60.
 31. Winton T, Livingston R, Johnson D, Rigas J, Johnston M, Butts C, et al. Vinorelbine plus cisplatin vs. observation in resected non-small-cell lung cancer. *N Engl J Med.* 2005;352:2589–97.
 32. Douillard JY, Rosell R, De Lena M, Carpagnano F, Ramlau R, Gonzales-Larriba JL, et al. Adjuvant vinorelbine plus cisplatin versus observation in patients with completely resected stage IB-IIIa non-small-cell lung cancer (Adjuvant Navelbine International Trialist Association [ANITA]): a randomised controlled trial. *Lancet Oncol.* 2006;7:719–27.
 33. Pignon JP, Tribodet H, Scagliotti GV, Douillard JY, Shepherd FA, Stephens RJ, et al. Lung adjuvant cisplatin evaluation: a pooled analysis by the LACE Collaborative Group. *J Clin Oncol.* 2008;26:3552–9.

薬理と治療 (JPT)

JAPANESE PHARMACOLOGY & THERAPEUTICS (国際文献略号 *Jpn Pharmacol Ther*)

2012年6月20日発行 vol.40 no.6 別刷

● パネルディスカッション3 /

これからの時代を担う IRB の機能と責務—中央 IRB (共同 IRB を含む) の普及に向けて— / 講演 4

中央 IRB 等への移行過程で生じた課題と その解決に向けた取り組み

花岡英紀 青柳玲子 松本和彦 吉澤弘久 小池竜司
大学病院臨床試験アライアンス推進室および中央 IRB 検討ワーキンググループ



ライフサイエンス出版

TEL (03) 3664-7900 (代表)

【禁 無断転載・複製】

● パネルディスカッション 3 /

これからの時代を担う IRB の機能と責務—中央 IRB (共同 IRB を含む) の普及に向けて— / 講演 4

中央 IRB 等への移行過程で生じた課題と その解決に向けた取り組み

花岡 英紀¹⁾ 青柳 玲子¹⁾ 松本 和彦²⁾ 吉澤 弘久³⁾ 小池 竜司⁴⁾
大学病院臨床試験アライアンス推進室および中央 IRB 検討ワーキンググループ

大学病院臨床試験アライアンスではいわゆる中央 IRB のあり方について検討を行うとともに、千葉大学、信州大学、新潟大学による中央 IRB の審議を 2010 年 7 月より試験的に一試験について開始した。本稿では大学病院臨床試験アライアンスで実施している中央 IRB 検討ワーキンググループ (以下 WG) およびその上部組織である推進室での議論、さらに欧州の中央 IRB のあり方をふまえた IRB のあるべき姿と、実際の取り組みについて述べるものである。また、この内容については、筆者らの私見にもとづいており、各大学でのコンセンサスを得られた段階に至っていない。

はじめに

近年、治験の活性化の中で、中央 IRB に向けた取り組みの必要性が謳われている。これは、多くの場合、治験を開始するにあたり、それぞれの IRB への申請に関して多くの時間と労力を必要とするため、その業務に当たる費用が高額となり、治験依頼者への負担が増しているという事実がある。また、一方で、経験の乏しい IRB において複雑な治験の審議をすることについて問題視する事実もある。このような状況のなか、GCP では 2 度の改訂を経ているにもかかわらず、いまだ中央 IRB の活動は活発ではない。平成 23 年 6 月には日本医師会治験促進センターによる治験等適正化作業班による提言がなされ、これを受け形でさらなる GCP の改訂がなされた。作業班では、中央 IRB の設置と IRB 事務局と治験事務局の外部委託による効率化も検討された。

大学病院臨床試験アライアンスは、国際共同試験をわが国で主導することを目的として関東信越 7 大学において先進的な取り組みを治験において実施してきた。そのなかで中央 IRB については、はたして実施にあたりどのような課題があるか具体的に検討

が必要との結論に至り、推進室の下部組織として信州大学を担当校とし各大学の担当者からなる WG を構成し議論を重ねてきた。

WG の検討内容について

そもそも中央 IRB の設置の前提としていくつかの疑問がある。治験を実施する各施設個別の状況を知らない中央 IRB において、①迅速性と効率性と、科学性と倫理性は共に成り立つにはどのようにしたらいいのか、②被験者を保護するという原則はどのようにしたら確保されるのか、という 2 課題は WG の検討以前においてそもそも重要な問題であり、IRB の審議の根幹を成すものである。中央 IRB で十分な議論ができる体制を維持すること、さらにそこでは、その施設固有の問題点を被験者保護という立場から話し合える状況にする必要があると考えている。

一方で、③審査手続きの共通化、④中央 IRB の設置場所、⑤具体的な審議の方法、という 3 課題については、技術的な側面が強い課題であり、解決可能な課題とした。

欧米の中央 IRB のあり方について

海外には中央 IRB がすでに多く設置され、活用法

1) 千葉大学 2) 信州大学 3) 新潟大学 4) 東京医科歯科大学

英国	3 機関の承認が必要 1.NHS National Health Service 行政区ごとの local Research Ethical Committee (REC) 2.Main REC 3.MHRA (Medicines and healthcare products Regulatory Agency)
フランス	被験者保護委員会 CCP 被験者保護に関する事項 臨床試験のプロトコルデザインに関する事項 施設に関する事項 医療製品保険安全局 AFSSAPS 試験薬等の品質と安全性に関する事項 被験者の安全性に関する事項 (適格基準、投与量、被験者のモニタリング、対照薬)
米国	All committees are registered with the Office for Human Research Protections (OHRP). FDA, pharmaceutical companies, contract research organizations, as well as AAHRPP perform regular inspections of our processes. An FDA inspection in July 2009 resulted in no findings. CERTIFICATION: Staff and board members have earned certifications, such as Certified IRB Professional (CIP), Certified Clinical Research Professional (CCRP) and Certified IRB Manager (CIM).

図1 欧米の中央IRBに関する大学病院臨床試験アライアンスによる訪問調査

IRB 開催日	IRB 審議資料		
	千葉大	新潟大	信州大
2010/7/20	新規実施の適否	新規実施の適否 (治験参加カード)	新規実施の適否 (治験参加カード)
2010/9/21	実施計画書 2.0 版 同意説明文書	実施計画書 2.0 版	実施計画書 2.0 版
2010/10/18		同意説明文書	
2010/11/15	実施計画書 2.1 版 同意説明文書 監査手順書 2.0 版	実施計画書 2.1 版 同意説明文書 監査手順書 2.0 版	実施計画書 2.1 版 同意説明文書 監査手順書 2.0 版
2011/2/21	院内の重篤有害事象	-	-
2011/3/22	継続審査、安全性情報	継続審査、安全性情報	継続審査、安全性情報
2011/4/18	実施計画書 2.2 版 治験薬概要書 2.0 版 同意説明文書	実施計画書 2.2 版 治験薬概要書 2.0 版	実施計画書 2.2 版 治験薬概要書 2.0 版
2011/5/16	同意説明文書	同意説明文書	同意説明文書
2011/9/20	実施計画書 2.3 版 同意説明文書 緊急脱退報告書	実施計画書 2.3 版 同意説明文書	実施計画書 2.3 版 同意説明文書

図3 3大学から千葉大学中央IRBへ提出され審議された内容

れている国々があるが、その設置の基準や役割はさまざまである。海外の状況 (6) を検討するためにわれわれは実際に規制当局や医療機関を訪問し、調査を行った (図1)。このなかで、施設固有の問題点を審議するローカルIRBと計画書の審議をする中央IRBがそれぞれ業務を分担し審議を行っていることを知った。さらに、規制当局との連携も重要な点であった。また、米国ではコマーシャルIRBが数多く設置されているが、それぞれのIRBではその質の高さと独立性を示すための取り組み (認定の取得、業務内容の公開、FDAの調査を受けた時期の公開) を行っていることが明らかになったが、一方で施設固有の問題点を審議することについては難しい状況であることが判明した。

医師主導治験の中央IRBについて

2010年7月千葉大学医学部附属病院治験審査委員

千葉大学、新潟大学、信州大学で実施中の中央IRB体制

手順

- 1) 治験実施の適否についての審査
中央IRB (千葉大IRB) にて審議 (施設固有事項含む) ③④
- 2) 治験の継続についての審査 (重篤な有害事象発生時など) ①②③④ (信州)




図2 WGで検討された中央IRBのフロー

会では、信州大学医学部附属病院長および新潟大学医歯薬総合病院長から千葉大学医学部附属病院長への医師主導治験の審議依頼を受け審議を開始した。これは国立大学病院としては初めての取り組みである。この審議を実施するにあたり3大学ではWGで検討された中央IRBの審議のフロー (図2) をもとにそれぞれ手順書の改訂 (7) 契約書の作成とこれに伴う審査費用の算定 (8) を行った。ここでは申請元の大学のIRBを「施設で発生した重篤有害事象を審議する専門IRB」と位置づけている。(9)3大学から中央IRB事務局へ提出される書類は内容がすべて同一でないことより中央IRBで承認された内容や時期が異なっている (図3)。このことは治験事務局と中央IRB事務局との独立性を意味し、推進する立場とこれを監督する立場のそれぞれの事務局の独立性が確保された状況となっている。(10) 医師会の作業班の提案する各大学の治験事務局を廃止し中央に委託することについては、作業の効率性に加え、技術的に千葉大学の事務局が各大学の業務を引き受けることは困難との結論に至った。

まとめ

およそ10の課題について検討を行いながら、具体的に中央IRBを運用するに至った。規制当局との連携については今までまったく議論をされてい内容である。つまり、その本質において単なる効率化ではなく真の意味での中央IRBの役割を考えることが重要であり、欧米での取り組みを十分考慮のうえ、進めて行く必要がある。大学病院臨床試験アライアンスではわが国を代表するグループとして欧米からも信頼される中央IRB体制の構築を実施する。

[原著] 脊髄障害性疼痛に対する顆粒球コロニー
刺激因子 (G-CSF) の治療効果

加藤 啓 山崎 正志 大河 昭彦 佐久間 毅
高橋 宏 橋本 光宏 林 浩一 川辺 純子
藤由 崇之 古矢 丈雄 山内 友規 門田 領
宮下 智大 萬納寺 誓人 染谷 幸男 鎌田 尊人
池田 修 橋本 将行 井上 雅俊 花岡 英紀¹⁾
国府田 正雄 高橋 和久

(2011年8月29日受付, 2011年9月28日受理)

要 旨

脊髄障害性疼痛を有する13例を対象として、顆粒球コロニー刺激因子 (G-CSF) を用いた神経保護療法の臨床試験を施行した。対象症例は、脊髄症の急性増悪と関連する疼痛を認めた6例 (脊髄症急性増悪群) および脊髄症の後遺症と考えられる疼痛を認めた7例 (脊髄症後遺症群) の二群に分けられた。G-CSFを5 μ g/kg/日 \times 5日間 (1例), 10 μ g/kg/日 \times 5日間 (11例), 10 μ g/kg/日 \times 3日間 (1例) の投与量・期間で点滴静注投与し、投与前後での疼痛の変化をVisual analogue scale (VAS) にて評価した。11例で投与後に疼痛が軽減し、2例では完全に消失した。疼痛軽減効果は全例、投与開始から1週間以内で認められた。脊髄症後遺症群では無効例が2例存在した。VASは、脊髄症急性増悪群で投与前平均63.3mmから投与後1週間で平均25.0mm ($P<0.05$) に、脊髄症後遺症群で投与前平均67.1mmから投与後1週間で平均44.3mm ($P<0.05$) に減少した。投与後6ヵ月間の観察期間中に7例で疼痛軽減効果の減弱が見られた。効果の減弱は投与後3ヵ月時に4例、6ヵ月時に3例で認められた。今回の検討から、G-CSFが脊髄障害性疼痛に対して疼痛軽減効果を有することが示唆された。

Key words: 脊髄障害性疼痛, 顆粒球コロニー刺激因子, 神経保護療法

千葉大学大学院医学研究院整形外科

¹⁾千葉大学医学部附属病院臨床試験部

Kei Kato, Masashi Yamazaki, Akihiko Okawa, Tsuyoshi Sakuma, Hiroshi Takahashi, Mitsuhiro Hashimoto, Koichi Hayashi, Junko Kawabe, Takayuki Fujiyoshi, Takeo Furuya, Tomonori Yamauchi, Ryo Kadota, Tomohiro Miyashita, Chikato Mannoji, Yukio Someya, Takahito Kamada, Osamu Ikeda, Masayuki Hashimoto, Masatoshi Inoue, Hideki Hanaoka¹⁾, Masao Koda and Kazuhisa Takahashi: Therapeutic effects of granulocyte-colony stimulating factor for neuropathic pain in patients with spinal cord lesions.

Department of Orthopedic Surgery, Graduate School of Medicine, Chiba University, Chiba 260-8670.

¹⁾Clinical Research Center, Chiba University Hospital, Chiba 260-8677.

Tel. 043-226-2117. Fax. 043-226-2116. E-mail: masashiy@faculty.chiba-u.jp

Received August 29, 2011, Accepted September 28, 2011.

I. 緒言

脊髄障害性疼痛とは、2009年以降に本邦で使用されるようになった新しい概念である。厳密な定義は定められてはいないが、ヘルニア・脊柱靭帯骨化症などによる圧迫性脊髄症や、急性脊髄損傷などの脊髄障害に起因すると考えられる多彩な痛みを総称している[1,2]。侵害受容性疼痛に比べ神経障害性疼痛の占める割合が大きいのが特徴であり、通常では痛みを起こさない程度の刺激が強い痛みを引き起こすアロディニア（異痛症）や、障害脊髄高位より尾側の締め付けられるような自発痛などに代表される。非ステロイド性抗炎症薬（NSAIDs）の効果は低く、最近では末梢神経障害性疼痛で得られたエビデンスをもとに、プレガバリンなどの抗てんかん薬や一部の抗うつ薬などの有効性が解明されてきたが[3-5]、多くの症例で必ずしも十分な除痛効果が得られるわけではない。病態が十分に解明されておらず、症例によって薬剤に対する反応性が異なるため、治療に難渋することがしばしばである。本邦では平成22年度に厚生労働省により脊髄障害性疼痛症候群が研究奨励分野に指定され、研究が進められている。

我々は、2008年から急性期脊髄損傷患者および圧迫性脊髄症急性増悪患者に対する顆粒球コロニー刺激因子（Granulocyte-colony stimulating factor: G-CSF）を用いた神経保護療法の臨床試験を進めている[6-9]。G-CSFは血球系に作用する増殖因子であり、顆粒球系細胞の分化・増殖・生存促進などの作用を有する[10]。本邦ではがん化学療法による好中球減少症や、末梢血幹細胞移植時の造血幹細胞の末梢血への動員などの目的で臨床使用されている。中枢神経系においては、骨髄細胞を脳・脊髄中へ動員する作用や[11]、脳卒中モデルに対する神経保護作用などが報告されており[12]、海外では脳梗塞に対する臨床試験が報告されている[13]。以上より、脊髄損傷に対してもG-CSFが治療効果を発揮しうる可能性が想定されたため、われわれはマウスおよびラットの脊髄損傷モデルを用いてG-CSFの有効性およびその作用機序につき報告してきた[14-18]。その結果、G-CSF投与により後肢運動機能が有意に改善した。その機序に関しては、①G-CSFによ

り動員された骨髄由来幹細胞が脊髄損傷部に生着する、②直接的に神経細胞死を抑制する、③Oligodendrocyteの細胞死を抑制し、髄鞘を保護する、④炎症性サイトカイン（TNF- α 、IL-1 β ）発現を抑制する、⑤血管新生を促進する、などが想定された[14-18]。これらの根拠から、慢性圧迫性脊髄症の急性増悪に対してもG-CSFが神経保護作用を有する可能性が示唆されたため、圧迫性脊髄症急性増悪患者に対する治療薬としてのG-CSFの安全性・有効性を確認するため臨床試験を開始した[6,7]。結果として、G-CSFの安全性が確認され、全例で神経症状の改善が認められた。さらに、予期せぬ効果として、G-CSF投与後に脊髄障害に起因する疼痛が軽減した例を少なからず経験した[19]。このことからG-CSFが脊髄障害性疼痛に対して何らかの疼痛軽減効果を有することが示唆された。今回は、臨床試験例におけるG-CSF投与後の脊髄障害性疼痛の変化を検討した。

II. 方法

2009年1月から2010年2月の期間に、脊髄障害に関連する疼痛を認めた16例に対し、G-CSFを用いた神経保護療法の臨床試験を施行した。今回は、その中で投与から6ヵ月以上経過観察しえた13例を対象として検討した（表1）。症例1～6では、後縦靭帯骨化症（OPLL）および黄色靭帯骨化症（OYL）による圧迫性脊髄症の急性増悪（日本整形外科学会頸髄症治療判定基準 [JOAスコア] が直近の1ヵ月で2点以上悪化）に対してG-CSFの投与が行われた。この6例での脊髄障害性疼痛は、脊髄症の急性増悪との関連が強く示唆されたため、脊髄症急性増悪群として分類し検討を進めた。G-CSFの投与量・期間は、症例1で5 μ g/kg/日 \times 5日間、症例2～6で10 μ g/kg/日 \times 5日間の点滴静注であった。6例全例で、G-CSF投与後1.5～8週で脊髄症に対する後方除圧固定術または椎弓切除術が行われた。

症例7では、頸椎症性脊髄症に伴うと考えられる疼痛が持続しており、症例8～13では、脊髄症に対する手術の術後に脊髄関連の疼痛が遺残していた。したがって、症例7～13を脊髄症後遺症群

表1 脊髄障害性疼痛に対するG-CSF投与症例

症例 No.	年齢 (歳) /性別	診断	脊髄症 急性増悪	責任高位	G-CSF 投与量 (μg/kg/日)	G-CSF 投与期間 (日)	手術術式 ^a [範囲]	G-CSF 投与～手術 (週)
1	68/男	後縦靱帯骨化症	+	T4/5	5	5	後方除圧固定 [T1-8]	1.5
2	46/男	後縦靱帯骨化症	+	T6/7	10	5	後方除圧固定 [T2-11]	8
3	75/男	後縦靱帯骨化症	+	C4/5	10	5	後方除圧固定 [C2-7]	7
4	64/男	後縦靱帯骨化症	+	C4/5	10	5	後方除圧固定 [C2-7]	2
5	32/男	後縦靱帯骨化症	+	T8/9	10	5	後方除圧固定 [T7-12]	4
6	67/男	黄色靱帯骨化症	+	T11/12	10	5	椎弓切除 [T10-12]	5
7	73/男	頸椎症性脊髄症	-	C4/5	10	5	-	-
8	58/男	後縦靱帯骨化症術後	-	C5/6	10	5	-	-
9	72/男	脊髄円錐症候群術後	-	T12/L1	10	5	-	-
10	71/男	後縦靱帯骨化症術後	-	C5/6	10	5	-	-
11	78/男	黄色靱帯骨化症術後	-	T10/11	10	5	-	-
12	70/女	後縦靱帯骨化症術後	-	C4/5	10	5	-	-
13	38/女	胸髄腫瘍術後	-	T11/12	10	3	-	-

a: G-CSF投与後の手術, T: Thoracic vertebrae (胸椎)/C: Cervical vertebrae (頸椎)

と分類して検討を進めた。G-CSFの投与量・期間は、症例7～12で10μg/kg/日×5日間の点滴静注であった。症例13でも同一条件でG-CSFの投与を開始したが、3日間の投与後に尿路感染に伴う発熱が生じたため、投与が中止となった。

疼痛部位と障害脊髄の位置関係から、脊髄障害性疼痛を2型に分類した。すなわち、障害髄節高位の痛みおよび障害髄節から2髄節以内の領域の痛みをat-level pain、髄節より3髄節以上尾側の領域の痛みをbelow-level painと定義した。疼痛の程度は、Visual analogue scale (VAS: 0～100mm)を用いて評価した。脊髄症の評価には、JOAスコア(頸髄症: 0～17点, 胸髄症: 0～11点)を用いた。上下肢の筋力に評価には、American Spinal Injury Association (ASIA) 運動スコア(0～100点)、上下肢体幹の感覚の評価にはASIA痛覚スコア(0～112点)を用いた。

統計学的評価にはフィッシャーの正確確率検定とウィルコクソンの符号付順位検定を用い、 $P < 0.05$ を有意差ありとした。

Ⅲ. 結 果

脊髄症急性増悪群6例における疼痛は、at-level painが4例、below-level painが2例であり、有痛期間は0.2～4年(平均1.5年)であった(表2)。脊髄症後遺症群7例における疼痛は、at-level painが3例、below-level painが4例であり、有痛期間は1～27年(平均10.9年)であった(表2)。脊髄症後遺症群における有痛期間が、有意に長かった($P < 0.01$)。

脊髄症急性増悪群では6例全例で10mm以上のVASの低下が認められ、投与後1週間以内に除痛効果が得られていた(表3A)。完全に疼痛が消失した例も1例(症例5)存在した。VASは投与前平均63.3mmから投与後1週間で平均25.0mmに減少した($P < 0.05$)。除痛効果は3例で投与後3ヵ月時に減弱が見られ、1例(症例3)では投与直前の疼痛に戻った。投与後6ヵ月の時点でのVASは平均33.3mmであった($P < 0.05$)。

脊髄症後遺症群では7例中5例で10mm以上のVASの低下が認められた(表3B)。完全に疼痛が消失した例も1例(症例11)認められた。無

表2 脊髄障害性疼痛の性状

症例 No.	疼痛の分類	疼痛の部位	疼痛の性質	有痛期間 (年)
1	below-level	両下肢全体	しびれ感の強い痛み	0.7
2	at-level	胸背部	絞め付けられるような痛み	0.8
3	below-level	両下腿	しびれ感の強い痛み	3
4	at-level	右手	しびれ感の強い痛み	0.3
5	at-level	背部 右胸部	しびれ感の強い痛み 刺すような痛み	4
6	at-level	両鼠径部	しびれ感の強い痛み	0.2
7	below-level	両足底部	しびれ感の強い痛み	13
8	at-level	左手指	しびれ感の強い痛み	1
9	below-level	両足底部	刺すような痛み	3
10	at-level	両手指 両下肢全体	しびれ感の強い痛み	3
11	below-level	両大腿内側	しびれ感の強い痛み	19
12	at-level	両前腕以下	灼熱感を伴う痛み	27
13	below-level	左下肢全体	しびれ感の強い痛み	10

表3 A G-CSF投与後のVASの変化 (脊髄症急性増悪群)

症例 No.	投与直前	VAS				VAS減少値 ^a			
		1週	1ヵ月	3ヵ月	6ヵ月	1週	1ヵ月	3ヵ月	6ヵ月
1	50	25	25	25	25	25	25	25	25
2	60	30	30	30	20	30	30	30	40
3	50	30	30	50	50	20	20	0	0
4	50	25	25	45	45	25	25	5	5
5	80	0	0	0	0	80	80	80	80
6	90	40	40	60	60	50	50	30	30
平均	63.3	25.0*	25.0*	35.0*	33.3*	38.3	38.3	28.3	30.0
± SD	± 17.5	± 13.4	± 13.4	± 21.4	± 22.3	± 22.9	± 22.9	± 28.4	± 28.8

VAS: 0 ~ 100mm

a: 投与直前のVASからの減少値

*: 投与直前に比し有意に減少 ($P < 0.05$)

表3 B G-CSF投与後のVASの変化 (脊髄症後遺症群)

症例 No.	投与直前	VAS				VAS減少値 ^a			
		1週	1ヵ月	3ヵ月	6ヵ月	1週	1ヵ月	3ヵ月	6ヵ月
7	70	70	70	70	70	0	0	0	0
8	50	40	40	40	50	10	10	10	0
9	90	90	90	90	90	0	0	0	0
10	80	50	50	50	50	30	30	30	30
11	60	0	0	60	60	60	60	0	0
12	60	20	20	20	30	40	40	40	30
13	60	40	40	40	60	20	20	20	0
平均	67.1	44.3*	44.3*	52.9	58.6	22.9	22.9	14.3	8.6
± SD	± 13.8	± 29.9	± 29.9	± 22.9	± 18.6	± 22.1	± 22.1	± 16.2	± 14.6

VAS: 0 ~ 100mm

a: 投与直前のVASからの減少値

*: 投与直前に比し有意に減少 ($P < 0.05$)

表 4 A G-CSF 投与後の JOA スコアの変化 (脊髄症急性増悪群)

症例 No.	JOA スコア (点)				改善率 (%)	
	投与 1 ヶ月前	投与直前	1 ヶ月後	6 ヶ月後	1 ヶ月後	6 ヶ月後
1	8.5/11	1.5/11	6/11	8/11	47.4	68.4
2	7.5/11	5.5/11	6.5/11	10/11	18.2	81.8
3	16/17	8.5/17	14.5/17	14.5/17	70.6	70.6
4	14/17	9.5/17	14.5/17	14.5/17	66.7	66.7
5	6/11	4/11	6/11	6/11	28.6	28.6
6	6/11	4/11	6.5/11	6.5/11	35.7	35.7
平均					44.5*	58.6*
± SD					± 21.0	± 21.3

JOA スコア: 日本整形外科学会頸髄症治療判定基準 (頸髄症: 0~17点, 胸髄症: 0~11点)
*: 脊髄症後遺症群 (表 4 B) と比べ有意に改善 ($P < 0.01$)

表 4 B G-CSF 投与後の JOA スコアの変化 (脊髄症後遺症群)

症例 No.	JOA スコア (点)				改善率 (%)	
	投与 1 ヶ月前	投与直前	1 ヶ月後	6 ヶ月後	1 ヶ月後	6 ヶ月後
7	13/17	13/17	13/17	13/17	0	0
8	5/11	5/11	6/11	6/11	16.7	16.7
9	5.5/11	5.5/11	5.5/11	5.5/11	0	0
10	4.5/17	4.5/17	8/17	8/17	28	28
11	4/11	4/11	6/11	6/11	28.6	28.6
12	11/17	11/17	11/17	11/17	0	0
13	3/11	3/11	3/11	3/11	0	0
平均					10.5	10.5
± SD					± 13.6	± 13.6

JOA スコア: 日本整形外科学会頸髄症治療判定基準 (頸髄症: 0~17点, 胸髄症: 0~11点)

効例が 2 例 (症例 7, 9) あり, 両者とも below-level pain の例であった。有効例では全例, 1 週間以内に除痛効果が得られていた。VAS は投与前平均 67.1mm から投与後 1 週間で平均 44.3mm に減少した ($P < 0.05$)。一方, 除痛効果は 1 例で投与後 3 ヶ月時に, 3 例で投与 6 ヶ月時に減弱が見られ, 経過観察期間中に 3 例で投与直前の疼痛に戻った。投与後 3 ヶ月の時点での VAS は平均 52.9mm と再度上昇していた。

JOA スコアは脊髄症急性増悪群の 6 例全例で G-CSF 投与後に増加が見られた (表 4 A)。投与後 1 ヶ月および 6 ヶ月時の改善率はそれぞれ平均 44.5%, 58.6% であった。急性増悪前と比較しても遜色ない程度まで回復していた。脊髄症後遺症群では 7 例中 3 例で JOA スコアの増加を認めたが, 投与後 1 ヶ月および 6 ヶ月時の改善率はいずれも平均 10.5% に過ぎなかった (表 4 B)。除痛効果のなかった 2 例 (症例 7, 9) では, JOA スコ

アの改善は得られていなかった。

脊髄症急性増悪群の ASIA スコアは, 運動, 痛覚とともに投与後 1 ヶ月および 6 ヶ月時で, 投与直前と比べ有意に改善を認めた (表 5 A)。一方, 脊髄症後遺症群の ASIA スコアは改善を認めなかった (表 5 B)。無効例の 2 例 (症例 7, 9) では, 運動, 痛覚スコアともに, 増加点数は 0 であった。

末梢血中の白血球数は投与開始後 1 日目から有意に増加した (表 6) が, 投与終了後速やかに減少し, 1 週間の時点では投与前の値に戻っていた。白血球の分画中, 顆粒球が著明に増加していたが, 単球の増加も有意であった。CRP も有意に増加していたが, 軽度の増加であり臨床的に問題となる値ではなかった。リンパ球数は投与前後で変化を認めなかった。投与期間中, および投与後に重篤な有害事象の発生は認めなかった。