

What's new?

MicroRNAs (miRNAs) play critical roles in carcinogenesis and may be valuable therapeutic targets for malignant disease. Here, the miRNA tumor suppressor *miR-29c* was found to be significantly downregulated in gastric cancer cells and to be reactivated by the selective cyclooxygenase-2 (COX-2) inhibitor celecoxib. Reactivation led to *miR-29c*-induced suppression of the anti-apoptotic protein Mcl-1. The data reveal that selective COX-2 inhibitors may have clinical promise for the treatment of gastric cancer via restoration of *miR-29c*.

which is commonly overexpressed in solid tumors including gastric cancer, and COX-1, which is expressed constitutively in normal tissues, suggesting that inhibition of COX-2 may reduce the risk of cancer development. Clinical trials have revealed that the selective COX-2 inhibitor celecoxib is effective for prevention of colorectal adenomas.^{15,16} Recent studies have also shown that celecoxib has a preventive effect against *Helicobacter pylori*-associated gastric cancer.^{17,18} Therefore, celecoxib has been suggested to have promise for the treatment of gastrointestinal tumors. However, the molecular mechanisms underlying the chemopreventive effects of selective COX-2 inhibitors are not fully understood.

To identify miRNAs that play critical roles in the development and progression of gastric cancer, we examined the miRNA expression profiles of gastric tumors including gastric adenomas and both early and advanced gastric cancers. We also investigated miRNAs that are targeted by celecoxib. Our findings revealed that *miR-29c* was downregulated in gastric cancers and activated by celecoxib. Recent studies have shown that *miR-29c* is downregulated in various human malignancies including gastric cancer and acts as a tumor suppressor.^{19–25} Herein, we show that the putative tumor suppressor *miR-29c* is an important miRNA in gastric carcinogenesis and is a potential therapeutic target for gastric cancer.

Material and Methods**Patients and tissue specimens**

A total of 53 clinical samples of gastric tumors were examined. The clinicopathological features of the patients are shown in Supporting Information Table 1. The average age of the patients was 67.7 years (male/female, 39/14). Thirty patients with gastric adenomas (atypical epithelium) or early gastric cancers underwent ESD at Keio University Hospital (Tokyo, Japan). This study was approved by the research ethics committee of Keio University School of Medicine (#19-68-5) and registered with the UMIN Clinical Trials Registry (UMIN 000001057). Informed consent was obtained from all patients before the examinations. Tissue specimens from gastric adenomas and early gastric cancers and the surrounding nontumor gastric mucosae were obtained by endoscopic biopsy and kept in RNAlater (Ambion, Austin, TX) at -80°C until RNA extraction. Tissue specimens from advanced gastric cancers and the surrounding nontumor gastric mucosae were obtained from materials surgically resected from 23 patients at the National Cancer Center Hospital (Tokyo, Japan). This study was approved by the Ethics Commit-

tee of the National Cancer Center and performed in accordance with the 1964 Declaration of Helsinki. All patients gave their informed consent for inclusion in this study.

Cell lines and treatment with celecoxib

The human gastric cancer cell lines AGS and MKN45 were used in this study. AGS was obtained from the American Type Culture Collection (Rockville, MD), and MKN45 was obtained from the Japan Health Science Foundation (Osaka, Japan). Cells were cultured in RPMI1640 medium supplemented with 10% fetal bovine serum and seeded at 1×10^5 cells per 100-mm dish 24 hr prior to treatment. Cells were treated with celecoxib (Pfizer, New York, NY) at 30 μM for 48 hr.

RNA extraction and microarray analysis

Total RNAs were extracted from both tissue specimens of gastric tumors resected by ESD and gastric cancer cell lines using the mirVana miRNA isolation kit (Ambion). Total RNA of tissue specimens from advanced gastric cancers and matched nontumor gastric mucosae was extracted using TRIzol reagent (Invitrogen, Carlsbad, CA).

Five hundred nanograms of total RNA from each of ten advanced gastric cancers were pooled, and the same was done with 10 matched nontumor gastric mucosae. miRNA microarray analysis of samples of advanced gastric cancer was conducted by Toray Industries (www.toray.com: Tokyo, Japan). miRNA microarray analysis of AGS cells treated with celecoxib was conducted by LC Sciences (www.lcsciences.com: Houston, TX). All data were submitted to the ArrayExpress database, under the accession numbers E-MEXP-2230 (gastric cancer) and E-MEXP-2231 (celecoxib). These microarray chips contain probe regions that detect 724 (Toray Industries) and 711 (LC Sciences) miRNA transcripts listed in Sanger miRBase Release 10.0 (<http://www.sanger.ac.uk>). These chips contain multiple probes for each miRNA (2 probes in the Toray Industries chip, 5 probes in the LC Sciences chip), and the average values of their signal intensities are shown in Tables 1 and 3.

Quantitative RT-PCR of *miR-29c*

Levels of miRNA expression were analyzed by quantitative RT-PCR using the TaqMan microRNA assay for *miR-29c* (Applied Biosystems, Foster City, CA) in accordance with the manufacturer's instructions. Expression levels were normalized to that of U6 RNA.

Table 1. Summary of the most downregulated miRNAs in advanced gastric cancers relative to non-tumorous gastric mucosae

No.	miRNAs	Non-tumor	Tumor	Fold change
1	<i>miR-193b</i>	1133.8	563.0	0.5
2	<i>miR-768-3p</i>	756.8	372.1	0.5
3	<i>miR-140-3p</i>	1438.6	690.7	0.5
4	<i>miR-923</i>	43834.8	20277.2	0.5
5	<i>miR-939</i>	3780.0	1655.2	0.4
6	<i>miR-487b</i>	173.0	72.2	0.4
7	<i>miR-378*</i>	95.3	39.6	0.4
8	<i>miR-29c</i>	1733.9	641.1	0.4
9	<i>miR-133b</i>	424.6	155.0	0.4
10	<i>miR-768-5p</i>	1499.5	518.1	0.3

Note: Data for Non-tumor and Tumor are average values of the signal intensities in microarray analysis. Fold change represents the ratio of the signal intensities for Tumor/Non-tumor.

In situ hybridization (ISH) of miR-29c

Locked nucleic acid (LNA)-modified probes for *miR-29c* were used (miRCURY-LNA detection probe, Exiqon, Vedbaek, Denmark). *In situ* hybridization was performed using the RiboMap *in situ* hybridization kit (Ventana Medical Systems, Tucson, AZ) on the Ventana Discovery automated *in situ* hybridization instrument (Ventana Medical Systems). *In situ* hybridization steps after deparaffinization were performed based on the standard protocol provided in the manufacturer's RiboMap application note (<http://www.ventanamed.com>). Hematoxylin-eosin (HE) was used for counterstaining.

Immunohistochemical examination of Mcl-1

Formalin-fixed and paraffin-embedded tissues were deparaffinized and rehydrated. For antigen retrieval, the sections were treated for 10 min at 105°C in an autoclave, and nonspecific reactions were blocked with a blocking reagent (Protein Block Serum-Free, Dako Cytomation, Glostrup, Denmark). The sections were incubated with the rabbit anti-human Mcl-1 polyclonal antibody (S-19, Santa Cruz Biotechnology, Santa Cruz, CA) overnight at 4°C followed by horseradish peroxidase labeled anti-rabbit IgG (Histofine, Simple stain MAX-PO, Nichirei, Tokyo, Japan) for 30 min at room temperature. Then, the sections were treated with 3,3'-diaminobenzidine tetrahydrochloride solution. All sections were counterstained with hematoxylin.

Immunoreactivity of Mcl-1 was confirmed in the germinal centers as described previously,²⁶ and this was defined as the internal positive control. The intensity of Mcl-1 immunoreactivity in each sample was graded as 0 (less than the internal positive control), 1 (equal to the internal positive control) and 2 (more than the internal positive control). Mcl-1 expression in randomly selected fields, two in gastric cancer tissues and two in nontumor gastric mucosal tissues, was examined in gastric adenomas and early ($n = 30$) and advanced ($n = 21$) gastric cancers at high magnification ($\times 400$). The Mcl-1 expression

score was determined as the product of the Mcl-1 immunoreactivity grade and the percentage of Mcl-1-positive cells.

Chromatin immunoprecipitation (ChIP) assay

The ChIP assay was performed as described previously.⁴ An antibody against C/EBP α (sc-61, Santa Cruz Biotechnology) was used. Quantitative analysis was performed by real-time PCR with the CYBR Premix Ex Taq (Takara Bio, Ohtsu, Japan) using the Thermal Cycler Dice Real-Time System (Takara Bio). The sequences of the primers used were as follows; Forward: 5'-CTAAGAGCAGACTGATGGTGTC-3', Reverse: 5'-CTATTTCTGTTGACTCCTAGCAGC-3'.

The fraction of immunoprecipitated DNA was calculated as follows: [immunoprecipitated DNA with C/EBP α antibody-nonspecific antibody control (NAC)]/(input DNA - NAC).

Western blotting

Protein extracts were separated by SDS/polyacrylamide gel electrophoresis and transferred onto nitrocellulose membranes. The membranes were hybridized with a rabbit anti-human Mcl-1 polyclonal antibody (S-19, Santa Cruz Biotechnology), and β -actin was used as the internal control.

Transfection of miR-29c precursor molecules and the specific miR-29c inhibitor

The *miR-29c* precursor molecules, the specific *miR-29c* inhibitor and negative control precursor miRNAs were purchased from Ambion. They were transfected into AGS and MKN45 cells at a final concentration of 100 nM each using oligofectamine (Invitrogen) in accordance with the manufacturer's instructions. Forty-eight hours after transfection, the cells were collected and their expression of Mcl-1 was analyzed by Western blotting as described above.

Apoptosis assay

The levels of apoptosis in AGS and MKN45 cells transfected with *miR-29c* and treated with celecoxib were evaluated using Annexin V kit (Beckman Coulter, Brea CA) in accordance with the manufacturer's instructions.

Statistics

Data were analyzed using the SPSS statistics 17.0 software package. Differences in miRNA expression levels and apoptosis levels between groups were analyzed using unpaired *t* test. Differences at $p < 0.05$ were considered significant.

Results

Downregulation of miR-29c is associated with progression of gastric cancer

To investigate the miRNA expression profile in gastric cancer, we compared the miRNA expression of gastric cancer tissues with that of the corresponding nontumorous gastric mucosae. The microarray data shown in Table 1 revealed that the putative tumor suppressor *miR-29c* was significantly




Figure 1. The expression levels of *miR-29c* in the 53 clinical gastric tumor samples and matched nontumorous gastric mucosae. All reactions were performed in duplicate with U6 as the internal control. The graph shows the ratio of *miR-29c* expression of tumor/nontumor on a logarithmic scale (mean \pm SD). (a) Gastric adenomas (atypical epithelium, $n = 8$). There was no significant association of *miR-29c* expression between tumor and nontumor. (b) Early gastric cancers ($n = 22$). The expression levels of *miR-29c* in early gastric cancers were significantly lower than those in the corresponding nontumorous gastric mucosae ($p < 0.005$). (c) Advanced gastric cancers ($n = 23$). The expression levels of *miR-29c* in advanced gastric cancers were significantly lower than those in the corresponding nontumorous gastric mucosae ($p < 0.0005$). (d) A representative example of ISH for *miR-29c* in normal gastric tissue. *miR-29c* is expressed in the cytoplasm of normal gastric epithelial cells (blue staining) but not detected in other cells such as fibroblasts and muscle cells. HE was used for counterstaining. (e) A representative example of ISH for *miR-29c* in gastric cancer tissue. *miR-29c* expression was not detected in gastric cancer cells. HE was used for counterstaining.

downregulated in gastric cancer tissues relative to nontumorous gastric mucosae. To confirm the microarray data, we examined the levels of *miR-29c* expression in tissue specimens of gastric adenomas (atypical epithelium), early gastric cancers and advanced gastric cancers by quantitative RT-PCR. As shown in Figure 1a, the expression level of *miR-29c* was reduced in 75% (6 of 8) of gastric adenomas compared with the corresponding nontumor gastric mucosae. There was no significant difference in the average level of *miR-29c* expression between gastric adenomas and the corresponding nontumor gastric mucosae. The expression level of *miR-29c* was reduced in 73% (16 of 22) of early gastric cancers compared with the corresponding nontumor gastric mucosae (Fig. 1b), and the expression levels of *miR-29c* in early gastric cancers were significantly lower than those in the corresponding nontumorous gastric mucosae ($p < 0.005$). Furthermore, the expression level of *miR-29c* was reduced in 91% (21 of 23) of advanced gastric cancers relative to the corresponding nontumorous gastric tissues (Fig. 1c), and this reduction was shown to be significant ($p < 0.0005$). The relative expression levels of *miR-29c* in nontumorous gastric mucosa and gastric tumors and the ratio of *miR-29c* expression on a linear scale are shown in Supporting Information Figure 1.

To confirm the localization of *miR-29c* in gastric cancer tissues, ISH for *miR-29c* was performed. Figures 1d and 1e show representative examples of ISH for *miR-29c*. *miR-29c* is expressed in the cytoplasm of normal gastric epithelial cells (blue staining) but not detected in other cells such as fibroblasts and muscle cells. *miR-29c* expression was not detected in gastric cancer cells. These results are consistent with the data for quantitative RT-PCR of *miR-29c*.

We also examined the association between the ratio of *miR-29c* expression and clinicopathological features such as tumor differentiation, tumor location, *H. pylori* infection and clinical stage. We found that downregulation of *miR-29c* was more prominent in undifferentiated than in differentiated advanced gastric cancers (Table 2, Supporting Information Fig. 1). There was no significant association between the ratio of *miR-29c* expression and other clinicopathological features such as tumor location, *H. pylori* infection and clinical stage (Table 2).

Increased expression of Mcl-1, a target gene of miR-29c, in gastric cancer tissues

We next investigated the target genes of *miR-29c*. Mott et al.¹⁹ have demonstrated that Mcl-1, encoding an antiapoptotic Bcl2 family protein, is one of the targets of *miR-29* miRNAs, and that *miR-29* miRNAs regulate apoptosis by targeting Mcl-1. The levels of Mcl-1 expression in gastric cancer tissues were examined by immunohistochemistry. Immunoreactivity of the Mcl-1 antibody was confirmed by staining of the germinal centers, as described previously (Fig. 2a).²⁶ Prominent staining of Mcl-1 was observed in the cytoplasm of gastric cancer cells. Expression of Mcl-1 was markedly

Table 2. Association between the ratio of *miR-29c* expression and clinicopathological features of patients with gastric cancer

	N	Ratio of <i>miR-29c</i> expression (T/N)	p-value
a) Early gastric cancer			
Tumor differentiation			
Differentiated	19	0.75 ± 0.13	$p = 0.98$
Undifferentiated	3	0.76 ± 0.27	
Tumor location			
Upper and Middle portion	3	0.98 ± 0.14	$p = 0.23$
Lower portion	19	0.72 ± 0.13	
<i>H. pylori</i> infection			
Positive	10	0.88 ± 0.23	$p = 0.51$
Negative	10	0.71 ± 0.11	
b) Advanced gastric cancer			
Tumor differentiation			
Differentiated	14	0.54 ± 0.14	$p = 0.04^1$
Undifferentiated	9	0.22 ± 0.04	
Tumor location			
Upper and Middle portion	13	0.45 ± 0.15	$p = 0.65$
Lower portion	10	0.37 ± 0.08	
T stage			
T1, T2	9	0.58 ± 0.21	$p = 0.25$
T3, T4	14	0.31 ± 0.05	
Clinical stage			
I, II, III	13	0.30 ± 0.05	$p = 0.22$
IV	10	0.56 ± 0.19	

Note: The ratios of *miR-29c* expression (Tumor/Non-tumor) are expressed as mean ± SE. Differences between groups were analyzed by unpaired t test.

¹There was a significant correlation between the ratio of *miR-29c* expression and tumor differentiation.

increased in advanced gastric cancer (Figs. 2c and 2d), whereas in nontumorous gastric tissues only faint staining was evident (Fig. 2b). Mcl-1 expression was also increased in early gastric cancer (Fig. 2e).

The degree of Mcl-1 overexpression was represented as the ratio of the Mcl-1 expression score between the tumor and nontumorous mucosa (T/N). The average value of the ratio of Mcl-1 expression (T/N) in advanced gastric cancers was significantly higher than that in gastric adenomas and early gastric cancers (Fig. 2f).

miR-29c is activated by the selective COX-2 inhibitor celecoxib and suppresses Mcl-1

To identify miRNAs that are differentially expressed upon treatment of AGS gastric cancer cells with the selective COX-2 inhibitor celecoxib, we performed miRNA microarray




Figure 2. Expression levels of Mcl-1 in gastric cancer tissues examined by immunohistochemistry. (a) Immunoreactivity of the Mcl-1 antibody was confirmed by staining of the germinal centers as described previously.²⁶ Immunohistochemical examinations of Mcl-1 in nontumorous gastric tissues (b), advanced gastric cancer (c, d) and early gastric cancer (e) are shown. (f) The average levels of the ratio of the Mcl-1 expression score (T/N) in gastric cancer. AE: atypical epithelium; AGC: advanced gastric cancer; EGC: early gastric cancer. * $p < 0.05$.

analysis. As shown in Table 3, *miR-29c*, which is downregulated in gastric cancer tissues, was significantly activated after treatment with celecoxib.

Figure 3a shows the expression levels of *miR-29c* in AGS and MKN45 gastric cancer cells treated with celecoxib. The expression levels of *miR-29c* in AGS and MKN45 cells treated with celecoxib were significantly higher than those in untreated cells ($p < 0.05$), being consistent with the microarray data. We also examined the expression levels of *miR-29c* in AGS and MKN45 cells after treatment with celecoxib in the presence of the specific *miR-29c* inhibitor. We confirmed that increased expression of *miR-29c* by celecoxib

was cancelled after transfection with the specific *miR-29c* inhibitor.

We next examined the expression levels of Mcl-1, a target of *miR-29c*, by Western blotting in AGS and MKN45 cells after transfection with the *miR-29c* precursor molecules and treatment with celecoxib with or without the specific *miR-29c* inhibitor. We measured the expression levels of *miR-29c* after its transfection and confirmed that *miR-29c* expression was markedly increased (over 100-fold) after transfection. The results of Western blotting showed that the expression levels of Mcl-1 in AGS and MKN45 cells treated with celecoxib and transfected with the *miR-29c* precursor molecules were

Table 3. Summary of the most upregulated miRNAs in AGS cells treated with celecoxib

No.	miRNAs	Control	Celecoxib	Fold change
1	<i>miR-663</i>	206.8	935.1	4.5
2	<i>miR-181b</i>	90.4	395.9	4.4
3	<i>miR-29c</i>	497.0	1972.0	4.0
4	<i>miR-141</i>	122.5	410.3	3.3
5	<i>miR-149*</i>	285.8	894.4	3.1
6	<i>miR-98</i>	392.9	1113.9	2.8
7	<i>miR-26b</i>	593.4	1340.3	2.3
8	<i>miR-195</i>	279.4	626.5	2.2
9	<i>miR-148a</i>	333.1	707.6	2.1
10	<i>miR-638</i>	3682.7	7073.3	1.9

Note: Data for Control and Celecoxib are average values of the signal intensities in microarray analysis. Fold change represents the ratio of the signal intensities for Celecoxib/Control.

lower than in the control. We confirmed that suppression of Mcl-1 by celecoxib was cancelled after transfection with the specific *miR-29c* inhibitor, suggesting that celecoxib-induced suppression of Mcl-1 is mediated by *miR-29c* (Fig. 3b)

Celecoxib activates miR-29c expression by enhancing the DNA binding of C/EBP α to the promoter

The results of microarray analysis revealed that celecoxib activates the expression of *miR-29c*, but not that of *miR-29b*, suggesting that celecoxib may modulate the binding of transcriptional factors between *miR-29b* and *miR-29c* (Fig. 3c). Using the database (www.gene-regulation.com/pub/programs.html), we found that there are C/EBP α binding sites in the promoter region of *miR-29c*. Moreover, Wu *et al.* have reported that celecoxib induces the expression of the invasion-suppressor CRMP-1 by enhancing the binding of C/EBP α DNA to the promoter.²⁷ These findings prompted us to perform a ChIP assay with the C/EBP α antibody in AGS and MKN45 cells treated with celecoxib. As shown in Figure 3c, the results of the ChIP assay showed that immunoprecipitation with the C/EBP α antibody was significantly increased in AGS and MKN45 cells treated with celecoxib, indicating that celecoxib activates the expression of *miR-29c* by enhancing the binding of C/EBP α DNA to the promoter.

Celecoxib and miR-29c induce apoptosis in gastric cancer cells

Since Mcl-1 is an antiapoptotic factor and suppressed by *miR-29c* in gastric cancer cells, we performed annexin V-FITC apoptosis assay in AGS and MKN45 cells. Fluorescence microscopy and flow cytometry analyses demonstrated that annexin V staining (green staining on the cell surface membrane) was markedly increased in AGS and MKN45 cells 48 hr after *miR-29c* transfection and celecoxib treatment, indicating that *miR-29c* and celecoxib induce apoptosis in gastric cancer cells (Fig. 4).

Discussion

We focused our study on *miR-29c*, because it was the only miRNA that was highlighted by the results of the two microarray analyses. Although the extent of these analyses was intrinsically limited, the results suggested that *miR-29c* is downregulated in gastric cancer and reactivated by celecoxib.

Gastric adenomas and early gastric cancers are clinically similar gastric neoplasms, and sometimes they are difficult to diagnose only by endoscopic findings even with biopsied pathological examination. Therefore, the significant reduction of *miR-29c* expression in early gastric cancers, but not in gastric adenomas, suggests that *miR-29c* could be a novel molecular marker of early gastric cancer. Moreover, reduction of *miR-29c* expression is more prominent in advanced gastric cancers than in gastric adenomas and early gastric cancers. These findings suggest that reduction of *miR-29c* expression is critical for the progression of gastric cancer.

Downregulation of *miR-29c* has been reported in various human malignancies including nasopharyngeal carcinoma,²⁰ bladder transitional cell carcinoma,²¹ esophageal cancer,²² chronic lymphocytic leukemia^{23,24} and gastric cancer.²⁵ Mott *et al.*¹⁹ have demonstrated that Mcl-1, encoding an antiapoptotic Bcl2 family protein, is one of the targets of *miR-29* miRNAs, and that *miR-29* miRNAs regulate apoptosis by targeting Mcl-1. Mcl-1 is an antiapoptotic protein originally isolated from the ML-1 human myeloid leukemia cell line during cell differentiation.²⁸ The biological relevance of Mcl-1 as an antiapoptotic protein promoting cell survival has been reported in various human malignancies.²⁹⁻³¹ Elevated expression of Mcl-1 and its association with poor prognosis have been reported in gastric cancer.^{32,33} In accordance with these reports, our results demonstrate that *miR-29c* is downregulated in gastric cancers and suppresses its target oncogene, *Mcl-1*. In addition, downregulation of *miR-29c* and overexpression of Mcl-1 are associated with the progression of gastric cancer. These findings suggest that *miR-29c* functions as a tumor suppressor by suppressing Mcl-1 in gastric epithelial cells. We and other groups have previously shown that other miRNAs such as *miR-512-5p* and transcription factors also regulate *Mcl-1* expression.⁷ Further studies are necessary to examine other miRNAs and/or transcription factors that regulate Mcl-1 expression in gastric epithelial cells.

Our data from this study showed that the selective COX-2 inhibitor celecoxib activated the expression of *miR-29c* and suppressed its target oncogene *Mcl-1*, resulting in induction of apoptosis in gastric cancer cells. We confirmed that suppression of Mcl-1 by celecoxib was cancelled after transfection with the specific *miR-29c* inhibitor. These results suggest that suppression of Mcl-1 and induction of apoptosis in gastric cancer cells by celecoxib are mediated by *miR-29c*. Since celecoxib is a selective COX-2 inhibitor frequently used for treatment of pain, fever and inflammation, it can prevent side effects of other anticancer drugs. Combination treatment with celecoxib and other anticancer drugs may be beneficial for reducing side effects and achieving more potent activity against gastric cancer.




Figure 3. *miR-29c* is activated by celecoxib and suppresses Mcl-1. (a) The expression levels of *miR-29c* in AGS and MKN45 cells treated with celecoxib with or without the specific *miR-29c* inhibitor. Cont: control; Cel: celecoxib; Cel+29cl: celecoxib with *miR-29c* inhibitor. * $p < 0.05$ compared with control. (b) Western blotting for expression levels of Mcl-1 in AGS and MKN45 cells after transfection with the *miR-29c* precursor molecules and treatment with celecoxib with or without the specific *miR-29c* inhibitor. 29c: *miR-29c*; Cel+29cl: celecoxib with *miR-29c* inhibitor; Cel: celecoxib; Cont: control. β -actin was used as the internal control. (c) ChIP assay with the C/EBP α antibody in AGS and MKN45 cells treated with celecoxib. The database shows that there are C/EBP α binding sites in the promoter region of *miR-29c* (www.gene-regulation.com/pub/programs.html). Arrows show the primers used for the ChIP assay. Underlining shows the C/EBP α binding sites. Immunoprecipitation with the C/EBP α antibody was significantly increased in AGS and MKN45 cells treated with celecoxib (* $p < 0.05$).

The miRNA expression profiles shown in this study, and in other reports, have demonstrated that *miR-29b* and *miR-29c* have different expression patterns, suggesting that *miR-29c* has a unique promoter in gastric cancer cells. The results of ChIP

assay indicated that celecoxib activates *miR-29c* expression by enhancing the binding of C/EBP α DNA at the promoter. Selective COX-2 inhibitors may activate the tumor suppressor *miR-29c* via C/EBP α and induce apoptosis of gastric cancer cells.




Figure 4. Celecoxib and *miR-29c* induce apoptosis in gastric cancer cells. Annexin V-FITC staining in AGS and MKN45 cells was analyzed by fluorescence microscopy (a) and flow cytometry (b). Annexin V staining (green staining on the cell surface membrane) was markedly increased in AGS and MKN45 cells 48 hr after *miR-29c* transfection and celecoxib treatment (* $p < 0.05$).

Further investigations will be needed to clarify the molecular mechanism underlying the recruitment of C/EBP α at the *miR-29c* promoter region by selective COX-2 inhibitors.

In conclusion, the tumor suppressor *miR-29c* is downregulated with overexpression of its target oncogene Mcl-1 in gastric cancers and plays important roles in cancer progression. Treatment of gastric cancer cells with the selective COX-2 inhibitor

celecoxib activates *miR-29c* expression, resulting in suppression of Mcl-1 and induction of apoptosis in the cells. Administration of selective COX-2 inhibitors may have promise for the treatment of gastric cancer via restoration of *miR-29c*.

Acknowledgements

The authors thank Kana Yamada for her technical assistance.

References

- He L, Hannon GJ. MicroRNAs: small RNAs with a big role in gene regulation. *Nat Rev Genet* 2004;5:522–31.
- Calin GA, Croce CM. MicroRNA signatures in human cancers. *Nat Rev Cancer* 2006;6:857–66.
- Calin GA, Croce CM. Chromosomal rearrangements and microRNAs: a new cancer link with clinical implications. *J Clin Invest* 2007;117:2059–66.
- Saito Y, Liang G, Egger G, et al. Specific activation of microRNA-127 with downregulation of the proto-oncogene BCL6 by chromatin-modifying drugs in human cancer cells. *Cancer Cell* 2006;9:435–43.
- Saito Y, Jones PA. Epigenetic activation of tumor suppressor microRNAs in human cancer cells. *Cell Cycle* 2006;5:2220–2.
- Saito Y, Friedman JM, Chihara Y, et al. Epigenetic therapy upregulates the tumor suppressor microRNA-126 and its host gene EGFL7 in human cancer cells. *Biochem Biophys Res Commun* 2009;379:726–31.
- Saito Y, Suzuki H, Tsugawa H, et al. Chromatin remodeling at Alu repeats by epigenetic treatment activates silenced microRNA-512-5p with downregulation of Mcl-1 in human gastric cancer cells. *Oncogene* 2009;28:2738–44.
- Saito Y, Suzuki H, Tsugawa H, et al. Dysfunctional gastric emptying with downregulation of muscle-specific microRNAs in *Helicobacter pylori*-infected mice. *Gastroenterology* 2011;140:189–98.
- Suzuki H, Hibi T, Marshall BJ. *Helicobacter pylori*: present status and future prospects in Japan. *J Gastroenterol* 2007;42:1–15.
- Suzuki H, Iwasaki E, Hibi T. *Helicobacter pylori* and gastric cancer. *Gastric Cancer* 2009;12:79–87.
- Fujishiro M. Endoscopic submucosal dissection for stomach neoplasms. *World J Gastroenterol* 2006;12:5108–12.
- Kakushima N, Fujishiro M. Endoscopic submucosal dissection for gastrointestinal neoplasms. *World J Gastroenterol* 2008;14:2962–7.
- Esquela-Kerscher A, Slack FJ. Oncomirs - microRNAs with a role in cancer. *Nat Rev Cancer* 2006;6:259–69.
- Saito Y, Suzuki H, Hibi T. The role of microRNAs in gastrointestinal cancers. *J Gastroenterol* 2009;44 Suppl 19:18–22.

15. Arber N, Eagle CJ, Spicak J, et al. Celecoxib for the prevention of colorectal adenomatous polyps. *N Engl J Med* 2006;355:885-95.
16. Bertagnolli MM, Eagle CJ, Zauber AG, et al. Celecoxib for the prevention of sporadic colorectal adenomas. *N Engl J Med* 2006;355:873-84.
17. Kuo CH, Hu HM, Tsai PY, et al. Short-term celecoxib intervention is a safe and effective chemopreventive for gastric carcinogenesis based on a Mongolian gerbil model. *World J Gastroenterol* 2009;15:4907-14.
18. Zhang LJ, Wang SY, Huo XH, et al. Anti-*Helicobacter pylori* therapy followed by celecoxib on progression of gastric precancerous lesions. *World J Gastroenterol* 2009;15:2731-8.
19. Mott JL, Kobayashi S, Bronk SF, et al. miR-29 regulates Mcl-1 protein expression and apoptosis. *Oncogene* 2007;26:6133-40.
20. Sengupta S, den Boon JA, Chen IH, et al. MicroRNA 29c is down-regulated in nasopharyngeal carcinomas, up-regulating mRNAs encoding extracellular matrix proteins. *Proc Natl Acad Sci USA* 2008;105:5874-8.
21. Friedman JM, Liang G, Liu CC, et al. The putative tumor suppressor microRNA-101 modulates the cancer epigenome by repressing the polycomb group protein EZH2. *Cancer Res* 2009;69:2623-9.
22. Guo Y, Chen Z, Zhang L, et al. Distinctive microRNA profiles relating to patient survival in esophageal squamous cell carcinoma. *Cancer Res* 2008;68:26-33.
23. Stamatopoulos B, Meuleman N, Haibe-Kains B, et al. microRNA-29c and microRNA-223 down-regulation has in vivo significance in chronic lymphocytic leukemia and improves disease risk stratification. *Blood* 2009;113:5237-45.
24. Mraz M, Malinova K, Kotaskova J, et al. miR-34a, miR-29c and miR-17-5p are downregulated in CLL patients with TP53 abnormalities. *Leukemia* 2009;23:1159-63.
25. Ueda T, Volinia S, Okumura H, et al. Relation between microRNA expression and progression and prognosis of gastric cancer: a microRNA expression analysis. *Lancet Oncol* 2010;11:136-46.
26. Krajewski S, Bodrug S, Gascoyne R, et al. Immunohistochemical analysis of Mcl-1 and Bcl-2 proteins in normal and neoplastic lymph nodes. *Am J Pathol* 1994;145:515-25.
27. Wu CC, Lin JC, Yang SC, et al. Modulation of the expression of the invasion-suppressor CRMP-1 by cyclooxygenase-2 inhibition via reciprocal regulation of Sp1 and C/EBPalpha. *Mol Cancer Ther* 2008;7:1365-75.
28. Kozopas KM, Yang T, Buchan HL, et al. MCL1, a gene expressed in programmed myeloid cell differentiation, has sequence similarity to BCL2. *Proc Natl Acad Sci USA* 1993;90:3516-20.
29. Shigemasa K, Katoh O, Shiroyama Y, et al. Increased MCL-1 expression is associated with poor prognosis in ovarian carcinomas. *Jpn J Cancer Res* 2002;93:542-50.
30. Taniai M, Grambihler A, Higuchi H, et al. Mcl-1 mediates tumor necrosis factor-related apoptosis-inducing ligand resistance in human cholangiocarcinoma cells. *Cancer Res* 2004;64:3517-24.
31. Zhou P, Qian L, Kozopas KM, et al. Mcl-1, a Bcl-2 family member, delays the death of hematopoietic cells under a variety of apoptosis-inducing conditions. *Blood* 1997;89:630-43.
32. Krajewska M, Fenoglio-Preiser CM, Krajewski S, et al. Immunohistochemical analysis of Bcl-2 family proteins in adenocarcinomas of the stomach. *Am J Pathol* 1996;149:1449-57.
33. Maeta Y, Tsujitani S, Matsumoto S, et al. Expression of Mcl-1 and p53 proteins predicts the survival of patients with T3 gastric carcinoma. *Gastric Cancer* 2004;7:78-84.

Overexpression of *miR-142-5p* and *miR-155* in Gastric Mucosa-Associated Lymphoid Tissue (MALT) Lymphoma Resistant to *Helicobacter pylori* Eradication

Yoshimasa Saito^{1,4}, Hidekazu Suzuki^{1*}, Hitoshi Tsugawa¹, Hiroyuki Imaeda¹, Juntaro Matsuzaki¹, Kenro Hirata¹, Naoki Hosoe¹, Masahiko Nakamura³, Makio Mukai², Hidetsugu Saito^{1,4}, Toshifumi Hibi¹

1 Division of Gastroenterology and Hepatology, Department of Internal Medicine, Shinjuku-ku, Tokyo, Japan, 2 Department of Pathology, Keio University School of Medicine, Shinjuku-ku, Tokyo, Japan, 3 School of Pharmaceutical Sciences, Kitasato University, Minato-ku, Tokyo, Japan, 4 Division of Pharmacotherapeutics, Keio University Faculty of Pharmacy, Minato-ku, Tokyo, Japan

Abstract

microRNAs (miRNAs) are small non-coding RNAs that can function as endogenous silencers of target genes and play critical roles in human malignancies. To investigate the molecular pathogenesis of gastric mucosa-associated lymphoid tissue (MALT) lymphoma, the miRNA expression profile was analyzed. miRNA microarray analysis with tissue specimens from gastric MALT lymphomas and surrounding non-tumor mucosae revealed that a hematopoietic-specific miRNA *miR-142* and an oncogenic miRNA *miR-155* were overexpressed in MALT lymphoma lesions. The expression levels of *miR-142-5p* and *miR-155* were significantly increased in MALT lymphomas which do not respond to *Helicobacter pylori* (*H. pylori*) eradication. The expression levels of *miR-142-5p* and *miR-155* were associated with the clinical courses of gastric MALT lymphoma cases. Overexpression of *miR-142-5p* and *miR-155* was also observed in *Helicobacter heilmannii*-infected C57BL/6 mice, an animal model of gastric MALT lymphoma. In addition, *miR-142-5p* and *miR-155* suppress the proapoptotic gene *TP53INP1* as their target. The results of this study indicate that overexpression of *miR-142-5p* and *miR-155* plays a critical role in the pathogenesis of gastric MALT lymphoma. These miRNAs might have potential application as therapeutic targets and novel biomarkers for gastric MALT lymphoma.

Citation: Saito Y, Suzuki H, Tsugawa H, Imaeda H, Matsuzaki J, et al. (2012) Overexpression of *miR-142-5p* and *miR-155* in Gastric Mucosa-Associated Lymphoid Tissue (MALT) Lymphoma Resistant to *Helicobacter pylori* Eradication. PLoS ONE 7(11): e47396. doi:10.1371/journal.pone.0047396

Editor: William C. S. Cho, Queen Elizabeth Hospital, Hong Kong

Received: June 2, 2012; **Accepted:** September 12, 2012; **Published:** November 28, 2012

Copyright: © 2012 Saito et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by Grant-in-Aid for Young Scientists A (23680090 to Y.S.) and Grant-in-Aid for Scientific Research B (22300169 to H.S.) from Japan Society for Promotion of Science (www.jsps.go.jp), Takeda Science Foundation (www.takeda-sci.or.jp; to Y.S.), Sagawa Foundation for Promotion of Cancer Research (sagawa-gan.or.jp; to Y.S.), Smoking Research Foundation (www.srf.or.jp; to H.S.), and Keio Gijuku Academic Development Funds (www.keio.ac.jp; to Y.S. and H.S.). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: hsuzuki@a6.keio.jp

Introduction

Extranodal marginal zone B-cell lymphoma of mucosa-associated lymphoid tissue (MALT) is a low-grade lymphoma characterized by histological features such as lymphoepithelial lesions (LELs). The stomach is the most common site of MALT lymphoma, accounting for almost half of all cases. Gastric MALT lymphomas are sometimes associated with chronic inflammation triggered by chronic infection with *Helicobacter pylori* (*H. pylori*), suggesting that the proliferation of MALT lymphoma cells may depend on immune responses to antigens. Indeed, *H. pylori* eradication therapy leads to complete remission in 60–80% of cases of gastric MALT lymphoma and has been used as a first-line treatment [1–3]. However, 20–40% of cases do not respond to *H. pylori* eradication therapy, and predictors of the response to antibiotic treatment as well as the appropriate length of the observation period before second-line treatment remain controversial.

The *API2-MALT1* fusion gene, which results from a t(11;18)(q21;q21) translocation, has been identified as the most frequent chromosome translocation in MALT lymphoma cells [4,5], and Liu *et al.* [6] have reported that it is a potential predictor

of resistance to *H. pylori* eradication therapy. Since *API2-MALT1* fusion transcripts lead to inhibition of apoptosis [7,8], they may confer a survival benefit on MALT lymphoma cells. Despite these reports, the molecular mechanism underlying the initiation and progression of gastric MALT lymphoma is not fully understood.

Many previous studies have focused mainly on aberrant expression of protein-coding genes in the pathogenesis of MALT lymphoma [9]. However, it has recently become apparent that non-coding genes including microRNAs (miRNAs) play important roles as tumor suppressor genes and oncogenes during human carcinogenesis. miRNAs are small (20–25 nucleotides) non-coding RNAs that function as endogenous silencers of target genes. miRNAs are expressed in a tissue-specific manner and play critical roles in cellular proliferation, apoptosis and differentiation [10]. It has been shown that aberrant expression of miRNAs contributes to the development of human malignancies, and that miRNA expression signatures are associated with prognostic factors of human diseases [11–15]. Moreover, we have recently proposed that epigenetic regulation of tumor suppressor miRNAs could be a novel therapeutic approach for the treatment of human malignancies [16–19].

Table 1. Clinicopathological and molecular features of gastric MALT lymphoma cases.

No.	Sex	Age	<i>API2-MALT1</i> fusion	<i>H.pylori</i> before Tx	infection after Tx	Remission after eradication	Months after eradication
1	M	74	+	+	–	NC	74
2	M	74	+	+	–	NC	32
3	M	61	+	–	–	NC	10
4	F	75	+	–	*	NC	*
5	F	59	+	–	*	NC	*
6	F	57	+	–	*	NC	*
7	M	77	–	+	–	NC	45
8	F	68	–	–	–	NC	3
9	F	58	–	+	–	CR	24
10	F	46	–	+	–	CR	10
11	F	66	–	+	–	CR	10
12	F	52	–	+	–	CR	8
13	M	64	–	+	–	CR	6
14	F	71	–	+	–	CR	4
15	F	70	–	+	–	CR	4
16	M	60	–	+	–	CR	4
17	F	56	–	+	–	CR	4
18	M	54	–	+	–	CR	3
19	F	68	–	+	–	CR	2
20	F	40	–	+	–	CR	1

CR, complete remission; NC, no change.

*Eradication therapy not used.

doi:10.1371/journal.pone.0047396.t001

Although recent studies have shown that *miR-155*, a potential oncogenic miRNA, is highly expressed in diffuse large B cell lymphoma (DLBCL) [20,21] and overexpression of *miR-155* is correlated with a poor outcome in patients with DLBCL [20], the miRNA expression profiles of low-grade MALT lymphoma have not yet been described. In the present study, therefore, the miRNA expression profiles and potential miRNA target genes of gastric MALT lymphoma were analyzed to clarify the molecular pathogenesis of this malignancy.

Methods

Patients and tissue specimens

Twenty patients with primary low-grade gastric MALT lymphomas who were diagnosed and treated at Keio University Hospital (Tokyo, Japan) were enrolled. This study was approved by the ethics committee of Keio University School of Medicine (No. 18-96-3) and was registered with the Clinical Trials Registry (UMIN 000000858). Written informed consent was obtained from all the patients before examination. The clinicopathological and molecular features of the patients are shown in Table 1. The *H. pylori* infection status was identified using the ¹³C-urea breath test. Some cases were confirmed by serological or histological examination in addition to the ¹³C-urea breath test. Tissue specimens from gastric MALT lymphomas and the corresponding non-tumor gastric mucosae were obtained from patients during an endoscopic biopsy and were stored in RNAlater (Ambion, Austin, TX) at –80°C until RNA extraction.

Fluorescence in situ hybridization (FISH) analysis

To detect the chromosome translocation t(11;18)(q21;q21) and the *API2-MALT1* fusion gene arising from it, FISH analysis using the LSI *API2-MALT1* t(11;18)(q21;q21) Dual Color, Dual Fusion Translocation Probe (Vysis/Abbot Laboratories Ltd., Maidenhead, Berkshire, United Kingdom) was performed by Mitsubishi Chemical Medience Corporation (Tokyo, Japan).

RNA extraction and microarray analysis

Total RNA, including small RNA, was extracted using a mirVana miRNA isolation kit (Ambion). The total RNAs from each of three gastric MALT lymphomas were pooled, as were the total RNAs from each of three matched samples of non-tumor gastric mucosa. The miRNA microarray analysis was performed by LC Sciences (www.lcsciences.com, Houston, TX). RNA from the MALT lymphomas was labeled using Cy5, while the RNA from the non-tumor gastric mucosa was labeled using Cy3. The microarray chip contains probe regions that detect 711 miRNA transcripts listed in Sanger miRBase Release 10.0 (<http://www.sanger.ac.uk>) and 5 probes for each miRNA. Detected signals greater than background plus 3 times the standard deviation were derived for each color channel; the mean and the co-variance (CV = standard deviation × 100 / replicate mean) of each probe having a detected signal was calculated. For two color experiments, the ratio of the two sets of detected signals and *p*-values of the *t*-test were calculated. Differentially detected signals were accepted as true when the *p*-values of the ratios were less than 0.01. The average values of their signal intensities are shown in

Table 2. miRNAs differentially expressed between MALT lymphomas and non-tumor gastric mucosae.

No.	microRNA	Non-tumor (Cy3 signal)	MALT (Cy5 signal)	Fold change
1	miR-142-5p	56	1236.3	22.1
2	miR-142-3p	25.4	440.5	17.3
3	miR-223	574.1	5330.9	9.3
4	miR-138	40.9	200.8	4.9
5	miR-155	3093.2	13503.9	4.4
6	miR-572	1581.7	368.2	0.2
7	miR-146b-5p	1029.1	4249.4	4.1
8	miR-141	6525.1	1588.8	0.2
9	miR-146a	2187.8	8374.4	3.8
10	miR-378	3218.7	933.6	0.3

The miRNA microarray analysis was performed by LC Sciences (www.lcsciences.com, Houston, TX). All data were submitted to the ArrayExpress database; the accession number was E-MEXP-1898.
doi:10.1371/journal.pone.0047396.t002

Table 2. All the data were submitted to the ArrayExpress database under the accession number E-MEXP-1898.

Quantitative RT-PCR of miRNAs

Levels of miRNA expression were analyzed using quantitative RT-PCR and the TaqMan microRNA assay for *miR-142-5p* and *miR-155* (Applied Biosystems, Foster City, CA), in accordance with the manufacturer's instructions. The expression levels were normalized to that of U6 RNA and expressed as the mean \pm standard deviation.

Infection of C57BL/6 mice with *H. heilmannii*

Ten months prior to the experiment, six-week-old C57BL/6 mice were inoculated with gastric mucosal homogenates from *H. heilmannii*-infected mice, as described previously [22]. Stool DNA was extracted using a QIAamp DNA Stool Mini Kit (Qiagen, Tokyo, Japan), and infection with *H. heilmannii* was confirmed by real-time PCR of stool DNA using specific primers (HeilF, 5'-AAGTCGAACGATGAAGCCTA-3' and HeilR, 5'-ATTTGG-TATTAATCACCATTTC-3'). RNA extraction and histological examination were performed using tissue specimens from the stomachs of *H. heilmannii*-infected mice and control mice. This study was approved by the Keio University Animal Research Committee (No. 08080).

Western blotting

Protein extracts were separated using SDS/polyacrylamide gel electrophoresis and transferred to a nitrocellulose membrane. The membranes were hybridized with the rabbit anti-human TP53INP1 polyclonal antibody (LifeSpan Biosciences, Seattle, WA). This antibody shows cross-reactivity with mouse TP53INP1. The signal intensities were analyzed using ImageJ software.

Immunohistochemistry

Formalin-fixed and paraffin-embedded tissues were deparaffinized and rehydrated. For antigen retrieval, the sections were treated for 20 min at 100°C in an autoclave and non-specific reactions were blocked with a blocking reagent (Protein Block Serum-Free, Dako Cytomation, Glostrup, Denmark). The sections were incubated with the rabbit anti-human TP53INP1 polyclonal

antibody (diluted 1:200; LifeSpan Biosciences) overnight at 4°C followed by horseradish peroxidase (HRP)-labeled anti-rabbit IgG (Histofine, Simple stain MAX-PO; Nichirei, Tokyo, Japan) for 30 min at room temperature. Then, the sections were treated with 3, 3'-diaminobenzidine tetrahydrochloride solution. All the sections were counterstained with HE.

Luciferase assay

Luciferase constructs were made by ligating oligonucleotides containing the wild-type or mutant target site of the *TP53INP1* 3' untranslated region (UTR) into the *Xba* I site of the pGL3-control vector (Promega, Madison, WI). The AGS human gastric cancer cell line was used in this study. AGS was obtained from the American Type Culture Collection (Rockville, MD). Cells were cultured in RPMI1640 medium supplemented with 10% fetal bovine serum. AGS cells were transfected with 0.4 μ g of firefly luciferase reporter vector containing the wild-type or mutant target site and 0.02 μ g of the control vector containing *Renilla* luciferase pRL-CMV (Promega) using lipofectamine 2000 (Invitrogen, Carlsbad, CA) in 24-well plates. The *miR-142-5p* and *miR-155* precursor molecules and negative control precursor miRNAs were purchased from Ambion. The molecules were transfected into AGS cells at a final concentration of 100 nM each. The luciferase assays were performed 24 hours after transfection using the Dual Luciferase Reporter Assay System (Promega). The activity of firefly luciferase was normalized to that of *Renilla* luciferase.

Statistics

Differences in the miRNA levels between the groups were analyzed using paired *t* test and unpaired *t* test. Differences at $p < 0.05$ were considered significant.

Results

Overexpression of *miR-142-5p* and *miR-155* in gastric MALT lymphoma

To identify miRNAs that play critical roles in the development of gastric MALT lymphoma, we performed miRNA microarray analysis using RNAs obtained from three gastric MALT lymphomas and three matched samples of non-tumor gastric mucosa. The miRNAs that were differentially expressed between non-tumor gastric mucosae and MALT lymphoma lesions are listed in Table 2. The miRNA expression profile revealed that a hematopoietic-specific miRNA, *miR-142* [23,24], and an oncogenic miRNA, *miR-155* [12,25], were overexpressed in MALT lymphoma lesions, relative to the levels of expression in the corresponding non-tumor mucosae.

To validate the microarray data, we performed quantitative RT-PCR for *miR-142-5p* and *miR-155* in 20 cases of gastric MALT lymphoma. The clinicopathological and molecular features of the 20 patients are shown in Table 1. All of the gastric MALT lymphoma cases in the present study were stage I_E according to the Ann Arbor staging system [26]. The average age of the patients was 62.5 years (male/female: 7/13). As shown in Figure 1, the expression levels of *miR-142-5p* and *miR-155* in the gastric MALT lymphoma lesions were significantly higher than those in the corresponding non-tumor gastric mucosae ($p < 0.05$ and $p < 0.05$, respectively).




Figure 1. Expression levels of miR-142-5p and miR-155 and responses to *H. pylori* eradication therapy in gastric MALT lymphoma cases. The levels of miR-142-5p and miR-155 expression were evaluated using quantitative RT-PCR and normalized to the expression of U6 RNA. The filled bars and blank bars indicate MALT lymphoma lesions and non-tumor gastric mucosae, respectively. The levels of miR-142-5p and miR-155 expression in gastric MALT lymphoma lesions were significantly increased relative to the corresponding non-tumor gastric mucosae ($p < 0.05$ and $p < 0.05$, respectively). The levels of miR-142-5p and miR-155 expression in MALT lymphoma lesions which confers resistance to *H. pylori* eradication were significantly higher than those in lesions lacking the *API2-MALT1* fusion gene, which exhibited CR after *H. pylori* eradication ($p < 0.0001$ and $p < 0.005$, respectively). CR, complete remission; NC, no change. doi:10.1371/journal.pone.0047396.g001


miR-142-5p and miR-155 as novel biomarkers of gastric MALT lymphoma

The patients with gastric MALT lymphoma were divided into two groups according to their response to *H. pylori* eradication therapy (Table 1). All the patients who were positive for *H. pylori* underwent eradication therapy and were subsequently confirmed to be *H. pylori*-negative. The use of *H. pylori* eradication therapy for patients with gastric MALT lymphoma who are *H. pylori*-negative is controversial. Nakamura *et al.* have recently reported the long-term clinical outcome of patients with gastric MALT lymphoma after *H. pylori* eradication [27]. In their study, 44 *H. pylori*-negative gastric MALT lymphoma patients underwent *H. pylori* eradication therapy, and 6 (14%) responded to *H. pylori* eradication. Akamatsu

et al. have also reported that 1 out of 9 (11%) *H. pylori*-negative patients showed complete regression of gastric MALT lymphoma [28]. On the other hand, Fischbach *et al.* have reported that most patients with histological residuals of gastric MALT lymphoma after successful *H. pylori* eradication (treatment failure cases) had a favorable disease course without additional treatment (Fischbach *et al.* Gut 2007; 56: 1685–7). For these patients, a watch and wait strategy with regular endoscopic examinations and biopsies appears to be safe. Our policy of *H. pylori* eradication therapy for gastric MALT lymphoma patients who were *H. pylori*-negative was decided in accordance with the preference of individual patients.

We also analyzed the chromosome translocation t(11;18)(q21;q21) with the *API2-MALT1* fusion gene by FISH

A



B




Figure 2. Expression levels of *miR-142-5p* and *miR-155* in gastric MALT lymphoma cases before and after *H. pylori* eradication therapy. (A) Representative cases of the FISH analyses for the detection of the *API2-MALT1* fusion gene are shown. The probes for *API2* and *MALT1* were labeled with green and red signals, respectively. The *API2-MALT1* fusion gene produces a yellow signal. (B) Patients #14 and #16, positive for *H. pylori* and negative for the *API2-MALT1* fusion gene. These patients received *H. pylori* eradication therapy and attained CR. The expression levels of *miR-142-5p* and *miR-155* were significantly lower after *H. pylori* eradication. * $p < 0.05$. Patient #1, positive for *H. pylori* and positive for the *API2-MALT1* fusion gene. This patient was resistant to *H. pylori* eradication therapy (NC). The expression levels of *miR-142-5p* and *miR-155* showed no significant differences.

doi:10.1371/journal.pone.0047396.g002

analysis. The probes for *API2* were labeled with green signals, and the probes for *MALT1* were labeled with red signals. In positive cases, the *API2-MALT1* fusion genes resulting from the chromosome translocation produced yellow signals (Figure 2A).

As shown in Figure 1, the levels of expression of *miR-142-5p* and *miR-155* in MALT lymphoma lesions that were resistant to *H. pylori* eradication were significantly higher than in cases showing complete remission (CR) after *H. pylori* eradication ($p < 0.0001$ and $p < 0.005$, respectively). Although the *API2-MALT1* fusion gene has been identified as a potential predictor of resistance to *H. pylori* eradication therapy, patients #7 and #8 with gastric MALT lymphoma lacking the *API2-MALT1* fusion gene were resistant to *H. pylori* eradication. These patients showed increased expression levels of *miR-142-5p* and *miR-155* (Figure 1).

In addition, we investigated the correlation between the levels of *miR-142-5p* and *miR-155* expression and the clinical courses of gastric MALT lymphoma cases. The expression data for patients #14 and #16 with gastric MALT lymphoma who achieved CR after *H. pylori* eradication therapy showed that the levels of *miR-142-5p* and *miR-155* expression were significantly reduced after

eradication (Figure 2B, $p < 0.05$ and $p < 0.05$, respectively). On the other hand, in patient #1 with gastric MALT lymphoma harboring the *API2-MALT1* fusion gene that was resistant to *H. pylori* eradication, there was no significant difference in the expression levels of *miR-142-5p* and *miR-155* (Figure 2B). Although the *H. pylori* infection was eradicated by antibiotic treatment, histological examination showed no regression of the MALT lymphoma lesion. This patient was followed up by observation without treatment, and no marked changes in the endoscopic or histological findings were observed. These findings indicate that the levels of *miR-142-5p* and *miR-155* expression have potential applicability as novel biomarkers of gastric MALT lymphoma.

Overexpression of *miR-142-5p* and *miR-155* in an animal model of gastric MALT lymphoma

To further confirm the molecular pathogenesis of gastric MALT lymphoma, we examined the expression levels of *miR-142-5p* and *miR-155* in an animal model of gastric MALT lymphoma. Infection with *H. heilmannii* induces gastric LELs that are consistent with low-grade MALT lymphomas in C57BL/6 mice [22].




Figure 3. Expression levels of *miR-142-5p* and *miR-155* in an animal model of gastric MALT lymphoma. (A) Macroscopic and light microscopic findings with HE staining of the stomach of control and *H. heilmannii*-infected mice (10 months after infection). Protrusive lesions in the fundic stomach of *H. heilmannii*-infected mice were observed. Light microscopic observations using HE staining revealed the presence of LELs consistent with low-grade MALT lymphoma. (B) Expression levels of *miR-142-5p* and *miR-155* in the stomach of control and *H. heilmannii*-infected mice. The levels of *miR-142-5p* and *miR-155* expression normalized to the level of U6 RNA were significantly increased in *H. heilmannii*-infected C57BL/6 mice. * $p < 0.05$. doi:10.1371/journal.pone.0047396.g003

Infection with *H. heilmannii* was confirmed using quantitative PCR analysis of stool DNA (data not shown). Ten months after infection with *H. heilmannii*, control ($n = 3$) and *H. heilmannii*-infected C57BL/6 mice ($n = 4$) were dissected, and this revealed protruding lesions in the gastric fundus in all of the latter. Light microscopic observations using hematoxylin and eosin (HE) staining revealed the presence of LELs consistent with low-grade MALT lymphomas (Figure 3A). Quantitative RT-PCR analyses showed that the levels of expression of *miR-142-5p* and *miR-155* were significantly higher in *H. heilmannii*-infected C57BL/6 mice ($p < 0.05$ and $p < 0.05$, respectively), being similar to the results for human gastric MALT lymphomas (Figure 3B).

miR-142-5p and *miR-155* suppress the proapoptotic gene *TP53INP1* as their target

Identification of miRNA target genes is essential for determining miRNA function. Recent studies have indicated that a single miRNA may regulate more than 200 target genes. A database for predicting target genes, TargetScan (<http://www.targetscan.org>), revealed that both *miR-142-5p* and *miR-155* are able to bind to the 3' UTR of the mRNA of the proapoptotic gene *TP53INP1* (Tumor Protein P53 Inducible Nuclear Protein 1). Moreover, *miR-155* has been shown to repress *TP53INP1*, which inhibits the development of pancreatic tumors [29]. Therefore, we focused on *TP53INP1* as a common target of *miR-142-5p* and *miR-155*.

To confirm the target specificity of *miR-142-5p* and *miR-155* for *TP53INP1*, we performed a luciferase reporter assay using a vector containing the putative *TP53INP1* 3' UTR target sites downstream of the luciferase reporter gene, which was transfected into AGS cells. The base pairing between *miR-142-5p* and *miR-155* and the wild-type (WT) or mutant (MUT) target sites in the 3' UTR of *TP53INP1* mRNA is shown in Figure 4A. The luciferase activities of the AGS cells transfected with the *TP53INP1*-WT construct were significantly lower after transfection with *miR-142-5p* and *miR-155* ($p < 0.05$ and $p < 0.005$, respectively), whereas those of cells transfected with the *TP53INP1*-MUT construct and the pGL3

control vector (empty vector) showed no significant differences (Figure 4B). It has been shown that conserved perfect 6- to 8-bp matches between the 5' end of the mature miRNA and the 3' UTR of the predicted target mRNA (called 'seed' matches) are the most important factor determining miRNA targets [30]. As shown in Fig. 4A, 'seed' matches between the 5' end of the miRNAs and the 3' UTR of *TP53INP1* were stronger in *miR-155* than in *miR-142*, suggesting that *miR-155* may have a more profound effect on suppression of *TP53INP1*.

We next examined the expression levels of *TP53INP1* by Western blot analysis in *H. heilmannii*-infected mice and immunohistochemistry in human gastric MALT lymphoma-like lesions. The expression of *TP53INP1* was suppressed in *H. heilmannii*-infected mice, relative to control mice (Figure 4C). We examined the expression levels of *TP53INP1* in human gastric tissues by immunohistochemistry. The gastric tissue shown in Fig. 4D contains both MALT lymphoma and non-MALT lymphoma tissue. The infiltrating lymphocytic lesion shown by HE staining is consistent with MALT lymphoma. *TP53INP1* staining is negative or faint in the MALT lymphoma lesion, whereas normal gastric glands around the MALT lymphoma show sufficient staining for *TP53INP1*. Thus *TP53INP1* staining was markedly reduced in MALT lymphoma lesions (Figure 4D). These findings suggest that *TP53INP1* is a common target of *miR-142-5p* and *miR-155* and is suppressed by overexpression of both *miR-142-5p* and *miR-155* in gastric MALT lymphoma lesions.

Discussion

A recent study has demonstrated that miRNA expression profiles can be used to classify the developmental lineages and differentiation stages of tumors, and are more accurate for tumor classification than conventional mRNA profiles [31]. Furthermore, miRNA expression signatures are associated with prognostic factors and disease progression in chronic lymphocytic leukemia [11] and lung cancer [32]. Thus miRNA expression is clinically promising as both a diagnostic tool and a prognostic marker for




Figure 4. miR-142-5p and miR-155 suppress TP53INP1 as their target. (A) Luciferase reporter constructs of wild-type (WT) and mutant (MUT) target sites in the 3' UTR of TP53INP1 mRNA. The base pairings between miR-142-5p and miR-155 and their putative target sites in the 3' UTR of TP53INP1 mRNA are shown. Underlining indicates mutant sequences. (B) Firefly luciferase activity was normalized to the Renilla luciferase activity of TP53INP1 wild-type (WT), mutant (MUT), and pGL3 control (empty vector) in AGS cells transfected with negative control (NC), miR-142-5p, and miR-155. The luciferase activities of the AGS cells transfected with the TP53INP1-WT construct were significantly lower after transfection of miR-142-5p and miR-155, whereas those transfected with the TP53INP1-MUT construct or the pGL3 control vector (empty vector) showed no significant differences. * $p < 0.05$ and ** $p < 0.005$, compared with negative control. (C) Western blot analysis of TP53INP1 in control and *H. heilmannii*-infected mice. β -actin was used as a loading control. The expression of TP53INP1 was suppressed in *H. heilmannii*-infected mice, compared with the expression level in control mice. (D) Immunohistochemistry for TP53INP1 in human gastric MALT lymphoma. Sections were counterstained with HE. TP53INP1 staining was markedly reduced in the MALT lymphoma lesions. doi:10.1371/journal.pone.0047396.g004

human malignancies. The results of microarray analysis revealed a unique miRNA expression profile in gastric MALT lymphoma. Recent studies have demonstrated that the expression level of miR-155 is significantly elevated in DLBCL, which is considered to represent high-grade transformation from MALT lymphoma [20,33]. miR-142, miR-155 and miR-223 have been reported to be hematopoiesis-specific miRNAs [24]. These findings are

consistent with our results, and in the present study we focused on miR-142 and miR-155, because miR-142 is the most up-regulated miRNA and miR-155 plays a critical role in the pathogenesis of B-cell lymphoma. The levels of miR-142-5p and miR-155 expression were associated with the clinical course of gastric MALT lymphoma, including the response to *H. pylori* eradication. The *API2-MALT1* fusion gene has been identified as a

potential predictor of resistance to *H. pylori* eradication therapy. In the present study, two cases of gastric MALT lymphoma resistant to *H. pylori* eradication lacked the *API2-MALT1* fusion gene (patients #7, #8). These cases showed increased expression of *miR-142-5p* and *miR-155*. Therefore, these miRNAs may be more useful markers than the *API2-MALT1* fusion gene in patients with gastric MALT lymphoma.

TP53INP1 is a proapoptotic stress-induced p53 target gene. p53 activates *TP53INP1* transcription, and overexpression of *TP53INP1* induces cell cycle arrest and apoptosis [34]. Gironella *et al.* [29] have shown that oncogenic *miR-155* is overexpressed in pancreatic ductal adenocarcinoma and suppresses its target, *TP53INP1*, resulting in cancer progression. In agreement with these findings, our results showed that *TP53INP1* was suppressed by both *miR-142-5p* and *miR-155*, possibly leading to inhibition of apoptosis and acceleration of MALT lymphoma cell proliferation. These findings suggest that overexpression of *miR-142-5p* and *miR-155* concomitant with suppression of *TP53INP1* reflect the increased proliferation of MALT lymphoma cells. The distinct connection between aberrant expression of *miR-142-5p* and *miR-155* and the progression of MALT lymphoma suggests that miRNAs could be potential therapeutic targets. A recent study has shown that chemically engineered oligonucleotides, termed

'antagomirs,' can work as specific inhibitors of endogenous miRNAs in mice [35], and might be potentially applicable to silence *miR-142-5p* and *miR-155* for the treatment of gastric MALT lymphomas that are resistant to *H. pylori* eradication therapy.

Conclusions


Overexpression of *miR-142-5p* and *miR-155* is presumed to play a critical role in the initiation and progression of gastric MALT lymphoma, suggesting that these miRNAs may be potentially useful as therapeutic targets and novel biomarkers for gastric MALT lymphomas. Inhibition of *miR-142-5p* and *miR-155* might be a novel approach for the prevention and treatment of gastric MALT lymphoma. Further studies involving more patients are warranted to explore the clinical use of *miR-142-5p* and *miR-155* for diagnosis and treatment of gastric MALT lymphoma.

Author Contributions

Conceived and designed the experiments: YS H. Suzuki TH. Performed the experiments: YS HT JM KH. Analyzed the data: YS H. Suzuki JM. Contributed reagents/materials/analysis tools: YS H. Suzuki HI JM KH NH MN MM H. Saito. Wrote the paper: YS H. Suzuki.

References

- Sackmann M, Morgner A, Rudolph B, Neubauer A, Thiede C, et al. (1997) Regression of gastric MALT lymphoma after eradication of *Helicobacter pylori* is predicted by endosonographic staging. MALT Lymphoma Study Group. *Gastroenterology* 113: 1087–1090.
- Ruskone-Fourmesttraux A, Lavergne A, Aegerter PH, Megraud F, Palazzo L, et al. (2001) Predictive factors for regression of gastric MALT lymphoma after anti-*Helicobacter pylori* treatment. *Gut* 48: 297–303.
- Suzuki H, Hibi T, Marshall BJ (2007) *Helicobacter pylori*: present status and future prospects in Japan. *J Gastroenterol* 42: 1–15.
- Akagi T, Motegi M, Tamura A, Suzuki R, Hosokawa Y, et al. (1999) A novel gene, MALT1 at 18q21, is involved in t(11;18)(q21;q21) found in low-grade B-cell lymphoma of mucosa-associated lymphoid tissue. *Oncogene* 18: 5785–5794.
- Dierlamm J, Baens M, Wlodarska I, Stefanova-Ouzounova M, Hernandez JM, et al. (1999) The apoptosis inhibitor gene API2 and a novel 18q gene, MLT, are recurrently rearranged in the t(11;18)(q21;q21) associated with mucosa-associated lymphoid tissue lymphomas. *Blood* 93: 3601–3609.
- Liu H, Ye H, Ruskone-Fourmesttraux A, De Jong D, Pileri S, et al. (2002) T(11;18) is a marker for all stage gastric MALT lymphomas that will not respond to *H. pylori* eradication. *Gastroenterology* 122: 1286–1294.
- Motegi M, Yonezumi M, Suzuki H, Suzuki R, Hosokawa Y, et al. (2000) API2-MALT1 chimeric transcripts involved in mucosa-associated lymphoid tissue type lymphoma predict heterogeneous products. *Am J Pathol* 156: 807–812.
- Hosokawa Y, Suzuki H, Suzuki Y, Takahashi R, Seto M (2004) Antiapoptotic function of apoptosis inhibitor 2-MALT1 fusion protein involved in t(11;18)(q21;q21) mucosa-associated lymphoid tissue lymphoma. *Cancer Res* 64: 3452–3457.
- Farinha P, Gascoyne RD (2005) Molecular pathogenesis of mucosa-associated lymphoid tissue lymphoma. *J Clin Oncol* 23: 6370–6378.
- He L, Hannon CJ (2004) MicroRNAs: small RNAs with a big role in gene regulation. *Nat Rev Genet* 5: 522–531.
- Calin GA, Ferracin M, Cimmino A, Di Leva G, Shimizu M, et al. (2005) A MicroRNA signature associated with prognosis and progression in chronic lymphocytic leukemia. *N Engl J Med* 353: 1793–1801.
- Calin GA, Croce CM (2006) MicroRNA signatures in human cancers. *Nat Rev Cancer* 6: 857–866.
- Saito Y, Suzuki H, Hibi T (2009) The role of microRNAs in gastrointestinal cancers. *J Gastroenterol* 44 Suppl 19: 18–22.
- Cho WC (2010) MicroRNAs: potential biomarkers for cancer diagnosis, prognosis and targets for therapy. *Int J Biochem Cell Biol* 42: 1273–1281.
- Cho WC (2010) MicroRNAs in cancer - from research to therapy. *Biochim Biophys Acta* 1805: 209–217.
- Saito Y, Liang G, Egger G, Friedman JM, Chuang JC, et al. (2006) Specific activation of microRNA-127 with downregulation of the proto-oncogene BCL6 by chromatin-modifying drugs in human cancer cells. *Cancer Cell* 9: 435–443.
- Saito Y, Jones PA (2006) Epigenetic activation of tumor suppressor microRNAs in human cancer cells. *Cell Cycle* 5: 2220–2222.
- Saito Y, Friedman JM, Chihara Y, Egger G, Chuang JC, et al. (2009) Epigenetic therapy upregulates the tumor suppressor microRNA-126 and its host gene EGF7 in human cancer cells. *Biochem Biophys Res Commun* 379: 726–731.
- Saito Y, Suzuki H, Tsugawa H, Nakagawa I, Matsuzaki J, et al. (2009) Chromatin remodeling at Alu repeats by epigenetic treatment activates silenced microRNA-512-5p with downregulation of Mcl-1 in human gastric cancer cells. *Oncogene* 28: 2738–2744.
- Eis PS, Tam W, Sun L, Chadburn A, Li Z, et al. (2005) Accumulation of miR-155 and BIC RNA in human B cell lymphomas. *Proc Natl Acad Sci U S A* 102: 3627–3632.
- Metzler M, Wilda M, Busch K, Viehmann S, Borkhardt A (2004) High expression of precursor microRNA-155/BIC RNA in children with Burkitt lymphoma. *Genes Chromosomes Cancer* 39: 167–169.
- Nakamura M, Murayama SY, Serizawa H, Sekiya Y, Eguchi M, et al. (2007) "Candidatus *Helicobacter heilmannii*" from a cynomolgus monkey induces gastric mucosa-associated lymphoid tissue lymphomas in C57BL/6 mice. *Infect Immun* 75: 1214–1222.
- Chen CZ, Li L, Lodish HF, Bartel DP (2004) MicroRNAs modulate hematopoietic lineage differentiation. *Science* 303: 83–86.
- Ramkissoon SH, Mainwaring LA, Ogasawara Y, Keyvanfar K, McCoy JP, Jr., et al. (2006) Hematopoietic-specific microRNA expression in human cells. *Leuk Res* 30: 643–647.
- Calin GA, Croce CM (2006) MicroRNAs and chromosomal abnormalities in cancer cells. *Oncogene* 25: 6202–6210.
- Lister TA, Crowther D, Sutcliffe SB, Glatstein E, Canellos GP, et al. (1989) Report of a committee convened to discuss the evaluation and staging of patients with Hodgkin's disease: Cotswolds meeting. *J Clin Oncol* 7: 1630–1636.
- Nakamura S, Sugiyama T, Matsumoto T, Iijima K, Ono S, et al. (2012) Long-term clinical outcome of gastric MALT lymphoma after eradication of *Helicobacter pylori*: a multicentre cohort follow-up study of 420 patients in Japan. *Gut* 61: 507–513.
- Akamatsu T, Mochizuki T, Okiyama Y, Matsumoto A, Miyabayashi H, et al. (2006) Comparison of localized gastric mucosa-associated lymphoid tissue (MALT) lymphoma with and without *Helicobacter pylori* infection. *Helicobacter* 11: 86–95.
- Gironella M, Seux M, Xie MJ, Cano C, Tomasini R, et al. (2007) Tumor protein 53-induced nuclear protein 1 expression is repressed by miR-155, and its restoration inhibits pancreatic tumor development. *Proc Natl Acad Sci U S A* 104: 16170–16175.
- Lewis BP, Burge CB, Bartel DP (2005) Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* 120: 15–20.
- Lu J, Getz G, Miska EA, Alvarez-Saavedra E, Lamb J, et al. (2005) MicroRNA expression profiles classify human cancers. *Nature* 435: 834–838.
- Yanaihara N, Caplen N, Bowman E, Seike M, Kumamoto K, et al. (2006) Unique microRNA molecular profiles in lung cancer diagnosis and prognosis. *Cancer Cell* 9: 189–198.
- Fang C, Zhu DX, Dong HJ, Zhou ZJ, Wang YH, et al. (2012) Serum microRNAs are promising novel biomarkers for diffuse large B cell lymphoma. *Ann Hematol* 91: 553–559.
- Tomasini R, Seux M, Nowak J, Bontemps C, Carrier A, et al. (2005) TP53INP1 is a novel p73 target gene that induces cell cycle arrest and cell death by modulating p73 transcriptional activity. *Oncogene* 24: 8093–8104.
- Krutzfeldt J, Rajewsky N, Braich R, Rajeev KG, Tuschl T, et al. (2005) Silencing of microRNAs in vivo with 'antagomirs'. *Nature* 438: 685–689.



Identification of a unique nsaid, fluoro-loxoprofen with gastroprotective activity^{☆,☆☆}

Shintaro Suemasu^{a,b}, Naoki Yamakawa^{a,b}, Tomoaki Ishihara^a, Teita Asano^a, Kayoko Tahara^a, Ken-ichiro Tanaka^{a,b}, Hirofumi Matsui^c, Yoshinari Okamoto^b, Masami Otsuka^b, Koji Takeuchi^d, Hidekazu Suzuki^e, Tohru Mizushima^{a,b,*}

^a Department of Analytical Chemistry, Faculty of Pharmacy, Keio University, Tokyo 105-8512, Japan

^b Faculty of Life Sciences, Kumamoto University, Kumamoto 862-0973, Japan

^c Graduate School of Comprehensive Human Sciences, University of Tsukuba, Tsukuba 305-8575, Japan

^d Division of Pathological Sciences, Department of Pharmacology and Experimental Therapeutics, Kyoto Pharmaceutical University, Kyoto 607-8414, Japan

^e Division of Gastroenterology and Hepatology, Department of Internal Medicine, Keio University School of Medicine, Tokyo 160-8582, Japan

ARTICLE INFO

Article history:

Received 6 August 2012

Accepted 18 September 2012

Available online 26 September 2012

Keywords:

NSAIDs

Gastrointestinal complications

Mucus

ABSTRACT

We previously proposed that direct cytotoxicity of NSAIDs due to their membrane permeabilization activity, together with their ability to decrease gastric prostaglandin E₂, contributes to production of gastric lesions. Compared to loxoprofen (LOX), fluoro-loxoprofen (F-LOX) has much lower membrane permeabilization and gastric ulcerogenic activities but similar anti-inflammatory activity. In this study, we examined the mechanism for this low ulcerogenic activity in rats. Compared to LOX, the level of gastric mucosal cell death was lower following administration of F-LOX. However, the gastric level of prostaglandin E₂ was similar in response to treatment with the two NSAIDs. Oral pre-administration of F-LOX conferred protection against the formation of gastric lesions produced by subsequent administration of LOX and orally administered F-LOX resulted in a higher gastric pH value and mucus content. In the presence of a stimulant of gastric acid secretion, the difference in the ulcerogenic activity of F-LOX and LOX was less apparent. Furthermore, an increase in the mucus was observed in gastric cells cultured in the presence of F-LOX in a manner dependent of increase in the cellular level of cAMP. These results suggest that low ulcerogenic activity of F-LOX involves its both low direct cytotoxicity and protective effect against the development of gastric lesions. This protective effect seems to be mediated through an increase in a protective factor (mucus) and a decrease in an aggressive factor (acid).

© 2012 Elsevier Inc. All rights reserved.

^{*} We thank Dr. H. Sakai (University of Toyama, Toyama, Japan) for helpful discussion on the membrane preparation and enzyme assay.

^{**} This work was supported by Grants-in-Aid of Scientific Research from the Ministry of Health, Labour, and Welfare of Japan, Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan, and Grants-in-Aid of the Japan Science and Technology Agency.

Abbreviations: ABTS, 2,2'-azinobis(3-ethylbenzothiazoline-6-sulfonic acid)-2NH₄; BAPTA-AM, 1,2-bis(2-aminophenoxy)ethane-*N,N,N',N'*-tetraacetic acid; COX, cyclooxygenase; DAPI, 4',6-diamidino-2-phenylindole dihydrochloride; EIA, enzyme immunoassay; ELLA, enzyme-linked lectin-binding assay; F-LOX, fluoro-loxoprofen; FBS, fetal bovine serum; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; H & E, hematoxylin and eosin; IBMX, 3-isobutyl-1-methylxanthine; LOX, loxoprofen; NSAID, non-steroidal anti-inflammatory drug; PG, prostaglandin; RGM1, rat normal gastric epithelial cell line; SD, Sprague-Dawley; SBA, soybean agglutinin; TCA, trichloroacetic acid; TUNEL, terminal deoxynucleotidyl transferase-mediated biotinylated UTP nick end labeling.

^{*} Corresponding author at: Department of Analytical Chemistry, Faculty of Pharmacy, Keio University, 1-5-30, Shibakoen, Minato-ku, Tokyo 105-8512, Japan. Tel./fax: +81 3 5400 2628.

E-mail address: mizushima-th@pha.keio.ac.jp (T. Mizushima).

1. Introduction

The balance between aggressive and defensive factors determines the development of gastric lesions, with either a relative increase in aggressive factors or a decrease in protective factors resulting in lesions. The gastric mucosa can be challenged by a variety of both endogenous and exogenous factors, including gastric acid, reactive oxygen species, ethanol, *Helicobacter pylori* and non-steroidal anti-inflammatory drugs (NSAIDs) [1]. In order to protect the mucosa, the body relies on defence systems such as the production of surface mucus and bicarbonate, and the regulation of gastric mucosal blood flow. Prostaglandin E₂ (PGE₂) also exerts a strong protective effect, inhibiting the secretion of gastric acid and stimulating the production of mucus [2].

NSAIDs, such as indomethacin, comprise a therapeutically valuable family of drugs [3]. An inhibitory effect of NSAIDs on cyclooxygenase (COX) activity is responsible for their anti-inflammatory actions, COX being an enzyme that is essential for

the synthesis of prostaglandins, which have a strong capacity to induce inflammation. However, as described above, NSAID use is also associated with gastrointestinal complications [4–7], which was thought to result from the inhibition of COX and a decrease in gastric PGE₂ level. In fact, NSAIDs have been reported to stimulate the secretion of gastric acid and inhibit the production of mucus through decreasing gastric PGE₂ level [8,9]. However, it is now believed that the production of gastric lesions by NSAIDs involves additional mechanisms, given that the increased incidence of gastric lesions and the decrease in PGE₂ levels induced by NSAIDs do not always occur in parallel [10,11]. We have recently demonstrated that NSAIDs induce cell death (apoptosis) in cultured gastric mucosal cells and at the gastric mucosa in a manner independent of COX inhibition [12–16]. With regards to the molecular mechanism governing this apoptosis, we have proposed the following pathway. Permeabilization of cytoplasmic membranes by NSAIDs stimulates Ca²⁺ influx and increases intracellular Ca²⁺ levels, which in turn induces the endoplasmic reticulum stress response [12,17,18]. During the course of this response, an apoptosis-inducing transcription factor, C/EBP homologous transcription factor, is induced and, as we have previously shown, this protein is essential for NSAID-induced apoptosis [13,19]. Furthermore, we have proposed that both COX inhibition and gastric mucosal cell death are important for the formation of NSAID-induced gastric lesions *in vivo* [16,20].

In 1991, two subtypes of COX, COX-1 and COX-2, which are responsible for the majority of COX activity at the gastrointestinal mucosa and in tissues subject to inflammation, respectively, were identified [21]. It is therefore not surprising that a reduced incidence of gastroduodenal lesions has been reported following treatment with selective COX-2 inhibitors [22–24]. However, a recently raised issue concerning the use of selective COX-2 inhibitors is the potential risk of cardiovascular thrombotic events [25,26]. This may be due to the fact that prostacyclin, a potent anti-aggregator of platelets and a vasodilator, is mainly produced by COX-2 [27–29]. Therefore, in order to minimize clinical complications, gastric safe NSAIDs other than selective COX-2 inhibitors need to be developed. Based on the hypothesis outlined above, we believe that NSAIDs with lower membrane permeabilization activity would represent an efficacious alternative, even if they had no selectivity for COX-2 [14].

In order to investigate this possibility, we screened for such compounds from a range of clinically used NSAIDs without COX-2 selectivity, and found that the membrane permeabilization activity and direct cytotoxicity of loxoprofen (LOX) (Fig. 1A) was relatively lower than that of the other NSAIDs tested [30]. LOX is a leading NSAID on the Japanese market, being widely used because clinical studies have suggested that it is safer than other traditional (non-selective) NSAIDs [31,32]. LOX is a pro-drug, which is converted (by reduction of the cyclopentanone moiety) to its active metabolite (the *trans*-alcohol metabolite, LOX-OH) by aromatic aldehyde-ketone reductase (Fig. 1A) [33]. We therefore synthesized a series of LOX derivatives, demonstrating that fluoro-loxoprofen (F-LOX) (Fig. 1A) has much lower membrane permeabilization and gastric ulcerogenic activities than LOX, but similar anti-inflammatory activity, suggesting that it is likely to be a therapeutically viable drug [34]. We suggested that F-LOX is also a pro-drug, which is converted to its active metabolite (the *trans*-alcohol metabolite, F-LOX-OH (Fig. 1A)), because the inhibitory effect of F-LOX-OH on COX is much more potent than that of F-LOX *in vitro* [34]. Although we concluded that the low membrane permeabilization activity of F-LOX is responsible for its low ulcerogenic activity [34], it remained possible that other mechanisms could also be involved. In this study, we therefore examined the mechanism governing the low ulcerogenic activity of F-LOX. Our results suggest that this effect is mediated not only by the low

direct cytotoxicity due to its low membrane permeabilization activity, but also by protection of the gastric mucosa. In contrast to LOX and other NSAIDs, oral administration of F-LOX led to an increase in gastric pH value and mucus, suggesting that these effects are also involved in the low ulcerogenic activity of this drug.

2. Experimental procedures

2.1. Chemicals and animals

LOX, LOX-OH, F-LOX and F-LOX-OH (Fig. 1A) were synthesized in our laboratory as previously described [34]. Methylcellulose and RPMI1640 were obtained from Wako Pure Chemical Industries (Osaka, Japan). Formaldehyde, paraformaldehyde, Alcian blue 8GX, mucin, cycloheximide, histamine, fetal bovine serum (FBS), Dulbecco's modified Eagle's medium nutrient mixture F-12 Ham, 2-methyl-8-(phenylmethoxy)imidazo[1,2-*a*]pyridine-3-acetonitrile (SCH 28080; an inhibitor of gastric H⁺,K⁺-ATPase), 3-isobutyl-1-methylxanthine (IBMX), omeprazole, forskolin and bovine serum albumin (BSA) were purchased from Sigma (St. Louis, MO). Mayer's hematoxylin, 1% eosin alcohol solution and mounting medium for histochemical analysis (malinol) were from MUTO Pure Chemicals (Tokyo, Japan). Terminal transferase was obtained from Roche Diagnostics (Mannheim, Germany). Biotin 14-ATP and streptavidin-conjugated Alexa Fluor 488 were purchased from Invitrogen (Carlsbad, CA). Mounting medium for the TdT-mediated biotinylated UTP nick end labeling (TUNEL) assay (VECTASHIELD) was from Vector Laboratories, Inc. (Burlingame, CA). 4',6-diamidino-2-phenylindole dihydrochloride (DAPI) and 1,2-bis(2-aminophenoxy)ethane-*N,N,N',N'*-tetraacetic acid (BAPTA-AM) were obtained from Dojindo (Kumamoto, Japan). SQ22536 was from CALBIOCHEM (San Diego, CA). ONO-8711 and ONO-AE2-227 were from our laboratory stocks. The prostaglandin E₂ enzyme immunoassay (EIA) kit was purchased from Cayman Chemical (Ann Arbor, MI). cAMP complete ELISA kit was from Enzo Life Sciences (Farmingdale, NY). Horseradish peroxidase-labeled soybean agglutinin (SBA-HRP) was from Seikagaku Biobusiness Co. (Tokyo, Japan). 2,2'-azinobis(3-ethylbenzothiazoline-6-sulfonic acid)-2NH₄ (ABTS) was obtained from KPL (Gaithersburg, MD). The RNeasy Mini kit was from QIAGEN (Valencia, CA), the first-strand cDNA synthesis kit was obtained from Takara (Kyoto, Japan), and iQ SsoFast EvaGreen Supermix was purchased from Bio-Rad (Hercules, CA). Wistar rats (3-week-old males) and Sprague-Dawley (SD) rats (6-week-old males) were obtained from Charles River Laboratories Japan (Yokohama, Japan). Guinea pigs (3-week-old males) were obtained from Japan SLC (Shizuoka, Japan). Animals were housed under conditions of controlled temperature (22–24 °C) and illumination (12 h light cycle starting at 8:00 AM) for 1 week before experiments. The experiments and procedures described here were performed in accordance with the Guide for the Care and Use of Laboratory Animals as adopted and promulgated by the National Institutes of Health, and were approved by the Animal Care Committee of Keio University and Kumamoto University. Totally, we used 287 Wistar rats, 182 SD rats and 4 guinea pigs for all experiments in this study.

2.2. Gastric and small intestinal damage assay

The gastric ulcerogenic response was examined as described previously [20,35] with some modifications. Wistar or SD rats fasted for 18 h were orally administered each NSAID and, after 8 h or 4 h, respectively, the animals were sacrificed, their stomachs were removed, and the gastric mucosal lesion area measured by an observer unaware of the treatment that the animals had received. For Wistar rats, calculation of the scores involved measuring the area of all the lesions in square millimeters and summing the




Fig. 1. Production of lesions in the stomach and small intestine by oral administration of F-LOX or LOX. Structures of drugs used in this study were shown (A). Wistar rats were orally administered either the indicated doses (B, E, –F), 200 or 213.4 mg/kg (C and D), or 30 or 32.0 mg/kg (G) of LOX or F-LOX, respectively. Rats were similarly administered LOX-OH or F-LOX-OH (E). Their stomachs (B–E) or small intestines (F, G) were then removed either 8 h later (B, D, E), at the indicated time-points (C, G), or 24 h later (F). The stomach and small intestine were scored for damage (B, C, E, F and G) and sections of gastric tissues were prepared and subjected to H & E staining. Images are magnified 2.5 times (lower panels) (D). Values are mean \pm S.E.M. ** $P < 0.01$; n.d., not detected. Scale bar, 50 μ m.