

(OS) of only 1 year [2–7]. Therefore, the development of novel anticancer agents or strategies for treating AGC is urgently required.

Human epidermal growth factor receptor 2 (HER2) is a growth factor receptor that is overexpressed in approximately 10–30 % of gastric cancers [8, 9]. The main function of HER2 is to mediate cell growth, differentiation, and survival [10, 11]. Therefore, HER2-positive tumors are expected to have more aggressive characteristics than HER2-negative tumors. Numerous studies have shown that HER2-positive breast cancer is associated with a poor prognosis when compared with HER2-negative breast cancer [11, 12]. Similarly, the prognostic value of HER2 status in gastric cancer has been evaluated in a number of reports. Although recent studies of gastric cancer have shown no impact of HER2 on survival [13, 14], the majority of the past studies have suggested that HER2-positive gastric cancer was associated with a poorer prognosis than HER2-negative gastric cancer [9, 15, 16].

Trastuzumab, a humanized monoclonal antibody that targets HER2, has recently been shown to improve the prognosis of HER2-positive AGC [17]. Combination chemotherapy consisting of 5-fluorouracil (5-FU) or capecitabine plus cisplatin with trastuzumab has shown a significantly higher response rate (47 vs. 35 %, $P = 0.0017$), a longer progression-free survival [PFS; 6.7 vs. 5.5 months; hazard ratio (HR) = 0.71, 95 % confidence interval (CI) 0.59–0.86], and a longer OS (13.8 vs. 11.1 months; HR = 0.74, 95 % CI 0.60–0.91) than chemotherapy alone [17]. This difference was more prominent in patients with immunohistochemical staining (IHC) of 3+ (IHC 3+) or IHC 2+ plus positive gene amplification according to fluorescence in situ hybridization (FISH) (16.0 vs. 11.8 months; HR = 0.65, 95 % CI 0.51–0.83). The risk reduction achieved with trastuzumab is almost comparable to that found in a pivotal study of breast cancer [18]. According to these results, chemotherapy with trastuzumab is considered to be standard chemotherapy for HER2-positive AGC. However, the question of whether trastuzumab improves the prognosis of HER2-positive AGC patients to the levels observed in HER2-negative disease still remains. HER2-positive breast cancer patients who receive trastuzumab have been reported to have a better prognosis (HR of 0.56) than women with HER2-negative breast cancer [19].

The purpose of this study was to evaluate the impact of HER2 status and trastuzumab treatment on the prognosis of patients with AGC. We evaluated the impact of trastuzumab on OS using a time-varying covariate (TVC) analysis.

Patients and methods

Patients

This retrospective study was designed to compare the OS of HER2-positive and HER2-negative AGC patients who received systemic chemotherapy. In this analysis, HER2 positivity was defined as IHC 3+ or IHC 2+ plus FISH positivity, because these criteria were considered to be indications for using trastuzumab by a subset analysis of the ToGA trial [17, 20]. The following additional principal inclusion criteria were used: the presence of histologically proven, inoperable gastric cancer in patients who had received systemic chemotherapy; an Eastern Cooperative Oncology Group performance status (ECOG PS) of 0–2; and sufficient bone marrow, liver, and renal function.

Between April 2005 and August 2011, 807 consecutive patients with AGC received systemic chemotherapy at our institution, 650 of whom met the inclusion criteria. Among those 650 patients, 364 patients were evaluated for HER2 to screen them for the ToGA study or for trastuzumab treatment, and these 364 patients were analyzed in this study. Although trastuzumab had not yet been approved in Japan at the time of our study, using trastuzumab for HER2-positive patients was approved by our institution based on the results of the ToGA study. We note that fluoropyrimidine (5-FU or the oral fluoropyrimidine S-1), cisplatin, docetaxel, and paclitaxel were approved for use and were consistently available throughout this period. Written informed consent for the chemotherapy was obtained from all of the patients.

Statistical analysis

We compared the OS according to HER2 status and trastuzumab exposure by dividing the patients into three groups (HER2-negative vs. HER2-positive treated with trastuzumab vs. HER2-positive treated without trastuzumab). The OS was estimated from the date of systemic chemotherapy initiation to the date of death or last known date of survival. The vital status and disease status were confirmed by checking the medical records from the date of the last follow-up visit. In the cases that were lost to follow up, the vital status was confirmed by the census record, which is conducted annually at our institution. The median OS rate was estimated by the Kaplan–Meier method. The baseline patient characteristics and exposure to each class of chemotherapeutic agent, fluoropyrimidine (5-FU or oral fluoropyrimidine), platinum agents (cisplatin or oxaliplatin), taxanes (docetaxel or paclitaxel), and irinotecan and trastuzumab, were evaluated. Because other agents (mitomycin C or methotrexate) were also commonly used as

salvage therapy at our institution, we also evaluated exposure to these treatments. The disease progression associated with each line of chemotherapy was measured from the beginning of treatment to the date of disease progression, as evaluated by the attending physician.

The differences in the OS of each group were evaluated by univariate and multivariate analyses using a Cox proportional hazards model and presented as an HR and a 95 % CI. A subset analysis was stratified by the characteristics of the patients with significant differences in the three groups. We also evaluated the association between the exposure to each class of chemotherapeutic agent and the OS using multivariate analyses, because differences in exposure to agents other than trastuzumab may have contributed to survival differences. Because the length of exposure to each agent class varied over time (i.e., between first-, second-, and third-line treatments), the analyses may have been compromised by lead-time bias, which would result in false-positive or false-negative associations between a larger number of chemotherapeutic lines and longer survival. To minimize this potential bias, the exposure to each agent class was analyzed as a TVC. In addition, because disease progression was the primary reason for proceeding to the next line of chemotherapy, tumor progression during each line of chemotherapy was included in the TVCs. Each TVC was constructed as a step function initially set at 0 and increased by 1 U each time the corresponding event was observed. This method has been used in several reports that have evaluated the impact of drug exposure on survival [21–23].

Other variables considered in the multivariate analyses were ECOG PS (0 vs. ≥ 1), gender, histological type (diffuse vs. intestinal), age (< 65 vs. ≥ 65 years), previous gastrectomy (no vs. yes), disease status (advanced vs. recurrent), prior adjuvant chemotherapy (no vs. yes), presence of liver metastasis (no vs. yes), presence of peritoneal metastasis (no vs. yes), number of metastatic sites (1 vs. ≥ 2), and first-line chemotherapy (monotherapy vs. combinations). The *P* values for testing differences in the baseline characteristics of each group were calculated with the χ^2 test for homogeneity or trend or with Fisher's exact test.

The statistical analyses were performed using STATA ver. 10 (Stata Corp LP, College Station, TX, USA). All of the tests were two-sided, and *P* values less than 0.05 were considered to be statistically significant.

Results

Patient characteristics

The patient characteristics according to HER2 status and trastuzumab use are shown in Table 1. Among the HER2-

evaluated patients, 58 (15.9 %) were HER2-positive. HER2-positivity was more frequently associated with intestinal-type histology, liver metastasis, and lymph node metastasis. By contrast, peritoneal metastasis was more common in the HER2-negative patients than in the HER2-positive patients (Table 1). Among the HER2-positive patients, 43 received trastuzumab; 23 patients started it as first-line therapy (9 of these patients were included in the ToGA study) and 20 started it as second- or further-line therapy. Among the patients who were treated with trastuzumab who were not in the ToGA study, 22 patients continued with trastuzumab beyond the date of progression. The median duration of trastuzumab treatment in the 43 patients receiving this agent was 8.4 months (range 0.5 to 25 +), and 28 patients were still continuing with trastuzumab treatment at the time of this analysis. The frequency of exposure to platinum agents tended to be higher in HER2-positive disease, although no other significant differences were observed between the 3 groups (Table 1).

Survival and multivariate analysis

The median follow up at the time of the analysis was 38.9 months. The median OS of the HER2-positive patients (*n* = 58; 24.1 months; 95 % CI, 14.9–31.1 months) was significantly longer than that of the HER2-negative patients (*n* = 304; 13.9 months; 95 % CI, 12.7–16.1 months) by univariate analysis (HR 0.68; 95 % CI, 0.46–0.99; *P* = 0.045), although the difference was not significant in the multivariate analysis (HR 0.67; 95 % CI, 0.44–1.02; *P* = 0.067). The median OS of the HER2-positive patients treated with trastuzumab (*n* = 43; 24.7 months; 95 % CI, 15.6–35.1 months) was significantly longer than that of the HER2-negative patients (13.9 months), as verified by univariate analysis (HR 0.57; 95 % CI, 0.36–0.90; *P* = 0.015, Fig. 1) and multivariate analysis (adjusted HR of 0.58; 95 % CI, 0.36–0.95; *P* = 0.03). By contrast, the HER2-positive patients not treated with trastuzumab (*n* = 15) showed survival similar to that of the HER2-negative patients (13.5 months; 95 % CI, 3.6 to not reached), as demonstrated by univariate analysis (HR 1.11; 95 % CI, 0.58–2.08; *P* = 0.76) and multivariate analysis (HR 1.04; 95 % CI, 0.52–2.11; *P* = 0.91). The subset analysis according to the characteristics of selected patients demonstrated consistently higher rates of survival among HER2-positive patients treated with trastuzumab than among HER2-negative patients, with no significant heterogeneity (Table 2).

According to the multivariate TVC analysis, exposure to trastuzumab was independently associated with an improved prognosis (HR 0.56, 95 % CI, 0.33–0.93; *P* = 0.026, Table 3). Additionally, exposure to fluoropyrimidine, taxanes, and irinotecan was associated with a better prognosis, and exposure to platinum agents was

Table 1 Patient characteristics

Characteristics	HER2-positive patients treated with trastuzumab (<i>n</i> = 43, %)	HER2-positive patients not treated with trastuzumab (<i>n</i> = 15, %)	HER2-negative patients (<i>n</i> = 306, %)	<i>P</i> value
Age (years)				
Median (range)	63 (30–78)	68 (45–80)	64 (30–93)	0.33
Gender				0.45
Male	28 (65)	12 (80)	196 (64)	
Female	15 (35)	3 (20)	110 (36)	
ECOG PS				0.52
0	23 (53)	7 (47)	123 (40)	
1	17 (40)	6 (40)	149 (49)	
2	3 (7)	2 (13)	34 (11)	
Histological type				<0.001
Diffuse	12 (28)	4 (27)	217 (71)	
Intestinal	31 (72)	11 (73)	99 (29)	
Site of primary tumor				<0.001
EG junction	8 (19)	5 (33)	28 (9)	
Gastric	35 (81)	10 (67)	278 (91)	
Disease status				0.49
Advanced	33 (77)	9 (60)	203 (66)	
Recurrent	10 (23)	6 (40)	103 (34)	
Previous gastrectomy				0.59
No	26 (60)	8 (53)	160 (52)	
Yes	17 (40)	7 (47)	146 (48)	
Adjuvant chemotherapy				0.81
No	34 (79)	13 (87)	248 (81)	
Yes	9 (21)	2 (13)	58 (19)	
Site of metastasis				
Lymph node	27 (63)	12 (80)	145 (47)	<u>0.01</u>
Peritoneum	12 (28)	5 (33)	172 (56)	<u>0.001</u>
Liver	21 (49)	6 (40)	71 (23)	<u>0.001</u>
Number of metastatic organs				0.11
1	15 (35)	8 (53)	174 (57)	
2 or more	28 (65)	7 (47)	132 (43)	
First-line CTx				<u>0.03</u>
Combination	35 (81)	12 (80)	192 (63)	
Monotherapy	8 (19)	3 (20)	114 (37)	
Initiation of CTx				0.1
2005–2007	10 (23)	8 (53)	96 (31)	
2008–2011	33 (77)	7 (47)	210 (69)	
Exposure to agents				
Fluoropyrimidine	38 (88)	13 (87)	292 (95)	0.08
Platinum	42 (98)	11 (73)	228 (75)	<u>0.003</u>
Taxane	33 (77)	12 (80)	206 (67)	0.29
Irinotecan	22 (51)	6 (40)	137 (45)	0.67
Others	5 (12)	1 (7)	49 (16)	0.49

Underlined *P* values are significant

HER2 human epidermal growth factor receptor 2, *PS* performance status, *ECOG* Eastern Cooperative Oncology Group, *CTx* chemotherapy, *EG* esophagogastric

associated with improved survival, although the significance was borderline. By contrast, the other agents had no impact on survival.

Discussion

In this study, we compared the OS in AGC patients according to HER2 status and exposure to trastuzumab. The OS of the HER2-positive patients who were treated

with trastuzumab was significantly longer than that of the HER2-negative patients, with an adjusted HR of 0.58. This result is similar to those observed in a large study of patients with breast cancer in which a similar degree of risk reduction was found [18]. In addition, the multivariate TVC analysis in our study showed that trastuzumab had a significant impact on OS. These results confirmed that trastuzumab therapy improves the prognosis of HER2-positive AGC patients beyond that of patients with HER2-negative AGC.

In this study, the HER2-positive patients not treated with trastuzumab showed survival rates similar to those of the HER2-negative patients. By contrast, the majority of previous studies have found poorer survival among patients with HER2-positive gastric cancer than among patients with HER2-negative tumors [9, 15, 16]. Most of these studies have been based on retrospective analyses in a single institution that combined patients with advanced-stage cancer and those with resected gastric cancer. Additionally, until the ToGA study, various definitions were used to evaluate HER2 status. Recently, Janjigian et al. [14] reported the results of a retrospective analysis of HER2 status among 381 patients who had enrolled in several clinical AGC studies. The HER2-positive patients (IHC 3 + or FISH +; $n = 78$) tended to have a better prognosis than the HER2-negative patients (13.9 vs. 11.4 months), although a multivariate analysis showed no impact of HER2 status on survival [14]. Additionally, an exploratory analysis of HER2 status in the AVAGAST study showed similar median OS in the HER2-negative and HER2-positive patients (10.5 vs. 9.8 months) who received capecitabine or 5-FU plus cisplatin [24]. Combined with our results, these findings suggest that HER2 status may




Fig. 1 The median OS of the HER2-positive patients treated with trastuzumab ($n = 43$; 24.7 months; 95 % CI, 15.6 to 35.1 months) was significantly longer than that of the HER2-negative patients $n = 306$; (13.9 months), as verified by univariate analysis (HR 0.57; 95 % CI, 0.36 to 0.90; $P = 0.015$) and multivariate analysis (adjusted HR of 0.58; 95 % CI, 0.36 to 0.95; $P = 0.03$). The HER2-positive patients not treated with trastuzumab ($n = 15$) showed survival similar to the HER2-negative patients (13.5 months; 95 % CI, 3.6 to not reached), as demonstrated by univariate analysis (HR 1.11; 95 % CI, 0.58 to 2.08; $P = 0.76$) and multivariate analysis (HR 1.04; 95 % CI, 0.52 to 2.11; $P = 0.91$)

Table 2 Impact of HER2 and trastuzumab use according to selected backgrounds

	HER2-negative	HER2-positive	
		Treated with trastuzumab HR (95 % CI)	Not treated with trastuzumab HR (95 % CI)
Histology			
Diffuse	1 (reference)	0.34 (0.12–0.99)	2.48 (0.67–8.90)
Intestinal	1 (reference)	0.53 (0.28–1.03)	0.90 (0.35–2.28)
<i>p-heterogeneity</i>		0.48	0.22
Site of metastasis			
Liver	1 (reference)	0.65 (0.28–1.51)	1.09 (0.24–4.74)
Lymph node	1 (reference)	0.68 (0.34–1.36)	0.78 (0.32–1.78)
Peritoneum	1 (reference)	0.76 (0.32–1.83)	1.00 (0.28–3.57)
<i>p-heterogeneity</i>		0.97	0.91
Initiation of chemotherapy			
Before 2008	1 (reference)	0.65 (0.27–1.59)	1.34 (0.56–3.24)
After 2008	1 (reference)	0.51 (0.25–1.06)	1.95 (0.25–15.27)
<i>p-heterogeneity</i>		0.69	0.87

Except for the stratifying factors, the HRs were adjusted for age, gender, ECOG PS, disease status, prior gastrectomy, adjuvant chemotherapy, and the number of metastatic sites
HR hazard ratio, CI confidence interval

Table 3 Results of the multivariate TVC analyses

Variable	Multivariate analysis with TVC ^a		
	HR	95 % CI	<i>P</i>
Exposure to agents			
Trastuzumab	0.56	0.33–0.93	<u>0.026</u>
Fluoropyrimidine	0.40	0.19–0.84	<u>0.015</u>
Platinum	0.63	0.38–1.03	0.06
Taxanes	0.37	0.25–0.57	<u><0.001</u>
Irinotecan	0.47	0.33–0.67	<u><0.001</u>
Other	0.85	0.55–1.38	0.39

Underlined *P* values are significant

TVC time-varying covariate, HR hazard ratio, CI confidence interval

^a Adjusted by age, gender, ECOG PS, disease status, histology, prior gastrectomy, adjuvant chemotherapy, presence of liver metastasis, presence of peritoneal metastasis, number of metastatic sites, time of initiation of chemotherapy, and tumor progression during each line of chemotherapy

have a small impact on survival in AGC. In addition, recent data from a large randomized study of resected gastric cancer found that HER2 status was not associated with the OS or with relapse-free survival in either the group who received surgery only or the group who also received adjuvant chemotherapy [13].

Interestingly, the risk reduction or median survival with trastuzumab treatment in our analysis was relatively better than the results of a subset analysis among those patients with IHC 3 + or IHC2 + plus FISH + in the ToGA study [17]. Several possibilities may explain this difference. First, the median exposure to trastuzumab in the ToGA study was reported to be 4.9 months [17]. Although the reasons for discontinuing the treatment were not reported in detail, continuing with trastuzumab beyond progression was not allowed in the ToGA study. By contrast, the median duration of exposure to trastuzumab in our study was longer (8.4 months). A number of patients who were treated outside of the ToGA study continued with trastuzumab beyond progression because this strategy has been found to contribute to improved treatment outcomes in randomized studies of breast cancer patients [25, 26]. A prospective study that evaluates the importance of trastuzumab beyond progression is warranted, as is further evaluation of AGC patients in a randomized study. Second, owing to the retrospective nature of the present study, unblinded bias may have overestimated the treatment effect. Regardless, our results reconfirmed the results of the ToGA study, which demonstrated that a new effective agent with the right target improved the prognosis of gastric cancer. Additionally, other agents targeting HER2 (such as lapatinib, which has already been shown to be effective in HER2-positive breast cancer) are under investigation in AGC, and the results are awaited.

It is important to note the methodological limitations of the present study. First, not all of the patients in this study period were evaluated for their HER2 status. Thus, the analysis may have been subject to some selection bias, although no significant differences other than PS were observed between the patients in the analysis and the patients not evaluated for HER2 status during the same period (data not shown). The median OS of the patients not evaluated for HER2 was 13.0 months, which was similar to that of the HER2-negative patients or HER2-positive patients not treated with trastuzumab in this analysis. Second, the study was a retrospective, non-randomized comparison of trastuzumab treatment in patients evaluated for HER2 status. Therefore, the differing characteristics of the patients and the differing treatments (other than with trastuzumab) in each group may have affected the results. Because the use of trastuzumab and the exposure to platinum agents differed between the groups, we used the TVC analysis to adjust the exposure to agents and comprehensively evaluate the impact of each agent class, regardless of the treatment line. However, TVC analysis is not necessarily adequate under the conditions of our study because its validity may depend on the assumption of a strong association between treatment selection at the time of the events and the history leading up to the events [27]. Other potential confounders, such as PS and the metastatic site, were also considered in the multivariate analyses; owing to the retrospective, non-randomized nature of the study, however, residual confounding effects caused by non-included factors cannot be completely ruled out. Nevertheless, because our survival results for the HER2-negative patients and the HER2-positive patients not treated with trastuzumab were quite similar to those of the ToGA trial and the results of other randomized controlled trials, the effect of this selection bias may have been small. Third, the small sample size and single-center population are other major limitations of this study.

In conclusion, we found that introducing trastuzumab improved the prognosis of HER2-positive AGC patients beyond that of HER2-negative patients. Because other targeted therapies are currently under study, more improvement in the prognosis of HER2-positive AGC patients is expected. Importantly, because HER2-positive patients are a minority of AGC patients (with a prevalence of less than 20 %), further investigation of other key AGC markers is warranted.

Conflict of interest None of the authors have financial or personal conflicts of interest to disclose.

References

1. International Agency for Research on Cancer. GLOBOCAN 2008: <http://www-dep.iarc.fr/CancerMondial.htm>.
2. Van Cutsem E, Moiseyenko VM, Tjulandin S, Majlis A, Constenla M, Boni C, et al. Phase III study of docetaxel and cisplatin

- plus fluorouracil compared with cisplatin and fluorouracil as first-line therapy for advanced gastric cancer: a report of the V325 Study Group. *J Clin Oncol.* 2006;24:4991–7.
3. Cunningham D, Starling N, Rao S, Iveson T, Nicolson M, Coxon F, et al. Capecitabine and oxaliplatin for advanced esophagogastric cancer. *N Engl J Med.* 2008;358:36–46.
 4. Koizumi W, Narahara H, Hara T, Takagane A, Akiya T, Takagi M, et al. S-1 plus cisplatin versus S-1 alone for first-line treatment of advanced gastric cancer (SPIRITS trial): a phase III trial. *Lancet Oncol.* 2008;9:215–21.
 5. Kang YK, Kang WK, Shin DB, Chen J, Xiong J, Wang J, et al. Capecitabine/cisplatin versus 5-fluorouracil/cisplatin as first-line therapy in patients with advanced gastric cancer: a randomised phase III noninferiority trial. *Ann Oncol.* 2009;20:666–73.
 6. Ajani JA, Rodriguez W, Bodoky G, Moiseyenko V, Lichinitser M, Gorbunova V, et al. Multicenter phase III comparison of cisplatin/S-1 with cisplatin/infusional fluorouracil in advanced gastric or gastroesophageal adenocarcinoma study: the FLAGS trial. *J Clin Oncol.* 2010;28:1547–53.
 7. Ohtsu A, Shah MA, Van Cutsem E, Rha SY, Sawaki A, Park SR, et al. Bevacizumab in combination with chemotherapy as first-line therapy in advanced gastric cancer: a randomized, double-blind, placebo-controlled phase III study. *J Clin Oncol.* 2011;29:3968–76.
 8. Hofmann M, Stoss O, Shi D, Büttner R, van de Vijver M, Kim W, et al. Assessment of a HER2 scoring system for gastric cancer: results from a validation study. *Histopathology.* 2008;52:797–805.
 9. Gravalos C, Jimeno A. HER2 in gastric cancer: a new prognostic factor and a novel therapeutic target. *Ann Oncol.* 2008;19:1523–9.
 10. Slamon DJ, Clark GM, Wong SG, Levin WJ, Ullrich A, McGuire WL. Human breast cancer: correlation of relapse and survival with amplification of the HER-2/neu oncogene. *Science.* 1987;235:177–82.
 11. Slamon DJ, Godolphin W, Jones LA, Holt JA, Wong SG, Keith DE, et al. Studies of the HER-2/neu proto-oncogene in human breast and ovarian cancer. *Science.* 1989;244:707–12.
 12. Seshadri R, Figgairi FA, Horsfall DJ, McCaul K, Setlur V, Kitchen P. Clinical significance of HER-2/neu oncogene amplification in primary breast cancer: the South Australian Breast Cancer Study Group. *J Clin Oncol.* 1993;11:1936–42.
 13. Terashima M, Ochiai A, Kitada K, Ichikawa W, Kurahashi I, Sakuramoto S, et al. Impact of human epidermal growth factor receptor (EGFR) and ERBB2 (HER2) expressions on survival in patients with stage II/III gastric cancer, enrolled in the ACTS-GC study. *J Clin Oncol.* 2011;29 (suppl; abstr 4013).
 14. Janjigian YY, Werner D, Pauligk C, Steinmetz K, Kelsen DP, Jäger E, et al. Prognosis of metastatic gastric and gastroesophageal junction cancer by HER2 status: a European and USA International collaborative analysis. *Ann Oncol.* 2012 [Epub ahead of print].
 15. Begnami MD, Fukuda E, Fregnani JH, Nonogaki S, Montagnini AL, Da Costa WL Jr, et al. Prognostic implications of altered human epidermal growth factor receptors (HERs) in gastric carcinomas: HER2 and HER3 are predictors of poor outcome. *J Clin Oncol.* 2011;29:3030–6.
 16. Kim KC, Koh YW, Chang HM, Kim TH, Yook JH, Kim BS, et al. Evaluation of HER2 protein expression in gastric carcinomas: comparative analysis of 1,414 cases of whole-tissue sections and 595 cases of tissue microarrays. *Ann Surg Oncol.* 2011;18:2833–40.
 17. Bang YJ, Van Cutsem E, Feyereislova A, Chung HC, Shen L, Sawaki A, et al. Trastuzumab in combination with chemotherapy versus chemotherapy alone for treatment of HER2-positive advanced gastric or gastro-oesophageal junction cancer (ToGA): a phase 3, open-label, randomised controlled trial. *Lancet.* 2010;376:687–97.
 18. Slamon DJ, Leyland-Jones B, Shak S, Fuchs H, Paton V, Bajamonde A, et al. Use of chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that overexpresses HER2. *N Engl J Med.* 2001;344:783–92.
 19. Dawood S, Broglio K, Buzdar AU, Hortobagyi GN, Giordano SH. Prognosis of women with metastatic breast cancer by HER2 status and trastuzumab treatment: an institutional-based review. *J Clin Oncol.* 2010;28:92–8.
 20. National Comprehensive Cancer Network: NCCN clinical practice guidelines in Oncology, Gastric Cancer V.2.2011. http://www.nccn.org/professionals/physician_gls/f_guidelines.asp.
 21. de Gramont A, Buyse M, Abrahantes JC, Burzykowski T, Quinaux E, Cervantes A, et al. Reintroduction of oxaliplatin is associated with improved survival in advanced colorectal cancer. *J Clin Oncol.* 2007;25:3224–9.
 22. Grothey A, Sugrue MM, Purdie DM, Dong W, Sargent D, Hedrick E, et al. Bevacizumab beyond first progression is associated with prolonged overall survival in metastatic colorectal cancer: results from a large observational cohort study (BRiTE). *J Clin Oncol.* 2008;26:5326–34.
 23. Shitara K, Matsuo K, Mizota A, Kondo C, Nomura M, Takahari D, et al. Association of fluoropyrimidines, platinum agents, taxanes, and irinotecan in any line of chemotherapy with survival in patients with advanced gastric cancer. *Gastric Cancer.* 2011;14:155–60.
 24. Shah MA, Kang YK, Ohtsu A, Delmar P, Foernzler D, Langer B, et al. Tumour biomarker analyses in the AVAGAST phase III randomized study of first-line bevacizumab + capecitabine/cisplatin in patients with advanced gastric cancer. *Ann Oncol.* 2010;21 suppl 8:viii67–viii68 (abstr 174PD).
 25. von Minckwitz G, du Bois A, Schmidt M, Maass N, Cufer T, de Jongh FE, et al. Trastuzumab beyond progression in human epidermal growth factor receptor 2-positive advanced breast cancer: a German Breast Group 26/Breast International Group 03–05 study. *J Clin Oncol.* 2009;27:1999–2006.
 26. Blackwell KL, Burstein HJ, Storniolo AM, Rugo H, Sledge G, Koehler M, et al. Randomized study of Lapatinib alone or in combination with trastuzumab in women with ErbB2-positive, trastuzumab-refractory metastatic breast cancer. *J Clin Oncol.* 2010;28:1124–30.
 27. Hernán MA, Brumback B, Robins JM. Marginal structural models to estimate the joint effect of non-randomized treatments. *J Am Stat Assoc.* 2001;96:440–8.

Preservation of peritoneal fibrinolysis owing to decreased transcription of plasminogen activator inhibitor-1 in peritoneal mesothelial cells suppresses postoperative adhesion formation in laparoscopic surgery

Manabu Shimomura, MD, Takao Hinoi, MD, Satoshi Ikeda, MD, Tomohiro Adachi, MD, Yasuo Kawaguchi, MD, Masakazu Tokunaga, MD, Tatsunari Sasada, MD, Hiroyuki Egi, MD, Kazuaki Tanabe, MD, Masazumi Okajima, MD, and Hideki Ohdan, MD, Hiroshima, Japan

Background. Postoperative adhesion formation is regulated by peritoneal fibrinolysis, which is determined by tissue-type plasminogen activator (tPA) and plasminogen activator inhibitor-1 (PAI-1). This study compared peritoneal fibrinolysis and adhesion formation after laparoscopic surgery (LAP) and open surgery (OP).

Methods. We divided 154 male rats into 3 groups after cecal cauterization: Control, no treatment; LAP, CO₂ pneumoperitoneum at 5 mmHg for 60 minutes; and OP, laparotomy for 60 minutes. Adhesions were quantified at day 7. The activity and mRNA level of tPA and PAI-1 were determined by enzyme-linked immunosorbent assay in plasma and peritoneal lavage and by real-time polymerase chain reaction in peritoneal mesothelial cells from omentum. We also examined peritoneal fibrinolysis in human gastric cancer patients treated with LAP (n = 14) or OP (n = 10).

Results. In the animal study, adhesion scores, PAI-1 activity in peritoneal lavage fluid, and PAI-1 mRNA levels in peritoneal mesothelium were significantly greater in the OP group than the control and LAP groups. In the human study, postoperative PAI-1 mRNA levels were significantly greater in the OP group than the LAP group. Additionally, PAI-1 mRNA levels and subsequent adhesion formation were induced by prolonged operative time in the OP group, but not the LAP group.

Conclusion. Preservation of peritoneal fibrinolysis owing to decreased PAI-1 expression at the transcriptional level in peritoneal mesothelial cells is associated with suppression of postoperative adhesion formation in LAP. PAI-1 mRNA levels and subsequent adhesion formation were not induced by prolonged operative time in LAP. These results highlight the less invasiveness nature of LAP. (Surgery 2013;153:344-56.)

From the Department of Gastroenterological and Transplant Surgery, Applied Life Sciences, Institute of Biomedical and Health Sciences, Hiroshima University, Hiroshima, Japan

IN RECENT DECADES, LAPAROSCOPIC TECHNIQUES have been integrated into common surgical practices. The benefits of laparoscopic surgery (LAP) in comparison with open surgery (OP) include

decreased pain, faster recovery, lesser hospital stay, possibly decreased immunosuppression, and decreased morbidity, such as small bowel obstruction.¹ Previous reports have investigated the influence of CO₂ pneumoperitoneum on the human body²; however, the effects of the lower invasiveness of LAP have not yet been fully determined.

Abdominal adhesion formation occurs in 67–93% of abdominal operations, representing a substantial clinical and financial problem.³ Postoperative adhesions have increased risk of long-term sequelae, including small bowel obstruction, chronic pain, infertility, and difficult reoperative surgery.⁴ Because laparoscopic procedures result

Accepted for publication July 25, 2012.

Reprint requests: Takao Hinoi, MD, Department of Gastroenterological and Transplant Surgery, Applied Life Sciences, Institute of Biomedical and Health Sciences, Hiroshima University, 1-2-3 Kasumi, Minami-ku, Hiroshima 734-8551, Japan. E-mail: thinoi@hiroshima-u.ac.jp.

0039-6060/\$ - see front matter

© 2013 Mosby, Inc. All rights reserved.

<http://dx.doi.org/10.1016/j.surg.2012.07.037>

in a more gentle manipulation of tissues, less bleeding, and less contamination with foreign bodies than conventional OP, as well as a smaller skin incision, LAP has potential theoretic advantages over OP in decreasing adhesion formation,⁵⁻⁸ but very few comparative studies are available to demonstrate this result.⁸

Substantial evidence indicates that formation of adhesions after surgery is determined by the balance between deposition and degradation of fibrin in the early phase of peritoneal tissue repair.^{4,9-12} Adhesions are formed when the peritoneum is damaged and the basal membrane of the mesothelial layer is exposed to the surrounding tissues. This injury to peritoneum elicits a local inflammatory response, which leads to the formation of fibrin-rich exudates as part of the hemostatic and traumatic process. In the presence of sufficient peritoneal fibrinolytic activity, resolution of the fibrin strands occurs. If peritoneal fibrinolytic activity is suppressed, however, organization of the fibrin matrix and cellular and vascular ingrowth ensues, leading to fibrous bands between the structures. Fibrinolysis is activated mainly by the plasminogen activator (tPA) and is inhibited by plasminogen activator inhibitor-1 (PAI-1) in the peritoneal cavity, both of which are produced primarily by peritoneal mesothelial cells.^{4,13}

Conventional abdominal surgery is accompanied by a rapid decrease in peritoneal fibrinolytic activity, which may lead to peritoneal adhesion formation.¹⁴ Several studies have investigated the alterations of peritoneal fibrinolysis during LAP.¹⁵⁻¹⁹ These reports indicated that peritoneal fibrinolysis was also disturbed by LAP, similar to conventional OP. A few clinical studies have aimed to clarify the difference in peritoneal fibrinolysis between OP and LAP; however, their results remain controversial.²⁰⁻²⁴

Given this background, we hypothesized that less impairment of peritoneal fibrinolysis after LAP could explain the fewer adhesions compared with OP. The aim of this study was to clarify the peritoneal fibrinolytic activity and subsequent adhesion formation after LAP compared with OP using an animal model and in human patients.

MATERIALS AND METHODS

Animal study. *Animals:* The study was performed using 8- to 10-week-old male F344 rats (CLEA Japan, Tokyo, Japan) that weighed 180–250 g. The animals were housed at a constant room temperature with 12-hour light and dark cycles and were provided standard rodent chow and water ad libitum. The animal study was approved

by the Hiroshima University Animal Care and Use Committee.

Rat model of operative adhesion formation: All operative procedures were performed under sterile conditions. We induced intestinal adhesion by cecal cauterization as described previously.¹² Briefly, rats were anesthetized with subcutaneous injection of 50 mg/kg pentobarbital sodium solution. The abdomen was shaved, and then laparotomy was performed with a 1.5-cm anterior midline incision. The cecum was isolated and cauterized using the coagulation mode of bipolar forceps (MERA; 30 W, 500 kHz) for 1 second at 1 point.

Operative interventions: Rats were divided into the following 3 groups after cecal cauterization: Control, LAP, and OP. In the control group, the incision was closed immediately after cecal cauterization without any other intervention. In the LAP group, the abdominal incision was closed after cecal cauterization. Subsequently, an 18-gauge intravenous catheter was inserted into the right lower quadrant, and the peritoneal space was insufflated with CO₂ to a pressure of 5 mmHg for 60 minutes using an automatic inflator (Olympus, Tokyo, Japan). A pneumoperitoneum pressure of 5 mmHg was selected for the laparoscopic model, because a previous experimental study demonstrated that 5 mmHg pressure is the optimal pressure in a rat model to induce the same physiologic changes seen in humans.²⁵ Another 18-gauge catheter was inserted into the left lower quadrant and connected to a water valve and the insufflation of the gas was balanced with its exit to maintain a constant pressure. At the end of the operation, catheters were removed after release of the inflated gas. In the OP group, the abdominal incision was extended to 4 cm from the xiphoid to pubis after cecal cauterization, and peritoneal cavities were exposed to the air for 60 minutes. Then, the abdominal incision was closed. We also examined pneumoperitoneum or laparotomy for 30 and 120 minutes for both LAP and OP groups to examine the influence of operative time.

Adhesion scoring: The rats were anesthetized and killed 7 days after cecal cauterization, and the abdomen was opened to assess adhesions. Each rat was evaluated according to the following standard scoring system, which has been used widely in this field: 0, no adhesion; 1, one thin filmy adhesion; 2, more than one thin adhesion; 3, thick adhesion with focal point; 4, thick adhesion with plantar attachment or >1 thick adhesion with focal point; and 5, very thick vascularized adhesion or >1 plantar adhesion. Images of adhesion formation

around the cecum were recorded at necropsy without revealing the incision size, and adhesion score was determined by 2 independent observers blinded to the experimental group.

Sample collection and storage. Blood and peritoneal lavage were harvested at the indicated time after cecal cauterization in this model. Blood was collected by aortic puncture. To harvest sufficient peritoneal lavage for examination, 4 mL of sterile phosphate-buffered saline was injected intraperitoneally 0.5 hours before killing. Samples were anticoagulated with 1:10 citrate buffer and centrifuged at $3,000 \times g$, and supernatants were stored at -70°C until analysis. tPA and PAI-1 activity were measured by enzyme-linked immunosorbent assay (ELISA). In addition, total RNA was extracted from omental tissue, and tPA and PAI-1 mRNA expression levels were assessed by quantitative real-time polymerase chain reaction (RT-PCR). Samples of omental tissue were placed immediately in RNAlater (Ambion) and stored at -70°C until RNA extraction, as described below.

Human study. Patients and surgery. We examined prospectively 24 consecutive patients who underwent LAP ($n=14$) or conventional OP ($n=10$) for gastric cancer with curative intent from February to July 2011 at the Hiroshima University Hospital. Only patients who underwent partial gastrectomy (distal or proximal gastrectomy) with standard lymph node resection were included. Exclusion criteria were emergency operation, previous history of abdominal surgery (except for appendectomy), metastatic disease, serosal invasion, and positive peritoneal cytology. In our institution, patients who were expected to have a tumor with invasion to the mucosa or submucosa without lymph node metastasis were selected for LAP. For LAP, 5 trocars were inserted for CO_2 pneumoperitoneum at a pressure of 10 mmHg. Removal of the stomach and reconstruction were performed through a 4-cm transverse incision of the left upper abdomen. For conventional OP, surgery was performed through a midline abdominal incision from the xiphoid to navel. All operative procedures were performed by the same surgical team. All patients were staged according to the American Joint Commission for Cancer Staging (AJCC/TNM, sixth edition) system. The study was approved by the local ethics committee, and written informed consent was obtained from all patients.

Sample collection and storage. Blood samples were collected just before (preoperative) and just after the operation (postoperative), and on the first postoperative day (POD1). Blood samples were anticoagulated with 1:10 citrate buffer and then

immediately put on melting ice. Plasma was separated by centrifugation at $3,000 \times g$ and stored at -70°C until analysis. tPA and PAI-1 activity were measured by ELISA. Omental tissues were harvested at the beginning and the end of the operation to extract RNA. Omental tissues were collected from sites demonstrating no mechanical injury, and resected specimens were not included. Samples of omental tissue were immediately placed in RNAlater and stored at -70°C until RNA extraction. tPA and PAI-1 mRNA levels were determined by quantitative RT-PCR.

Isolation and culture of human peritoneal mesothelial cells. For patients in the OP group ($n=3$), 5-cm^2 samples of omental tissue were harvested at the beginning and end of the operation, and then human peritoneal mesothelial cells (HPMCs) were isolated as described previously.^{26,27} Briefly, pieces of omentum were washed with sterile phosphate-buffered saline and then incubated with shaking in 0.125% trypsin for 10 minutes at 37°C . The tissues were removed, trypsin solution containing free mesothelial cells was centrifuged at $100 \times g$ for 10 minutes, and then the supernatant was discarded. RNA was extracted immediately from this cell pellet, and tPA and PAI-1 mRNA expression was determined by quantitative RT-PCR. In addition, the obtained cell pellets were confirmed to contain HPMCs by the following method: The cell pellets were suspended in M199 medium (Sigma-Aldrich, St. Louis, MO) supplemented with 10% fetal calf serum, 100 U/mL penicillin, and 2 mmol/L L-glutamine and cultured on 10-cm^2 tissue culture flasks coated with rat type I collagen (Nitta Zerin Co. Ltd, Osaka, Japan). We then confirmed that HPMCs were cultured 1 week after tissue harvest. Cells were identified as HPMCs by their typical cobblestone appearance when confluent (see Fig 6, A), as well as by positive staining for cytokeratin and vimentin plus negative staining for factor VIII on immunocytochemistry, as described previously.^{26,27} In this manner, we could determine mRNA levels in peritoneal mesothelium, with minimal contamination by other cell types. We isolated HPMCs from omentum harvested at the beginning and the end of operation in the same way; therefore, we were able to compare mRNA levels between these time points without consideration of phenotypic change owing to isolation.

ELISA: The tPA and PAI-1 activity in plasma and peritoneal lavage was measured with rat and human tPA and PAI-1 activity ELISA kits (Innovative Research, Novi, MI) according to the manufacturer's specifications.




Fig 1. A, Adhesion scores of the 3 rat treatment groups at day 7 after cecal cauterization. The adhesion scores of the OP group were greater than those of both the control and LAP groups (* $P < .05$). B, Relative tPA mRNA level 6 hours after cecal cauterization for the 3 experimental groups. The tPA mRNA level of the LAP and OP groups was greater than that of the control group (* $P < .05$). No difference was observed between the LAP and OP groups. All data are expressed as a ratio of the mean value for the control group (set at 1). C,

Quantitative RT-PCR: Total RNA was extracted from rat and human samples with the RNeasy Mini Kit (Qiagen, Venlo, The Netherlands), and cDNA was synthesized using ReverTraAce qPCR RT kit (Toyobo, Osaka, Japan). We quantified gene expression with TaqMan Gene Expression Assays (Applied Biosystems, Carlsbad, CA). Results were expressed as relative expression standardized to the expression of the gene encoding GAPDH. Specific primers used for quantitative RT-PCR were as follows: PAI-1 (Serp1; Assay ID Rn01481341_m1), tPA (Plat; Assay ID Rn01482584_m1), and GAPDH (Assay ID Rn99999916_s1) for the animal study; and PAI-1 (Serp1; Assay ID, Hs01126606_m1), tPA (Plat; Assay ID Hs00263492_m1), and GAPDH (Assay ID Hs99999905_m1) for the clinical study (Applied Biosystems).

Statistical analysis: All data are expressed as the mean value \pm standard deviation. Differences between 2 groups were analyzed by the Mann-Whitney test for nonpaired comparison and by paired t -test for paired comparison. In all analyses, statistical significance was set at a $P < .05$. All analyses were performed using JMP 8 software (version 8.02, SAS Institute Inc; SAS, Inc, Cary, NC).

RESULTS

Animal study. Adhesion scoring: After cecal cauterization, 27 rats were divided into 3 groups: control ($n = 9$), LAP ($n = 9$), and OP ($n = 9$). We killed the rats 7 days later, and then each rat was evaluated according to the previously described adhesion scoring system. No rats died during the experiment or demonstrated intestinal perforation at the time of study. The adhesion scores of the OP group were greater than those of both the control ($P < .05$) and LAP groups ($P < .05$; Fig 1, A). These results suggest that the assessment of postoperative adhesion formation in this model is reproducible and optimal to compare fibrinolytic activity between LAP and OP groups.

Measurement of tPA and PAI-1 activity in plasma and peritoneal lavage by ELISA: Another 52 rats were divided into control ($n = 16$), LAP ($n = 18$), and OP ($n = 18$) groups after cecal cauterization. All rats were killed, and blood and peritoneal lavage were collected at the end of the operative

Relative PAI-1 mRNA level 6 hours after cecal cauterization for the 3 experimental groups. The PAI-1 mRNA level of the OP group was greater than that of both the control (** $P < .01$) and LAP groups (* $P < .05$). All data are expressed as a ratio of the mean value for the control group (set at 1).




Fig 2. The results of ELISA (animal study). (A) Plasma tPA activity. Plasma tPA activity at 1, 6, and 24 hours was decreased compared with that of non-op in the control, LAP, and OP groups ($*P < .05$). No significant differences were observed between groups at each time point. (B) tPA activity of peritoneal lavage. The tPA activity in peritoneal lavage was low in all 3 groups at each time point. (C) Plasma PAI-1 activity. Plasma PAI-1 activity was increased from non-op to 6 hours in the LAP group ($P < .05$) and OP group ($P < .01$). At 6 hours, PAI-1 activity was greater in the OP group than the control group ($*P < .05$). (D) PAI-1 activity of peritoneal lavage. The PAI-1 activity of peritoneal lavage increased from 1 to 6 hours in the control ($P < .05$), LAP ($P < .01$), and OP groups ($P < .01$). At 6 hours, PAI-1 activity was greater in the OP group than in the control ($**P < .01$) and LAP groups ($*P < .05$).

intervention (1 hour after cecal cauterization; $n = 5$ for each group; 6 hours after cecal cauterization, $n = 6$ for the control group, $n = 8$ for the LAP and OP groups; and 24 hours after cecal cauterization, $n = 5$ for each group). Blood and omental tissue were collected as nonoperated control samples from another 5 rats (non-op; $n = 5$).

Plasma tPA activity at 1, 6, and 24 hours was decreased compared with that of non-op controls in all 3 groups ($P < .05$). No differences were observed between each of the 3 groups at each time point (Fig 2, A). The tPA activity of peritoneal lavage was low in all 3 groups at all time points (Fig 2, B). By contrast, plasma PAI-1 activity increased from non-op to 6 hours in both the LAP ($P < .05$) and OP groups ($P < .01$), and then decreased from 6 to 24 hours. At 6 hours, the PAI-1 activity of the OP group was greater than that of the control group ($P < .05$; Fig 2, C). The PAI-1 activity of peritoneal lavage increased from 1 to 6 hours (control group, $P < .05$; LAP and OP groups, $P < .01$), and then decreased at 24 hours. At 6 hours, the PAI-1 activity of the OP group was greater than that of both the control ($P < .01$) and LAP groups ($P < .05$; Fig 2, D). The PAI-1 activity of

peritoneal lavage demonstrated a dramatic change compared with that of plasma samples, which suggested that induction of PAI-1 activity in the peritoneal cavity was more important for the development of adhesion formation than systemic changes in this model.

Assessment of tPA and PAI-1 gene expression in peritoneal mesothelial cells from omentum: To investigate the transcriptional changes of tPA and PAI-1 in the peritoneal cavity owing to these operative interventions, we examined mRNA expression in omental tissues, which contain high levels of peritoneal mesothelial cells. Peritoneal mesothelial cells are thought to be the primary producer of tPA and PAI-1 in the peritoneal cavity,⁴ and the method for extraction of peritoneal mesothelial cells from omental tissue has been established.^{26,27} We examined first the time course of PAI-1 mRNA expression changes in response to this operative intervention. We measured PAI-1 mRNA levels from omental tissue extracted from non-op control rats ($n = 5$) and at 1 ($n = 5$), 3 ($n = 5$), 6 ($n = 8$), 9 ($n = 5$), and 24 hours ($n = 3$), and 7 days ($n = 3$) after cecal cauterization in the OP group. The PAI-1 mRNA level gradually increased from




Fig 3. Time course analysis of relative PAI-1 mRNA level after cecal cauterization in the OP group. The PAI-1 mRNA level gradually increased from non-op to 6 hours and then decreased to day 7. All data are expressed as a ratio of the mean value for non-op (set at 1).

non-op to 6 hours, and then gradually decreased up to day 7 (Fig 3).

We then examined omental tPA and PAI-1 mRNA levels at 6 hours in each group ($n = 6$ for the control group; $n = 8$ for the LAP and OP groups). The tPA mRNA level of the LAP and OP groups was greater than that of the control group ($P < .05$; Fig 1, B); however, no difference was observed between the LAP and OP groups. By contrast, the PAI-1 mRNA level of the OP group was greater than that of both the control ($P < .01$) and LAP groups ($P < .05$; Fig 1, C). This difference was consistent with the adhesion score results, which suggested a correlation between local PAI-1 mRNA expression levels and postoperative adhesion formation. These results indicated that peritoneal fibrinolysis was better preserved in the LAP group owing to decreased transcription of PAI-1 than the OP group, which contributed to the decrease in postoperative adhesion formation in the rat model.

Correlation between peritoneal fibrinolysis and treatment time. We next examined the relationship between peritoneal fibrinolytic activity and treatment time. We performed pneumoperitoneum or laparotomy for 30 and 120 minutes in both LAP (30 minutes, $n = 8$; 120 minutes, $n = 8$) and OP groups (30 minutes, $n = 8$; 120 minutes, $n = 8$) and determined the PAI-1 mRNA levels in omental tissues at 6 hours after cecal cauterization. PAI-1 mRNA levels were greater in the OP group than the LAP group in the 60- and 120-minute (both $P < .05$) treatment groups. PAI-1 mRNA levels were greater in the 120-minute treatment

group than the 30-minute treatment group ($P < .01$); however, this induction was not observed in the LAP group (Fig 4, A).

Additionally, we examined the relationship between adhesion scores and operative time using another group of rats. We performed pneumoperitoneum or laparotomy for 30 and 120 minutes for both the LAP (30 minutes, $n = 6$; 120 minutes, $n = 8$) and OP groups (30 minutes, $n = 6$; 120 minutes, $n = 8$); adhesion scores were determined at day 7 after cecal cauterization. Adhesion scores were greater in the OP group than the LAP group in the 60- and 120-minute (both $P < .05$) treatment groups. In the OP group, the adhesion score of the 120-minute treatment group was greater than that of the 30-minute treatment group ($P < .05$). This induction, however, was not observed in the LAP group (Fig 4, B). PAI-1 gene expression was highly induced in a time-dependent manner in the OP group, resulting in a greater adhesion score at day 7, whereas its expression was suppressed even with the greater operative time in the LAP group, resulting in a lesser adhesion score.

Human study. Patient characteristics: We next investigated peritoneal fibrinolytic activity after LAP compared with conventional OP in clinical samples. Patient characteristics are summarized in Table I. No difference was observed in age or gender; however, patients with early stage disease were more common in the LAP group than the OP group ($P < .05$). Increased plasma PAI-1 levels have been shown to associate with increased risk of cardiovascular disease, diabetes mellitus, and the metabolic syndrome. Therefore, we compared preoperative body mass index, serum HbA1c level, total cholesterol, and triglyceride between the 2 groups. No difference was observed in these factors. No patients had an anticoagulant disorder, and prothrombin time did not differ between the 2 groups. Both preoperative and POD1 white blood cell count and C-reactive protein level did not differ between the 2 groups. No patients experienced intraoperative complications. No patients in the LAP group were converted to laparotomy.

tPA and PAI-1 activity in plasma determined by ELISA: Blood samples were collected just before (preoperative) and after the operation (postoperative), and on POD1. The results of ELISA are summarized in Table II. Both preoperative tPA activity and PAI-1 activity did not differ between the 2 groups. Postoperative and POD1 tPA concentration demonstrated no change compared with preoperative levels (Table II). By contrast, PAI-1




Fig 4. A, The correlation between PAI-1 mRNA level and treatment time for LAP and OP groups in the rat cecal cauterization model. PAI-1 mRNA level was greater in the OP group than the LAP group in the 60- ($*P < .05$) and 120-minute ($*P < .05$) treatment groups, respectively. PAI-1 mRNA level was greater in the 120-minute treatment group than the 30-minute treatment group ($**P < .05$) in the OP group; however, this induction was not observed in the LAP group. All data are expressed as a ratio of the mean value for the control group (set at 1). B, The correlation between adhesion score and treatment time for the LAP and OP groups. Adhesion scores were greater in the OP group than the LAP group for both the 60- ($*P < .05$) and 120-minute ($*P < .05$) treatment groups. Adhesion scores were greater in the 120-minute treatment group than the 30-minute treatment group in the OP group ($**P < .05$); however, this induction was not observed in the LAP group. Closed bar (■), LAP group; open bar (□), OP group.

Table I. Patient characteristics

	LAP (n = 14)	OP (n = 10)	P value
Age	60 ± 15	69 ± 10	NS
Gender (male/female)	7/7	4/6	NS
BMI	23 ± 3	22 ± 3	NS
Stage I/II/III	13/1/0	5/3/2	<.05
TG	126 ± 63	126 ± 85	NS
T-Chol	214 ± 37	192 ± 36	NS
HbA1c	5.6 ± 0.8	5.6 ± 0.4	NS
PT	99 ± 8	102 ± 11	NS
WBC count (preoperative)	6,300 ± 1,900	5,600 ± 1,300	NS
CRP (preoperative)	0.05 ± 0.06	0.08 ± 0.08	NS
WBC count (POD1)	8,900 ± 2,500	9,200 ± 1,600	NS
CRP (POD1)	5.7 ± 2.1	7.0 ± 2.6	NS
Operative time (min)	308 ± 52	224 ± 32	<.001
Bleeding (mL)	48 ± 95	168 ± 138	<.05

All data are expressed as mean values ± standard deviation.

BMI, Body mass index; CRP, C-reactive protein; T-Chol, total cholesterol; TG, triglyceride; PT, prothrombin time; WBC, white blood cell.

Table II. ELISA results (human study)

	Preoperative	Postoperative	POD1
tPA activity (IU/mL)			
LAP	0.1 ± 0.1	0.1 ± 0.1	0.1 ± 0.1
OP	0.1 ± 0.1	0.1 ± 0.1	0.1 ± 0.1
PAI-1 activity (ng/mL)			
LAP	8.3 ± 15.5	20.9 ± 32.9*	13.4 ± 14.9*
OP	2.9 ± 4.4	21.8 ± 30.7*	7.3 ± 7.3*

*Versus the preoperative value, $P < .05$.

All data are expressed as mean values ± standard deviation.

activity increased between preoperative and postoperative, and then decreased at POD1 ($P < .05$; Table II); however, the LAP and OP groups did not differ at each time point.

tPA and PAI-1 gene expression in omentum determined by quantitative RT-PCR: Omental tissues were harvested at the beginning and the end of the operation to examine tPA and PAI-1 mRNA expression levels. For both groups, the tPA mRNA level was unchanged after the operation. Both preoperative and postoperative tPA mRNA levels also demonstrated no significant difference between the 2 groups (Fig 5, A). By contrast, PAI-1 mRNA levels were significantly increased from pre- to postoperative for both the LAP and OP groups (both $P < .01$). Preoperative PAI-1 mRNA expression did not differ between the 2 groups;




Fig 5. Results of quantitative RT-PCR (human study). *A*, Relative tPA mRNA level. The tPA mRNA expression did not differ between the 2 groups at both pre- and postoperative specimens. All data are expressed as a ratio of the mean preoperative value for the OP group (set at 1). *B*, Relative PAI-1 mRNA level. PAI-1 mRNA level was increased from pre- to postoperative for both the LAP and OP groups (** $P < .001$). The 2 groups did not significantly differ preoperatively; however, the postoperative PAI-1 mRNA level of the OP group was greater than that of the LAP group (* $P < .05$). All data are expressed as a ratio of the mean preoperative value for the OP group (set at 1).

however, the postoperative PAI-1 mRNA level of the OP group was greater than that of the LAP group ($P < .05$; Fig 5, *B*).




Fig 6. *A*, Phase-contrast micrographs of isolated human peritoneal mesothelial cells (HPMCs). *B*, Relative tPA mRNA level of isolated HPMCs. The tPA mRNA level did not differ between pre- and postoperative specimens. All data are expressed as a ratio of the mean preoperative value (set at 1). *C*, Relative PAI-1 mRNA level for isolated HPMCs. Postoperative PAI-1 mRNA level tended to be greater than the preoperative level, to a similar extent as that observed in omental tissues. All data are expressed as a ratio of the mean preoperative value (set at 1).

tPA and PAI-1 gene expression in isolated HPMCs from omental tissue. To confirm the induction of PAI-1 mRNA in HPMCs from omental tissue in

response to abdominal operation, we harvested 5-cm² samples of omental tissue at the beginning and the end of the operation and then extracted HPMCs from each sample from patients in the OP group ($n = 3$). We only examined the OP group, because obtaining a sufficient amount of omental tissue to extract the HPMCs was difficult in the LAP group owing to the small skin incision. We extracted total RNA from HPMCs and analyzed tPA and PAI-1 mRNA expression by quantitative RT-PCR. tPA mRNA expression did not change after the operation (Fig 6, B). Consistent with the induction of PAI-1 mRNA expression in omentum, PAI-1 mRNA expression seemed to be upregulated in HPMCs after the operation (Fig 6, C).

Correlation between PAI-1 gene expression and operative time. We investigated the possible correlation of postoperative PAI-1 mRNA levels with several clinical parameters and found that high PAI-1 mRNA levels were correlated with operative time in the OP group ($P = .02$; $r = 0.67$; Fig 7). No correlation was observed in the LAP group ($P = .56$; $r = 0.17$; Fig 7).

Consistent with our findings in the animal study, these results suggest that achieving a more prolonged hypofibrinolytic state by inhibition of PAI-1 upregulation in peritoneal mesothelial cells during LAP may predispose patients to less intestinal adhesions than OP.

DISCUSSION

A considerable body of data now supports the notion that formation of adhesions after abdominal operations is determined by peritoneal fibrinolytic activity.^{1,4,12,15,16,18,19,28} The specific condition of a pneumoperitoneum, especially the warm and humid intra-abdominal atmosphere, has been hypothesized to preserve peritoneal fibrinolytic activity in LAP better than in conventional OP^{1,8}; however, the extent of this hypofibrinolysis after LAP compared with OP has remained controversial.

The underlying molecular mechanisms that control the peritoneal fibrinolytic system have been clarified in vivo and in vitro. Molinas et al¹⁷ reported that PAI-1 was induced by pneumoperitoneum and identified upregulation of PAI-1 as a mechanism for adhesion formation, because pneumoperitoneum-enhanced adhesions were not observed in *PAI-1* knockout mice. Using an in-vitro pneumoperitoneum model, Bergstrom et al¹⁶ found that cultured HPMCs demonstrated increased synthesis and release of PAI-1 on exposure to CO₂. Ziprin et al¹⁸ reported that incubation of HPMCs with CO₂ enhances mesothelial cell




Fig 7. Scatter plots of relative PAI-1 mRNA level versus operative time in the OP and LAP groups. PAI-1 mRNA expression level and operative time were correlated in the OP group ($r = 0.67$, $P = .02$). By contrast, PAI-1 mRNA expression level and operative time were not correlated in the LAP group ($r = 0.17$, $P = .56$). Closed circle (●) and solid line, LAP group; open circle (○) and dotted line, OP group.

fibrinolytic activity because of a decrease in PAI-1 concentration as compared with standard culture conditions.

Some clinical studies have compared the alteration of peritoneal fibrinolytic activity between LAP and OP and concluded that both operative techniques had similar effects.²¹⁻²⁴ Brokelman et al²⁴ reported that peritoneal hypofibrinolysis via decreased tPA activity seemed to induce more rapidly during conventional surgery than LAP. All these previous reports analyzed only protein levels from plasma or intraoperative peritoneal biopsies; transcriptional regulation of tPA and PAI-1 in peritoneal mesothelial cells has not been analyzed previously. Because tissue samples to assess the peritoneal fibrinolytic activity in these studies were obtained from peritoneum of the abdominal wall during operation, the quality and quantity of mesothelial cells, which are more abundant in omentum, may have been limited. Moreover, PAI-1 activity and PAI-1 antigen/protein concentration after transcription/translation of the PAI-1 gene may not change significantly, because we detected in the rapid increase of PAI-1 mRNA during operation, presumably owing to the limited duration of the time course in these studies. Further potential biases may have occurred as a result of the operative procedure, skill of the surgeon, blood loss, and the selection of patients suitable for LAP in these clinical studies. By examining transcriptional




Fig 8. Schema of transcriptional regulation of PAI-1 in peritoneal mesothelial cells for postoperative adhesion formation. *A*, Postoperative adhesions were administered by peritoneal hypofibrinolysis, which induced transcriptional upregulation of PAI-1 in peritoneal mesothelial cells in response to surgical stress caused by OP. *B*, Decreased transcription of PAI-1 in peritoneal mesothelial cells suppresses postoperative adhesion formation in LAP. Although the mechanism of transcriptional decrease in PAI-1 in peritoneal mesothelial cells in LAP has not yet been clarified, pneumoperitoneum using CO₂ may affect peritoneal mesothelial cells. *FDP*, Fibrin degradation products.

levels of tPA and PAI-1 in peritoneal mesothelial cells, we first demonstrated that decreased expression of PAI-1 transcript in peritoneal mesothelial cells suppresses postoperative adhesion formation in LAP using animal and clinical samples. Our proposed schema based on our findings is indicated in Fig 8. Although the mechanism of decreased PAI-1 transcript expression in peritoneal mesothelial cells after LAP has not yet been clarified, pneumoperitoneum using CO₂ may affect the reduction of peritoneal mesothelial cells.

In the present study, we utilized a unique model of intestinal adhesion formation,¹² with cecal cauterization to establish intestinal adhesions in rats. This model could minimize the technical bias of cecal cauterization and provided better reproducibility than previously described models of intestinal adhesion formation, such as the cecal abrasion model (in which the cecum is abraded by scrubbing with surgical gauze).^{29,30} Our data demonstrated that quantification of adhesion at day 7 reflects the status of small intestinal adhesion with good reproducibility. To compare solely the difference in peritoneal fibrinolytic activity

between pneumoperitoneum and laparotomy, we aimed to minimize the effect of technical bias between LAP and OP for cecal cauterization. For this reason, we induced cecal cauterization in the same manner with a small skin incision across all experimental groups, and we defined rats treated with pneumoperitoneum after cecal cauterization as the LAP group. We defined adequately the control group (only cecal cauterization without any intervention); therefore, we were able to directly compare the difference between pneumoperitoneum and laparotomy after the same extent of peritoneal injury through the use of this control group. The adhesion score, PAI-1 activity in peritoneal lavage, and PAI-1 mRNA expression in peritoneal mesothelium were greater in the OP group than the control and LAP groups. By contrast, tPA was activated at the transcriptional level in peritoneal mesothelial cells; however, tPA activity in plasma and peritoneal lavage was suppressed by PAI-1, which demonstrated a more drastic effect on fibrinolysis than tPA as shown in Fig 2. Moreover, PAI-1 mRNA expression level and subsequent adhesion formation were induced by prolonged

operative time in the OP group but not the LAP group in this model. To the best of our knowledge, this is the first animal model to demonstrate that preservation of peritoneal fibrinolysis by a decrease in transcription of PAI-1 in peritoneal mesothelial cells after LAP may contribute to the decrease of postoperative adhesion formation (Fig 8). Consistent with this rat model, tPA mRNA expression levels were similar in both OP and LAP groups in human patients, although PAI-1 mRNA expression level was significantly greater in the OP group. Furthermore, PAI-1 mRNA levels and subsequent adhesion formation were significantly induced by prolonged operative time in the OP group but not in the LAP group. To further confirm our hypotheses in this clinical study, we extracted HPMCs from omentum during operation. Few molecular markers that indicate clearly the efficacy of LAP compared with OP have been reported previously. We identified PAI-1 mRNA in peritoneal mesothelium as such a molecular marker and believe this finding to be of clinical importance.

Based on our animal model without manual manipulation during laparotomy or pneumoperitoneum, physiologic and/or biologic responses under pneumoperitoneum may lead to the observed effects on peritoneal mesothelial cells directly or via indirect effects by changing arterial or venous blood flow or pH levels in vivo. Sawdey et al³¹ reported that the treatment of mice with lipopolysaccharide, tumor necrosis factor- α , or transforming growth factor- β increased the steady-state levels of PAI-1 mRNA within 3 hours in all tissues examined. Other studies reported that statins (HMG-CoA reductase inhibitors) decreased postoperative adhesions by increasing peritoneal fibrinolytic activity.^{32,33} In a rat model, lovastatin and atorvastatin significantly decreased adhesion formation without affecting anastomotic burst pressure by upregulating tPA expression and activity in the peritoneum. Statins increased tPA and decreased PAI-1 production in cultured human mesothelial cells, and these effects were reversed by their inhibitors, although this effect may be mediated by inactivation of RhoA, a small GTPase, leading to increased tPA expression and activity, not PAI-1 expression. The antioxidant NAC (*N*-acetyl-L-cysteine) is a clinically relevant antioxidant that donates its L-cysteine moiety to the synthesis of the intracellular antioxidant glutathione. In a rat model, intraperitoneal administration of NAC increased the tPA/PAI-1 protein ratio in peritoneal tissue and peritoneal fluid through a decrease in

PAI-1 mRNA and protein expression, suggesting that decreased peritoneal oxidative stress downregulates PAI-1 gene expression to result in a fibrinolytic state.³⁴ In endothelial cells, oxidative stress upregulates PAI-1 expression through a NADPH oxidase-dependent nuclear factor- κ B signaling pathway.

PAI-1 is well known as the main regulator of fibrinolysis, and thus, increased plasma PAI-1 levels are associated with the formation or progression of atherosclerosis.³⁵ In the present study, therefore, we confirmed that preoperative body mass index, HbA1c, total cholesterol, triglyceride, and prothrombin time values demonstrated no difference between LAP and OP patients. In addition, elevated PAI-1 was found to be related to poor prognosis of several malignancies.³⁶ In the present human study, the LAP group contained significantly more patients with early stage disease than the OP group, owing to their indication for operation. To minimize this effect, we excluded patients with metastatic diseases, serosal invasion, and positive peritoneal cytology. PAI-1 mRNA levels in patients with advanced stage disease were not greater than those with early stage disease in the present study (data not shown); therefore, this difference in disease stage was not a substantive confounder of our results.

In our institution, only patients who were expected to have invasion to the mucosa or submucosa without lymph node metastasis were selected for LAP (T1 N0 stage IA and T2 N0 stage IB). Therefore, in this study, we examined prospectively consecutive patients according to this selection criterion who were treated by the same surgical team without selection bias by the surgeons. In addition, in the present human study we examined samples obtained prospectively from relatively early-stage gastric cancer patients who underwent standard partial gastrectomy. Further exploration is required to determine whether patients undergoing gastric cancer resection differ from those undergoing operation (laparoscopic or open) for other types of abdominal operations.

In conclusion, the results of the present study suggest strongly that achieving a more prolonged hypofibrinolytic state by inhibition of PAI-1 upregulation during LAP may predispose patients to less intestinal adhesion than OP. Elucidation of the precise mechanisms underlying the downregulation of PAI-1 gene expression during LAP may reveal a potential therapeutic target without inhibiting the fibrin formation that is indispensable for wound healing in response to operative trauma.

The authors thank Dr Yukio Yokoyama (Department of Nephrology, Hiroshima University Hospital) for instruction in the isolation and culture of HPMCs, and the Analysis Center of Life Science, Hiroshima University, for the use of their facilities.


REFERENCES

1. Brokelman WJ, Lensvelt M, Borel Rinkes IH, Klinkenbijl JH, Reijnen MM. Peritoneal changes due to laparoscopic surgery. *Surg Endosc* 2011;25:1-9.
2. Yoshida M, Ikeda S, Sumitani D, Takakura Y, Yoshimitsu M, Shimomura M, et al. Alterations in portal vein blood pH, hepatic functions, and hepatic histology in a porcine carbon dioxide pneumoperitoneum model. *Surg Endosc* 2010;24:1693-700.
3. Ellis H, Moran BJ, Thompson JN, Parker MC, Wilson MS, Menzies D, et al. Adhesion-related hospital readmissions after abdominal and pelvic surgery: a retrospective cohort study. *Lancet* 1999;353:1476-80.
4. Cheong YC, Laird SM, Li TC, Shelton JB, Ledger WL, Cooke ID. Peritoneal healing and adhesion formation/ref-ormation. *Hum Reprod Update* 2001;7:556-66.
5. Krahenbuhl L, Schafer M, Kuzinkovas V, Renzulli P, Baer HU, Buchler MW. Experimental study of adhesion formation in open and laparoscopic fundoplication. *Br J Surg* 1998;85:826-30.
6. Garrard CL, Clements RH, Nanney L, Davidson JM, Richards WO. Adhesion formation is reduced after laparoscopic surgery. *Surg Endosc* 1999;13:10-3.
7. Dowson HM, Bong JJ, Lovell DP, Worthington TR, Karanjia ND, Rockall TA. Reduced adhesion formation following laparoscopic versus open colorectal surgery. *Br J Surg* 2008;95:909-14.
8. Gutt CN, Oniu T, Schemmer P, Mehrabi A, Buchler MW. Fewer adhesions induced by laparoscopic surgery? *Surg Endosc* 2004;18:898-906.
9. Falk K, Bjorquist P, Stromqvist M, Holmdahl L. Reduction of experimental adhesion formation by inhibition of plasminogen activator inhibitor type 1. *Br J Surg* 2001;88:286-9.
10. Hellebrekers BW, Kooistra T. Pathogenesis of postoperative adhesion formation. *Br J Surg* 2011;98:1503-16.
11. Holmdahl L. Making and covering of surgical footprints. *Lancet* 1999;353:1456-7.
12. Kosaka H, Yoshimoto T, Fujimoto J, Nakanishi K. Interferon-gamma is a therapeutic target molecule for prevention of postoperative adhesion formation. *Nat Med* 2008;14:437-41.
13. van Hinsbergh VW, Kooistra T, Scheffer MA, Hajo van Bockel J, van Muijen GN. Characterization and fibrinolytic properties of human omental tissue mesothelial cells. Comparison with endothelial cells. *Blood* 1990;75:1490-7.
14. Holmdahl L. The role of fibrinolysis in adhesion formation. *Eur J Surg Suppl* 1997;577:24-31.
15. Nagelschmidt M, Gerbecks D, Minor T. The impact of gas laparoscopy on abdominal plasminogen activator activity. *Surg Endosc* 2001;15:585-8.
16. Bergstrom M, Falk P, Holmdahl L. CO2 promotes plasminogen activator inhibitor type 1 expression in human mesothelial cells. *Surg Endosc* 2003;17:1818-22.
17. Molinas CR, Elkalan O, Campo R, Luttun A, Carmeliet P, Koninckx PR. Role of the plasminogen system in basal adhesion formation and carbon dioxide pneumoperitoneum-enhanced adhesion formation after laparoscopic surgery in transgenic mice. *Fertil Steril* 2003;80:184-92.
18. Ziprin P, Ridgway PF, Peck DH, Darzi AW. Laparoscopic-type environment enhances mesothelial cell fibrinolytic activity in vitro via a down-regulation of plasminogen activator inhibitor-1 activity. *Surgery* 2003;134:758-65.
19. Bergstrom M, Falk P, Holmdahl L. Effect of acidosis on expression of mesothelial cell plasminogen activator inhibitor type-1. *Surg Endosc* 2006;20:1448-52.
20. Bergstrom M, Ivarsson ML, Holmdahl L. Peritoneal response to pneumoperitoneum and laparoscopic surgery. *Br J Surg* 2002;89:1465-9.
21. Neudecker J, Junghans T, Ziemer S, Raue W, Schwenk W. Effect of laparoscopic and conventional colorectal resection on peritoneal fibrinolytic capacity: a prospective randomized clinical trial. *Int J Colorectal Dis* 2002;17:426-9.
22. Neudecker J, Junghans T, Ziemer S, Raue W, Schwenk W. Prospective randomized trial to determine the influence of laparoscopic and conventional colorectal resection on intravasal fibrinolytic capacity. *Surg Endosc* 2003;17:73-7.
23. Neudecker J, Junghans T, Raue W, Ziemer S, Schwenk W. Fibrinolytic capacity in peritoneal fluid after laparoscopic and conventional colorectal resection: data from a randomized controlled trial. *Langenbecks Arch Surg* 2005;390:523-7.
24. Brokelman W, Holmdahl L, Falk P, Klinkenbijl J, Reijnen M. The peritoneal fibrinolytic response to conventional and laparoscopic colonic surgery. *J Laparoendosc Adv Surg Tech A* 2009;19:489-93.
25. Avital S, Itah R, Szomstein S, Rosenthal R, Inbar R, Sckornik Y, et al. Correlation of CO₂ pneumoperitoneal pressures between rodents and humans. *Surg Endosc* 2009;23:50-4.
26. Yokoyama Y, Masaki T, Kiribayashi K, Nakashima A, Kokoroishi K, Ogawa T, et al. 15-Deoxy-delta12,14-prostaglandin J2 inhibits angiotensin II-induced fibronectin expression via hepatocyte growth factor induction in human peritoneal mesothelial cells. *Ther Apher Dial* 2010;14:43-51.
27. Ogata S, Yorioka N, Kohno N. Glucose and prednisolone alter basic fibroblast growth factor expression in peritoneal mesothelial cells and fibroblasts. *J Am Soc Nephrol* 2001;12:2787-96.
28. Matsuzaki S, Botchorishvili R, Jardon K, Maleysson E, Canis M, Mage G. Impact of intraperitoneal pressure and duration of surgery on levels of tissue plasminogen activator and plasminogen activator inhibitor-1 mRNA in peritoneal tissues during laparoscopic surgery. *Hum Reprod* 2011;26:1073-81.
29. Chung DR, Chitnis T, Panzo RJ, Kasper DL, Sayegh MH, Tzianabos AO. CD4+Tcells regulate surgical and postinfectious adhesion formation. *J Exp Med* 2002;195:1471-8.
30. Holsti MA, Chitnis T, Panzo RJ, Bronson RT, Yagita H, Sayegh MH, et al. Regulation of postsurgical fibrosis by the programmed death-1 inhibitory pathway. *J Immunol* 2004;172:5774-81.
31. Sawdey MS, Loskutoff DJ. Regulation of murine type 1 plasminogen activator inhibitor gene expression in vivo. Tissue specificity and induction by lipopolysaccharide, tumor necrosis factor-alpha, and transforming growth factor-beta. *J Clin Invest* 1991;88:1346-53.
32. Haslinger B, Kleemann R, Toet KH, Kooistra T. Simvastatin suppresses tissue factor expression and increases fibrinolytic activity in tumor necrosis factor-alpha-activated human peritoneal mesothelial cells. *Kidney Int* 2003;63:2065-74.
33. Aarons CB, Cohen PA, Gower A, Reed KL, Leeman SE, Stucchi AF, et al. Statins (HMG-CoA reductase inhibitors)

decrease postoperative adhesions by increasing peritoneal fibrinolytic activity. *Ann Surg* 2007;245:176-84.

34. Chu DI, Lim R, Heydrick S, Gainsbury ML, Abdou R, D'Addese L, et al. N-acetyl-L-cysteine decreases intra-abdominal adhesion formation through the upregulation of peritoneal fibrinolytic activity and antioxidant defenses. *Surgery* 2011;149:801-12.
35. Kohler HP, Grant PJ. Plasminogen-activator inhibitor type 1 and coronary artery disease. *N Engl J Med* 2000;342:1792-801.
36. Sakakibara T, Hibi K, Koike M, Fujiwara M, Kodera Y, Ito K, et al. Plasminogen activator inhibitor-1 as a potential marker for the malignancy of gastric cancer. *Cancer Sci* 2006;97:395-9.

I.V. League



"No, we don't have a clue how the operation really works but I recommend it strongly"



TECHNICAL INNOVATIONS

Open Access

Single-port laparoscopic colectomy versus conventional laparoscopic colectomy for colon cancer: a comparison of surgical results

Hiroyuki Egi^{1,2*}, Minoru Hattori², Takao Hinoi¹, Yuji Takakura¹, Yasuo Kawaguchi¹, Manabu Shimomura¹, Masakazu Tokunaga¹, Tomohiro Adachi¹, Takashi Urushihara³, Toshiyuki Itamoto³ and Hideki Ohdan¹

Abstract

Background: Single-port laparoscopic surgery is a new technique that leaves no visible scar. This new technique has generated strong interest among surgeons worldwide. However, single-port laparoscopic colon surgery has not yet been standardized. Our aim in this study was to evaluate the feasibility of single-port laparoscopic colectomy compared with conventional laparoscopic colectomy for colon cancer.

Methods: We conducted a case-matched, controlled study comparing single-port laparoscopic colectomy to conventional laparoscopic colectomy for right-sided colon cancer.

Results: A total of ten patients were included for the single-port laparoscopic colectomy (S-LAC) group and ten patients for the conventional laparoscopic colectomy (C-LAC) group. The length of the skin incision in the S-LAC group was significantly shorter than that of the C-LAC group.

Conclusion: Our early experiences indicated that S-LAC for right-sided colon cancer is a feasible and safe procedure and that S-LAC results in a better cosmetic outcome.

Keywords: Single-port laparoscopic surgery, Single-incision laparoscopic surgery, Conventional laparoscopic surgery, Laparoscopic colectomy, Colon cancer, Gelpert

Background

Laparoscopic surgery has been a standard strategy for a variety of gastrointestinal diseases. The first report about laparoscopic colectomy was published by Jacobs *et al.* [1] two decades ago. Since then the use of laparoscopic colectomy for colon cancer has gradually increased, and it is now acceptable treatment not only for early colon cancer but also for advanced cases because of its oncological safety and feasibility [2,3]. Recently, natural orifice transluminal endoscopic surgery (NOTES) has been studied as the next generation of minimally invasive surgery. This new technique was described for the first time by Kalloo *et al.*, who introduced their work performing transgastric peritoneoscopy in a porcine model [4]. Marescaux *et al.*

also reported successful NOTES in a clinical case [5]. However, the feasibility and safety of NOTES have not been evaluated. Single-port laparoscopic surgery is also a new technique which leaves no visible scar. This new technique has generated interest among surgeons worldwide. Although the use of single-port laparoscopic cholecystectomy has spread rapidly, single-incision laparoscopic colon surgery has not yet been standardized. Our aim in this study was to evaluate the feasibility of single-port laparoscopic colectomy compared with conventional laparoscopic colectomy for colon cancer which requires D2 lymph node dissection.

Methods

This study was performed with permission of the Ethics Committee of the Hiroshima University.

We conducted a case-matched, controlled study comparing single-port laparoscopic colectomy to conventional laparoscopic colectomy for right-sided colon

* Correspondence: hiroegi@yahoo.co.jp

¹Department of Surgery, Division of Frontier Medical Science, Programs for Biomedical Research, Graduate School of Biomedical Sciences, Hiroshima University, 1-2-3 Kasumi, Minami-ku, Hiroshima 734-8551, Japan
Full list of author information is available at the end of the article