

Table 2 Response

	CR	PR	SD	PD	Overall response
Total	5	13	5	0	18 (78.3 %)
TC	1	3	0	0	4 (100 %)
DC	4	10	5	0	14 (73.7 %)

CR complete response, PR partial response, SD stable disease, PD progressive disease, TC paclitaxel + carboplatin, DC docetaxel + carboplatin

Table 3 Adverse events of TC/DC therapy

<i>n</i> = 23	Grade					≥3 (%)
	1	2	3	4		
Leukopenia	2	9	11	1	12 (52.2)	
Neutropenia	1	1	7	14	21 (91.3)	
Thrombocytopenia	11	0	0	0	0	
Anemia	11	12	0	0	0	
Nausea	11	3	1	0	1 (4.3)	
Vomiting	5	3	0	0	0	
Diarrhea	2	0	0	0	0	
Neurotoxicity	18	0	0	0	0	
Dyspnea	3	0	0	0	0	
Fibrile neutropenia	0	0	2	0	2 (8.7)	

TC paclitaxel + carboplatin, DC docetaxel + carboplatin

Surgery completion and adjuvant therapy

Radical hysterectomy after NAC was completed in 18 of the 23 patients, giving a surgery completion rate of 78.3 %. Adjuvant therapy after radical hysterectomy consisted of no treatment in 3 cases (13.0 %), radiotherapy in 2 cases (8.7 %), chemotherapy in 15 cases (65.2 %), and CCRT in 3 cases (13.0 %).

Survival

The median follow-up period was 31 months (range 9–90 months). The median progression-free survival period was 26 months (95 % CI, 13.5–38.5 months), and the median overall survival period was 35 months (95 % CI,

20.9–49.1 months) (Fig. 1). The 5 patients in whom surgery was not complete died of their primary disease within 35 months. Their median PFS and OS were 8 months (3–12 months) and 21 months (10–35 months), respectively.

Discussion

The incidence of non-squamous cell carcinoma of the uterine cervix has been steadily rising in Japan, currently accounting for approximately 10–15 % of all cervical cancer cases. Lymph node metastasis is more frequent with this disease, compared with invasive squamous cell carcinoma [11], and its sensitivity to radiotherapy and chemotherapy is considered to be lower [12]. Thus, squamous and non-squamous cell carcinomas must be analyzed separately. It is advisable and desirable to try new therapeutic strategies in non-squamous cell carcinoma, but the number of published studies involving this type of cervical cancer is small, with the number of cases analyzed in these reports also small. Thus, no high-level evidence regarding treatment has been obtained for this type of cervical carcinoma.

The response rates of adenocarcinoma are reportedly 20 % to cisplatin [13], 15 % to ifosfamide [14], 14 % to 5-fluorouracil [15], and 12 % to oral etoposide [16]; these response rates are lower than those of squamous cell carcinoma. According to Curtin et al. [17], however, the response rate of adenocarcinoma to paclitaxel is as high as 31 %, even when the agent is used independently. Docetaxel has also been attracting interest as an agent of NAC. Nagao et al. evaluated the efficacy of combined chemotherapy using a DC regimen (docetaxel 60 mg/m² and carboplatin at AUC 6 on day 1, repeating the combination every 21 days) in 17 patients with advanced or recurrent cervical cancer, including 6 with adenocarcinoma and 1 with adenosquamous carcinoma. A partial response was obtained in 6 of the 7 cases with adenocarcinoma (including the case of adenosquamous carcinoma); the response rate was 86 % [18]. Considering these findings, we conducted a pilot study involving standard regimens of TC and DC, conventionally used for the treatment of ovarian cancer.

Fig. 1 Kaplan–Meier curves for progression-free survival (a) and overall survival (b). The median PFS for all patients was 26 months (95 % CI, 13.5–38.5 months), and the median OS was 35 months (95 % CI, 20.9–49.1 months)




Table 4 Treatment results and outcomes of all patients

Patients	Age	Stage	Cell type	Regimen	Cycles	Responses	Surgery	Adjuvant	Follow-up period (months)	PFS (months)	OS (months)	Outcome
1	52	Ib	ASC	DC	2	CR	Incomplete	CCRT	21	12	21	DOD
2	50	Ib2	MAC	TC	2	CR	Complete	NT	90	90	90	NED
3	55	Ib	ASC	DC	2	CR	Complete	CT	62	62	62	NED
4	39	Ib2	MAC	DC	3	CR	Complete	CT	51	11	51	AWD
5	36	Ib2	MAC	DC	2	CR	Complete	CT	22	22	22	NED
6	32	Ib	ASC	DC	3	PR	Incomplete	NT	19	11	19	DOD
7	49	Ib2	MAC	DC	2	PR	Complete	CCRT	78	78	78	NED
8	60	Ib	ASC	DC	2	PR	Complete	NT	68	30	68	AWD
9	54	Ib	EDC	TC	1	PR	Complete	CT	68	68	68	NED
10	40	Ib2	MAC	TC	2	PR	Complete	CT	67	67	67	NED
11	38	Ib	MAC	DC	2	PR	Complete	CT	9	6	9	DOD
12	63	Ib	CCC	DC	2	PR	Complete	RT	48	48	48	NED
13	50	Ib	EDC	DC	2	PR	Complete	CT	35	35	35	NED
14	53	Ib2	EDC	DC	2	PR	Complete	CT	38	38	38	NED
15	54	Ib	MAC	DC	2	PR	Complete	CT	29	29	29	NED
16	52	Ib	MAC	TC	3	PR	Incomplete	CT	27	7	27	DOD
17	45	Ib	EDC	DC	2	PR	Complete	CT	10	10	10	NED
18	51	Ib	EDC	DC	3	PR	Complete	CT	13	13	13	NED
19	45	Ib	MAC	DC	2	SD	Incomplete	CT	10	3	10	DOD
20	52	Ib	ASC	DC	2	SD	Complete	CT	13	6	13	DOD
21	56	Ib	ASC	DC	2	SD	Complete	CCRT	56	56	56	NED
22	61	Ib	ASC	DC	3	SD	Incomplete	RT	35	8	35	DOD
23	45	Ib	MAC	DC	2	SD	Complete	CT	26	26	26	NED

ASC adenosquamous cell carcinoma, MAC mucinous adenocarcinoma, EDC endometrioid adenocarcinoma, CCC clear cell adenocarcinoma, DC docetaxel + carboplatin, TC paclitaxel + carboplatin, CR complete response, PR partial response, SD stable disease, NT no treatment, CT chemotherapy, RT radiotherapy, CCRT concurrent chemoradiation therapy, PFS progression-free survival, OS overall survival, NED no evidence of disease, AWD alive with disease, DOD died of disease

In the analysis of adverse events, severe neutropenia developed in 91.3 % of patients, but subsided in response to short-term treatment with a G-CSF preparation. During the first course of DC therapy, grade 3 febrile neutropenia developed in 2 cases; the dose of both agents was reduced for the next course of treatment. All signs, specific to taxanes, of peripheral neuropathy were grade 1 or less, allowing for continuation of treatment while preserving the quality of life of the individual patients. No serious adverse events occurred, and the response rate was 78.3 %. This study demonstrated a high response rate of bulky non-squamous cell carcinoma of the cervix to NAC using taxanes (paclitaxel or docetaxel) and carboplatin. It also demonstrated the safety of the medications in this regimen. The completion rate of radical hysterectomy, however, was only 78.3 %; thus, the treatment outcomes in this study were not satisfactory. Possible reasons for the low surgery completion rate include the rapid progression of non-squamous cell carcinoma, frequent invasion of tissues and organs surrounding the uterus, and frequent lymph node metastasis.

The treatment results and outcomes of all patients were shown in Table 4. Unfortunately, all patients with incomplete surgery ultimately experienced disease recurrence and died of their primary disease. Thus, the significance of NAC at present may not be to prolong survival time. Instead, in our view, NAC should be performed to fully optimize patients' conditions with its antitumor effect in order to improve the chances of complete surgery. Further study is needed regarding the long-term outcomes of NAC.

Conflict of interest The authors have no conflict of interest to declare.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- Whitney CW, Sause W, Bundy BN, Malfetano JH, Hannigan EV, Fowler WC Jr, Clarke-Pearson DL, Liao SY (1999) Randomized

- comparison of fluorouracil plus cisplatin versus hydroxyurea as an adjunct to radiation therapy in stage IIB-IVA carcinoma of the cervix with negative para-aortic lymph nodes: a Gynecologic Oncology Group and Southwest Oncology Group study. *J Clin Oncol* 17:1339–1348
2. Morris M, Eifel PJ, Lu J, Grigsby PW, Levenback C, Stevens RE, Rotman M, Gershenson DM, Mutch DG (1999) Pelvic radiation with concurrent chemotherapy compared with pelvic and para-aortic radiation for high-risk cervical cancer. *N Engl J Med* 340:1137–1143
 3. Rose PG, Bundy BN, Watkins EB, Thigpen JT, Deppe G, Maiman MA, Clarke-Pearson DL, Insalaco S (1999) Concurrent cisplatin-based radiotherapy and chemotherapy for locally advanced cervical cancer. *N Engl J Med* 340:1144–1153
 4. Keys HM, Bundy BN, Stehman FB, Muderspach LI, Chafe WE, Suggs CL III, Walker JL, Gersell D (1999) Cisplatin, radiation, and adjuvant hysterectomy compared with radiation and adjuvant hysterectomy for bulky stage 1B cervical carcinoma. *N Engl J Med* 340:1154–1161
 5. Peters WA III, Liu PY, Barrett RJ II, Stock RJ, Monk BJ, Berek JS, Souhami L, Grigsby P, Gordon W Jr, Alberts DS (2000) Concurrent chemotherapy and pelvic radiation therapy compared with pelvic radiation therapy alone as adjuvant therapy after radical surgery in high-risk early-stage cancer of the cervix. *J Clin Oncol* 18:1606–1613
 6. Thomas GM (1999) Improved treatment for cervical cancer—concurrent chemotherapy and radiotherapy. *N Engl J Med* 340:1198–1200
 7. Green JA, Kirwan JM, Tierney JF, Symonds P, Fresco L, Collingwood M, Williams CJ (2001) Survival and recurrence after concomitant chemotherapy and radiotherapy for cancer of the uterine cervix: a systematic review and meta-analysis. *Lancet* 358:781–786
 8. Sugiyama T, Nishida T, Kumagai S, Nishino S, Fujivoshi K, Okura N, Yakushiji M, Hiura M, Umetsuki N (1999) Combination chemotherapy with irinotecan and cisplatin as neoadjuvant in locally advanced cervical cancer. *Br J Cancer* 81:95–98
 9. Shoji T, Takatori E, Hatayama S, Omi H, Kagabu M, Honda T, Kumagai S, Morohara Y, Miura F, Yoshizaki A, Sugiyama T (2010) Phase II study of tri-weekly cisplatin and irinotecan as neoadjuvant chemotherapy for locally advanced cervical cancer. *Oncol Lett* 1:515–519
 10. Benedetti-Panici P, Greggi S, Colombo A, Amoroso M, Smaniotto D, Giannarelli D, Amunni G, Raspagliesi F, Zola P, Mangioni C, Landoni F (2002) Neoadjuvant chemotherapy and radical surgery versus exclusive radiotherapy in locally advanced squamous cell cervical cancer: results from the Italian multicenter randomized study. *J Clin Oncol* 20:179–188
 11. Aoki Y, Sato T, Watanabe M, Sasaki M, Tsuneki I, Tanaka K (2001) Neoadjuvant chemotherapy using low-dose consecutive intraarterial infusion of cisplatin combined with 5FU for locally advanced cervical adenocarcinoma. *Gynecol Oncol* 83:496–499
 12. Landoni F, Maneo A, Colombo A, Placa F, Milani R, Perego P, Favini G, Ferri L, Mangioni C (1997) Randomised study of radical surgery versus radiotherapy for stage Ib-IIa cervical cancer. *Lancet* 350:535–540
 13. Thigpen JT, Blessing JA, Fowler WC Jr, Hatch K (1986) Phase II trials of cisplatin and piperazinedione as single agents in the treatment of advanced or recurrent non-squamous cell carcinoma of the cervix: a Gynecologic Oncology Group Study. *Cancer Treat Rep* 70:1097–1100
 14. Sutton GP, Blessing JA, DiSaia PJ, McGuire WP (1993) Phase II study of ifosfamide and mesna in nonsquamous carcinoma of the cervix: a Gynecologic Oncology Group study. *Gynecol Oncol* 49:48–50
 15. Look KY, Blessing JA, Valea FA, McGehee R, Manetta A, Webster KD, Andersen WA (1997) Phase II trial of 5-fluorouracil and high-dose leucovorin in recurrent adenocarcinoma of the cervix: a Gynecologic Oncology Group study. *Gynecol Oncol* 67:255–258
 16. Rose PG, Blessing JA, Buller RE, Mannel RS, Webster KD (2003) Prolonged oral etoposide in recurrent or advanced non-squamous cell carcinoma of the cervix: a Gynecologic Oncology Group study. *Gynecol Oncol* 89:267–270
 17. Curtin JP, Blessing JA, Webster KD, Rose PG, Mayer AR, Fowler WC Jr, Malfetano JH, Alvarez RD (2001) Paclitaxel, an active agent in nonsquamous carcinomas of the uterine cervix: a Gynecologic Oncology Group Study. *J Clin Oncol* 19:1275–1278
 18. Nagao S, Fujiwara K, Oda T, Ishikawa H, Koike H, Tanaka H, Kohno I (2005) Combination chemotherapy of docetaxel and carboplatin in advanced or recurrent cervix cancer. A pilot study. *Gynecol Oncol* 96:805–809

Efficacy of neoadjuvant chemotherapy followed by radical hysterectomy in locally advanced non-squamous carcinoma of the uterine cervix: a retrospective multicenter study of Tohoku Gynecologic Cancer Unit

T. Shoji¹, S. Kumagai¹, A. Yoshizaki¹, Y. Yokoyama², T. Fujimoto³,
T. Takano⁴, N. Yaegashi⁴, K. Nakahara⁵, H. Nishiyama⁶, T. Sugiyama¹

¹Department of Obstetrics and Gynecology, Iwate Medical University School of Medicine, Morioka

²Department of Obstetrics and Gynecology, Hirosaki University Graduate School of Medicine, Hirosaki

³Department of Obstetrics and Gynecology, Akita University School of Medicine, Akita

⁴Department of Obstetrics and Gynecology, Tohoku University School of Medicine, Aoba-ku, Sendai

⁵Department of Obstetrics and Gynecology, Yamagata University School of Medicine, Yamagata

⁶Department of Obstetrics and Gynecology, School of Medicine Fukushima Medical University, Fukushima (Japan)

Summary

Objective: Radical hysterectomy (RH) is a standard treatment for locally advanced non-squamous cell carcinoma (N-SCC) of the uterine cervix, but there have been no reports on whether neoadjuvant chemotherapy (NAC) followed by radical hysterectomy could improve the outcome of patients with this disease. **Materials and Methods:** This multicenter retrospective study enrolled 77 patients with Stage IB2 to IIB N-SCC of the uterine cervix. Of these, 27 patients were treated with NAC prior to radical hysterectomy (NAC group) and 50 with RH alone (RH group). The two-year recurrence-free survival (RFS) rate, progression-free survival (PFS), and overall survival (OS) were compared between the two groups. Clinical parameters such as clinical stage, histological type, and postoperative treatment were also examined between the groups. **Results:** While the two-year RFS rates were 81.5% and 70.0% in NAC and RH groups, respectively ($p = 0.27$) and the median PFS was 51 months and 35 months in NAC and RH groups, respectively ($p = 0.35$), the median OS was 58 months and 48 months in NAC and RH groups, respectively, which was significant ($p = 0.0014$). The median OS of patients with mucinous adenocarcinoma in NAC group was significantly higher than that in RH group: 58 months versus 37 months ($p = 0.03$). **Conclusion:** NAC prior to RH may offer the prognostic advantage of patients with locally advanced N-SCC of the uterine cervix, especially mucinous adenocarcinoma.

Key words: Uterine cervical carcinoma; Non-squamous cell carcinoma; Neoadjuvant chemotherapy; Radical hysterectomy; Outcome.

Introduction

Radical hysterectomy and radiotherapy are a traditional therapeutic modality for invasive carcinoma of the uterine cervix in Japan. Since some observations showed that chemo-radiotherapy with cisplatin offered the advantage of clinical outcome in locally advanced carcinoma of the uterine cervix, chemotherapy has become the treatment of preference of uterine cervical carcinoma [1-7]. The Italian multicenter randomized study, which enrolled patients with locally advanced Stage IB2 to IIB squamous cell carcinoma of the uterine cervix, showed that NAC prior to RH improved the patient outcome as compared to conventional radiation therapy alone [8]. Combination of docetaxel and carboplatin in the neoadjuvant setting for patients with advanced or recurrent uterine cervical malignancy showed complete or partial response in all of patients with uterine cervical adenocarcinoma, suggesting that the combination may be quite promising for treatment of uterine cervical adenocarcinoma [9]. However, there is no evidence that NAC improves the outcome of

patients with uterine cervical adenocarcinoma. The aim of this multicenter study was to retrospectively evaluate whether NAC can improve the outcome of patients with locally advanced N-SCC of the uterine cervix.

Materials and Methods

This study enrolled 77 patients with Stage IB2 to IIB N-SCC of the uterine cervix who underwent RH at the institutions belonging to the Tohoku Gynecologic Cancer Unit (TGCU) between January 1996 and December 2008. Of these, 27 patients were treated with NAC prior to RH (NAC group), and 50 patients were treated with RH alone (RH group). The two-year recurrence-free survival (RFS) rate, progression-free survival (PFS), and overall survival (OS) were compared between the two groups. Clinical parameters, such as: clinical stage, histological type, and postoperative treatment were also examined between the groups.

The PFS and OS in the two groups were calculated by the Kaplan-Meier method, and the statistical significance of differences in the cumulative curves between the two groups was evaluated by log-rank test. Categorical variables comparisons were conducted by two-tailed Chi square or Mann-Whitney U test where appropriate. A result was deemed significant at $p < 0.05$.

Revised manuscript accepted for publication July 8, 2011

Results

Patient characteristics

The median age was 49 and 45 years in NAC and RH groups, respectively. Eleven (40.7%) and 29 (58.0%) patients had Stage IB2 disease in NAC and RH groups, respectively, and 16 (59.3%) and 21 (42.0%) patients had Stage II disease in NAC and RH groups, respectively. In regard to the histological type, 13 patients had mucinous adenocarcinoma, four had endometrioid adenocarcinoma, three had clear cell carcinoma, and seven had adenosquamous carcinoma in the NAC group, while 27 patients had mucinous adenocarcinoma, nine had endometrioid adenocarcinoma, two had clear cell carcinoma, nine had adenosquamous carcinoma, and three had other types in RH group. Of the 27 patients in NAC group and 50 in RH group, 19 (70.4%) and 40 (80.0%) underwent any postoperative treatments, respectively (Table 1).

NAC regimens and number of cycles

Because this was a retrospective and multicenter study, the combination of anti-cancer agents utilized was heterogeneous as shown in Table 2. Of the 27 patients in NAC group, eight received DC: seven patients received two cycles and one patient received three cycles. Five patients received cisplatin alone. Four patients received MEP: one patient received one cycle, two patients received two cycles, and one patient received three cycles. Three patients received TC of two cycles. Three patients received FCAP: one patient received one cycle and two patients received three cycles. Other four patients received cisplatin/CPT-11 of two cycles, cisplatin/Adriamycin of two cycles, cisplatin/mitomycin C of three cycles, and carboplatin/actinomycin D of three cycles, respectively.

Comparison of clinical outcome between NAC and RH groups

The two-year RFS rate was 81.5% in NAC group and 70.0% in RH group ($p = 0.27$, Table 3). The median PFS was 51 months (range, 14-157 months) in NAC group and 35 months (range, 4-157 months) in RH group ($p = 0.35$, Table 3). On the other hand, the median OS was 58 months (range, 15-157 months) in NAC group and 48 months (range, 9-157 months) in RH group, which was significant ($p = 0.0014$, Table 3 and Figure 1A).

Comparison of clinical outcome according to clinical parameters

There were no significant differences in the median PFS and OS between NAC and RH groups according to stage, histological type and adjuvant therapy, except mucinous adenocarcinoma (Table 4). While the median PFS of patients with mucinous adenocarcinoma was 58 months (range, 8-124 months) in NAC group and 33 months (range, 4-125 months) in RH group ($p = 0.34$), the

Table 1. — Patient characteristics.

Variable	NAC (n = 27)	RH (n = 50)	p value
Median age in years [range]	49 [30-63]	45 [25-76]	$p = 0.85^*$
Stage			
IB2	11 (40.7)	29 (58.0)	$p = 0.15^{**}$
II			
IIA	0	6 (12.0)	
IIB	16 (59.3)	15 (30.0)	
Histological type			
Adenocarcinoma			
mucinous	13 (48.1)	27 (54.0)	$p = 0.98^{**}$
endometrioid	4 (14.9)	9 (18.0)	
clear cell	3 (11.1)	2 (4.0)	
Adenosquamous carcinoma	7 (25.9)	9 (18.0)	
Others	0	3 (6.0)	
Adjuvant therapy administered	8 (29.7)	10 (20.0)	$p = 0.34^{**}$
not administered			
Chemotherapy	9 (33.3)	16 (32.0)	
Chemoradiation therapy	5 (18.5)	14 (28.0)	
Radiotherapy	5 (18.5)	10 (20.0)	

*Mann-Whitney U test, **Chi-square test, numbers of parenthesis represent %.

Table 2. — List of NAC regimens.

Regimen	No. of patients
DC (Docetaxel 70 mg/m ² , carboplatin AUC6 day 1 q21 days)	8
Cisplatin alone (total 200 mg/body for 3 days)	5
MEP (MMC 10 mg/m ² day 1, etoposide 100 mg/m ² days 1,3,5, cisplatin 50 mg/m ² day 1, q28 days)	4
TC (Paclitaxel 175 mg/m ² , carboplatin AUC6 day 1 q21 days)	3
FCAP (5-FU 200 mg/body, CPM100 mg/body, cisplatin 20 mg/m ² days 1-7, ADM 35 mg/m ² day 7)	3
Cisplatin/CPT-11 (cisplatin 70 mg/m ² day 1, CPT-11 70 mg/m ² days 1,8 q21 days)	1
Cisplatin/ADM (cisplatin 100 mg/body, ADM 40 mg/body days 1,2 q21 days)	1
Cisplatin/MMC (cisplatin 50 mg/body, MMC 4 mg/body day 1 q21 days)	1
Carboplatin/Actinomycin D (Carboplatin 300 mg/body, Actinomycin D 1.5 mg/body day 1 q21 days)	1

MMC: mitomycin C; CPM: cyclophosphamide; ADM: adriamycin.

Table 3. — Comparison of the clinical outcome between the two groups.

	NAC (n = 27)	RH (n = 50)	p value
Two-year RFS rate	81.5% (22/27)	70.0% (35/50)	$p = 0.27$
Median PFS (range)	51 months (14-157)	35 months (4-157)	$p = 0.35$
Median OS (range)	58 months (15-157)	48 months (9-157)	$p = 0.0014$

RFS: recurrence free survival; PFS: progression-free survival; OS: overall survival.

median OS of those with mucinous adenocarcinoma was 58 months (range, 24-124 months) in NAC group and 37 months (range, 9-125 months) in RH group, which was significant ($p = 0.03$) (Table 4 and Figure 1B).

Clinical outcome according to therapeutic modality after NAC and radical surgery

The outcome of patients who underwent chemotherapy or chemoradiotherapy or radiotherapy after NAC and RH were compared. As shown in Table 5, chemotherapy after NAC and surgery prolonged PFS and OS, and increased

Table 4. — Comparison of the clinical outcome according to clinical parameters.

Clinical parameters	Median PFS			Median OS		
	NAC	RH	<i>p</i> value	NAC	RH	<i>p</i> value
Stage						
IB2	64 (11-157)	37 (9-157)	0.26	64 (15-157)	54 (12-157)	0.26
II	33 (4-124)	45 (4-92)	0.45	39 (16-124)	45 (9-92)	0.40
Histological type						
Adenocarcinoma						
mucinous	58 (8-124)	33 (4-125)	0.34	58 (24-124)	37 (9-125)	0.03
endometrioid	31 (10-97)	70 (14-97)	0.29	31 (10-97)	70 (20-97)	0.49
clear cell	22 (4-108)	64 (12-106)	0.89	22 (16-108)	83 (60-106)	0.41
Adenosquamous	36 (12-157)	45 (12-157)	0.11	43 (21-157)	46 (12-157)	0.31
Adjuvant therapy						
administered	77 (25-157)	31 (5-157)	0.18	77 (35-157)	32 (12-157)	0.24
not administered	33 (4-124)	47 (4-106)	0.66	39 (16-124)	51 (9-106)	0.61

Numbers show months: Parenthesis means range. PFS: progression-free survival; OS: overall survival.

Table 5. — Clinical outcome according to therapeutic modality after NAC and radical surgery.

	Chemotherapy (n = 9)	Chemoradiotherapy (n = 5)	Radiotherapy (n = 5)
PFS (months)	42 (10-108)	30 (4-76)	22 (8-97)
OS (months)	42 (19-108)	30 (16-76)	31 (22-97)
Two-year RFS rate	88.9%	60.0%	60.0%

PFS: progression-free survival; OS: overall survival; RFS: recurrence free survival.

the two-year RFS rate compared to chemoradiotherapy or radiotherapy after NAC and surgery, although they did not reach significance.

Discussion

Numerous phase II studies have reported the favorable effects of NAC in the treatment of locally advanced carcinoma of the uterine cervix. The authors have previously reported the efficacy and safety of NAC with cisplatin plus irinotecan in this disease [10]. However, few randomized clinical trials (RCT) have evaluated the effect of NAC in the clinical outcome of patients with this disease. Sardi *et al.* reported a significant improvement of the seven-year survival rate in patients treated by NAC and radical surgery and radiotherapy (65%), as compared with that in those treated by radical surgery and radiotherapy (41%) in a four-arm randomized controlled trial (RCT) (NAC and radical surgery and radiotherapy, radical surgery and radiotherapy, radiotherapy alone, and NAC and radiotherapy) [11]. However, the retrospective study did not show obvious improvement of the five-year survival rate in patients with Stage IB2 carcinoma of the uterine cervix treated by NAC prior to surgery, as compared with that in those treated by surgery alone (80% versus 69%) [12]. These two reports were conducted in patients with SCC of the uterine cervix. Since N-SCC of the uterine cervix has recently increased in Japan, it is an important issue to evaluate the effectiveness of NAC in the outcome of patients with N-SCC of the uterine cervix. Some evidence showed that the outcome of patients with N-SCC of the uterine cervix was poorer than that of

patients with SCC of the uterine cervix [13, 14], because of the higher incidence of lymph node metastases at a relatively early stage of the disease, and a lower sensitivity to radiotherapy in N-SCC of the uterine cervix [15, 16]. Chemotherapy is therefore expected to have a greater beneficial effect on the outcome of patients with N-SCC than radiotherapy or chemoradiation therapy. Because the present study was conducted retrospectively in multicenters, the combination of anti-cancer agents used was heterogeneous. The NAC regimens used in this study invariably included one of platinum derivatives, such as cisplatin and carboplatin, so platinum agents seem favorable for chemotherapy prior to surgery in N-SCC of the uterine cervix.

The two-year RFS rate and the median PFS were better in NAC group than in RH group, which were not significant, whereas the median OS in NAC group was significantly longer than in RH group (*p* = 0.0014). Furthermore, prognostic analysis in clinical parameters showed that the median OS of patients with mucinous adenocarcinoma in NAC group was significantly longer than in RH group (*p* = 0.03), although other histological types and postoperative treatment did not significantly affect the prognosis of patients between NAC and RH groups. These results suggest that NAC may offer the prognostic advantage of patients with locally advanced N-SCC of the uterine cervix, especially mucinous adenocarcinoma. Because mucinous adenocarcinoma accounts for approximately 70% out of adenocarcinomas of the uterine cervix, NAC may improve prognosis of patients with N-SCC of the uterine cervix, although NAC should be used individually at the present time.

The present study showed that chemotherapy after NAC and surgery prolonged PFS and OS, compared to chemoradiotherapy or radiotherapy after NAC and surgery, which did not reach significance. Tattersall *et al.* reported that primary chemotherapy followed by radiotherapy significantly decreased the survival rate of patients with uterine cervical carcinoma compared to those who were treated by radiotherapy alone [17]; furthermore, meta-analysis showed that chemotherapy followed by radiotherapy did not improve the survival time

Fig. 1A




Fig. 1B




Figure 1. — 1A: overall survival in all patients who underwent neoadjuvant chemotherapy followed by radical hysterectomy (NAC) or radical hysterectomy alone (RH). 1B: overall survival in patients with mucinous adenocarcinoma who underwent neoadjuvant chemotherapy followed by radical hysterectomy (NAC) or radical hysterectomy alone (RH).

in uterine cervical carcinoma [18]. Considering these reports together with the present results, chemoradiotherapy or radiotherapy after NAC and surgery may contribute to unfavorable outcome of the patients with uterine cervical adenocarcinoma compared to chemotherapy after NAC and surgery, although further investigation is necessary to confirm the appropriate therapeutic modality following NAC and surgery.

The recent reports demonstrated that taxanes were used effectively in NAC for uterine cervical adenocarcinoma [19, 20]. Most of the institutions joining TGCU had adopted cisplatin-based regimens in the 1990s, and switched to the regimens combining taxanes and platinum derivatives after 2000. Despite the diverse NAC regimens and the small sample size, the authors believe that the present results have provided constructive ideas for the development of new therapeutic strategy for N-SCC of the uterine cervix. An effective chemotherapeutic regimen for N-SCC of the uterine cervix should be urgently integrated in a phase II study and then a RCT that compares a new single NAC and radical surgery with radical surgery alone, is warranted to confirm the present results.

Acknowledgements

The authors thank the medical staff in the institutions joining Tohoku Gynecologic Cancer Unit for clinical review of this study.

References

- [1] Whitney C.W., Sause W., Bundy B.N., Malfetano J.H., Hannigan E.V., Fowler W.C. Jr. *et al.*: "Randomized comparison of fluorouracil plus cisplatin versus hydroxyurea as an adjunct to radiation therapy in Stage IIB-IVA carcinoma of the cervix with negative para-aortic lymph nodes: a Gynecologic Oncology Group and Southwest Oncology Group study". *J. Clin. Oncol.*, 1999, 17, 1339.
- [2] Morris M., Eifel P.J., Lu J., Grigsby P.W., Levenback C., Stevens R.E. *et al.*: "Pelvic radiation with concurrent chemotherapy compared with pelvic and para-aortic radiation for high-risk cervical cancer". *N. Engl. J. Med.*, 1999, 340, 1137.
- [3] Rose P.G., Bundy B.N., Watkins E.B., Thigpen J.T., Deppe G., Maiman M.A. *et al.*: "Concurrent cisplatin-based radiotherapy and chemotherapy for locally advanced cervical cancer". *N. Engl. J. Med.*, 1999, 340, 1144.
- [4] Keys H.M., Bundy B.N., Stehman F.B., Mudderspach L.I., Chafe W.E., Suggs C.L. 3rd *et al.*: "Cisplatin, radiation, and adjuvant hysterectomy compared with radiation and adjuvant hysterectomy for bulky Stage 1B cervical carcinoma". *N. Engl. J. Med.*, 1999, 340, 1154.
- [5] Peters W.A. 3rd, Liu P.Y., Barrett R.J. 2nd, Stock R.J., Monk B.J., Berek J.S. *et al.*: "Concurrent chemotherapy and pelvic radiation therapy compared with pelvic radiation therapy alone as adjuvant therapy after radical surgery in high-risk early-stage cancer of the cervix". *J. Clin. Oncol.*, 2000, 18, 1606.
- [6] Thomas G.M.: "Improved treatment for cervical cancer—concurrent chemotherapy and radiotherapy". *N. Engl. J. Med.*, 1999, 340, 1198.
- [7] Green J.A., Kirwan J.M., Tierney J.F., Symonds P., Fresco L., Collingwood M. *et al.*: "Survival and recurrence after concomitant chemotherapy and radiotherapy for cancer of the uterine cervix: a systematic review and meta-analysis". *Lancet*, 2001, 358, 781.
- [8] Benedetti-Panici P., Greggi S., Colombo A., Amoroso M., Smaniotto D., Giannarelli D. *et al.*: "Neoadjuvant chemotherapy and radical surgery versus exclusive radiotherapy in locally advanced squamous cell cervical cancer: results from the Italian multicenter randomized study". *J. Clin. Oncol.*, 2002, 20, 179.
- [9] Nagao S., Fujiwara K., Oda T., Ishikawa H., Koike H., Tanaka H. *et al.*: "Combination chemotherapy of docetaxel and carboplatin in advanced or recurrent cervix cancer. A pilot study". *Gynecol. Oncol.*, 2005, 96, 805.
- [10] Shoji T., Takatori E., Hatayama S., Omi H., Kagabu M., Honda T. *et al.*: "Phase II Study of Tri-weekly cisplatin and irinotecan as neoadjuvant chemotherapy for locally advanced cervical cancer". *Oncol. Lett.*, 2010, 1, 515.
- [11] Sardi J.E., Giaroli A., Sananes C., Ferreira M., Soderini A., Bermudez A. *et al.*: "Long-term follow-up of the first randomized trial using neoadjuvant chemotherapy in Stage 1b squamous carcinoma of the cervix: The final results". *Gynecol. Oncol.*, 1997, 67, 61.
- [12] Serur E., Mathews R.P., Gates J., Levine P., Maiman M., Remy J.C.: "Neoadjuvant chemotherapy in stage 1B2 squamous cell carcinoma of the cervix". *Gynecol. Oncol.*, 1997, 65, 348.

- [13] Shingleton H.M., Gore H., Bradley D.H., Soong H.J.: "Adenocarcinoma of the cervix : I. Clinical evaluation and pathologic features". *Am. J. Obstet. Gynecol.*, 1981, 139, 799.
- [14] Fu Y.S., Reagan J.W., Hsui J.G., Storaasli J.P., Wentz W.B.: "Adenocarcinoma and mixed carcinoma of the cervix: A clinico-pathologic study". *Cancer*, 1982, 49, 2560.
- [15] Berek J.S., Hacker N.F., Fu Y., Sokale J.R., Leucher R.C., Lagasse L.D.: "Adenocarcinoma of the uterine cervix: histologic variables associated with lymph node metastasis and survival". *Obstet. Gynecol.*, 1985, 65, 46.
- [16] Moberg P.J., Einhorn N., Silfverswärd C., Söderberg G.: "Adenocarcinoma of the cervix". *Cancer*, 1986, 57, 407.
- [17] Tattersall M.H., Lorvidhaya V., Vootiprux V., Cheirsilpa A., Wong F., Azhar T. *et al.*: "Randomized trial of epirubicin and cisplatin chemotherapy followed by pelvic radiation in locally advanced cervical cancer. Cervical Cancer Study Group of the Asian Oceanian Clinical Oncology Association". *J. Clin. Oncol.*, 1995, 13, 444.
- [18] Neoadjuvant Chemotherapy for Locally Advanced Cervical Cancer Meta-analysis Collaboration: "Neoadjuvant chemotherapy for locally advanced cervical cancer: a systematic review and meta-analysis of individual patient data from 21 randomized trials". *Eur. J. Cancer*, 2003, 39, 2470.
- [19] Oguri H., Maeda N., Fukaya T.: "Cervical adenocarcinoma treated with docetaxel and carboplatin". *Int. J. Gynaecol. Obstet.*, 2003, 83, 209.
- [20] Huang X., Lan C., Huang H., Zhang Y., Huang H., Cao X. *et al.*: "Neoadjuvant docetaxel combined with cisplatin and followed by radical surgery for the treatment of locally advanced (Stage IB2 - IIB) cervical cancer: preliminary results of a single-institution experience". *Expert Opin. Pharmacother.*, 2011, 12, 165.

Address reprint requests to:

T. SHOJI, M.D.

Department of Obstetrics and Gynecology
Iwate Medical University School of Medicine
19-1 Uchimaru
Morioka 020-8505 (Japan)
e-mail: tshoji@iwate-med.ac.jp



RESEARCH

Open Access

Are platinum agents, paclitaxel and irinotecan effective for clear cell carcinoma of the ovary? DNA damage detected with γ H2AX induced by anticancer agents

Eriko Takatori^{1†}, Tadahiro Shoji^{1†}, Seisuke Kumagai^{1†}, Takashi Sawai^{2†}, Akira Kurose^{3†} and Toru Sugiyama^{1†}

Abstract

Objectives: Differences in the incidences and types of DNA damage induced by antitumor agents for clear cell carcinoma (CCC) were determined in 2 ovarian CCC cell lines using γ H2AX.

Material and methods: The antitumor activity of anticancer agents, CDDP, CBDCA, PTX and SN-38, was examined using ovarian clear cell carcinoma cultured cell lines (OVISE and RMG-I). After culture, each cell line was treated with each anticancer agent, the cells were collected, fixed, and then reacted with the anti- γ H2AX antibody. γ H2AX and nuclear DNA were then simultaneously detected by flow cytometry using FITC and propidium iodide, respectively, to determine γ H2AX in each cell cycle phase.

Results: After administration of CDDP, DNA damage was frequent in S-phase cells, while cell-cycle arrest occurred in the G1 and G2/M phases and γ H2AX did not increase in CDDP-resistant cells. Sensitivities to CDDP and CBDCA differed between the two cell lines. The antitumor effect of PTX is induced by G2/M arrest, and combination treatment with CBDCA, inducing DNA damage in G2/M-phase cells, might be effective.

Conclusions: This is the first study in Japan to evaluate the antitumor activity of anticancer agents by focusing on the relationship between the cell cycle and DNA damage using γ H2AX as an indicator. The immunocytochemical method used in this study detects γ H2AX, which indicates DNA damage even at very low concentrations and with high sensitivity. Therefore, a promising method of easily and rapidly identifying agents potentially effective against CCC.

Keywords: γ H2AX, Clear cell carcinoma, Ovarian cancer, DNA damage, Apoptosis, Chemotherapy

Introduction

Clear cell adenocarcinoma (CCC), a subtype of epithelial ovarian cancer, is less sensitive to chemotherapy and is thus classified as a refractory ovarian cancer [1]. It has been shown that a combination of carboplatin (CBDCA) and paclitaxel (PTX), a standard therapy for ovarian cancer [2,3], is effective against serous adenocarcinoma and endometrioid adenocarcinoma, with a response rate of approximately 75%, while CCC has lower response rates

ranging from 18% to 50% [4]. The incidence of CCC has been increasing and is now 25% in Japan, while that in Europe is 5-6%. As yet, no treatment for this histological subtype of ovarian cancer has been established. Histopathology remains the gold standard for classifying epithelial ovarian cancer subgroups; however, there is emerging evidence indicating different genetic and molecular profiles. Consequently, there is no international consensus regarding the necessity of establishing treatment strategies based on histological subtypes. In fact, global clinical trials of CCC and mucinous adenocarcinoma have already begun. Although which cytotoxic agents have true efficacy against CCC remains unknown, small trials in Japan and basic studies have suggested the efficacy of irinotecan

* Correspondence: tshoji@iwate-med.ac.jp

[†]Equal contributors

¹Department of Obstetrics and Gynecology, Iwate Medical University School of Medicine, 19-1, Uchimaru, Morioka, Iwate 020-8505, Japan

Full list of author information is available at the end of the article



(CPT-11) [5-7]. The Japanese Gynecologic Oncology Group (JGOG) started an international randomized controlled trial (RCT) of cisplatin (CDDP)/CPT-11 therapy with a control arm of CBDCA/TXL (TC) therapy (JGOG3017/GCIG); patient accrual is ongoing and approximately 560 patients had been enrolled in the trial as of July 2010. In addition, an ongoing translational study, as part of the JGOG3017/GCIG trial, also aims to establish an updated treatment strategy.

Nucleosomes, units of chromatin, consist of core histones wrapped in 146 bp of DNA and linker DNA. Core histones are octamers designated H2A, H2B, H3 and H4. Histone H2AX is a variant of histone H2A and accounts for 10-15% of all variants. When DNA damage occurs, serine 139 of histone H2AX in chromatin on both sides of a damaged site is phosphorylated by two enzymes: ataxia telangiectasia mutated (ATM) protein kinase and by ATM and Rad3 related (ATR) protein kinase [8,9]. Phosphorylated histone H2AX is called γ H2AX. Dot γ H2AX, which is detectable using γ H2AX-specific antibody, is considered to correspond to specific DNA damage. Therefore, DNA damage can be immunocytochemically detected [10]. DNA damage in individual cells has been detected employing a single-cell DNA gel electrophoresis technique (comet assay), in which the extent and length of the comet's tail indicate the severity of DNA damage. Recently, however, it has become apparent that phosphorylation of histone H2AX, one of the variants of the nucleosome core histone H2A, can provide a sensitive and reliable marker of DNA damage. More specifically, DNA damage, particularly that involving the formation of DNA double-strand breaks (DSBs), induces phosphorylation of histone H2AX on Ser-139; phosphorylated H2AX is defined as γ H2AX. The phosphorylation takes place on H2AX molecules on both sides of DSBs along a megabase length of DNA. Although DSBs generated during DNA fragmentation in the course of apoptosis also induce γ H2AX, the degree of γ H2AX induction in apoptotic cells is much greater than that in primary DSBs induced by antitumor drugs or radiation. The presence of γ H2AX in cells can be detected immunocytochemically in the form of distinct nuclear γ H2AX immunofluorescent foci and each focus is considered to correspond to a single DSB. This immunocytochemical approach has made it possible to assay DNA damage and *in situ* repair of the chromatin of individual cells. The immunocytochemical approach is significantly more sensitive than the comet assay. The use of multi-parameter flow cytometry in measurements of γ H2AX immunofluorescence allows DNA damage to be correlated with cellular DNA content and, therefore, the cell-cycle phase. Determination of the cell-cycle phase targeted by the drug is of importance in elucidating the mechanism of antitumor drug activity.

In the present study, we conducted flow cytometric bivariate analyses of γ H2AX and DNA contents in two different cell lines of CCC treated with CDDP, CBDCA, PTX or CPT-11 (SN-38), which have been used in the aforementioned international clinical randomized trial targeting CCC, and examined effects of these drugs with regard to the induction of DNA damage, apoptosis and cell-cycle progression vis-à-vis the cell-cycle phase.

Materials and methods

Cell culture

We used two CCC cell lines (OVISe and RMG-I) were obtained from the Health Science Research Resources Bank (Osaka, Japan). OVISe was established from a patient with metastatic disease after completion of six cycles of platinum combination therapy, and was grown in dishes (Becton Drive, Franklin Lakes, NJ, USA) in RPMI1640 medium (Sigma Chemical Co., St Louis, MO, USA) with 10% fetal bovine serum. RMG-I was established from a chemotherapy-naïve patient with ascites, and was reported to be primary platinum resistant (Table 1) [11]. RMG-I was grown in dishes (Becton Drive) in Ham F-12 medium supplemented with 10% fetal bovine serum. The media for the two cell lines were supplemented with 100 U/ml penicillin and 100 μ g/ml streptomycin (Meiji Seika Co., Ltd., Tokyo, Japan). All cell lines were maintained at 37°C in a humidified atmosphere of 5% CO₂ in air.

Drugs

CDDP, PTX and SN-38 (CPT-11) were dissolved in dimethyl sulfoxide (DMSO, Sigma); the final concentration of DMSO in the culture medium never exceeded 0.1% (w/v). CBDCA was dissolved in phosphate-buffered saline (PBS). The concentration of each agent was set to correspond to the blood concentration at a standard clinical dose (Table 2).

Immunohistochemistry

Both cells floating in the medium and the cells that remained attached after trypsinization were collected and fixed with 1% methanol-free formaldehyde (Polysciences Inc., Warrington, PA, USA) in PBS at 0 °C for 15 minutes and post-fixed with 80% ethanol for at least 2 hours at -20 °C. The fixed cells were washed twice in PBS and

Table 1 Clinical biological characteristics of the cell line

Cell Line	Source	Histopathology	Pretreatment	Median doubling time
OVISe	Solid metastatic	CCC	CAP \times 6 courses	60 hours
RMG-1	Ascites	CCC	No	60 hours

CCC; clear cell carcinoma

CAP; cyclophosphamide, doxorubicin, cisplatin

Table 2 Minimum effective concentration (MEC)

	Cmax	MEC
PTX	10 µg/ml	50 ng/ml
CDDP	7 µg/ml	1 µg/ml
CBDCA	55 µg/ml	10 µg/ml
SN-38	30 µg/ml	1 ng/ml

Cmax; clinical maximum drug concentration.

suspended in a 1% (w/v) solution of bovine serum albumin (BSA) (Sigma) in PBS to suppress non-specific antibody binding. The cells were then incubated in 100 µl of 1% BSA containing 1:100 diluted anti-phosphohistone H2AX (Ser-139) monoclonal antibody (Upstate, Lake Placid, NY, USA) for 2 hours at room temperature, washed twice with PBS and resuspended in 100 µl of 1:20 diluted fluorescein isothiocyanate (FITC)-conjugated F(ab')² fragment of goat anti-mouse immunoglobulin (Dako, Glostrup, Denmark) for 30 minutes at room temperature in the dark. The cells were then counterstained with 5 µg/ml propidium iodide (PI) (Sigma) in the presence of 100 µl of RNaseA (Sigma) for 30 minutes.

Fluorescence measurements by flow cytometry

The FITC (green) and PI (red) fluorescence of individual cells in suspension induced by excitation with a 488-nm argon ion laser was measured using a FACScan flow cytometer (Becton-Dickinson, San Jose, CA, USA). The green and red fluorescence from each cell was separated and quantified using standard optics and Cell Quest software (Becton-Dickinson). Ten thousand cells were measured per sample. All experiments were repeated at least three times.

After γH2AX and DNA staining, the DNA content was represented on the x axis and the γH2AX content on the y axis using flow cytometry. The γH2AX in each cell cycle was determined, thereby allowing the relationships between cell kinetics and DNA damage induced by antitumor agents to be examined.

Results

Platinum Agents

CDDP

OVICE cells showed an increase in the number of distinct green dots (γH2AX foci) after exposure to 10 µg/ml CDDP for 24 h, which indicates that CDDP caused DNA damage (Figure 1). Using flow cytometry, DNA damage was evident from the increase in γH2AX. After 24-hour treatment with CDDP, DNA damage in OVICE and RMG-I was seen gradually in the S phase at concentrations of 100 ng/ml and 1 ng/ml (Figure 2A). In both cell lines, treatment with 100 ng/ml or more CDDP for 24 hours caused DNA damage throughout the cell cycle. The cells with DNA damage gradually underwent apoptosis, as was evident by the presence of cells with markedly elevated γH2AX and fractional




Figure 1 Immunohistochemistry. Representative microscopic images of OVICE cell line after exposure to 10 µg/ml CDDP for 24 h. γH2AX foci in a nuclear are stained green and red, respectively. An increase in green dots, indicating elevation of γH2AX, can be seen after exposure to CDDP. Thus, DNA damage is visually recognizable in each nucleus. Apoptotic bodies (insert) are distinguishable by the entire cell being intensely positive for γH2AX.

(sub-G1) DNA contents. In OVICE, DNA damage in the S and G2/M phases after treatment with 100 ng/ml CDDP was seen for 24 and 72 hours, respectively (Figure 2B). Although in RMG-I, 100 ng/ml CDDP caused DNA damage in the S phase, in other phases of the cell cycle it was not apparent even with longer treatment. In both cell lines, the cells with damaged DNA underwent apoptosis. The number of cells in the G2/M phase in OVICE decreased gradually indicating S-phase arrest. On the other hand, in RMG-I showed G1 and G2/M-phases arrest. RMG-I was found to be less susceptible to DNA damage and subsequent apoptosis than OVICE.

CBDCA

DNA damage in the S phase was seen gradually after exposure to CBDCA for 24 hours in OVICE and RMG-I lines at 1 µg/ml and 10 µg/ml, respectively (Figure 3A). Subsequently, the cells with damaged DNA underwent apoptosis. Gradually both cell lines showed DNA damage in the G2/M phase and underwent apoptosis. OVICE showed S and G2/M-phases arrest, while RMG-I G2/M-phase arrest (Figure 3B).

PTX

Exposure to 10 ng/ml or more PTX for 24 hours caused apoptosis without primary DNA damage in both cell lines. (Figure 4A). Although, further apoptotic effects were not seen at doses exceeding 50 ng/ml. PTX induced both cell lines G2/M-phase arrest, but some cells remained 120 hours after exposure without primary DNA damage (Figure 4B).




Figure 2 Bivariate distributions (DNA content vs. γ H2AX) of CCC cell lines, OVISE and RMG-1, treated with CDDP (Upper, OVISE; Lower, RMG-1). The dotted lines indicate the upper level of γ H2AX immunofluorescence for 95% of cells in the untreated (control) culture. Arrow heads indicate elevation of γ H2AX that means DNA damage. Arrows indicate apoptotic cell populations with marked increase in γ H2AX and gradual decrease in DNA. (A) Both cell lines treated with various concentrations of CDDP for 24 h. OVISE and RMG-1 cell lines exhibited DNA damage in S-phase cells at minimum concentrations of 100 ng/ml and 1 μ g/ml, respectively. Both cell lines were subjected to DNA damage concentration-dependently at every cell cycle, and apoptosis was induced at concentrations of 5 μ g/ml or higher. More cells in RMG-1 remained free of DNA damage as compared to OVISE. (B) Both cell lines were treated with 100 ng/ml, the minimum concentration inducing DNA damage in either cell line, for various reaction times. S-phase cells of OVISE showing DNA damage progressed to apoptosis after 48 h. In addition, S-phase arrest was observed. DNA damage was induced in S-phase cells of RMG-1 after 48 h. Cells with DNA damage progressed to apoptosis after 72 h. Furthermore, cell-cycle arrest occurred in all cells.






Figure 3 Bivariate distributions (DNA content vs. γ H2AX) of OVISE and RMG-1 cell lines, treated with CBDCA (Upper, OVISE; Lower, RMG-1). The dotted lines indicate control. Arrow heads and arrows indicate DNA damage and apoptosis, respectively. (A) Both cell lines treated with various concentrations of CBDCA for 24 h. DNA damage was observed in the S-phase cells at 1 μ g/ml and 10 μ g/ml concentrations in OVISE and RMG-1, respectively. DNA damage was found in both cell lines at every cell cycle as the concentration increased, and apoptosis occurred at a concentration of 50 μ g/ml. More cells remained free of DNA damage in RMG-1 than in OVISE. (B) Both cell lines were treated with 1 μ g/ml, the minimum concentration inducing DNA damage in either cell line, for various reaction times. In OVISE, S-phase cells with DNA damage progressed to apoptosis after 48 h. DNA damage was also found in G2/M-phase cells after 48 h, but most did not progress to apoptosis. S and G2/M-phase arrests were observed. DNA damage was found in S and G2/M-phase cells after 48 h in RMG-1. The S-phase cells with DNA damage progressed to apoptosis 78 h later, but G2/M-phase cells showing DNA damage remained. S and G2/M-phase arrests were observed.



SN-38

After treatment with 0.5 ng/ml or more of SN-38 for 24 hours, both cell lines distinguished DNA damage in the S phase. Nevertheless even 10 ng/ml which is nearly the clinical maximum blood concentration did not cause

apoptosis (Figure 5A). Apoptosis was seen barely 72 hours after exposure to 0.5 ng/ml SN-38 in a portion of OVISE while in RMG-I after 96 hours (Figure 5B). However, the cells with significant DNA damage remained in the S phase over 120 hours after exposure.



Discussion

After CDDP administration, DNA damage was observed mainly in the S phase. It is reasonable to assume that the DNA was structurally altered by CDDP, leading to DNA replication fork arrest and ultimately resulting in

apoptosis. This result was consistent with a known pharmacological effect of CDDP [12]. In RMG-I, apoptotic cells were minimally increased in the S phase, moreover the cells showing arrest in the G1 and G2/M phases without DNA damage were increased as

compared with OVISe. Therefore, the results of the present study support the clinical experience that RMG-I is CDDP-resistant [11,13]. After CBDCA administration, DNA damage was seen in the S and G2/M phases in both cell lines. OVISe contained a remarkable cell population rescued from apoptosis and surviving with DNA damage. On the other hand, most RMG-I cells with DNA damage underwent apoptosis. These results suggest that cell lines respond differently to platinum agents, i.e., RMG-I was CDDP-resistant but responded to CBDCA. PTX directly induced apoptosis in M-phase cells but not via DNA damage, an observation consistent with a known pharmacological effect of PTX, i.e. microtubule inhibition [14]. PTX was confirmed to induce apoptosis through a p53-independent pathway; it was, therefore, expected to have an effect on CCC, in which the p53 mutation is rare [15,16]. The mechanism underlying the antitumor effect of PTX is G2/M arrest. Therefore, the combination with CBDCA, an agent inducing DNA damage, in G2/M-phase arrested cells might be effective, at least theoretically. As shown in this study, it is noteworthy that sensitivities to CDDP and CBDCA differed between the CCC cell lines. In practice, CCC is less sensitive to CBDCA/PTX treatment [4,6,17], which is the standard regimen for ovarian cancer. Since the effect of PTX was independent of both the concentration and the response time, these results raise the possibility that repeated administration of PTX at a low dose increases the antitumor effect more than a single administration. These findings support the results of the JGOG3016, i.e. that weekly CBDCA (AUC6)/PTX (80 mg/m², weekly × 3) is more effective than tri-weekly CBDCA (AUC 6)/PTX (175 mg/m²) treatment [18].

On the other hand, after administration of SN-38, DNA damage occurred in S-phase cells, followed by apoptosis. This confirmed that SN-38 acts as a type I topoisomerase inhibitor [19]. Furthermore, it appears that SN-38 had an effect on the cell cycle because S-phase arrest continued for more than 120 hours. It is, therefore, possible that the improved administration method for SN-38 increases its antitumor effect. Cells rescued from apoptosis remained in S phase with DNA damage; consequently, the efficacy of combining SN-38 with CDDP, which induces DNA damage mainly in S-phase cells, was supported.

In conclusion, the present results suggest that an effective treatment for CCC with a slow growth rate and a low ratio of S-phase cells would be a combination of agents arresting the cell cycle, thereby causing accumulation of cells in the S phase or the G2/M phase, and agents specifically inducing DNA damage in S-phase cells. The method used in this study allows immunocytochemical detection of γ H2AX, which indicates DNA damage even at very low concentrations and has high

sensitivity in comparison with the comet assay. Employing this method, we were able to analyze relationships between anti-tumor effects and cell cycle perturbations. Therefore, γ H2AX detection is a promising method of simply and rapidly identifying agents potentially effective against CCC.

Competing interest

The authors have no conflicts of interest to report.

Author details

¹Department of Obstetrics and Gynecology, Iwate Medical University School of Medicine, 19-1, Uchimaru, Morioka, Iwate 020-8505, Japan. ²Department of Pathology, Iwate Medical University School of Medicine, Uchimaru, Morioka, Iwate 020-8505, Japan. ³Department of Anatomic Pathology, Hirosaki University Graduate School of Medicine, 5, Zaifu-cho, Hirosaki, Aomori 036-8562, Japan.

Authors' contributions

ET participated in all aspects of the study, from design to clinical and laboratory performance, and manuscript writing. TS participated in design, data analysis and drafting of the manuscript. AK participated in the design of the study and technical assistance. SK, TS and TS participated in design and analysis of clinical data. All authors have read and approved the manuscript.

Received: 20 April 2012 Accepted: 12 June 2012

Published: 12 June 2012

References

1. Sugiyama T, Kamura T, Kigawa J, Terakawa N, Kikuchi Y, Kita T, Suzuki M, Sato I, Taguchi K: Clinical characteristics of clear cell carcinoma of the ovary. *Cancer* 2000, **88**:2584–2589.
2. McGuire WP, Hoskins WJ, Brady MF, Kucera PR, Partridge EE, Look KY, Clarke-Pearson DL, Davidson M: Cyclophosphamide and cisplatin compared with paclitaxel and cisplatin in patients with stage III and stage IV ovarian cancer. *N Engl J Med* 1996, **334**:1–6.
3. Ozols RF, Bundy BN, Greer BE, Fowler JM, Clarke-Pearson D, Burger RA, Mannel RS, DeGeest K, Hartenbach EM, Baergen R, Gynecologic Oncology Group: Phase III trial of carboplatin and paclitaxel compared with cisplatin and paclitaxel in patients with optimally resected stage III ovarian cancer: a Gynecologic Oncology Group study. *J Clin Oncol* 2003, **21**:3194–3200.
4. Sugiyama T, Fujiwara K: Clear cell carcinoma of the ovary. In *American Society of Clinical Oncology 2007 educational book*. Edited by Govindan R. Alexandria, VA; 2007:318–322.
5. Takano M, Kikuchi Y, Yaegashi N, Suzuki M, Tsuda H, Sagae S, Udagawa Y, Kuzuya K, Kigawa J, Takeuchi S, Tsuda H, Moriya T, Sugiyama T: Adjuvant chemotherapy with irinotecan hydrochloride and cisplatin for clear cell carcinoma of the ovary. *Oncol Rep* 2006, **16**:1301–1306.
6. Takano M, Sugiyama T, Yaegashi N, Suzuki M, Tsuda H, Sagae S, Udagawa Y, Kuzuya K, Kigawa J, Takeuchi S, Tsuda H, Moriya T, Kikuchi Y: Progression-free survival and overall survival of patients with clear cell carcinoma of the ovary treated with paclitaxel-carboplatin or irinotecan-cisplatin: retrospective analysis. *Int J Clin Oncol* 2007, **12**:256–260.
7. Nishino K, Aoki Y, Amikura T, Obata H, Sekine M, Yahata T, Fujita K, Tanaka K: Irinotecan hydrochloride (CPT-11) and mitomycin C as the first line chemotherapy for ovarian clear cell adenocarcinoma. *Gynecol Oncol* 2005, **97**:893–897.
8. Dickey JS, Redon CE, Nakamura AJ, Baird BJ, Sedelnikova OA, Bonner WM: H2AX: functional roles and potential applications. *Chromosoma* 2009, **118**:683–692.
9. Fragkos M, Jurvansuu J, Beard P: H2AX is required cell cycle arrest via the p53/p21 pathway. *Mol Cell Biol* 2009, **29**:2828–2840.
10. Bonner WM, Redon CE, Dickey JS, Nakamura AJ, Sedelnikova OA, Solier S, Pommier Y: γ H2AX and cancer. *Nat Rev Cancer* 2008, **8**:957–967.
11. Itamochi H, Kigawa J, Sultana H, Iba T, Akeshima R, Kamazawa S, Kanamori Y, Terakawa N: Sensitivity to anticancer agents and resistance mechanisms in clear cell carcinoma of the ovary. *Jpn J Cancer Res* 2002, **93**:723–728.

12. Zwelling LA, Kohn KW: Mechanism of action of cis-Dichlorodiammineplatinum (II). *Cancer Treat Rep* 1979, **63**:1439-1444.
13. Okuma Y, Kiguchi K, Koshitaka Y, Okamura A, Ishiwata I, Kondo H, Ishizuka B, Tadokoro M: Correlation between expression of oncogene products and resistance to anticancer drugs in cultured ovarian cancer cell lines. *Hum Cell* 2003, **16**:131-139.
14. Rowinsky EK, Donehower RC, Jones RJ, Tucker RW: Microtubule changes and cytotoxicity in leukemic cell lines treated with Taxol. *Cancer Res* 1988, **48**:4093-4100.
15. Takahashi M, Kigawa J, Minagawa Y, Itamochi H, Shimada M, Kamazawa S, Sato S, Akeshima R, Terakawa N: Sensitivity to paclitaxel is not related to p53-dependent apoptosis in ovarian cancer cells. *Eur J Cancer* 2000, **36**:1863-1868.
16. Ho ES, Lai CR, Hsieh YT, Chen JT, Lin AJ, Hung MH, Liu FS: p53 mutation is infrequent in clear cell carcinoma of the ovary. *Gynecol Oncol* 2001, **80**:189-193.
17. Enomoto T, Kuragaki C, Yamasaki M, Sugita N, Otsuki Y, Ikegami H, Matsuzaki M, Yamada T, Wakimoto A, Murata Y: Is clear cell carcinoma and mucinous carcinoma of the ovary sensitive to combination chemotherapy with paclitaxel and carboplatin? *Proc Am Soc Clin Oncol* 2003, **22**:447(#1797).
18. Katsumata N, Yasuda M, Takahashi F, Isonishi S, Jobo T, Aoki D, Tsuda H, Sugiyama T, Kodama S, Kimura E, Ochiai K, K Noda, Japanese Gynecologic Oncology Group: Dose-dense paclitaxel once a week in combination with carboplatin every 3 weeks for advanced ovarian cancer: a phase 3, open-label, randomized controlled trial. *Lancet* 2009, **374**:1331-1338.
19. Hsiang YH, Liu LF, Wall ME, Wani MC, Nicholas AW, Manikumar G, Kirschenbaum S, Silber R, Potmesil M: DNA topoisomerase-I mediate DNA cleavage and cytotoxicity of camptothecin analogs. *Cancer Res* 1989, **49**:4385-4389.

doi:10.1186/1757-2215-5-16

Cite this article as: Takatori et al.: Are platinum agents, paclitaxel and irinotecan effective for clear cell carcinoma of the ovary? DNA damage detected with γ H2AX induced by anticancer agents. *Journal of Ovarian Research* 2012 **5**:16.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



Case of peptide-YY-producing strumal carcinoid of the ovary: A case report and review

Eriko Takatori, Tadahiro Shoji, Jiyu Miura, Satoshi Takeuchi, Akira Yoshizaki and Toru Sugiyama

Department of Obstetrics and Gynecology, Iwate Medical University, Morioka, Japan

Abstract

Ovarian carcinoid is a rare tumor accounting for approximately 0.1% of all ovarian malignancies. We describe a case of peptide-YY-producing strumal carcinoid of the ovary associated with severe constipation. A 48-year-old woman was found to have a pelvic mass on ultrasonography when she visited her primary doctor for a health check-up. She was thus referred to our department. Magnetic resonance imaging revealed a solid right ovarian tumor 60 × 50 mm in size. The patient underwent a right adnexectomy and was histopathologically diagnosed as having strumal carcinoid of the ovary. On immunohistochemical examination, the tumor cells were positive for peptide YY. The patient's constipation resolved rapidly after surgery. Based on her clinical course, her constipation was considered to have been caused by the strumal carcinoid of the ovary. The clinical course of this case supports the previously recognized correlation between peptide-YY-producing ovarian carcinoid and constipation.

Key words: constipation, new carcinoid syndrome, peptide YY, strumal carcinoid.

Introduction

Carcinoid is a tumor originating from argyrophil cells (peripheral endocrine cells) located in organs throughout the body including the mucosa of the gastrointestinal tract. Ovarian carcinoid is relatively rare, accounting for approximately 1.3% of all carcinoids. Only about 150 cases have been reported worldwide, and 30 of these cases in Japan. Strumal carcinoid is histologically characterized by an intimate mixture of follicles containing thyroid tissue and carcinoid components, and only about 60 such cases have been reported worldwide. Ovarian carcinoid is a rare tumor accounting for approximately 0.1% of ovarian tumors. It is generally considered to be a borderline malignancy. This tumor is often accompanied by other types of tumors, such as teratoma and mucinous adenoma, and tends to be identified incidentally when specimens

from these tumors are examined. The clinical characteristics and histogenesis of ovarian carcinoids remain unclear. We describe a case of strumal carcinoid of the ovary with severe constipation (a newly recognized carcinoid syndrome), with a review of the relevant literature.

Case Report

The patient was a 48-year-old woman who had a history of 2 gravitas, 2 para, an inguinal hernia at age 17 and appendectomy at age 34. Her family history was unremarkable. The patient visited her primary doctor for a health check-up. She was found to have a pelvic mass on ultrasonography and was thus referred to our department. On pelvic examination, the uterus was egg-sized and a movable eagle egg-sized mass was palpated at the right side of the uterus. Transvaginal

Received: November 10 2011.

Accepted: January 3 2012.

Reprint request to: Dr Tadahiro Shoji, Department of Obstetrics and Gynecology, Iwate Medical University, Uchimarui 19-1, Morioka, Iwate 020-8505, Japan. Email: tshoji@iwate-med.ac.jp

Figure 1 (a) Transvaginal ultrasonography and (b) magnetic resonance imaging of the pelvic cavity showed a 5-cm diameter mass in the right ovary comprised of solid and cystic parts.




Figure 2 Macroscopic finding of right ovary. (a) Macroscopic appearance of a smooth-surfaced, firm, mobile, unruptured right ovarian tumor measuring approximately 6 cm at its largest diameter. (b) On excision, the yellow, gray, and white part was found to contain solid components, whereas the cystic part contained brown fluid.



ultrasonography revealed a mass shadow 51 × 39 mm in size on the right side of the uterus (Fig. 1a). There was no ascites in the Pouch of Douglas. Levels of the tumor markers were 1.5 ng/mL for carcinoembryonic antigen (CEA), 14.1 U/mL for carbohydrate antigen (CA)19-9, and 14.6 IU/mL for CA125. Magnetic resonance imaging revealed a mass, 60 × 50 mm in size, consisting of a solid component with a wide range of signal intensities (Fig. 1b). The patient underwent laparotomy based on a suspicion of ovarian malignancy. Intraoperatively, a fist-sized multilocular ovarian mass with a smooth surface was identified (Fig. 2a). The cut surface of the tumor was solid, showing a mixture of yellow, gray, and white (Fig. 2b). Since ascites was minimal and no abnormalities were found in the left adnexa, the uterus, or the abdominal cavity, the patient underwent right uterine adnexectomy only.

On histopathological examination of the isolated tumor, most of the tissue was arranged in a trabecular formation and the tumor cells had medium-sized round nuclei, showing a pattern consistent with trabecular carcinoid (Fig. 3). On immunohistological examination, the tumor cells were positive for synaptophysin and chromogranin A (Fig. 4a,b). The tumor was diagnosed as a strumal carcinoid because there was a

mixture of thyroid tissue and carcinoid components. The carcinoid tumor cells were positive for thyroglobulin (Fig. 4c). Furthermore, the tumor cells were strongly positive for peptide YY (Fig. 4d). The postoperative pathological diagnosis was strumal carcinoid of the ovary stage Ia (pT1A, NX, M0), and ascites cytology was negative. As no other lesions were identified on gastrointestinal endoscopy or computed tomography, the patient was diagnosed as having primary strumal carcinoid of the ovary. After this diagnosis had been made, a more detailed medical history was obtained from the patient, and it was found that she had suffered severe constipation (a newly recognized carcinoid syndrome). After adnexectomy, the patient recovered from this constipation. She has remained alive and well, without any evidence of recurrence, for 18 months, to date, since surgery.

Discussion

Carcinoids usually arise in the gastrointestinal tract, lungs, bronchial tubes, thymus gland, and mediastinum. Ovarian carcinoid is relatively rare, accounting for approximately 1.3% of all carcinoids.¹ Furthermore, ovarian carcinoid accounts for approximately 0.1% of




Figure 3 Hematoxylin-eosin staining of the strumal carcinoid tumor. (a) Carcinoid tumor composed of solid parts and showing a trabecular growth pattern (original magnification $\times 10$). (b) Carcinoid cells with oblong nuclei and columnar cytoplasm presented with long, wavy, parallel ribbon-like arrangement (original magnification $\times 40$).




Figure 4 Immunohistochemical staining of strumal carcinoid showing positive staining for (a) thyroglobulin, $\times 4$ (b) synaptophysin, $\times 10$ (c) chromogranin A, $\times 10$ and (d) anti-peptide YY antibody, $\times 10$. Most of the cytoplasm of the carcinoid cells was stained with the anti-PYY antibody.

all ovarian malignancies.² According to the General Rules for Clinical and Pathological Management of Ovarian Tumors, carcinoid, strumal carcinoid, and mucinous carcinoid fall under the heading of 'monodermal and highly specialized teratomas,' a category which includes germ cell tumors. Carcinoids are classified into insular and trabecular types, based on their histopathological patterns.³ Strumal carcinoid is histologically characterized by an intimate mixture of follicles containing thyroid tissue and carcinoid components. According to the report by Robboy *et al.* on 50 cases ranging in age from 21 to 77 years with strumal carcinoids of the ovary, half the cases showed trabecular carcinoids while two had insular carcinoids, and the others showed a mixture in which either the trabe-

cular or the insular pattern predominated.⁴ The thyroid component was either normal thyroid tissue or follicular adenoma.

Patients with insular carcinoids present with carcinoid syndrome (flushing, diarrhea, and so on), whereas those with a trabecular or strumal carcinoid rarely develop carcinoid syndrome, although some do reportedly have severe constipation.⁵⁻¹⁰

Peptide YY is a 36-amino acid gastrointestinal hormone localized in endocrine cells of the distal small bowel. This peptide exerts an inhibitory effect on the peristaltic actions of the jejunum and colon. The reported correlation between peptide-YY-producing ovarian carcinoids and constipation is assumed to reflect this inhibitory effect on peristalsis.⁵⁻¹⁰ Matsuda