

Table 5. Characteristics of study sample – prevalence (%) by occupational group.

Country/ Occupational Group	Sex	Age (years)				Age finished full time education (years)				Years in current job	Hours worked/week		
		Males	20–29	30–39	40–49	50–59	<14	14–16	17–19		20+	>5	<30
Brazil													
Nurses	11.4	15.7	24.9	43.8	15.7	32.6	38.6	13.6	15.2	90.3	5.6	87.2	7.3
Office workers	21.7	1.4	23.1	57.3	18.1	36.9	35.0	17.9	10.2	86.6	50.5	44.7	4.8
Other workers	94.6	32.3	34.4	23.7	9.7	59.1	21.6	12.5	6.8	57.1	0.0	100.0	0.0
Ecuador													
Nurses	0.0	6.8	17.8	33.8	41.6	1.8	2.3	29.7	66.2	78.5	73.5	26.5	0.0
Office workers	0.0	11.9	19.8	44.9	23.5	0.4	0.0	35.8	63.8	77.0	3.3	90.5	6.2
Other workers	0.0	43.6	41.4	11.9	3.1	52.0	19.4	11.9	16.7	39.6	2.2	90.3	7.5
Colombia													
Office workers	37.0	27.2	44.6	25.0	3.3	0.0	6.5	17.4	76.1	64.1	26.1	64.1	9.8
Costa Rica													
Nurses	33.6	32.3	28.2	25.9	13.6	2.3	3.2	22.2	72.2	65.1	0.5	72.1	27.4
Office workers	38.1	32.7	27.8	25.6	13.9	0.5	1.4	21.2	77.0	63.3	1.4	94.6	4.1
Other workers	36.6	49.8	23.4	16.1	10.7	0.0	0.5	27.9	71.6	49.0	16.1	82.4	1.5
Nicaragua													
Nurses	3.2	7.4	34.0	37.9	20.6	0.4	2.5	10.7	86.4	88.3	1.1	91.4	7.5
Office workers	27.4	33.3	35.1	22.1	9.5	0.7	4.6	7.4	87.4	57.9	5.3	93.3	1.4
Other workers	54.8	51.8	37.1	7.1	4.1	9.6	24.4	35.0	31.0	21.8	0.0	100.0	0.0
UK													
Nurses	10.1	24.5	37.4	26.1	12.1	0.0	23.7	31.9	44.4	73.4	27.6	72.4	0.0
Office workers	44.7	14.7	31.3	32.1	21.8	0.0	11.1	21.6	67.4	62.5	1.6	94.1	4.3
Other workers	62.4	5.4	19.9	36.8	37.8	0.8	31.5	33.3	34.4	85.5	21.8	70.9	7.3
Spain													
Nurses	9.9	25.0	29.2	29.4	16.4	0.3	7.8	15.4	76.5	72.4	11.8	87.3	0.9
Office workers	16.4	16.7	37.7	34.7	11.0	0.0	2.5	21.7	75.8	67.4	11.6	88.1	0.2
Italy													
Nurses	16.4	17.5	34.9	32.5	15.1	3.5	11.2	19.4	65.9	79.3	13.1	86.1	0.8
Other workers	28.1	5.0	36.0	37.4	21.6	16.5	33.1	40.3	10.1	83.2	9.6	90.4	0.0
Greece													
Nurses	12.1	5.8	67.0	27.2	0.0	0.0	0.4	18.3	81.3	92.0	0.5	97.3	2.3
Office workers	25.1	7.0	46.2	32.7	14.1	0.0	0.0	20.1	79.9	86.4	16.1	71.9	12.1
Other workers	82.9	1.4	12.1	57.9	28.6	2.9	2.1	66.4	28.6	88.6	2.9	92.9	4.3
Estonia													
Nurses	0.5	15.1	31.3	26.1	27.5	0.3	10.3	46.7	42.7	70.0	5.8	86.4	7.8
Office workers	15.3	17.3	31.2	27.7	23.8	0.0	0.0	20.5	79.5	66.3	5.0	89.0	6.0
Lebanon													
Nurses	33.7	57.6	31.0	9.8	1.6	0.5	0.0	4.9	94.6	48.4	0.0	97.3	2.7
Office workers	42.4	20.3	31.4	30.2	18.0	0.0	1.2	15.1	83.7	70.9	0.0	85.5	14.5
Other workers	52.6	53.3	29.9	12.4	4.4	26.3	29.2	29.9	14.6	47.4	0.0	70.8	29.2
Iran													
Nurses	18.3	32.5	46.7	17.9	2.8	0.0	0.8	12.2	87.0	68.7	0.8	65.9	33.3
Office workers	35.2	49.5	34.6	14.8	1.1	0.5	0.5	30.8	68.1	50.0	1.1	63.7	35.2
Pakistan													
Nurses	25.7	72.2	23.0	3.7	1.1	0.0	4.3	29.0	66.7	36.4	0.5	26.7	72.7
Office workers	82.2	53.9	34.4	10.6	1.1	0.0	1.7	17.4	80.9	48.0	1.1	35.0	63.9
Other workers	100.0	9.9	22.5	53.6	14.0	0.9	7.8	25.1	66.2	86.9	16.7	77.5	5.9

Table 5. Cont.

Country/ Occupational Group	Sex	Age (years)				Age finished full time education (years)				Years in current job >5	Hours worked/week		
		Males	20–29	30–39	40–49	50–59	<14	14–16	17–19		20+	<30	30–49
Sri Lanka													
Nurses	0.0	46.2	38.6	12.7	2.5	0.0	0.8	38.6	60.6	50.4	0.0	34.3	65.7
Office workers	71.7	75.7	19.1	2.6	2.6	0.0	0.0	12.5	87.5	30.9	0.0	36.8	63.2
Other workers (1)	100.0	0.4	8.4	46.0	45.2	3.6	65.2	28.0	3.2	81.6	0.0	21.6	78.4
Other workers (2)	0.0	67.5	17.9	10.6	4.0	2.6	29.1	47.0	21.2	40.4	0.0	25.8	74.2
Japan													
Nurses	3.4	43.1	32.6	13.5	10.8	0.0	0.0	10.1	89.9	62.5	5.7	59.6	34.7
Office workers	56.5	4.5	36.1	32.9	26.5	0.0	1.3	13.2	85.5	73.9	13.1	50.7	36.3
Other workers (1)	99.6	20.9	40.4	27.4	11.3	0.0	5.7	65.8	28.5	78.3	14.3	15.3	70.5
Other workers (2)	93.2	29.0	50.1	17.7	3.1	0.0	1.4	4.8	93.8	78.3	8.8	12.7	78.5
South Africa													
Nurses	3.6	16.2	31.6	37.2	15.0	0.0	0.8	18.0	81.2	69.6	0.0	100.0	0.0
Office workers	32.3	42.8	28.4	20.5	8.3	0.4	11.2	62.3	26.0	41.9	0.0	100.0	0.0
Australia													
Nurses	6.8	13.2	29.6	29.2	28.0	0.0	6.8	31.3	61.8	57.8	43.1	48.4	8.5
New Zealand													
Nurses	5.6	8.5	21.5	35.6	34.5	0.6	14.7	37.3	47.5	75.7	32.2	62.7	5.1
Office workers	6.2	4.1	12.4	40.0	43.4	0.7	40.7	49.0	9.7	71.7	31.7	64.8	3.5
Other workers	33.6	18.6	17.7	31.0	32.7	0.0	37.2	46.0	16.8	54.9	47.3	51.8	0.9

doi:10.1371/journal.pone.0039820.t005

12 months. These restrictions were set when the CUPID study was first planned, the latter because some outcomes of interest from the baseline survey, such as sickness absence in the past 12 months, would otherwise be difficult to interpret.

The questions used in the baseline and follow-up surveys were for the most part well-established, having been used successfully in previous studies. In particular, the items on mental health and somatising tendency were taken from validated instruments, and have previously demonstrated predictive validity for the incidence and persistence of musculoskeletal symptoms [7]. Similarly, the questions on fear avoidance beliefs were based on a validated questionnaire [25], and have shown predictive validity in a longitudinal study [7]. The questions on occupational physical activities have been successfully used in earlier studies [7,13,23,24], and the consistency of answers with expectation (e.g. the high prevalence of prolonged keyboard use in office workers) supports their validity. There is no reliable standard against which to assess the accuracy with which subjective symptoms such as pain are reported, but the questions about pain and disability had again been used successfully in earlier studies. Moreover, the style of our questions about symptoms was similar to that of the Nordic questionnaire, which has been shown to have acceptable reliability [28].

Ensuring the accuracy with which the questionnaire was translated into local languages was a challenge. Care was taken to check the accuracy of translation by independent back-translation to English, and this revealed a number of problems. One was the distinction between “stairs” and “flights of stairs”, and despite attempts to resolve this problem, it is not certain that the term “30 flights of stairs” was always interpreted correctly. Therefore, this question will be ignored in future analyses based on the full dataset. Another difficulty arose with questions of the form

“Do you expect that your back pain will be a problem in 12 months time”. In some languages this became “Do you expect your back pain will be a problem over the next 12 months”. Attempts were made to correct this misunderstanding, but it is possible that they were not fully successful.

In addition, terms such as “pain” may be understood differently in different languages even though translated as closely as possible. For this reason, when comparing countries, differences in the relative frequency of pain at different anatomical sites may be particularly revealing – there should have been little ambiguity in the understanding of anatomical sites since they were depicted clearly in diagrams. Interpretation should also be assisted by the questions that were asked about associated difficulty with tasks of daily living, since these were probably understood more uniformly.

Another difficulty that had not been expected was in the use of dates. It emerged that some participants in Iran and Japan used different numbering for calendar years, and where this occurred, corrections had to be made.

Some local investigators opted to include extra questions in addition to the core questions prescribed by CUPID. However, these additions were relatively minor and generally followed after the core questions. Thus, it seems unlikely that they will have influenced answers to the core questions importantly.

Ideally, all questionnaires would have been completed in the same way (interview or self-administration) by all participants. However, this proved impractical. Some occupational groups (especially manual workers in developing countries) would have had great difficulty in answering a written questionnaire, while some employers were unwilling to release their staff for interviews. Moreover, in New Zealand, where nurses and office workers were recruited from across the country, interviews would have been prohibitively expensive.

Table 6. Physical activities in an average working day – prevalence (%) by occupational group.

Country/Occupational Group	Activity ^a					
	Use keyboard >4 hours	Other repeated wrist/hand movement >4 hours	Repeated elbow bending >1 hour	Hands above shoulder height >1 hr	Lifting ≥25 kg by hand	Kneeling/squatting >1 hour
Brazil						
Nurses	9.7	51.9	68.1	11.9	49.7	34.1
Office workers	70.8	70.8	81.5	12.5	10.3	13.2
Other workers	0.0	100.0	100.0	0.0	0.0	100.0
Ecuador						
Nurses	8.2	82.6	89	36.1	68.0	62.6
Office workers	84.0	78.6	84.8	39.1	5.3	16.0
Other workers	11.5	92.1	95.2	82.4	21.1	79.3
Colombia						
Office workers	90.2	62.0	72.8	18.5	6.5	4.3
Costa Rica						
Nurses	10.9	66.4	82.7	30.9	63.6	44.1
Office workers	96.0	76.2	84.8	19.3	5.4	9.4
Other workers	99.0	86.3	88.3	20.5	4.9	4.9
Nicaragua						
Nurses	0.7	78.4	83.0	35.8	42.2	50.0
Office workers	89.8	91.6	84.9	46.0	13.3	17.2
Other workers	4.1	73.6	81.7	26.4	13.2	14.7
UK						
Nurses	12.8	44.0	54.9	8.9	28.4	18.7
Office workers	88.9	31.1	27.1	1.3	4.2	0.5
Other workers	4.1	81.9	91.2	51.8	12.2	9.8
Spain						
Nurses	18.9	59.4	93.7	52.5	82.2	70.5
Office workers	96.8	71.0	91.8	27.4	2.1	14.8
Italy						
Nurses	4.9	55.4	80.2	24.6	60.6	17.0
Other workers	10.1	84.2	85.6	29.5	26.6	4.3
Greece						
Nurses	2.7	71.4	88.8	29.0	70.1	30.4
Office workers	87.4	58.8	74.9	6.0	7.0	6.5
Other workers	1.4	83.6	96.4	65.7	47.1	22.1
Estonia						
Nurses	18.1	64.4	72.5	21.0	56.6	28.6
Office workers	94.6	40.6	51.0	8.4	2.5	2.5
Lebanon						
Nurses	3.3	97.3	96.2	42.9	51.6	34.2
Office workers	85.5	73.8	77.3	13.4	14.5	7.0
Other workers	1.5	98.5	97.1	45.3	44.5	25.5
Iran						
Nurses	10.2	63.0	81.3	43.1	24.8	49.6
Office workers	97.3	89.6	81.3	40.1	7.1	18.7
Pakistan						
Nurses	54.5	93.6	64.2	90.9	73.3	23.0
Office workers	91.7	95.6	35.6	83.9	24.4	10.0
Other workers	7.2	78.4	30.2	77.5	25.7	7.2

Table 6. Cont.

Country/Occupational Group	Activity*					
	Use keyboard >4 hours	Other repeated wrist/hand movement >4 hours	Repeated elbow bending >1 hour	Hands above shoulder height >1 hr	Lifting \geq 25 kg by hand	Kneeling/squatting >1 hour
Sri Lanka						
Nurses	1.3	60.6	43.2	14.4	36.9	9.3
Office workers	100.0	94.7	72.4	11.8	25.7	17.1
Other workers (1)	0.0	95.6	95.6	95.6	0.0	0.0
Other workers (2)	0.7	86.1	60.9	25.2	4.6	29.1
Japan						
Nurses	23.5	23.8	72.8	12.5	66.9	48.5
Office workers	89.0	12.9	22.6	1.6	3.2	2.3
Other workers (1)	2.4	32.8	77.8	33.7	83.3	52.3
Other workers (2)	27.9	10.1	30.1	4.2	9.3	12.1
South Africa						
Nurses	11.3	76.1	85.0	53.4	80.2	26.3
Office workers	100.0	76.9	78.6	26.2	4.8	1.3
Australia						
Nurses	25.6	32.8	47.6	8.4	25.2	15.2
New Zealand						
Nurses	26.6	32.8	42.4	4.0	31.6	14.1
Office workers	91.7	40.0	44.8	0.7	2.1	0.0
Other workers	10.6	87.6	91.2	34.5	51.3	5.3

doi:10.1371/journal.pone.0039820.t006

To explore whether the two methods of answering the questionnaire might lead to systematic differences in answers, we therefore elected to interview a random subset of UK participants while collecting data from the remainder by self-administration. Comparison of responses using the two approaches (Table 10) suggests that no major bias will have occurred as a consequence using both interviews and self-administration. However, if appropriate, method of data collection can be taken into account in statistical analyses.

Participation rates among subjects eligible for study were mostly high, but were less than 50% in five occupational groups (Table 2). We have no reason to expect that those who elected to take part were importantly unrepresentative in the prevalence of pain and its associations with risk factors. However, in future work it may be appropriate to carry out sensitivity analyses, excluding the occupational groups with the lowest response rates. The incomplete response to the baseline questionnaire will be less of a concern in longitudinal analyses based on the follow-up questionnaire.

The numbers of participants by occupational group that were suitable for analysis ranged from 92 to 1018 with a mean of 264. At the outset, our aim was to recruit at least 200 subjects in each group, and this was for the most part achieved (only 7 groups provided fewer than 150 subjects). Furthermore, the occupational groups studied varied substantially in their employment conditions (Table 3), access to healthcare (Table 4), and prevalence of psychosocial risk factors (Tables 7, 8, and 9). When exploring possible reasons for differences in the prevalence of pain and disability between occupational groups, it will be important to

investigate these group-level characteristics as well as individual-level risk factors such as mental health and somatising tendency. The heterogeneity in their distribution should enhance statistical power to address their impact.

As might be expected, the demographic constitution of occupational groups also varied. In particular, many of the samples of nurses were largely or completely female, whereas some groups of "other workers" were all men. This reflects the nature of the occupations of interest. However, it should not be a major problem in interpretation of comparisons since there were an adequate number of occupational groups with a fairly even distribution of sex and age. Moreover, the occurrence of common musculoskeletal complaints appears not to vary greatly between men and women or between older and younger adults of working age [13,23,24].

In summary, the CUPID study is a major resource for the investigation of cultural and psychological determinants of common musculoskeletal disorders and associated disability. Although the data collected have inevitable limitations, the large differences in psychosocial risk factors (including knowledge and beliefs about MSDs) between occupational groups carrying out similar physical tasks in different countries should allow the study hypothesis to be addressed effectively. It will also allow exploration of differences in patterns of musculoskeletal complaint between the three categories of occupation examined, and the consistency of these differences across countries.

Table 7. Psychosocial aspects of work – prevalence (%) by occupational group.

Country/Occupational Group	Incentives ^a	Time pressure ^b	Lack of choice ^c	Lack of support ^d	Job dissatisfaction ^e	Perceived job insecurity ^f
Brazil						
Nurses	25.4	65.4	13.5	4.9	7.6	20.0
Office workers	13.9	49.8	9.6	11.7	19.2	24.9
Other workers	100.0	96.8	96.8	2.2	5.4	90.3
Ecuador						
Nurses	29.2	69.4	39.7	51.6	1.8	30.1
Office workers	37.0	63.4	10.7	63.4	4.5	29.2
Other workers	45.8	65.2	52.0	63.4	11.5	50.7
Colombia						
Office workers	50.0	56.5	2.2	40.2	2.2	25.0
Costa Rica						
Nurses	48.2	92.7	24.5	36.8	12.7	17.7
Office workers	63.2	77.6	8.1	28.7	10.8	18.4
Other workers	67.8	77.6	50.7	29.3	17.1	26.3
Nicaragua						
Nurses	16.0	72.3	10.3	41.5	13.5	22.7
Office workers	26.0	80.0	19.3	43.2	9.5	23.2
Other workers	86.8	60.9	37.1	41.1	6.1	31.0
UK						
Nurses	6.2	75.1	9.7	10.1	14.8	17.9
Office workers	0.5	76.6	6.8	7.9	7.9	5.0
Other workers	19.2	79.5	37.8	17.4	15.5	35.8
Spain						
Nurses	21.0	80.1	19.9	77.7	12.0	16.5
Office workers	26.3	54.3	32.4	78.5	6.6	13.7
Italy						
Nurses	11.6	80.6	13.2	8.2	17.4	21.5
Other workers	19.4	82.7	53.2	34.5	51.8	41.7
Greece						
Nurses	6.3	97.3	8.9	14.7	33.9	29.0
Office workers	6.5	83.4	1.5	9.5	7.0	12.6
Other workers	2.1	97.9	15.0	40.7	18.6	17.9
Estonia						
Nurses	7.8	66.6	23.7	27.0	6.2	14.3
Office workers	4.0	64.4	2.0	8.4	5.9	23.3
Lebanon						
Nurses	81.0	95.1	6.0	6.5	20.1	38.6
Office workers	11.6	75.6	7.6	12.2	16.9	25.0
Other workers	75.9	76.6	29.9	6.6	16.8	41.6
Iran						
Nurses	28.9	90.2	24.8	23.6	29.3	54.9
Office workers	29.7	74.2	18.7	26.9	26.4	66.5
Pakistan						
Nurses	62.0	96.3	40.1	7.5	9.1	56.7
Office workers	68.3	96.1	45.6	7.8	7.8	53.9
Other workers	11.7	95.0	68.0	7.7	9.0	14.9
Sri Lanka						
Nurses	56.8	91.5	5.9	7.2	4.7	11.4
Office workers	18.4	87.5	10.5	5.3	8.6	43.4

Table 7. Cont.

Country/Occupational Group	Incentives ^a	Time pressure ^b	Lack of choice ^c	Lack of support ^d	Job dissatisfaction ^e	Perceived job insecurity ^f
Other workers (1)	100.0	100.0	0.0	0.0	2.8	1.6
Other workers (2)	95.4	94.0	17.2	11.9	4.0	33.8
Japan						
Nurses	4.4	63.0	20.9	5.7	44.4	41.2
Office workers	3.2	35.5	18.1	12.6	70.3	43.5
Other workers (1)	30.7	81.1	28.0	20.1	41.9	64.5
Other workers (2)	9.9	41.4	4.5	5.4	69.6	49.6
South Africa						
Nurses	21.1	80.2	23.1	13.8	34.8	29.6
Office workers	52	95.2	37.6	21.8	43.7	66.4
Australia						
Nurses	4.4	66.8	3.2	7.6	8.8	10.8
New Zealand						
Nurses	1.7	58.2	9.0	8.5	13.6	22.0
Office workers	2.1	58.6	4.8	18.6	8.3	17.9
Other workers	34.5	80.5	23.9	14.2	8.8	20.4

^aEither a) piecework or b) payment of a bonus if more than an agreed number of articles/tasks are finished in a day.

^bEither a) a target number of articles or tasks to be finished in the day or b) working under pressure to complete tasks by a fixed time.

^cChoice seldom or never in all of: a) how work is done, b) what is done at work, and c) work timetable and breaks.

^dSupport from colleagues or supervisor/manager seldom or never.

^eDissatisfied or very dissatisfied overall.

^fFeel job would be rather unsafe or very unsafe if off work for three months with significant illness.

doi:10.1371/journal.pone.0039820.t007

Table 8. Awareness of repetitive strain injury (RSI) work related upper limb disorder (WRULD) or cumulative trauma syndrome (CTS) – prevalence (%) by occupational group.

Country/Occupational Group	Proportion (%) of participants reporting awareness of				
	RSI, WRULD or CTS	Someone outside work with pain in past 12 months in			
		Low back	Neck	Upper limb	Knee
Brazil					
Nurses	94.6	62.7	49.2	53.0	55.1
Office workers	94.3	60.9	49.1	52.7	50.2
Other workers	0.0	60.2	12.9	36.6	14.0
Ecuador					
Nurses	52.1	42.9	34.7	30.1	42.5
Office workers	28.0	50.6	46.1	37.0	42.4
Other workers	24.2	48.0	27.3	39.2	32.2
Colombia					
Office workers	43.5	40.2	34.8	32.6	39.1
Costa Rica					
Nurses	54.1	55.9	43.6	42.7	46.4
Office workers	26.9	61.0	49.3	48.4	45.7
Other workers	36.1	74.6	65.9	65.9	61.5
Nicaragua					
Nurses	56.0	71.6	57.8	58.2	62.8
Office workers	34.0	60.4	54.0	51.2	48.8
Other workers	29.4	41.6	28.4	31.5	26.9
UK					
Nurses	76.3	59.1	30.0	35.0	41.2
Office workers	93.7	60	31.8	33.4	42.6
Other workers	47.9	42.5	21.0	26.7	35.0
Spain					
Nurses	67.9	82.6	73.1	49.8	55.9
Office workers	59.8	82.9	80.2	45.3	50.6
Italy					
Nurses	84.7	82.3	75.6	56.0	55.4
Other workers	77.0	69.8	66.9	54.0	51.1
Greece					
Nurses	21.4	82.6	62.5	56.3	50.4
Office workers	24.6	81.4	68.3	64.8	51.3
Other workers	15.7	70.7	50	43.6	36.4
Estonia					
Nurses	66.6	69.0	55.3	46.9	57.1
Office workers	49.5	65.8	59.4	47.0	51.5
Lebanon					
Nurses	67.9	70.1	58.2	39.1	57.6
Office workers	67.4	56.4	40.7	36.6	32.6
Other workers	34.3	38.7	27.7	16.1	29.2
Iran					
Nurses	45.5	76.8	53.3	59.3	69.5
Office workers	25.3	67.0	46.7	54.4	63.2
Pakistan					
Nurses	36.9	44.4	23.5	31.0	52.4
Office workers	17.8	39.4	15.0	20	41.1
Other workers	32.4	30.6	19.8	18.9	26.6

Table 8. Cont.

Country/Occupational Group	Proportion (%) of participants reporting awareness of				
	RSI, WRULD or CTS	Someone outside work with pain in past 12 months in			
		Low back	Neck	Upper limb	Knee
Sri Lanka					
Nurses	48.3	53.0	40.3	45.8	61.0
Office workers	51.3	45.4	36.8	37.5	47.4
Other workers (1)	82.4	57.2	27.6	36.0	57.2
Other workers (2)	36.4	37.1	20.5	25.2	45.0
Japan					
Nurses	72.3	59.5	27.4	35.8	33.6
Office workers	69.4	53.5	28.7	33.5	35.8
Other workers (1)	35.9	51.6	17.5	22.5	20.5
Other workers (2)	70.7	60.8	23.4	27.0	26.8
South Africa					
Nurses	47.0	51.4	36.4	34.8	53.8
Office workers	7.0	55.0	38.4	39.3	40.2
Australia					
Nurses	78.0	71.6	49.2	49.6	53.2
New Zealand					
Nurses	84.7	72.3	53.1	58.2	57.6
Office workers	95.9	64.1	44.8	47.6	54.5
Other workers	86.7	46.9	27.4	37.2	42.5

doi:10.1371/journal.pone.0039820.t008

Table 9. Adverse health beliefs regarding low back and arm pain – prevalence (%) by occupational group.

Country/ Occupational Group	Low back pain			Arm pain		
	Commonly caused by people's work ^a	Physical activity is harmful ^b	Poor prognosis ^c	Commonly caused by people's work ^a	Physical activity is harmful ^b	Poor prognosis ^c
Brazil						
Nurses	25.9	5.9	29.7	31.9	7.0	31.4
Office workers	32.7	7.5	31.3	42.7	6.0	31.0
Other workers	0.0	1.1	0.0	0.0	1.1	0.0
Ecuador						
Nurses	53.9	25.1	20.5	52.1	18.7	20.5
Office workers	37.9	18.9	10.7	33.7	16.0	9.9
Other workers	77.1	36.1	4.0	76.2	27.3	5.3
Colombia						
Office workers	12.0	1.1	13.0	13.0	1.1	13.0
Costa Rica						
Nurses	30.0	10.9	17.7	35.0	10.5	19.1
Office workers	13.9	4.0	24.2	11.7	2.7	22.0
Other workers	16.1	2.9	25.9	18.0	2.0	21.5
Nicaragua						
Nurses	36.2	23.8	15.2	35.5	21.3	14.5
Office workers	29.1	11.9	9.5	32.3	12.6	9.1
Other workers	38.1	22.3	10.7	36.5	16.8	8.6
UK						
Nurses	23.7	9.3	5.8	15.2	3.5	2.7
Office workers	9.2	2.9	4.7	10.8	1.3	3.2
Other workers	25.6	10.4	8.8	20.7	5.2	5.7
Spain						
Nurses	46.8	23.8	28.2	36.1	13.8	18.3
Office workers	22.4	15.5	22.1	19.6	9.6	15.3
Italy						
Nurses	34.1	3.2	6.9	24.1	0.9	4.5
Other workers	36.0	7.9	15.8	40.3	3.6	16.5
Greece						
Nurses	73.2	49.1	14.7	68.3	33.5	12.9
Office workers	40.2	31.2	10.6	44.2	18.6	12.6
Other workers	78.6	68.6	20.0	76.4	47.1	12.9
Estonia						
Nurses	27.5	9.2	7.5	25.9	5.9	5.9
Office workers	15.8	2.5	11.4	21.3	0.5	10.9
Lebanon						
Nurses	77.7	43.5	27.2	62.5	23.9	9.8
Office workers	36.6	24.4	15.1	36.0	11.0	7.6
Other workers	66.4	77.4	14.6	59.9	57.7	6.6
Iran						
Nurses	31.7	11	2.8	24.8	4.1	1.6
Office workers	24.2	12.1	4.9	22.0	2.7	1.6
Pakistan						
Nurses	51.9	50.3	5.9	47.1	26.2	4.8
Office workers	54.4	43.3	3.9	38.9	29.4	1.7
Other workers	40.5	31.5	5.9	36.9	28.4	6.3

Table 9. Cont.

Country/ Occupational Group	Low back pain			Arm pain		
	Commonly caused by people's work ^a	Physical activity is harmful ^b	Poor prognosis ^c	Commonly caused by people's work ^a	Physical activity is harmful ^b	Poor prognosis ^c
Sri Lanka						
Nurses	5.9	6.4	9.3	9.7	3.0	11.4
Office workers	13.8	10.5	4.6	19.7	4.6	3.9
Other workers (1)	4.0	36.0	10.4	3.6	11.2	8.0
Other workers (2)	20.5	9.9	7.3	20.5	6.0	6.0
Japan						
Nurses	46.6	14.7	18.2	24.3	5.7	9.3
Office workers	16.5	19.7	14.2	11.6	9.0	7.4
Other workers (1)	47.2	25.6	21.8	33.2	11.7	10.1
Other workers (2)	21.4	23.7	17.5	12.4	16.1	6.5
South Africa						
Nurses	37.7	5.3	7.7	36.0	3.6	6.1
Office workers	24.9	6.6	4.8	22.7	3.1	3.5
Australia						
Nurses	19.2	2.8	6.8	12.4	2.4	2.4
New Zealand						
Nurses	20.3	2.8	2.3	11.9	1.1	4.0
Office workers	6.2	2.1	2.8	9.0	2.1	4.1
Other workers	21.2	14.2	6.2	29.2	12.4	5.3

^aCompletely agree that such pain is commonly caused by people's work.

^bCompletely agree that for someone with such pain, a) physical activity should be avoided as it might cause harm, and b) rest is needed to get better.

^cCompletely agree that for someone with such pain, rest is needed to get better, and completely disagree that such problems usually get better within three months.

doi:10.1371/journal.pone.0039820.t009

Table 10. Comparison of UK participants who provided information by interview and by self-administered questionnaire.

	Nurses		Office workers		Other workers	
	Interview	Self-administered questionnaire	Interview	Self-administered questionnaire	Interview	Self-administered questionnaire
Number selected	190	500	200	851	240	1329
Number (%) participated	91 (48)	199 (40)	88 (44)	388 (46)	122 (51)	320 (24)
Number of subjects analysed	78	179	66	314	110	276
Prevalence (%) of activities in an average working day						
Use keyboard >4 hr	6.4	15.6	84.9	89.8	1.8	5.1
Other repeated wrist/hand movement >4 hr	46.2	43.0	22.7	32.8	86.4	80.1
Repeated elbow bending >1 hr	60.3	52.5	13.6	29.9	96.4	89.1
Hands above shoulder height >1 hr	7.7	9.5	1.5	1.3	55.5	50.4
Lifting ≥25 kg by hand	28.2	28.5	9.1	3.2	12.7	12.0
Kneeling/squatting >1 hr	21.8	17.3	1.5	0.3	15.5	7.6
Prevalence (%) of pain in past month						
Low back	26.9	36.3	28.8	26.8	34.6	34.4
Neck	14.1	20.1	21.2	22.9	20.9	20.7
Shoulder	9.0	21.8	21.2	20.7	33.6	31.2
Elbow	2.6	2.8	12.1	8.0	14.6	15.2
Wrist/hand	14.1	15.6	19.7	17.5	24.6	21.7
Knee	12.8	18.4	27.3	22.3	21.8	24.6

doi:10.1371/journal.pone.0039820.t010

Supporting Information

Appendix S1 Committees which provided ethical approval for the cupid study.
(DOCX)

Appendix S2 Baseline questionnaire.
(DOCX)

Appendix S3 Follow-up questionnaire.
(DOCX)

Acknowledgments

We thank: Pietro Muñoz, Patricio Oyos, Gonzalo Albuja, María Belduma and Francisco Lara for their assistance with data collection in Ecuador; Patrica Monge, Melania Chaverri and Freddy Brenes, who helped with data collection in Costa Rica; Aurora Aragón, Alberto Berrios, Samaria Balladares and Martha Martínez who helped with data collection in Nicaragua; Alfredo José Jirón who assisted with data entry in Nicaragua; Catalina Torres for translation and piloting of the questionnaire in Spain; Ben and Marie Carmen Coggon for back translation of the Spanish questionnaire; Cynthia Alcantara, Xavier Orpella, Josep Anton Gonzalez, Joan Bas, Pilar Peña, Elena Brunat, Vicente San José, Anna Sala March, Anna Marquez, Josefina Lorente, Cristina Oliva, Montse Vergara and Eduard Gaynés for their assistance with data collection in Spain; Natale Battevi, Lorenzo Bordini, Marco Conti and Luciano Riboldi who carried out data collection in Italy; Paul Maurice Conway for back translation of the Italian questionnaire; Tiina Freimann, who helped with data collection

References

- Lötters F, Burdorf A, Kuiper J, Miedema H (2003) Model for the work-relatedness of low-back pain. *Scand J Work Environ Health* 29: 431–40.
- Palmer KT, Smedley J (2007) Work relatedness of chronic neck pain with physical findings – a systematic review. *Scand J Work Environ Health* 33: 165–91.
- Palmer KT (2011) Carpal tunnel syndrome: the role of occupational factors. *Best Pract Res Clin Rheumatol* 25: 15–29.
- Palmer KT, Harris EC, Coggon D (2007) Compensating occupationally related tenosynovitis and epicondylitis: a literature review. *Occup Med* 57: 67–74.
- Linton SJ (2000) A review of psychological risk factors in back and neck pain. *Spine* 25: 1148–56.
- Macfarlane GJ, Hunt IM, Silman AJ (2000) Role of mechanical and psychosocial factors in the onset of forearm pain: prospective population based study. *Br Med J* 321: 676–9.
- Palmer KT, Reading I, Linaker C, Calnan M, Coggon D (2008) Population-based cohort study of incident and persistent arm pain: role of mental health, self-rated health and health beliefs. *Pain* 136: 30–37.
- Macfarlane GJ, Pallewatte N, Paudyal P, Biyth FM, Coggon D, et al. (2009) Evaluation of work-related psychosocial factors and regional musculoskeletal pain: result from a EULAR Task Force. *Ann Rheum Dis* 68: 885–91.
- Palmer K, Calnan M, Wainwright D, Poole J, O'Neill C, et al. (2005) Disabling musculoskeletal pain and its relation to somatization: A community-based postal survey. *Occup Med* 55: 612–617.
- Clinical Standards Advisory Group (1994) *Epidemiology review: the epidemiology and cost of back pain*. London: HMSO.
- Gun RT (1990) The incidence and distribution of RSI in South Australia 1980–81 and 1986–87. *Med J Aust* 153: 376–80.
- Coggon D (2005) Occupational Medicine at a turning point. *Occup Environ Med* 62: 281–3.
- Madan I, Reading I, Palmer KT, Coggon D (2008) Cultural differences in musculoskeletal symptoms and disability. *Int J Epidemiol* 37: 1181–1189.
- Ramond A, Bouton C, Richard I, Roquelaire Y, Baufreton C, et al. (2011) Psychosocial risk factors for chronic low back pain in primary care – a systematic review. *Fam Pract* 28: 12–21.
- Buchbinder R, Jolley D, Wyatt M (2001) Population based intervention to change back pain beliefs and disability: three part evaluation. *Br Med J* 322: 1516–1520.
- Ferrari AL, Baptista PCP, Felli VEA, Coggon D (2010) Translation, adaptation and validation of the “Cultural and Psychosocial Influences on Disability (CUPID) Questionnaire” for use in Brazil. *Rev Latino-Am Enfermagem* 18: 1092–8.
- Harcombe H, McBride D, Derrett S, Gray A (2009) Prevalence and impact of musculoskeletal disorders in New Zealand nurses, postal workers and office workers. *Aust N Z J Public Health* 33: 437–41.
- Harcombe H, McBride D, Derrett S, Gray A (2010) Physical and psychosocial risk factors for musculoskeletal disorders in New Zealand nurses, postal workers and office workers. *Injury Prevention* 16: 96–100.
- Hoe VCW, Kelsall HL, Urquhart DM, Sim MR (2011) Risk factors for musculoskeletal symptoms of the neck or shoulder alone or neck and shoulder among hospital nurses. *Occup Environ Med* 69: 198–204.
- Matsudaira K, Palmer KT, Reading I, Hirai M, Yoshimura N, et al. (2011) Prevalence and correlates of regional pain and associated disability in Japanese workers. *Occup Environ Med* 68: 191–196.
- Solidaki E, Chatzi L, Bitsios P, Markatzi I, Plana E, et al. (2010) Work related and psychological determinants of multi-site musculoskeletal pain. *Scand J Work Environ Health* 2010;36: 54–61.
- Warnakulasariya S, Peiris-John R, Ntani G, Coggon D, Sathiakumar N, et al. (In press) Musculoskeletal pain in four occupational populations in Sri Lanka. *Occupational Medicine*.
- Palmer KT, Walsh K, Bendall H, Cooper C, Coggon D (2000) Back pain in Britain: Comparison of two prevalence surveys at an interval of ten years. *Br Med J* 320:1577–8.
- Palmer KT, Walker-Bone K, Griffin MJ, Syddall H, Pannett B (2001) The prevalence and occupational associations of neck pain in the British population. *Scand J Work Environ Health* 27: 49–56.
- Waddell G, Newton M, Henderson I, Somerville D, Main CJ (1993) A Fear-Avoidance Beliefs Questionnaire (FABQ) and the role of fear-avoidance beliefs in chronic low back pain and disability. *Pain* 52: 157–68.
- Derogatis LR, Melisaratos N (1983) The Brief Symptom Inventory: an introductory report. *Psychol Med* 13: 595–605.
- Ware JE, Sherbourne CD (1992) The MOS 36-item short-form health survey (SF-36). *Med Care* 30: 473–83.
- Kuorinka I, Jonsson B, Kilbom A, Vinterberg H, Biering-Sørensen F, et al. (1987) Standardised Nordic questionnaires for the analysis of musculoskeletal symptoms. *Appl Ergon*;18: 233–7.

Available online at www.sciencedirect.com

Public Health

journal homepage: www.elsevier.com/puhe

Original Research

Prevalence of vertebral fracture in Asian men and women: Comparison between Hong Kong, Thailand, Indonesia and Japan

A.W.L. Kwok^{a,*}, J.C.S. Leung^a, A.Y.H. Chan^a, B.S.K. Au^a, E.M.C. Lau^b, H. Yurianto^c, P. Yuktanandana^d, N. Yoshimura^e, S. Muraki^f, H. Oka^e, T. Akune^f, P.C. Leung^a

^aJockey Club Centre for Osteoporosis Care and Control, The Chinese University of Hong Kong, Hong Kong, China

^bCenter for Clinical and Basic Research, Hong Kong, China

^cFaculty of Medicine, Hasanuddin University, Indonesia

^dDepartment of Orthopedics, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand

^eDepartment of Joint Disease Research, 22nd Century Medical and Research Centre, University of Tokyo, Tokyo, Japan

^fDepartment of Clinical Motor System Medicine, 22nd Century Medical and Research Centre, University of Tokyo, Tokyo, Japan

ARTICLE INFO

Article history:

Received 14 December 2010

Received in revised form

31 January 2012

Accepted 12 March 2012

Available online 5 May 2012

Keywords:

Vertebral fracture

Asia

Elderly

Quality of life

SUMMARY

Objectives: Little is known about the prevalence of vertebral fracture among Asians. This study investigated the prevalence of radiographically defined vertebral fracture, and identified associated risk factors in the aged population of four Asian countries.

Study design: In total, 1588 males and females aged ≥ 65 years were recruited from Hong Kong, Thailand, Indonesia and Japan.

Methods: Standard X-rays for the spine were taken and vertebral heights were measured. Vertebral fracture was defined as a reduction of >3 standard deviations in vertebral height ratio. Bone mineral density (BMD) of the hip was measured by dual energy X-ray absorptiometry, and anthropometric measurements were taken in Hong Kong and Japan. Other relevant data were entered in a standard questionnaire.

Results: The prevalence of vertebral fracture for both males and females was highest in Japan for younger (65–74 years) and older (≥ 75 years) age groups (36.6% and 37.6% for males; 18.8% and 28.7% for females). Lower hip BMD was associated with vertebral fracture in both sexes. Older age, lower quality of life score on Short Form-12 (physical), past longest occupation as a farmer, and history of cataract were significantly associated with vertebral fracture in females. However, smoking did not appear to be an important risk factor for vertebral fracture.

Conclusions: Radiographic assessments for vertebral fracture were performed in all four Asian countries. The prevalence of vertebral fracture was highest in Japan. Lower hip BMD, poorer physical condition and past longest occupation as a farmer were associated with vertebral fracture.

© 2012 The Royal Society for Public Health. Published by Elsevier Ltd. All rights reserved.

* Corresponding author. Jockey Club Centre for Osteoporosis Care and Control, The Chinese University of Hong Kong, Room 316, 3/F, School of Public Health, Prince of Wales Hospital, Shatin, N.T., Hong Kong SAR China. Tel.: +852 2252 8830; fax: +852 2649 2447.

E-mail addresses: anthonykwok@cuhk.edu.hk (A.W.L. Kwok), jason-leung@cuhk.edu.hk (J.C.S. Leung).

0033-3506/\$ – see front matter © 2012 The Royal Society for Public Health. Published by Elsevier Ltd. All rights reserved.
doi:10.1016/j.puhe.2012.03.002

Introduction

Osteoporosis is a condition characterized by low bone mass and micro-architectural deterioration of bone tissues causing increased bone fragility and susceptibility to fracture. The disease is becoming a major health problem in Asia where a rapidly increasing aged population will be accompanied by increasing incidence of hip fracture.¹ In Hong Kong, for instance, over the last two decades, the number of cases of hip fracture has increased two-fold due to the increase in the ageing population. Of every 1000 people who are aged ≥ 80 years, one man and two women will have a fractured hip each year.² The age-adjusted rate in Hong Kong is comparable to that in Singapore and higher than that in Japan. It has been projected that over half of all hip fractures in the world will occur in Asia by 2050.¹

While there are many data on the prevalence and risk factors for hip fracture, little is known of the prevalence and risk factors for vertebral fracture in Asia. In a study by Lau *et al.*, the prevalence of vertebral fracture was estimated to be 29% in women in Hong Kong.³ Vertebral deformity, back pain and psychological problems were associated symptoms in elderly Chinese women, and the use of analgesics was common in men. Overseas studies have demonstrated associations between vertebral fracture and back pain, disability and the need for healthcare support.^{4–6} The survival rate of elderly people who had experienced a clinical vertebral fracture was found to be lower than that of elderly people who had not experienced a fracture.⁷ Women with severe vertebral deformities were found to have a consistently higher risk of back pain and height loss. In addition, genetic and lifestyle factors may influence vertebral fracture. However, the risk factors for Asian men are still unclear.

The aim of this study was to investigate the prevalence of radiographically defined vertebral fracture, and to identify underlying risk factors associated with vertebral fracture in the aged population in four Asian countries (Hong Kong, Thailand, Indonesia and Japan). Existing data from Beijing, China were used for comparison.

Methods

Subjects

In total, 1588 subjects were recruited from Hong Kong, Thailand, Indonesia and Japan. Each country was required to recruit 400 ambulatory community-dwelling subjects (200 males and 200 females), half of whom were aged 65–74 years and the other half were aged ≥ 75 years. They all were ethnic Asians. Subjects were recruited in community centres (recreation centres for group activities, social support and public information) for the elderly. Recruitment notices were placed at these centres explaining the purpose and procedures of this study in different districts, including urban and rural areas. Moreover, community activities and community advertisements were used to enhance the representativeness of the whole region.

Radiographs and digitization

Radiographic films of the lateral thoracic and lumbar spine were taken with a tube-to-film distance of 100 cm, with thoracic films centred at T8 and lumbar films centred at L3. The radiographs were evaluated by morphometry using a backlight translucent digitizing table (GTCO, Rockville, MD, USA) and cursor. Six points were marked on the radiographs with a wax pencil for each vertebral body T4–L4, and the X,Y co-ordinates for each point were recorded on an electronic grid with a resolution of 0.1 mm. In total, 120 and 68 subjects lacked T4 and T5, respectively. Standardized procedures were adopted. The six points corresponded to the four corners of the vertebral body and the midpoints of the end plates⁸ (Fig. 1). Vertebral height ratios (VHR) were calculated [anterior to posterior (Ha/Hp), middle to posterior (Hm/Hp) and posterior above to posterior below (Hp/Hp–1 and Hp/Hp+1)]. Prevalent vertebral fracture was defined as a reduction of >3 standard deviations (SD) among any one of the VHRs.^{6,8} Mean (SD) VHRs are shown in Table 1. X-ray films for Hong Kong, Thailand and Indonesia were measured in Hong Kong by three trained members of staff, all with >10 years of experience and trained by the radiologist-in-charge with standardized procedures. Fifteen films were selected at random for reliability analysis. Whereas X-ray films from Japan were measured in Japan, 30 were selected at random and sent to Hong Kong for reliability analysis. The analysis was performed by two members of staff (one from Hong Kong and one from Japan).

Questionnaire

Subjects were interviewed using a standardized and structured questionnaire in Hong Kong, Thailand and Indonesia by trained staff. Data collected included demographic information, Mini-Mental Status Examination score, medical history, fall history, smoking habit, alcohol consumption, Physical Activity Screening for Elderly score, quality of life,




Fig. 1 – Measurement of vertebral height. Hp, posterior height; Hm, mid-body height; Ha, anterior height.

Table 1 – Adjusted means^a and standard deviations (SD) of vertebral height ratios in Chinese male and female subjects.

Level	Ha/Hp		Hm/Hp		Hp/Hp-1		Hp/Hp+1	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Male^b								
T4	0.87	0.06	0.91	0.04	–	–	1.00	0.06
T5	0.87	0.07	0.92	0.04	1.00	0.06	0.98	0.06
T6	0.87	0.07	0.92	0.04	1.03	0.06	0.98	0.06
T7	0.86	0.07	0.92	0.04	1.02	0.06	1.00	0.06
T8	0.89	0.07	0.93	0.04	1.00	0.06	0.98	0.06
T9	0.91	0.07	0.94	0.04	1.02	0.06	0.95	0.06
T10	0.92	0.07	0.94	0.04	1.05	0.06	0.91	0.06
T11	0.86	0.08	0.91	0.05	1.10	0.07	0.93	0.05
T12	0.84	0.07	0.90	0.04	1.08	0.06	0.95	0.05
L1	0.84	0.06	0.89	0.04	1.06	0.05	1.01	0.04
L2	0.88	0.06	0.91	0.03	1.00	0.04	1.01	0.04
L3	0.92	0.06	0.94	0.04	0.99	0.04	1.06	0.05
L4	0.97	0.07	1.00	0.05	0.95	0.04	–	–
Female^c								
T4	0.90	0.08	0.92	0.06	–	–	0.99	0.09
T5	0.88	0.08	0.91	0.06	1.03	0.09	0.99	0.07
T6	0.86	0.07	0.91	0.05	1.03	0.07	0.99	0.07
T7	0.86	0.08	0.91	0.05	1.02	0.07	0.99	0.06
T8	0.88	0.07	0.92	0.05	1.01	0.06	0.99	0.06
T9	0.92	0.08	0.94	0.05	1.01	0.06	0.97	0.06
T10	0.92	0.08	0.94	0.05	1.04	0.07	0.95	0.06
T11	0.87	0.08	0.91	0.05	1.06	0.07	0.94	0.06
T12	0.86	0.09	0.90	0.06	1.07	0.07	0.96	0.07
L1	0.87	0.08	0.89	0.05	1.05	0.08	1.00	0.05
L2	0.93	0.08	0.91	0.05	1.01	0.06	1.01	0.06
L3	0.98	0.08	0.95	0.06	0.99	0.06	1.04	0.07
L4	1.02	0.10	1.01	0.07	0.96	0.07	–	–

Ha/Hp, anterior height of vertebrae/posterior height of vertebrae; Hm/Hp, middle height of vertebrae/posterior height of vertebrae; Hp/Hp-1, posterior height of vertebrae/posterior height of vertebrae above; Hp/Hp+1, posterior height of vertebrae/posterior height of vertebrae below.

a Adjusted means were estimated as the mode on the log-frequency curve after trimming 5% of the values from each end of the distribution. The adjusted SDs were estimated by quantile-quantile plot after 10% of values were trimmed from each end of the distribution.

b Data used by Lau et al.³⁰

c Data published by Lau et al.³

fracture history, back pain and musculoskeletal symptoms, medications, diet history, Geriatric Depression Scale score, and past longest occupation in their life.

Dual X-ray densitometry and anthropometric measurement

Bone mineral density (BMD), height and weight were measured in Hong Kong and Japan. Hip and spine BMD were measured by dual energy X-ray absorptiometry (DXA) using a Hologic QDR-4500W densitometer (Hologic, Waltham, MA, USA). The same type of DXA machine was used in both Hong Kong and Japan. The coefficient of variation (CV) in the laboratory was 0.7% and 0.9% for hip and spine, respectively. Height was measured using a Holtain Harpenden stadiometer (Holtain Ltd., Crosswell, UK), and weight was

measured using the Physician Beam Balance Scale (Healthometer, IL, USA) with subjects wearing a light gown.

Study flow

The study started in Hong Kong, Indonesia and Thailand. Radiographs and questionnaires were performed in these three countries. Japan joined the study at a later date; radiographs, BMD and anthropometric measurements were conducted but no questionnaire was used. As different information was collected from the four countries, the results are shown in three parts. First, the prevalence of vertebral fracture was compared between the four countries. Second, lifestyle and medical risk factors for vertebral fracture were analysed for Hong Kong, Indonesia and Thailand. Finally, anthropometric factors in Japan were compared with those for Hong Kong.

Statistical analysis

Inter-rater reliability of each parameter was assessed using the intraclass correlation coefficient (ICC). The prevalence of deformity was calculated based on the number of individuals with at least one vertebral deformity, and this was compared between countries using Chi-squared test. Data were analysed separately by sex. Continuous variables were compared between the countries using t-test or analysis of variance, while categorical variables were compared using Chi-squared test. Logistic regression (with adjustment for age) was used to calculate odds ratios (OR) and 95% confidence intervals (CI) of various lifestyle and anthropometric factors for definite vertebral deformity. Significant factors were collated to form the final models. All statistical analyses were performed using SAS Version 9.1 (SAS Institute, Inc., Cary, NC, USA). An α level of 5% was used as the level of significance.

Results

Inter-rater reliabilities of vertebral measurements were fair to good. ICC (3,1) ranged from 0.54 to 0.94 for anterior heights, from 0.70 to 0.96 for middle heights, and from 0.60 to 0.92 for posterior heights among 15 randomly selected subjects from Hong Kong, Thailand and Indonesia. ICC (3,1) ranged from 0.73 to 0.93 for anterior heights, from 0.69 to 0.93 for middle heights, and from 0.76 to 0.95 for posterior heights among 30 Japanese subjects.

Part 1: prevalence of vertebral fracture in four Asian countries

Seven hundred and seventy males (mean age 72.9 years) and 818 females (mean age 72.4 years) were recruited. The prevalence of vertebral fracture in Hong Kong, Indonesia, Thailand and Japan is shown in Table 2. Among males, the prevalence of fracture was highest in Japan (36.6% for age 65–74 years, 37.6% for age ≥ 75 years); the difference between Japan and the other three countries was significant. The prevalence of fracture was lowest in Hong Kong males (9.2%

Table 2 – Prevalence of vertebral fracture.

	Prevalence of vertebral fracture (%)				P-value of Chi-square
	Hong Kong (H)	Indonesia (I)	Thailand (T)	Japan (J)	
Male					
Age 65–74 years	(n = 98)	(n = 127)	(n = 105)	(n = 101)	
Radiological fracture					
Definite vertebral deformity, <mean – 3 SD	9.2	15.0	17.1	36.6 ^{a,b,c}	<0.0001
Severe vertebral deformity, <mean – 4 SD	3.1	7.1	13.3 ^a	20.8 ^{a,b}	0.0003
Reported fracture	0.0	0.0	2.9	–	0.0402
Age ≥75 years	(n = 100)	(n = 40)	(n = 98)	(n = 101)	
Radiological fracture					
Definite vertebral deformity, <mean – 3 SD	18.0	20.0	24.5	37.6 ^{a,b,c}	0.0102
Severe vertebral deformity, <mean – 4 SD	9.0	17.5	20.4 ^a	20.8 ^a	0.0926
Reported fracture	2.0	0.0	1.0	–	0.6125
Female					
Age 65–74 years	(n = 99)	(n = 170)	(n = 98)	(n = 101)	
Radiological fracture					
Definite vertebral deformity, <mean – 3 SD	2.0	7.6	8.2 ^a	18.8 ^{a,b,c}	0.0010
Severe vertebral deformity, <mean – 4 SD	1.0	5.9	6.1	10.9 ^a	0.0338
Reported fracture	1.0	0.0	3.0	–	0.2930
Age ≥75 years	(n = 98)	(n = 54)	(n = 97)	(n = 101)	
Radiological fracture					
Definite vertebral deformity, <mean – 3 SD	15.3	13.0	17.5	28.7 ^{a,b}	0.0727
Severe vertebral deformity, <mean – 4 SD	10.2	9.3	15.5	18.8	0.2312
Reported fracture	1.0	0.0	3.0	–	0.2930

SD, standard deviation.

Note: a higher number of reported fractures was significantly associated with a higher number of radiological fractures (Chi-squared = 7.078, P-value = 0.0078).

a P-value < 0.05 comparing Indonesia (I), Thailand (T) or Japan (J) with Hong Kong (H), by Chi-squared test.

b P-value < 0.05 comparing Thailand (T) or Japan (J) with Indonesia (I), by Chi-squared test.

c P-value < 0.05 comparing Japan (J) with Thailand (T), by Chi-squared test.

for age 65–74 years; 18% for age ≥75 years). In females, the prevalence of fracture was also highest in Japan (18.8% for age 65–74 years; 28.7% for age ≥75 years), and was lowest in Hong Kong and Indonesia (2% for age 65–74 years; 13% for age ≥75 years). The reported fracture rates ranged from 0 to 2.97%, which were much lower than radiological fracture rates (0–24.5%) (Table 2). More reported fractures were significantly associated with more radiological fractures (Chi-squared = 7.078, P-value = 0.0078).

Part 2: lifestyle and medical factors of vertebral fracture in Hong Kong, Indonesia and Thailand

Results for Hong Kong, Indonesia and Thailand are shown in Tables 3 and 4. Subjects' characteristics are shown in Table 3. Males and females in Hong Kong had significantly higher quality of life [Short Form(SF)-12 (mental)]; higher prevalence of hypertension, heart disease and myocardial infarction/angina; and lower levels of education compared with Indonesia and Thailand. The percentage of respondents who stated that their longest occupation had been as a farmer was highest in Thailand.

Risk factors of vertebral fracture are shown in Table 4. For males, older age, lower SF-12 (physical) score, and history of stroke were significantly associated with vertebral fracture. For females, older age, lower SF-12 (physical) score, history of cataract, and past longest occupation as a farmer were

significantly associated with vertebral fracture. In logistic regression, a lower SF-12 (physical) score (OR 0.68, 95% CI 0.55–0.84) was associated with vertebral fracture among the males. Older age (OR 1.33, 95% CI 1.03–1.71), lower SF-12 (physical) score (OR 0.73, 95% CI 0.56–0.95), history of cataract (OR 2.86, 95% CI 1.48–5.53), and past longest occupation as a farmer (OR 2.47, 95% CI 1.07–5.72) were associated with vertebral fracture in females.

Part 3: anthropometric factors of vertebral fracture in Hong Kong and Japan

To investigate the difference in fracture rate between Japan and Hong Kong, 25 males and 27 females in Japan were selected at random as subsamples to obtain anthropometric factors: weight and height and bone density. Results are shown in Tables 5 and 6. Differences in characteristics between Hong Kong and Japan are shown in Table 5. Subjects in Hong Kong were heavier and taller (marginally in males but significantly in females). The body mass index of Hong Kong females was also significantly higher. Logistic regressions of vertebral fractures are shown in Table 6. Lower hip BMD (OR 1.47, 95% CI 1.001–2.17 per SD reduction) was associated with vertebral fracture in males, while in females, older age (OR 1.9, 95% CI 1.13–3.2), heavier weight (OR 1.6, 95% CI 1.1–2.3) and lower hip BMD (OR 3.23, 95% CI 1.72–6.25 per SD reduction) were associated with vertebral fracture.

Table 3 – Characteristics of subjects by country.

Variable	Male			P-value ^a	Female			P-value ^a
	Mean (SD)/%				Mean (SD)/%			
	Hong Kong (n = 198)	Indonesia (n = 167)	Thailand (n = 203)		Hong Kong (n = 197)	Indonesia (n = 224)	Thailand (n = 195)	
Age	73.7 (5.4)	70.3 (5.1)	73.8 (6.0)	0.9610	73.8 (5.5)	70.3 (5.8)	73.7 (5.5)	0.9078
SF-12 (physical)	49.7 (9.0)	47.2 (10.4)	46.0 (10.9)	0.0003	45.3 (9.0)	41.8 (11.6)	43.6 (11.6)	0.1085
SF-12 (mental)	56.4 (5.9)	52.1 (8.1)	53.1 (8.7)	<0.0001	55.7 (8.1)	52.3 (8.3)	52.7 (9.1)	0.0005
Physical activity (PASE score)	104.8 (49.8)	102.7 (67.9)	93.0 (80.8)	0.0880	81.3 (30.7)	70.8 (42.7)	70.5 (47.6)	0.0081
Marital status: married	84.1%	82.4%	81.3%	0.7564	57.7%	62.3%	50.5%	0.0456
Education				<0.0001				<0.0001
Primary or below	53.2%	41.9%	64.5%		76.6%	55.1%	72.9%	
Secondary	33.8%	43.8%	14.2%		18.9%	34.8%	15.1%	
Tertiary or above	12.9%	14.4%	21.3%		4.5%	10.1%	12.0%	
Demented (MMSE < 24)	11.0%	13.4%	15.8%	0.3634	36.3%	22.9%	25.0%	0.0046
Diabetes	14.9%	10.5%	11.3%	0.3690	9.0%	14.2%	13.0%	0.2243
Reported osteoporosis	5.0%	1.7%	3.5%	0.2361	9.0%	11.6%	10.0%	0.6492
Stroke	6.0%	2.3%	4.4%	0.2263	3.5%	2.2%	2.0%	0.5768
Hypertension	39.3%	19.2%	22.7%	<0.0001	42.3%	26.7%	25.0%	0.0002
Heart disease	19.9%	2.3%	4.9%	<0.0001	14.4%	3.5%	4.5%	<0.0001
Myocardial infarction/angina	18.4%	1.2%	4.9%	<0.0001	11.4%	4.3%	3.5%	0.0014
Congestive heart failure	2.5%	0.6%	2.5%	0.3150	3.5%	1.3%	1.0%	0.1328
Chronic obstructive pulmonary disease	15.9%	5.2%	3.0%	<0.0001	6.5%	4.3%	1.5%	0.0424
Cataracts	33.3%	18.6%	22.2%	0.0024	44.8%	23.3%	27.0%	<0.0001
Gastrectomy	9.0%	8.1%	11.3%	0.5425	4.0%	14.2%	7.0%	0.0005
Cancer	4.5%	0.6%	1.5%	0.0268	3.5%	2.2%	1.5%	0.4115
Fall in past 12 months	22.9%	4.1%	21.7%	<0.0001	20.9%	14.2%	31.5%	<0.0001
Smoker	9.5%	29.2%	19.4%	<0.0001	0.5%	1.4%	1.0%	0.6472
Use of alcohol at least 12 times in past 12 months	22.9%	5.5%	36.1%	<0.0001	0.5%	0.5%	6.5%	<0.0001
Fracture after 50 years of age	6.0%	5.5%	9.4%	0.2739	16.9%	5.6%	7.6%	0.0002
Past longest occupation as a farmer	1.0%	14.3%	44.8%	<0.0001	3.1%	5.7%	40.0%	<0.0001

SF-12, Short Form-12; PASE, physical activity screening for elderly; MMSE, Mini-Mental State Examination.

Bold: P-value < 0.05.

a P-value of Chi-squared for categorical variables or analysis of variance for continuous variables where appropriate.

Discussion

The occurrence of osteoporotic vertebral fracture is particularly common in older postmenopausal Asian women. The common sites involved are T12 and L1 as these levels are, biomechanically, the most vulnerable regiments of the thoracolumbar spine.⁹ The first vertebral fracture is an important indicator for subsequent vertebral fracture, and possibly non-vertebral fracture including hip fracture.^{10,11} Understanding the prevalence of vertebral fracture and deformity is critical for prevention.

Fracture identification can easily be affected by differences in radiographic methods. Fracture definition and methods of morphological assessment also affect the determination of fracture rates. Degenerative spondylosis and old asymptomatic spinal fractures may lead to spuriously high BMD at the site of involvement, thus diminishing the sensitivity in the prediction of fracture. In this study, all the radiographs were

taken according to a standardized protocol, and were analysed by trained and experienced assessors to minimize possible observer bias and to acquire more precise and accurate results. The number of reported vertebral fractures was generally lower in both males and females than the number of radiologically determined vertebral fractures in all four countries. As elsewhere, vertebral fractures were often under-reported. Vertebral fractures could also be overlooked when patients present with back pain arising from various spinal conditions.

This study found that the prevalence of vertebral fracture was significantly higher in Japan compared with Hong Kong, Thailand and Indonesia for both sexes. BMD is an important factor,^{12,13} but in this study, no difference was noted between Hong Kong and Japan. This may be explained by the smaller body size of Japanese subjects. In a systematic review, Ruysen-Witrand *et al.*¹⁴ showed that vertebral size was an independent risk factor for vertebral fracture. After controlling for well-known confounding factors such as age, height,

Table 4 – Characteristics in subjects with vertebral fracture and control.

Variable	Unit	Male		Female	
		Age-adjusted model OR (95% CI)	Multivariate model OR (95% CI)	Age-adjusted model OR (95% CI)	Multivariate model OR (95% CI)
Age	5	1.24 (1.03–1.49) ^a	1.17 (0.96–1.44)	1.48 (1.20–1.84) ^a	1.33 (1.03–1.71)
Country					
Hong Kong	—	1	1	1	1
Indonesia	Indonesia/Hong Kong	1.44 (0.79–2.62)	1.28 (0.68–2.40)	1.36 (0.67–2.72)	1.20 (0.53–2.73)
Thailand	Thailand/Hong Kong	1.66 (0.97–2.82)	1.45 (0.84–2.51)	1.58 (0.82–3.05)	1.36 (0.61–2.99)
SF-12 – Physical	10	0.65 (0.53–0.80)	0.68 (0.55–0.84)	0.67 (0.53–0.85)	0.73 (0.56–0.95)
SF-12 – Mental	10	0.99 (0.75–1.32)		0.86 (0.64–1.17)	
Physical activity (PASE score)	50	1.00 (0.84–1.19)		0.72 (0.50–1.03)	
Marital status – married	Yes/no	1.78 (0.91–3.46)		0.68 (0.40–1.18)	
Education					
Primary or below	—	1		1	
Secondary	Secondary/primary	0.63 (0.36–1.11)		0.96 (0.48–1.91)	
Tertiary or above	Tertiary/primary	1.02 (0.55–1.90)		0.37 (0.09–1.57)	
Demented (MMSE < 24)	Yes/no	1.36 (0.74–2.49)		1.34 (0.76–2.37)	
Diabetes	Yes/no	1.09 (0.56–2.14)		0.67 (0.26–1.75)	
Reported osteoporosis	Yes/no	1.26 (0.41–3.91)		2.01 (0.95–4.24)	
Stroke	Yes/no	2.61 (1.08–6.31)	2.32 (0.94–5.71)	1.43 (0.31–6.60)	
Hypertension	Yes/no	1.08 (0.66–1.77)		0.72 (0.39–1.31)	
Heart disease	Yes/no	0.80 (0.36–1.77)		0.94 (0.35–2.53)	
Myocardial infarction/angina	Yes/no	0.94 (0.42–2.08)		0.88 (0.29–2.60)	
Congestive heart failure	Yes/no	1.67 (0.43–6.47)		1.74 (0.37–8.26)	
Chronic obstructive pulmonary disease	Yes/no	1.02 (0.46–2.27)		—	
Cataracts	Yes/no	1.21 (0.73–1.98)		1.80 (1.05–3.10)	2.86 (1.48–5.53)
Gastroctomy	Yes/no	0.83 (0.38–1.83)		0.72 (0.25–2.09)	
Cancer	Yes/no	—		0.64 (0.08–5.04)	
Fall in past 12 months	Yes/no	0.90 (0.49–1.63)		1.39 (0.76–2.51)	
Smoker	Yes/no	0.62 (0.33–1.20)		—	
Use of alcohol at least 12 times in past 12 months	Yes/no	1.30 (0.78–2.17)		2.12 (0.45–9.95)	
Fracture after 50 years of age	Yes/no	2.00 (0.93–4.31)		1.36 (0.62–2.98)	
Past longest occupation as a farmer	Yes/no	1.19 (0.68–2.08)		2.14 (1.13–4.05)	2.47 (1.07–5.72)

SF-12, Short Form-12; PASE, Physical Activity Screening for Elderly; MMSE, Mini-Mental State Examination; OR, odds ratio; CI, confidence interval.
 Bold: P-value < 0.05.
 a Crude OR.

weight and BMD, small vertebral dimensions including area, cross-sectional area and volume were associated with vertebral fracture.^{14–16} Moreover, there is a difference in the nutrient intake of these two populations; dietary calcium intake in the Japanese population is less than that of the Hong Kong population (550 mg/day in men and 519 mg/day in women vs 628 mg/day and 569 mg/day for people aged ≥65 years).^{17,18} Kobayashi reported that the calcium content of river water in Japan was lower than that in most European countries.¹⁹

For Asian women, a review of vertebral fracture revealed that the prevalence of vertebral fracture increases steeply with age in Beijing: the prevalence of vertebral fracture in Chinese women aged 50–59 years was 3.9%, compared with 10.5%, 15% and 31.2% at ages 60–69, 70–79 and ≥80 years.^{6,20,21} A similar trend has also been observed among Japanese women residing in Japan and Hawaii^{22,23}; the rates were slightly higher for those living in Japan (8%, 25%, 38% and 43% for ages 65–70, 70–75, 75–80 and 80–85 years, compared

with 5%, 15%, 25% and 27% for those living in Hawaii in 1996), suggesting that environmental factors are more influential than genetic factors.²⁴ Horikawa et al. reported a prevalence of 22.6% for Japanese women aged 65–92 years living in a fishing/farming village located in Nansei-cho in 2001.²⁵ Data for Japan in the present study showed similar rates. These data are similar to findings among Caucasian women in the USA, where the prevalence of vertebral fracture was 22% for women aged 70–79 years, and 34% for women aged ≥80 years. Hence, observations from the present study suggest that vertebral fracture is as prevalent in Asian women as in Caucasian women.²⁰

The prevalence of vertebral fracture among Chinese women in 1996 in Hong Kong was 29% at age 70–79 years.^{1,3} The current study found a prevalence of vertebral fracture of 2% at age 65–74 years and 15% at age ≥75 years among Hong Kong Chinese women. The prevalence of vertebral fracture appears to have decreased significantly over the years between these studies. This may reflect a proper

Table 5 – Comparison of characteristics between Hong Kong and Japan.

Variable	Mean (SD)		P-value ^a
	Hong Kong	Japan	
Male	<i>n</i> = 198	<i>n</i> = 25	
Age (years)	73.75 (5.40)	73.28 (4.31)	0.6784
Weight (kg)	61.86 (9.35)	58.34 (6.66)	0.0692
Height (cm)	162.64 (5.75)	160.50 (7.70)	0.0925
Body mass index (kg/m ²)	23.35 (3.13)	22.72 (2.73)	0.3315
Hip BMD (g/cm ²)	0.851 (0.129)	0.823 (0.105)	0.2956
Spine BMD (g/cm ²)	0.950 (0.183)	0.961 (0.151)	0.7742
Female	<i>n</i> = 197	<i>n</i> = 27	
Age (years)	73.76 (5.51)	72.33 (4.67)	0.2012
Weight (kg)	55.01 (9.01)	48.62 (7.98)	0.0006
Height (cm)	151.48 (4.99)	148.61 (6.23)	0.0072
Body mass index (kg/m ²)	23.93 (3.44)	21.98 (3.05)	0.0056
Hip BMD (g/cm ²)	0.697 (0.113)	0.691 (0.112)	0.7654
Spine BMD (g/cm ²)	0.757 (0.141)	0.787 (0.137)	0.3028

BMD, bone mineral density; SD, standard deviation.
^a P-value of t-test for continuous or Chi-squared test for categorical variables.

osteoporosis prevention regime, health promotion and raised public awareness of osteoporosis in Hong Kong since 2001. The Hong Kong experience may set an example to other Asian countries for reference in combating osteoporosis.

Lau et al.¹³ reported that the prevalence of vertebral fracture among Chinese men aged 70–79 years in 1998 in Hong Kong was 16%. The present study shows a similar result; 9.2% and 18% for men aged 65–74 and ≥75 years, respectively.

This study found that both male and female subjects aged ≥75 years with a lower hip BMD have more vertebral fractures. Most vertebral fractures are not caused by direct trauma²⁶ but are linked with low BMD.^{10,12} Previous studies have shown that US women of African descent have the highest bone mass in the spine, while Caucasian women have intermediate values, and Asian women have the lowest values (unadjusted BMD values). An analysis and comparison of different ethnic populations residing in the USA suggested that the BMD of Asian women differed little from that of Caucasian women after adjustment for body size.²⁷ Unadjusted lumbar spine and femoral neck BMDs were 7–12% and 14–24% higher in African-American women compared with Caucasian, Japanese or Chinese women. Among women of comparable weight of <70 kg, there were no differences in lumbar spine BMD among African-American, Chinese and Japanese women, all of whom have higher BMDs than Caucasians.²⁰ This study also noted that there were no differences in lumbar spine and hip BMDs between Hong Kong and Japanese women or men. Kung reported that the pattern and magnitude of age-related bone loss in the lumbar spine was similar between female Asian and Caucasian populations.²⁰ Subjects with a spinal fracture had a spine or hip BMD that was 20–30% lower compared with normal healthy controls. After adjusting for age and body size, each SD reduction in bone density in the Hong Kong Chinese women increased the risk of vertebral fracture approximately two-fold for both spine and femoral neck measurements, whereas in Beijing Chinese women, each SD reduction in BMD at the spine or hip was associated with a 2.5-fold increased risk of vertebral fracture.²⁰ In the present study, among Hong Kong and Japanese women, each SD reduction in hip BMD was associated with a 3.2-fold increased risk of vertebral fracture after adjustment for age and body size, which is consistent with the previous finding. In men, each SD reduction had a 1.5-fold increased risk, showing less

Table 6 – Logistic regression of vertebral fracture.

Variable	Mean (SD)/frequency (%)		Unit	Adjusted OR (95% CI)
	Control	Vertebral fracture		
Male	<i>n</i> = 187	<i>n</i> = 36		
Age	73.57 (5.27)	74.61 (5.29)	5	1.15 (0.81–1.65)
Country				
Hong Kong	171 (91.4%)	27 (75.0%)	–	1
Japan	16 (8.6%)	9 (25.0%)	Japan/Hong Kong	3.67 (1.42–9.49)
Weight (kg)	61.48 (9.17)	60.94 (9.21)	5	1.27 (0.95–1.69)
Height (cm)	162.59 (5.77)	161.07 (7.14)	5	0.77 (0.55–1.09)
Hip BMD (g/cm ²)	0.855 (0.126)	0.816 (0.125)	–0.1	1.47 (1.001–2.17)
Female	<i>n</i> = 201	<i>n</i> = 23		
Age	73.09 (5.31)	77.43 (4.81)	5	1.90 (1.13–3.20)
Country				
Hong Kong	180 (89.6%)	17 (73.9%)	–	1
Japan	21 (10.5%)	6 (26.1%)	Japan/Hong Kong	5.99 (1.58–22.68)
Weight (kg)	54.45 (9.10)	53.42 (9.69)	5	1.61 (1.11–2.33)
Height (cm)	151.51 (5.01)	148.55 (6.39)	5	0.68 (0.39–1.17)
Hip BMD (g/cm ²)	0.707 (0.109)	0.608 (0.115)	–0.1	3.23 (1.72–6.25)

BMD, bone mineral density; SD, standard deviation; OR, odds ratio; CI, confidence interval.
Bold: P-value < 0.05.

influence than that in women. However, low BMD is still a causal factor of vertebral fracture.¹²

Lifestyle and medical factors are other important determinants of vertebral fracture. Women who had jobs involving heavy physical labour had a lower prevalence of vertebral fracture than women who had had more sedentary jobs. This suggests that strenuous physical activity during young adulthood is protective against vertebral fracture.⁶ Gregg et al.²⁸ found that physical activity reduced the risk of hip fracture but not vertebral or wrist fractures. However, in this study, subjects with vertebral fracture had significantly lower SF-12 (physical) scores, which implies that poor physical strength may lead to more vertebral fractures. Stroke is another risk factor for vertebral fracture. Kim et al. reported that in the acute stages of stroke, bone loss progressed rapidly. Risks of osteoporosis and vertebral fracture were higher among subjects who had experienced a stroke.²⁹ People with cataracts were more likely to fall, which increases the risk of fracture.³⁰ The prevalence of vertebral fracture was higher in farmers. This may be explained by the lower intake of calcium. Kamiyama et al. performed a survey in various areas in Japan, and found that calcium intake in three farm villages was lower than that in the city (512, 532 and 601 mg vs 633 mg). The prevalence of osteoporosis was also higher in farm villages.³¹

Subjects from Hong Kong had the highest prevalence of hypertension, heart disease and myocardial infarction/angina, but the association with vertebral fracture was insignificant. This implies that hypertension, heart disease and myocardial infarction/angina are not direct risk factors for vertebral fracture. However, Lau et al.³² reported that hypertension was associated with higher BMD; this may explain the fact that the lowest prevalence of vertebral fracture was seen in Hong Kong.

There were more male smokers in Indonesia. However, smoking in Indonesia did not appear to lead to adverse effects resulting in vertebral fracture. Grazio et al.³³ found that current smoking was a risk factor for vertebral fracture, but a study by Bensen et al.³⁴ did not. Similarly, the rate of falls was lowest in Indonesia, but this did not reflect on the prevalence of vertebral fracture, implying that falls are not a strong risk factor for vertebral fracture. This differed from hip fracture, which was always associated with falls and trauma²⁰ and 90% of cases resulted from falls.³⁵

This study was a joint venture between Hong Kong, Thailand, Indonesia and Japan. Radiographic assessments of vertebral fractures were completed carefully in all four countries using the same methodology. As subjects were recruited using community activities and community advertisements, the subjects were members of the general population and hence the results are representative. The percentages of people who worked as farmers in Hong Kong, Indonesia and Thailand were 2.1%, 9.3% and 42.4% in this study, which is close to the population figures: 0.9%, 12.4% and 39%, respectively.^{36–38} Unfortunately, the Indonesian and Thai collaborators in this study did not possess a DXA machine; otherwise data correlation between the groups could be more comprehensive. Likewise, the questionnaire was not used in Japan, which influenced the completeness of this study. Moreover, it would be more convincing if a study with a larger sample size was conducted. Despite the obvious limitations, this study provides

useful data for prevention planning. Some of these data are similar to the results of previous studies, but some are unique (e.g. strong association between low SF-12 (physical) score and vertebral fracture among Asians; smoking habit was not an important risk factor for vertebral fracture).

Acknowledgements

Ethical approval

The institutional review board at each centre approved the study protocol, and written informed consent was obtained from all participants.

Funding

None declared.

Conflict of interest

None declared.

REFERENCES

1. Lau EMC, Cooper C. The epidemiology of osteoporosis: the oriental perspective in a world context. *Clin Orthop Relat R* 1996;323:65–74.
2. Lau EMC, Woo J, Leung PC, Swaminathan R, Leung D. Low bone mineral density, grip strength and skinfold thickness are important risk factors for hip fracture in Hong Kong Chinese. *Osteoporos Int* 1993;3:66–70.
3. Lau EMC, Chan HHL, Woo J, Lin F, Black D, Nevitt M, et al. Normal ranges for vertebral height ratios and prevalence of vertebral fracture in Hong Kong Chinese: a comparison with American Caucasians. *J Bone Miner Res* 1996;11:1364–8.
4. Gold DT. The clinical impact of vertebral fractures: quality of life in women with osteoporosis. *Bone* 1996;18:S185–9.
5. Rowe SM, Jung ST, Lee JY. Epidemiology of osteoporosis in Korea. *Osteoporos Int* 1997;7:S88–90.
6. Xu L, Cummings SR, Qin M, Zhao X, Chen X, Nevitt M, et al. Vertebral fractures in Beijing, China: the Beijing osteoporosis project. *J Bone Miner Res* 2000;15:2019–25.
7. Huang C, Ross PD, Wasnich RD. Vertebral fractures and other predictors of back pain among older women. *J Bone Miner Res* 1996;11:1026–32.
8. Black DM, Cummings SR, Stone K, Hudes E, Palermo L, Steiger P. A new approach to defining normal vertebral dimensions. *J Bone Miner Res* 1991;6:883–92.
9. Cooper C, Atkinson EJ, O'Fallon WM, Melton 3rd LJ. Incidence of clinically diagnosed vertebral fractures: a population-based study in Rochester, Minnesota, 1985–1989. *J Bone Miner Res* 1992;7:221–7.
10. Cauley JA, Hochberg MC, Lui LY, Palermo L, Ensrud KE, Hillier TA, et al. Long-term risk of incident vertebral fractures. *JAMA* 2007;298:2761–7.
11. Burger H, van Daele PL, Algra D, Hofman A, Grobbee DE, Schutte HE, et al. Vertebral deformities as predictors of non-vertebral fractures. *BMJ* 1994;309:991–2.
12. World Health Organization. *Assessment of fracture risk and application to screening for postmenopausal osteoporosis*. In: WHO technical report series. Geneva: WHO; 1994. pp. 1–129.
13. Lau EM, Chan YH, Chan M, Woo J, Griffith J, Chan HH, et al. Vertebral deformity in Chinese men: prevalence, risk factors,

- bone mineral density, and body composition measurements. *Calcif Tissue Int* 2000;66:47–52.
14. Ruysse-Witrand A, Gossec L, Kolta S. Vertebral dimensions as risk factor of vertebral fracture in osteoporotic patients: a systematic literature review. *Osteoporos Int* 2007;18:1271–8.
 15. Gilsanz V, Boechat MI, Gilsanz R, Loro ML, Roe TF, Goodman WG. Gender differences in vertebral sizes in adults: biomechanical implications. *Radiology* 1994;190:678–82.
 16. Duan Y, Parfitt A, Seeman E. Vertebral bone mass, size, and volumetric density in women with spinal fractures. *J Bone Miner Res* 1999;14:1796–802.
 17. Ministry of Health, Labour and Welfare. *The report of National Health and Nutrition Survey of Japan*. Available at: <http://www.mhlw.go.jp/houdou/2009/11/dl/h1109-1b.pdf>; 2008 [28.10.2010, date last accessed].
 18. Tang NL, Liao CD, Ching JK, Suen EW, Chan IH, Orwoll E, et al. Sex-specific effect of pirin gene on bone mineral density in a cohort of 4000 Chinese. *Bone* 2010;46:543–50.
 19. Kobayashi J. On geographical relationship between the chemical nature of river water and death-rate from apoplexy. *Berichte des Ohara Instituts fur landwirtschafliche Biologie* 1957;11:12–21.
 20. Kung AWC. Epidemiology and diagnostic approaches to vertebral fractures in Asia. *J Bone Miner Metab* 2004;22:170–5.
 21. Liu ZH, Zhao YL, Ding GZ, Zhou Y. Epidemiology of primary osteoporosis in China. *Osteoporos Int* 1997;7:S84–7.
 22. Ross PD, Wasnich RD, Davis JW, Vogel JM. Vertebral dimension differences between Caucasian populations, and between Caucasians and Japanese. *Bone* 1991;12:107–12.
 23. Ross PD, Fujiwara S, Huang C, Davis J, Epstein RS, Wasnich RD, et al. Vertebral fracture prevalence in women in Hiroshima compared to Caucasians or Japanese in the US. *Int J Epidemiol* 1995;24:1171–7.
 24. Huang C, Ross PD, Fujiwara S, Davis JW, Epstein RS, Kodama K, et al. Determinants of vertebral fracture prevalence among native Japanese women and women of Japanese descent living in Hawaii. *Bone* 1996;18:437–42.
 25. Horikawa K, Kasai Y, Yamakawa T, Sudo A, Uchida A. Prevalence of osteoarthritis, osteoporotic vertebral fractures, and spondylolisthesis among the elderly in a Japanese village. *J Orthop Surg* 2006;14:9–12.
 26. Haczynski J, Jakimiuk A. Vertebral fractures: a hidden problem of osteoporosis. *Med Sci Monit* 2001;7:1108–17.
 27. Finkelstein JS, Lee ML, Sowers M, Ettinger B, Neer RM, Kelsey JL, et al. Ethnic variation in bone density in premenopausal and early perimenopausal women: effects of anthropometric and lifestyle factors. *J Clin Endocrinol Metab* 2002;87:3057–67.
 28. Gregg EW, Cauley JA, Seeley DG, Ensrud KE, Bauer DC. Physical activity and osteoporotic fracture risk in older women. Study of Osteoporotic Fractures Research Group. *Ann Intern Med* 1998;129:81–8.
 29. Kim HW, Kang E, Im S, Ko YJ, Im SA, Lee JI. Prevalence of pre-stroke low bone mineral density and vertebral fracture in first stroke patients. *Bone* 2008;43:183–6.
 30. Ivers RQ, Cumming RG, Mitchell P, Peduto AJ. Diabetes and risk of fracture: the Blue Mountains Eye study. *Diabetes Care* 2001;24:1198–203.
 31. Kamiyama S, Kobayashi S, Abe S, Takahashi E, Wakamatsu E. Osteoporosis prevalence and nutritional intake among the people in farm, fishing and urban districts. *Tohoku J Exp Med* 1972;107:387–94.
 32. Lau EM, Leung PC, Kwok T, Woo J, Lynn H, Orwoll E, et al. The determinants of bone mineral density in Chinese men—results from mr. os (Hong Kong), the first cohort study on osteoporosis in Asian men. *Osteoporos Int* 2006;17:297–303.
 33. Grazio S, Korsic M, Jajic I. Effects of smoking and alcohol consumption on vertebral deformity in the elderly – an epidemiological study; evaluation of easily measured risk factors in the prediction of osteoporotic fractures. *Coll Antropol* 2005;29:567–72.
 34. Bensen R, Adachi JD, Papaioannou A, Ioannidis G, Olszynski WP, Sebaldt RJ, et al. Evaluation of easily measured risk factors in the prediction of osteoporotic fractures. *BMC Musculoskel Disord* 2005;6:47.
 35. Tinetti ME. Clinical practice. Preventing falls in elderly persons. *N Engl J Med* 2003;348:42.
 36. Census and Statistics Department The Government of the Hong Kong Special Administrative Region. 2006 population by-census thematic report: older persons. Available at: www.censtatd.gov.hk [accessed 02.01.12].
 37. Hasoloan MA. *Country report the Indonesian labor market. The OECD forum on the restated OECD jobs strategy in Tokyo*. Available at: www.oecd.org/dataoecd/20/36/37873500.pdf; 2006 [accessed 02.01.12].
 38. Darawuttimaprakorn N, Punpuing S. *Living arrangements and elderly depression*. Kanchanaburi DSS: Population Association of America 2010 Annual Meeting Program; 2010.