

Table 2 Incidence of de-novo degenerative lumbar scoliosis in participants, grouped by sex and age

Age at 1st survey	Male	Female	Total
40–49	3/25 (12.0)	1/32 (3.1)	4/57 (7.0)
50–59	2/30 (6.7)	7/43 (16.3)	9/73 (12.3)
60–69	4/23 (17.4)	11/33 (33.3)	15/56 (26.8)
70–79	1/3 (33.3)	4/5 (80.0)	5/8 (62.5)
Total	10/81 (12.3)	23/113 (20.4)	33/194 (17.0)

Percentage incidence are given in parenthesis

Table 3 Radiographic features of de-novo degenerative lumbar scoliosis

Case	Age at the 2nd survey	Gender	Cobb angle at the 2nd survey	Convexity of the curve	Location of the apex	Rotation angle of the apex
1	61	M	26	L	L3	5
2	77	F	12	L	L3	5
3	72	F	10	L	L3	5
4	71	F	12	L	L2	5
5	57	F	12	R	L3	15
6	85	F	10	L	L3	0
7	79	F	10	L	L3	5
8	83	F	17	L	L3	5
9	82	F	24	L	L3	5
10	67	M	10	L	L3	0
11	76	M	21	L	L2	10
12	76	F	10	R	L4	5
13	57	M	10	R	L3	0
14	75	M	18	L	L3	5
15	67	F	15	L	L3	10
16	86	F	19	R	L4	0
17	71	F	10	L	L3	5
18	85	M	10	R	L3	5
19	81	F	14	L	L3	5
20	70	F	11	R	L3	5
21	79	F	12	L	L3	5
22	77	M	11	L	L4	0
23	81	F	13	R	L3	15
24	90	F	13	L	L3	5
25	78	F	12	L	L3	5
26	73	F	11	L	L2	15
27	82	F	12	R	L3	5
28	89	F	18	L	L3	5
29	72	M	10	L	L3	5
30	76	M	18	L	L3	5
31	71	F	10	L	L3	5
32	69	F	13	R	L3	5
33	63	M	12	L	L3	5

Table 4 Results from measurement of radiographic features comparing the S-group (subjects who developed de-novo degenerative lumbar scoliosis during the follow-up period) and the C-group (the subjects who did not develop scoliosis)

	S-group (n = 33)	C-group (n = 161)	P value
Tilting angle			
L1	3.06 ± 2.12	1.19 ± 1.45	<0.01
L2	2.42 ± 2.03	1.13 ± 1.43	<0.01
L3	2.00 ± 1.82	1.23 ± 1.46	<0.01
L4	3.15 ± 3.38	1.20 ± 1.38	<0.01
L5	2.33 ± 1.81	0.98 ± 1.29	<0.01
Intervertebral disc angle			
L1/2	1.03 ± 1.56	0.3 ± 0.67	<0.01
L2/3	1.56 ± 1.88	0.48 ± 0.83	<0.01
L3/4	1.34 ± 1.33	0.71 ± 1.13	<0.01
L4/5	0.78 ± 1.16	0.69 ± 1.17	0.69
L5/S	0.13 ± 0.49	0.30 ± 0.85	0.87
Lateral spondylolisthesis (case)			
L1	0	0	–
L2	0	0	–
L3	5	1	<0.01
L4	0	2	0.52
L5	0	0	–
Axial rotation (case)			
L1	2	5	0.39
L2	5	11	0.12
L3	12	8	<0.01
L4	1	1	0.23
L5	0	2	0.52
Lateral osteophyte difference			
L1/2	1.03 ± 1.56	0.3 ± 0.67	<0.01
L2/3	1.56 ± 1.88	0.48 ± 0.83	<0.01
L3/4	1.34 ± 1.33	0.71 ± 1.13	<0.01
L4/5	0.78 ± 1.16	0.69 ± 1.17	0.69
L5/S	0.13 ± 0.49	0.30 ± 0.85	0.87
Cobb angle	3.66 ± 2.53	1.79 ± 2.03	<0.01
Pelvic tilt (case)	12	33	0.07
Coronal spinal balance	4.03 ± 3.54	3.42 ± 3.50	0.36
Sagittal spinal balance	9.0 ± 16.4	10.1 ± 15.4	0.71
Lumbar lordosis	26.7 ± 14.0	27.1 ± 10.6	0.85

Discussion

A 15-year follow-up study of lumbar radiographic examinations was conducted in a community-based cohort of 194 adults, 40 years of age or older, at the beginning of the study. The cumulative incidence of DNDLS over the 15-year study period was 33/194 (17.0 %) in this cohort. Compared with childhood scoliosis, which affects

approximately 2 % of the population, DNDLS seems to be more common in the elderly population. There was a tendency for female predominance, and the frequency of the disorder increased with age. However, the severity of these curves was relatively low, and none of the curves developed a Cobb angle of greater than 30°, with most in the range 10°–20°.

There are few epidemiological data on DNDLS in the elderly. To our knowledge, only 2 longitudinal studies have been published describing community-based cohorts. Robin et al. reported that 55 of 554 subjects (10 %) developed a spinal deformity during the 7–13-year follow-up. None of the reported curves exceeded 20° for the Cobb angle and most were in the 1°–9° range [20]. Kobayashi et al. reported that 22 of 60 subjects without any scoliosis (36.6 %) developed de-novo scoliosis curves of 10° or more during the follow-up period, with a mean of 12.0 years. None of the scoliosis curvatures exceeded 20° in the Cobb angle, which ranged from 10° to 18°, and the mean Cobb angle for de-novo scoliosis was 13.0° [18]. These data indicate that incidence varies substantially, from 10 to 36.6 %, depending on the study sample. The radiographic features of DNDLS reported in these studies are similar to those reported in our study. Taken together, these studies show that DNDLS is a common condition that is characterized by low-grade curve magnitude and slow progression in the elderly and that the prognosis of the clinical course of DNDLS seems to be benign.

Berner and Ehni [2] reported that asymmetric degeneration of the intervertebral space was the key risk factor for curve progression. Kobayashi et al. also suggested that asymmetric intervertebral disc degeneration leads to the development of scoliosis [18]. By contrast, in our study the presence of an intervertebral disc angle, which indicates asymmetric intervertebral disc degeneration, occurred often in the lower lumbar segments in the S-group, but the disc angle at L4–5 and L5–S1 did not differ significantly between the S and C-groups. This study did not identify asymmetric intervertebral disc degeneration in the lower lumbar segments as a key risk factor for curve progression.

Some authors have proposed that rotatory subluxation is the initial event in the induction of de-novo scoliosis [6, 21]. Our results strongly support this hypothesis, because rotatory deformity and lateral spondylolisthesis of L3 were observed significantly more often in the S-group than in the C-group. Although rotatory subluxation could not be judged from the combination of anteroposterior and lateral radiographs alone, these findings suggest the existence of rotatory subluxation.

We offer the following hypothesis to explain the pathological mechanism responsible for induction of DNDLS in the elderly. Asymmetric anatomical changes in the lower

lumbar segments are generally compensated by the relatively undegenerated upper lumbar segments, so maintaining normal spinal balance in the early stage of spinal degeneration. However, this compensatory mechanism cannot compensate when disc degeneration advances to the upper lumbar segments. As a result, the mechanical stress concentrates at the L3–4 intervertebral disc level, located at the junction between the fixed, unbalanced, lower lumbar segments and the mobile, upper lumbar segments. DNDLS progresses when rotatory subluxation occurs at L3–4, because of decompensation during the advanced stages of spinal degeneration.

The rotatory deformation and translation of the L3 vertebra observed in the standing, anteroposterior lumbar X-rays may indicate the initiation of DNDLS, and rotatory subluxation at L3–4 may be a predisposing factor for curve progression in the elderly. In this study, the inclination angles of all segments of the lumbar spine were significantly larger in the S-group. These findings may also indicate the beginning of decompensation and its effects on spinal balance in the aging spine, although total spinal balance was maintained fairly well at this point.

Much has been written about lumbar scoliosis detected during childhood, and the risk factors for pain and curve progression have been clearly demonstrated by radiographic follow-up studies. A Cobb angle greater than 30°, loss of lumbar lordosis, apical rotation greater than Nash–Moe grade II, lateral listhesis greater than 5 mm, and an intercrestal line through or below the L4–5 disc space are regarded as potential risk factors for curve progression [1, 20, 22–25]. Although none of the participants in our study met these criteria after the 15-year follow-up period, we cannot conclude that DNDLS does not carry a potential risk of progression to a high-grade deformity, on the basis of the results from this longitudinal study. It is possible that some of the participants with DNDLS in this study may progress to a high-grade deformity in the future. It would be interesting to follow these subjects over a longer period to determine the percentage of individuals who develop high-grade curve magnitudes with disabling pain and a collapsing spine.

This study has several limitations. First, the study sample is not large, so the cumulative incidence determined in this study may not be generalized to the Japanese population. Second, follow-up of this cohort was relatively low. Although the main reason for drop-out is presumed to be death, the authors were unable to identify all of the reasons for study drop-out. Therefore, the real cumulative incidence may be different from that reported in this study. Nonetheless, the long-term observations of the progression of new lumbar curves in the elderly are noteworthy. The results obtained in this study are helpful for understanding the etiology and natural history of DNDLS in the elderly,

and further follow-up on this cohort will provide additional useful information.

Conclusion

The radiographic features of DNDLS in this cohort of elderly subjects revealed mild scoliosis with minimal rotatory deformity. Spinal decompensation by the upper lumbar segments to the asymmetric anatomical deformity in the lower lumbar segments may induce de-novo lumbar scoliosis. The rotatory deformity and lateral spondylolisthesis of the L3 vertebra may be a prognostic factor for de-novo lumbar curve progression in the elderly.

Conflict of interest The authors declare that they have no conflict of interest.

References

- Vanderpool DW, James JIP. Scoliosis in the elderly. *J Bone Joint Surg [Am]*. 1969;51:446–55.
- Berner B, Ehni G. Degenerative lumbar scoliosis. *Spine*. 1979;4:548–52.
- Epstein JA, Epstein BS, Jones MD. Symptomatic lumbar scoliosis with degenerative changes in the elderly. *Spine*. 1979;4:542–7.
- Ploumis A, Transfeldt EE, Denis F. Degenerative lumbar scoliosis associated with spinal stenosis. *Spine J*. 2007;7:428–36.
- Berven SH, Lowe T. The scoliosis research society classification for adult spinal deformity. *Neurosurg Clin N Am*. 2007;18:207–13.
- Marty-Poumarat C, Scattin L, Marpeau M, de Loubresse CG, Aegerter P. Natural history of progressive adult scoliosis. *Spine*. 2007;32:1227–34.
- Schwab F, El-Fegoun AB, Gamez L, Goodman H, Farcy JP. A lumbar classification of scoliosis in the adult patient: preliminary approach. *Spine*. 2005;30:1670–3.
- Weinstein SL, Ponseti IV. Curve progression in idiopathic scoliosis. *J Bone Joint Surg [Am]*. 1983;65:447–55.
- Newton PO, Parent S, Wenger DR. Treatment algorithm: observation, bracing, surgery. In: Newton PO, editor. *Adolescent Idiopathic Scoliosis*. Rosemont: American Academy of Orthopedic Surgeons; 2004. pp. 23–8.
- Kasamatsu T, Morioka S, Hashimoto T, Kinoshita H, Yamada H, Tamaki T. Epidemiological study on the bone mineral density of inhabitants in Miyama Village, Wakayama Prefecture (Part I) Background of study population and sampling method. *J Bone Miner Metab*. 1991;9:50–5.
- Kinoshita H, Danjoh S, Yamada H, Tamaki T, Kasamatsu T, Ueda A, Hashimoto T. Epidemiological study on the bone mineral density of inhabitants in Miyama village, Wakayama prefecture (part II) Bone mineral density of the spine and proximal femur. *J Bone Miner Metab*. 1991;9:56–60.
- Yoshimura N. Incidence of fast bone losers and factors affecting changes in bone mineral density: a cohort study in a rural Japanese community. *J Bone Miner Metab*. 1996;14:171–7.
- Muraki S, Oka H, Akune T, Mabuchi A, En-Yo Y, Yoshida M, Saika A, Suzuki T, Yoshida H, Ishibashi H, Yamamoto S, Nakamura K, Kawaguchi H, Yoshimura N. Prevalence of radiographic lumbar spondylosis and its association with low back pain in the elderly of population-based cohorts: the ROAD study. *Ann Rheum Dis*. 2008;68:1401–6.
- Muraki S, Oka H, Akune T, Mabuchi A, En-Yo Y, Yoshida M, Saika A, Suzuki T, Yoshida H, Ishibashi H, Yamamoto S, Nakamura K, Kawaguchi H, Yoshimura N. Prevalence of radiographic knee osteoarthritis and its association with knee pain in the elderly of Japanese population-based cohorts: the ROAD study. *Osteoarthr Cartil*. 2009;17:1137–43.
- Yoshimura N, Muraki S, Oka H, Mabuchi A, En-Yo Y, Yoshida M, Saika A, Suzuki T, Yoshida H, Ishibashi H, Yamamoto S, Nakamura K, Kawaguchi H, Yoshimura N. Prevalence of knee osteoarthritis, lumbar spondylosis and osteoporosis in Japanese men and women: the Research on Osteoarthritis/osteoporosis Against Disability (ROAD). *J Bone Miner Metab*. 2009;27:620–8.
- Yoshimura N, Muraki S, Oka H, Kawaguchi H, Nakamura K, Akune T. Cohort profile: research on osteoarthritis/osteoporosis against disability (ROAD) study. *Int J Epidemiol*. 2010;39:988–95.
- Pedriolle R, Vidal J. Thoracic idiopathic scoliosis: curve evolution and prognosis. *Spine*. 1985;10:758–90.
- Kobayashi T, Atsuta Y, Takemitsu M, Matsuno T, Takeda N. A prospective study of de novo scoliosis in a community based cohort. *Spine*. 2006;31:178–82.
- Kawakami M, Tamaki T, Ando M, Yamada H, Hashizume H, Yoshida M. Lumbar sagittal balance influences the clinical outcome after decompression and posterolateral spinal fusion for degenerative lumbar spondylolisthesis. *Spine*. 2002;27:59–64.
- Robin GC, Span Y, Steinberg R, Makin M, Menczel J. Scoliosis in the elderly: a follow-up study. *Spine*. 1982;7:355–35.
- Korovessis P, Piperos G, Sidiripoulos P, Dimas A. Adult idiopathic lumbar scoliosis. A formula for prediction of progression and review of the literature. *Spine*. 1994;19:1926–32.
- Grubb SA, Lipscomb HJ. Diagnostic findings in painful adult scoliosis. *Spine*. 1992;17:518–27.
- Pritchett JW, Bortel DT. Degenerative symptomatic lumbar scoliosis. *Spine*. 1993;18:700–3.
- Sapkas G, Efstathiou P, Badekas AT, Antoniadis A, Kyrtatzoulis J, Meleteas L. Radiological parameters associated with the evolution of degenerative scoliosis. *Bull Hosp Jt Dis*. 1996;55:40–5.
- Thevenon A, Pollez B, Cantegrit F, Tison-Muchery F, Marchandise X, Duquesnoy B. Relationship between kyphosis, scoliosis, and osteoporosis in the elderly population. *Spine*. 1987;12:744–5.

Incidence and Risk Factors for Radiographic Knee Osteoarthritis and Knee Pain in Japanese Men and Women

A Longitudinal Population-Based Cohort Study

Shigeyuki Muraki,¹ Toru Akune,¹ Hiroyuki Oka,¹ Yuyu Ishimoto,² Keiji Nagata,² Munehito Yoshida,² Fumiaki Tokimura,³ Kozo Nakamura,⁴ Hiroshi Kawaguchi,⁴ and Noriko Yoshimura¹

Objective. To examine the incidence and progression of radiographic knee osteoarthritis (OA) and the incidence of knee pain, and their risk factors in Japan, using the large-scale population of the nationwide cohort study ROAD (Research on Osteoarthritis/osteoporosis Against Disability).

Methods. Subjects from the ROAD study who had been recruited in 2005–2007 were followed up with knee radiography 3 years later. A total of 2,262 paired radiographs (74.4% of the original sample) were scored using the Kellgren/Lawrence (K/L) grading system, and the incidence and progression rate of knee OA was

examined. The incidence rate of knee pain was also examined. In addition, risk factors were tested for their association with incident and progressive radiographic knee OA and incident knee pain.

Results. Given the ~3.3-year followup, the rate of incident K/L grade ≥ 2 radiographic knee OA was 6.9% and 11.9% in men and women, respectively, while that of K/L grade ≥ 3 knee OA was 8.4% and 13.9% in men and women, respectively. The rate of progressive knee OA was 17.8% and 22.3% in men and women, respectively. The incident rate of knee pain was 21.2% and 27.3% in men and women, respectively. Female sex was a risk factor for incident K/L grade ≥ 2 knee OA, but was not associated with incident K/L grade ≥ 3 knee OA or progressive knee OA. Knee pain was a risk factor for incident and progressive knee OA. Previous knee injury was a risk factor for knee pain but not for radiographic knee OA.

Conclusion. The present longitudinal study revealed a high incidence of radiographic knee OA in Japan.

Knee osteoarthritis (OA), characterized by pathologic features including joint space narrowing and osteophytosis, is a major public health issue causing chronic pain and disability in the elderly in most developed countries (1). The prevalence of radiographic knee OA is high in Japan (2), with 25,300,000 subjects ages ≥ 40 years estimated to experience radiographic knee OA (3). According to the recent National Livelihood Survey of the Ministry of Health, Labor, and Welfare in Japan, OA is ranked fourth among diseases that cause

Supported by grants-in-aid for Scientific Research (S19109007, B20390182, C20591737, C20591774), for Young Scientists (A18689031), and for Exploratory Research (19659305) from the Japanese Ministry of Education, Culture, Sports, Science, and Technology; grants H17-Men-eki-009, H18-Chouju-037, H20-Chouju-009, and H21-Chouju-Wakate-011 from the Ministry of Health, Labor, and Welfare; research funding from the Japanese Orthopaedic Association (JOA-Subsidized Science Project Research 2006-1 and 2010-2); and grant 166 from the Japan Orthopaedics and Traumatology Foundation.

¹Shigeyuki Muraki, MD, PhD, Toru Akune, MD, PhD, Hiroyuki Oka, MD, Noriko Yoshimura, MD, PhD: 22nd Century Medical and Research Center and University of Tokyo, Tokyo, Japan; ²Yuyu Ishimoto, MD, Keiji Nagata, MD, Munehito Yoshida, MD, PhD: Wakayama Medical University, Wakayama, Japan; ³Fumiaki Tokimura, MD, PhD: Tokyo Metropolitan Geriatric Medical Center, Tokyo, Japan; ⁴Kozo Nakamura, MD, PhD, Hiroshi Kawaguchi, MD, PhD: University of Tokyo, Tokyo, Japan.

Address correspondence to Shigeyuki Muraki, MD, PhD, Department of Clinical Motor System Medicine, 22nd Century Medical and Research Center, Faculty of Medicine, University of Tokyo, 7-3-1, Hongo, Bunkyo-ku, Tokyo 113-8655, Japan. E-mail: murakis-ort@h.u-tokyo.ac.jp.

Submitted for publication February 8, 2011; accepted in revised form November 22, 2011.

disabilities that subsequently require support with activities of daily living (4). Despite the urgent need of strategies for the prevention and treatment of this condition, there have only been a few studies of rates of incidence and progression of knee OA (5–9). Furthermore, to the best of our knowledge, there has been no study of the incidence or progression of knee OA in Asians, although large differences in prevalence of knee OA exist among the different races (2,10,11).

Previous studies that investigated risk factors for knee OA (12–23) showed that obesity, previous knee injury, female sex, muscle strength, occupational activities, and older age were risk factors for knee OA mainly in Caucasians. However, in addition to the differences in prevalence among different ethnic groups, anthropometric measurements and environmental situations varied substantially in different countries. Thus, findings in Caucasians cannot be applied to different ethnic groups. Few population-based cohort studies have examined risk factors for knee OA in Asians.

The principal clinical symptom of knee OA is knee pain (24). Ours and other previous studies have demonstrated that the impact of knee pain on quality of life is disproportionate to the radiographic changes themselves in the knee (25,26). Several cross-sectional studies have investigated the factors associated with knee pain (25,27), but in most of those studies, radiographic findings were not included in the analysis, although radiographic severity of knee OA is an important factor for knee pain (2). Furthermore, there have been few longitudinal studies of the incidence of knee pain that include radiographic findings in the analysis.

The objective of the present study was to clarify the incidence and progression rate of radiographic knee OA as well as the incidence of knee pain in Japanese men and women using the large-scale, population-based cohort study known as the Research on Osteoarthritis/osteoporosis Against Disability (ROAD). In addition, we examined the risk factors for the incidence and progression of knee OA as well as for the incidence of knee pain.

SUBJECTS AND METHODS

Subjects. The ROAD study was a nationwide prospective study of bone and joint diseases (with osteoarthritis and osteoporosis as the representative bone and joint diseases) constituting population-based cohorts established in several communities in Japan. As a detailed profile of the ROAD study has already been described elsewhere (2,3,28), a brief summary is provided here. In 2005–2007, we created a baseline database that included the clinical and genetic information for

3,040 inhabitants (1,061 men, 1,979 women) in the age range of 23–95 years (mean 70.6 years), recruited from listings of resident registrations in 3 communities: an urban region in Itabashi, Tokyo; a mountainous region in Hidakagawa, Wakayama; and a coastal region in Taiji, Wakayama. Participants in the urban region were recruited from a randomly selected cohort from the Itabashi-ward residents' registration database (29). The participation rate was 75.6%. Participants in mountainous and coastal regions were also recruited from the resident registration lists, and the participation rates in these 2 areas were 56.7% and 31.7%, respectively. The inclusion criteria, apart from residence in the communities mentioned above, were the ability to walk to the survey site, report data, and understand and sign an informed consent form. All participants provided written informed consent, and the study was conducted with the approval of the ethics committees of the University of Tokyo and the Tokyo Metropolitan Geriatric Medical Center.

Participants completed an interviewer-administered questionnaire of 400 items that included lifestyle information such as smoking habit, alcohol consumption, family history, medical history, and previous knee injury history. Anthropometric measurements included height and weight, from which the body mass index (BMI) (weight [kg]/height [m²]) was calculated. Grip strength was measured on bilateral sides using a TOEI LIGHT handgrip dynamometer, and the better measurement was used to represent maximum muscle strength. Furthermore, all participants were interviewed by well-experienced orthopedists regarding pain in both knees, who asked "Have you experienced right knee pain on most days in the past month, in addition to now?" and "Have you experienced left knee pain on most days in the past month, in addition to now?" Subjects who answered "yes" were defined as having knee pain.

In 2008–2010, we attempted to trace and review all 3,040 subjects; they were invited to attend a followup interview and undergo repeat radiography. The interviews, which included questions about current knee pain, were conducted by the same trained orthopedists who undertook the baseline study in 2005–2007. Anthropometric measurements also included height, weight, and grip strength at followup.

Radiographic assessment. All participants underwent radiographic examination of both knees using an anteroposterior view with weight-bearing and foot map positioning. Fluoroscopic guidance with a horizontal anteroposterior x-ray beam was used to properly visualize the joint space. Knee radiographs at baseline and followup were read in pairs without knowledge of the participant's clinical status by a single well-experienced orthopedist (SM), and the Kellgren/Lawrence (K/L) grade was defined using the K/L radiographic atlas for overall knee radiographic grades (30). In the K/L grading system, radiographs are scored from grade 0 to grade 4, with the higher grades being associated with more severe OA. To evaluate the intraobserver variability of the K/L grading, 100 randomly selected radiographs of the knee were scored by the same observer >1 month after the first reading. One hundred other radiographs were also scored by 2 experienced orthopedic surgeons (SM and HO) using the same atlas for interobserver variability. The intra- and intervariabilities evaluated for K/L grade (0–4) were confirmed by kappa

Table 1. Baseline characteristics of the participants*

	Overall (n = 2,262)	Men (n = 763)	Women (n = 1,499)
Age, years	68.7 ± 11.3	69.6 ± 11.1	68.2 ± 11.4†
Height, cm	154.6 ± 8.8	163.1 ± 6.6	150.3 ± 6.4†
Weight, kg	55.4 ± 10.2	62.2 ± 9.9	51.9 ± 8.4†
BMI, kg/m ²	23.1 ± 3.3	23.3 ± 3.0	23.0 ± 3.4†
Grip strength, kg	26.2 ± 9.2	34.3 ± 8.8	22.1 ± 6.2†
OA prevalence, no. (%)			
K/L grade ≥1	1,898 (83.9)	586 (76.8)	1,312 (87.5)‡
K/L grade ≥2	1,164 (51.5)	296 (38.8)	868 (57.9)‡
K/L grade ≥3	355 (15.7)	75 (9.8)	280 (18.7)‡
Knee pain, no. (%)	478 (21.1)	111 (14.6)	367 (24.5)‡
Previous knee injury, no. (%)	217 (9.6)	52 (6.8)	165 (11.0)‡
Smoking, no. (%)	212 (9.4)	165 (21.6)	47 (3.1)‡
Alcohol, no. (%)	831 (36.7)	483 (63.3)	348 (23.2)‡

* Except where indicated otherwise, values are the mean ± SD. BMI = body mass index; OA = osteoarthritis; K/L = Kellgren/Lawrence.

† P < 0.05 versus men, by Student's unpaired t-test.

‡ P < 0.05 versus men, by chi-square test.

analysis to be sufficient for assessment ($\kappa = 0.86$ and $\kappa = 0.80$, respectively).

For the purposes of this study, we defined 4 knee OA outcomes. First, a subject was defined as having incident K/L grade ≥1 radiographic knee OA if both knees had less than grade 1 disease at baseline and if at least one knee had grade 1 or higher disease at followup. Second, a subject was defined as having incident K/L grade ≥2 radiographic knee OA if both knees had less than grade 2 disease at baseline and if at least one knee had grade 2 or higher disease at followup. Third, incident K/L grade ≥3 radiographic knee OA was defined as less than grade 3 disease in both knees at baseline and grade 3 or higher disease in at least one knee at followup. Fourth, progressive knee OA was defined as K/L grade 2 and K/L grade 3 knee OA at baseline (because K/L grade 4 knee OA cannot progress) and an increase by at least 1 grade in the affected knee at followup.

Statistical analysis. Odds ratios (ORs) and 95% confidence intervals (95% CIs) are provided. The differences in age, height, weight, BMI, and grip strength between men and women were examined using Student's unpaired t-test. To compare the prevalence of radiographic knee OA and knee pain between men and women, we performed the chi-square test. A logistic regression analysis was used to determine the association of incident radiographic knee OA, progressive knee OA, and incident knee pain with age in men and women. To determine risk factors for incident knee OA and progressive knee OA, a univariate generalized estimating equation (GEE) logistic regression analysis was used. Furthermore, to determine independent risk factors, multiple GEE logistic regression analysis was used with significant risk factors in a univariate GEE logistic regression analysis model as independent variables. Incident knee pain was defined as no knee pain in both knees at baseline and knee pain in at least one knee at followup. To determine risk factors for incident knee pain, a univariate GEE logistic regression analysis was used. Furthermore, to determine independent risk factors, multiple GEE

logistic regression analysis was used with significant risk factors in a univariate logistic regression analysis model in addition to regions as independent variables. When we analyzed risk factors for incident knee OA, progressive knee OA, and incident knee pain, K/L grade at baseline, knee pain at baseline, and previous knee injury were defined for the knee which had the incident OA, progressive OA, and incident pain, respectively. Data analyses were performed using SAS software, version 9.0 (SAS Institute).




Figure 1. Number (N) of male and female subjects by Kellgren/Lawrence (K/L) grade at baseline and followup.

Table 2. Incidence of radiographic knee OA, progressive knee OA, and knee pain according to sex*

	K/L grade ≥ 1		K/L grade ≥ 2		K/L grade ≥ 3		Progressive knee OA		Knee pain	
	No. at risk	Cumulative incidence, no. (%)	No. at risk	Cumulative incidence, no. (%)	No. at risk	Cumulative incidence, no. (%)	No. at risk	Cumulative incidence, no. (%)	No. at risk	Cumulative incidence, no. (%)
Overall	364	70 (19.2)	1,098	107 (9.7)	1,907	228 (12.0)	1,084	229 (21.1)	1,784	447 (25.1)
Men	177	35 (19.8)	467	32 (6.9)	688	58 (8.4)	276	49 (17.8)	652	138 (21.2)
Women	187	35 (18.7)	631	75 (11.9) [†]	1,219	169 (13.9) [†]	808	180 (22.3)	1,132	309 (27.3) [†]

* See Table 1 for definitions.

[†] $P < 0.05$ versus men, by chi-square test.

RESULTS

Of the 3,040 subjects in the baseline study in 2005–2007, 125 (4.1%) had died by the time of the review 3 years later, 123 (4.0%) did not participate in the followup study due to bad health, 69 (2.3%) had moved away, 83 (2.7%) declined the invitation to attend the followup study, and 155 (5.1%) did not participate in the followup study for other reasons. Among the 2,485 subjects who did participate in the followup study, 175 (5.8%) did not undergo plain radiography and 18 (0.6%) provided incomplete pain questionnaires; these were excluded. We also excluded 30 subjects (1.0%) who underwent total knee arthroplasty before the followup study, leaving a total of 2,262 subjects (74.4%) (763 men and 1,499 women) from whom paired radiographs and complete pain histories were obtained. Their mean \pm SD age at followup was 72.2 ± 11.4 years. The mean \pm SD duration of followup between initial and second radiographs was 3.3 ± 0.6 years. Those participating in the followup study were younger than those who did not survive or who did not participate for other reasons (mean age 68.6 years for responders versus 75.1 years for nonresponders; $P < 0.0001$). The followup study participants also were more likely to be women (66.3% of responders were women and 61.8% of nonresponders were women; $P = 0.03$) and were less likely to have knee OA at the baseline examination than either those who did not survive to followup or those who did not participate for other reasons (51.5% of responders versus 60.9% of nonresponders; $P < 0.0001$).

The characteristics of the 2,262 participants at baseline in the ROAD study are shown in Table 1. Men were significantly older than women, while BMI was higher in men than in women. The prevalence of knee OA and knee pain was significantly higher in women than in men at baseline. The proportion of previous knee injuries was also higher in women than in men. The number of subjects with each K/L grade at baseline and at followup is shown in Figure 1.


**Figure 2.** Percentage of subjects with incident radiographic knee osteoarthritis (OA) (Kellgren/Lawrence [K/L] grade ≥ 1 , ≥ 2 , or ≥ 3), progressive knee OA, and incident knee pain in each age stratum (<50 years, 50–59 years, 60–69 years, 70–79 years, and ≥ 80 years).

Table 3. Risk factors for incident radiographic knee osteoarthritis*

	K/L grade ≥ 2			K/L grade ≥ 3		
	No./total no. (%) of subjects	Crude OR (95% CI)	Adjusted OR (95% CI)	No./total no. (%) of subjects	Crude OR (95% CI)	Adjusted OR (95% CI)
Age (+5 years)	-	1.26 (1.15-1.39)	1.31 (1.15-1.49)	-	1.33 (1.23-1.44)	1.25 (1.13-1.39)
BMI (+5 kg/m ²)	-	2.00 (1.49-2.69)	2.43 (1.76-3.39)	-	1.67 (1.36-2.04)	1.68 (1.35-2.11)
Grip strength (+1 kg)	-	0.96 (0.94-0.98)	1.01 (0.97-1.04)	-	0.95 (0.94-0.97)	1.00 (0.97-1.02)
Sex						
Men	32/467 (6.9)	Referent	Referent	58/688 (8.4)	Referent	Referent
Women	75/631 (11.9)	1.83 (1.20-2.86)	2.76 (1.50-5.18)	169/1,219 (13.9)	1.75 (1.28-2.41)	1.42 (0.88-2.29)
K/L grade at baseline						
0	-	Referent	Referent	-	Referent	Referent
1	-	4.11 (2.33-7.83)	2.48 (1.35-4.87)	-	1.91 (0.69-5.43)	1.29 (0.45-3.80)
2	-	-	-	-	5.69 (2.38-14.30)	5.94 (1.07-35.83)
Knee pain at baseline						
No	-	Referent	-	-	Referent	Referent
Yes	-	0.91 (0.32-2.24)	-	-	3.77 (2.44-5.73)	2.53 (1.59-4.00)
Previous knee injury						
No	-	Referent	-	-	Referent	-
Yes	-	4.08 (0.66-18.8)	-	-	1.24 (0.45-3.11)	-
Smoking						
No	99/958 (10.3)	Referent	-	213/1,713 (12.4)	Referent	Referent
Yes	8/140 (5.7)	0.53 (0.23-1.04)	-	14/194 (7.2)	0.55 (0.30-0.93)	1.07 (0.55-1.94)
Alcohol use						
No	68/627 (10.9)	Referent	-	158/1,171 (13.5)	Referent	Referent
Yes	39/471 (8.3)	0.74 (0.49-1.12)	-	69/736 (9.4)	0.66 (0.49-0.89)	0.96 (0.67-1.36)

* Adjusted odds ratios (ORs) were calculated by multiple generalized estimating equation logistic regression analysis after adjustment for all other variables in addition to regions. We show all variables we analyzed in the present study. K/L = Kellgren/Lawrence; 95% CI = 95% confidence interval; BMI = body mass index.

Table 2 shows the rates of incident and progressive knee OA and incident knee pain in the overall population and subgroups classified by sex. The incidences of K/L grade ≥ 2 and K/L grade ≥ 3 knee OA and knee pain were significantly higher in women than in men, while there were no significant differences in the incidence of K/L grade ≥ 1 knee OA and the progression of knee OA between men and women. The incidence and progression rate of knee OA tended to increase with age in men and women (for 5-year increase: K/L grade ≥ 1 , OR 1.22 [95% CI 1.06-1.43] and OR 1.52 [95% CI 1.29-1.84], respectively; K/L grade ≥ 2 , OR 1.35 [95% CI 1.12-1.67] and OR 1.29 [95% CI 1.15-1.45], respectively; K/L grade ≥ 3 , OR 1.34 [95% CI 1.15-1.58] and OR 1.35 [95% CI 1.24-1.49], respectively; progressive knee OA, OR 1.15 [95% CI 0.95-1.42] and OR 1.15 [95% CI 1.0-1.28], respectively) (Figure 2). Interestingly, the incidence rate of knee pain was age-dependent in women (OR 1.10 [95% CI 1.04-1.17]), while it was not age-dependent in men (OR 0.97 [95% CI 0.90-1.06]). Furthermore, in subjects age < 60 years, the incidence of knee pain was similar between women and men (OR 1.12 [95% CI 0.88-1.42]), while in subjects age > 60 years, the incidence of knee pain was significantly higher

in women than in men (OR 0.78 [95% CI 0.68-0.88]) (Figure 2).

Table 3 shows the baseline risk factors for incident radiographic knee OA. Univariate logistic regression analysis showed that age, BMI, grip strength, sex, and K/L grade were associated with incident K/L grade ≥ 2 knee OA. Age, BMI, grip strength, sex, K/L grade, knee pain at baseline, previous knee injury, smoking, and alcohol consumption were associated with incident K/L grade ≥ 3 knee OA. We then determined independent risk factors using a multiple logistic regression analysis that included the above significant factors in the univariate model in addition to regions as independent variables. The results showed that age and BMI were risk factors for incident K/L grade ≥ 2 and incident K/L grade ≥ 3 knee OA. Female sex was a risk factor for incident K/L grade ≥ 2 knee OA, while being female was not significantly associated with incident K/L grade ≥ 3 knee OA. A more severe K/L grade at baseline was strongly associated with incident K/L grade ≥ 2 and incident K/L grade ≥ 3 knee OA. Knee pain at baseline was significantly associated with incident K/L grade ≥ 3 knee OA.

Univariate logistic regression analysis showed

Table 4. Risk factors for progressive knee OA and incident knee pain*

	Progressive knee OA			Knee pain		
	No./total no. (%) of subjects	Crude OR (95% CI)	Adjusted OR (95% CI)	No./total no. (%) of subjects	Crude OR (95% CI)	Adjusted OR (95% CI)
Age (+5 years)	–	1.14 (1.04–1.25)	1.17 (1.05–1.30)	–	1.05 (1.01–1.10)	1.01 (0.95–1.07)
BMI (+5 kg/m ²)	–	1.47 (1.20–1.80)	1.43 (1.16–1.77)	–	1.60 (1.37–1.88)	1.54 (1.30–1.82)
Grip strength (+1 kg)	–	0.98 (0.96–1.00)	0.99 (0.96–1.01)	–	0.98 (0.97–1.00)	1.00 (0.98–1.02)
Sex						
Men	49/276 (17.8)	Referent	–	138/652 (21.2)	Referent	Referent
Women	180/808 (22.3)	1.33 (0.94–1.90)	–	309/1,132 (27.3)	1.40 (1.11–1.76)	1.32 (0.94–1.84)
K/L grade at baseline						
0 and 1	–	–	–	–	Referent	Referent
2	–	–	–	–	1.89 (0.80–4.49)	1.58 (0.65–3.85)
3 and 4	–	–	–	–	3.17 (1.95–5.17)	2.54 (1.52–4.24)
Knee pain at baseline						
No	–	Referent	Referent	–	–	–
Yes	–	2.87 (1.99–4.14)	2.63 (1.81–3.81)	–	–	–
Previous knee injury						
No	–	Referent	–	–	Referent	Referent
Yes	–	0.79 (0.31–1.86)	–	–	3.09 (1.34–7.23)	2.91 (1.26–6.82)
Smoking						
No	219/1,016 (21.6)	Referent	–	411/1,603 (25.6)	Referent	–
Yes	10/68 (14.7)	0.63 (0.30–1.19)	–	36/181 (19.9)	0.72 (0.49–1.04)	–
Alcohol use						
No	168/746 (22.4)	Referent	–	281/1,093 (25.7)	Referent	–
Yes	61/338 (18.1)	0.76 (0.54–1.04)	–	166/691 (24.0)	0.91 (0.73–1.14)	–

* Adjusted ORs were calculated by multiple generalized estimating equation logistic regression analysis after adjustment for all other variables in addition to regions. We show all variables we analyzed in the present study. OA = osteoarthritis (see Table 3 for other definitions).

that age, BMI, grip strength, and knee pain at baseline were associated with progressive knee OA. We then included age, BMI, grip strength, and knee pain at baseline in addition to regions as independent variables in a multiple logistic regression analysis to determine independent risk factors (Table 4). Age and BMI at baseline were risk factors, but their adjusted ORs for progressive knee OA were lower than those for incident K/L grade ≥ 2 knee OA (Table 4). Knee pain was significantly associated with progressive knee OA.

We further investigated risk factors for incident knee pain (Table 4). Univariate logistic regression analysis showed that age, BMI, grip strength, sex, K/L grade, and previous knee injury were associated with incident knee pain. To determine independent risk factors for knee pain, multiple logistic regression analysis was used with age, BMI, grip strength, sex, K/L grade, and previous knee injury in addition to regions as independent variables. BMI was significantly associated with incident knee pain, but female sex was not associated with incident knee pain. Subjects with K/L grade ≥ 3 knee OA at baseline had an ~ 2.5 -fold increased risk for incidence of knee pain compared with K/L grade 0 and K/L grade 1 knees. Previous knee injury was also significantly associated with incident knee pain.

DISCUSSION

This is the first population-based study to examine the incidence and progression of knee OA and risk factors for incident and progressive knee OA among Japanese men and women. We also examined the incident rate of knee pain and its risk factors. The present study showed high rates of incident knee OA, progressive knee OA, and incident knee pain.

Few population-based studies have examined incident radiographic knee OA (6–9). In the Framingham Osteoarthritis Study (6), given the ~ 8.1 -year followup, the incident rate of K/L grade ≥ 2 knee OA was 11.1% and 18.1% (1.4% and 2.2% per year) in Caucasian men and women, respectively. A population-based study in the UK (18) showed that given the ~ 5.1 -year followup, the incident rate of K/L grade ≥ 2 knee OA was 18.5% (2.3% per year), but men and women were not separated in the analysis. In the present study, the incidence of K/L grade ≥ 2 OA was 2.0% and 3.7% per year in men and women, respectively, which is a little higher than that in previous epidemiologic studies in the US and Europe (6,8), implying that the incidence is higher among Japanese than in Caucasians. This is compatible with our findings regarding prevalence of K/L grade ≥ 2 knee OA

in our previous study (2), which showed that the prevalence of K/L grade ≥ 2 knee OA was much higher in Japanese people than in Caucasians (10,11).

For incident K/L grade ≥ 3 knee OA, to the best of our knowledge no population-based studies have been previously reported. In the Chingford Study (7), knee OA was not defined according to K/L grade but according to osteophytosis and joint space narrowing. The Chingford Study showed that given the ~ 4 -year followup, the incidence of joint space narrowing was 12.6% (3.2% per year) in women, which may be comparable to our results for incident K/L grade ≥ 3 knee OA, considering the K/L grade definition; however, a closer comparison provides quite limited accuracy.

In the present study, the incident rate of K/L grade ≥ 3 knee OA was 4.1% per year in Japanese women, which was also a little higher than that seen in Caucasian women. However, this higher incident rate of K/L grade ≥ 3 may be partly explained by the definition of K/L grade ≥ 3 knee OA, because by considering any knees that start at K/L grade < 3 as eligible for this outcome, we combined incident (e.g., knees starting at K/L grade 0–1) and progressive (knees starting at K/L grade 2) disease. In the present study, we also examined progression of knee OA, and we found that the progression rate of knee OA was 5.2% and 6.3% per year in Japanese men and women, respectively, which was also higher than that in other studies in the US and the UK (2.2–3.9%) (6,8). The higher incidence of radiographic knee OA in Japan could also be due to lifestyle factors, because the traditional Japanese lifestyle includes sitting on the heels on a mat and using Japanese-style lavatories, requiring squatting and kneeling, which are associated with knee OA (31–33). These positions may cause mechanical stress to the knee joint and possibly lead to the acceleration of OA.

Although the rate of incident radiographic knee OA and progressive knee OA increased with age in both sexes, that of knee pain was age-dependent in women but not in men. This may be due to the fact that elderly men generally retire from their occupations at age ~ 60 –70 years, and thus the load on the knees may be lighter in men age > 60 years compared with those age < 60 years, whereas women must often continue to do household chores even after age 70 years, and thus the load on the knees may not be lighter in women age > 70 years compared with those age < 70 years.

The present study also showed that age and BMI are risk factors for incident radiographic knee OA, consistent with findings of previous epidemiologic studies (5,7,8). Previous studies have shown that obesity is a

strong risk factor for incident knee OA (34), possibly due to the accumulation of mechanical stress on the knee joint. More severe K/L grade was also a risk factor for incident radiographic knee OA in the present study, which is also consistent with findings of previous studies (7,8). Female sex was also a strong risk factor for incident K/L grade ≥ 2 knee OA, as in previous studies (6,8), possibly implicating the involvement of muscle strength to compensate for mechanical stress, as women are known to have less muscle strength than men in all decades of life (35). However, female sex was not a significant risk factor for incident K/L grade ≥ 3 knee OA or progressive knee OA. Furthermore, age and BMI at baseline were risk factors for progressive knee OA, but their ORs for progressive knee OA were lower than those for incident K/L grade ≥ 2 knee OA.

This discordance between the determinants of incidence of K/L grade ≥ 2 and K/L grade ≥ 3 knee OA or between those of incidence and progression of knee OA using K/L grade suggests that different mechanisms might influence the initiation of osteophytosis (the principal abnormality in K/L grade 2 disease) and joint space narrowing (the principal abnormality in K/L grade 3 disease). However, since K/L grade was defined by a categorical method, which is comparably insensitive to change, this discordance might simply be a function of the scoring system. Nevertheless, there is also accumulating evidence from previous studies that osteophytosis and joint space narrowing have distinct etiologic mechanisms. A recent cross-sectional study has shown that osteophytosis is unrelated not only to joint space narrowing on plain radiographs, but also to cartilage loss measured by quantitative magnetic resonance imaging (36). Furthermore, our study of an experimental mouse model of OA has identified a cartilage-specific molecule, carminerin, that regulates osteophytosis without affecting joint cartilage destruction during OA progression (37,38). Our most recent findings have implications for our understanding of the pathogenesis of knee OA, as well as for preventive strategies.

In the present study, knee pain was a risk factor for incident K/L grade ≥ 3 knee OA and progressive knee OA. Subjects with knee pain may tend not to go out or exercise because of the pain, which may lead to lower quadriceps strength. This may be one of the reasons why knee pain is a risk factor for incident and progressive knee OA, as quadriceps weakness has been previously associated with radiographic knee OA (39).

For incidence of knee pain, age was not a risk factor after adjustment for BMI, sex, and K/L grade at baseline. Knee pain occurrence may be mainly due to

environmental factors rather than individual factors. As described above, elderly men generally retire from their occupations at ages 60–70 years, and thus the load on the knees may be lighter in men age <60 years compared with those age >60 years, which may partly explain the lack of significant association between age and incidence of knee pain. BMI was a risk factor for incident knee pain even after adjustment for K/L grade at baseline, indicating that obesity is a strong risk factor not only for incident radiographic knee OA but also for incident knee pain. In addition, knee OA at baseline was a risk factor for knee pain, but the ORs for knee pain of K/L grade 2 knee OA and K/L grades 3 and 4 knee OA were just 1.6 and 2.5, respectively. In fact, the proportion of subjects with knee pain of those with K/L grade 2 knee OA and K/L grades 3 and 4 knee OA was just 28.0% and 47.1%, respectively, indicating that ~70% of subjects with K/L grade 2 knee OA who had no knee pain at baseline and ~50% of subjects with K/L grades 3 and 4 knee OA who had no knee pain at baseline also had no knee pain after 3 years.

Previous cross-sectional studies have also demonstrated that correlation of knee pain with radiographic severity of knee OA is not as strong as one would expect (2,40–42), most likely because knee pain may arise from a variety of structures other than joint cartilage, such as the menisci, synovium, ligaments, bursae, bone, and bone marrow (43–47). Hence, comprehensive mechanistic studies of knee pain taking various tissues in and around the knee joint into consideration will be needed to elucidate the relationship between radiographic OA and symptomatic OA.

We were unable to detect an association between knee injury and incident OA in the present study. Other cross-sectional studies of OA prevalence have observed strong association with previous knee injury (48), while the incidence data from the Zoetermeer Study, Framingham Study, and Chingford Study (5–7) also showed a slight increase in risk with interim knee injury but were based on small numbers; no significant association with past knee injury was seen in those groups. In the present study, K/L grade ≥ 2 knee OA in subjects with previous knee injury was not significantly associated with previous knee injury, which suggests that the association of incident radiographic knee OA with previous knee injury may be weak, although the number of subjects with incident K/L grade ≥ 2 knee OA who had previous knee injury was just 12. Thus, the small number may partly explain the lack of statistical significance. The present study showed that previous knee injury is a risk factor for incident knee pain. As mentioned above, the correlation

of knee pain with radiographic severity of knee OA is not as strong as expected (2,40–42), as knee pain may arise from a variety of structures other than joint cartilage, such as menisci, synovium, ligaments, bursae, bone, and the bone marrow (43–47), and these tissues may have been damaged by a previous knee injury, which may lead to the incident knee pain.

We were unable to detect an association between smoking/drinking alcohol and incident knee OA or knee pain. The association between smoking and incident knee OA is controversial. The Zoetermeer Study showed that smoking has no association with incident knee OA (5), while incidence data from the Framingham Study showed that smoking protects against incident knee OA (49). We were also unable to show any effect of physical activity in this incidence study. However, the numbers and power were too low to examine this group and to confirm or exclude such effects on incidence.

The present study has several limitations. First, the radiographic investigators did not have readers calibrate their readings to those from other studies. Although we reported a higher incidence of radiographic knee OA than in previous studies, differences in radiographic acquisition, scoring techniques, and methodology across studies limit strict comparisons between our results and previous reports. Differences across studies in the thresholds used by readers to define osteophytes may have had a substantial impact on their incidence. The high incidence of knee OA in our study compared to that in other populations may be due to such differences. Second, our analysis did not include patellofemoral joint radiographs, which would likely increase the concordance between radiographic knee OA and its pain. Third, we defined knee pain as present or absent, rather than as a continuous measure such as the Western Ontario and McMaster Universities Osteoarthritis Index (50) or visual analog scale score. Categorical methods are statistically less powerful than continuous methods. Thus, the association between knee pain and other variables might have been underestimated in the present study.

In conclusion, the present longitudinal study, using a large-scale population from the ROAD study, revealed a high incidence of radiographic knee OA in Japan compared with previous studies. Age, BMI, and female sex influence incidence more than radiographic progression of knee OA, indicating that different mechanisms might influence the initiation of osteophytosis and joint space narrowing. Knee pain was a risk factor for radiographic knee OA. Knee injury was not signifi-

cantly associated with radiographic knee OA, but was a risk factor for incident knee pain. Further progress, along with continued longitudinal surveys within the ROAD study, will elucidate the environmental and genetic backgrounds of knee OA.

ACKNOWLEDGMENTS

The authors thank Drs. Seizo Yamamoto, Hideaki Ishibashi, and Yorito Anamizu and members of the Department of Orthopedics, and Mr. Yuji Kutsuma and other members of the Department of Radiology at Tokyo Metropolitan Geriatric Medical Center; Mrs. Tomoko Takijiri and other members of the Public Office in Hidakagawa, Japan; and Mrs. Tamako Tsutsumi, Mrs. Kanami Maeda, and other members of the Public Office in Taiji, Japan, for their assistance in the location and scheduling of participants for examinations.

AUTHOR CONTRIBUTIONS

All authors were involved in drafting the article or revising it critically for important intellectual content, and all authors approved the final version to be published. Dr. Muraki had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study conception and design. Muraki, Akune, Oka, Ishimoto, Nagata, Yoshida, Tokimura, Nakamura, Kawaguchi, Yoshimura.

Acquisition of data. Muraki, Akune, Oka, Ishimoto, Nagata, Yoshimura.

Analysis and interpretation of data. Muraki, Akune, Oka, Yoshimura.

REFERENCES

- Sharma L, Kapoor D. Epidemiology of osteoarthritis. In: Moskowitz RW, Altman RD, Hochberg MC, Buckwalter JA, Goldberg VM, editors. *Osteoarthritis: diagnosis and medical/surgical management*. 4th ed. Philadelphia: Lippincott Williams & Wilkins; 2007. p. 3–26.
- Muraki S, Oka H, Akune T, Mabuchi A, En-yo Y, Yoshida M, et al. Prevalence of radiographic knee osteoarthritis and its association with knee pain in the elderly of Japanese population-based cohorts: the ROAD study. *Osteoarthritis Cartilage* 2009;17:1137–43.
- Yoshimura N, Muraki S, Oka H, Mabuchi A, En-yo Y, Yoshida M, et al. Prevalence of knee osteoarthritis, lumbar spondylosis and osteoporosis in Japanese men and women: the Research on Osteoarthritis/osteoporosis Against Disability study. *J Bone Miner Metab* 2009;27:620–8.
- Ministry of Health, Labor, and Welfare. The outline of the results of National Livelihood Survey 2007. URL: <http://www.mhlw.go.jp/toukei/list/20-19-1.html>. In Japanese.
- Schouten JS. A 12 year follow up study on osteoarthritis of the knee in the general population [thesis]. Rotterdam: Erasmus University Medical School; 1991.
- Felson DT, Zhang Y, Hannan MT, Naimark A, Weissman BN, Aliabadi P, et al. The incidence and natural history of knee osteoarthritis in the elderly: the Framingham Osteoarthritis Study. *Arthritis Rheum* 1995;38:1500–5.
- Hart DJ, Doyle DV, Spector TD. Incidence and risk factors for radiographic knee osteoarthritis in middle-aged women: the Chingford Study. *Arthritis Rheum* 1999;42:17–24.
- Cooper C, Snow S, McAlindon TE, Kellingray S, Stuart B, Coggon D, et al. Risk factors for the incidence and progression of radiographic knee osteoarthritis. *Arthritis Rheum* 2000;43:995–1000.
- Lohmander LS, Gerhardsson de Verdier M, Roloff J, Nilsson PM, Engstrom G. Incidence of severe knee and hip osteoarthritis in relation to different measures of body mass: a population-based prospective cohort study. *Ann Rheum Dis* 2009;68:490–6.
- Jordan JM, Helmick CG, Renner JB, Luta G, Dragomir AD, Woodard J, et al. Prevalence of knee symptoms and radiographic and symptomatic knee osteoarthritis in African Americans and Caucasians: the Johnston County Osteoarthritis Project. *J Rheumatol* 2007;34:172–80.
- Zhang Y, Xu L, Nevitt MC, Aliabadi P, Yu W, Qin M, et al. Comparison of the prevalence of knee osteoarthritis between the elderly Chinese population in Beijing and whites in the United States: the Beijing Osteoarthritis Study. *Arthritis Rheum* 2001;44:2065–71.
- Davis MA, Ettinger WH, Neuhaus JN, Cho SA, Hauck WW. The association of knee injury and obesity with unilateral and bilateral osteoarthritis of the knee. *Am J Epidemiol* 1989;130:278–88.
- Spector TD. The fat on the joint: osteoarthritis and obesity. *J Rheumatol* 1990;17:283–4.
- Cooper C, McAlindon T, Snow S, Vines K, Young P, Kirwan J, et al. Mechanical and constitutional risk factors for symptomatic knee osteoarthritis: differences between medial, tibiofemoral and patellofemoral disease. *J Rheumatol* 1994;21:307–13.
- Hart DJ, Doyle DV, Spector TD. The association between metabolic factors and knee osteoarthritis in women: the Chingford Study. *J Rheumatol* 1995;22:1118–23.
- Felson DT, Zhang Y, Hannan MT, Naimark A, Weissman B, Aliabadi P, et al. Risk factors for incident radiographic knee osteoarthritis in the elderly: the Framingham Study. *Arthritis Rheum* 1997;40:728–33.
- Slemenda C, Heilman DK, Brandt KD, Katz BP, Mazucca SA, Braunstein EM, et al. Reduced quadriceps strength relative to body weight: a risk factor for knee osteoarthritis in women? *Arthritis Rheum* 1998;41:1951–9.
- Ledingham J, Regan M, Jones A, Doherty M. Factors affecting radiographic progression of knee osteoarthritis. *Ann Rheum Dis* 1995;54:53–8.
- McAlindon TE, Wilson PW, Aliabadi P, Weissman B, Felson DT. Level of physical activity and the risk of radiographic and symptomatic knee osteoarthritis in the elderly: the Framingham Study. *Am J Med* 1999;106:151–7.
- Wilder FV, Hall BJ, Barrett JP Jr, Lemrow NB. History of acute knee injury and osteoarthritis of the knee: a prospective epidemiological assessment. The Clearwater Osteoarthritis Study. *Osteoarthritis Cartilage* 2002;10:611–6.
- Lachance L, Sowers MF, Jamadar D, Hochberg M. The natural history of emergent osteoarthritis of the knee in women. *Osteoarthritis Cartilage* 2002;10:849–54.
- Reijman M, Pols HA, Bergink AP, Hazes JM, Belo JN, Lievens A, et al. Body mass index associated with onset and progression of osteoarthritis of the knee but not of the hip: the Rotterdam Study. *Ann Rheum Dis* 2007;66:158–62.
- Blagojevic M, Jinks C, Jeffery A, Jordan KP. Risk factors for onset of osteoarthritis of the knee in older adults: a systematic review and meta-analysis. *Osteoarthritis Cartilage* 2010;18:24–33.
- Linaker CH, Walker-Bone K, Palmer K, Cooper C. Frequency and impact of regional musculoskeletal disorders. *Baillieres Clin Rheumatol* 1999;13:197–215.
- Jordan JM, Luta G, Renner JB, Linder GF, Dragomir A, Hochberg MC, et al. Self-reported functional status in osteoarthritis of the knee in a rural southern community: the role of sociodemographic factors, obesity, and knee pain. *Arthritis Care Res* 1996;9:273–8.

26. Muraki S, Akune T, Oka H, En-yo Y, Yoshida M, Saika A, et al. Association of radiographic and symptomatic knee osteoarthritis with health-related quality of life in a population-based cohort study in Japan: the ROAD study. *Osteoarthritis Cartilage* 2010; 18:1227-34.
27. Bergenudd H, Nilsson B, Lindgarde F. Knee pain in middle age and its relationship to occupational work load and psychosocial factors. *Clin Orthop Relat Res* 1989;245:210-5.
28. Yoshimura N, Muraki S, Oka H, Kawaguchi H, Nakamura K, Akune T. Cohort profile: Research on Osteoarthritis/osteoporosis Against Disability (ROAD) study. *Int J Epidemiol* 2010;39: 988-95.
29. Shimada H, Lord SR, Yoshida H, Kim H, Suzuki T. Predictors of cessation of regular leisure-time physical activity in community-dwelling elderly people. *Gerontology* 2007;53:293-7.
30. Kellgren JH, Lawrence JS, editors. *The epidemiology of chronic rheumatism: atlas of standard radiographs of arthritis*. Oxford: Blackwell Scientific; 1963.
31. Felson DT, Hannan MT, Naimark A, Berkeley J, Gordon G, Wilson PW, et al. Occupational physical demands, knee bending, and knee osteoarthritis: results from the Framingham Study. *J Rheumatol* 1991;18:1587-92.
32. Cooper C, McAlindon T, Coggon D, Egger P, Dieppe P. Occupational activity and osteoarthritis of the knee. *Ann Rheum Dis* 1994;53:90-3.
33. Muraki S, Akune T, Oka H, Mabuchi A, En-yo Y, Yoshida M, et al. Association of occupational activity with radiographic knee osteoarthritis and lumbar spondylosis in elderly patients of population-based cohorts: a large-scale population-based study. *Arthritis Rheum* 2009;61:779-86.
34. Altman RD, Fries JF, Bloch DA, Carstens J, Cooke TD, Genant H, et al. Radiographic assessment of progression in osteoarthritis. *Arthritis Rheum* 1987;30:1214-25.
35. Sinaki M, Nwaogwugwu NC, Phillips BE, Mokri MP. Effect of gender, age, and anthropometry on axial and appendicular muscle strength. *Am J Phys Med Rehabil* 2001;80:330-8.
36. Jones G, Ding C, Scott F, Glisson M, Cicuttini F. Early radiographic osteoarthritis is associated with substantial changes in cartilage volume and tibial bone surface area in both males and females. *Osteoarthritis Cartilage* 2004;12:169-74.
37. Yamada T, Kawano H, Koshizuka Y, Fukuda T, Yoshimura K, Kamekura S, et al. Carminerin contributes to chondrocyte calcification during endochondral ossification. *Nat Med* 2006;12:665-70.
38. Kamekura S, Kawasaki Y, Hoshi K, Shimoaka T, Chikuda H, Maruyama Z, et al. Contribution of runt-related transcription factor 2 to the pathogenesis of osteoarthritis in mice after induction of knee joint instability. *Arthritis Rheum* 2006;54:2462-70.
39. Slemenda C, Brandt KD, Heilman DK, Mazucca S, Braunstein EM, Katz BP, et al. Quadriceps weakness and osteoarthritis of the knee. *Ann Intern Med* 1997;127:97-104.
40. Summers MN, Haley WE, Reveille JD, Alarcon GS. Radiographic assessment and psychologic variables as predictors of pain and functional impairment in osteoarthritis of the knee or hip. *Arthritis Rheum* 1988;31:204-9.
41. Cicuttini FM, Baker J, Hart DJ, Spector TD. Association of pain with radiological changes in different compartments and views of the knee joint. *Osteoarthritis Cartilage* 1996;4:143-7.
42. Wluka AE, Wolfe R, Stuckey S, Cicuttini FM. How does tibial cartilage volume relate to symptoms in subjects with knee osteoarthritis? *Ann Rheum Dis* 2004;63:264-8.
43. Saito T, Koshino T. Distribution of neuropeptides in synovium of the knee with osteoarthritis. *Clin Orthop Relat Res* 2000;376: 172-82.
44. Bollet AJ. Edema of the bone marrow can cause pain in osteoarthritis and other diseases of bone and joints. *Ann Intern Med* 2001;134:591-3.
45. Teichtahl AJ, Wluka AE, Morris ME, Davis SR, Cicuttini FM. The relationship between the knee adduction moment and knee pain in middle-aged women without radiographic osteoarthritis. *J Rheumatol* 2006;33:1845-8.
46. Thorp LE, Sumner DR, Wimmer MA, Block JA. Relationship between pain and medial knee joint loading in mild radiographic knee osteoarthritis. *Arthritis Rheum* 2007;57:1254-60.
47. Felson DT, Niu J, Guermazi A, Roemer F, Aliabadi P, Clancy M, et al. Correlation of the development of knee pain with enlarging bone marrow lesions on magnetic resonance imaging. *Arthritis Rheum* 2007;56:2986-92.
48. Davies MA, Ettinger WH, Neuhaus JM, Cho SA, Hauck WW. The association of knee injury and obesity with unilateral and bilateral osteoarthritis of the knee. *Am J Epidemiol* 1989;130:278-88.
49. Felson DT, Anderson JJ, Naimark A, Hannan MT, Kannel WB, Meenan RF. Does smoking protect against osteoarthritis? *Arthritis Rheum* 1989;32:166-72.
50. Bellamy N, Buchanan WW, Goldsmith CH, Campbell J, Stitt LW. Validation study of WOMAC: a health status instrument for measuring clinically important patient relevant outcomes to anti-rheumatic drug therapy in patients with osteoarthritis of the hip or knee. *J Rheumatol* 1988;15:1833-40.

Osteoarthritis and Cartilage

Incidence and risk factors for radiographic lumbar spondylosis and lower back pain in Japanese men and women: the ROAD study

S. Muraki †*, T. Akune †, H. Oka ‡, Y. Ishimoto §, K. Nagata §, M. Yoshida §, F. Tokimura ||, K. Nakamura ¶, H. Kawaguchi #, N. Yoshimura ‡

† Department of Clinical Motor System Medicine, 22nd Century Medical & Research Center, Faculty of Medicine, the University of Tokyo, Tokyo, Japan

‡ Department of Joint Disease Research, 22nd Century Medical & Research Center, Faculty of Medicine, the University of Tokyo, Tokyo, Japan

§ Department of Orthopaedic Surgery, Wakayama Medical University, Wakayama, Japan

|| Department of Orthopaedic Surgery, Tokyo Geriatric Medical Center, Tokyo, Japan

¶ Rehabilitation Services Bureau, National Rehabilitation Center for Persons with Disabilities, Saitama, Japan

Department of Sensory & Motor System Medicine, Faculty of Medicine, the University of Tokyo, Tokyo, Japan

ARTICLE INFO

Article history:

Received 16 November 2011

Accepted 14 March 2012

Keywords:

Spondyloarthritis

Epidemiology

Lower back pain

SUMMARY

Objective: To determine the incidence of radiographic lumbar spondylosis (LS) and lower back pain, and their risk factors in Japan using a large-scale population from the nationwide cohort Research on Osteoarthritis/osteoporosis Against Disability (ROAD) Study.

Methods: Participants in the ROAD study who had been recruited between 2005 and 2007 were followed up with lumbar spine radiography for 3 years. A total of 2,282 paired radiographs (75% of the original sample) were scored using Kellgren and Lawrence (KL) grades, and the incidence and progression rate of radiographic LS was analyzed. The incidence of lower back pain was also examined. In addition, associations between risk factors and incident and progressive radiographic LS as well as incident lower back pain were tested.

Results: Given a 3.3-year follow-up, the incidence of KL ≥ 2 radiographic LS was 50.0% and 34.4% (15.3% and 10.5% per year), while that of KL ≥ 3 LS was 15.3% and 23.7% (4.6% and 7.2% per year) in men and women, respectively. The progression rate of LS was 20.5% and 27.4% (6.2% and 8.3% per year) in men and women, respectively. In addition, the incidence of lower back pain was 28.3% and 31.2% (8.6% and 9.5% per year) in men and women. Lower back pain was not significantly associated with incident radiographic LS, while a more severe KL grade at baseline was associated with incident lower back pain.

Conclusion: The present longitudinal study revealed a high incidence of radiographic LS in Japan.

© 2012 Osteoarthritis Research Society International. Published by Elsevier Ltd. All rights reserved.

Introduction

Lumbar spondylosis (LS) and lower back pain are considered a major public health issue causing chronic disability of the elderly in most developed countries^{1,2,3–8}. The prevalence of radiographic LS is high in Japan³, with an estimated 37,900,000 individuals aged ≥ 40 years being affected by radiographic LS⁹. According to the recent National Livelihood Survey of the Ministry of Health, Labour and Welfare in Japan, lower back pain is rated first among symptoms that send men to the hospital¹⁰. Despite the urgent need for

strategies to prevent and treat this condition, several cross sectional studies have investigated the prevalence of radiographic LS^{3,9,11–16}, but only a few have examined the incidence and progression of radiographic LS or their risk factors^{17–21}. In addition, although lower back pain is believed to be the principal clinical symptom of LS, the correlation is not as close as would be expected, and the findings of cross sectional studies have often indicated a disconnect between them^{3,11}. However, the incidences of radiographic LS and lower back pain have never been simultaneously analyzed in a longitudinal study.

The objective of the present study was to clarify the incidence and progression rate of radiographic LS as well as the incidence rate of lower back pain in Japanese men and women using the large-scale, population-based cohort study known as Research on Osteoarthritis/osteoporosis Against Disability (ROAD). In addition, we examined risk factors for the incidence and progression of LS as well as the incidence of lower back pain.

* Address correspondence and reprint requests to: S. Muraki, Department of Clinical Motor System Medicine, 22nd Century Medical & Research Center, Faculty of Medicine, the University of Tokyo, Tokyo, Japan. Tel: 81-3-5800-9178; Fax: 81-3-5800-9179

E-mail address: murakis-ort@h.u-tokyo.ac.jp (S. Muraki).

Subjects and methods

Subjects

The ROAD study was a nationwide prospective study of bone and joint diseases (with osteoarthritis and osteoporosis as representative bone and joint diseases) in population-based cohorts established in several communities in Japan. A detailed profile of the ROAD study has already been described elsewhere^{3,10,22,23}, and thus a brief summary is provided here. Between 2005 and 2007, we created a baseline database that included clinical and genetic information about 3,040 inhabitants (1,061 men; 1,979 women) in the age range of 23–95 years (mean, 70.6 years), recruited from listings of resident registrations in three communities: an urban region in Itabashi, Tokyo; a mountainous region in Hidakagawa, Wakayama; and a coastal region in Taiji, Wakayama. All participants provided written informed consent, and the study proceeded with the approval of ethics committees of the University of Tokyo and the Tokyo Metropolitan Institute of Gerontology. Participants completed an interviewer-administered questionnaire of 400 items that included lifestyle information such as smoking habits, alcohol consumption, family history, and medical history. Anthropometric measurements included height and weight, from which the body mass index (BMI) (weight [kg]/height² [m²]) was calculated. Experienced orthopaedists also asked all participants the question regarding lower back pain: "Have you experienced lower back pain on most days during the past month, in addition to now?" Those who answered "yes" were defined as having lower back pain based on previous studies³.

Between 2008 and 2010, we attempted to trace and review all 3,040 participants by inviting them to attend a follow-up interview and undergo repeat radiography. The interviews included questions about current lower back pain and were conducted by the same trained orthopaedists who undertook the baseline study between 2005 and 2007. Anthropometric data including height and weight, were also obtained at follow-up.

Radiographic assessment

Plain radiographs of the lumbar spine at baseline and follow-up were taken in anteroposterior and lateral positions, and the images were downloaded into Digital Imaging and Communication in Medicine (DICOM) files to assess radiographic spondylosis. We used contrast-adjusted images to detect osteophytes and intervertebral spaces when the original images were obscure. Osteophytes were analyzed at endplates. LS at baseline and follow-up was read in pairs according to the Kellgren and Lawrence (KL) grading²⁴ (without blinding to baseline and follow-up status) at each intervertebral level from L1/2 to L5/S by a single experienced orthopaedist (S.M.), who was blinded to the background of each patient. The KL scale defines radiographic OA in 5 categories: KL grade 0, no radiographic features of OA; KL grade 1, minimal osteophytosis only; KL grade 2, definite osteophytosis with some sclerosis of the anterior part of the vertebral plate; KL grade 3, marked osteophytosis and sclerosis of the vertebral plates with slight narrowing of the disc space; and KL grade 4, large osteophytes, marked sclerosis of the vertebral plates, and marked narrowing of the disc space. To evaluate the intraobserver variability of the KL grading, 100 randomly selected radiographs of the lumbar spine were scored by the same observer more than 1 month after the first reading. Furthermore, 100 other radiographs were scored by two experienced orthopaedic surgeons (S.M. and H.O.) using the same radiographic atlas to determine interobserver variability. Intra- and interobserver variability was evaluated by kappa

analysis. These variabilities in the KL grading on lumbar radiographs were sufficient for assessment (0.84 and 0.76, respectively).

For the purposes of this study, we defined three LS outcomes. Incident KL ≥ 2 radiographic LS was defined if all vertebral interspaces had less than grade 2 disease at baseline, and if at least one vertebral interspace had grade ≥ 2 disease at follow-up. Incident KL ≥ 3 radiographic LS was defined if all vertebral interspaces had less than grade 3 disease at baseline, and if at least one vertebral interspace had \geq grade 3 disease at follow-up. Progressive LS was defined as KL ≥ 2 LS at baseline, excluding subjects with KL = 4 LS at all vertebral interspaces because it cannot progress, and an increase of at least one grade in the affected vertebral interspace at follow-up.

Statistical analysis

Differences in age, height, weight, and BMI between men and women were examined using a non-paired Student's *t*-test. The prevalence of radiographic LS and lower back pain between men and women was compared the chi-squared test. We determined risk factors for incident and progressive LS and incident lower back pain using a univariate logistic regression analysis. Independent risk factors were determined by multiple logistic regression analysis with significant risk factors in a univariate logistic regression analysis with age, gender and BMI, as independent variables. Incident lower back pain was defined as no lower back pain at baseline and lower back pain at follow-up. Associations between the number of affected vertebral interspaces and incident lower back pain were assessed using the Cochran-Armitage test for trends. The odds ratio (OR) and 95% confidence interval (CI) of the number of affected vertebral interspaces with incident lower back pain compared with no affected vertebral interspaces was determined using a logistic regression analysis with adjustment for age and BMI. Data were analyzed using SAS version 9.0 software (SAS Institute Inc., Cary, NC).

Results

Of the 3,040 participants in the baseline study between 2005 and 2007, 125 (4.1%) had died by the time of the review 3 years later, 123 (4.0%) did not participate in the follow-up study due to bad health, 69 (2.3%) had moved, 83 (2.7%) declined the invitation to attend the follow-up study and 155 (5.1%) did not participate for other reasons. Among the 2,485 individuals who participated in the follow-up study, we excluded 186 (6.1%) who did not undergo plain radiography and 17 (0.6%) who provided incomplete pain questionnaires, leaving a total of 2,282 (74.4%; 758 men; 1,524 women) from whom paired radiographs and complete pain histories were obtained. Their median age at follow-up was 72.1 \pm 11.5 years. The duration of follow-up between the initial and second radiographs was 3.3 \pm 0.6 (mean \pm SD) years. Those participating in the follow-up study were younger than those who did not survive or who did not participate for other reasons (responders 68.8 years, nonresponders 74.8 years; $P < 0.0001$). The follow-up study participants were also more likely to be women (responders 66.8% women, nonresponders 60.0% women; $P = 0.0007$) and were less likely to have LS at the baseline examination than either those who did not survive to follow-up or those who did not participate for other reasons (responders 68.1%, nonresponders 77.5%; $P < 0.0001$). The prevalence of lower back pain was similar between responders and nonresponders (responders 19.0%, nonresponders 18.7%; $P = 0.91$).

Table I shows the characteristics of the 2,282 participants at baseline in the ROAD study. Men were significantly older than women, and the BMI was higher in men than women. The prevalence of KL ≥ 2 LS was significantly higher in men than women at

Table I
Characteristics of participants at baseline

	Men	Women	P-values
Number of subjects	758	1,524	
Age at baseline, years	69.8 ± 11.0	68.3 ± 11.3	0.003
Height at baseline, cm	163.0 ± 6.6	150.4 ± 6.4	<0.0001
Weight at baseline, kg	62.0 ± 9.7	52.1 ± 8.6	<0.0001
BMI at baseline, kg/m ²	23.3 ± 3.0	23.0 ± 3.4	0.054
Grip strength at baseline, kg	34.3 ± 8.7	22.2 ± 6.1	<0.0001
Prevalence at baseline			
KL ≥ 2 (%)	79.9	62.3	<0.0001
KL ≥ 3 (%)	43.1	44.6	0.531
Lower back pain (%)	16.9	20.0	0.073
Smoking (%)	21.5	3.2	<0.0001
Alcohol (%)	63.2	23.0†	<0.0001

Except where indicated otherwise, values represent mean ± SD.

* $P < 0.05$ vs. men by non-paired Student's *t*-test; † $P < 0.05$ vs men by chi squared test.

baseline, while that of KL ≥ 3 LS and lower back pain was similar between men and women.

Table II shows the rates of incident and progressive radiographic LS as well as that of incident lower back pain. Given the 3.3-year follow-up, the rates of incident KL ≥ 2 and ≥ 3 LS and progressive LS, and incident lower back pain were 38%, 21%, 25%, and 30%, respectively. The incidence of KL ≥ 2 LS was significantly lower, but that of KL ≥ 3 LS was significantly higher in women than in men. The rate of progressive LS was also significantly higher in women than men. The rate of incident and progressive LS increased with age in men and women ($P < 0.05$) (Fig. 1). The rate of incident lower back pain was not age-dependent in either men or women ($P = 0.44$ and 0.85 , respectively) (Fig. 1). We also showed incidence and progression of LS at each vertebral interspace in Supplementary Table. Among the vertebral interspaces, the incident rate of KL ≥ 2 LS was highest at the L2/3 interspace. While, the incident rate of KL ≥ 3 LS was highest at the L4/5 interspace.

Table III shows baseline risk factors for radiographic LS. Multiple logistic regression analysis showed that age was a risk factor for KL ≥ 2 and KL ≥ 3 LS and that higher BMI was a risk factor for KL ≥ 2, but not for KL ≥ 3. Female gender was a protective factor against the incidence of KL ≥ 2 LS but was a risk factor for the incidence of KL ≥ 3 LS. A higher KL grade at baseline was a risk factor for KL ≥ 3 LS. Lower back pain at baseline, smoking and alcohol consumption were not associated with incident KL ≥ 2 or KL ≥ 3 LS. We further examined the risk factors for progressive LS in individuals with KL ≥ 2 LS, excluding those with KL = 4 LS at all vertebral interspaces (Table IV). Age and female gender were also risk factors for progressive LS, whereas BMI, lower back pain at baseline, smoking and alcohol consumption were not associated with progressive LS. A grade of KL ≥ 3 at baseline was a risk factor for progressive LS compared with KL = 2.

We next examined the risk factors for incident lower back pain (Table IV). KL ≥ 3 LS was associated with incident lower back pain

compared with KL = 0 or 1, whereas age, BMI, gender, smoking and alcohol consumption were not associated with incident lower back pain. We next examined the association between KL grade at each vertebral interspace and incident lower back pain (Table V). In women, KL ≥ 3 LS at L2/3, 3/4, 4/5, and 5/S and the most severely affected interspaces were significantly associated with incident lower back pain compared with KL < 3 at the corresponding interspaces. KL ≥ 3 LS at L2/3, 3/4, 4/5 and 5/S in men tended to be associated with incident lower back pain compared with KL < 3 at the corresponding interspaces, but these findings did not reach statistical significance except for the L3/4 and L5/S interspaces. KL ≥ 3 LS at the L1/2 interspace was not associated with incident lower back pain in men or women. Thus, we further examined the number of KL ≥ 3 vertebral interspaces among L2/3, 3/4, 4/5 and 5/S interspaces (Supplementary Fig. 1). The Cochran-Armitage test for trends showed that the incidence rate of lower back pain significantly increased as the number of affected vertebral interspaces increased in women ($P < 0.001$), but not in men ($P = 0.09$). In addition, multiple logistic regression analysis after adjustment for age and BMI showed that having three or more KL ≥ 3 vertebral interspaces was significantly associated with incident lower back pain in men (OR 1.69 95% CI 1.03–2.76) and in women (OR 1.77, 95% CI 1.34–2.34).

Discussion

This is the first population-based study to examine the rates of incident and progressive radiographic LS as well as incident lower back pain, and their risk factors in Japanese men and women. We found high rates of incident and progressive LS and incident lower back pain in Japanese men and women.

Few population-based studies have examined incident radiographic LS^{17,18}. Symmons *et al.* examined radiographic changes in the lumbar spines of Dutch women (mean age, 54 years) using KL grade¹⁷ and found that 4.2% per year of individuals with no disc degeneration (KL grade 0/1) but with recurrent back pain, and 3.2% per year of those with no disc degeneration and no back pain at baseline, had disc degeneration at follow-up. The present study found a 27.6% incidence rate of KL ≥ 2 LS in women aged in their 50s over a period of 3.3 years (9.0% per year), and thus the incidence of KL ≥ 2 LS is apparently considerably higher in Japanese than Caucasian women, although a strict comparison may be limited because of differences in definition of the incidence of LS. Considering the definition of the KL grade, this may suggest that the incidence of osteophytosis is higher in Japanese women than in Caucasian women.

Regarding progression of radiographic LS, Symmons *et al.* reported that 63.1% (7.0% per year) of individuals with disc degeneration and with recurrent back pain, and 55.4% (6.2% per year) of those with disc degeneration but without back pain at baseline, had worse disc degeneration at follow-up¹⁷. The present

Table II
Incidence of radiographic LS and progressive LS as well as incidence of lower back pain

	KL ≥ 2 LS		KL ≥ 3 LS		Progressive LS		Lower back pain	
	No. at risk	Incidence (%)	No. at risk	Incidence (%)	No. at risk	Incidence (%)	No. at risk	Incidence (%)
Overall	727	274 (37.7)	1,276	266 (20.8)	1,530	378 (24.7)	1,849	558 (30.2)
Men	152	76 (50.0)	431	66 (15.3)	599	123 (20.5)	630	178 (28.3)
Women	575	198 (34.4)*	845	200 (23.7)*	931	255 (27.4)*	1,219	380 (31.2)

Incident KL ≥ 2 and ≥ 3 radiographic LS at the overall vertebral interspace was defined as all vertebral interspaces having less than grade 2 or 3 disease at baseline, and if at least one vertebral interspace was grade 2 or higher or grade 3 or higher at follow-up, respectively.

Progressive LS in the overall inter spaces was defined as KL ≥ 2 LS at baseline, excluding subjects with KL = 4 LS at all vertebral interspaces because it cannot progress, and an increase by at least 1 grade in the affected vertebral interspace at follow-up.

Incident lower back pain was defined as no lower back pain at baseline and lower back pain at follow-up.

* $P < 0.05$ vs men by chi square test.




Fig. 1. Ratios (%) of individuals with incident radiographic LS (KL ≥ 2 and KL ≥ 3), progressive LS and incident lower back pain in each age stratum (<50, 50–59, 60–69, 70–79 and ≥80 years). Data in brackets are the number of individuals in each group.

study found that the progression rate of LS in women aged in their 50s was 20.9% over a period of 3.3 years (6.3% per year) and thus the progression rate of LS appears similar between Japanese and Caucasian women. In the present study, progression of radiographic LS was defined as KL ≥ 2 LS at baseline and an increase of at least one grade in the affected vertebral interspace at follow-up. Considering the definition of the KL grade, progression of radiographic LS may mean incidence or progression of disc space narrowing in subjects with osteophytosis, thus, our results may

indicate that the incidence or progression of disc space narrowing is similar between Japanese and Caucasian women.

Furthermore, the present study included an investigation of KL ≥ 3 LS. To the best of our knowledge, incident KL ≥ 3 LS has never been investigated in a population-based study. LS was not defined according to KL grade but according to osteophytosis and disk space narrowing in the Chingford study¹⁸. That study showed that the progression of disk space narrowing was 3.2% per year in women whose mean age was 54 years at baseline. Our results regarding incident KL ≥ 3 LS might be comparable to these, considering the definition of KL grade, although a detailed comparison provides only limited accuracy. The incidence rate of KL ≥ 3 LS was 15.0% (4.5% per year) in Japanese women aged in their 50s at baseline in the present study, which was also higher than that in Caucasian women. This might in part be related to ethnic variations.

The incidence of KL ≥ 2 spondylosis was notably higher in men than in women, while that of KL ≥ 3 spondylosis was higher in women in the present study. Considering the definition of KL grade, this might mean that the incidence of osteophytosis is higher in men, whereas the incidence of disk space narrowing is higher in women. In fact, osteophytosis of the lumbar spine is more common in men than in women^{11,12}, whereas disk space narrowing is more prevalent in women¹². A cross-sectional study that investigated the extent, prevalence and distribution of spinal spondylosis in women also showed that osteophytosis and disk space narrowing significantly correlated, but each predicted only 19% of the variation in the other¹³. This discordance suggests that different mechanisms influence the initiation of osteophytosis (the principal abnormality in KL grade 2 disease) and disk space narrowing (a principal abnormality in KL grade 3 disease). Our findings have implications for understanding of the pathogenesis of LS, as well as for designing preventive strategies.

In the present study, age, BMI, gender and KL grade at baseline were significantly associated with incident LS; this result differed from the findings of previous studies^{19–21}. The UK twin spine study¹⁹ using magnetic resonance imaging (MRI) showed that age, BMI and gender had no detectable effect on the progression of lumbar disc degeneration. The Finnish twin spine study also showed that body weight was not associated with progression of lumbar disc degeneration²⁰. These differences may be explained not only by the differences in the definition of progressive LS, but also the ages of the subjects between these previous studies and the present study. The subjects in the UK twin study and Finnish twin study were quite younger at baseline than those in the present study (55 years, 49 years and 69 years, respectively). The association of these factors with LS may change among the age strata. In addition, racial differences may exist in the association of these factors with LS, because the prevalence or incidence of LS is different among races³. Age, BMI and female gender were not risk factors for lower back pain in the present study. Lower back pain occurrence might be mainly due to environmental, rather than to individual factors. Elderly men in particular generally retire at around age 60–70 years, and thus the load on the lower back might be greater in men aged below 60 years compared with those over 60 years, which might partly explain the lack of a significant association between age and the incidence of lower back pain. KL ≥ 3 LS was significantly associated with incident lower back pain compared with the absence of LS. Cross sectional studies have shown that the correlation between LS and lower back pain is not as strong as would be expected, and they are often disconnected^{3,11}. However, this longitudinal study discovered that severe radiographic LS is a risk factor for lower back pain. We further found that the association between the number of KL ≥ 3 vertebral interspaces and the incidence of lower back pain differed between men and

Table III
Baseline risk factors for incident radiographic LS.

	KL ≥ 2					KL ≥ 3				
	No (%)	Crude OR	95% CI	Adjusted OR	95% CI	No (%)	Crude OR	95% CI	Adjusted OR	95% CI
Age, years		1.05	1.03–1.06	1.05	1.03–1.06		1.05	1.04–1.07	1.05	1.03–1.06
BMI, kg/m ²		1.07	1.02–1.12	1.07	1.02–1.13		1.01	0.97–1.06		
Gender										
Men	76/152 (50.0)	1.00	Reference	1.00	Reference	66/431 (15.3)	1.00	Reference	1.00	Reference
Women	198/575 (34.4)	0.53	0.37–0.76	0.50	0.34–0.72	200/845 (23.7)	1.71	1.27–2.34	2.19	1.54–3.17
Low back pain										
No	223/607 (36.7)	1.00	Reference			219/1078 (20.3)	1.00	Reference		
Yes	51/120 (42.5)	1.27	0.85–1.89			47/198 (23.7)	1.22	0.85–1.74		
Smoking										
No	244/661 (36.9)	1.00	Reference			246/1136 (21.7)	1.00	Reference	1.00	Reference
Yes	30/66 (45.5)	1.42	0.85–2.37			20/140 (14.3)	0.60	0.36–0.97	1.01	0.58–1.69
Alcohol										
No	184/476 (38.7)	1.00	Reference			185/774 (23.9)	1.00	Reference	1.00	Reference
Yes	90/251 (35.9)	0.89	0.64–1.22			81/502 (16.1)	0.61	0.46–0.82	0.87	0.63–1.20
KL grade										
KL = 0 or 1							1.00	Reference	1.00	Reference
KL = 2							1.66	1.27–2.19	1.67	1.24–2.25

The adjusted ORs were calculated by multiple logistic regression analysis after adjustment for all other significant variables without adjustment. We did not include KL grade in the analysis of incident KL ≥ 2 LS, because all subjects had KL = 0 or 1.

women. The incidence of lower back pain increased as the number of KL ≥ 3 vertebral interspaces increased in women, whereas the incidence was similar in men with 0, 1 and 2 KL ≥ 3 vertebral interspaces, and having 3 or more KL ≥ 3 vertebral interspaces suddenly increased the incidence of lower back pain.

There were several limitations in this study. First, we did not read the X-rays for osteophytes and joint space narrowing scored separately. Furthermore, in the KL classification, atrophic and degenerative features of LS, which likely have different aetiology, were combined; thus, the differences in associations with pain between these features may have been obscured. We are developing a computer-aided diagnostic program to enable fully automated measurements of the major features of LS, including disc space narrowing and osteophytosis on plain radiographs. The second limitation of our study was that a single orthopaedist read both films in pairs without being blinded to baseline and follow-up status. This may likely have caused the reader to over-read progression (i.e., inflate sensitivity) and therefore confer bias. This may be one reason for the higher incidence of LS in the present study compared with other studies. Third, we used only plain radiography to assess LS, although computed tomography (CT)/MRI is standard practice for evaluating nonspecific lower back pain in

many countries. In addition, plain films can be affected by scoliosis, positioning and multiple other factors, which may have affected our results. Fourth, although experienced orthopaedists asked all participants the question regarding lower back pain based on previous studies^{3,8}, we defined lower back pain as present or absent, rather than as a continuous validated measure of pain, such as assessed by the Oswestry Disability Index²⁵. Categorical methods are statistically less powerful than continuous methods. In addition, severity of lower back pain was not assessed in the present study. The association between lower back pain and other variables might have been underestimated in the present study. Furthermore, although the psychosocial dimension is an important factor for lower back pain²⁶, we did not include this in our analysis. Fifth, in the follow-up study, the responders was younger, more likely to be women and less likely to have LS at baseline compared with the nonresponders, which may have affected the results in the present study, because age, gender and KL grade were found to be associated with incident LS in the present study.

In conclusion, the present longitudinal study using a large-scale population from the ROAD study revealed a high incidence of radiographic LS in Japan. Gender seems to be distinctly associated with incident KL ≥ 2 and KL ≥ 3 LS, indicating that different

Table IV
Baseline risk factors for progressive LS and incident lower back pain

	Progressive LS					Lower back pain				
	No (%)	Crude OR	95% CI	Adjusted OR	95% CI	No (%)	Crude OR	95% CI	Adjusted OR	95% CI
Age, years		1.05	1.04–1.07	1.05	1.04–1.07		1.00	0.99–1.01	1.00	0.99–1.01
BMI, kg/m ²		1.01	0.98–1.05				1.01	0.98–1.04	1.01	0.98–1.04
Gender										
Men	123/599 (20.5)	1.00	Reference	1.00	Reference	178/630 (28.3)	1.00	Reference	1.00	Reference
Women	255/931 (27.4)	1.46	1.14–1.87	1.44	1.10–1.91	380/1219 (31.2)	1.15	0.93–1.42	1.12	0.90–1.39
Low back pain										
No	302/1225 (24.7)	1.00	Reference							
Yes	76/305 (24.9)	1.01	0.76–1.35							
Smoking										
No	348/1385 (25.1)	1.00	Reference			503/1677 (30.0)	1.00	Reference		
Yes	30/145 (20.7)	0.78	0.50–1.17			55/172 (32.0)	1.10	0.78–1.53		
Alcohol										
No	253/958 (26.4)	1.00	Reference			360/1162 (31.0)	1.00	Reference		
Yes	125/572 (21.9)	0.78	0.61–0.99			198/687 (28.8)	0.90	0.73–1.11		
KL grade										
KL = 0 or 1						177/607 (29.2)	1.00	Reference	1.00	Reference
KL = 2	103/549 (18.8)	1.00	Reference			118/471 (25.1)	0.81	0.62–1.06	0.86	0.64–1.14
KL ≥ 3	275/981 (28.0)	1.69	1.31–2.18			263/771 (34.1)	1.26	1.00–1.58	1.32	1.03–1.69

The adjusted ORs were calculated by multiple logistic regression analysis after adjustment for all other significant variables without adjustment. We did not include KL grade in the analysis of incident KL ≥ 2 LS, because all subjects had KL = 0 or 1.

Table V
Association of KL ≥ 3 LS at baseline with incident lower back pain by each vertebral interspace and the severest space in 1,849 subjects with no lower back pain at baseline

	L1/2		L2/3		L3/4		L4/5		L5/S		The severest	
	No. (%)	OR (95% CI)	No. (%)	OR (95% CI)	No. (%)	OR (95% CI)	No. (%)	OR (95% CI)	No. (%)	OR (95% CI)	No. (%)	OR (95% CI)
Men	KL < 3 154/552 (27.9)	1.00	142/528 (26.9)	1.00	136/512 (26.6)	1.00	117/424 (27.6)	1.00	130/496 (26.2)	1.00	98/368 (26.6)	1.00
N = 630	KL ≥ 3 24/78 (30.8)	1.20 (0.70–2.01)	36/102 (35.3)	1.57 (0.98–2.48)	42/118 (35.6)	1.62 (1.04–2.50)	61/206 (29.6)	1.15 (0.79–1.67)	48/134 (35.8)	1.65 (1.09–2.49)	80/262 (30.5)	1.26 (0.88–1.81)
Women	KL < 3 331/1,083 (30.6)	1.00	298/1,007 (29.6)	1.00	284/960 (29.6)	1.00	236/828 (28.5)	1.00	284/971 (29.3)	1.00	197/710 (27.8)	1.00
N = 1,219	KL ≥ 3 49/136 (36.0)	1.28 (0.87–1.87)	82/212 (38.7)	1.52 (1.11–2.10)	96/259 (37.1)	1.43 (1.06–1.92)	144/391 (36.8)	1.50 (1.15–1.97)	96/248 (38.7)	1.56 (1.15–2.10)	183/509 (36.0)	1.51 (1.16–1.95)

Multiple logistic regression analysis after adjustment for age was used to calculate OR and 95% CI.

mechanisms might influence the initiation of osteophytosis and joint space narrowing. Lower back pain was not significantly associated with incident radiographic LS, whereas radiographic severe LS was a risk factor for incident lower back pain. Further progress, along with continued longitudinal surveys of the ROAD study, will elucidate the environmental and genetic background of LS.

Author contributions

All authors have made substantial contributions to all three of sections (1), (2) and (3) below;

- (1) The conception and design of the study, or acquisition of data, or analysis and interpretation of data
- (2) Drafting the article or revising it critically for important intellectual content
- (3) Final approval of the version to be submitted

Conflicts of interest

There are no conflicts of interest.

Acknowledgements

This study was supported by Grants-in-Aid for Scientific Research (S19109007, B20390182, C20591737, C20591774), for Young Scientists (A18689031), and for Exploratory Research (19659305) from the Japanese Ministry of Education, Culture, Sports, Science and Technology, H17-Men-eki-009, H18-Choujyu-037, H20-Choujyu-009, and H21-Chouju-Wakate-011 from the Ministry of Health, Labour and Welfare, Research Aid from the Japanese Orthopaedic Association (JOA-Subsidized Science Project Research 2006-1 & 2010-2), and Grant No.166 from the Japan Orthopaedics and Traumatology Foundation.

The authors thank Drs Seizo Yamamoto, Hideaki Ishibashi, Yorito Anamizu and members of Department of Orthopaedics, as well as members of the Department of Radiology at Tokyo Metropolitan Geriatric Medical Center. We also appreciate Mrs Tomoko Takijiri and other members of the Public Office in Hidakagawa, Mrs Tamako Tsutsumi, Mrs Kanami Maeda, and other members of the Public Office in Taiji, for assistance with the location and scheduling of participants for examinations.

Supplementary material

Supplementary data related to this article can be found online at doi:10.1016/j.joca.2012.03.009

References

1. Hadjipavlou AG, Simmons JW, Pope MH, Necessary JT, Goel VK. Pathomechanics and clinical relevance of disc degeneration and annular tear: a point-of-view review. *Am J Orthop* 1999;28:561–71.
2. Emery SE, Ringus VM. Osteoarthritis of the spine. In: Moskowitz RW, Altman RD, Hochberg MC, Buckwalter JA, Goldberg VM, Eds. *Osteoarthritis: Diagnosis and Medical/Surgical Management*. 4th edn. Philadelphia: Lippincott Williams & Wilkins; 2007:427–52.
3. Muraki S, Oka H, Akune T, Mabuchi A, En-Yo Y, Yoshida M, et al. Prevalence of radiographic lumbar spondylosis and its association with lower back pain in the elderly of

- population-based cohorts: the ROAD study. *Ann Rheum Dis* 2008;68:1401–6.
4. Horng YS, Hwang YH, Wu HC, Liang HW, Mhe YJ, Twu FC, et al. Predicting health-related quality of life in patients with lower back pain. *Spine* 2005;30:551–5.
 5. Kovacs FM, Abraira V, Zamora J, Teresa Gil del Real M, Llobera J, Fernández C, et al. Correlation between pain, disability, and quality of life in patients with common lower back pain. *Spine* 2004;29:206–10.
 6. Leidig-Bruckner G, Minne HW, Schlaich C, Wagner G, Scheidt-Nave C, Bruckner T, et al. Clinical grading of spinal osteoporosis: quality of life components and spinal deformity in women with chronic lower back pain and women with vertebral osteoporosis. *J Bone Miner Res* 1997;12:663–75.
 7. Silverman SL, Piziak VK, Chen P, Misurski DA, Wagman RB. Relationship of health related quality of life to prevalent and new or worsening back pain in postmenopausal women with osteoporosis. *J Rheumatol* 2005;32:2405–9.
 8. Muraki S, Akune T, Oka H, En-yo Y, Yoshida M, Saika A, et al. Impact of knee and lower back pain on health-related quality of life in Japanese women: the Research on Osteoarthritis Against Disability (ROAD). *Modern Rheum* 2010;20:444–51.
 9. Yoshimura N, Muraki S, Oka H, Mabuchi A, En-yo Y, Yoshida M, et al. Prevalence of knee osteoarthritis, lumbar spondylosis and osteoporosis in Japanese men and women: The Research on Osteoarthritis/osteoporosis Against Disability (ROAD). *J Bone Miner Metab* 2009;27:620–8.
 10. Ministry of Health, Labour and Welfare. The outline of the results of National Livelihood Survey 2007 available at <http://www.mhlw.go.jp/toukei/list/20-19-1.html>
 11. O'Neill TW, McCloskey EV, Kanis JA, Bhalla AK, Reeve J, Reid DM, et al. The distribution, determinants, and clinical correlates of vertebral osteophytosis: a population based survey. *J Rheumatol* 1999;26:842–8.
 12. Yoshimura N, Dennison E, Wilman C, Hashimoto T, Cooper C. Epidemiology of chronic disc degeneration and osteoarthritis of the lumbar spine in Britain and Japan: a comparative study. *J Rheumatol* 2000;27:429–33.
 13. Kramer PA. Prevalence and distribution of spinal osteoarthritis in women. *Spine* 2006;31:2843–8.
 14. Kellgren JH, Lawrence JS. Osteo-arthrosis and disk degeneration in an urban population. *Ann Rheum Dis* 1958;17:388–97.
 15. Lawrence JS. Disc degeneration. Its frequency and relationship to symptoms. *Ann Rheum Dis* 1969;28:121–38.
 16. van Saase JL, van Romunde LK, Cats A, Vandenbroucke JP, Valkenburg HA. Epidemiology of osteoarthritis: Zoetermeer survey. Comparison of radiological osteoarthritis in a Dutch population with that in 10 other populations. *Ann Rheum Dis* 1989;48:271–80.
 17. Symmons DP, van Hemert AM, Vandenbroucke JP, Valkenburg HA. A longitudinal study of back pain and radiological changes in the lumbar spines of middle aged women. I. Clinical findings. *Ann Rheum Dis* 1991;50:158–61.
 18. Hassett G, Hart DJ, Manek NJ, Doyle DV, Spector TD. Risk factors for progression of lumbar spine disc degeneration: the Chingford Study. *Arthritis Rheum* 2003;48:3112–7.
 19. Williams FM, Popham M, Sambrook PN, Jones AF, Spector TD, MacGregor AJ. Progression of lumbar disc degeneration over a decade: a heritability study. *Ann Rheum Dis* 70: 1203–7.
 20. Videman T, Battie MC, Parent E, Gibbons LE, Vainio P, Kaprio J. Progression and determinants of quantitative magnetic resonance imaging measures of lumbar disc degeneration: a five-year follow-up of adult male monozygotic twins. *Spine (Phila Pa 1976)* 2008;33:1484–90.
 21. Videman T, Battie MC, Ripatti S, Gill K, Manninen H, Kaprio J. Determinants of the progression in lumbar degeneration: a 5-year follow-up study of adult male monozygotic twins. *Spine (Phila Pa 1976)* 2006;31:671–8.
 22. Muraki S, Oka H, Akune T, Mabuchi A, En-yo Y, Yoshida M, et al. Prevalence of radiographic knee osteoarthritis and its association with knee pain in the elderly of Japanese population-based cohorts: the ROAD study. *Osteoarthritis Cartilage* 2009;17:1137–43.
 23. Yoshimura N, Muraki S, Oka H, Kawaguchi H, Nakamura K, Akune T. Cohort Profile: Research on Osteoarthritis/osteoporosis Against Disability (ROAD) Study. *Int J Epidemiol* 2010;39:988–95.
 24. Kellgren JH, Lawrence JS, Eds. The epidemiology of chronic rheumatism: atlas of standard radiographs of arthritis. Oxford, England: Blackwell Scientific; 1963.
 25. Fairbank J, Couper J, Davies J, O'Brien JP. The Oswestry low back pain questionnaire. *Physiotherapy* 1980;66:271–3.
 26. Hartvigsen J, Frederiksen H, Christensen K. Physical and mental function and incident low back pain in seniors: a population-based two-year prospective study of 1387 Danish Twins aged 70 to 100 years. *Spine (Phila Pa 1976)* 2006;31:1628–32.