

### Falls assessment

In 2008–2010, we attempted to trace and review all 3,040 subjects; they were invited to attend a follow-up interview. All subjects were interviewed with regard to falls by experienced interviewers and were asked the following questions: “Have you experienced falls during 3 years of follow-up, and if yes, how many falls did you experience?” According to a previous study on falls [35], a fall is defined as a sudden, unintentional change in position causing an individual to land at a lower level on an object, the floor, or the ground, other than as a consequence of a sudden onset of paralysis, epileptic seizure, or overwhelming external force.

### Pain assessment

All subjects were interviewed by experienced orthopedists with regard to knee pain and lower back pain at baseline and were asked the following questions based on previous studies [17, 18]: “Have you experienced knee pain on most days in the past year, in addition to now?” and “Have you experienced lower back pain on most days in the past year, in addition to now?” Those who answered yes were defined as having pain.

### Radiographic assessment

At baseline, all participants underwent radiographic examination of both knees using anteroposterior and lateral views with weight-bearing and foot map positioning; radiographic examination of the anteroposterior and lateral views of the lumbar spine, including intervertebral levels L1/2 to L5/S, was also performed. Knee and lumbar spine radiographs were read without the knowledge of participant clinical status by a single, experienced orthopedist (S.M.) using the Kellgren–Lawrence (KL) radiographic atlas [36] to determine the severity of KL grading. Radiographs were scored as grade 0 through 4, with higher grades being associated with more severe OA. We defined knee OA and LS as  $KL \geq 3$  in at least one knee and one intervertebral level, respectively. To evaluate the intraobserver variability of KL grading, 100 randomly selected radiographs of the knee and the lumbar spine were scored by the same observer more than 1 month after the first reading. One hundred other radiographs were also scored by two experienced orthopedic surgeons (S.M. and H.O.) using the same atlas for interobserver variability. The intraobserver and interobserver variabilities evaluated were confirmed by kappa analysis to be sufficient for assessment (0.86 and 0.80 for knee OA and 0.84 and 0.76 for LS, respectively).

### Physical performance

Anthropometric measurements included height, weight, and body mass index (BMI) (weight [in kilograms]/height<sup>2</sup> [in

square meters]) at baseline. Grip strength was also measured on bilateral sides using a TOEI LIGHT handgrip dynamometer (TOEI LIGHT CO., LTD., Saitama, Japan) at baseline, and the best measurement was used to characterize maximum muscle strength. To measure physical performance, the time taken to walk 6 m at normal walking speed in a hallway was recorded. Subjects were told to walk from a marked starting line to a 6-m mark as if they were walking down their hallway at home. Time was measured in seconds with a stopwatch and rounded to the nearest hundredth of a second. These walking speed trial measurements are considered highly reliable in community-dwelling elderly subjects [34, 37–39].

### Statistical analyses

The differences in age, anthropometric measurements, and physical performance measurements between men and women and between nonfallers and fallers were examined by a nonpaired Student's *t* test. The incidence of falls was also compared between men and women, among subjects with no severe knee OA (KL=0, 1, or 2) and KL=3 or 4 knee OA, among subjects with no severe LS (KL=0, 1, or 2) and KL=3 or 4 LS, among subjects with and without knee pain, and among subjects with and without lower back pain using the chi-square test. Multiple logistic regression analysis after adjustment for age and BMI was used to determine the association of anthropometric measurements, physical performance, radiographic knee OA and LS defined as KL=3 or 4, and knee and lower back pain and with falls compared with nonfalls in men and women. Further, to determine an independent association of physical performance, radiographic knee OA, and knee pain with falls compared with nonfalls, we used multiple logistic regression analysis with age, BMI, walking speed, radiographic knee OA, and knee pain as independent variables. Data analyses were performed using SAS version 9.0 (SAS Institute Inc., Cary, NC, USA).

### Results

Of the 3,040 subjects in the baseline study in 2005–2007, 125 (4.1 %) had died by the time of the review 3 years later, 123 (4.0 %) did not participate in the follow-up study due to bad health, 69 (2.3 %) had moved away, 83 (2.7 %) declined the invitation to attend the follow-up study, and 155 (5.1 %) did not participate in the follow-up study for other reasons. Among the 2,485 subjects who did participate in the follow-up study, 182 (6.0 %) provided incomplete fall questionnaires. In addition, 15 (0.5 %) provided incomplete pain questionnaires; these were excluded. We also excluded 14 (0.5 %) subjects who had undergone total knee arthroplasty at baseline. Further, 59 (1.9 %) subjects did not measure

walking speed, leaving a total of 2,215 (72.9 %) subjects (745 men and 1,470 women) from whom radiographs at baseline and complete fall and pain histories were obtained. The mean  $\pm$  SD duration of follow-up between initial and second surveys was  $3.3\pm 0.6$  years.

Table 1 shows the age, anthropometric measurements, physical performance, and prevalence of radiographic knee OA and LS as well as knee and lower back pain of participants at baseline. Regarding physical performance, grip strength and walking speed were significantly better in men than in women. The prevalence of radiographic knee OA and knee pain was significantly higher in women than in men, whereas that of LS and lower back pain was not different between men and women.

During the approximately 3-year follow-up, 141 (18.9 % [95 % confidence interval [CI], 16.3–21.9]) men and 362 (24.6 % [95 % CI, 22.5–26.9]) women reported at least one fall. Chi-square test showed that the incidence of falls were significantly different between men and women ( $p=0.0025$ ). With increasing age, the incidence of falls tended to increase in men and women (Fig. 1).

Table 2 shows the age, anthropometric measurements, and physical performance at baseline between nonfallers and fallers. Age was significantly higher in fallers than nonfallers in men and women. Height was higher in fallers than in nonfallers in women, whereas weight and BMI was not significantly different between nonfallers and fallers in men and women. Grip strength and walking speed were worse in fallers than nonfallers in men and women.

Figure 2 shows the incidence rate of falls according to knee OA, knee pain, LS, and lower back pain. The incidence rate of falls was higher in subjects with knee OA than those without knee OA in men (27.9 and 18.0 %,  $p<0.05$ ,

respectively) and women (33.1 and 22.6 %,  $p<0.05$ , respectively). The incidence rate of falls was also higher in subjects with knee pain than those without knee pain in men (30.4 and 17.1 %,  $p<0.05$ , respectively) and women (32.6 and 22.1 %,  $p<0.05$ , respectively). There were no significant differences in incidence rate of falls between subjects with and without LS in men (20.5 and 17.8 %,  $p=0.35$ , respectively) and women (25.5 and 23.5 %,  $p=0.39$ , respectively). Men with lower back pain had significantly higher incidence rate of falls than men without lower back pain (25.6 and 17.6 %,  $p<0.05$ , respectively), whereas women with lower back pain did not (23.8 and 24.8 %,  $p=0.76$ , respectively).

In men, multiple logistic regression analysis after adjustment for age and BMI showed that slower walking speed ( $p<0.001$ ) and knee pain ( $p=0.0046$ ) were risk factors for falls, but grip strength ( $p=0.4903$ ), radiographic knee OA ( $p=0.1569$ ), LS ( $p=0.8312$ ), and lower back pain ( $p=0.0553$ ) were not (Table 3). In women, multiple logistic regression analysis after adjustment for age and BMI showed that walking speed ( $p=0.013$ ), knee OA ( $p=0.0218$ ), and knee pain ( $p=0.0021$ ) were risk factors for falls, whereas grip strength ( $p=0.1209$ ) and lower back pain ( $p=0.5293$ ) were not. LS was not significantly associated with falls in the crude model ( $p=0.3890$ ). To determine independent associations of walking speed, radiographic knee OA, and knee pain, we used multiple logistic regression analysis with age, BMI, walking speed, radiographic knee OA, and knee pain as independent variables and found that slower walking speed was an independent risk factor for falls in men and women ( $p<0.0001$  and  $p=0.0104$ , respectively). Knee pain was an independent risk factor for falls in women ( $p=0.0305$ ), but not in men ( $p=0.0632$ ).

**Table 1** Characteristics of participants

	Overall	Men	Women
Number of subjects	2,215	745	1,470
Age (years)	68.5 $\pm$ 11.3	69.4 $\pm$ 11.1	68.1 $\pm$ 11.4*
Height (cm)	154.7 $\pm$ 8.8	163.2 $\pm$ 6.6	150.4 $\pm$ 6.3*
Weight (kg)	55.5 $\pm$ 10.2	62.2 $\pm$ 9.9	52.0 $\pm$ 8.5*
BMI (kg/m <sup>2</sup> )	23.1 $\pm$ 3.3	23.3 $\pm$ 3.0	23.0 $\pm$ 3.4*
Grip strength (kg)	26.3 $\pm$ 9.3	34.5 $\pm$ 8.8	22.1 $\pm$ 6.2*
Walking speed (m/s)	1.24 $\pm$ 0.34	1.26 $\pm$ 0.35	1.23 $\pm$ 0.33*
Radiographic knee OA (%)	15.8	9.1	19.1**
Radiographic LS (%)	43.7	42.6	44.2
Knee pain (%)	20.8	13.7	24.4**
Lower back pain (%)	18.7	16.8	19.7

Values are presented as the mean  $\pm$  SD, except where indicated

BMI body mass index, OA osteoarthritis

\* $p<0.05$  vs. men by nonpaired Student's *t* test; \*\* $p<0.05$  vs. men by chi-square test

## Discussion

The present study is a large-scale, population-based cohort study regarding the incidence of falls and their association with physical performance and radiographic knee OA and LS as well as pain in Japanese men and women. We found that slower walking speed was a risk factor for falls in men and women and knee pain was a risk factor for falls in women only.

The present population-based longitudinal study determined whether radiographic knee OA is a risk factor for falls in Japanese men and women. Jones et al. showed that individuals with self-reported arthritis had an increased tendency to fall [8]. In the present study, after adjustment for age and BMI, radiographic knee OA was a risk factor for falls in women, but not in men. The sex differences identified in the association between radiographic knee OA and falls may be partly explained by the weaker quadriceps




Fig. 1 Incidence rate of falls (95 % CI) by gender and age

muscles and increased postural sway associated with knee OA [8, 40], both of which are known to be independent risk factors for falls [7, 41]. In men, muscle strength is higher than that in women in all decades [42], which may obscure the association between radiographic knee OA and falls. LS was not a risk factor for falls in this study. Thus, falls may be more strongly associated with problems of the lower limbs rather than the trunk.

After adjustment for age, BMI, walking speed, and radiographic knee OA, knee pain was independently associated with the incidence of falls in women. Given that the significant association of radiographic knee OA with falls disappeared after adjustment, falls may occur due to symptoms such as pain caused by radiographic knee OA rather than radiographic changes in the knee itself. Our study and other previous cross-sectional studies also suggested that knee pain was significantly associated with falls [6, 24]. In addition, a prospective study also showed that knee pain increases in falls risk in Tasmanian men and women [10]. Jones et al. showed that, for the hand, the presence of pain is what weakens grip strength [43]. In a similar way, knee pain may weaken leg strength, leading to falls. In other words,

falls may be preventable when pain is relieved by medical care, even if subjects have radiographic knee OA.


In the present study, after adjustment for knee OA and knee pain, slower walking speed was an independent risk factor for falls in men and women. Verghese et al. also showed that risk for falls increased to approximately 7 % as walking speed decreased per 0.1 m/s [44], although bone and joint diseases were not included and men and women were not separately analyzed in the study. In the present study, multiple logistic regression analysis after adjustment for knee OA and knee pain showed that, as walking speed decreased per 0.1 m/s, the risk for falls were 15 and 5 % higher in men and women, respectively, indicating that slower walking speed may more strongly affect the risk of falls in men than women. Although dependent on the availability of equipment, quantitative gait measures can be easily and quickly collected in clinical and research settings without requiring attachment of monitoring devices or extensive training. The present study may indicate that walking speed is a simple and quick option for measuring fall risk, particularly in men.

The present study has several limitations. First, our subjects lived in the community, and thus, our findings may not

Table 2 Comparison of characteristics among nonfallers and fallers in men and women

	Men			Women		
	Nonfallers	Fallers	<i>p</i> value	Nonfallers	Fallers	<i>p</i> value
Number of subjects	604	141		1,108	362	
Age (years)	68.9±11.2	71.8±10.2	0.003	67.3±11.4	70.3±10.8	<0.001
Height (cm)	163.3±6.9	162.6±5.4	0.18	150.8±6.2	149.0±6.5	<0.001
Weight (kg)	62.2±10.0	62.1±9.8	0.92	52.1±8.6	51.7±8.2	0.34
BMI (kg/m <sup>2</sup> )	23.3±3.0	23.5±3.3	0.51	22.9±3.4	23.3±3.4	0.06
Grip strength (kg)	34.8±8.9	33.0±8.2	0.02	22.4±6.2	21.1±6.1	<0.001
Walking speed (m/s)	1.30±0.36	1.11±0.28	<0.001	1.25±0.33	1.15±0.33	<0.001

Values are presented as the mean ± SD, except where indicated. Nonpaired Student's *t* test was used to determine the differences in age, height, weight, BMI, grip strength, and walking speed between nonfallers and fallers  
BMI body mass index



**Fig. 2** Incidence of falls by knee OA, knee pain, LS, and lower back pain. \* $p < 0.05$  vs. subjects without knee OA, LS, knee pain, and lower back pain, respectively, by chi-square test

apply to elderly persons residing in institutions. Second, we did not include other weight-bearing OAs such as hip OA in the analysis, although this disorder also affect falls [45]. However, the prevalence of KL=3 or 4 hip OA is 1.4 and 3.5 % in Japanese men and women [46], respectively, which is smaller than that of KL=3 or 4 knee OA in the present

study. Thus, it is possible that hip OA would not strongly affect the results of the present study.

In conclusion, the present longitudinal analysis using a large-scale population from the ROAD study revealed the incidence and risk factors for falls in men and women. Slower walking speed was a risk factor for falls in men

**Table 3** Association of physical performance and bone and joint diseases with the incidence of falls in men and women

	Men			Women		
	Crude OR (95 % CI)	Adjusted OR <sub>1</sub> (95 % CI)	Adjusted OR <sub>2</sub> (95 % CI)	Crude OR (95 % CI)	Adjusted OR <sub>1</sub> (95 % CI)	Adjusted OR <sub>2</sub> (95 % CI)
Grip strength (5-kg decrease)	1.14 (1.02–1.27)	1.05 (0.92–1.20)	–	1.20 (1.09–1.33)	1.10 (0.98–1.25)	–
Walking speed (0.1-m/s decrease)	1.19 (1.11–1.25)	1.16 (1.10–1.25)	1.15 (1.09–1.23)	1.10 (1.05–1.14)	1.06 (1.02–1.11)	1.05 (1.01–1.10)
Radiographic knee OA	1.76 (0.98–3.06)	1.52 (0.83–2.67)	1.12 (0.59–2.08)	1.69 (1.27–2.24)	1.43 (1.05–1.93)	1.21 (0.87–1.66)
Knee pain	2.12 (1.31–3.36)	1.99 (1.22–3.18)	1.63 (0.96–2.70)	1.71 (1.31–2.22)	1.54 (1.17–2.02)	1.38 (1.03–1.84)
LS	1.19 (0.83–1.73)	1.04 (0.71–1.52)	–	0.90 (0.71–1.14)	0.74 (0.57–0.94)	–
Low back pain	1.61 (1.02–2.51)	1.59 (0.99–2.49)	–	0.95 (0.79–1.27)	0.91 (0.67–1.23)	–

Multiple logistic regression analysis was used to calculate the odds ratio (OR) and 95 % confidence interval (CI) compared with nonfallers. Adjusted OR<sub>1</sub> was calculated using multiple logistic regression analysis after adjustment for age and BMI. Adjusted OR<sub>2</sub> was calculated using multiple logistic regression analysis with age, BMI, walking speed, radiographic knee OA, and knee pain as independent variables. Radiographic knee OA and LS were defined as KL grade 3 or 4

OA osteoarthritis

and women. Knee pain was a risk factor for falls in women. Further studies, along with continued longitudinal surveys in the ROAD study, will help elucidate the background of knee OA and LS and their relationship with falls.

**Acknowledgments** This work was supported by Grants-in-Aid for Scientific Research (S19109007, B20390182, C20591737, and C20591774), for Young Scientists (A18689031), and for Exploratory Research (19659305) from the Japanese Ministry of Education, Culture, Sports, Science and Technology; H17-Men-eki-009, H18-Chouju-037, H20-Chouju-009, H21-Chouju-Wakate-011, and H22-Chouju-Wakate-007 from the Ministry of Health, Labor and Welfare; Research Aid from the Japanese Orthopaedic Association (JOA-Subsidized Science Project Research 2006-1); and Grant No. 166 from the Japan Orthopaedics and Traumatology Foundation. The authors thank Dr. Yamamoto, Dr. Ishibashi, Dr. Anamizu, and members of the Department of Orthopaedics and Mr. Kutsuma and other members of the Department of Radiology at Tokyo Metropolitan Geriatric Medical Center; Mrs. Tomoko Takijiri and other members of the Public Office in Hidakagawa Town; and Mrs. Tamako Tsutsumi, Mrs. Kanami Maeda, and other members of the Public Office in Taiji Town for their assistance in the location and scheduling of participants for examinations.


**Conflicts of interest** None.

## References

- Baker S, O'Neill B, Karpf RS (1984) The injury fact book. Lexington Books, Lexington, Mass
- Fife D, Barancik JI CMS (1984) Northeastern Ohio Trauma Study: II. Injury rates by age, sex and cause. *Am J Public Health* 74:473–478
- Ministry of Health, Labour and Welfare. The outline of the results of National Livelihood Survey 2007. Available at <http://www.mhlw.go.jp/toukei/list/20-19-1.html>
- Dargent-Molina P, Favier F, Grandjean H, Baudoin C, Schott AM, Hausherr E, Meunier PJ, Breart G (1996) Fall-related factors and risk of hip fracture: the EPIDOS prospective study. *Lancet* 348:145–149
- Tinetti ME, Speechley M, Ginter SF (1988) Risk factors for falls among elderly persons living in the community. *N Engl J Med* 319:1701–1707
- Arden NK, Crozier S, Smith H, Anderson F, Edwards C, Raphael H, Cooper C (2006) Knee pain, knee osteoarthritis, and the risk of fracture. *Arthritis Rheum* 55:610–615
- Nevitt MC, Cummings SR, Kidd S, Black D (1989) Risk factors for recurrent nonsyncopal falls. A prospective study. *JAMA* 261:2663–2668
- Jones G, Nguyen T, Sambrook PN, Lord SR, Kelly PJ, Eisman JA (1995) Osteoarthritis, bone density, postural stability, and osteoporotic fractures: a population based study. *J Rheumatol* 22:921–925
- Foley SJ, Lord SR, Srikanth V, Cooley H, Jones G (2006) Falls risk is associated with pain and dysfunction but not radiographic osteoarthritis in older adults: Tasmanian Older Adult Cohort study. *Osteoarthr Cartil* 14:533–539
- Scott D, Blizzard L, Fell J, Jones G (2012) Prospective study of self-reported pain, radiographic osteoarthritis, sarcopenia progression, and falls risk in community-dwelling older adults. *Arthritis Care Res (Hoboken)* 64:30–37
- Jackson DW, Simon TM, Aberman HM (2001) Symptomatic articular cartilage degeneration: the impact in the new millennium. *Clin Orthop Relat Res* 391(Suppl):S14–S25
- Reginster JY (2002) The prevalence and burden of arthritis. *Rheumatology (Oxford)* 41(Suppl 1):S3–S6
- Buckwalter JA, Saltzman C, Brown T (2004) The impact of osteoarthritis: implications for research. *Clin Orthop Relat Res* 427(Suppl):S6–S15
- Sharma L, Kapoor D (2007) Epidemiology of osteoarthritis. In: Moskowitz RW, Altman RD, Hochberg MC, Buckwalter JA, Goldberg VM (eds) *Osteoarthritis: diagnosis and medical/surgical management*, 4th edn. Lippincott Williams & Wilkins, Philadelphia, pp 3–26
- Hadjipavlou AG, Simmons JW, Pope MH, Necessary JT, Goel VK (1999) Pathomechanics and clinical relevance of disc degeneration and annular tear: a point-of-view review. *Am J Orthop* 28:561–571
- Emery SE, Ringus VM (2007) Osteoarthritis of the spine. In: Moskowitz RW, Altman RD, Hochberg MC, Buckwalter JA, Goldberg VM (eds) *Osteoarthritis: diagnosis and medical/surgical management*, 4th edn. Lippincott Williams & Wilkins, Philadelphia, pp 427–452
- Muraki S, Oka H, Akune T, Mabuchi A, En-yo Y, Yoshida M, Saika A, Suzuki T, Yoshida H, Ishibashi H, Yamamoto S, Nakamura K, Kawaguchi H, Yoshimura N (2009) Prevalence of radiographic knee osteoarthritis and its association with knee pain in the elderly of Japanese population-based cohorts: the ROAD study. *Osteoarthr Cartil* 17:1137–1143
- Muraki S, Oka H, Mabuchi A, Akune T, En-yo Y, Yoshida M, Saika A, Suzuki T, Yoshida H, Ishibashi H, Yamamoto S, Nakamura K, Kawaguchi H, Yoshimura N (2009) Prevalence of radiographic lumbar spondylosis and its association with low back pain in the elderly of population-based cohorts: the ROAD study. *Ann Rheum Dis* 68:1401–1406
- Yoshimura N, Muraki S, Oka H, Mabuchi A, En-yo Y, Yoshida M, Saika A, Yoshida H, Suzuki T, Yamamoto S, Ishibashi H, Kawaguchi H, Nakamura K, Akune T (2009) Prevalence of knee osteoarthritis, lumbar spondylosis and osteoporosis in Japanese men and women: the Research on Osteoarthritis/Osteoporosis Against Disability (ROAD). *J Bone Miner Metab* 27:620–628
- Linaker CH, Walker-Bone K, Palmer K, Cooper C (1999) Frequency and impact of regional musculoskeletal disorders. *Baillieres Clin Rheumatol* 13:197–215
- Summers MN, Haley WE, Reveille JD, Alarcon GS (1988) Radiographic assessment and psychologic variables as predictors of pain and functional impairment in osteoarthritis of the knee or hip. *Arthritis Rheum* 31:204–209
- Cicutini FM, Baker J, Hart DJ, Spector TD (1996) Association of pain with radiological changes in different compartments and views of the knee joint. *Osteoarthr Cartil* 4:143–147
- Wluka AE, Wolfe R, Stuckey S, Cicutini FM (2004) How does tibial cartilage volume relate to symptoms in subjects with knee osteoarthritis? *Ann Rheum Dis* 63:264–268
- Muraki S, Akune T, Oka H, En-yo Y, Yoshida M, Nakamura K, Kawaguchi H, Yoshimura N (2011) Prevalence of falls and its association with knee osteoarthritis and lumbar spondylosis as well as knee and lower back pain in Japanese men and women. *Arthritis Care Res* 63:1425–1431
- Ganz DA, Bao Y, Shekelle PG, Rubenstein LZ (2007) Will my patient fall? *JAMA* 297:77–86
- Studenski S, Perera S, Wallace D, Chandler JM, Duncan PW, Rooney E, Fox M, Guralnik JM (2003) Physical performance measures in the clinical setting. *J Am Geriatr Soc* 51:314–322
- Cesari M, Kritchevsky SB, Penninx BW, Nicklas BJ, Simonsick EM, Newman AB, Tylavsky FA, Brach JS, Satterfield S, Bauer DC, Visser M, Rubin SM, Harris TB, Pahor M (2005) Prognostic value of usual gait speed in well-functioning older people—results from the Health, Aging and Body Composition Study. *J Am Geriatr Soc* 53:1675–1680

28. Campbell AJ, Borrie MJ, Spears GF (1989) Risk factors for falls in a community-based prospective study of people 70 years and older. *J Gerontol* 44:M112–M117
29. Lipsitz LA, Jonsson PV, Kelley MM, Koestner JS (1991) Causes and correlates of recurrent falls in ambulatory frail elderly. *J Gerontol* 46:M114–M122
30. Wolfson L, Whipple R, Amerman P, Tobin JN (1990) Gait assessment in the elderly: a gait abnormality rating scale and its relation to falls. *J Gerontol* 45:M12–M19
31. Luukinen H, Koski K, Laippala P, Kivela SL (1995) Predictors for recurrent falls among the home-dwelling elderly. *Scand J Prim Health Care* 13:294–299
32. Chan BK, Marshall LM, Winters KM, Faulkner KA, Schwartz AV, Orwoll ES (2007) Incident fall risk and physical activity and physical performance among older men: the Osteoporotic Fractures in Men Study. *Am J Epidemiol* 165:696–703
33. Yoshimura N, Muraki S, Oka H, Kawaguchi H, Nakamura K, Akune T (2010) Cohort profile: Research on Osteoarthritis/osteoporosis Against Disability (ROAD) Study. *Int J Epidemiol* 39:988–995
34. Shimada H, Lord SR, Yoshida H, Kim H, Suzuki T (2007) Predictors of cessation of regular leisure-time physical activity in community-dwelling elderly people. *Gerontology* 53:293–297
35. Tinetti M, Baker D, Dutcher J (1997) Reducing the risk of falls among older adults in the community. Peacable Kingdom Press, Berkeley
36. Kellgren JH, Lawrence JS (eds) (1963) The epidemiology of chronic rheumatism: atlas of standard radiographs of arthritis. Blackwell Scientific, Oxford
37. Judge JO, Davis RB 3rd, Ounpuu S (1996) Step length reductions in advanced age: the role of ankle and hip kinetics. *J Gerontol A Biol Sci Med Sci* 51:M303–M312
38. Judge JO, Lindsey C, Underwood M, Winsemius D (1993) Balance improvements in older women: effects of exercise training. *Phys Ther* 73:254–264
39. Steffan TM, Hacker TA, Mollinger L (2002) Age- and gender-related test performance in community-dwelling older people: six-minute walk test, Berg balance scale, timed up and go test, and gait speeds. *Phys Ther* 82:128–137
40. Wegener L, Kisner C, Nichols D (1997) Static and dynamic balance responses in persons with bilateral knee osteoarthritis. *J Orthop Sports Phys Ther* 25:13–18
41. Campbell AJ, Borrie MJ, Spears GF (1989) Risk factors for falls in a community-based prospective study of people 70 years and older. *J Gerontol* 44:M112–M117
42. Sinaki M, Nwaogwugwu NC, Phillips BE (2001) Mokri MP (2001) Effect of gender, age, and anthropometry on axial and appendicular muscle strength. *Am J Phys Med Rehabil* 80:330–338
43. Jones G, Cooley HM, Bellamy N (2001) A cross-sectional study of the association between Heberden's nodes, radiographic osteoarthritis of the hands, grip strength, disability and pain. *Osteoarthritis Cartil* 9:606–611
44. Verghese J, Holtzer R, Lipton RB, Wang C (2009) Quantitative gait markers and incident fall risk in older adults. *J Gerontol* 64:896–901
45. Arden NK, Nevitt MC, Lane NE, Gore LR, Hochberg MC, Scott JC, Pressman AR, Cummings SR (1999) Osteoarthritis and risk of falls, rates of bone loss, and osteoporotic fractures. Study of Osteoporotic Fractures Research Group. *Arthritis Rheum* 42:1378–1385
46. Inoue K, Wicart P, Kawasaki T, Huang J, Ushiyama T, Hukuda S (2000) Prevalence of hip osteoarthritis and acetabular dysplasia in French and Japanese adults. *Rheumatology (Oxford)* 39:745–748

Provided for non-commercial research and education use.  
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at SciVerse ScienceDirect

Bone

journal homepage: [www.elsevier.com/locate/bone](http://www.elsevier.com/locate/bone)

Original Full Length Article

## Risk factors for falls in a longitudinal population-based cohort study of Japanese men and women: The ROAD Study

Shigeyuki Muraki <sup>a,\*</sup>, Toru Akune <sup>a</sup>, Yuyu Ishimoto <sup>b</sup>, Keiji Nagata <sup>b</sup>, Munehito Yoshida <sup>b</sup>, Sakae Tanaka <sup>c</sup>, Hiroyuki Oka <sup>d</sup>, Hiroshi Kawaguchi <sup>c</sup>, Kozo Nakamura <sup>e</sup>, Noriko Yoshimura <sup>d</sup>

<sup>a</sup> Department of Clinical Motor System Medicine, 22nd Century Medical and Research Center, Faculty of Medicine, the University of Tokyo, 7-3-1, Hongo, Bunkyo-ku, Tokyo 113-8655, Japan

<sup>b</sup> Department of Orthopaedic Surgery, Wakayama Medical University, 811, Kimiidera, Wakayama-shi, Wakayama 641-8509, Japan

<sup>c</sup> Department of Sensory and Motor System Medicine, Faculty of Medicine, the University of Tokyo, 7-3-1, Hongo, Bunkyo-ku, Tokyo 113-8655, Japan

<sup>d</sup> Department of Joint Disease Research, 22nd Century Medical and Research Center, Faculty of Medicine, the University of Tokyo, 7-3-1, Hongo, Bunkyo-ku, Tokyo 113-8655, Japan

<sup>e</sup> Rehabilitation Services Bureau, National Rehabilitation Center for Persons with Disabilities, 4-1, Namiki, Tokorozawa-shi, Saitama 359-8555, Japan

### ARTICLE INFO

#### Article history:

Received 20 April 2012

Revised 17 October 2012

Accepted 19 October 2012

Available online 24 October 2012

Edited by: Toshio Matsumoto

#### Keywords:

Falls

Longitudinal study

Pain

Osteoarthritis

Physical performance

### ABSTRACT

The objective of this study was to clarify the associations of physical performance and bone and joint diseases with single and multiple falls in Japanese men and women using a population-based longitudinal cohort study known as Research on Osteoarthritis/osteoporosis Against Disability (ROAD). A total of 452 men and 896 women were analyzed in the present study (mean age, 63.9 years). A questionnaire was used to assess the number of falls during the 3-year follow-up. Grip strength, 6-m walking time, and chair stand time were measured at baseline. Knee osteoarthritis (OA) and lumbar spondylosis were defined as Kellgren Lawrence = 2, 3 or 4. Vertebral fracture (Vfx) was assessed with the Japanese Society of Bone and Mineral Research criteria. Osteoporosis was defined by bone mineral density using dual energy X-ray absorptiometry based on World Health Organization criteria. Knee and lower back pain were estimated by an interview. During a 3-year follow-up, 79 (17.4%) men and 216 (24.1%) women reported at least one fall, and 54 (11.9%) men and 111 (12.4%) women reported multiple falls. Knee pain was a risk factor for multiple falls in women, but not in men. Vfx tended to be associated with multiple falls in women, but not in men. A longer 6-m walking time was a risk factor for multiple falls in women, whereas a longer chair stand time was a risk factor for multiple falls in men. We found gender differences in risk factors for falls.

© 2012 Elsevier Inc. All rights reserved.

### Introduction

Falls are one of the main causes of injury, disability, and death among the elderly [1,2]. In Japan, according to the recent National Livelihood Survey of the Ministry of Health, Labour and Welfare, falls and fractures are ranked fifth among diseases that cause disabilities and subsequently require support with activities of daily living [3]. However, few population-based studies have been performed on the incidence of falls based on sex and age. Furthermore, in terms of factors associated with falls, muscle strength, balance, vision, functional capacities, and cognitive impairment are traits that diminish with aging, and these factors have been suggested as predictive risk factors for falls and fractures [4,5]. However, the association of bone and joint diseases, especially osteoarthritis (OA), with falls remains unclear.

The representative sites of OA are the knee and lumbar spine. Knee OA and lumbar spondylosis (LS) are major public health issues because

they cause chronic pain and disability [6,7]. The prevalence rates of radiographic knee OA and LS are 54.6% and 70.2%, respectively, in persons aged 40 years and older in Japan, which indicates that 25,300,000 and 37,900,000 persons aged 40 years and older are estimated to experience radiographic knee OA and LS, respectively [10]. The National Livelihood Survey ranked OA fourth among diseases that cause disabilities and subsequently require support with activities of daily living [3], but there have been few studies of the association between falls and OA [11,12]. In previous studies, knee OA was assessed only by interview and not by radiography. The principal clinical symptom of knee OA is pain [13], but its correlation with the radiographic severity of knee OA is not as strong as expected [8]. In fact, in a study in Japan, approximately 20% of persons without knee OA had knee pain, and 30% of persons with severe knee OA had no knee pain [8]. Thus, knee OA diagnosed by interview could be limited by variable accuracy. In addition, men and women were not examined separately in these previous studies, although sex differences have been found in the prevalence of knee OA [8]. Our previous study showed that knee pain is significantly associated with falls in women [14], but that study used a cross-sectional design; thus, a causal relationship remains unclear. Regarding LS, to the best of our knowledge, no population-based studies have been performed

\* Corresponding author at: Department of Clinical Motor System Medicine, 22nd Century Medical and Research Center, Faculty of Medicine, University of Tokyo, 7-3-1, Hongo, Bunkyo-ku, Tokyo 113-8655, Japan. Fax: +81 3 5800 9179.

E-mail address: [murakis-ort@h.u-tokyo.ac.jp](mailto:murakis-ort@h.u-tokyo.ac.jp) (S. Muraki).



regarding its association with falls except for our previous cross-sectional study [14], which showed that LS is not significantly associated with falls. In addition, among fractures due to osteoporosis (OP), vertebral fracture (VFX) is the most likely to lead to marked public health problems. VFX is reportedly associated with functional impairment [15], back pain, kyphosis [16,17], esophageal reflux [18], depressive mood [19], respiratory dysfunctions [20], and mortality [21]. However, whether VFX is an independent risk factor for the incidence of falls remains unclear.

Measuring walking speed is a simple way to assess health and function in older adults [22,23]. Walking speed has been found to be associated with falls in a few studies [4,24–26], although most studies were limited by a small sample size, a cross-sectional design [24,25], or evaluation of a single sex [4,26]. In addition, although walking abnormalities indicative by a slower walking speed are significantly associated with bone and joint diseases such as knee OA, LS, and their associated pain [14], no longitudinal studies have been performed to determine the associations of falls with bone and joint diseases and walking abnormalities at the same time. Furthermore, measuring the chair stand time is also reported to be a simple and established method to assess health and function in the elderly [27,28], but to the best of our knowledge, no longitudinal studies have been performed to determine the associations of falls with chair stand time.

Previous studies have shown that associations between individual risk factors and a single fall are few in number and weak compared to risk factors for multiple falls [12], indicating that single and multiple falls may have different backgrounds. Thus, to determine factors associated with falls, single and multiple falls should be analyzed separately.

The objective of this study was to clarify the associations of physical performance and bone and joint diseases with the incidence of single and multiple falls in Japanese men and women using a population-based longitudinal cohort study known as Research on Osteoarthritis/osteoporosis Against Disability (ROAD).

## Methods

### Participants

The ROAD study is a nationwide, prospective study designed to establish epidemiologic indices for evaluation of clinical evidence for the development of a disease-modifying treatment for bone and joint diseases (OP and OA are the representative bone and joint diseases, respectively). ROAD consists of population-based cohorts in three communities in Japan. A detailed profile of the ROAD study has been described elsewhere [8–10,29]; a brief summary is provided here. To date, we have completed the creation of a baseline database that includes clinical and genetic information for 3,040 participants (1,061 men and 1,979 women) ranging in age from 23 to 95 years (mean, 70.6 years) who were recruited from resident registration listings in three communities: an urban region in Itabashi, Tokyo; a mountainous region in Hidakagawa, Wakayama; and a coastal region in Taiji, Wakayama.

Residents of these regions were recruited from the resident registration list of the relevant region. Participants in the urban region were recruited from a randomly selected cohort from the Itabashi-ward residents' registration database [30]. The participation rate was 75.6%. Participants in mountainous and coastal regions were also recruited from the resident registration lists, and the participation rates in these two areas were 56.7% and 31.7%, respectively. The inclusion criteria, apart from residence in the communities mentioned above, were the ability to (1) walk to the survey site, (2) report data, and (3) understand and sign an informed consent form. The baseline survey of the ROAD study was completed in 2006. All participants provided written informed consent, and the study was conducted with the approval of the ethics committees of the University of Tokyo and the Tokyo Metropolitan Institute of Gerontology.

### Assessment of falls

Three years after the baseline data were obtained, we attempted to trace and review all 3,040 participants between 2008 and 2010; they were invited to attend a follow-up interview. All participants were interviewed with regard to falls by experienced interviewers and were asked the following questions: "Have you experienced falls during the 3-year follow-up, and if yes, how many falls did you experience"? At baseline, all participants were also interviewed regarding falls by experienced interviewers and were asked the following questions: "Have you experienced falls during the 12 months preceding baseline, and if yes, how many falls did you experience"? According to a previous study on falls [31], a fall is defined as a sudden, unintentional change in position causing an individual to land at a lower level on an object, the floor, or the ground, other than as a consequence of a sudden onset of paralysis, epileptic seizure, or overwhelming external force.

### Pain assessment

All participants were interviewed by experienced orthopedists regarding knee pain and lower back pain at baseline and were asked the following questions based on previous studies [8,9]: "Have you experienced knee pain on most days in the past month, in addition to now?" and "Have you experienced lower back pain on most days in the past month, in addition to now"? Those who answered "yes" were defined as having pain. Buttock pain and sciatica were not included as lower back pain in the present study.

### Radiographic assessment

At baseline, all participants underwent radiographic examination of both knees using anteroposterior and lateral views with weight-bearing and foot-map positioning; radiographic examination of the anteroposterior and lateral views of the lumbar spine, including intervertebral levels L1/2 to L5/S, was also performed. VFX was assessed by lateral radiographs of the lumbar spine (L1–L5) in terms of a wedge, biconcave, or crush appearance according to the Japanese Society of Bone and Mineral Research criteria [32]. The films were marked up, and morphometric measurements of anterior, middle, and posterior heights on lateral radiography of the thoracic and lumbar spine were made. Wedge appearance was defined as a site at which the anterior height of the vertebra was  $\leq 75\%$  of the posterior height. Biconcave appearance occurred if the height of the central part of the vertebra was  $\leq 80\%$  of that of the anterior or posterior parts of the vertebra. Crush appearance occurred if the height of the anterior, central, and posterior parts of an axial vertebra were all reduced to  $\leq 80\%$  of the normal value (Supplementary Fig. 1). Knee and lumbar spine radiographs were also read without knowledge of the participant's clinical status by a single, experienced orthopedist (S.M.) using the Kellgren Lawrence (KL) radiographic atlas [33] to determine the severity of KL grading. Radiographs were scored as grade 0–4, with higher grades associated with more severe OA. We defined knee OA and LS as KL  $\geq 2$  in at least one knee and one intervertebral level, respectively. To evaluate the intraobserver variability of KL grading, 100 randomly selected radiographs of the knee and lumbar spine were scored by the same observer more than 1 month after the first reading. One hundred other radiographs were also scored by two experienced orthopedic surgeons (S.M. and H.O.) using the same atlas for interobserver variability. The intra- and interobserver variabilities evaluated were confirmed by kappa analysis to be sufficient for assessment (0.86 and 0.80 for knee OA, and 0.84 and 0.76 for LS, respectively).

### Bone mineral density (BMD) measurement

BMD was measured at the lumbar spine (L2–4) and the proximal femur using dual energy X-ray absorptiometry (DXA) (Hologic

Discovery; Hologic, Waltham, MA, USA) at baseline. For quality control, the same DXA equipment was used, and the same spine phantom was scanned daily to monitor the machine's performance in study populations at different regions. The BMD of the phantom was adjusted to  $1.032 \pm 0.016 \text{ g/cm}^2$  ( $\pm 1.5\%$ ) during all examinations. In addition, the same physician (N.Y.) examined all participants to prevent observer variability. Coefficient of variance (CV) for L2–L4 in the phantom was 0.35%, and CVs for L2–L4, the proximal femur, Ward's triangle, and the trochanter examined in five volunteers were 0.61–0.90, 1.02–2.57, 1.97–5.45, and 1.77–4.17%, respectively [34].

OP was defined based on World Health Organization (WHO) criteria in which OP was diagnosed as T-scores of BMD  $\leq 2.5$  standard deviations (SDs) lower than peak bone mass [35]. Mean L2–4 BMD (SD) for young adult men and women measured using the Hologic QDR devices in Japan is reportedly  $1.011 \text{ g/cm}^2$  ( $0.119 \text{ g/cm}^2$ ) [36]. Mean femoral neck BMD (SD) in Japan is reported to be  $0.863 \text{ g/cm}^2$  ( $0.127 \text{ g/cm}^2$ ) for young men and  $0.787$  ( $0.109$ ) for young women [36]. The present study therefore defined OP using these indices as lumbar spine BMD  $< 0.714 \text{ g/cm}^2$  for both men and women, and as femoral neck BMD  $< 0.546 \text{ g/cm}^2$  for men and  $< 0.515 \text{ g/cm}^2$  for women.

#### Physical performance

At baseline, anthropometric measurements were taken, including height and weight, and body mass index (BMI) ( $\text{weight [kg]/height}^2 \text{ [m}^2\text{]}$ ) was estimated based on the measured height and weight. Grip strength was measured on bilateral sides using a TOEI LIGHT handgrip dynamometer (TOEI LIGHT CO., LTD, Saitama, Japan), and the best measurement was used to characterize maximum muscle strength. To measure physical performance, the time taken to walk 6 m at normal walking speed in a hallway was recorded. Participants were told to walk from a marked starting line to a 6-m mark as if they were walking down their hallway at home. Time was measured in seconds with a stopwatch and rounded to the nearest hundredth of a second. The average of two trials was recorded. These gait-speed trial measurements are considered highly reliable in community-dwelling elderly persons [37]. The time taken for five consecutive chair rises without the use of hands was also recorded. Hands were folded in front of the chest with feet flat on the floor, following the protocol described by Guralnik et al. [27] and used by other researchers [28]. Time was measured in seconds with a stopwatch and rounded to the nearest hundredth of a second. Timing began with the command "Go" and ended when the buttocks contacted the chair on the fifth landing. The reliability of this protocol is adequate [27].

#### Cognition assessment

At baseline, cognition was also evaluated for all participants using a Mini-Mental State Examination, and a cut-off score of  $< 24$  was used to select participants with cognitive impairment [38].

#### Statistical analyses

The differences in age and anthropometric measurements between the responders (those who completed the study) and non-responders (those lost to follow-up or who did not complete the study as described below) and between men and women were examined with a non-paired Student's *t*-test. Differences in physical performance measurements between the responders and non-responders and between men and women were examined with Wilcoxon signed-rank test. Differences in age and anthropometric measurements, among non-fallers, single fallers, and multiple fallers, were examined with one-way analysis of variance. Differences in physical performance measurements among non-fallers, single fallers, and multiple fallers were examined with the Kruskal–Wallis test. The prevalence of bone and joint diseases and cognitive impairment was compared between men

and women and among non-fallers, single fallers, and multiple fallers with the chi square test. Multinomial logistic regression analysis after adjusting for age and BMI was used to determine the association of anthropometric measurements, physical performance, bone and joint diseases, and cognitive impairment with single and multiple falls compared with the absence of falls in men and women. Further, to determine an independent association of physical performance with single and multiple falls compared with the absence of falls, we used multinomial logistic regression analysis with age, BMI, 6-m walking time, and chair stand time as explanatory variables. To determine independent risk factors for single and multiple falls, we used multinomial logistic regression analysis with age, BMI, physical performance, bone and joint diseases, and cognitive impairment as explanatory variables. Data analyses were performed using SAS version 9.0 (SAS Institute Inc., Cary, NC, USA).

#### Results

Of the 1,690 participants in the mountainous and seaside cohorts at baseline in 2006 and 2007, 40 (2.4%) had died by the time of the review 3 years later, 97 (5.7%) did not participate in the follow-up study due to poor health, 16 (0.9%) had moved away, 51 (3.0%) declined the invitation to attend the follow-up study, and 47 (2.8%) did not participate in the follow-up study for other reasons. Among the 1,439 volunteers who did participate in the follow-up study, 68 (4.0%) provided incomplete fall questionnaires. In addition, six (0.4%) provided incomplete pain questionnaires; these were excluded. We also excluded eight (0.5%) participants who had undergone total knee arthroplasty before baseline. An additional nine (1.9%) participants did not perform the 6-m walking time or chair stand time, leaving a total of 1,348 (79.8%) participants (452 men and 896 women) from whom radiographs at baseline and complete fall and pain histories were obtained. The mean followup time was  $2.93 \pm 0.12$  years, ranging from 2.65 to 3.22 years. Table 1 shows characteristics of responders and non-responders. The responders were significantly younger than the non-responders (63.9 and 70.7 years, respectively). Physical performance measurements were better in responders than non-responders. Prevalence of knee OA, LS and knee pain was lower in responders (47.0, 61.6 and 9.7%,

**Table 1**  
Baseline characteristics of responders and non-responders.

	Overall	Responders	Non-responders
Number of participants	1,690	1,348	342
Female (%)	64.7	66.5	57.9***
Age (years)	$65.2 \pm 12.0$	$63.9 \pm 11.8$	$70.7 \pm 11.4^*$
Height (cm)	$155.2 \pm 9.3$	$155.6 \pm 9.0$	$153.6 \pm 10.1^*$
Weight (kg)	$55.6 \pm 10.8$	$56.1 \pm 10.7$	$53.7 \pm 10.8^*$
BMI ( $\text{kg/m}^2$ )	$23.0 \pm 3.4$	$23.1 \pm 3.4$	$22.7 \pm 3.4$
Grip strength (kg) (median [IQR])	$26.0 [21.0–33.0]$	$26.0 [21.0–34.0]$	$24.0 [18.0–30.0]**$
6-m walking time (s) (median [IQR])	$5.0 [4.0–7.0]$	$5.0 [4.0–6.0]$	$7.0 [5.0–9.0]**$
Chair stand time (s) (median [IQR])	$9.0 [7.0–12.0]$	$9.0 [7.0–11.0]$	$12.0 [8.25–15.0]**$
Cognitive impairment (%)	4.5	2.8	11.4***
Radiographic knee OA (%)	50.4	47.0	63.8***
Radiographic LS (%)	63.2	61.6	69.1***
Radiographic VFX	10.1	9.7	12.0
Knee pain (%)	24.3	22.2	32.6***
Lower back pain (%)	21.1	20.6	22.9
Previous falls (%)	17.3	16.3	21.0***

Values are mean  $\pm$  SD, except where indicated.

BMI: body mass index, OA: osteoarthritis, LS: lumbar spondylosis, VFX: vertebral fracture, IQR: interquartile range.

\*  $p < 0.05$  vs. responders by non-paired Student's *t*-test.

\*\*  $p < 0.05$  vs. men by Wilcoxon signed-rank test.

\*\*\*  $p < 0.05$  vs. men by chi square test.

**Table 2**  
Baseline characteristics of participants.

	Men	Women
Number of participants	452	896
Age (years)	64.9 ± 11.7	63.3 ± 11.8*
Height (cm)	164.0 ± 7.0	151.3 ± 6.6*
Weight (kg)	63.3 ± 10.7	52.5 ± 8.7*
BMI (kg/m <sup>2</sup> )	23.5 ± 3.2	22.9 ± 3.4*
Grip strength (kg) (median [IQR])	37.0 [32.0–42.5]	23.5 [20.0–23.5]**
6-m walking time (s) (median [IQR])	5.0 [4.0–6.0]	5.0 [4.0–6.0]
Chair stand time (s) (median [IQR])	8.5 [7.0–11.0]	9.0 [7.0–11.0]
Cognitive impairment (%)	3.6	2.4
Radiographic knee OA (%)	37.4	51.9***
Radiographic LS (%)	76.1	54.2
Radiographic Vfx	8.9	10.1
Knee pain (%)	15.3	25.7***
Lower back pain (%)	18.8	21.5
Previous falls (%)	13.1	18.0***

Values are mean ± SD, except where indicated.

BMI: body mass index, OA: osteoarthritis, LS: lumbar spondylosis, Vfx: vertebral fracture, IQR: interquartile range.

\*  $p < 0.05$  vs. men by non-paired Student's *t*-test.

\*\*  $p < 0.05$  vs. men by Wilcoxon signed-rank test.

\*\*\*  $p < 0.05$  vs. men by chi square test.

respectively) than non-responders (63.8, 69.1 and 12.0, respectively). Prevalence of previous falls was significantly lower in responders than non-responders (16.3 and 21.0%, respectively).

Table 2 shows the age, anthropometric measurements, physical performance, and prevalence of cognitive impairment, bone and joint diseases, and previous falls of participants at baseline in men and women. Regarding physical performance, grip strength and chair stand time were significantly better in men (37.0 kg and 8.5 s, respectively) than in women (23.5 kg and 9.0 s, respectively), but the 6-m walking time was not (5.0 s and 5.0 s, respectively). The prevalence of radiographic knee OA and knee pain was significantly higher in women (51.9% and 25.7%, respectively) than in men (37.4% and 15.3%, respectively), whereas that of LS and lower back pain was not different between men and women. The prevalence of previous falls was significantly higher in women than in men (18.0% and 13.1%, respectively).

During the 3-year follow-up, 79 (17.4% [95% confidence interval [CI] 14.3–21.2]) men and 216 (24.1% [95% CI 21.4–27.0]) women reported at least one fall, and 54 (11.9% [95% CI 9.3–15.3]) men and 111 (12.4% [95% CI 10.4–14.7]) women reported multiple falls. The chi square test showed that the incidence of falls was significantly different between men and women ( $p = 0.0011$ ). The incidence of single and multiple falls was significantly higher in the mountainous regions (11.5% and

17.4%, respectively) than coastal regions (8.1% and 7.8%, respectively). With increasing age, the incidence of falls increased in women, but the incidence of falls was similar in men in their 60s and 70s (Fig. 1).

Table 3 shows the age, anthropometric measurements, physical performance, and BMD at baseline between non-fallers, single fallers, and multiple fallers. Age and BMI were significantly higher in female fallers than non-fallers, but this was not the case in men. Grip strength was worse in female fallers than non-fallers, but this was not the case in men. The 6-m walking time and chair stand time were longer in both male and female fallers than in non-fallers. LS and neck BMD were significantly lower in female fallers than non-fallers, but this was not the case in men.

We next examined the incidence rate of falls during the 3-year follow-up according to previous falls at baseline in men and women (Supplementary Fig. 2). The incidence rates of multiple falls were 7.9%, 22.7%, and 48.7% in men and 8.8%, 20.4%, and 43.1% in women among non-fallers, single fallers, and multiple fallers, respectively. The incidence rates of single falls were 5.9%, 9.1%, and 0.0% in men and 12.5%, 7.8%, and 8.6% in women among non-fallers, single fallers, and multiple fallers, respectively. The chi square test showed that the incidence of falls during the 3-year follow-up was significantly associated with previous falls at baseline in men and women ( $p < 0.0001$ ).

Fig. 2 shows the incidence rate of falls during the 3-year follow-up according to the presence of bone and joint diseases and cognitive impairment. The incidence rates of multiple falls were 16.6% and 9.2% in men and 14.8% and 9.7% in women in those with and without knee OA, respectively. The incidence rates of a single fall were 8.3% and 3.9% in men and 14.2% and 9.1% in women in those with and without knee OA, respectively. The chi square test showed that knee OA at baseline was significantly associated with the incidence rate of falls during the 3-year follow-up in men and women ( $p < 0.0001$ ). Regarding knee pain, the incidence rates of multiple falls were 18.8% and 10.7% in men and 18.7% and 10.2% in women in those with and without knee pain, respectively. The incidence rates of a single fall were 8.7% and 5.0% in men and 10.4% and 10.4% in women in those with and without knee OA, respectively. The chi square test showed that knee pain at baseline was significantly associated with the incidence of falls during the 3-year follow-up in men and women ( $p < 0.0001$ ). LS and lower back pain were not significantly associated with the incidence of falls in men ( $p = 0.52$  and  $0.77$ , respectively) or in women ( $p = 0.45$  and  $0.58$ , respectively). Vfx at baseline was significantly associated with the incidence of falls in women (multiple falls 22.2% and 11.3%, single falls 14.4% and 11.4%, in those with and without Vfx, respectively,  $p = 0.005$ ), but not in men ( $p = 0.06$ ). OP defined by L2–4 and femoral neck BMD was not associated with the incidence of falls in men and women. Cognitive impairment




Fig. 1. Incidence rate of falls (error bars represent 95% confidence intervals) by gender and age strata.

**Table 3**  
Comparison of characteristics among non-fallers, single fallers, and multiple fallers in men and women.

	Men			p value	Women			p value
	Non-fallers	Single fallers	Multiple fallers		Non-fallers	Single fallers	Multiple fallers	
Number of participants	373	25	54		680	105	111	
Age (years)	64.4 (11.7)	67.2 (13.2)	67.6 (10.1)	0.10	62.4 (11.6)	66.0 (12.6)	66.7 (11.4)	<0.001
BMI (kg/m <sup>2</sup> )	23.4 (3.1)	24.6 (3.9)	23.7 (3.3)	0.16	22.8 (3.5)	22.7 (3.1)	23.8 (3.5)	0.01
Grip strength (kg)	37.0 (median [IQR]) [32.0–43.0]	37.0 (median [IQR]) [30.0–41.5]	35.0 (median [IQR]) [28.8–40.0]	0.08	24.0 (median [IQR]) [20.0–27.0]	23.0 (median [IQR]) [19.5–27.0]	22.0 (median [IQR]) [18.0–26.0]	0.01
6-m walking time (s)	4.5 (median [IQR]) [4.0–6.0]	5.5 (median [IQR]) [4.6–7.3]	6.2 (median [IQR]) [5.0–6.6]	<0.0001	5.0 (median [IQR]) [4.0–6.0]	5.0 (median [IQR]) [4.0–6.5]	5.5 (median [IQR]) [4.0–7.5]	<0.0001
Chair stand time (s)	8.0 (median [IQR]) [7.0–10.0]	11.0 (median [IQR]) [9.0–12.0]	10.0 (median [IQR]) [8.0–13.0]	<0.0001	9.0 (median [IQR]) [7.0–11.0]	9.0 (median [IQR]) [8.0–12.0]	10.0 (median [IQR]) [8.0–12.25]	0.0001
LS BMD	1.05 (0.20)	1.05 (0.20)	1.05 (0.15)	0.99	0.89 (0.18)	0.85 (0.16)	0.86 (0.17)	0.04
Neck BMD	0.75 (0.13)	0.77 (0.12)	0.75 (0.10)	0.79	0.65 (0.13)	0.61 (0.11)	0.63 (0.11)	0.003

Values are the means (standard deviation), except where indicated.

One-way analysis of variance was used to determine the differences in age, height, weight and BMI among non-fallers, single fallers, and multiple fallers.


Kruskal–Wallis test was used to determine the differences in grip strength, 6-m walking time and chair stand time among non-fallers, single fallers, and multiple fallers.

The chi square test was used to determine the differences in the prevalence of cognitive impairment among non-fallers, single fallers, and multiple fallers.

BMI: body mass index, LS: lumbar spondylosis, BMD: bone mineral density.

was associated with the incidence of falls in men (multiple falls 31.3% and 10.9%, single falls 18.8% and 5.1%, in those with and without cognitive impairment, respectively,  $p=0.002$ ), but not in women ( $p=0.19$ ).

In men, multinomial logistic regression analysis after adjusting for age and BMI showed that a longer 6-m walking time, longer chair stand time, and previous falls were risk factors for falls, but grip strength, bone and joint diseases, and cognitive impairment were not



**Fig. 2.** Incidence of single and multiple falls by bone and joint diseases and cognitive impairment. \* $p < 0.05$  vs. participants without each disease or pain, respectively, according to the chi square test. OA, osteoarthritis; LS, lumbar spondylosis; VFX, vertebral fracture.

**Table 4**  
Risk factors for single and multiple falls in men.

	Crude OR (95% CI)		Adjusted OR (95% CI)	
	Single falls	Multiple falls	Single falls	Multiple falls
Grip strength (5 kg increase)	0.90 (0.71–1.14)	0.84 (0.71–0.99)	1.14 (1.01–1.29)	0.88 (0.72–1.08)
6-m walking time (1 s increase)	1.12 (0.98–1.27)	1.13 (1.03–1.26)	1.11 (0.95–1.25)	1.11 (1.01–1.23)
Chair stand time (1 s increase)	1.17 (1.03–1.32)	1.21 (1.11–1.33)	1.15 (1.00–1.32)	1.21 (1.09–1.33)
LS BMD (0.1 mg/cm <sup>2</sup> increase)	1.00 (0.80–1.22)	1.00 (0.86–1.16)	0.92 (0.73–1.15)	0.97 (0.83–1.13)
Neck BMD (0.1 mg/cm <sup>2</sup> increase)	1.10 (0.81–1.47)	0.98 (0.78–1.21)	1.07 (0.73–1.51)	1.01 (0.77–1.30)
Knee OA	2.44 (1.09–5.56)	2.08 (1.18–3.70)	2.07 (0.84–5.21)	1.77 (0.95–3.33)
Knee pain	2.04 (0.72–5.09)	2.05 (0.99–4.00)	1.65 (0.57–4.21)	1.78 (0.85–3.55)
VFx	2.58 (0.82–6.85)	0.40 (0.06–1.36)	2.48 (0.75–7.04)	0.32 (0.05–1.13)
Cognitive impairment	6.19 (1.29–23.1)	4.83 (1.41–15.1)	13.48 (0.98–178.64)	3.17 (0.44–21.99)
<i>Previous falls</i>				
Single fall	–	–	–	3.52 (1.07–9.97)
Multiple falls	1.18 (0.25–4.61)	9.54 (3.15–30.08)	–	12.6 (5.80–27.97)

Multinomial logistic regression analysis was used to calculate the crude odds ratio (OR) and 95% confidence interval (CI) compared with non-fallers.

Adjusted OR was calculated using multinomial logistic regression analysis after adjusting for age and body mass index (BMI).

OA: osteoarthritis, VFx: vertebral fracture, BMD: bone mineral density, LS: lumbar spondylosis.

Radiographic knee OA was defined as Kellgren Lawrence grade 3 or 4.

(Table 4). Previous falls were significantly associated with the incidence of multiple falls. In women, multinomial logistic regression analysis after adjusting for age and BMI showed that a longer 6-m walking time was a risk factor for multiple, but not single falls (Table 5). Chair stand time also tended to be associated with the incidence of single and multiple falls. Regarding bone and joint diseases, knee pain was a risk factor for single and multiple falls. VFx also tended to be associated with multiple falls, but radiographic knee OA was not associated with falls. Cognitive impairment was a risk factor for multiple falls, but not for single falls. A history of previous falls was a risk factor for multiple, but not single falls.

To determine the independent association of each physical performance parameter with the incidence of falls, multinomial logistic regression analysis was performed with age, BMI, 6-m walking time, and chair stand time as explanatory variables. We found that a longer chair stand time was an independent risk factor for multiple falls (OR 1.18, 95% CI 1.06–1.32), but a longer 6-m walking time was not (OR 1.05, 0.93–1.16). In women, a longer 6-m walking time tended to be associated with the incidence of multiple falls (OR 1.09, 95% CI 0.98–1.22), but a longer chair stand time was not (OR 1.01, 95% CI 0.94–1.07). After adjusting for previous falls, the independent association of a longer chair stand time with the incidence of falls remained in men (OR 1.15,

95% CI 1.02–1.30), and the independent association of a longer 6-m walking time with the incidence of falls remained in women (OR 1.12, 95% CI 1.00–1.25). In addition, knee pain and cognitive impairment in women were also significantly associated with falls, and VFx tended to be associated with falls with multinomial logistic regression analysis after adjusting for age and BMI. Thus, to determine the independent association of physical performance, bone and joint diseases, and cognitive impairment, multinomial logistic regression analysis was used with age, BMI, 6-m walking time, knee pain, VFx, and cognitive impairment as explanatory variables. We found that a longer 6-m walking time was an independent risk factor for multiple falls (OR 1.08, 95% CI 1.00–1.18), but the significant association of knee pain, VFx, and cognitive impairment with the incidence of falls disappeared (OR 1.47, 95% CI 0.91–2.35, OR 1.52, 95% CI 0.80–2.81, and OR 1.16, 95% CI 0.35–3.24, respectively).

## Discussion

The present study is the first longitudinal population-based cohort study to examine whether physical performance, bone and joint diseases, and cognitive impairment are risk factors for single and multiple falls in men and women. We found gender differences in risk factors for

**Table 5**  
Risk factors for single and multiple falls in women.

	Crude OR (95% CI)		Adjusted OR (95% CI)	
	Single falls	Multiple falls	Single falls	Multiple falls
Grip strength (5 kg increase)	0.84 (0.70–0.99)	0.81 (0.68–0.95)	0.94 (0.77–1.11)	0.91 (0.75–1.08)
6-m walking time (1 s increase)	1.10 (1.01–1.19)	1.16 (1.08–1.25)	1.04 (0.94–1.14)	1.11 (1.02–1.20)
Chair stand time (1 s increase)	1.07 (1.02–1.12)	1.07 (1.03–1.12)	1.04 (0.99–1.10)	1.04 (0.99–1.09)
LS BMD (0.1 mg/cm <sup>2</sup> increase)	0.88 (0.78–1.00)	0.90 (0.80–1.01)	0.96 (0.83–1.11)	0.92 (0.80–1.06)
Neck BMD (0.1 mg/cm <sup>2</sup> increase)	0.75 (0.63–0.90)	0.85 (0.72–1.01)	0.79 (0.62–1.01)	0.87 (0.69–1.10)
Knee OA	1.79 (1.18–2.78)	1.75 (1.16–2.63)	1.52 (0.94–2.50)	1.12 (0.79–1.82)
Knee pain	1.83 (1.17–2.83)	2.22 (1.44–3.37)	1.62 (1.00–2.60)	1.60 (1.00–2.54)
VFx	1.54 (0.78–2.85)	2.40 (1.35–4.12)	1.15 (0.57–2.20)	1.81 (0.98–3.24)
Cognitive impairment	0.42 (0.02–2.12)	2.12 (0.68–5.60)	0.73 (0.19–2.61)	4.95 (1.50–16.08)
<i>Previous falls</i>				
Single fall	0.55 (0.16–1.74)	1.51 (0.33–5.41)	0.70 (0.30–1.43)	2.48 (1.40–4.28)
Multiple falls	0.86 (0.39–1.81)	8.55 (3.80–19.20)	1.06 (0.35–2.62)	6.93 (3.76–12.72)

Multinomial logistic regression analysis was used to calculate the crude odds ratio (OR) and 95% confidence interval (CI) compared with non-fallers.

Adjusted OR was calculated using multinomial logistic regression analysis after adjusting for age and body mass index (BMI).

OA: osteoarthritis, VFx: vertebral fracture, BMD: bone mineral density, LS: lumbar spondylosis.

Radiographic knee OA was defined as Kellgren Lawrence grade 3 or 4.

falls. Regarding physical performance, a longer chair stand time was an independent risk factor for falls in men, whereas a longer 6-m walking time was an independent risk factor for falls in women. Knee pain, VFX, and cognitive impairment were associated with falls in women, but not in men.

The present study is a population-based longitudinal study to determine whether bone and joint diseases are risk factors for falls in Japanese men and women. After adjusting for age and BMI, knee pain was a risk factor for falls in women, but not in men. The sex differences regarding the association of knee pain with falls may be partly explained by the weaker quadriceps muscles in women, which is known to be an independent risk factor for falls [16]. Muscle strength is higher in men than in women in all decades [39], which may obscure the association of knee pain with falls. In addition, given the insignificant association of radiographic knee OA with falls, falls may occur due to symptoms such as pain rather than radiographic changes in the knee itself. Our study and other previous cross-sectional studies also suggested that knee pain is significantly associated with falls [11]. In other words, falls may be preventable when pain is relieved by medical care, even if patients have radiographic knee OA.

In the present study, LS and lower back pain were not associated with falls, whereas VFX was associated with falls. Lower BMD was not associated with falls in the present study, and thus, radiographic changes but not OP may be associated with falls. Studies of patients with VFX have reported increased kyphosis angles [16,17], which is an independent risk factor for injurious falls [40]. Previous studies [41,42] have demonstrated that people with kyphosis have greater balance abnormalities as assessed by computerized dynamic posturography. Specifically, they reported that women with OP-related kyphosis had greater mediolateral displacement and increased mediolateral velocity compared to controls [42]. In addition, lateral spontaneous sway amplitude has been reported to be the single best predictor of future risk of falls [43]. These observations may partly explain the association between VFX and falls.

In the present study, after adjusting for age and BMI, both a longer 6-m walking time and a longer chair stand time were associated with falls in men and women. A previous study also showed that slower walking speed is a risk factor for falls [44], although men and women were not separately analyzed in the study. To determine the independent association of the 6-m walking time and chair stand time, we further used multinomial logistic regression analysis with age, BMI, 6-m walking time, and chair stand time as explanatory factors, and found that in men, a longer chair stand time was an independent risk factor for multiple falls, but a longer 6-m walking time was not. In women, a longer 6-m walking time was associated with the incidence of multiple falls, whereas a longer chair stand time was not. This indicates that slower walking speed may more strongly affect the risk of falling in women than in men, whereas a longer chair stand time may more strongly affect the risk of falling in men than in women. The walking time and chair stand time can be easily and quickly measured in clinical and research settings without requiring monitoring devices or extensive training. The present study may indicate that walking time is a simple and quick option for measuring the risk of falling, particularly in women, and measuring the chair stand time is a simple and quick option for estimating the risk of falling, particularly in men.

The present study has several limitations. First, our participants lived in the community, and thus, our findings may not apply to elderly persons residing in institutions. Second, we did not include other anatomical locations of weight-bearing OA such as hip OA in the analysis, although this disorder also affects falls [45]. However, the prevalence of KL=3 or 4 hip OA is 1.4% and 3.5% in Japanese men and women [46], respectively, which is lower than that of KL=3 or 4 knee OA (12.2% and 21.0% in men and women, respectively) in the present study. Thus, it is possible that hip OA would not strongly affect the results of the present study. Third, non-responders were older, had

lower physical performance and higher prevalence of knee pain, which were risk factors for falls. This means that the incidence of falls in the present study may have been underestimated. Fourth, the accuracy and reliability of recall of falls over the past 3 years was not assessed in the present study. Previous studies have shown that 13–32% of elderly subjects with confirmed falls did not recall falling over a 12-month period [47], even when excluding subjects with cognitive impairment. Therefore, the incidence of falls may be underestimated, particularly in older subjects and those with cognitive impairment. In addition, individuals are more likely to recall a fall that resulted in injury, which may have influenced the results of this study.

## Conclusion

The present longitudinal analysis using a large-scale population from the ROAD study revealed gender differences in risk factors for falls. A longer walking time was a risk factor for falls in women, whereas a longer chair stand time was a risk factor for falls in men. Knee pain and VFX were risk factors for falls in women, but not in men. Further studies, along with continued longitudinal surveys in the ROAD study, will help elucidate the background of bone and joint diseases and their relationship with falls.

## Acknowledgments

This work was supported by Grants-in-Aid for Scientific Research (S19109007, MB20390182, C20591737, C20591774), Young Scientists (A18689031), and Exploratory Research (19659305) from the Japanese Ministry of Education, Culture, Sports, Science and Technology; H17-Men-eki-009, H18-Choujyu-037, H20-Choujyu-009, H21-Chouju-Wakate-011, and H22-Chouju-Wakate-007 from the Ministry of Health, Labour and Welfare; Research Aid from the Japanese Orthopaedic Association (JOA-Subsidized Science Project Research 2006–1 and 2010–2); and Grant No. 166 from the Japan Orthopaedics and Traumatology Foundation.

The authors thank Tomoko Takijiri and other members of the Public Office in Hidakagawa Town, and Mrs. Tamako Tsutsumi, Mrs. Kanami Maeda, and other members of the Public Office in Taiji Town, for their assistance in the location and scheduling of participants for examinations.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.bone.2012.10.020>.

## References

- [1] Baker S, O'Neill B, Karpf RS. *The Injury Fact Book*. Lexington, Mass: Lexington Books; 1984.
- [2] Fife D, Barancik JI, Chatterjee MS. Northeastern Ohio Trauma Study, II: injury rates by age, sex and cause. *Am J Public Health* 1984;74:473–8.
- [3] Ministry of Health, Labour and Welfare. The outline of the results of National Liveliness Survey 2007. <http://www.mhlw.go.jp/toukei/list/20-19-1.html>.
- [4] Dargent-Molina P, Favier F, Grandjean H, Baudoin C, Schott AM, Hausherr E, et al. Fall-related factors and risk of hip fracture: the EPIDOS prospective study. *Lancet* 1996;348:145–9.
- [5] Tinetti ME, Speechley M, Ginter SF. Risk factors for falls among elderly persons living in the community. *N Engl J Med* 1988;319:1701–7.
- [6] Sharma L, Kapoor D. Epidemiology of osteoarthritis. In: Moskowitz RW, Altman RD, Hochberg MC, Buckwalter JA, Goldberg VM, editors. *Osteoarthritis: Diagnosis and Medical/Surgical Management*. 4th ed. Philadelphia: Lippincott Williams & Wilkins; 2007. p. 3–26.
- [7] Emery SE, Ringus VM. Osteoarthritis of the spine. In: Moskowitz RW, Altman RD, Hochberg MC, Buckwalter JA, Goldberg VM, editors. *Osteoarthritis: Diagnosis and Medical/Surgical Management*. 4th ed. Philadelphia: Lippincott Williams & Wilkins; 2007. p. 427–52.
- [8] Muraki S, Oka H, Akune T, Mabuchi A, En-yo Y, Yoshida M, et al. Prevalence of radiographic knee osteoarthritis and its association with knee pain in the elderly of Japanese population-based cohorts: the ROAD study. *Osteoarthritis Cartilage* 2009;17:1137–43.

- [9] Muraki S, Oka H, Mabuchi A, Akune T, En-yo Y, Yoshida M, et al. Prevalence of radiographic lumbar spondylosis and its association with low back pain in the elderly of population-based cohorts: the ROAD study. *Ann Rheum Dis* 2009;68:1401–6.
- [10] Yoshimura N, Muraki S, Oka H, Mabuchi A, En-yo Y, Yoshida M, et al. Prevalence of knee osteoarthritis, lumbar spondylosis and osteoporosis in Japanese men and women: the Research on Osteoarthritis/Osteoporosis Against Disability (ROAD). *J Bone Miner Metab* 2009;27:620–8.
- [11] Arden NK, Crozier S, Smith H, Anderson F, Edwards C, Raphael H, et al. Knee pain, knee osteoarthritis, and the risk of fracture. *Arthritis Rheum* 2006;55:610–5.
- [12] Nevitt MC, Cummings SR, Kidd S, Black D. Risk factors for recurrent nonsyncopal falls. A prospective study. *JAMA* 1989;261:2663–8.
- [13] Linaker CH, Walker-Bone K, Palmer K, Cooper C. Frequency and impact of regional musculoskeletal disorders. *Baillieres Clin Rheumatol* 1999;13:197–215.
- [14] Muraki S, Akune T, Oka H, En-yo Y, Yoshida M, Nakamura K, et al. Prevalence of falls and its association with knee osteoarthritis and lumbar spondylosis as well as knee and lower back pain in Japanese men and women. *Arthritis Care Res* 2011;63:1425–31.
- [15] Burger H, Van Daele PL, Grashuis K, Hofman A, Grobbee DE, Schutte HE, et al. Vertebral deformities and functional impairment in men and women. *J Bone Miner Res* 1997;12:152–7.
- [16] Ross PD. Clinical consequences of vertebral fractures. *Am J Med* 1997;103:30S–42S [discussion 42S–43S].
- [17] Nevitt MC, Ettinger B, Black DM, Stone K, Jamal SA, Ensrud K, et al. The association of radiographically detected vertebral fractures with back pain and function: a prospective study. *Ann Intern Med* 1998;128:793–800.
- [18] Yamaguchi T, Sugimoto T, Yamada H, Kanzawa M, Yano S, Yamauchi M, et al. The presence and severity of vertebral fractures is associated with the presence of esophageal hiatal hernia in postmenopausal women. *Osteoporos Int* 2002;13:331–6.
- [19] Gold DT, Lyles KW, Shipp KM, Drezner MK. Osteoporosis and its nonskeletal consequences: their impact on treatment decisions. In: Marcus R, Feldman D, Kelsey J, editors. *Osteoporosis*. 2nd ed. San Diego, California, USA: Academic Press; 2001. p. 819–29.
- [20] Leech JA, Dulberg C, Kellie S, Pattee L, Gay J. Relationship of lung function to severity of osteoporosis in women. *Am Rev Respir Dis* 1990;141:68–71.
- [21] Johnell O, Kanis JA, Oden A, Sernbo I, Redlund-Johnell I, Pettersson C, et al. Mortality after osteoporotic fractures. *Osteoporos Int* 2004;15:38–42.
- [22] Studenski S, Perera S, Wallace D, Chandler JM, Duncan PW, Rooney E, et al. Physical performance measures in the clinical setting. *J Am Geriatr Soc* 2003;51:314–22.
- [23] Cesari M, Kritchevsky SB, Penninx BW, Nicklas BJ, Simonsick EM, Newman AB, et al. Prognostic value of usual gait speed in well-functioning older people — results from the Health, Aging and Body Composition Study. *J Am Geriatr Soc* 2005;53:1675–80.
- [24] Lipsitz LA, Jonsson PV, Kelley MM, Koestner JS. Causes and correlates of recurrent falls in ambulatory frail elderly. *J Gerontol* 1991;46:M114–22.
- [25] Wolfson L, Whipple R, Amerman P, Tobin JN. Gait assessment in the elderly: a gait abnormality rating scale and its relation to falls. *J Gerontol* 1990;45:M12–9.
- [26] Chan BK, Marshall LM, Winters KM, Faulkner KA, Schwartz AV, Orwoll ES. Incident fall risk and physical activity and physical performance among older men: the Osteoporotic Fractures in Men Study. *Am J Epidemiol* 2007;165:696–703.
- [27] Guralnik JM, Simonsick EM, Ferrucci L, Glynn RJ, Berkman LF, Blazer DG, et al. A short physical performance battery assessing lower extremity function: association with self-reported disability and prediction of mortality and nursing home admission. *J Gerontol* 1994;49:M85–94.
- [28] Bohannon RW. Sit-to-stand test for measuring performance of lower extremity muscles. *Percept Mot Skills* 1995;80:163–6.
- [29] Yoshimura N, Muraki S, Oka H, Kawaguchi H, Nakamura K, Akune T. Cohort profile: Research on Osteoarthritis/Osteoporosis Against Disability (ROAD) Study. *Int J Epidemiol* 2010;39:988–95.
- [30] Shimada H, Lord SR, Yoshida H, Kim H, Suzuki T. Predictors of cessation of regular leisure-time physical activity in community-dwelling elderly people. *Gerontology* 2007;53:293–7.
- [31] Tinetti M, Baker D, Dutcher J. Reducing the risk of falls among older adults in the community. Berkeley, CA: Peacable Kingdom Press; 1997.
- [32] Inoue T. Clinical features and findings: osteoporosis. *Bone* 1990;4:39–47 [in Japanese].
- [33] Kellgren JH, Lawrence JS, editors. *The Epidemiology of Chronic Rheumatism: Atlas of Standard Radiographs of Arthritis*. Oxford: Blackwell Scientific; 1963.
- [34] Yoshimura N, Kakimoto T, Nishioka M, Kishi T, Iwasaki H, Niwa T, et al. Evaluation of reproducibility of bone mineral density measured by dual energy X-ray absorptiometry (DPX-L). *J Wakayama Med Soc* 1997;48:461–6.
- [35] World Health Organization. Assessment of fracture risk and its application to screening for postmenopausal osteoporosis. WHO technical report series, 843. Geneva: WHO; 1994.
- [36] Orimo H, Hayashi Y, Fukunaga M, Sone T, Fujiwara S, Shiraki M, et al. Osteoporosis Diagnostic Criteria Review Committee: Japanese Society for Bone and Mineral Research. Diagnostic criteria for primary osteoporosis: year 2000 revision. *J Bone Miner Metab* 2001;19:331–7.
- [37] Steffan TM, Hacker TA, Mollinger L. Age- and gender-related test performance in community-dwelling older people: six-minute walk test, Berg balance scale, timed up and go test, and gait speeds. *Phys Ther* 2002;82:128–37.
- [38] Folstein MF, Folstein SE, McHugh PR. "Mini-mental state." A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975;12:129–98.
- [39] Sinaki M, Nwaogwugwu NC, Phillips BE, Mokri MP. Effect of gender, age, and anthropometry on axial and appendicular muscle strength. *Am J Phys Med Rehabil* 2001;80:330–8.
- [40] Kado DM, Huang MH, Nguyen CB, Barrett-Connor E, Greendale GA. Hyperkyphotic posture and risk of injurious falls in older persons: the Rancho Bernardo Study. *J Gerontol A Biol Sci Med Sci* 2007;62:652–7.
- [41] Lynn SG, Sinaki M, Westerlind KC. Balance characteristics of persons with osteoporosis. *Arch Phys Med Rehabil* 1997;78:273–7.
- [42] Sinaki M, Brey RH, Hughes CA, Larson DR, Kaufman KR. Balance disorder and increased risk of falls in osteoporosis and kyphosis: significance of kyphotic posture and muscle strength. *Osteoporos Int* 2005;16:1004–10.
- [43] Maki BE, Holliday PJ, Topper AK. A prospective study of postural balance and risk of falling in an ambulatory and independent elderly population. *J Gerontol Med Sci* 1994;49:M72–84.
- [44] Vergheze J, Holtzer R, Lipton RB, Wang C. Quantitative gait markers and incident fall risk in older adults. *J Gerontol* 2009;64:896–901.
- [45] Arden NK, Nevitt MC, Lane NE, Gore LR, Hochberg MC, Scott JC, et al. Osteoarthritis and risk of falls, rates of bone loss, and osteoporotic fractures. Study of Osteoporotic Fractures Research Group. *Arthritis Rheum* 1999;42:1378–85.
- [46] Inoue K, Wicart P, Kawasaki T, Huang J, Ushiyama T, Hukuda S. Prevalence of hip osteoarthritis and acetabular dysplasia in French and Japanese adults. *Rheumatology (Oxford)* 2000;39:745–8.
- [47] Cummings SR, Nevitt MC, Kidd S. Forgetting falls. The limited accuracy of recall of falls in the elderly. *J Am Geriatr Soc* 1988;36:613–6.

## IV. 資料



## 第2回 お達者整形外来問診票

このたびは、第2回お達者整形外来にご参加いただきありがとうございます。われわれ東京大学22世紀医療センターでは、東京都老人医療センターおよび東京都老人総合研究所と共同で、変形性関節症を中心としたお達者整形外来を東京都老人医療センターにて開催させていただいて参りました。

今回は、3年ぶりの開催になりますが、この3年の間に膝、腰椎、股関節がどのように変化をしたかを知るよい機会になると考えております。つきましては、下記問診票にご協力いただけましたら幸いです。

プライバシーの保持には万全を期しております。あなた様にご迷惑がかかるようなことは絶対にございませんで、どうぞよろしく願いいたします。

生年月日：明治・大正・昭和 年 月 日

性別： 男 ・ 女

お名前：

ご住所：

今日の日付：平成 年 月 日

インタビューサイン



東京大学医学部附属病院22世紀医療センター  
 関節疾患総合研究講座  
 臨床運動器医学講座  
 東京都健康長寿医療センター整形外科  
 東京都健康長寿医療センター研究所

## 1. 今までの骨折や骨の病気のこと。

今までに骨折や骨の病気がなかったかどうか、お教えてください。あてはまるものの番号に○をおつけください。

**1** 今までに骨折したことがありますか？

- 1) はい      2) いいえ

1) はい の場合は以下の質問にお答えください。2) いいえ の場合は **11** へお進みください。

**2** 骨折の場所はどこですか？（もし2ヶ所以上の場合はすべてお答えください。）

- 1) 右肩      (      ) 歳  
2) 左肩      (      ) 歳  
3) 右手首    (      ) 歳  
4) 左手首    (      ) 歳  
5) 右股関節(足の付け根) (      ) 歳

→ 骨折したとき手術をしましたか？

- ① した    ② しない    ③ わからない

- 6) 左股関節(足の付け根) (      ) 歳

→ 骨折したとき手術をしましたか？

- ① した    ② しない    ③ わからない

- 7) 背骨、腰骨 (      ) 歳

- 8) その他 ① (      )

(      ) 歳

- ② (      )

(      ) 歳

**3** 骨折の原因はなんですか？詳しくお教えてください。（もし2回以上の場合はすべてお答えください。）

(      )

(      )

**4** 骨折したときの場所はどこでしたか？

- 1) 屋内  
2) 屋外  
3) その他 (      )

**5** 骨折したときは、どのくらいの明るさでしたか

- 1) 昼間のような明るさで      2) 薄明かりで      3) 暗闇で

- 6** 骨折したときの服装は  
 1) 和服  
 2) 洋服  
 3) その他（具体的に \_\_\_\_\_ )
- 7** そのときの履き物は（下の番号からお選びください。いくつ選んでも結構です。）  
 1) くつ                      2) げた                      3) ぞうり、サンダル                      4) スリッパ  
 5) くつした、たび          6) はだし                      7) その他（具体的に \_\_\_\_\_ )
- 8** そのときになにか別の病気にかかっていましたか？  
 （下の番号からお選びください。いくつ選んでも結構です。）  
 1) 脳卒中                                      2) 心臓の病気                                      3) ふらつき、めまい  
 4) 膝などが固く動きにくい              5) 興奮していた                                      6) ゆううつだった  
 7) 目がかすんでいた                      8) その他（ \_\_\_\_\_ )  
 9) 特になし
- 9** そのとき、次の薬や飲み物を飲んでいましたか？  
 （下の番号からお選びください。いくつ選んでも結構です。）  
 0) 飲まなかった  
 1) 精神安定剤・睡眠薬  
 2) 降圧剤（血圧の薬）  
 3) くしゃみ止めや風邪薬  
 4) アルコール  
 5) その他（具体的に \_\_\_\_\_ )
- 10** 今も骨折部が痛いなど骨折の影響は今もありますか？  
 1) はい →（具体的に \_\_\_\_\_ )  
 2) いいえ
- 11** 今までに、骨粗鬆症（骨が弱い、もろい）といわれましたか。  
 1) はい  
 2) いいえ
- 12** 今までに骨粗鬆症の治療（骨が強くなる）をしたことがありますか。  
 1) はい（1. 筋注      2. 静注      3. 服薬 → 薬の名前 \_\_\_\_\_ )  
 2) いいえ
- 13** （女性のみ）閉経されましたか。また、された方は何歳でしたか？  
 1) はい（ \_\_\_\_\_ ）歳  
 2) いいえ

