

- 330-338
- 5) Yang JG, Li J, Lu C, Hasimu B, Yang Y, Hu D: Chronic kidney disease, all-cause mortality and cardiovascular mortality among Chinese patients with established cardiovascular disease. *J Atheroscler Thromb*, 2010; 17: 395-401
 - 6) Kubozono T, Miyata M, Ueyama K, Nagaki A, Hamasaki S, Kusano K, Kubozono O, Tei C: Association between arterial stiffness and estimated glomerular filtration rate in the Japanese general population. *J Atheroscler Thromb*, 2009; 16: 840-845
 - 7) Damsgaard EM, Froland A, Jorgensen OD, Mogensen CE: Microalbuminuria as predictor of increased mortality in elderly people. *BMJ*, 1990; 300: 297-300
 - 8) Wattanakit K, Folsom AR, Criqui MH, Kramer HJ, Cushman M, Shea S, Hirsch AT: Albuminuria and peripheral arterial disease: results from the multi-ethnic study of atherosclerosis (MESA). *Atherosclerosis*, 2008; 201: 212-216
 - 9) Wu CK, Yang CY, Tsai CT, Chiu FC, Huang YT, Lee JK, Cheng CL, Lin LY, Lin JW, Hwang JJ, Chiang FT: Association of low glomerular filtration rate and albuminuria with peripheral arterial disease: the National Health and Nutrition Examination Survey, 1999-2004. *Atherosclerosis*, 2010; 209: 230-234
 - 10) Arima H, Tanizaki Y, Kiyohara Y, Tsuchihashi T, Kato I, Kubo M, Tanaka K, Ohkubo K, Nakamura H, Abe I, Fujishima M, Iida M: Validity of the JNC VI recommendations for the management of hypertension in a general population of Japanese elderly: the Hisayama study. *Arch Intern Med*, 2003; 163: 361-366
 - 11) Tsuchikura S, Shoji T, Kimoto E, Shinohara K, Hatsuda S, Koyama H, Emoto M, Nishizawa Y: Brachial-ankle pulse wave velocity as an index of central arterial stiffness. *J Atheroscler Thromb*, 2010; 17: 658-665
 - 12) National Kidney Foundation: KDOQI Clinical Practice Guidelines and Clinical Practice Recommendations for Diabetes and Chronic Kidney Disease. *Am J Kidney Dis*, 2007; 49: S1-S180
 - 13) Arima H, Kubo M, Yonemoto K, Doi Y, Ninomiya T, Tanizaki Y, Hata J, Matsumura K, Iida M, Kiyohara Y: High-sensitivity C-reactive protein and coronary heart disease in a general population of Japanese: the Hisayama study. *Arterioscler Thromb Vasc Biol*, 2008; 28: 1385-1391
 - 14) DeLong ER, DeLong DM, Clarke-Pearson DL: Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. *Biometrics*, 1988; 44: 837-845
 - 15) Cao JJ, Barzilay JI, Peterson D, Manolio TA, Psaty BM, Kuller L, Wexler J, Bleyer AJ, Cushman M: The association of microalbuminuria with clinical cardiovascular disease and subclinical atherosclerosis in the elderly: the Cardiovascular Health Study. *Atherosclerosis*, 2006; 187: 372-377
 - 16) Yudkin JS, Forrest RD, Jackson CA: Microalbuminuria as predictor of vascular disease in non-diabetic subjects. *Islington Diabetes Survey*. *Lancet*, 1988; 2: 530-533
 - 17) Hlatky MA, Greenland P, Arnett DK, Ballantyne CM, Criqui MH, Elkind MS, Go AS, Harrell FE Jr, Hong Y, Howard BV, Howard VJ, Hsue PY, Kramer CM, McConnell JP, Normand SL, O'Donnell CJ, Smith SC Jr, Wilson PW: Criteria for evaluation of novel markers of cardiovascular risk: a scientific statement from the American Heart Association. *Circulation*, 2009; 119: 2408-2416
 - 18) Deckert T, Feldt-Rasmussen B, Borch-Johnsen K, Jensen T, Kofoed-Enevoldsen A: Albuminuria reflects widespread vascular damage. The Steno hypothesis. *Diabetologia*, 1989; 32: 219-226
 - 19) Weir MR: Microalbuminuria and cardiovascular disease. *Clin J Am Soc Nephrol*, 2007; 2: 581-590
 - 20) Festa A, D'Agostino R, Howard G, Mykkanen L, Tracy RP, Haffner SM: Inflammation and microalbuminuria in nondiabetic and type 2 diabetic subjects: The Insulin Resistance Atherosclerosis Study. *Kidney Int*, 2000; 58: 1703-1710
 - 21) Boger RH, Bode-Boger SM, Thiele W, Junker W, Alexander K, Frolich JC: Biochemical evidence for impaired nitric oxide synthesis in patients with peripheral arterial occlusive disease. *Circulation*, 1997; 95: 2068-2074
 - 22) Hogan SL, Vupputuri S, Guo X, Cai J, Colindres RE, Heiss G, Coresh J: Association of cigarette smoking with albuminuria in the United States: the third National Health and Nutrition Examination Survey. *Ren Fail*, 2007; 29: 133-142
 - 23) Tylicki L, Puttinger H, Rutkowski P, Rutkowski B, Horl WH: Smoking as a risk factor for renal injury in essential hypertension. *Nephron Clin Pract*, 2006; 103: c121-c128
 - 24) Yanbaeva DG, Dentener MA, Creutzberg EC, Wesseling G, Wouters EF: Systemic effects of smoking. *Chest*, 2007; 131: 1557-1566
 - 25) Argacha JF, Adamopoulos D, Gujic M, Fontaine D, Amyai N, Berkenboom G, van de Borne P: Acute effects of passive smoking on peripheral vascular function. *Hypertension*, 2008; 51: 1506-1511

The first part of the document discusses the importance of maintaining accurate records of all transactions. It emphasizes that every entry should be supported by a valid receipt or invoice. This not only helps in tracking expenses but also ensures compliance with tax regulations.

In the second section, the author provides a detailed breakdown of the company's revenue streams. This includes sales from various product lines and services. The analysis shows that while one product line is currently the primary source of income, diversification into new markets is necessary for long-term growth.

The third section addresses the company's financial health and liquidity. It highlights the need for a robust cash flow management strategy to ensure that all operational needs are met. The author suggests implementing regular financial reviews to identify potential risks and opportunities early on.

Finally, the document concludes with recommendations for future strategic planning. It suggests that the company should focus on innovation and customer engagement to stay competitive in a rapidly changing market. The author also notes the importance of maintaining strong relationships with suppliers and partners.