

Table 1 JBCRG trials closed/in follow-up

Trial	Design	No. of pts	Primary endpoints	Regimen	Enrollment start date
Neoadjuvant setting					
1	Phase II	202	Clinical response, safety	FEC100 q3w×4 → Doc75 q3w×4	Jun 02
2	Phase II	31	Clinical response, safety	FEC100 q3w×4 → Doc100 q3w×4	Aug 04
2'	Validation	19	Clinical response, histological effects, safety	FEC100 q3w×4 → Doc100 q3w×4	Dec 05
3	Phase II	130	Histological effects, safety	Doc75 q3w×4 → FEC100 q3w×4	Oct 05
5	Phase II	33	Response rates	Doc75 q3w×4 → letrozole 12 (-18) w	Sep 07
6	Phase II	40	Response rates	Letrozole 12 (-18) w	Sep 07
7	Phase II	40	Response rates	Letrozole + cyclophosphamide 24 w	Oct 07
10	Randomized phase II	180	Pathological CR rate	(1) FEC×4 → TCH×4, (2) TCH×4 → FEC×4, (3) TCH×6	Jun 09
13	Phase II	40	Pathological CR rate	Metronomic PCX 4 → FEC×4	Jan 10
Postoperative setting					
4 (CREATE-X)	Phase III	900	Disease-free survival	Any preoperative systemic therapy ± capecitabine	Feb 07
SOLE with BIG	Phase III	4,800	Disease-free survival	Intermittent or continuous letrozole	Apr 10
8 ALTTO	Phase III	140	Disease-free survival	Lapatinib and/or trastuzumab	Jul 07
Metastatic setting					
M01	Phase I	6	MTD, DLT, RD	CPT11 + S1	Jul 06
M01	Phase II	37	Response rates, clinical efficacy	CPT11 + S1	Jul 06
M02	Phase II	50	Response rates	Letrozole	Nov 06
Cohort study					
C01	Cohort	1,500	Disease-free survival	Trastuzumab	Sep 07

Data correct as of 31 March 2012

CR complete response, MTD maximum tolerated dose, DLT dose limiting toxicity, RD recommended dose, FEC 5-fluorouracil + epirubicin + cyclophosphamide, Doc docetaxel, TCH docetaxel + cyclophosphamide + trastuzumab, PCX paclitaxel + cyclophosphamide + capecitabine

Neoadjuvant pharmacotherapy

The first clinical trial conducted by the JBCRG was JBCRG-01, a phase II trial of preoperative systemic therapy (PST) using fluorouracil, epirubicin, and cyclophosphamide (FEC) followed by docetaxel (Doc) in patients with primary operable breast cancer [1–3]. Subsequently, JBCRG-02 study was conducted using FEC followed by Doc100 to investigate the safety and feasibility of 100 mg/m² Doc as PST. JBCRG-03 was a study to clarify the most effective sequence of FEC and Doc75 [4]. From the results of these studies, we defined new criteria of pathological response to PST, quasi pathological complete response (QpCR), total or near total disappearance of the invasive tumor in the removed breast. QpCR following preoperative chemotherapy predicts favorable disease-free survival (DFS). HER2 overexpression and clinical response to FEC predict QpCR [5, 6].

JBCRG-01

JBCRG-01 was started in 2002 [1–3]. This multicenter phase II study examined the impact of pathological effect on survival after preoperative chemotherapy in Japanese women with early-stage breast cancer (ESBC). Prior to surgery, patients received four cycles of FEC (fluorouracil 500 mg/m², epirubicin 100 mg/m², cyclophosphamide 500 mg/m² q3w) followed by four cycles of docetaxel (75 mg/m² q3w). The primary endpoint was 3-year DFS stratified by the absence or presence of QpCR (absence of invasive tumor or only focal residual tumor cells). Secondary endpoints were predictors for QpCR, clinical response, breast conservation rate, and safety. Between June 2002 and June 2004, 202 women were enrolled. Among 191 assessable patients, 25 % achieved QpCR. With 40 months median follow-up, 3-year DFS was estimated at 91 % for all patients. The 3-year DFS for patients

Table 2 JBCRG trials open to recruitment

Trial	Design	No. of pts	Primary endpoints	Regimen	Enrollment start date
Neoadjuvant setting					
9	Randomized phase II	195	Histological response	TC×6, FEC×3 → TC×3,TC×3 → FEC×3	Sep 09
11CPA	Phase II	55	Response rates	Letrozole ± low dose cyclophosphamide	Oct 10
11TC	Phase II	60	Clinical response	Exemestane 12w or exemestane 12w+TC×4	Oct 10
Postoperative setting					
15	Phase II	30	Pharmacokinetics	Toremifene	Mar 09
SUPREMO with IBCSG	Phase III	3,700	Overall survival	Chest wall radiation	Jul 09
Metastatic setting					
12	Phase II	200	CYP2D6 and pharmacokinetics	Tamoxifen and toremifene	Jan 10
Cohort study					
C02	Cohort	100	Progression-free survival	Trastuzumab	Jul 09

Data correct as of 31 March 2012

TC docetaxel + cyclophosphamide, FEC 5-fluorouracil + epirubicin + cyclophosphamide

with QpCR was 98 vs. 89 % for those without QpCR (hazard ratio 0.38 [95 % confidence interval 0.09–0.84], $P = 0.0134$). HER2 status and response to FEC were independent predictors of QpCR. The overall clinical response rate was 75 %; 85 % of patients achieved breast conservation. Grade 3/4 neutropenia was the most common adverse event, observed in 44 and 35 % of patients during FEC and docetaxel treatment, respectively. Treatment-related side effects were manageable; there were no treatment-related fatalities.

JBCRG-02

The JBCRG-02 study was conducted to evaluate the safety and clinical and histologic effects of primary systemic chemotherapy using FEC followed by docetaxel in primary breast cancer. The primary endpoints were safety and clinical and histologic effects. Secondary endpoints were breast-conserving rate and DFS. Fluorouracil 500 mg/m², epirubicin 100 mg/m², and cyclophosphamide 500 mg/m², q3w ×4 cycles, were followed by docetaxel 100 mg/m², q3w ×4 cycles, as primary systemic chemotherapy. Among patients receiving this regimen, 19.5 % experienced a pathological complete response and 9.7 % had a near pathological complete response, resulting in a QpCR of 29.2 %.

JBCRG-03

JBCRG-03 was a multicenter, open-label, single-arm, phase II study assessing the efficacy of a neoadjuvant chemotherapy with docetaxel (75 mg/m² q3w) followed by

5-fluorouracil 500 mg/m², epirubicin 100 mg/m², and cyclophosphamide 500 mg/m² q3w in patients with ESBC [4]. The primary endpoint was the pathological complete response (pCR) rate defined for the breast alone, assessed by a central review committee. Secondary endpoints included clinical response and safety. Of the 132 patients assessable for pathologic response, 23 % experienced a pCR and 6 % had a near pathological complete response (few remaining cancer cells), resulting in a QpCR of 29 %. Clinical response rate following the initial docetaxel regimen was 64 %. The overall clinical response rate was 79 %. Breast-conserving surgery was performed in 79 % of patients. More patients with triple-negative disease experienced a pCR (14/29, 48 %) versus those with other molecular subtypes. The safety profile was acceptable.

Oncotype DX

The 21-gene signature has been intensively studied and incorporated into major guidelines for treatment decision in early breast cancer. However, it remains to be examined whether this system is applicable to Asian populations.

Retrospective analysis

Toi et al. [7] were the first report to show that the 21-gene signature has value in providing prognostic information in Asian populations with estrogen receptor (ER)-positive, lymph node (LN)-negative breast cancer. A total of 325 tumor tissues were collected from ER-positive primary breast cancer patients who had undergone surgery and were

treated with tamoxifen between 1992 and 1998. The tissues were analyzed for the 21-gene signature, and the patients were classified into groups of low, intermediate, or high risk on the basis of the recurrence score. A total of 280 patients were eligible, with adequate reverse transcription polymerase chain reaction profiles for the recurrence score. Of those, 200 and 80 patients had LN-negative and LN-positive disease, respectively. The proportions of LN-negative patients categorized as being at low, intermediate, or high risk were 48, 20, and 33 %, respectively. In LN-negative patients, the Kaplan–Meier estimates of the distant recurrence rate at 10 years were 3.3 % (95 % CI 1.1–10.0 %), 0 %, and 24.8 % (95 % CI 15.7–37.8 %) for those in the low-risk, intermediate-risk, and high-risk groups, respectively. The risk of distant recurrence in the low-risk group was significantly lower than that in the high-risk group when the entire Kaplan–Meier plots were compared ($P < 0.001$, log-rank test). There was a significant difference for overall survival between the low-risk and the high-risk groups ($P = 0.008$, log-rank test).

Economic evaluation

JBCRG-TR03

This study evaluates the cost-effectiveness of two scenarios designed to include the assay into Japan's social health insurance benefit package: one for LN–, ER+, ESBC and another for LN±, ER+, ESBC [8]. An economic decision tree and Markov model under Japan's health system from the societal perspective is constructed with new evidence from the Japanese validation study. Incremental cost-effectiveness ratios are estimated as ¥384,828 (US\$3,848) per quality-adjusted life year (QALY) for the LN– scenario and ¥568,533 (US\$5,685) per QALY for the LN± scenario. Both estimates are not more than the suggested social willingness-to-pay for one QALY gain from an innovative medical intervention in Japan, ¥5,000,000/QALY (US\$50,000/QALY). Sensitivity analyses show that this result is plausibly robust, because the incremental cost effectiveness ratios (ICERs) do not exceed the threshold despite various changes of assumptions made and values employed. Therefore, the inclusion of the assay in Japan's social health insurance benefit package for not only LN– diseases but also LN+ diseases is cost-effective. Such a decision can be justifiable as an efficient use of finite resources for health care.

Toxicity

Steroids and H(2) blockers are commonly used as supportive care for taxane-containing chemotherapy, but they also affect docetaxel's primary metabolizer, cytochrome

P(450) 3A4. Kawaguchi et al. [9] performed a retrospective observational study to better understand the effects of these compounds on docetaxel-induced skin toxicities, specifically hand-foot syndrome (HFS) and facial erythema (FE), a relationship that is currently poorly understood. Member institutions of the JBCRG were invited to complete a questionnaire on the occurrence of grade 2 or higher HFS and FE among patients treated between April 2007 and March 2008 with docetaxel as an adjuvant or neoadjuvant chemotherapeutic treatment for breast cancer. We obtained data for 993 patients from 20 institutions. Twenty percent received H(2) blockers, and all patients received dexamethasone. Univariate and multivariate analyses revealed that H(2) blockers are associated with a significantly higher incidence of both HFS and FE. The incidence of FE was significantly higher for the docetaxel + cyclophosphamide (TC) regimen than for non-TC regimens combined. Dexamethasone usage did not affect the incidence of either HFS or FE. In conclusion, the use of H(2) blockers as premedication in breast cancer patients receiving docetaxel significantly increases the risk of both HFS and FE.

International study

The JBCRG is a member of the international breast cancer research group BIG. The JBCRG has joined in with several international clinical studies.

JBCRG-04 (CREATE-X)

This study aims to investigate the efficacy and safety of capecitabine, as a postoperative adjuvant chemotherapy, for breast cancer patients who were pathologically demonstrated to have residual cancer cells after the preoperative chemotherapy. In addition, the cost-effectiveness of capecitabine is to be investigated. The primary objective is DFS and secondary ones are overall survival, safety, and cost-effectiveness. Eligible patients had stage I–IIIB at the first diagnosis (curable breast cancer) and were non-pCR after preoperative chemotherapy including at least two cycles anthracycline agents; that is, they were confirmed pathologically by surgical and/or histological tests to have residual cancer cells. The patients had also been confirmed to be HER2 negative.

JBCRG-08 (ALTO)

JBCRG-08 was a randomized, multicenter, open-label, phase III study of adjuvant lapatinib, trastuzumab, their sequence, and their combination in patients with HER2/ErBB2-positive primary breast cancer (BIG 2-06/N063D/EGF 106708.). The objective of this study was to compare DFS in patients with HER2 overexpressing and/or amplified

breast cancer randomized to trastuzumab for 1 year versus lapatinib for 1 year versus trastuzumab (12 weeks) followed by a 6-week treatment-free interval followed by lapatinib (34 weeks) versus trastuzumab in combination with lapatinib for 1 year. Endpoints were DFS, overall survival (OS), time to recurrence (TTR), time to distant recurrence (TTDR), safety and tolerability, cumulative incidence of brain metastases as the first site of breast cancer recurrence, presence or absence of cMyc gene amplification, expression levels of PTEN, and presence or absence of p95 HER2 domain. Trial periods were between July 2007 and February 2011 (registration, 2 years; follow-up study, 5 years). Target sample size was 140 from 15 institutions.

SOLE trial

SOLE trial is a phase III trial evaluating the role of continuous letrozole versus intermittent letrozole following 4–6 years of prior adjuvant endocrine therapy for postmenopausal women with hormone receptor-positive, node-positive ESBC. The JBCRG is collaborating with the International Breast Cancer Study Group (IBCSG) on this trial. A total of 4,800 patients are expected to be enrolled in this study. The primary endpoint is DFS, and secondary ones are OS, distant DFS, breast cancer-free interval, sites of first failure, second (non-breast) malignancies, deaths without prior cancer events, and adverse events.

SUPREMO trial

The SUPREMO trial is a randomized phase III trial assessing the role of chest wall irradiation in women with intermediate-risk breast cancer following mastectomy conducted by BIG. Postoperative radiotherapy is routinely given to patients at higher risk of recurrence with 4 or more LNs or large tumor(s). In patients with less than 4 LNs under the armpit involved by cancer or with no LNs involved but other features of the cancer that increase the risk of recurrence, it is not clear whether postoperative radiotherapy is needed. Eligibility criteria are a postoperative breast cancer patient who has had a mastectomy, and who has an intermediate risk of the cancer returning. An intermediate risk is diagnosed when there are less than 4 LNs under the armpit involved by cancer or there are no LNs involved, but there are other features of the cancer that mean it is more likely to come back. The trial will involve 1,600 women.

Conclusion

The JBCRG was founded in order to perform good-quality multicenter studies, and related clinical trials in close liaison with research institutions in other countries and

regions, as well as in Japan. The JBCRG has performed a variety of studies, including primary pharmacotherapy, pharmacotherapy for recurrent breast cancer, clinical trials on postoperative pharmacotherapy, prediction of prognosis in hormone receptor-positive breast cancer, and prediction of the effect of chemotherapeutic drugs. The JBCRG has reported a number of outcomes to academic societies and in journals, and has obtained a good reputation. The incidence of breast cancer in Japan has increased yearly; thus, more attention has been given to breast cancer treatment, among all cancers. In order to save as many breast cancer patients as possible, and to improve their QOL, we will develop new diagnostic methods, treatments, and prophylaxes for breast cancer.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- Ohno S, Toi M, Kuroi K, Nakamura S, Iwata H, Kusama M, et al. Update results of FEC followed by docetaxel neoadjuvant trials for primary breast cancer. *Biomed Pharmacother*. 2005;59:S323–3234.
- Iwata H, Nakamura S, Toi M, Shin E, Masuda N, Ohno S, et al. Interim analysis of a phase II trial of cyclophosphamide, epirubicin and 5-fluorouracil (CEF) followed by docetaxel as preoperative chemotherapy for early stage breast carcinoma. *Breast Cancer*. 2005;12:99–103.
- Toi M, Nakamura S, Kuroi K, Iwata H, Ohno S, Masuda N, et al. Phase II study of preoperative sequential FEC and docetaxel predicts of pathological response and disease free survival. *Breast Cancer Res Treat*. 2008;110:531–9.
- Iwata H, Sato N, Masuda N, Nakamura S, Yamamoto N, Kuroi K, et al. Docetaxel followed by fluorouracil/epirubicin/cyclophosphamide as neoadjuvant chemotherapy for patients with primary breast cancer. *Jpn J Clin Oncol*. 2011;41:867–75.
- Kuroi K, Toi M, Tsuda H, Kurosumi M, Akiyama F, Japan Breast Cancer Research Group (JBCRG). Issues in the assessment of the pathologic effect of primary systemic therapy for breast cancer. *Breast Cancer*. 2005;13:38–48.
- Kuroi K, Toi M, Tsuda H, Kurosumi M, Akiyama F. Un argued issues on the pathological assessment of response in primary systemic therapy for breast cancer. *Biomed Pharmacother*. 2005;59:S387–92.
- Toi M, Iwata H, Yamanaka T, Masuda N, Ohno S, Nakamura S, et al. Clinical significance of the 21-gene signature (Oncotype DX) in hormone receptor-positive early stage primary breast cancer in the Japanese population. *Cancer*. 2010;1:3113–8.
- Kondo M, Hoshi S, Yamanaka T, Ishiguro H, Toi M. Economic evaluation of the 21-gene signature (Oncotype DX®) in lymph node-negative/positive, hormone receptor-positive early-stage breast cancer based on Japanese validation study (JBCRG-TR03). *Breast Cancer Res Treat*. 2011;127:739–49.
- Kawaguchi K, Ishiguro H, Morita T, Nakamura S, Ohno S, Masuda N, et al. Correlation between docetaxel-induced skin toxicity and the use of steroids and H2 blockers: a multi-institution survey. *Breast Cancer Res Treat*. 2011;130:627–34.

