

Table 2. Results of the JCOG trials for adult T-cell leukemia–lymphoma

Protocol	Regimen	Phase	No. of patients	%CR	MST (months)	Survival (%)	Reference
JCOG7801	VEPA	II	18	17	5	0 (3 years)	11
JCOG8101		III	54	28	8	8.3 (4 years)	12
	VEPA		24	17	NA	NA	
	VEPA-M		30	37	NA	NA	
JCOG8701	LSG4	II	42	43	8	12 (5 years)	13
JCOG9109	LSG11	II	60	28	7	16 (2 years)	31
JCOG9303	LSG15	II	93	36	13	31 (2 years)	32
JCOG9801		III	118				33
	mLSG15		57	40	13	24 (3 years)	
	CHOP-14		61	25	11	13 (3 years)	

For abbreviations, see Table 1. LSG 11 consists of 2'-deoxycoformycin, VCR, ETP, PSL and DOX; LSG15 consists of VCAP (VCR, CPA, PSL and DOX), AMP [DOX, raimustine (MCNU), VECP [VDS, ETP, carboplatin (CBDCA) and PSL], intrathecal MTX + PSL, with each intensified by the prophylactic use of G-CSF (granulocyte colony-stimulating factor); mLSG15 is a modified LSG15.

Table 3. Results of the JCOG trials for lymphoblastic lymphoma/acute lymphoblastic leukemia

Protocol	Regimen	Phase	No. of patients	%CR	PFS (%)	MST (months)	Survival (%)	Reference
JCOG8702	LSG 5	II	46	78	NA	14	15 (7 years)	38
JCOG9004	LSG10	II	143	83	26 (5 years)	26	32 (7 years)	39
JCOG9402	LSG16	II	108	81	28 (5 years)	21	28 (7 years)	40

For abbreviations, see Tables 1 and 2. PFS, progression-free survival; LSG5 consists of VEPA-L [VEPA with L-asparaginase (L-ASP) and intrathecal (IT) MTX/PSL] and M-VEPA (moderate-dose methotrexate plus VEPA); LSG10 consists of induction by LSG5/consolidation by DCMP (DOX, AraC, VDS, PSL, IT-MTX/PSL)/MEVP (mitoxantron, ETP, VCR, PSL, IT-MTX/PSL)/maintenance by 6-mercaptopurine (6-MP)/MTX, with allowing HSCT; LSG16 consists of induction by VEPA-L/consolidation by DCMP and CCMOL (CPA, AraC, 6-MP, VCR, L-ASP with IT-MTX/PSL)/intensified maintenance with allowing HSCT.

Table 4. Results of the JCOG trials for advanced Hodgkin's lymphoma

Protocol	Regimen	Phase	No. of patients	%CR	PFS (%)	Survival (%)	Reference
JCOG8905	C-MOPP/ABVd	II	79	84	73 (4 years)	85 (5 years)	41
JCOG9305	ABVd	II	128	81	78 (5 years)	91 (5 years)	42
JCOG9705	ABV + R	II	72 ^a	72	49 (2 years)	92 (2 years)	44

For abbreviations, see Tables 1–3. C-MOPP consists of CPA, VLB, PCZ and PDN; ABVd consists of DOX, BLM, VLB and dacarbazine (DTIC); ABV + R consists of DOX, BLM, VLB with radiation.

^aNo. of enrolled patients with eligibility until the early termination.

lymphoid malignancies since 1978 with the help of the Central Pathology Review, the Radiation Therapy Quality Assurance and the Central CT Review Committees.

The research on treatments for lymphoid malignancies by JCOG-LSG is now supported by four grants for the principal investigators of the LSG studies by MHLW and Grants-in-Aid for Cancer Research (23A-17). JCOG-LSG

has conducted more than 30 clinical trials including 10 randomized trials for several entities of lymphoid malignancies, meta-analyses of them, and correlative epidemiological and pathological studies on HTLV-1 and T/B-cell phenotype, respectively, providing several standard treatments, classifications and prognostic indexes for lymphoid malignancies as shown in the following sections.

Table 5. Results of the JCOG trials for advanced multiple myeloma

Protocol	Regimen	Phase	No. of patients	%RR (no)	Median PFS	MST (months)	Survival (%)	Reference
JCOG8906	COP/MP	II	69	51	13	39	51/27 (3/5 years)	48
JCOG9301		III	210					49
	MCNU-COP/MP		107	56	23	50	38 (5 years)	
	mCOP/MP		103	44	16	44	40 (5 years)	
JCOG0005-DI	VAD and up-front auto-HSCT	II	16 ^a	NA	NA	NA	NA	NA
JCOG0112	MP/VAD with IFN + PSL versus PSL	III	34 ^a					50
	VAD		16	44	NA	NA	NA	
	MP		17	47	NA	NA	NA	

For abbreviations, see Tables 1–4. DI, Data Center independent; IFN, interferon- α ; COP consists of CPA, VCR and PSL; MP consists of melphalan and PSL; mCOP/MP is a modified COP/MP; VAD consists of VCR, DOX and dexamethasone.

^aNo. of enrolled patients until the early termination.

Table 6. Results of the JCOG trials for indolent B-cell lymphomas and localized nasal natural killer/T-cell lymphoma

Protocol	Regimen	Diseases	Phase	No. of patients	%CR/CRu (no)	PFS (%)	Survival (%)	Reference
JCOG0203		Indolent B	III	300				52
	CHOP-14			151	76	43 (6 years)	88 (6 years)	
	CHOP-21			149	78	41 (6 years)	87 (6 years)	
JCOG0211-DI	DEVIC/50 Gy	Nasal NK/T	I/II	33	77 (20/26)	67 (2 years)	78 (2 years)	56

For abbreviations, see Tables 1–5. DEVIC consists of DEX, ETP, ifosfamide (IFM) and CBDCA.

Table 7. Summaries of the JCOG-LSG correlative studies on trials for malignant lymphomas

Protocol	Trials	Disease	No. of patients	Reference
JCOG0108-A	9305, 0705	Hodgkin	167	45
JCOG0108-A	9002, 9203, 9505, 9506, 9508, 9809	NHL	1141	55
		DLBCL		NA
		T/NK	136	55
JCOG0103-A		NHL	499	59

For abbreviations, see Tables 1–6. NHL, non-Hodgkin’s lymphoma; DLBCL, diffuse large B-cell lymphoma; T/NK, peripheral T and NK-cell lymphomas.

CONSECUTIVE AND ONGOING TRIALS FOR MAJOR LYMPHOID MALIGNANCIES BY JCOG-LSG

ADVANCED-STAGE, AGGRESSIVE NHL

Since 1978, chemotherapy trials have been consecutively conducted for patients with advanced-stage, aggressive NHL

in JCOG-LSG (Table 1 and Fig. 1). After the completion of JCOG7801, a Phase II study of VEPA therapy (vincristine, cyclophosphamide, prednisone and doxorubicin), with promising results, JCOG-LSG started in 1981 a randomized Phase III trial (JCOG8101) to evaluate VEPA versus VEPA-M [VEPA plus methotrexate (MTX)] for advanced-stage NHL (11,12). The difference in survival between the two arms was not significant; however, unique pretreatment variables predictive for efficacy were found. Three factors, leukemic change, poor performance status (PS) and T-cell phenotype, were negatively associated with both the complete remission rate (%CR) and overall survival (OS). In addition, ATL was found to have a much poorer in prognosis than non-ATL peripheral T-cell lymphoma (26).

In 1987, JCOG-LSG initiated a Phase II study (JCOG8701) of a multiagent combination chemotherapy (LSG4 protocol) for advanced aggressive NHL (10). The LSG4 protocol consisted of three regimens: (i) VEPA-B (VEPA plus bleomycin), (ii) M-FEPA (MTX, vindesine, cyclophosphamide, prednisone and doxorubicin) and (iii) VEPP-B (vincristine, etoposide, procarbazine, prednisone and bleomycin). A central pathology review revealed 84 patients with T-NHL, including 42 with ATL, 151 with B-NHL and 33 with NHL of undetermined lineage




Figure 1. Consecutive studies by JCOG-LSG. HL, Hodgkin's lymphoma; LBL/ALL, lymphoblastic lymphoma/acute lymphoblastic leukemia; Agg-NHL, aggressive non-Hodgkin's lymphoma; Ind-B-NHL, indolent B-NHL; ATL, adult T-cell leukemia-lymphoma; NK/T ML, localized nasal natural killer/T-cell lymphoma; MM, multiple myeloma; DLBCL, diffuse large B-cell lymphoma; MCL, mantle cell lymphoma. LSG1, VEPA, consists of vincristine (VCR), cyclophosphamide (CPA), prednisone (PSL) and doxorubicin (DOX); LSG2, VEPA-M, consists of VEPA plus methotrexate (MTX); LSG4 consists of VEPA-B, M-FEPA and VEPP-B, where VEPA-B consists of VEPA plus Bleomycin (BLM), M-FEPA consists of a moderate dose of MTX, vindesine (VDS), CPA, PSL and DOX, and VEPP-B consists of VCR, CPA, PSL and procarbazine (PCZ); LSG5 consists of VEPA-L [VEPA with L-asparaginase (L-ASP) and intrathecal (IT) MTX/PSL] and M-VEPA (moderate-dose methotrexate plus VEPA); LSG6 consists of C-MOPP/ABVd; C-MOPP consists of CPA, VLB, PCZ and PDN; ABVd consists of DOX, BLM, VLB and dacarbazine (DTIC); LSG8 consists of COP/MP; COP consists of CPA, VCR and PSL; MP consists of melphalan and PSL; mLSG8 is a modified LSG8; LSG9 consists of dose-intensified mLSG4; LSG10 consists of induction by LSG5/consolidation by DCMP (DOX, AraC, VDS, PSL, IT-MTX/PSL)/MEVP (mitoxantron, ETP, VCR, PSL, IT-MTX/PSL)/maintenance by 6-mercaptopurine (6-MP)/MTX, with HSCT; LSG11 consists of DCF, VCR, ETP, PSL and DOX; LSG12 consists of VEPA/FEPP, where FEPP consists of vindesine, etoposide, procarbazine and prednisolone; LSG13 consists of raimustine (MCNU)-COP/MP; LSG14 consists of ABVd; LSG15 consists of VCAP (VCR, CPA, PSL, DOX), AMP (DOX, MCNU, PSL), VECF [VDS, ETP, carboplatin (CBDCA), PSL], intrathecal MTX + PLS, with each intensified by the prophylactic use of G-CSF (granulocyte colony-stimulating factor); mLSG15 is a modified LSG15; LSG16 consists of induction by VEPA-L/consolidation by DCMP and CCMOL (CPA, cytarabine, 6-MP, VCR, L-ASP with IT-MTX/PSL)/intensified maintenance with allowing HSCT; LSG17, CHOP, consists of CPA, DOX, VCR and PSL; LSG18 consists of CHOP-14 followed by up-front autologous hematopoietic stem cell transplantation (auto-HSCT); LSG19 consists of CHOP-14; LSG22, ABV + R, consists of DOX, BLM, VLB with radiation; RT/DeVIC in JCOG0005DI consisting of VAD (VCR, DOX and DEX) followed by up-front auto-HSCT; Maint-IFN/PSL in JCOG0112 consisting of MP/VAD induction therapy followed by maintenance therapy with interferon plus PSL versus PSL; R-CHOP and R-CHOP-14 in JCOG0203 consisting of rituximab plus CHOP and rituximab plus CHOP-14, respectively; MCL-ASCT in JCOG0406 consisting of R-high-CHOP followed by CHASER, LEED and auto-HSCT; DLBCL-R-CHOP in JCOG0601 consisting of weekly rituximab plus CHOP versus R-CHOP; BD/TD in JCOG0904 consisting of bortezomib plus dexamethasone versus thalidomide plus dexamethasone; ATL-allo-HSCT in JCOG0907 consisting of mLSG15 followed by allo-HSCT; DLBCL-ASCT in JCOG0908 consists of R-CHOP-14 versus R-CHOP-14 followed by CHASER as induction therapy prior to LEED and auto-HSCT.

(U-NHL). After a median follow-up of 56 months, the estimated overall 5-year OS rate was 48%: 60% in B-NHL, 45% in U-NHL, 35% in PTCL and 12% in ATL (Fig. 2). Unfavorable factors influencing OS that remained independently significant in Cox's analyses were clinical diagnosis of ATL, total number of involved lesions ≥ 4 , C-reactive protein-positivity and Eastern Cooperative Oncology Group PS ≥ 2 .

JCOG8701 led to the following conclusions: (i) T-cell phenotype was an important pretreatment variable for aggressive NHL in Japan, and (ii) LSG4 protocol was effective against B-NHL. Since the clinical diagnosis of ATL was an independent unfavorable factor, ATL patients were excluded from subsequent JCOG trials for aggressive NHL, but LSG has started clinical trials specialized for ATL since then.




Figure 2. The Kaplan–Meier estimate of overall survival for all patients treated with LSG4 in relation to disease entity. ATL, adult T-cell leukemia–lymphoma.

JCOG9002, a randomized Phase III study, evaluated the dose-intensification strategy for doxorubicin and cyclophosphamide in the third-generation multiagent combination chemotherapy, LSG9 (VEPA-B/FEPP-AB/M-FEPA every 10 weeks; three courses, 28 weeks in total), when compared with second-generation combination chemotherapy, modified LSG4 (mLSG4) (VEPA-B/FEPP-B/M-FEPA every 14 weeks; four courses, 54 weeks in total) (14). Planned dose intensity (DI)/week of DOX and CPA was 1.9- and 1.5-fold higher in LSG9 than in mLSG4, respectively. Median actual DIs of DOX and CPA were 1.6- and 1.2-fold higher in LSG9 than in mLSG4, respectively, with no difference in 5-year OS and the %CR, revealing no survival benefit of the DI strategy.

In 1993, an intergroup US Phase III study revealed that CHOP remained the standard therapy for aggressive NHL when compared with second- and third-generation regimens (15). Also, the international prognostic index (IPI) for patients with aggressive NHL was developed (16). Based on these findings, JCOG-LSG changed the treatment strategy for aggressive NHL from the multiagent chemotherapies to the dose intensification of key agents, and initiated several Phase II studies of regimens based on CHOP for patients divided by IPI risk grouping. Among them, JCOG9508, a Phase II study of standard CHOP every 3 weeks for low and low-intermediate (L/LI)-risk patients with advanced aggressive NHL, revealed that the full dose of CHOP was feasible and effective for Japanese patients as for westerners (17).

JCOG9505, a randomized Phase II study of CHOP every 2 weeks (CHOP-14) and dose-escalated CHOP both supported with the prophylactic use of G-CSF in high-intermediate and high (HI/H)-risk aggressive NHL, revealed that the former was more promising with similar %CR and progression-free survival (PFS) rates, but lower toxicity (18). Following the results of JCOG9505, a randomized Phase III (JCOG9809) study comparing CHOP-14 with CHOP-21 in patients newly diagnosed with advanced-stage aggressive NHL at all IPI risk was conducted (19,20). A planned interim analysis revealed that dose intensification with

interval shortening of CHOP did not prolong PFS in advanced, aggressive NHL, resulting in an early stop to the study (19), and long-term follow-up confirmed the results (20). There were no remarkable differences in PFS or OS between the two arms. Secondary malignancies, including myelodysplastic syndrome, were significantly more frequent in the CHOP-14 arm.

Since around 2000, rituximab (R), an anti-CD20 monoclonal antibody, has changed the treatment strategy for all CD20-expressing B-cell neoplasms including diffuse large B-cell lymphoma (DLBCL), follicular lymphoma (FL) and chronic lymphocytic leukemia (21). Six to eight courses of rituximab plus CHOP (R-CHOP) every 3 weeks (R-CHOP-21) is now the standard therapy for DLBCL of all risk groups (22). To evaluate the efficacy of DI of rituximab, a randomized Phase II/III study, JCOG0601, is now ongoing comparing the standard with weekly R-CHOP for DLBCL, based on the results of a pharmacokinetic analysis of rituximab monotherapy in a Phase II study for relapsed or refractory aggressive B-NHL (23).

High-dose chemotherapy (HDC) supported with auto-HSCT has been evaluated after induction chemotherapy to improve the prognosis for poor-risk aggressive NHL. However, the results have been controversial in the era before rituximab (24). The US intergroup has conducted a randomized Phase III study evaluating the efficacy of the addition of HDC/auto-SCT after R-CHOP for high-risk DLBCL (25). Considering the next step in the clinical trial for poor-risk DLBCL, JCOG-LSG is now conducting a randomized Phase II study of R-CHOP every 2 weeks (R-CHOP-14) versus R-CHOP-14 followed by CHASER (cyclophosphamide, cytarabine, dexamethasone, etoposide, rituximab) as induction therapy prior to HDC, LEED (melphalan, cyclophosphamide, etoposide, dexamethasone) and auto-HSCT in poor-risk DLBCL (JCOG0908).

ADULT T-CELL LEUKEMIA–LYMPHOMA

JCOG-LSG has consecutively studied the treatment of ATL, which was discovered as a new disease entity just before the establishment of LSG. The earlier trials revealed a poor prognosis of ATL when compared with the other aggressive NHL (JCOG7801, 8101, 8701) (11,13,26). Then, a nationwide survey in 854 patients with ATL in Japan revealed that the major prognostic factors were advanced PS, high lactic dehydrogenase (LDH) level, age of 40 years or more, more than three involved lesions and hypercalcemia by multivariate analysis (27). Also, a classification of clinical subtypes into acute, lymphoma, chronic and smoldering types was proposed based on prognostic factors and clinical features of the disease (28). This subtype classification has been reported to be reproducible for predicting prognosis and has been widely applied for treatment decisions. Recently, a treatment strategy based on the clinical subtype classification and prognostic factors was suggested, including a watchful waiting approach, chemotherapy, antiviral therapy,

allo-HSCT and targeted therapies for clinical trials and clinical practice (29).

The disappointing results with conventional chemotherapies in the 1980s and the proposal for a subtype classification of ATL have led to a search for new active agents focusing on aggressive ATL in JCOG-LSG. The first Phase II study of combination chemotherapy with pentostatin (2'-deoxycoformycin, an inhibitor of adenosine deaminase) was conducted exclusively against aggressive ATL, based on the promising results of pentostatin monotherapy for relapsed or refractory ATL patients (30). However, the results were disappointing with a median survival time (MST) of 7 months similar to previous studies by JCOG-LSG (31). The next Phase II trial (JCOG9303) consisting of vincristine, cyclophosphamide, doxorubicin and prednisone (VCAP); doxorubicin, ranimustine and prednisone (AMP); and vindesine, etoposide, carboplatin and prednisone (VECP) intensified with the prophylactic use of G-CSF revealed a promising response rate and MST superior to those obtained by our previous trials, despite considerable hematological toxicity (32). Based on the promising results of JCOG9303, we conducted a Phase III trial comparing modified (m)LSG15 (VCAP-AMP-VECP) with CHOP-14 both supported with G-CSF and intrathecal prophylaxis. The longer survival at 3 years and higher %CR with VCAP-AMP-VECP compared with CHOP-14 suggest that the former is a more effective regimen at the expense of greater toxicity, providing the basis for future investigations in the treatment of ATL (33). However, the MST of 13 months still compares unfavorably to other hematologic malignancies.

Allo-HSCT is now recommended for the treatment of young patients with aggressive ATL (29). To evaluate the promising efficacy of allo-HSCT, possibly associated with a graft-versus-ATL effect, especially in view of a comparison with intensive chemotherapy, a prospective multicenter Phase II study of mLSG15 chemotherapy followed by allo-HSCT, comparing the results with historical control in JCOG9801, has been initiated as JCOG0907.

A combination of interferon- α (IFN) and zidovudine (AZT) was reported as promising for the treatment of ATL in small Phase II trials in 1995 from the USA and Europe (34-36). Recently, in a worldwide retrospective analysis, it was reported that this combination might be effective especially for indolent ATL when compared with watchful waiting (37). A prospective Phase III study evaluating the efficacy of IFN/AZT when compared with watchful waiting for indolent ATL is to be initiated (JCOG PC908) under the highly advanced medical technology assessment system because IFN and AZT are not covered for ATL by the National Health Insurance in Japan.

LYMPHOBLASTIC LYMPHOMA/ACUTE LYMPHOBLASTIC LEUKEMIA

Lymphoblastic lymphoma (LBL)/acute lymphoblastic leukemia (ALL) is a malignancy of immature T/B lymphoblasts and takes an acute and aggressive course affecting relatively

young individuals. Treatment of child ALL/LBL has much advanced. In contrast, advances for adults have been modest.

JCOG7801 and JCOG8101 revealed that T-LBL and ATL had a poor prognosis compared with other NHLs. Then, a Phase II study of a short-term, combination chemotherapy without maintenance therapy (JCOG8702) for LBL/ALL revealed that a fraction of adult patients with the disease were curable with a short-term, six-drug chemotherapy regimen (38). The next Phase II study (JCOG9004), G-CSF-supported, intensive post-remission chemotherapy and subsequent allo/auto-SCT, revealed that survival and PFS were improved from JCOG8702 in adult ALL and LBL (39). The next chemotherapeutic regimen with the intensified induction and post-remission chemotherapy with auto/allo-HSCT in JCOG9402 was feasible; however, this study failed to show improvements in long-term follow-up results when compared with the historical control JCOG9004 (40).

To further improve the therapeutic outcomes of adults with LBL/ALL, novel strategies are warranted such as risk-adapted treatment for bcr-abl-positive poor prognostic ALL with abl inhibitors. Partly because of the relatively low incidence of LBL/ALL, JCOG-LSG never activated clinical studies after JCO9402.

HODGKIN'S LYMPHOMA

HL is the most chemo/radio-sensitive malignancy among malignant lymphomas, and clinical trials for the disease have steadily produced standard therapies. However, trials are less frequently conducted in Japan and other Asian countries because of a low incidence. Sequential Phase II studies for advanced HL (JCOG8905 and 9305) of C-MOPP (cyclophosphamide, vincristine, procarbazine and prednisone)/ABVd (doxorubicin, bleomycin, vinblastine and dacarbazine) and ABVd, respectively, both with a dose reduction of dacarbazine (250 mg/m²) because of severe emesis in previous studies in Japanese, confirmed a similar efficacy to those from the USA and Europe (41-43). Safety and efficacy profiles of dacarbazine included in C-MOPP/ABVd and ABVd led to the approval of dacarbazine for clinical use covered through the National Health Insurance by MHLW in Japan without industrial trials.

The next Phase II study of ABV deleting dacarbazine with increased dose of doxorubicin followed by IF-RT (JCOG9705) revealed at the interim analysis that the 2-year PFS was significantly inferior to JCOG9305 (ABVd), suggesting that dacarbazine is a key agent for the treatment of HL (44).

A recent meta-analysis of the two JCOG studies in HL revealed two independent factors for OS, male and an elevated serum LDH, after a multivariate analysis (JCOG0108A) (45). Partly because of the low incidence of HL in Japanese, JCOG-LSG never conducted clinical studies after JCO9705. Recent studies from westerners revealed the efficacy of further risk-adaptive treatment for HL, lower dose of chemo/radio-therapy for those at low risk and more

intensive chemotherapy for those at high risk (46). New agents for HL include anti-CD30 monoclonal antibodies. Furthermore, therapy adjustment after an interim analysis of the response by F-fluorodeoxyglucose positron emission tomography-computed tomography (PET-CT) is now suggested. JCOG-LSG is now planning a new trial for HL including PET/CT scans.

MULTIPLE MYELOMA

MM is a progressive and incurable malignancy of plasma cells affecting mainly aged individuals. Alkylators and steroids have been the key drugs for remission induction, but MST was around 3 years without a plateau in the survival curve (47). A Phase II study of COP (cyclophosphamide, prednisolone)-MP (melphalan, prednisolone) for untreated overt MM patients (JCOG8906) revealed a similar efficacy to those from the USA and Europe (48). A subsequent randomized Phase III study comparing modified (m)COP-MP with/without ranimustine for untreated overt MM (JCOG9301) revealed that addition of ranimustine to mCOP/MP has no benefit for survival, despite improving the response rate and PFS, similar to findings of other studies evaluating the addition of new agents to alkylators and steroids in MM (49).

Both Phase III and II studies on untreated overt MM patients who were ineligible for HDC/auto-HSCT and eligible, respectively (JCOG0112 and JCOG0005-DI), were terminated early because of poor patient accrual and the results of a planned interim analysis, respectively (50). The planned interim analysis of JCOG0005-DI, when 16 of the 50 planned patients were enrolled, revealed that the primary endpoint, response rate, was less than the lower threshold associated with violation in two patients who underwent allo-HSCT instead of scheduled auto-SCT because of availability of HLA-matched sibling donors. Since 2000, several promising new agents have been incorporated in standard therapy for the disease (51). Following the results, LSG is now conducting a randomized Phase II study of bortezomib, a proteasome inhibitor, plus dexamethasone versus thalidomide, an immune modulator, plus dexamethasone for relapsed or refractory MM (JCOG0904).

INDOLENT B-CELL NHL

Advanced FL and other low-grade B-cell lymphomas are clinically indolent but non-curable diseases in most patients. The prognosis for lymphomas has been improved by adding rituximab to chemotherapy (21). However, the optimal combination schedule of chemotherapy and rituximab has not been elucidated. We attempted to determine whether patients with indolent B-cell NHL would have long-term benefits from G-CSF-supported, dose-dense immune-chemotherapy which potentiates the antibody-dependent cell-mediated cytotoxicity of rituximab by comparing R-CHOP-21 versus R-CHOP-14 (JCOG0203) (52). However, the dose-dense

strategy failed to improve PFS at the median follow-up time of 5.2 years. We are now planning to follow-up the patients enrolled in this study to further evaluate the long-term prognosis of this indolent disease and potential late complications including secondary malignancies.

MANTLE CELL LYMPHOMA

Mantle cell lymphoma (MCL) is a progressive, non-curable and relatively rare B-cell lymphoma derived from mantle zone B-cell with BCL1 translocation. In contrast to DLBCL and FL, addition of rituximab to CHOP did not improve the survival of MCL and HDC/ASCT has been reported as promising (53). Therefore, LSG is now conducting a single-arm Phase II study of R-high-CHOP followed by CHASER and HDC, LEED and auto-HSCT for previously untreated advanced-stage MCL (JCOG0406).

LOCALIZED NASAL NK/T-CELL LYMPHOMA

Localized nasal NK/T-cell lymphoma is a refractory lymphoma relatively frequent in East Asia. Both the international project on PTCL and JCOG meta-analysis on T-NHL (JCOG0902A) revealed that the diagnosis of NK/T-cell lymphoma was poor (54,55). A Phase I/II study (JCOG0211-DI) of concurrent radiotherapy (50 Gy) and three courses of dexamethasone, etoposide, ifosfamide and carboplatin (DeVIC) consisting of multidrug resistance-non-related agents revealed that 2/3 dose of DeVIC and radiation was a safe and effective treatment when compared with a historical control of radiotherapy alone (56). A correlative study is ongoing to elucidate risk factors for relapse because PFS was not sufficient.

FUTURE ISSUES FOR JCOG-LSG

JCOG-LSG has conducted clinical trials for aggressive NHL since 1970, which has been divided into DLBCL, MCL, LBL/ALL, ATL and NK/T-NHL later, HL, MM and indolent-B-NHL, as shown in Fig. 1, to evaluate combined modality, dose intensification and incorporation of new agents in multidisciplinary treatment for lymphoid malignancies. LSG, now consisting of 47 institutions, is a relatively large group in JCOG, using much of the resources of the JCOG Data Center and Committees. JCOG-LSG initiated several studies independent from the JCOG-Data Center and supported by its own data center. However, numbers of patient enrollment in LSG have decreased in the last several years mainly due to the small number of ongoing trials. It takes longer to activate LSG protocols when compared with those by other cancer groups in JCOG mainly because of many disease entities, diverse prognosis and complex response criteria in each major disease entities, such as DLBCL, ATL and MM. Lymphoid malignancies are relatively rare; however, the spectrum is diverse consisting of 81

disease entities from indolent to aggressive in the WHO classification of 2008 (1). One way to conduct future LSG trials is to focus more on each disease entity as in the case of ATL and ENK/TML. The other is grouping the entities by treating modalities as in the case of several CD20-expressing low-grade B-cell lymphomas with rituximab-containing chemotherapy. It is desirable that clinical trials in LSG be based on disease entity, and if possible with risk grouping as in the case of trials for DLBCL and ATL. However, most of the diseases are rare and some of them take similar clinical courses including prognosis and response to therapies. On this issue, peripheral T-cell lymphomas other than ATL and T/NK ML, and low-grade B-cell lymphomas are the major two categories of disease-entity grouping.

Not only the complexity in lymphoma classification mentioned above, but also that in response criteria for lymphoma, ATL and MM which are, respectively, distinct from RECIST (response evaluation criteria in solid tumors) for other solid cancers has made the trials in LSG difficult. Recent revised criteria for lymphoma, which are applied in the JCOG0601 study, incorporating the PET/CT scan for decision of CR, might reduce the difficulty (57).

As described, JCOG-LSG has conducted clinical trials for establishing standard therapies. To keep and further upgrade the originality of JCOG-LSG trials in relation to similar cooperative study groups in the USA and Europe, several points are important including major target diseases, risk grouping for stratification and major phase of the trials. Since its establishment, JCOG-LSG has consecutively focused on diseases relatively common in Japan such as DLBCL, MM, ATL and NK/T-NHL. Recent advances in molecular-targeting therapy introduced many promising new agents for the diseases and other lymphoid malignancies. This promotes research on lymphoid malignancies for the early development of a new-standard combination therapy with the new agents in Japan. However, for the evaluation of new agent-combining treatment in Japan, JCOG-LSG should go side by side with those through industry-supported new agent trials to contribute to further improvement in the treatment of lymphoid malignancies with less lag from foreign developments. For instance, the highly advanced medical technology assessment system, which was enacted recently, would be one way of reducing the lag in Japan. Multigroup trials, including global ones and bridging studies, are another way.

Correlative studies in clinical trials have changed the next step of stratified treatments. For instance, ATL and T-LBL patients were excluded from subsequent JCOG trials for aggressive NHL since their clinical diagnosis was found to be poor in early trials (11–13). Future correlative studies in JCOG-LSG, retrospective and prospective and pathological and molecular analysis, should change the stratification of the clinical trials in future for risk-adaptive treatment. For that purpose, a banking system, which is now being established in JCOG for blood and tissue samples, is warranted.

Fortunately, in the case of lymphoid malignancies, relatively easy access to samples of the neoplasm can promote correlative studies.

Lastly, as a member of cancer groups in JCOG, LSG will continue efforts to produce valuable and reliable evidence for the improvement of therapy for patients with lymphoid malignancies as rapidly as possible.

Acknowledgements

We are grateful to Drs K. Tajima and H. Fukuda, and the members of the JCOG Statistical/Data Center, for data management and analyses of JCOG trials. We are greatly appreciative of all members of the JCOG-LSG.

Funding

JCOG-LSG has been supported by Grants-in-Aid for Cancer Research from the Ministry of Health, Labor and Welfare of Japan since 1978 (53S-1, 56S-1, 59S-1, 62S-1, 2S-1, 5S-1, 8S-1, 11S-1, 14S-1, 17S-1, 20S-1, 23A-17).

Conflict of interest statement

None declared.

References

1. Swerdlow SH, Campo E, Harris NL, Jaffe ES, Pileri SA, Stein H, et al., editors. *WHO Classification of Tumour of Haematopoietic and Lymphoid Tissues*. 4th edn. Lyon: IARC Press 2008;281–4.
2. Uchiyama T, Yodoi J, Sagawa K, Takatsuki K, Uchino H. Adult T-cell leukemia: clinical and hematologic features of 16 cases. *Blood* 1977;50:481–92.
3. Poesz BJ, Ruscetti FW, Gazdar AF, Bunn PA, Minna JD, Gallo RC. Detection and isolation of type C retrovirus particles from fresh and cultured lymphocytes of a patient with cutaneous T-cell lymphoma. *Proc Natl Acad Sci USA* 1980;77:7415–9.
4. Hinuma Y, Nagata K, Hanaoka M, Nakai M, Matsumoto T, Kinoshita KI, et al. Adult T-cell leukemia: antigen in an ATL cell line and detection of antibodies to the antigen in human sera. *Proc Natl Acad Sci USA* 1981;78:6476–80.
5. Yoshida M, Miyoshi I, Hinuma Y. Isolation and characterization of retrovirus from cell lines of human adult T-cell leukemia and its implication in the disease. *Proc Natl Acad Sci USA* 1982;79:2031–5.
6. Fisher RI. Advanced insights into the biology of malignant lymphomas. *J Clin Oncol* 2011;29:1799–800.
7. Richardson P. Multiple myeloma and amyloidosis: an evolving treatment paradigm toward the goal of tailored therapy. *J Clin Oncol* 2011;29:1801–2.
8. Shimoyama M, Fukuda H, Saijo N, Yamaguchi N. Japan Clinical Oncology Group (JCOG). *Jpn J Clin Oncol* 1998;28:158–62.
9. Tobinai K, Hotta T. Clinical trials for malignant lymphoma in Japan. *Jpn J Clin Oncol* 2004;34:369–78.
10. Fukuda H. Development of cancer cooperative groups in Japan. *Jpn J Clin Oncol* 2010;40:881–90.
11. Shimoyama M, Ichimaru M, Yunoki K, Oota K, Ogawa M. Final results of cooperative study of VEPA [vincristine, cyclophosphamide (endoxan), prednisolone and adriamycin] therapy in advanced adult non-Hodgkin's lymphoma: relation between T- or B-cell phenotype and response. *Jpn J Clin Oncol* 1982;12:227–37.

12. Shimoyama M, Ota K, Kikuchi M, Yunoki K, Konda S, Takatsuki K, et al. Chemotherapeutic results and prognostic factors of patients with advanced non-Hodgkin's lymphoma treated with VEPA or VEPA-M. *J Clin Oncol* 1988;6:128-41.
13. Tobinai K, Shimoyama M, Minato K, Shirakawa S, Kobayashi T, Hotta T, et al. Japan Clinical Oncology Group phase II trial of second-generation 'LSG4 protocol' in aggressive T- and B-lymphoma: a new predictive model for T- and B-lymphoma. *Proc Am Soc Clin Oncol* 1994;13:378a (Abstract).
14. Kinoshita T, Hotta T, Tobinai K, Kobayashi T, Ishizuka N, Tomonaga M, et al. A randomized controlled trial investigating the survival benefit of dose-intensified multidrug combination chemotherapy (LSG9) for intermediate- or high-grade non-Hodgkin's lymphoma: Japan Clinical Oncology Group Study 9002. *Int J Hematol* 2004;80:341-50.
15. Fisher RI, Gaynor ER, Dahlborg S, Oken MM, Grogan TM, Mize EM, et al. Comparison of a standard regimen (CHOP) with three intensive chemotherapy regimens for advanced non-Hodgkin's lymphoma. *N Engl J Med* 1993;328:1002-6.
16. The International Non-Hodgkin's Lymphoma Prognostic Factors Project. A predictive model for aggressive non-Hodgkin's lymphoma. *N Engl J Med* 1993;329:987-94.
17. Ogura M, Kagami Y, Itoh K, Tobinai K, Fukuda H, Mukai K, et al. Standard CHOP therapy for low (L) or low-intermediate (L-I) risk patients (pts) with aggressive non-Hodgkin's lymphoma (NHL): A multicenter phase II study by Japan Clinical Oncology Group (JCOG 9508). *Proc Am Soc Hem* 2004;22:6703 (Abstract).
18. Itoh K, Ohtsu T, Fukuda H, Sasaki Y, Ogura M, Morishima Y, et al. Randomized phase II study of biweekly CHOP and dose-escalated CHOP with prophylactic use of lenograstim (glycosylated G-CSF) in aggressive non-Hodgkin's lymphoma: Japan Clinical Oncology Group Study 9505. *Ann Oncol* 2002;13:1347-55.
19. Hotta T, Shimakura Y, Ishizuka N, Fukuda H, Itoh K, Morishima Y, et al. The Lymphoma Study Group of the JCOG. Randomized phase III study of standard CHOP (S-CHOP) versus biweekly CHOP (Bi-CHOP) in aggressive non-Hodgkin's lymphoma (NHL): Japan Clinical Oncology Group study, JCOG 9809. *Proc Am Soc Clin Oncol* 2003;21:2271a (Abstract).
20. Ohmachi K, Tobinai K, Kobayashi Y, Itoh K, Nakata M, Shibata T, et al. Phase III trial of CHOP-21 versus CHOP-14 for aggressive non-Hodgkin's lymphoma: final results of the Japan Clinical Oncology Group Study, JCOG 9809. *Ann Oncol* 2011;22:1382-91.
21. Czuczman MS, Gregory SA. The future of CD20 monoclonal antibody therapy in B-cell malignancies. *Leuk Lymphoma* 2010;51:983-94.
22. Coiffier B, Lepage E, Briere J, Herbrecht R, Tilly H, Bouabdallah R, et al. CHOP chemotherapy plus rituximab compared with CHOP alone in elderly patients with diffuse large-B-cell lymphoma. *N Engl J Med* 2002;346:235-42.
23. Tobinai K, Igarashi T, Itoh K, Kobayashi Y, Taniwaki M, Ogura M, et al. Japanese multicenter phase II and pharmacokinetic study of rituximab in relapsed or refractory patients with aggressive B-cell lymphoma. *Ann Oncol* 2004;15:821-30.
24. Shipp MA, Abeloff MD, Antman KH, Carroll G, Hagenbeek A, Loeffler M, et al. International Consensus Conference on High-Dose Therapy with Hematopoietic Stem Cell Transplantation in Aggressive Non-Hodgkin's Lymphomas: report of the jury. *J Clin Oncol* 1999;17:423-9.
25. Stiff PJ, Unger JM, Cook J, Constine LS, Couban S, Shea TC, et al. Randomized phase III U.S./Canadian intergroup trial (SWOG S9704) comparing CHOP ± R for eight cycles to CHOP ± R for six cycles followed by autotransplant for patients with high-intermediate (H-Int) or high IPI grade diffuse aggressive non-Hodgkin's lymphoma (NHL). *Proc Am Soc Clin Oncol* 2011;29:504a (Abstract).
26. Shimoyama M, Ota K, Kikuchi M, Yunoki K, Konda S, Takatsuki K, et al. Major prognostic factors of adult patients with advanced T-cell lymphoma/leukemia. *J Clin Oncol* 1988;6:1088-97.
27. Major prognostic factors of patients with adult T-cell leukemia-lymphoma: a cooperative study. Lymphoma Study Group (1984-1987). *Leuk Res* 1991;15:81-90.
28. Shimoyama M. Diagnostic criteria and classification of clinical subtypes of adult T-cell leukaemia-lymphoma: a report from the Lymphoma Study Group (1984-87). *Br J Haematol* 1991;79:428-37.
29. Tsukasaki K, Hermine O, Bazarbachi A, Ratner L, Ramos JC, Harrington W, Jr, et al. Definition, prognostic factors, treatment, and response criteria of adult T-cell leukemia-lymphoma: a proposal from an international consensus meeting. *J Clin Oncol* 2009;27:453-9.
30. Tobinai K, Shimoyama M, Inoue S, Takayasu S, Mikuni C, Kozuru M, et al. Phase I study of YK-176 (2-deoxycoformycin) in patients with adult T-cell leukemia-lymphoma. *Jpn J Clin Oncol* 1992;22:164-71.
31. Tsukasaki K, Tobinai K, Shimoyama M, Kozuru M, Uike N, Yamada Y, et al. Deoxycoformycin-containing combination chemotherapy for adult T-cell leukemia-lymphoma: Japan Clinical Oncology Group study (JCOG9109). *Int J Hematol* 2003;77:164-70.
32. Yamada Y, Tomonaga M, Fukuda H, Hanada S, Utsunomiya A, Tara M, et al. A new G-CSF-supported combination chemotherapy, LSG15, for adult T-cell leukemia-lymphoma (ATL): Japan Clinical Oncology Group (JCOG) Study 9303. *Br J Haematol* 2001;113:375-82.
33. Tsukasaki K, Utsunomiya A, Fukuda H, Shibata T, Fukushima T, Takatsuka Y, et al. VCAP-AMP-VECP compared with biweekly CHOP for adult T-cell leukemia-lymphoma: Japan Clinical Oncology Group Study JCOG9801. *J Clin Oncol* 2007;25:5458-64.
34. Gill PS, Harrington W, Kaplan MH, Ribeiro RC, Bennett JM, Liebman HA, et al. Treatment of adult T-cell leukemia-lymphoma with a combination of interferon alfa and zidovudine. *N Engl J Med* 1995;332:1744-8.
35. Hermine O, Bloussard D, Gessain A, Turlure P, Leblond V, Franck N, et al. Treatment of adult T-cell leukemia-lymphoma with zidovudine and interferon alfa. *N Engl J Med* 1995;332:1749-51.
36. Tobinai K, Kobayashi Y, Shimoyama M. Interferon alfa and zidovudine in adult T-cell leukemia-lymphoma. *N Engl J Med* 1995;333:1285-6 (Correspondence).
37. Bazarbachi A, Plumelle Y, Carlos Ramos J, Tortevoye P, Otroek Z, Taylor G, et al. Meta-analysis on the use of zidovudine and interferon-alfa in adult T-cell leukemia/lymphoma showing improved survival in the leukemic subtypes. *J Clin Oncol* 2010;28:4177-83.
38. Kobayashi T, Tobinai K, Shimoyama M, Mikuni C, Konda S, Kozuru M, et al. Long-term follow-up results of adult patients with acute lymphocytic leukemia or lymphoblastic lymphoma treated with short-term, alternating non-cross-resistant chemotherapy: Japan Clinical Oncology Group Study 8702. Lymphoma Study Group. *Jpn J Clin Oncol* 1999;29:340-8.
39. Tobinai K, Takeyama K, Arima F, Aikawa K, Kobayashi T, Hanada S, et al. Phase II study of chemotherapy and stem cell transplantation for adult acute lymphoblastic leukemia or lymphoblastic lymphoma: Japan Clinical Oncology Group study 9004. *Cancer Science* 2007;98:1350-7.
40. Azuma T, Kensei Tobinai K, Takeyama K, Shibata T, Fukuda H, Kawano F, et al. Phase II study of chemotherapy (Cx) and stem cell transplantation (SCT) for adult acute lymphoblastic leukemia (ALL) and lymphoblastic lymphoma (LBL): long-term follow-up results of Japan Clinical Oncology Group (JCOG) Study 9402. *Proc Am Soc Hem* 2009;114:3087 (Abstract).
41. Takenaka T, Mikuni C, Miura A, Sasaki T, Suzuki H, Hotta T, et al. Alternating combination chemotherapy C-MOPP (cyclophosphamide, vincristine, procarbazine, prednisone) and ABVd (adriamycin, bleomycin, vinblastine, dacarbazine) in clinical stage II-IV Hodgkin's disease: a multicenter phase II study (JCOG 8905). *Jpn J Clin Oncol* 2000;30:146-52.
42. Ogura M, Itoh K, Kinoshita T, Fukuda H, Takenaka T, Ohtsu T, et al. Phase II study of ABVd therapy for newly diagnosed clinical stage II-IV Hodgkin lymphoma: Japan Clinical Oncology Group study (JCOG 9305). *Int J Hematol* 2010;92:713-24.
43. Canellos GP, Anderson JR, Propert KJ, Nissen N, Cooper MR, Henderson ES, et al. Chemotherapy of advanced Hodgkin's disease with MOPP, ABVD, or MOPP alternating with ABVD. *N Engl J Med* 1992;327:1478-84.
44. Ogura M, Morishima Y, Itoh K, Ishizawa K, Kobayashi Y, Tobinai K, et al. Dacarbazine cannot be deleted from ABVD therapy for advanced Hodgkin lymphoma: Japan Clinical Oncology Group (JCOG) Study 9705. *Proc Am Soc Clin Oncol* 2003;22:573a (Abstract).
45. Itoh K, Kinoshita T, Watanabe T, Yoshimura K, Okamoto R, Chou T, et al. Prognostic analysis and a new risk model for Hodgkin lymphoma in Japan. *Int J Hematol* 2010;91:446-55.
46. Blum KA. Upcoming diagnostic and therapeutic developments in classical Hodgkin's lymphoma. *Hematology Am Soc Hematol Educ Program* 2010;2010:93-100.

47. Gregory WM, Richards MA, Malpas JS. Combination chemotherapy versus melphalan and prednisolone in the treatment of multiple myeloma: an overview of published trials. *J Clin Oncol* 1992;10:334–42.
48. Takenaka T, Shirakawa S, Mikuni C, Sai T, Shimamoto Y, Araki K, et al. Alternating combination chemotherapy COP (cyclophosphamide, vincristine, prednisone) and MP (melphalan, prednisone) in multiple myeloma: a multicenter phase II study (JCOG8906). Lymphoma Study Group of the Japan Clinical Oncology Group. *Jpn J Clin Oncol* 1999;29:485–9.
49. Takenaka T, Itoh K, Suzuki T, Utsunomiya A, Matsuda S, Chou T, et al. Phase III study of ranimustine, cyclophosphamide, vincristine, melphalan, and prednisolone (MCNU-COP/MP) versus modified COP/MP in multiple myeloma: a Japan clinical oncology group study, JCOG 9301. *Int J Hematol* 2004;79:165–73.
50. Chou T, Tobinai K, Uike N, Asakawa T, Saito I, Fukuda H, et al. Melphalan–prednisolone and vincristine–doxorubicin–dexamethasone chemotherapy followed by prednisolone/interferon maintenance therapy for multiple myeloma: Japan Clinical Oncology Group Study, JCOG0112. *Jpn J Clin Oncol* 2011;41:586–9.
51. Palumbo A, Anderson K. Multiple myeloma. *N Engl J Med* 2011;364:1046–60.
52. Watanabe T, Tobinai K, Shibata T, Tsukasaki K, Morishima Y, Maseki N, et al. Phase II/III study of R-CHOP-21 versus R-CHOP-14 for untreated indolent B-cell non-Hodgkin's lymphoma: JCOG 0203 trial. *J Clin Oncol* 2011;29:3990–8.
53. Williams ME, Connors JM, Dreyling MH, Gascoyne RD, Kahl BS, Leonard JP, et al. Mantle cell lymphoma: report of the 2010 Mantle Cell Lymphoma Consortium Workshop. *Leuk Lymphoma* 2011;52:24–33.
54. Vose J, Armitage J, Weisenburger D. International T-Cell Lymphoma Project. International peripheral T-cell and natural killer/T-cell lymphoma study: pathology findings and clinical outcomes. *J Clin Oncol* 2008;26:4124–30.
55. Watanabe T, Kinoshita T, Itoh K, Yoshimura K, Ogura M, Kagami Y, et al. Pretreatment total serum protein is a significant prognostic factor for the outcome of patients with peripheral T/natural killer-cell lymphomas. *Leuk Lymphoma* 2010;51:813–21.
56. Yamaguchi M, Tobinai K, Oguchi M, Ishizuka N, Kobayashi Y, Isobe Y, et al. Phase I/II study of concurrent chemoradiotherapy for localized nasal natural killer/T-cell lymphoma: Japan Clinical Oncology Group Study JCOG0211. *J Clin Oncol* 2009;27:5594–600.
57. Cheson BD, Pfistner B, Juweid ME, Gascoyne RD, Specht L, Horning SJ, et al. Revised response criteria for malignant lymphoma. *J Clin Oncol* 2007;25:579–86.
58. Mizoroki F, Hirose Y, Sano M, Fukuda H, Tobinai K, Nakata M, et al. A phase II study of VEPA/FEPP chemotherapy for aggressive lymphoma in elderly patients: Japan Clinical Oncology Group Study JCOG9203. *Int J Hematol* 2006;83:55–62.
59. Yoshino T, Matsuno Y, Makamura S, Ohshima K, Mukai K, Kikuchi M, et al. Effectiveness of an educational program for improving the accuracy of histopathologic diagnosis of malignant lymphoma. *Proc Jpn Soc Lymphoreticular Tissue Res* 2003;43:88 (Abstract).

Phase II Study of Intensive Post-remission Chemotherapy and Stem Cell Transplantation for Adult Acute Lymphoblastic Leukemia and Lymphoblastic Lymphoma: Japan Clinical Oncology Group Study, JCOG9402

Teruhisa Azuma^{1,11}, Kensei Tobinai^{1,*}, Kunihiko Takeyama^{1,12}, Taro Shibata², Michihiro Hidaka³, Mitsutoshi Kurosawa⁴, Masaharu Kasai⁵, Takaaki Chou⁶, Noriyasu Fukushima⁷, Kiyoshi Mukai^{8,13}, Kunihiro Tsukasaki⁹, Masanori Shimoyama^{1,10} and Japan Clinical Oncology Group-Lymphoma Study Group (JCOG-LSG), Tokyo, Japan

¹National Cancer Center Hospital, Tokyo, ²JCOG Data Center, National Cancer Center, Tokyo, ³Kumamoto Medical Center, Kumamoto, ⁴Hokkaido Cancer Center, Sapporo, ⁵Sapporo Hokuyu Hospital, Sapporo, ⁶Niigata Cancer Center, Niigata, ⁷Saga University, Saga, ⁸Tokyo Medical University, Tokyo, ⁹Nagasaki University, Nagasaki and ¹⁰Nagoya Medical Center, Nagoya, Japan

¹¹Present address: Tenri Hospital, Tenri, Japan.

¹²Present address: Bristol-Myers Squibb, Tokyo, Japan.

¹³Present address: Saiseikai Central Hospital, Tokyo, Japan.

*For reprints and all correspondence: Kensei Tobinai, Department of Hematology, and Hematopoietic Stem Cell Transplantation, National Cancer Center Hospital, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan. E-mail: ktobinai@ncc.go.jp

Received August 1, 2011; accepted February 13, 2012

Objective: To evaluate the efficacy and safety of intensive post-remission chemotherapy for untreated patients aged 15–69 years with adult acute lymphoblastic leukemia and lymphoblastic lymphoma in a multicenter Phase II study.

Methods: The chemotherapy regimen consisted of induction, post-remission and maintenance for 2 years. The primary endpoint was 5-year progression-free survival, and secondary endpoints included complete remission rate, overall survival and adverse events. Among 115 patients enrolled, 108 eligible patients [median age, 33.5 years (range, 15–69)] including 96 acute lymphoblastic leukemia and 12 lymphoblastic lymphoma were assessed. Other major characteristics were male 50%, T-cell phenotype 21%, Philadelphia chromosome 22%, B-symptom+ 35% and performance status 2/3 22%.

Results: Eighty-seven patients achieved complete remission (81%; 95% confidence interval 72–88%), while five (5%) died during the chemotherapy protocol. The median overall survival and progression-free survival were 1.8 years (95% confidence interval, 1.5–2.6) and 1.2 years (95% confidence interval, 0.8–1.6), respectively. Their 5-year overall survival and progression-free survival were 29 and 28%, respectively. The 5-year overall survival of 31 patients who underwent allogeneic ($n = 19$) or autologous ($n = 12$) stem cell transplantation during first complete response was 51%. Major non-hematologic toxicities of Grade 3 or greater were infections (21%) and pulmonary complications (6%). When compared with the investigators' previous Phase II trials, JCOG9402 improved progression-free survival and overall survival when compared with JCOG8702; however, it did not show improvement when compared with JCOG9004.

Conclusions: Although the intensified induction and post-remission chemotherapy was feasible and 28% of the patients with adult acute lymphoblastic leukemia or lymphoblastic lymphoma achieved long-term progression-free survival, JCOG9402 did not show improvement.

Key words: chemo-haematopoietic – clinical trials – hematol-leukemia/lymphoma – hematol-transplantation

INTRODUCTION

Acute lymphoblastic leukemia (ALL) and lymphoblastic lymphoma (LBL) are categorized as precursor B- or T-cell malignancy (1) and show overlapping clinical features, including leukemic manifestation and a high prevalence of central nervous system (CNS) involvement. ALL and LBL of the T-cell phenotype are frequently accompanied by a mediastinal mass (1,2). Within each cell lineage, there is clinical and biological overlap between ALL and LBL. Based on these aspects, patients with ALL and those with LBL have been treated with similar chemotherapy regimens (3–7). Generally, however, the prognosis of adult patients is unfavorable compared with pediatric patients (3–11).

Previously, the Lymphoma Study Group of the Japan Clinical Oncology Group (JCOG-LSG) conducted two Phase II studies against ALL and LBL, JCOG8702 and JCOG9004 (12,13). JCOG8702 was designed to improve the therapeutic results of adult ALL and LBL, using VEPA [vincristine (VCR), cyclophosphamide (CPA), prednisolone (PSL) and doxorubicin (DXR)], L-asparaginase (L-asp) and intermediate-dose methotrexate (MTX) without maintenance therapy with oral 6-mercaptopurine (6-MP) and MTX. Although JCOG8702 showed a satisfactory complete remission (CR) rate (%CR) (78%, 36/46), the long-term survival [7-year overall survival (OS), 15%] was inferior to that of the published programs incorporating maintenance therapy (12). The investigators concluded that simplified JCOG8702 therapy without maintenance therapy does not deserve further investigation.

Subsequently, JCOG9004 was designed to mainly evaluate intensive post-remission therapy. The following three strategies were undertaken: (i) intensive post-remission therapy with the prophylactic use of granulocyte colony-stimulating factor (G-CSF); (ii) long-term maintenance therapy including 6-MP and MTX; and (iii) allogeneic and autologous stem cell transplantation (SCT).

Various strategies have been evaluated in the treatment of adult ALL. The Japan Adult Leukemia Study Group (JALSG) evaluated a response-oriented individualized induction therapy in their ALL87 and ALL90 studies (14,15) and intensified induction and consolidation therapy by dose intensification of DXR in their ALL93 and ALL97 studies (16,17), respectively; however, the advantage of these treatment strategies remain unclear. In the early 1990s, especially

at the time of designing this study, the significance of post-remission therapy on the outcome of ALL patients had been recognized (3). In particular, autologous and allogeneic SCT had been thought to be promising (18).

Based on these findings, we planned to conduct this multicenter Phase II study, JCOG9402, to further improve the outcome of adult patients with ALL and LBL by the following strategies: (i) consolidation including cytosine arabinoside (Ara-C); (ii) prophylactic intrathecal MTX (IT-MTX) eight times to prevent a CNS relapse; (iii) interim maintenance with MTX and 6-MP; (iv) intensification consisting of DXR, VCR, PSL, CPA, Ara-C and 6-MP; (v) maintenance consisting of vindesine (VDS), CPA, DXR, PSL, 6-MP and MTX for 60 weeks; (vi) allogeneic SCT in first CR; and (vii) autologous SCT in LBL patients. We evaluated the efficacy and safety of this sequential chemotherapy for untreated adult patients with ALL and LBL, focusing on the long-term follow-up results. The feasibility and efficacy of autologous and allogeneic SCT were also assessed.

PATIENTS AND METHODS

PATIENTS

Patients aged 15–69 years were eligible. ALL and LBL were diagnosed according to the French-American-British (FAB) Classification (19,20) and Working Formulation (21), respectively. Patients with LBL at any stage were eligible. Patients having the following disorders were ineligible: preceding myelodysplastic syndrome, performance status (PS) 4 according to the Eastern Cooperative Oncology Group (ECOG) scale (22), liver dysfunction (liver cirrhosis, total bilirubin ≥ 2.0 mg/dl or hepatic transaminase $\geq 4 \times$ upper normal limits), renal dysfunction (serum creatinine ≥ 2.0 mg/dl), hypoxemia ($\text{PaO}_2 < 60$ mmHg), diabetes mellitus requiring insulin treatment, severe infection, active tuberculosis, cardiac failure, ischemic heart disease, cardiomyopathy, active concurrent cancer, human immunodeficiency virus infection, severe psychiatric disorder, pregnancy or other severe complications. Patients who had received anticancer agents until enrollment were ineligible, but those receiving radiation or glucocorticoids were eligible.

Immunophenotyping for tumor cells was performed at each institution to distinguish B-cell or T-cell origin. Flow

cytometric analysis was conducted on suspended cells. Samples were judged positive if more than 20% of cells showed specific labeling above those of controls. For biopsied lymph node or tumor samples, immunohistochemical analysis was conducted. Even if the biopsied specimen showed histopathologic findings consistent with LBL, patients having 25% tumor cells or more in their bone marrow (BM) were diagnosed as having ALL (2). It was recommended that a pre-treatment peripheral blood (PB), BM or biopsied specimen be subjected to cytogenetic analysis. Biopsy specimens of patients diagnosed with LBL at each institution were submitted to central pathology review. The data of flow cytometric analyses conducted at each institution were utilized in the central pathology review.

Registration was done by a telephone call or fax from participating physicians to the JCOG Statistical or Data Center (1994–97: Statistical Center; 1998–: Data Center) (23). When a patient achieving first CR met the following eligibility criteria for SCT, the physician was requested to register. For allogeneic SCT, patients had to meet the following criteria: (i) age 15–49 years; (ii) human leukocyte antigen (HLA)-matched sibling donor; (iii) continuous CR after consolidation A; (iv) PS of 0 or 1; (v) PaO₂ ≥ 70 mmHg; (vi) cardiac ejection fraction ≥ 50%; (vii) creatinine clearance ≥ 50 ml/min; (viii) written informed consent from each patient and his/her donor. For autologous SCT, patients should meet the following criteria: (i) histopathologic LBL; (ii) no matched sibling donor; (iii) no tumor cells in BM or PB at stem cell harvest, in addition to the above-described criteria (i) and (iii)–(vii) for allogeneic SCT.

The study protocol was approved by the Protocol Review Committee of the JCOG and by the institutional review board at each institution. Before treatment, informed consent was obtained from each patient or his/her family member.

TREATMENTS

CHEMOTHERAPY

The total treatment plan is shown in Fig. 1, and drug doses and administration schedules are listed in Table 1. The treatment regimens consisted of one cycle of induction, one cycle of consolidation A and consolidation B, CNS prophylaxis, interim maintenance, intensification and five cycles of maintenance. During and after administration of L-asparaginase in induction therapy, it was recommended that fresh-frozen plasma or anti-thrombin be given to patients showing a marked decrease in fibrinogen and anti-thrombin III. The total planned duration of treatment was 102 weeks. For patients who did not achieve CR with induction therapy, consolidation A and consolidation B were given. For those who did not achieve CR by then, protocol treatment was terminated. Testicular irradiation or CNS irradiation was not administered prophylactically, but irradiation of a mass lesion initially presenting as over 5 cm in diameter was recommended after consolidation therapy. Because severe complications such as pancreatitis and hyperglycemia occurred, L-asparaginase was withdrawn from consolidation B according to the recommendation of the JCOG Data and Safety Monitoring Committee in September 1996.




Figure 1. Total treatment scheme of the JCOG9402 protocol. ALL, acute lymphoblastic leukemia; LBL, lymphoblastic lymphoma; HLA, human leukocyte antigen; CR, complete remission; IF-RT, involved-field radiotherapy; CNS, central nervous system; BMT, bone marrow transplantation; PBSCT, peripheral blood stem cell transplantation.

Table 1. Drug doses and administration schedules in JCOG9402

Agent	Route	Dose (≥ 60 years old)	Schedule (≥ 60 years old)
Induction			
Vincristine	IV	1.4 mg/m ² up to 2.0 mg total	Days 1, 8, 15, 22, 29
Cyclophosphamide	IV	600 mg/m ² (400 mg/m ²)	Days 1, 15, 29
Prednisolone	PO	40 mg/m ²	Days 1–28, then taper (days 1–7, no taper) ^a
Doxorubicin	IV	40 mg/m ² (30 mg/m ²)	Days 1, 8, 22
L-Asparaginase	IV	6000 U/m ² (4000 U/m ²)	Days 22–28
Methotrexate	IT	15 mg/patient	Day 15
Prednisolone	IT	20 mg/patient	Day 15
Consolidation A			
Daunorubicin	IV	40 mg/m ² (30 mg/m ²)	Days 1, 2
Cytosine arabinoside	CI	75 mg/m ²	Days 1–7
Vindesine	IV	2.4 mg/m ² (2.0 mg/m ²)	Day 1
Prednisolone	PO	40 mg/m ²	Days 1–7, no taper
Methotrexate	IT	15 mg/patient	Day 1
Prednisolone	IT	20 mg/patient	Day 1
Consolidation B			
Cyclophosphamide	IV	1000 mg/m ² (700 mg/m ²)	Days 1, 29
Cytosine arabinoside	CI	75 mg/m ²	Days 1–7, 29–35
6-Mercaptopurine	PO	60 mg/m ²	Days 1–7, 29–35 (no allopurinol)
Vincristine	IV	1.4 mg/m ² up to 2.0 mg total	Days 15, 22, 43, 50
L-Asparaginase ^b	IM/SQ	6000 U/m ² (4000 U/m ²)	Days 15, 18, 22, 25, 43, 46, 50, 53
Methotrexate	IT	15 mg/patient	Days 1, 29
Prednisolone	IT	20 mg/patient	Days 1, 29
CNS prophylaxis and interim maintenance			
6-Mercaptopurine	PO	60 mg/m ²	Days 1–70 (no allopurinol)
Methotrexate	PO	20 mg/m ²	Days 29, 36, 43, 50, 57, 64
Methotrexate	IT	15 mg/patient	Days 1, 8, 15, 22
Prednisolone	IT	20 mg/patient	Days 1, 8, 15, 22
Intensification			
Doxorubicin	IV	30 mg/m ²	Days 1, 8, 15
Vincristine	IV	1.4 mg/m ² up to 2.0 mg total	Days 1, 8, 15
Prednisolone	PO	40 mg/m ²	Days 1–14, then taper (days 1–7, no taper) ^a
Cyclophosphamide	IV	1000 mg/m ² (700 mg/m ²)	Day 29
Cytosine arabinoside	CI	75 mg/m ²	Days 29–35
6-Mercaptopurine	PO	60 mg/m ²	Days 29–35 (no allopurinol)
Maintenance^c			
Vindesine	IV	2.4 mg/m ² (2.0 mg/m ²)	Days 1, 22
Cyclophosphamide	IV	600 mg/m ² (400 mg/m ²)	Day 1
Prednisolone	PO	40 mg/m ²	Days 1–3, 22–24, no taper
Doxorubicin	IV	40 mg/m ² (30 mg/m ²)	Day 1
Methotrexate	PO	20 mg/m ²	Days 22, 29, 36, 43, 50, 57

Continued

Table 1. Continued

Agent	Route	Dose (≥ 60 years old)	Schedule (≥ 60 years old)
6-Mercaptopurine	PO	60 mg/m ²	Days 22–56 (no allopurinol)

The doses and schedules in parentheses indicate those for patients aged 60 years or older.

IV, intravenously; PO, per os (orally); IT, intrathecally; CI, continuous infusion; IM, intramuscularly; SQ, subcutaneously.

^aAlso given to post-menopausal women aged <60 years old.

^bL-Asparaginase was withdrawn from consolidation B regimen in September 1996 because of severe adverse events including pancreatitis.

^cThis maintenance regimen was repeated at least five cycles every 3 months until 2 years after induction chemotherapy.

STEM CELL TRANSPLANTATION

For patients aged 15–49 years with an HLA-matched related donor who achieved CR and met the eligibility criteria for SCT, allogeneic SCT was recommended after consolidation A. For LBL patients aged 15–49 years without HLA-matched sibling donors, non-purged autologous SCT (PB or BM) was recommended. For conditioning before SCT, one of the following regimens was taken: busulfan (1 mg/kg \times 4/day orally for 4 days on days –7 to –4) and CPA (60 mg/kg/day on days –3 and –2) (24), or CPA (60 mg/kg/day on consecutive 2 days until day –2) and total body irradiation (TBI; 12 Gy by 4–6 fractions on consecutive 2–3 days until day –1) were given. For prophylaxis of acute graft-versus-host disease, cyclosporine A and short-term MTX were given (25); however, the doses of MTX were reduced to 10 mg/m² on day 1 and 7 mg/m² on days 3 and 5 (26).

MODIFICATION OF DOSES AND SCHEDULES

For patients aged 60 years or older, the dose reductions and schedule modifications were performed as shown in Table 1. Treatment was postponed if the aspartate aminotransferase (AST) [glutamic oxaloacetic transaminase (GOT)] or alanine aminotransferase (ALT) [glutamic pyruvic transaminase (GPT)] level was over 200 U or if the total bilirubin was over 2.0 mg/dl. DXR or daunorubicin was omitted when cardiotoxicities such as tachycardia, arrhythmia, ejection fraction $<40\%$ and cardiomyopathy developed. In patients seropositive for hepatitis B virus surface antigen or hepatitis C virus antibody, PSL was omitted. In patients developing an infection of Grade 3 or greater, treatment was postponed, and the doses of 6-MP and MTX in the interim maintenance therapy were reduced to 50% for $<750/\text{mm}^3$ of neutrophils or $<50\,000/\text{mm}^3$ of platelets, or skipped for $<500/\text{mm}^3$ of neutrophils, $<30\,000/\text{mm}^3$ of platelets or over three times higher than the upper normal limit of AST or ALT. After recovery of those toxicities, 6-MP and MTX were given again at the 50% dose and increased to the full dose if possible. In patients who developed constipation of Grade 3 or greater caused by vinca alkaloids, their administration was skipped. In patients who developed Grade 4 paralytic ileus, all treatments were discontinued. In patients suffering from VCR-induced neurotoxicity, VCR could be changed to VDS.

In those who developed CPA-induced hemorrhagic cystitis, CPA was omitted in subsequent cycles. Hyperglycemia was treated with insulin, but in uncontrollable patients, PSL was reduced in doses or discontinued. In patients with psychosis, osteoporosis or hypertension exaggerated by PSL, PSL was reduced in doses or discontinued. In patients with paralysis, convulsion or psychosis developing 72 h after IT-MTX, it was withdrawn. In patients developing arachnoiditis associated with IT-MTX, it was switched to IT-Ara-C (Ara-C 40 mg + PSL 20 mg).

SUPPORTIVE THERAPY

Trimethoprim/sulfamethoxazole was recommended for the prophylaxis of *Pneumocystis jirovecii* pneumonia. G-CSF was prophylactically used when the neutrophil count decreased to $<1000/\text{mm}^3$. Appropriate antibiotics were given for neutropenic infection. When they were not effective, the addition of anti-fungal agents was recommended. The prophylactic use of amphotericin-B syrup for fungal infection of the oral cavity and esophagus was recommended. In patients with suspected varicella zoster virus infection or herpes simplex virus infection, acyclovir was given immediately. Adequate drugs for peptic ulcer were given during PSL administration. Total parenteral nutrition was recommended when patients' intake decreased by chemotherapy. Attention was paid to patients with cough, dyspnea and fever to make early diagnosis of interstitial pneumonia and to start appropriate treatment.

CRITERIA FOR RESPONSE

Patients with ALL were judged to have achieved CR when the results of a BM examination became normal (blasts $\leq 5\%$) along with the normalization of all abnormal laboratory findings such as absolute neutrophil count $>1000/\text{mm}^3$ and platelets $>100\,000/\text{mm}^3$, and all extramedullary diseases had resolved. Patients with LBL were judged to have achieved CR when the tumor or swollen lymph nodes disappeared for more than 4 weeks, according to the WHO response criteria (27).

TOXICITY

Toxicity was assessed according to the JCOG toxicity criteria, an expanded version of the Common Toxicity Criteria version 1.0 by the National Cancer Institute (28).

ENDPOINT

The primary endpoint was 5-year progression-free survival (PFS). Secondary endpoints included CR rate (%CR), OS and adverse events.

STATISTICAL ANALYSIS

A required sample size to evaluate chemotherapy was 97 patients with one-sided α of 0.05, β of 0.20, an expected %5-year PFS of 30% and a threshold %5-year PFS rate of 14.3% (29). Considering ineligible patients, a planned sample size was determined as 120. %CR was evaluated in 108 eligible patients. OS and PFS were evaluated in 108 eligible patients, 87 patients who achieved CR, Philadelphia (Ph) chromosome-negative ALL patients (72 patients) and adolescent and young adult (AYA) patients aged 25 years old or younger (41 patients). Survival curves were estimated with the Kaplan–Meier method. The difference between survival curves according to each characteristic was analyzed using the log-rank test. The Cox proportional hazards regression analysis was performed to assess the association between each patient characteristic and OS. All analyses were carried out with SAS release 9.1 (SAS Institute Inc., Cary, NC, USA).

RESULTS

PATIENT CHARACTERISTICS

Between November 1994 and December 1999, 115 newly diagnosed patients with ALL or LBL aged 15–69 years were registered from 27 institutions, and 108 (94%) were eligible. Case report forms of one patient were lost. Two enrolled patients under the institutional diagnosis of LBL were judged ineligible by the central pathology review, because each diagnosis was changed to Burkitt's lymphoma. The remaining five patients, including one patient who was ineligible by the central pathology review, were ineligible due to the following reasons; a diagnosis of diffuse large B-cell lymphoma, small cell lung cancer and granulocytic sarcoma, prior therapy and coexistence of miliary tuberculosis at enrollment. In five other eligible patients, the clinical diagnosis was changed to ALL because the neoplastic cells in BM were 25% or more (2). The 108 eligible patients ranged in age from 15 to 69 years, with a median of 33.5 years (38.5 years in ALL and 20.5 years in LBL, respectively). Immunophenotypic analysis was adequately performed in 100 (93%) of 108 eligible cases (Table 2). Among those cases, 69 (64%) were categorized as B-lineage, whereas 23

Table 2. Characteristics of 108 eligible patients

	ALL (96)	LBL (12)	Total (n, %)
Immunophenotype			
T-lineage	13 (14)	10 (83)	23 (21)
B-lineage	69 (72)	0 (0)	69 (64)
Others	8 (8)	0 (0)	8 (7)
NA	6 (6)	2 (17)	8 (7)
FAB subtype			
L1	23 (24)	NA	23 (21)
L2	69 (72)	NA	69 (64)
L3	4 (4)	NA	4 (4)
Chromosome			
Philadelphia chromosome	24 (25)	0	24 (22)
t(4;11)	2 (2)	0	2 (2)
Other abnormalities	22 (23)	0	22 (20)
Normal	29 (30)	3 (25)	32 (30)
Not evaluable	12 (13)	4 (33)	16 (15)
Not examined	7 (7)	5 (42)	12 (11)
PS			
0	32 (33)	3 (25)	35 (32)
1	46 (48)	3 (25)	49 (45)
2	15 (16)	5 (42)	20 (19)
3	3 (3)	1 (8)	4 (4)
B-symptom			
Absence	58 (60)	10 (83)	68 (63)
Presence	36 (38)	2 (17)	38 (35)
Not available	2 (2)	0	2 (2)

ALL, acute lymphoblastic leukemia; LBL, lymphoblastic lymphoma; NA, not applicable; FAB, French–American–British Classification; PS, performance status.

(21%) as T-lineage. Cytogenetic analysis was performed in 96 (89%) of 108 cases. Ph chromosome and t(4;11) were detected in 24 (22%) and 2 (2%) cases, respectively. Other chromosomal abnormalities were reported in 22 cases (20%), normal karyotypes in 32 (30%) and indeterminate results in 16 (15%).

THERAPEUTIC RESULTS

Eighty-seven [81%; 95% confidence interval (CI), 72–88%] of the 108 eligible patients achieved CR. Among those who achieved CR, 31 patients proceeded to SCT. At the time of the last follow-up, 29 patients remained alive without progression. The OS and PFS curves of the 108 eligible patients are shown in Fig. 2. The median PFS and OS of all 108 eligible patients were 14 months (95% CI, 10–19 months) and 21 months (95% CI, 18–31 months), respectively. As shown in Fig. 3, the estimated probability of their 5-year OS and




Figure 2. Progression-free survival (PFS) in blue and overall survival (OS) in red of 108 eligible patients with ALL and LBL. The median PFS and OS of the 108 eligible patients were 14 months [95% confidence interval (CI), 10–19 months] and 21 months (95% CI, 18–31 months). Estimated probability of PFS and OS at 5 years were 28% (95% CI, 19–36%) and 29% (95% CI, 21–38%).




Figure 3. PFS in blue and OS in red of the 87 patients with ALL and LBL who achieved CR. Estimated probability of PFS and OS at 5 years were 32% (95% CI, 22–42%) and 34% (95% CI, 24–44%).

PFS of 87 patients who achieved CR was 34% (95% CI, 24–44%) and 32% (95% CI, 22–42%), respectively.

Eleven patients including one ineligible patient who achieved CR were registered and underwent SCT according to the JCOG9402 protocol, and high-dose busulfan and CPA, or TBI and high-dose CPA as the conditioning regimen; however, several advances in this area during the study period have enabled treating physicians to choose a TBI-based conditioning regimen and an unrelated donor. As a result, 21 patients who achieved CR underwent SCT outside the protocol. To exploratory assess the role of SCT, 31 patients who underwent SCT in the first CR were evaluated separately from the protocol results, as shown in Table 3. Their median age was 24 years. Twenty-four patients (77%) with ALL and seven (23%) with LBL were included. Autologous and allogeneic stem cells were used in 12 (PB in 12 patients) and 19 patients (related donor BM in 11, unrelated donor BM in 6 and related donor PB in 2), respectively. The OS of the 31 patients is shown in Fig. 4, and

Table 3. Characteristics of 31 patients who received stem cell transplantation during first complete remission and main therapeutic results

Patient characteristics	n
Registered	10
Not registered	21
Median age (years, range)	24 (16–67)
Gender (male/female)	17/14
Disease	
ALL	24
LBL	7
Chromosome abnormality	
Philadelphia chromosome	3
t(4;11)	0
Conditioning regimen	
BU + CPA	6
CPA + TBI	18
Others	4
Unknown	3
Stem cell source	
Allo-PBSCT	2
Autologous (BM/PB)	0/12
Allogeneic (sibling/unrelated)	11/6
Sibling (BM/PB)	11/2
Acute GVHD prophylaxis	
CSP + short-term MTX	15
Others	3
Unknown	1
Results	
Died <100 days after stem cell transplantation	3
5-year overall survival	51% (95% CI, 32–67)
Alive at last follow-up	16

BU, busulfan; CPA, cyclophosphamide; TBI, total body irradiation; PBSCT, peripheral blood stem cell transplantation; BM, bone marrow; PB, peripheral blood; GVHD, graft-versus-host disease; CSP, cyclosporine; MTX, methotrexate; CI, confidence interval.

the estimated probability of OS at 5 years was 51% (95% CI, 32–67%). The estimated survival probabilities of autologous and allogeneic SCT at 5 years were 58% (95% CI, 27–80%) and 47% (95% CI, 24–67%), respectively.

SUBGROUP ANALYSIS

OS was compared between subsets according to the following eight variables: age (≤ 40 versus > 40 years), sex (male versus female), clinical diagnosis (ALL versus LBL), ECOG-PS (0, 1 versus 2–4), leukocytes at initial presentation ($< 10\,000$ versus $\geq 10\,000/\mu\text{l}$), immunophenotype of neoplastic cells (non-T versus T), C-reactive protein




Figure 4. OS of the 31 patients who received stem cell transplantation in first complete remission. Estimated probability of OS at 5 years was 51% (95% CI, 32–67%).




Figure 5. OS of patients with Philadelphia chromosome-negative ALL. Estimated probability at 5 years was 37% (95% CI, 26–48%).

Table 4. Univariate analysis of factors affecting overall survival

Variables	Category	Distribution (n)	P value ^a	HR ^b (95% CI)
Age	40/>40	60/48	0.02	1.702 (1.088–2.663)
Gender	M/F	54/54	0.488	0.853 (0.545–1.335)
Disease	ALL/LBL	96/12	0.434	0.747 (0.359–1.553)
PS	0, 1/2–4	84/24	0.554	0.847 (0.488–1.469)
WBC count (× 10 ⁴ /μl)	<1.0/1.0	49/56	0.092	1.481 (0.938–2.338)
Immunophenotype	Non-T/T	85/23	0.132	0.64 (0.358–1.143)
CRP (mg/dl)	0.3/>0.3	25/77	0.078	1.719 (0.942–3.139)
Ph or t(4;11)	-/+	82/26	0	2.567 (1.568–4.203)

HR, hazard ratio; CRP, C-reactive protein; Ph, Philadelphia chromosome.

^aP value by log-rank test.

^bHazard ratio by the Cox proportional hazards regression analysis.

(CRP: ≤0.3 versus >0.3 mg/dl) and chromosome anomaly of Ph chromosome or t(4;11) (not detected versus detected); small P values in univariate analysis using the log-rank test were obtained in the following subsets: age ≤40 years versus older and chromosome anomaly of Ph chromosome and t(4;11) in not detected versus detected (Table 4). Multivariate analysis showed that only the detection of Ph chromosome or t(4;11) was significantly associated with shortened survival (P = 0.0197), although age >40 years (P = 0.0654) and male sex (P = 0.0724) showed tendencies toward shorter survival. As shown in Figs 5 and 6, the 5-year OS of patients with Ph chromosome-negative ALL and AYA patients aged 25 years or younger was 37% (95% CI; 26–48%) and 41% (95% CI; 26–55%), respectively.

TOXICITY

The number of patients with treatment-related death (TRD) was six; three patients died in the induction phase and the




Figure 6. OS of adolescent and young adult ALL patients aged 25 years or younger. Estimated probability at 5 years was 41% (95% CI, 26–55%).

remaining three in the consolidation A, CNS prophylaxis and SCT phases, respectively. The causes of death were infection in four patients, TRD in one and hemorrhage in one. The numbers of patients who developed non-hematologic toxicities of Grade 3 or greater are shown in Table 5. The major non-hematologic toxicities of Grade 3 or greater in induction therapy were ALT (GPT) elevation (28%), infection (14%), pulmonary event (6%) and peripheral neuropathy (2%). Among the 96 patients treated with consolidation A, 7 (7%) developed ALT (GPT) elevation and 6 (6%) infection of Grade 3 or greater. Pancreatitis (Grade 4), and hyperglycemia (Grade 4) and diarrhea (Grade 3) occurred in one patient each after administration of L-asparaginase in consolidation B, which lead to the exclusion of L-asparaginase from consolidation B. Thereafter, no serious adverse events associated with L-asparaginase were observed.

DISCUSSION

We conducted a multicenter Phase II study of adult ALL and LBL, JCOG9402, to evaluate an intensified post-remission chemotherapy. The primary endpoint was a 5-year PFS of 28%, which almost achieved an estimated one of 30%. It is, however, unsatisfactory in terms of similar results of 5-year OS (29%) and the %CR to our previous Phase II

Table 5. Non-hematologic toxicities of Grade 3 or greater

Regimen	Induction (n, %)	Consolidation A (n, %)	Consolidation B (n, %)
Total number of treated patients	115	98	83
Total number of evaluated courses	113	96	81
Toxicity			
GPT elevation	31 (28%)	7 (7%)	12 (15%)
GOT elevation	12 (11%)	2 (2%)	3 (4%)
Fever (non-infectious)	5 (4%)	3 (3%)	3 (4%)
Infection	16 (14%)	6 (6%)	5 (6%)
Pulmonary	7 (6%)	0 (0%)	0 (0%)
Diarrhea	5 (4%)	0 (0%)	3 (4%)
Constipation	3 (3%)	0 (0%)	1 (1%)
Cardiac dysfunction	3 (3%)	0 (0%)	0 (0%)
Arrhythmia	2 (2%)	0 (0%)	0 (0%)
Peripheral neuropathy	2 (2%)	0 (0%)	0 (0%)

GPT, glutamic pyruvic transaminase; GOT, glutamic oxaloacetic transaminase.

study, JCOG9004 (13), despite of an intensified chemotherapy. In JCOG9402, 81% of all eligible patients achieved CR. When compared with the %CR (83%) in JCOG9004, the %CR was almost the same. These results are equivalent to those from other multicenter trials (6,9–11). In JCOG9402, Ara-C was incorporated into consolidation A to intensify the consolidation phase with the support of G-CSF. Between consolidation and maintenance therapy, both CNS prophylaxis with interim maintenance therapy and intensification therapy were also incorporated. In maintenance therapy, long-term administration of MTX and 6-MP was combined with the intermittent use of VDS, CPA, PSL and DXR. In JCOG9402, five patients (5%) died during the protocol chemotherapy, whereas in the previous JCOG9004, using a similar strategy to JCOG9402 in view of the intensified induction and post-remission chemotherapy with the support of G-CSF, 17 patients (12%) died during the protocol chemotherapy. These results suggest that the intensified induction and post-remission chemotherapy in JCOG9402, especially including the revised consolidation B regimen excluding L-asp, could be performed more safely than those in JCOG9004. Compared with the important role of L-asp in the treatment of pediatric ALL, that of adult ALL might be limited partly because L-asp is thought to be more toxic especially in elderly patients than in pediatric patients (30). Regarding the long-term outcome, however, the median survival time in JCOG9402 (21 months) did not improve compared with JCOG9004 (26 months) (13), and the estimated probability of survival at 5 years (29%) also did not improve. Thus, it is unlikely that the strategies of

JCOG9402, especially further intensification of post-remission chemotherapy, improved the overall therapeutic results, when compared with JCOG9004 (13).

In the univariate analysis of prognostic factors affecting OS, older age and chromosomal abnormality of Ph chromosome or t(4;11) were unfavorable factors. In recent years, regimens for pediatric ALL have been applied for AYA ALL (31). As far as AYA in our study, the 5-year OS (41%) was unsatisfactory compared with the previous AYA trials with pediatric regimens (31). The proportion of patients with Ph chromosome anomaly (24/108, 22%) was not different from other reports, and they showed an unfavorable prognosis, similar to other studies (3,6,10,11,16,17,32). Since it was not planned to analyze the fusion gene of *BCR-ABL* in the present study, there remains the possibility that a fraction of patients who were judged not to have Ph chromosome by conventional cytogenetics may have the fusion gene of *BCR-ABL*, as Jinnai precisely discussed in the JALSG-ALL97 study (17). The 5-year OS of Ph chromosome-negative ALL patients (37%) appears to be unsatisfactory, comparing with recent survival improvement of Ph chromosome-positive ALL (Ph+ ALL) patients with imatinib (32). It is, however, not conclusive because our study period is different from those of other studies, in which more allogeneic SCTs are conducted and imatinib is available (32). CRP did not have an impact in the present study, in contrast to the authors' previous trial, JCOG8702 (12). Female sex was not a significant prognostic factor different from JCOG9004 (13). Similar to JCOG9004, leukocytes at initial presentation were not significant. Other prognostic factors identified in this study showed consistent results with previous reports, although there were some discrepancies. In an exploratory multivariate analysis (data not shown), chromosomal abnormality was the only significant unfavorable prognostic factor. Although age of >40 years and male sex were statistically marginal, both are considered to have a clinically meaningful impact on poor outcome (hazard ratio, 1.604 in >40 years and 1.582 in male). Patients with LBL did not show a more favorable prognosis than those with ALL, as shown in Table 4; however, this is not conclusive because of the small number of LBL patients ($n = 12$) enrolled in the present study.

Because there is considerable heterogeneity in the stem cell source, conditioning regimen etc., in the 31 patients who underwent SCT in first CR, it is impossible to draw meaningful conclusions; however, the results of OS suggest that SCT, particularly allogeneic SCT, in first CR is worthy of further investigation as well as JCOG9004 (13). In the recent meta-analysis, allogeneic SCT proved to be superior to autologous SCT or chemotherapy for patients with ALL in first CR (33).

The therapeutic outcome of JCOG9402 is still unsatisfactory. One of the plausible reasons for the unfavorable therapeutic outcome is age. In the present study, the median age of the 108 eligible patients was 33.5 years, and that of the 96 eligible patients with ALL was 38.5 years. When this was

compared with the median age of 25–40 years in the reported results (3,6,9–12,16,32), the median age in the present study, especially that of ALL, is rather older. Considering that age is a strong prognostic factor in most studies of adult ALL and LBL (34,35), the therapeutic results in this study could be worse. In addition, the median age of eligible patients in the present JCOG9402 study (33.5 years) was lower than in JCOG9004 (41 years). Therefore, it seems that intensification of induction and post-remission chemotherapy by available agents with the support of G-CSF for adult ALL and LBL had reached its limit (36). To further improve the therapeutic results, application of a dose-intensive pediatric regimen (7,37–40), incorporation of effective new agents such as nelarabine for T-ALL and T-LBL (41), imatinib and new tyrosine kinase inhibitors for Ph+ ALL (42–47) and other new targeted agents including monoclonal antibodies (48) may be needed, in addition to the risk-adapted SCT (48), considering the status of minimal residual disease (MRD) (48,49).

In conclusion, JCOG9402 did not show improvement in long-term follow-up results in adult patients with ALL and LBL when compared with the investigators' latest historical control, JCOG9004, although intensified post-remission chemotherapy, especially its revised version excluding L-asparaginase from the consolidation B regimen, was performed safely in most patients with an achievement of the primary endpoint of estimated 5-year PFS. To further improve the therapeutic outcomes of adult patients with ALL and LBL, new strategies including that for non-AYA ALL with Ph-negative chromosome using pediatric regimens, MRD-guided strategy and developing of new agents such as liposomal-doxorubicin (50), peg-asparaginase (51) and clofarabine (52) must be considered.

Acknowledgements

We are greatly appreciative of all members of JCOG-LSG for their excellent cooperation in this long-term follow-up study. We thank the treating physicians, nurses, laboratory technicians and patients. We wish to thank Ms Yuko Watanabe (JCOG Data Center) for data management.

Funding

This study was supported by Grants-in-Aid for Cancer Research from the Ministry of Health, Labor and Welfare of Japan (5S-1, 8S-1, 11S-1, 11S-4, 14S-1, 14S-4, 17S-1, 17S-5, 20S-1, 20S-6, 1-1, 4-5, 7-29, 9-10) and by the National Cancer Center Research and Development Fund (23-A-16, 23-A-17) (1993–present).

Conflict of interest statement

None declared.

References

1. Swerdlow SH, Campo E, Harris NL, et al., editors. *World Health Organization Classification of Tumours of Haematopoietic and Lymphoid Tissues*. Lyon: IARC Press 2008.
2. Slater DE, Mertelsmann R, Koziner B, et al. Lymphoblastic lymphoma in adults. *J Clin Oncol* 1986;4:57–67.
3. Larson RA, Dodge RK, Burns CP, et al. A five-drug remission induction regimen with intensive consolidation for adults with acute lymphoblastic leukemia: Cancer and Leukemia Group B study 8811. *Blood* 1995;85:2025–37.
4. Hoelzer D, Gokbuget N, Digel W, et al. Outcome of adult patients with T-lymphoblastic lymphoma treated according to protocols for acute lymphoblastic leukemia. *Blood* 2002;99:4379–85.
5. Thomas DA, O'Brien S, Cortes J, et al. Outcome with hyper-CVAD regimens in lymphoblastic lymphoma. *Blood* 2004;104:1624–30.
6. Kantarjian H, Thomas D, O'Brien S, et al. Long-term follow-up results of hyperfractionated cyclophosphamide, vincristine, doxorubicin, and dexamethasone (Hyper-CVAD), a dose-intensive regimen, in adult acute lymphocytic leukemia. *Cancer* 2004;101:2788–801.
7. Huguet F, Leguay T, Raffoux E, et al. Pediatric-inspired therapy in adults with Philadelphia chromosome-negative acute lymphoblastic leukemia: the GRAALL-2003 study. *J Clin Oncol* 2009;27:911–8.
8. Rivera GK, Pinkel D, Simone JV, Hancock ML, Crist WM. Treatment of acute lymphoblastic leukemia—30 years' experience at St. Jude Children's Research Hospital. *N Engl J Med* 1993;329:1289–95.
9. Hoelzer D, Thiel E, Loftier H, et al. Prognostic factors in a multicenter study for treatment of acute lymphoblastic leukemia in adults. *Blood* 1988;71:123–31.
10. Larson RA, Dodge RK, Linker CA, et al. A randomized controlled trial of filgrastim during remission induction and consolidation chemotherapy for adults with acute lymphoblastic leukemia: CALGB study 9111. *Blood* 1998;92:1556–64.
11. Thomas X, Boiron JM, Huguet F, et al. Outcome of treatment in adults with acute lymphoblastic leukemia: analysis of the LALA-94 trial. *J Clin Oncol* 2004;22:4075–86.
12. Kobayashi T, Tobinai K, Shimoyama M, et al. Long-term follow-up results of adult patients with acute lymphocytic leukemia or lymphoblastic lymphoma treated with short-term, alternating non-cross-resistant chemotherapy: Japan Clinical Oncology Group study 8702. *Jpn J Clin Oncol* 1999;29:340–8.
13. Tobinai K, Takeyama K, Arima F, et al. Phase II study of chemotherapy and stem cell transplantation for adult acute lymphoblastic leukemia or lymphoblastic lymphoma: Japan Clinical Oncology Group study 9004. *Cancer Sci* 2007;98:1350–7.
14. Tanimoto M, Miyawaki S, Ino T, et al. Response-oriented individualized induction therapy followed by intensive consolidation and maintenance for adult patients with acute lymphoblastic leukemia: the ALL-87 study of the Japan Adult Leukemia Study Group (JALSG). *Int J Hematol* 1998;68:421–9.
15. Ueda T, Miyawaki S, Asou N, et al. Response-oriented individualized induction therapy with six drugs followed by four courses of intensive consolidation, 1 year maintenance and intensification therapy: the ALL90 study of the Japan Adult Leukemia Study Group. *Int J Hematol* 1998;68:279–89.
16. Takeuchi J, Kyo T, Naito K, et al. Induction therapy by frequent administration of doxorubicin with four other drugs, followed by intensive consolidation and maintenance therapy for adult acute lymphoblastic leukemia: the JALSG ALL93 study. *Leukemia* 2002;16:1259–66.
17. Jinnai I, Sakura T, Tsuzuki M, et al. Intensified consolidation therapy with dose-escalated doxorubicin did not improve the prognosis of adults with acute lymphoblastic leukemia: the JALSG-ALL97 study. *Int J Hematol* 2010;92:490–502.
18. Fiere D, Lepage E, Sebban C, et al. Adult acute lymphoblastic leukemia: a multicentric randomized trial testing bone marrow transplantation as postremission therapy: the French Group on therapy for adult acute lymphoblastic leukemia. *J Clin Oncol* 1993;11:1990–2001.
19. Bennett JM, Catovsky D, Daniel MT, et al. Proposals for the classification of acute leukemias. *Br J Haematol* 1976;33:451–8.
20. Bennett JM, Catovsky D, Daniel MT, et al. Proposal for the recognition of minimally differentiated acute myeloid leukaemia (AML-M0). *Br J Haematol* 1991;78:325–9.

21. The Non-Hodgkin's Lymphoma Pathologic Classification Project: National Cancer Institute sponsored study of classifications of non-Hodgkin's lymphoma: summary and description of a Working Formulation for clinical usage. *Cancer* 1982;49:2112–35.
22. Oken M, Creech R, Tormey D, et al. Toxicity and response criteria of the Eastern Cooperative Oncology Group. *Am J Clin Oncol* 1982;5:649–55.
23. Shimoyama M, Fukuda H, Saijo N, Yanaguchi N. Japan Clinical Oncology Group (JCOG). *Jpn J Clin Oncol* 1998;28:158–62.
24. Tutschka PJ, Copelan EA, Klein JP. Bone marrow transplantation for leukemia following a new busulfan and cyclophosphamide regimen. *Blood* 1987;70:1382–8.
25. Storb R, Deeg HJ, Whitehead J, et al. Methotrexate and cyclosporine compared with cyclosporine alone for prophylaxis of acute graft versus host disease after marrow transplantation for leukemia. *N Engl J Med* 1986;314:729–35.
26. Morishima Y, Morishita Y, Tanimoto M, et al. Low incidence of acute graft-versus-host disease by the administration of methotrexate and cyclosporine in Japanese leukemia patients after bone marrow transplantation from human leukocyte antigen compatible siblings; possible role of genetic homogeneity. *Blood* 1989;74:2252–6.
27. *WHO Handbook for Reporting Results of Cancer Treatment*. WHO offset publication no. 48. Geneva: World Health Organization 1979.
28. Tobinai K, Kohno A, Yasuhiro S, et al. Toxicity grading criteria of the Japan Clinical Oncology Group. *Jpn J Clin Oncol* 1993;23:250–7.
29. Makuch RW, Simon RM. Sample size considerations for non-randomized comparative studies. *J Chr Dis* 1980;33:175–81.
30. Douer D. Is asparaginase a critical component in the treatment of acute lymphoblastic leukemia? *Best Pract Res Clin Haematol* 2008;21:647–58.
31. Wendy S. Adolescents and young adults with acute lymphoblastic leukemia. *Hematology Am Soc Hematol Educ Program* 2010;2010:489–93.
32. Kantarjian H, O'Brien S, Smith TL, et al. Results of treatment with Hyper-CVAD, a dose-intensive regimen, in adult acute lymphocytic leukemia. *J Clin Oncol* 2000;18:547–61.
33. Ram R, Gafter-Gvili A, Vidal L, et al. Management of adult patients with acute lymphoblastic leukemia in first complete remission: systematic review and meta-analysis. *Cancer* 2010;116:3447–57.
34. Marks DI. Treating the 'older' adult with acute lymphoblastic leukemia. *Hematology Am Soc Hematol Educ Program* 2010;2010:13–20.
35. Larson RA. Acute lymphoblastic leukemia: older patients and newer drugs. *Hematology Am Soc Hematol Educ Program* 2005;131–6.
36. Bassan R. Randomized clinical trials in adult acute lymphoblastic leukemia: which is the question? *Haematologica* 1998;83:193–6.
37. DeAngelo DJ. The treatment of adolescents and young adults with acute lymphoblastic leukemia. *Hematology Am Soc Hematol Educ Program* 2005;123–30.
38. Stock W, La M, Sanford B, et al. What determines the outcomes for adolescents and young adults with acute lymphoblastic leukemia treated on cooperative group protocols? A comparison of Children's Cancer Group and Cancer and Leukemia Group B studies. *Blood* 2008;112:1646–54.
39. Litzow MR. Evolving paradigms in the therapy of Philadelphia chromosome-negative acute lymphoblastic leukemia in adults. *Hematology Am Soc Hematol Educ Program* 2009;362–70.
40. Nachman JB, La MK, Hunger SP, et al. Young adults with acute lymphoblastic leukemia have an excellent outcome with chemotherapy alone and benefit from intensive postinduction treatment: a report from the Children's Oncology Group. *J Clin Oncol* 2009;27:5189–94.
41. Berg SL, Blaney SM, Devidas M, et al. Phase II study of nelarabine (compound 506U78) in children and young adults with refractory T-cell malignancies: a report from the Children's Oncology Group. *J Clin Oncol* 2005;23:3376–82.
42. Yanada M, Takeuchi J, Sugiura I, et al. High complete remission rate and promising outcome by combination of imatinib and chemotherapy for newly diagnosed BCR-ABL-positive acute lymphoblastic leukemia: a phase II study by the Japan Adult Leukemia Study Group. *J Clin Oncol* 2006;24:460–6.
43. Wassmann B, Pfeifer H, Goekbuget N, et al. Alternating versus concurrent schedules of imatinib and chemotherapy as front-line therapy for Philadelphia positive acute lymphoblastic leukemia (Ph+ ALL). *Blood* 2006;108:1469–77.
44. Bassan R, Rossi G, Pogliani EM, et al. Chemotherapy-phased imatinib pulses improve long-term outcome of adult patients with Philadelphia chromosome-positive acute lymphoblastic leukemia: Northern Italy Leukemia Group protocol 09/00. *J Clin Oncol* 2010;28:3644–52.
45. Talpaz M, Shah NP, Kantarjian H, et al. Dasatinib in imatinib-resistant Philadelphia chromosome-positive leukemias. *N Engl J Med* 2006;354:2531–41.
46. Kantarjian H, Giles F, Wunderle L, et al. Nilotinib in imatinib-resistant CML and Philadelphia chromosome-positive ALL. *N Engl J Med* 2006;354:2542–51.
47. Ottmann OG, Pfeifer H. Management of Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph+ ALL). *Hematology Am Soc Hematol Educ Program* 2009;371–81.
48. Bassan R, Hoelzer D. Modern therapy of acute lymphoblastic leukemia. *J Clin Oncol* 2011;29:532–43.
49. Bassan R, Spinelli O, Oldani E, et al. Improved risk classification for risk-specific therapy based on the molecular study of minimal residual disease (MRD) in adult acute lymphoblastic leukemia (ALL). *Blood* 2009;113:4153–62.
50. Hunault BM, Leguay T, Thomas X, et al. A randomized study of pegylated liposomal doxorubicin versus continuous-infusion doxorubicin in elderly patients with acute lymphoblastic leukemia: the GRAALL-SAI study. *Haematologica* 2011;96:245–52.
51. Wetzler M, Sanford BL, Kurtzberg J, et al. Effective asparagine depletion with pegylated asparaginase results in improved outcomes in adult acute lymphoblastic leukemia: Cancer and Leukemia Group B Study 9511. *Blood* 2007;109:4164–7.
52. Hijjiya N, Thomson B, Isakoff MS, et al. Phase 2 trial of clofarabine in combination with etoposide and cyclophosphamide in pediatric patients with refractory or relapsed acute lymphoblastic leukemia. *Blood* 2011;118:6043–9.

Appendix

List of participants

K. Aikawa and M.Ku. (Hokkaido Cancer Center); M.Ka. and Y. Kiyama (Sapporo Hokuyu Hospital); I. Miura and A. Miura (Akita University); T. Sai (Iwaki Kyoritsu General Hospital); S. Matsuda (Ohta Nishinouchi Hospital); Y. Sasaki and K. Itoh (National Cancer Center Hospital East); K.To., K. Minato, T. Takenaka, K.Ta. and A. Kohno (National Cancer Center Hospital); T. Miwa and A. Togawa (International Medical Center of Japan); H. Yamada and S. Iwase (Jikei University of Medicine, Aoto Hospital); T. Hotta and K. Ohmachi (Tokai University); H. Nagoshi (St Marianna University Yokohama City West Hospital); T.C. (Niigata Cancer Center Hospital); S. Konda and Y. Masaki (Kanazawa Medical University); R. Ueda and S. Iida (Nagoya City University); M.S. (Nagoya Medical Center); S. Shirakawa and M. Yamaguchi (Mie University); T. Suzuki (Shiga Medical Center for Adults); T. Ohno (Ohtsu Red Cross Hospital); H. Ohno (Kyoto University); S. Fukuhara (Kansai Medical University); Y. Ohno (Tenri Hospital); S. Okamoto (National Kyusyu Medical Center); K. Fujita and Y. Izumi (Kokura Memorial Hospital); M. Sano, Y. Shimamoto and H. Fukushima (Saga University); M.H. and F. Kawano (Kumamoto Medical Center); S. Hanada and K. Uozumi (Kagoshima University); M. Tara (Kagoshima City Hospital); and K. Araki and I. Ohshiro (University of the Ryukyus).