

REFERENCES

- Lynch TJ, Bell DW, Sordella R, et al. Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. *N Engl J Med* 2004;350:2129–2139.
- Paez JG, Jänne PA, Lee JC, et al. EGFR mutations in lung cancer: correlation with clinical response to gefitinib therapy. *Science* 2004;304:1497–1500.
- Pao W, Miller V, Zakowski M, et al. EGF receptor gene mutations are common in lung cancers from “never smokers” and are associated with sensitivity of tumors to gefitinib and erlotinib. *Proc Natl Acad Sci U S A* 2004;101:13306–13311.
- Mitsudomi T, Yatabe Y. Mutations of the epidermal growth factor receptor gene and related genes as determinants of epidermal growth factor receptor tyrosine kinase inhibitors sensitivity in lung cancer. *Cancer Sci* 2007;98:1817–1824.
- Calvo E, Baselga J. Ethnic differences in response to epidermal growth factor receptor tyrosine kinase inhibitors. *J Clin Oncol* 2006;24:2158–2163.
- Mitsudomi T, Morita S, Yatabe Y, et al; West Japan Oncology Group. Gefitinib versus cisplatin plus docetaxel in patients with non-small-cell lung cancer harbouring mutations of the epidermal growth factor receptor (WJTOG3405): an open label, randomised phase 3 trial. *Lancet Oncol* 2010;11:121–128.
- Maemondo M, Inoue A, Kobayashi K, et al. Gefitinib or chemotherapy for non-small-cell lung cancer with mutated EGFR. *N Engl J Med* 2010;362:2380–2388.
- Mok TS, Wu YL, Thongprasert S, et al. Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma. *N Engl J Med* 2009;361:947–957.
- Jackman D, Pao W, Riely GJ, et al. Clinical definition of acquired resistance to epidermal growth factor receptor tyrosine kinase inhibitors in non-small-cell lung cancer. *J Clin Oncol* 2010;28:357–360.
- Herbst RS, Heymach JV, Lippman SM. Lung cancer. *N Engl J Med* 2008;359:1367–1380.
- Kobayashi S, Boggon TJ, Dayaram T, et al. EGFR mutation and resistance of non-small-cell lung cancer to gefitinib. *N Engl J Med* 2005;352:786–792.
- Pao W, Miller VA, Politi KA, et al. Acquired resistance of lung adenocarcinomas to gefitinib or erlotinib is associated with a second mutation in the EGFR kinase domain. *PLoS Med* 2005;2:e73.
- Engelman JA, Zejnullahu K, Mitsudomi T, et al. MET amplification leads to gefitinib resistance in lung cancer by activating ERBB3 signaling. *Science* 2007;316:1039–1043.
- Yano S, Wang W, Li Q, et al. Hepatocyte growth factor induces gefitinib resistance of lung adenocarcinoma cells with EGF receptor mutations. *Cancer Res* 2008;68:9479–9487.
- Matsumoto K, Nakamura T, Sakai K, et al. Hepatocyte growth factor and Met in tumor biology and therapeutic approach with NK4. *Proteomics* 2008;8:3360–3370.
- Turke AB, Zejnullahu K, Wu YL, et al. Preexistence and clonal selection of MET amplification in EGFR mutant NSCLC. *Cancer Cell* 2010;17:77–88.
- Bean J, Brennan C, Shih JY, et al. MET amplification occurs with or without T790M mutations in EGFR mutant lung tumors with acquired resistance to gefitinib or erlotinib. *Proc Natl Acad Sci U S A* 2007;104:20932–20937.
- Onitsuka T, Uramoto H, Nose N, et al. Acquired resistance to gefitinib: the contribution of mechanisms other than the T790M, MET, and HGF status. *Lung Cancer* 2010;68:198–203.
- Kosaka T, Yatabe Y, Endoh H, et al. Analysis of epidermal growth factor receptor gene mutation in patients with non-small cell lung cancer and acquired resistance to gefitinib. *Clin Cancer Res* 2006;12:5764–5769.
- Herbst RS, Sun Y, Eberhardt WE, et al. Vandetanib plus docetaxel versus docetaxel as second-line treatment for patients with advanced non-small-cell lung cancer (ZODIAC): a double-blind, randomised, phase 3 trial. *Lancet Oncol* 2010;11:619–626.
- Yatabe Y, Hida T, Horio Y, et al. A rapid, sensitive assay to detect EGFR mutation in small biopsy specimens from lung cancer. *J Mol Diagn* 2006;8:335–341.
- Cappuzzo F, Hirsch FR, Rossi E, et al. Epidermal growth factor receptor gene and protein and gefitinib sensitivity in non-small-cell lung cancer. *J Natl Cancer Inst* 2005;97:643–655.
- Hirsch FR, Herbst RS, Olsen C, et al. Increased EGFR gene copy number detected by fluorescent in situ hybridization predicts outcome in non-small-cell lung cancer patients treated with cetuximab and chemotherapy. *J Clin Oncol* 2008;26:3351–3357.
- Toyooka S, Date H, Uchida A, et al. The epidermal growth factor receptor D761Y mutation and effect of tyrosine kinase inhibitor. *Clin Cancer Res* 2007;13:3431. author reply: 3431–3432.
- Wang W, Li Q, Yamada T, et al. Crosstalk to stromal fibroblasts induces resistance of lung cancer to EGFR tyrosine kinase inhibitors. *Clin Cancer Res* 2009;15:6630–6638.
- Yamada T, Matsumoto K, Wang W, et al. Hepatocyte growth factor reduces susceptibility to an irreversible epidermal growth factor receptor inhibitor in EGFR-T790M mutant lung cancer. *Clin Cancer Res* 2010;16:174–183.
- Okamoto W, Okamoto I, Tanaka K, et al. TAK-701, a humanized monoclonal antibody to hepatocyte growth factor, reverses gefitinib resistance induced by tumor-derived HGF in non-small cell lung cancer with an EGFR mutation. *Mol Cancer Ther* 2010;9:2785–2792.
- Date K, Matsumoto K, Kuba K, et al. Inhibition of tumor growth and invasion by a four-kringle antagonist (HGF/NK4) for hepatocyte growth factor. *Oncogene* 1998;17:3045–3054.
- Prat M, Crepaldi T, Gandino L, et al. C-terminal truncated forms of Met, the hepatocyte growth factor receptor. *Mol Cell Biol* 1991;11:5954–5962.
- Gandino L, Di Renzo MF, Giordano S, et al. A tyrosine protein kinase activated by bombesin in normal fibroblasts and small cell carcinomas. *Oncogene* 1990;5:721–725.
- Qi J, McTigue MA, Rogers A, et al. Multiple mutations and bypass mechanisms can contribute to development of acquired resistance to MET inhibitors. *Cancer Res* 2011;71:1081–1091.
- Katayama R, Khan TM, Benes C, et al. Therapeutic strategies to overcome crizotinib resistance in non-small cell lung cancers harboring the fusion oncogene EML4-ALK. *Proc Natl Acad Sci U S A* 2011;108:7535–7540.
- Sequist LV, Besse B, Lynch TJ, et al. Neratinib, an irreversible pan-ErbB receptor tyrosine kinase inhibitor: results of a phase II trial in patients with advanced non-small-cell lung cancer. *J Clin Oncol* 2010;28:3076–3083.
- Jänne PA, Schellens JH, Engelman JA, et al. Preliminary activity and safety results from a phase I clinical trial of PF-00299804, an irreversible pan-HER inhibitor, in patients (pts) with NSCLC. *J Clin Oncol* 2008;26:S20(abstr 8027).
- Balak MN, Gong Y, Riely GJ, et al. Novel D761Y and common secondary T790M mutations in epidermal growth factor receptor-mutant lung adenocarcinomas with acquired resistance to kinase inhibitors. *Clin Cancer Res* 2006;12:6494–6501.
- Sharma SV, Lee DY, Li B, et al. A chromatin-mediated reversible drug-tolerant state in cancer cell subpopulations. *Cell* 2010;141:69–80.
- Zhou W, Ercan D, Chen L, et al. Novel mutant-selective EGFR kinase inhibitors against EGFR T790M. *Nature* 2009;462:1070–1074.
- Kasahara K, Arai T, Sakai K, et al. Impact of serum hepatocyte growth factor on treatment response to epidermal growth factor receptor tyrosine kinase inhibitors in patients with non-small cell lung adenocarcinoma. *Clin Cancer Res* 2010;16:4616–4624.
- Tanaka H, Kimura T, Kudoh S, et al. Reaction of plasma hepatocyte growth factor levels in non-small cell lung cancer patients treated with EGFR-TKIs. *Int J Cancer* 2011;129:1410–1416.

Clinical Science

Outcome of surgical resection for recurrent pulmonary metastasis from colorectal carcinoma

Ryu Kanzaki, M.D.^{a,*}, Masahiko Higashiyama, M.D.^a, Kazuyuki Oda, M.D.^b, Ayako Fujiwara, M.D.^a, Toshiteru Tokunaga, M.D.^a, Jun Maeda, M.D.^a, Jiro Okami, M.D.^a, Koji Tanaka, M.D.^c, Tatsushi Shingai, M.D.^c, Shingo Noura, M.D.^c, Masayuki Ohue, M.D.^c, Ken Kodama, M.D.^a

^aDepartment of Thoracic Surgery, Osaka Medical Center for Cancer and Cardiovascular Diseases, 1-3-3 Nakamichi, Higashinari-ku, Osaka 5378511, Japan; ^bDepartment of Surgery, Sakai Municipal Hospital, 1-1-1 Minamiyasui-cho, Sakai-ku, Sakai, Osaka 5900064, Japan; ^cDepartment of Surgery, Osaka Medical Center for Cancer and Cardiovascular Diseases, 1-3-3 Nakamichi, Higashinari-ku, Osaka 5378511, Japan

KEYWORDS:

Pulmonary metastasis;
Colorectal cancer;
Repeat thoracotomy

Abstract

BACKGROUND: The outcomes after repeat pulmonary resection for colorectal cancer (CRC) and the factors associated with the prognosis of these patients remain uncharacterized.

METHODS: Data on 156 patients who underwent curative resection of pulmonary metastasis from CRC were reviewed. Repeat pulmonary resection was performed in 25 patients; the present study examined the outcomes and factors associated with prognosis after repeat pulmonary resection.

RESULTS: The 5-year survival rate after the first pulmonary resection was 56.2%. A multivariate analysis identified a histological type other than well-differentiated adenocarcinoma, a high prethoracotomy serum carcinoembryonic antigen (CEA) level, and the presence of hilar or mediastinal lymph node metastasis as poor prognostic factors for the first pulmonary resection. The 5-year survival rate after repeat pulmonary resection was 42.1%. Hilar or mediastinal lymph node metastasis at the time of the repeat resection was significantly associated with poor survival.

CONCLUSIONS: Repeat pulmonary resection for metastatic CRC provides satisfactory outcomes. Hilar or mediastinal lymph node involvement is consistently associated with a poor prognosis after the first and repeat pulmonary resections.

© 2011 Elsevier Inc. All rights reserved.

Colorectal cancer (CRC) remains one of the leading causes of cancer death in Western countries. More than two thirds of these patients undergo primary curative resection; however, more than half of the resected patients eventually succumb to the disease.¹ The most common sites of recur-

rence after resection of primary CRC are liver and lung. Patients with untreated metastatic CRCs have a median survival time of less than 10 months and a 5-year survival frequency of less than 5%.² Recently, antiangiogenic therapy with bevacizumab combined with oxaliplatin-based chemotherapy was reported to improve the survival time of patients with CRC.³ However, few patients achieved complete remission using these new treatments, and most patients therefore exhibit disease progression.

Therefore, surgery remains the best treatment for patients with pulmonary metastases from CRC if potentially curative

* Corresponding author. Tel.: +81-06-6972-1181; fax: +81-06-6981-8055.

E-mail address: rkanzaki@tj8.so-net.ne.jp

Manuscript received July 21, 2009; revised manuscript August 13, 2010

resection is expected and is an established treatment modality in patients with metastatic CRCs. The reported 5-year survival rates after pulmonary metastasectomy for CRC are 24% to 71.2%.²⁻¹⁵

Repeat pulmonary resection is also effective for recurrent pulmonary metastasis.¹⁶ Several authors^{5,9,11,13-15,17} have advocated this treatment for patients with metastatic CRCs; however, the outcomes after repeat pulmonary resection for CRC and the factors associated with the prognosis of these patients remain uncharacterized. The present study examined the outcome of surgical resection for recurrent pulmonary metastasis from CRCs and determined the prognostic factors compared with those of the first pulmonary resection.

Patients and Methods

Patient selection

From January 1980 to December 2008, a total of 156 patients with previous CRCs underwent curative pulmonary resection at the Osaka Medical Center for Cancer and Cardiovascular Diseases. Curative resection was defined as follows: no additional extrapulmonary sites of metastatic disease or already resected if present, no locoregional recurrence, and no residual macroscopic tumor tissue after the resection. Histopathological evaluations of the resected lung specimens confirmed CRC metastases in all patients.

Patients were selected for resection of pulmonary metastases after meeting the following criteria: (1) pulmonary metastases were deemed to be completely resectable by preoperative radiological examination, (2) absence of apparent hilar or mediastinal lymph node metastases determined by preoperative radiologic examination, (3) metastatic disease limited to the lungs or extrapulmonary distant metastasis(es) that was controlled or controllable if present, (4) locoregional control of the primary CRC was achieved or achievable, and (5) good general condition and adequate respiratory function to tolerate lung resection.

Before 2006, lymph node involvement was generally assessed by computed tomography (CT) scanning, with lymph nodes diagnosed as positive if they extended more than 10 mm across the short-axis diameter. Since 2006, lymph node involvement is generally assessed by F18-fluorodeoxyglucose positron emission tomography/CT (FDG-PET/CT) in our hospital.

Patient characteristics

Clinical information was obtained from the medical records in our hospital. The median time interval between resection of primary CRC and first pulmonary resection was 27 months (range, 0-109 months). The mean age at the time of first pulmonary resection was 62 years (range, 39-83 years of age). Thirty-eight patients had previously under-

gone resection for extrapulmonary metastases or local recurrences before the pulmonary resection. Of these, 29 patients underwent liver metastasectomy, 6 patients underwent resection of a local recurrence of the primary tumor, 2 patients underwent inguinal lymph node metastasectomy, and 1 patient underwent resection of a para-aortic lymph node metastasis. Three patients underwent simultaneous resection of metastatic CRCs to the lung and either to an extrathoracic site or local recurrence: thyroid metastasis in 1 patient, brain metastasis in 1 patient, and local recurrence of primary tumor in 1 patient. Perioperative chemotherapy at thoracotomy, including preoperative and/or postoperative adjuvant therapy, was performed in 76 patients as follows: 5-fluorouracil or its derivatives were administered in 61 patients; tegafur in 23 patients; doxifluridine in 21 patients; fluorouracil in 4 patients; carmofur in 3 patients; UFT in 9 patients; and S-1 in 1 patient. Cisplatin-, irinotecan-, and oxaliplatin-based chemotherapy were administered to 6, 4, and 5 patients, respectively. Table 1 summarizes the patient characteristics.

Repeat pulmonary resection

If new nodules had evolved after the first pulmonary resection, a second resection was defined as a repeat pulmonary resection. Planned staged thoracotomy for bilateral metastases was counted as a single operation and was therefore excluded from this study definition. In the survival analysis of patients who underwent a planned staged thoracotomy, the date when the earlier surgery was performed was recognized as the starting point. Repeat pulmonary metastasectomies were also performed if the patient met the criteria for the first pulmonary resection as described earlier. Repeat pulmonary resections were performed in 25 patients; 24 of these patients underwent second pulmonary resections, and 1 patient underwent a third pulmonary resection.

Follow-up schedule

Follow-up generally involved a chest x-ray or a chest and abdominal CT scan, a physical examination, and blood chemistry performed every 6 to 12 months after the first pulmonary resection. Follow-up information was obtained from the medical records in our hospital, letters from the patient's general practitioner, or from the death certificates of the Osaka Cancer Registry. Patients or their families were contacted by phone or by letter if necessary.

Statistical analysis

The statistical analyses were performed using the StatView 5.0 software program (SAS Institute, Berkeley, CA). The overall survival after the first and repeat pulmonary resections was analyzed by the Kaplan-Meier method using the dates of the first and second pulmonary resections, respectively, as the starting points. Significance of differ-

Table 1 Patient characteristics and details of the first pulmonary resection

Characteristics	n
Sex	
Male	91
Female	65
Age at the first pulmonary resection (y)	
Mean	62
Range	39–83
Stage of primary tumor	
Dukes A	2
Dukes B	37
Dukes C	85
Dukes D	21
Unknown	11
Location of primary tumor	
Colon	74
Rectum	82
Histology of primary tumor	
Well-differentiated adenocarcinoma	65
Moderately differentiated adenocarcinoma	77
Others*	11
Unknown	3
Interval between primary resection and first pulmonary resection (months)	
Median	27
Range	0–109
Prethoracotomy serum CEA level (ng/mL)	
<5	90
≥5	66
History of surgical treatment of extrathoracic recurrence	
Yes	41
No	115
Repeat pulmonary resection	
Yes	25
No	131
Number of resected metastases	
1	100
2	32
≥3	24
Site of metastasis	
Unilateral	130
Bilateral	26
Maximum tumor size (mm)	
≤30	115
>30	41
Type of resection	
Sublobar resection	99
Lobectomy or pneumonectomy	57
Hilar or mediastinal lymph node metastasis	
Yes	15
No	141

*Others include poorly differentiated adenocarcinoma and mucinous adenocarcinoma.

Results

Details of first pulmonary resection

There was no operative major morbidity or mortality. Sublobar resection (wide-wedge resection or segmentectomy) was performed in 99 patients, lobectomy in 56, and pneumonectomy in 1 patient. One hundred patients had a solitary metastasis, and 56 patients had multiple metastases. The details of the first pulmonary resections are shown in Table 1. The median time interval between the first pulmonary resection and death or the latest follow-up examination in the present series was 43 months (range, 4–270 months).

Clinical course after the first pulmonary resection

Ninety-three patients developed recurrence after the first pulmonary resection (Fig. 1). The initial pattern of recurrence after lung resection was pulmonary metastasis in 39 patients, including 7 patients with radiologically apparent mediastinal involvement, surgical margin relapse in 5 patients including 1 patient with radiologically apparent mediastinal involvement, pleuritis carcinomatosa in 2 patients, pulmonary metastasis and extrathoracic recurrence in 7 patients, and extrathoracic recurrence in 40 patients. Twenty-five patients underwent a second pulmonary resection; 13 of these subsequently experienced recurrent disease as pulmonary metastasis in 10 patients and extrathoracic recurrence in 3 patients. Only 1 patient underwent a third pulmonary resection. Currently, 69 patients are alive with no evidence of disease, 14 patients are alive with disease, 8 patients died of another disease, and 65 patients succumbed to the disease.

Overall survival of patients after the first pulmonary resection

The cumulative 3-, 5-, and 10-year survival rates after the first pulmonary resection were found to be 71.4%, 56.2%, and 44.0%, respectively (Fig. 2).




Figure 1 Clinical course after the first pulmonary resection. AWD, alive with disease; DOAD, died of another disease; DOD, died of disease; NED: no evidence of disease.

ences between subgroups was calculated using the log-rank test. The multivariate analysis of prognostic factors was performed using the Cox multivariate proportional hazard model. A *P* value of less than .05 was considered statistically significant. Data are expressed as the mean ± standard deviation or median values.




Figure 2 Overall survival of patients after the first pulmonary resection. The cumulative 3-year, 5-year, and 10-year survival rates after the first pulmonary resection were 71.4%, 56.2%, and 44.0%, respectively.

Analysis of prognostic factors for the first pulmonary resection

The following factors were selected for the univariate analysis of survival: sex, age, pathological stage of the primary CRC according to the Dukes classification, the location of the primary tumor, the histology of the primary tumor (well-differentiated adenocarcinoma/moderately differentiated adenocarcinoma or other/unknown), the interval between primary resection and first pulmonary resection (<24 or \geq 24 months), the prethoracotomy serum carcinoembryonic antigen (CEA) level (<5 or \geq 5 ng/mL), a history of surgical treatment for extrathoracic recurrence, repeat pulmonary resection (yes/no), the number of resected metastases, the site of metastasis (unilateral/bilateral), the maximum tumor size (\leq 3 or >30 mm), the hilar or mediastinal lymph node metastasis (yes/no), and the type of resection (sublobar resection/lobectomy/pneumonectomy). Table 2 shows the results

Table 3 Multivariate analysis of prognosis after the first pulmonary resection

Variable	P value	Risk ratio	95% confidence interval
Histology of primary tumor: moderately differentiated adenocarcinoma or others	.03	1.75	1.06–2.89
Prethoracotomy serum CEA level: \geq 5 ng/mL	.01	1.87	1.16–3.03
Maximum tumor size: >30 mm	.1	1.53	.92–2.52
Hilar or mediastinal lymph node metastasis: yes	.04	2.12	1.05–4.29

of the univariate analysis. Significant relationships ($P < .05$) were found between survival and the following factors: histology of the primary tumor, prethoracotomy serum CEA levels, the maximum tumor size, the type of resection (sublobar resection, lobectomy, or pneumonectomy), and hilar or mediastinal lymph node metastasis. The significant factors identified by the univariate analysis were then subjected to a multivariate analysis (Table 3). A moderately differentiated carcinoma or other histological type, an elevated prethoracotomy serum CEA level (\geq 5 ng/mL), and hilar or mediastinal lymph node metastasis were found to be independent and significant determinants of a poor prognosis.

Details of repeat pulmonary resections

Twenty-one patients who had pulmonary metastasis and 4 patients who had surgical margin relapse after the first pulmonary resection underwent repeat pulmonary resections. There was no operative major morbidity or

Table 2 Univariate analysis of the prognosis after the first pulmonary resection

Factors	n	5-year survival	P value
Sex (male/female)	91/65	49.7/65.2	NS
Age at first pulmonary resection (<63/ \geq 63 y)	76/80	61.2/50.5	NS
Stage of primary tumor (Dukes ABC/D/unknown*)	124/21/11	55.3/49.6	NS
Location of primary tumor (colon/rectum)	74/82	56.6/55.5	NS
Histology of primary tumor (Well/moderately differentiated adenocarcinoma or others/unknown*)	65/88/3	61.9/51.6	.04
Interval between primary resection and first pulmonary resection (<24/ \geq 24 months)	64/92	53.3/58.1	NS
Prethoracotomy serum CEA level (<5/ \geq 5 ng/mL)	90/66	65.2/43.1	.006
History of surgical treatment of extrathoracic recurrence (yes/no)	41/115	46.3/59.2	NS
Repeat pulmonary resection (yes/no)	25/131	64.0/54.5	NS
Number of resected metastases (solitary/multiple)	100/56	60.0/49.9	NS
Site of metastasis (unilateral/bilateral)	130/26	61.3/33.5	NS
Maximum tumor size (\leq 30/>30 mm)	115/41	59.9/46.1	.01
Type of resection (sublobar resection/lobectomy or pneumonectomy)	99/57	57.7/53.7	NS
Hilar or mediastinal lymph node metastasis (yes/no)	15/141	28.1/59.4	.006

NS = not significant.

*Excluding unknown cases.

Table 4 Patient characteristics and details of repeat pulmonary resections

Characteristics	n
Sex (male/female)	
Male	17
Female	8
Age at second pulmonary resection (y)	
Mean	61
Range	47–79
Stage of primary tumor	
Dukes B	6
Dukes C	11
Dukes D	5
Unknown	3
Location of primary tumor	
Colon	8
Rectum	17
Histology of primary tumor	
Well-differentiated adenocarcinoma	11
Moderately differentiated adenocarcinoma	11
Others	2
Unknown	1
Interval between first and second pulmonary resection (months)	
Median	20
Range	5–57
Serum CEA level (ng/mL) before second pulmonary resection	
<5	18
≥5	7
Number of resected metastases at second pulmonary resection	
1	16
2	4
≥3	5
Site of metastasis	
Unilateral	23
Bilateral	2
Site of recurrence	
Pulmonary metastasis	21
Surgical margin relapse	4
Maximum tumor size (mm) at second pulmonary resection	
≤30	20
>30	5
Type of resection at second pulmonary resection	
Sublobar resection	16
Lobectomy or pneumonectomy	9
Hilar or mediastinal lymph node metastasis at second pulmonary resection	
Yes	23
No	2

mortality in the present patients. The surgical mode of second pulmonary resection was a wedge resection in 8 patients, segmentectomy in 8 patients, completion lobectomy in 4 patients, lobectomy in 3 patients, and completion pneumonectomy in 2 patients. All repeat pulmonary metastasectomies were curative resections. The median time interval between the first and repeat pulmonary resection was 20 months (range, 5–57 months). The mean patient age at the time of second pulmonary resection was

61 years (range, 47–79 years of age). Table 4 summarizes the patient characteristics and details of repeat pulmonary resections. The median time interval between second pulmonary resection and either death or the latest follow-up in the present series was 20 months (range, 1–238 months).

Overall survival of the patients after second pulmonary resection

The cumulative 3- and 5-year survival rates after the second pulmonary resection were 54.1% and 42.1%, respectively (Fig. 3).

Analysis of the prognostic factors for repeat pulmonary resection

The following factors were selected for a univariate analysis of survival for repeat pulmonary resection: sex, age at second pulmonary resection, pathological stage of the primary CRC according to the Dukes classification, location of primary tumor, histology of primary tumor, interval between first and second pulmonary resection (<20 or ≥20 months), serum CEA level before second pulmonary resection (<5 or ≥5 ng/mL), the number and site of resected metastases at the second pulmonary resection, site of recurrence (pulmonary metastasis/surgical margin relapse), maximum tumor size at second pulmonary resection (≤30 or >30 mm), type of resection at second pulmonary resection (sublobar resection, lobectomy, or pneumonectomy), and hilar or mediastinal lymph node metastasis at the time of the second pulmonary resection (Table 5). Significant relationships were identified only between hilar or mediastinal lymph node metastasis at the second pulmonary resection and survival after the repeat pulmonary resection.

Comments

In our hospital, pulmonary metastasectomies have been performed for various diseases including CRC, soft-tissue




Figure 3 Overall survival rates of patients after the second pulmonary resection. The cumulative 3- and 5-year survival rates after the second pulmonary resection were 54.1% and 42.1%, respectively.

Table 5 Univariate analysis of the prognosis for repeat pulmonary resections

Factors	N	5-year survival	P value
Sex (male/female)	17/8	46.2/33.3	NS
Age at second pulmonary resection (<61 y/ \geq 61 y)	13/12	45.7/40.0	NS
Stage of primary tumor (dukes BC/D/unknown*)	17/5/3	41.7/25.0	NS
Location of primary tumor (Colon/rectum)	8/17	42.9/41.0	NS
Histology of primary tumor (well/moderately differentiated adenocarcinoma or others/unknown*)	11/13/1	44.4/41.6	NS
Interval between first and second pulmonary resection (<20/ \geq 20 months)	12/13	55.6/25.0	NS
Serum CEA level before second pulmonary resection (<5/ \geq 5 ng/mL)	18/7	36.1/53.6	NS
Number of resected metastases at second pulmonary resection (solitary/multiple)	16/9	47.6/33.3	NS
Site of metastasis at second pulmonary resection (unilateral/bilateral)	23/2	44.3/0	NS
Site of recurrence (pulmonary metastasis/surgical margin relapse)	21/4	54.9/0	NS
Maximum tumor size at second pulmonary resection (\leq 30/ $>$ 30 mm)	20/5	43.0/40.0	NS
Type of resection at second pulmonary resection (sublobar resection/lobectomy or pneumonectomy)	16/9	55.9/25.0	NS
Hilar or mediastinal lymph node metastasis at second pulmonary resection (yes/no)	2/23	0/46.8	.048

NS = not significant.

*Excluding unknown cases.

sarcoma, transitional cell carcinoma, and hepatocellular carcinoma according to the general eligibility criteria described in the Patients and Methods section and with good surgical outcomes.^{18–23} The present study analyzed the outcomes in patients undergoing pulmonary metastasectomy from CRCs. The overall 5-year survival rate was 56.2%, which was consistent with the 24% to 71.2% range previously reported.^{4–15}

Twenty-five patients who met the criteria described for this study underwent a second pulmonary metastasectomy and had a 5-year survival rate after the second resection of 42.1%. Table 6 compares the outcomes in studies involving more than 10 patients undergoing repeat pulmonary resection of metastatic CRCs.^{5,9,11,13–15,17} The overall 5-year survival rates after the second pulmonary resection in these previous reports ranged from 23% to 52.1%. Considering the results of the current study, good surgical outcomes were achieved by repeat pulmonary resections of metastatic CRCs.

Many studies^{5,8,9,14,24} showed no association between repeat pulmonary resection and poor survival by a multivariate analysis for survival after the first pulmonary resection. In addition, previous studies^{5,13} of repeat pulmonary resection patients identified no increase in the risk of morbidity or mortality with a repeat pulmonary resection compared with the initial resection. In the present study, there was no major operative morbidity or mortality for patients undergoing repeat pulmonary resection even in 2 patients who had undergone completion pneumonectomies. These findings indicated that repeat pulmonary resection for metastatic CRC patients is a safe procedure that provides satisfactory patient outcomes.

In the present study, the poor prognostic factors for the first pulmonary resection of metastatic CRC included a moderately differentiated carcinoma or other histological type, an elevated prethoracotomy serum CEA level (\geq 5 ng/mL), and hilar or mediastinal lymph node metastasis

according to a multivariate analysis. The prethoracotomy serum CEA levels were the most consistently reported prognostic factor for pulmonary metastasectomy from CRC.^{4–6,9,12,19,24–28} We previously reported that the prethoracotomy CEA level was the most useful prognostic factor, and an elevated serum CEA level is associated with extrathoracic metastasis after pulmonary metastasectomy from CRCs. A number of reports^{4,5,25,29} associated the presence of hilar or mediastinal lymph node metastases with a poor patient prognosis. Several authors^{7,25} have shown results consistent with the present study, showing that a histological diagnosis for the primary CRC of well-differentiated adenocarcinoma is an independent significant prognostic factor after the first pulmonary resection of CRC.

Although prognostic factors for the first pulmonary resection of metastatic CRC have been well studied, the predictive factors for repeat pulmonary resection have not been sufficiently investigated. Ogata et al¹³ reported that patients with extrathoracic recurrence before a second pulmonary metastasectomy or with mediastinal lymph node metastasis had a poorer prognosis. Welter et al¹⁴ identified an increasing number of metastases as a poor prognostic factor for repeat pulmonary resection for metastatic CRC patients by a multivariate analysis, whereas Park et al¹⁵ showed an association between elevated serum preoperative CEA levels and a poor prognosis by a univariate analysis.

Interestingly, the present study showed that the prognostic factors for the first and repeat pulmonary resection are different. Hilar or mediastinal lymph node involvement is consistently associated with a poor prognosis for both the first and repeat pulmonary resections, whereas the histological type of the primary CRC and the prethoracotomy serum CEA level does not significantly affect patient survival after second pulmonary resection. The differences in the prognostic factors for the first and repeat pulmonary resections depend on the differences between the characteristics of these 2 cohorts. The characteristics between these 2 cohorts

Table 6 Comparison of the surgical outcome of studies of patients undergoing repeat pulmonary resections for metastatic CRC

Author	Year	Total number of patients	Number of patients who underwent repeat pulmonary resection	#1	#2	Prognostic factor for second pulmonary resection	#3
Saito	2002	165	23	54.6	52.1	NA	NA
Pfannschmidt	2003	167	24	24.5*	NA	NA	NA
Vogelsang	2004	75	18	NA	NA	NA	NA
Ogata	2005	76	14	NA	23	Extrathoracic recurrence before second pulmonary metastasectomy, mediastinal lymph node metastasis is associated with poor prognosis	7/11
Welter	2007	169	33	53.8	37.1	Number of metastases by a multivariate analysis	NA
Lee	2007	59	13	NA	NA	NA	NA
Kim	2008	69	28	29	NA	NA	NA
Park	2009	202	48	79.3	NA	Elevated preoperative serum CEA level by univariate analysis	28/NA
Our series	2010	156	25	64	42.1	Lymph node involvement by a univariate analysis	10/13

NA = not available.

*Curative resection only.

#1: Five-year survival rate of patients who underwent repeat pulmonary resections after the first pulmonary resection.

#2: Five-year survival rate of patients who underwent repeat pulmonary resections after the second pulmonary resection.

#3: Total number of patients who experienced recurrence after second pulmonary resections.

reflect different patterns of recurrence after pulmonary resection. The proportion of pulmonary metastasis of patients who experienced recurrence after repeat resection was higher than after the first resection in the present study. Pulmonary metastasis after the second pulmonary resection was found in 10 patients (77%) out of 13 patients who experienced recurrence and in 39 of 93 patients (42%) who experienced recurrence after the first pulmonary resection in the present study. This higher proportion of pulmonary metastasis of patients who experienced recurrence after repeat resection is also observed in Ogata's series¹³ in which the proportions was 64% (7/11) and 33% (25/75). It was speculated that patients who underwent repeat pulmonary resection are highly selected patients whose disease tended to metastasize to the lung without extrathoracic lesions after the first pulmonary resection.

Based on the data of the present study, it is speculated that repeat pulmonary resection for metastatic CRC patients with lymph node involvement should be avoided. Therefore, the preoperative assessment of lymph node involvement is important for repeat pulmonary resection. In our hospital, patients with apparent mediastinal lymph node metastases as determined by preoperative radiologic examinations were excluded as candidates for pulmonary metastasectomy. Before 2006, lymph node involvement was generally assessed by a CT scan. In 2006, FDG-PET/CT was introduced in our hospital for hilar and mediastinal lymph node staging in patients with metastatic lung tumors. One hundred thirty patients of 156 patients underwent a first pulmonary resection during this period in the present study. Fourteen of 130 patients (11%) in the CT era had lymph node involvement, whereas only 1 of 26 patients (4%) in the FDG-PET/CT era had lymph node involvement. Although

the superior diagnostic performance of FDG-PET/CT compared with CT could not be shown because of the retrospective nature of the present study, we recommend the use of FDG-PET/CT for the preoperative assessment of patients with metastatic CRC.

The present study had several limitations. The analyses were performed on a large number of patients treated over several decades with changing chemotherapeutic regimens, and it is possible that reported outcomes were partly influenced by the medical treatments. Patients with recurrence after pulmonary resection who received more recently developed treatments, such as antiangiogenic therapy with bevacizumab combined with oxaliplatin-based chemotherapy,³ may survive for a longer period than those who received an older regimen of chemotherapy regardless of the other factors that were analyzed. However, the significance and influence of chemotherapy on patient outcomes remains difficult to establish in the present study because of its retrospective nature.

Conclusions

Repeat pulmonary resection for metastatic CRC is a safe procedure that provides satisfactory outcomes. The prognostic factors for the first and repeat pulmonary resection are different. Prethoracotomy serum CEA levels and the histological type of primary CRC affect survival after the first pulmonary resection but do not affect survival after repeat pulmonary resection. Hilar or mediastinal lymph node involvement is consistently associated with a poor prognosis after the first and repeat pulmonary resections.

The preoperative assessment of lymph node involvement is important for both the first and repeat pulmonary resection. Aggressive repeat resection is justified for carefully selected patients.

References

- Rosen M, Chan L, Beart RW Jr, et al. Follow-up of colorectal cancer: a meta-analysis. *Dis Colon Rectum* 1998;41:1116–26.
- Simmonds PC. Palliative chemotherapy for advanced colorectal cancer: systematic review and meta-analysis. *Colorectal Cancer Collaborative Group. BMJ* 2000;321:531–5.
- Giantonio BJ, Catalano PJ, Meropol NJ, et al. Bevacizumab in combination with oxaliplatin, fluorouracil, and leucovorin (FOLFOX4) for previously treated metastatic colorectal cancer: results from the Eastern Cooperative Oncology Group Study. *J Clin Oncol* 2007;25:1539–44.
- Pfannschmidt J, Muley T, Hoffmann H, et al. Prognostic factors and survival after complete resection of pulmonary metastases from colorectal carcinoma: experiences in 167 patients. *J Thorac Cardiovasc Surg* 2003;126:732–9.
- Saito Y, Omiya H, Kohno K, et al. Pulmonary metastasectomy for 165 patients with colorectal carcinoma: a prognostic assessment. *J Thorac Cardiovasc Surg* 2002;124:1007–13.
- Rena O, Casadio C, Viano F, et al. Pulmonary resection for metastases from colorectal cancer: factors influencing prognosis. Twenty-year experience. *Eur J Cardiothorac Surg* 2002;21:906–12.
- Ike H, Shimada H, Ohki S, et al. Results of aggressive resection of lung metastases from colorectal carcinoma detected by intensive follow-up. *Dis Colon Rectum* 2002;45:468–73.
- Moore KH, McCaughan BC. Surgical resection for pulmonary metastases from colorectal cancer. *ANZ J Surg* 2001;71:143–6.
- Lee WS, Yun SH, Chun HK, et al. Pulmonary resection for metastases from colorectal cancer: prognostic factors and survival. *Int J Colorectal Dis* 2007;22:699–704.
- McAfee MK, Allen MS, Trastek VF, et al. Colorectal lung metastases: results of surgical excision. *Ann Thorac Surg* 1992;53:780–5.
- Kelvin AW, Faber LP, Warren WH, et al. Repeat pulmonary resection for metachronous colorectal carcinoma is beneficial. *Surgery* 2008;144:712–7.
- Watanabe K, Nagai K, Kobayashi A, et al. Factors influencing survival after complete resection of pulmonary metastases from colorectal cancer. *Br J Surg* 2009;96:1058–65.
- Ogata Y, Matono K, Hayashi A, et al. Repeat pulmonary resection for isolated recurrent lung metastases yields results comparable to those after first pulmonary resection in colorectal cancer. *World J Surg* 2005;29:363–8.
- Welter S, Jacobs J, Krbek T, et al. Long-term survival after repeated resection of pulmonary metastases from colorectal cancer. *Ann Thorac Surg* 2007;84:203–10.
- Park JS, Kim HK, Choi YS, et al. Outcomes after repeated resection for recurrent pulmonary metastases from colorectal cancer. *Ann Oncol* 2009;21:1285–9.
- Groeger AM, Kandioler D, Mueller MR, et al. Survival after surgical treatment of recurrent pulmonary metastases. *Eur J Cardiothorac Surg* 1997;12:703–5.
- Vogelsang H, Haas S, Hierholzer C, et al. Factors influencing survival after resection of pulmonary metastases from colorectal cancer. *Br J Surg* 2004;91:1066–71.
- Higashiyama M, Kodama K, Takami K, et al. Intraoperative lavage cytologic analysis of surgical margins as a predictor of local recurrence in pulmonary metastasectomy. *Arch Surg* 2002;137:469–74.
- Higashiyama M, Kodama K, Higaki N, et al. Surgery for pulmonary metastases from colorectal cancer: the importance of prethoracotomy serum carcinoembryonic antigen as an indicator of prognosis. *Jpn J Thorac Cardiovasc Surg* 2003;51:289–96.
- Kodama K, Doi O, Higashiyama M, et al. Surgical management of lung metastases. Usefulness of resection with the neodymium: yttrium-aluminum-garnet laser with median sternotomy. *J Thorac Cardiovasc Surg* 1991;101:901–8.
- Kodama K, Doi O, Higashiyama M, et al. Surgery for multiple lung metastases from alveolar soft-part sarcoma. *Surg Today* 1997;27:806–11.
- Tomimaru Y, Sasaki Y, Yamada T, et al. The significance of surgical resection for pulmonary metastasis from hepatocellular carcinoma. *Am J Surg* 2006;192:46–51.
- Kanzaki R, Higashiyama M, Fujiwara A, et al. Outcome of surgical resection of pulmonary metastasis from urinary tract transitional cell carcinoma. *Interact Cardiovasc Thorac Surg* 2010;11:60–4.
- Sakamoto T, Tsubota N, Iwanaga K, et al. Pulmonary resection for metastases from colorectal cancer. *Chest* 2001;119:1069–72.
- Kanemitsu Y, Kato T, Hirai T, et al. Preoperative probability model for predicting overall survival after resection of pulmonary metastases from colorectal cancer. *Br J Surg* 2004;91:112–20.
- Watanabe I, Arai T, Ono M, et al. Prognostic factors in resection of pulmonary metastasis from colorectal cancer. *Br J Surg* 2003;90:1436–40.
- Iizasa T, Suzuki M, Yoshida S, et al. Prediction of prognosis and surgical indications for pulmonary metastasectomy from colorectal cancer. *Ann Thorac Surg* 2006;82:254–60.
- Headrick JR, Miller DL, Nagorney DM, et al. Surgical treatment of hepatic and pulmonary metastases from colon cancer. *Ann Thorac Surg* 2001;71:975–9.
- Welter S, Jacobs J, Krbek T, Poettgen C, et al. Prognostic impact of lymph node involvement in pulmonary metastases from colorectal cancer. *Eur J Cardiothorac Surg* 2007;31:167–72.



Occult mediastinal lymph node metastasis in NSCLC patients diagnosed as clinical N0-1 by preoperative integrated FDG-PET/CT and CT: Risk factors, pattern, and histopathological study

Ryu Kanzaki^{a,*}, Masahiko Higashiyama^a, Ayako Fujiwara^a, Toshiteru Tokunaga^a, Jun Maeda^a, Jiro Okami^a, Takenori Kozuka^b, Takuya Hosoki^c, Yoshihisa Hasegawa^c, Motohisa Takami^c, Yasuhiko Tomita^d, Ken Kodama^a

^a Department of Thoracic Surgery, Osaka Medical Center for Cancer and Cardiovascular Diseases, 1-3-3 Nakamichi, Higashinari-ku, Osaka 537-8511, Japan

^b Department of Diagnostic Radiology, Osaka Medical Center for Cancer and Cardiovascular Diseases, 1-3-3 Nakamichi, Higashinari-ku, Osaka 537-8511, Japan

^c Morinomiya Clinic PET Center, 1-3-15 Nakamichi, Higashinari-ku, Osaka 537-0025, Japan

^d Department of Pathology, Osaka Medical Center for Cancer and Cardiovascular Diseases, 1-3-3 Nakamichi, Higashinari-ku, Osaka 537-8511, Japan

ARTICLE INFO

Article history:

Received 27 March 2010

Received in revised form 7 June 2010

Accepted 10 June 2010

Keywords:

Computed tomography

Lung cancer

Diagnosis

Positron emission tomography

ABSTRACT

Background: Integrated F18-fluorodeoxyglucose positron emission tomography/computed tomography (FDG-PET/CT) is widely used for mediastinal lymph node (MLN) staging in patients with non-small cell lung cancer (NSCLC). However, FDG-PET/CT has certain limitations. Prediction of occult MLN metastasis could allow selection of candidates for preoperative cervical mediastinoscopy or endobronchial ultrasound-guided transbronchial needle aspiration. This study defined risk factors for occult MLN metastasis in patients with NSCLC patients who were diagnosed as clinical N0-1 by preoperative integrated FDG-PET/CT and CT.

Methods: Consecutive patients with NSCLC who underwent staging using integrated FDG-PET/CT as an adjunct to CT prior to lung resection from October 2006 to September 2009 were evaluated retrospectively. The prevalence of MLN metastasis in patients diagnosed as clinical N0-1 was analyzed according to clinicopathological factors such as tumor location, tumor size, histology, and FDG uptake by the primary tumor. Risk factors for occult MLN metastasis were defined by multivariate analysis. Patterns of occult MLN metastasis were also analyzed and the involved MLNs were further examined histopathologically.

Results: The incidence of MLN metastasis was 11% (24 patients of 224). Multivariate analysis identified adenocarcinoma ($P=0.04$), tumors located in upper or middle lobe ($P=0.02$), tumor size >3 cm ($P=0.01$), and SUV_{max} of primary tumor >4.0 g/ml ($P=0.04$) as significant risk factors for MLN metastasis. The pattern of occult MLN metastasis was typical for NSCLC cases. The size of metastatic foci were small, with 68% of foci smaller than 4.0 mm.

Conclusions: The present study demonstrated that adenocarcinoma, tumors located in the upper or middle lobe, tumor size >3 cm, and SUV_{max} of primary tumor >4.0 g/ml are risk factors for occult MLN metastasis in patients with NSCLC who were diagnosed as clinical N0-1 by preoperative integrated FDG-PET/CT and CT. Patients with tumors located in the right upper or middle lobe are considered candidates for cervical mediastinoscopy because the involved metastatic mediastinal lymph nodes are easily accessible by these modalities.

© 2010 Elsevier Ireland Ltd. All rights reserved.

1. Introduction

Lung cancer is the leading cause of cancer-related deaths in the Western world [1]. Despite advances in surgical management, chemoradiotherapy, and early diagnosis, the prognosis for patients with lung cancer remains poor. Accurate staging for non-small cell

lung cancer (NSCLC) patients is important for both assessing prognosis and selecting the optimal therapy. Mediastinal lymph node (MLN) staging is particularly critical because survival is improved in patients with stage IIIA disease who undergo chemoradiotherapy followed by surgery, compared to those undergoing with surgery alone [2–4].

Chest computed tomography (CT) and cervical mediastinoscopy have been the gold-standard modalities for MLN staging in NSCLC patients. F18-fluorodeoxyglucose positron emission tomography (FDG-PET) has become increasingly utilized for MLN staging in

* Corresponding author. Tel.: +81 06 6972 1181; fax: +81 06 6981 8055.
E-mail address: rkanzaki@tj8.so-net.ne.jp (R. Kanzaki).

NSCLC patients. To overcome the inherent disadvantages of FDG-PET such as poor quality of the anatomical information, new imaging systems using integrated FDG-PET/CT were developed recently [5]. This integrated approach provides higher sensitivity compared with CT alone in MLN staging for NSCLC.

A practical approach adopted in many centers is scheduling patients with negative mediastinal uptake by integrated FDG-PET/CT for resection [6]. However, the incidence of occult MLN metastasis in NSCLC patients showing negative uptake by FDG-PET/CT is 7–16% [6–8]. Several studies therefore investigated factors associated with occult MLN metastasis in NSCLC patients [6–8], since a successful prediction could select candidates for either preoperative cervical mediastinoscopy or endobronchial ultrasound-guided transbronchial needle aspiration. However, these studies were limited in that no histological examination of the involved MLNs was undertaken nor was a source of the negative result by integrated FDG-PET/CT discussed sufficiently.

The aim of the present study was to define risk factors for occult MLN metastasis in patients with NSCLC diagnosed as clinical N0–1 by preoperative integrated FDG-PET/CT and CT. Histopathological examination of the involved MLNs was conducted to examine why these lymph nodes were diagnosed as negative. The patterns of occult MLN metastasis were also analyzed.

2. Patients and methods

2.1. Patient eligibility and staging

The study retrospectively evaluated patients with NSCLC in our hospital who underwent staging by integrated FDG-PET/CT as an adjunct to CT from October 2006 to September 2009. No patient underwent preoperative mediastinoscopy in this period. The following patients were excluded from the present study: those who underwent limited resection (wide-wedge resection or segmentectomy; $n = 144$), patients with diabetes mellitus ($n = 39$), patients who received neo-adjuvant chemotherapy ($n = 11$), and patients with positive mediastinal uptake on integrated PET-CT or enlarged MLNs on CT (i.e., clinical N2/N3; $n = 39$). The remaining 224 patients were staged as clinical N0 or N1 by integrated FDG-PET/CT and CT, and underwent resection with systematic lymph node dissection. The preoperative CT, integrated FDG-PET/CT, and pathological findings were reviewed. Preoperative clinical staging and postoperative pathological staging was based on the 1997 update of the TNM staging system [9]. The study group comprised 122 men and 102 women age from 30 to 83 years (mean age of 65.5 years). Table 1 details the patient characteristics and preoperative tumor evaluations.

2.2. CT imaging

All studies (CT and integrated FDG-PET/CT) were interpreted independently. CT examination was performed using a helical scanner (LightSpeed VCT, General Electric, Waukesha, WI; Aquilion 16, Toshiba, Tokyo, Japan), and all patients had contrast-enhanced CT. Lymph nodes were interpreted as positive if >1 cm across the short-axis diameter. All positive nodes were localized according to the lymph node stations, based on the classification by Naruke et al. [10].

A tumor was deemed central if its center was located in the inner 1/3 of the lung parenchyma (adjacent to the mediastinum) on transverse CT imaging. Non-centrally located tumors were identified as those centered in the outer 2/3 of the lung parenchyma on transverse CT imaging. All CT images were performed within 4 weeks of surgery.

Table 1
Characteristics of patients and tumors ($n = 224$).

Characteristics	Distribution (%)
Sex	
Male	122 (54)
Female	102 (46)
Age (years)	
Mean \pm SD	65.5 \pm 8.9
Range	30–83
Smoking status	
Smoker	131 (58)
Never smoker	93 (42)
Concurrent lung disease^a	
Present	40 (18)
Absent	184 (82)
Elevated serum CEA level (>5 ng/ml)	
Yes	68 (30)
No	156 (70)
Lobar distribution of the tumor	
RUL	100 (45)
RML	12 (5)
RLL	38 (17)
LUL	45 (20)
LLL	29 (13)
Location of the tumor	
Central	18 (8)
Non-central	206 (92)
Positive N1 node on PET/CT	
Yes	39 (17)
No	185 (83)
Tumor size (cm)	
Median	3.0
Mean \pm SD	3.1 \pm 1.5
Range	0.8–9.0
SUV_{max} of primary tumor	
Median	4.0
Mean \pm SD	5.6 \pm 5.1
Range	0–22.8

^a Concurrent lung disease includes interstitial lung disease, chronic obstructive pulmonary disorder, bronchial asthma, and tuberculosis.

2.3. Integrated FDG-PET/CT imaging

Patients were asked to fast, except for glucose-free oral hydration, for at least 5 h before the injection of ^{18}F -FDG (3.5 MBq/kg body weight). After injection of the tracer, patients were kept lying comfortably on a bed. No urinary bladder catheterization was performed and no oral muscle relaxants were administered. Whole-body PET/CT fusion scanning was performed 1 h after the injection, using a PET/CT system (Discovery LS, General Electric, Waukesha, WI; Biograph Duo LSD, Siemens-Asahi Medical Technologies, Tokyo, Japan). PET, CT, and integrated PET/CT images were available for review, displayed in axial, coronal, and sagittal planes. The FDG uptake of tumor was visually compared with that of the surrounding tissue in areas devoid of prominent artifacts and overlapping increased FDG uptake organs. A team of experienced radiologists reviewed the integrated FDG-PET/CT images independently from the CT data. Nodal uptake with a standardized uptake value (SUV_{max}) >2.5 were interpreted as positive. All integrated FDG-PET/CT imaging was performed within 4 weeks of surgery.

2.4. Surgical resection

All of the surgical resections and mediastinal nodal dissections were conducted by thoracic surgeons at Osaka Medical Center for Cancer and Cardiovascular Diseases. Systematic lymph node dissection was carried out.

Table 2
Postoperative pathological evaluation of the tumors.

Type of operation	
Lobectomy	213 (95%)
Bilobectomy	10 (4%)
Pneumonectomy	1 (1%)
Histopathological type	
Adenocarcinoma	180 (80%)
Squamous cell carcinoma	37 (17%)
Other types ^a	7 (3%)
Lymph node metastasis	
pN0	168 (75%)
pN1	32 (14%)
pN2	24 (11%)
Pathological stage after surgery ^b	
IA	101 (46%)
IB	64 (29%)
IIA	10 (4%)
IIB	21 (9%)
IIIA	23 (10%)
IIIB	5 (2%)

^a Other histopathological types of NSCLC included adeno-squamous carcinoma, large cell carcinoma.

^b Stage of disease was defined according to the 1997 update of TNM criteria established by UICC.

2.5. Pathological examination and the size of metastases

All resected tumor specimens were examined by experienced pulmonary pathologists. Histological classification of NSCLC was based on the WHO classification [11]. The dissected lymph nodes were examined histologically following hematoxylin and eosin staining, and the long-axis diameters of the metastatic foci in all involved lymph nodes were measured. The lymph node with the largest metastatic foci was selected as a representative foci of each station containing metastases.

2.6. Statistical analysis

Statistical analysis was performed with Dr.SPSS II software (SPSS Japan, Tokyo, Japan). Univariate data analysis was conducted using Fischer's exact test or Pearson's chi-square test. Multivariate analysis was conducted using the logistic regression (backwards stepwise) method. *P*-values were considered statistically significant if <0.05.

3. Results

3.1. The incidence and pattern of MLN metastasis

The incidence of MLN metastasis in this study was 11% (24 of 224 patients). Table 2 details the postoperative pathological evaluation of the tumors. Of 24 patients with mediastinal node metastasis, multistation MLN metastasis was found in 12 patients (50%), while the other 12 patients showed on a single station (50%). Skip metastasis was found in 10 patients (42%). Forty-four MLN stations in total were involved. Table 3 indicated the patterns of MLN involvement. In patients whose tumors were located in the right upper lobe (*n* = 12), 9 had metastasis in the superior lymph nodes (#1, #2, #3, #4) and 3 patients had metastases in both the superior and inferior (#7) lymph nodes. One patient whose tumor was located in the right middle lobe had metastasis in #3 and #7 lymph nodes, while a patient with a tumor in the right lower lobe had metastasis in #7 lymph node. In patients whose tumors were located in left upper lobes (*n* = 9), 7 patients had metastasis in the aortic lymph nodes (#5, #6), 1 patient had metastasis in #4 lymph nodes, and 1 patient had metastasis in #4 and #5 lymph nodes. Finally, a patient

Table 3
Pattern of mediastinal lymph node involvement.

Case	Lobar distribution	pN2 station	Pathological N1 node
1	RUL	1,3	Negative
2	RUL	2	Negative
3	RUL	3	Positive
4	RUL	3	Negative
5	RUL	3,4	Positive
6	RUL	3,4	Positive
7	RUL	3,4	Negative
8	RUL	4	Negative
9	RUL	1,2,3,4	Positive
10	RUL	1,2,4,7	Positive
11	RUL	1,3,4,7	Positive
12	RUL	1,3,4,7	Positive
13	RML	3,7	Negative
14	RLL	7	Negative
15	LUL	4	Positive
16	LUL	4,5	Negative
17	LUL	5	Positive
18	LUL	5	Positive
19	LUL	5	Positive
20	LUL	5	Negative
21	LUL	5	Negative
22	LUL	5,6	Positive
23	LUL	5,6	Positive
24	LLL	7	Positive

LLL, left lower lobe; LUL, left upper lobe; RLL, right lower lobe; RML, right middle lobe; RUL, right upper lobe.

whose tumor was located in the left lower lobe had metastasis in #7 lymph node.

3.2. Analysis of risk factors associated with MLN metastasis

Table 4 summarizes the results of univariate analysis for factors associated with MLN metastasis. Factors that are significantly associated with MLN metastasis are: never smoked (*P* = 0.03), tumor located in the upper or middle lobe (*P* = 0.01), tumor >3 cm (*P* = 0.005), SUV_{max} of primary tumor >4.0 g/ml (*P* = 0.0498), and adenocarcinoma (*P* = 0.04). The multivariate risk-factor analysis (Table 5) identified adenocarcinoma (*P* = 0.04), tumors located in the upper or middle lobe (*P* = 0.02), tumor >3 cm (*P* = 0.01), and SUV_{max} of primary tumor >4.0 g/ml (*P* = 0.04) as risk factors for MLN metastasis.

3.3. A Pathological examination and the size of metastases

The size of metastatic foci across 44 stations ranged from 0.5 to 9 mm with a mean value of 3.7 ± 2.0 mm (±SD). Thirty of the 44 involved stations (68%) had metastatic foci smaller than 4.0 mm. Table 6 details the size distribution of the metastatic foci.

4. Discussion

The present study defined risk factors for occult metastasis in patients with NSCLC diagnosed as clinical N0–1 by preoperative integrated FDG-PET/CT and CT. The patterns of occult MLN metastasis were also analyzed, and the involved MLNs were examined histologically. The incidence of MLN metastasis in our series of patients was 11%. This finding was concordant with comparable previous studies; Al-Sarraf et al. [6] reported the incidence of occult N2 disease in similar patients as 16%, while Cerfolio et al. [12] reported a 14% incidence, although when they limited the analysis to clinical stage I patients, the incidence decreased to 7% [7]. Risk factors for occult N2 disease reported in these previous studies were as follows: adenocarcinoma [7], tumors located in right upper lobe [6], larger tumor size [7], a high SUV_{max} of the primary tumor [7,12], centrally located tumors [6,7], positive N1 nodes on PET [6],

Table 4
Univariate analysis for factors associated with mediastinal lymph node metastasis.

Variable	Pathological N2 (n=24)	Pathological N0-1 (n=200)	P-value
Sex			
Male	9	113	N.S.
Female	15	87	
Age (years)			
≥65	13	114	N.S.
<65	11	86	
Smoking status			
Smoker	9	122	0.03
Never smoker	15	78	
Concurrent lung disease			
Present	2	38	N.S.
Absent	22	162	
Elevated serum CEA level (>5 ng/ml)			
Yes	7	61	N.S.
No	17	139	
Lobar distribution of the tumor ^a			
Upper or middle lobe	22	135	0.01
Lower lobe	2	65	
Location of the tumor			
Central	2	16	N.S.
Non-central	22	184	
Positive N1 node on PET/CT			
Yes	3	36	N.S.
No	21	164	
Tumor size (cm)			
>3.0	18	90	0.005
≤3.0	6	110	
SUV _{max} of primary tumor			
>4.0	16	91	0.0498
≤4.0	8	109	
Histopathological type			
Adenocarcinoma	23	157	0.04
Non-adenocarcinoma	1	43	

^a Upper or middle lobe includes right upper lobe, right middle lobe, and left upper lobe. Lower lobe includes right lower lobe, left lower lobe.

Table 5
Multivariate analysis for risk factors for mediastinal lymph node metastasis.

Variable	Odds ratio	Confidence interval	P-value
Adenocarcinoma	9.26	1.13–76.9	0.04
Located in upper or middle lobe	6.29	1.36–29.4	0.02
Tumor size >3 cm	4.18	1.48–11.76	0.01
SUV _{max} of primary tumor >4.0	2.79	1.04–7.52	0.04

and poorly differentiated histology [12]. A limitation of these studies was the lack of histological examination of the involved MLNs and insufficient consideration of why the involved lymph nodes showed as negative by integrated FDG-PET/CT.

Adenocarcinoma was also identified as a risk factor for occult MLN metastasis in the present patient cohort. Interestingly, Lee et al. [7] reported similar data in that all of their 16 patients with pathological N2 disease showed adenocarcinoma as the primary tumor cell type. On the contrary, in the Al-Sarraf and co-workers' series [6], the primary tumor cell type did not affect the incidence of occult MLN metastasis. A major difference between their series and

ours is the inclusion of patients with enlarged lymph nodes (>1 cm) on CT in our analysis. In the comparative study [6], 8 of 25 patients with pathological N2 disease in their series had enlarged lymph nodes (>1 cm) on CT, and 12 of the 25 had non-adenocarcinoma tumors. It is reported that lymph node metastases from adenocarcinoma were of normal size (1 cm ≤ across the short-axis diameter) more frequently than those from squamous cell carcinoma [13,14]. Therefore, we speculate that patients with MLN metastasis from squamous cell carcinoma tended to be excluded in our series and included in the series of patients reported on by Al-Sarraf and co-workers [6].

Tumor location in the upper or middle lobe also proved to be a risk factor for occult MLN metastasis in the present study. A predisposition of lobar distribution was also observed in the previous study by Al-Sarraf and co-workers, with the right upper lobe dominating in their series. The other risk factors for occult MLN metastasis were identified as tumor size >3 cm, SUV_{max} of primary tumor >4.0 g/ml. It is well known that the incidence of MLN metastasis increased as the tumor size increased [15]. The relationship between SUV_{max} of the primary tumor and incidence of lymph node metastasis has also been investigated previously. Downey and colleagues [16] reported higher SUV_{max} of the primary tumors in patients with pathological nodal involvement than in patients without nodal involvement, while Cerfolio and colleagues [8] showed that SUV_{max} of lung tumors increased as the cancer progressed from N0 to N3.

In contrast to the report of Al-Sarraf et al. [6], positive N1 nodes on PET was not a risk factor for MLN metastasis in the present study. This difference might be attributable to the relatively high incidence of skip metastasis in the present series (42%).

To overcome a limitation of similar previous studies, we conducted histological examination of the involved MLNs. This additional analysis demonstrated that metastatic foci of occult MLN metastases were small; of 44 stations, the size of foci ranged from 0.5 to 9 mm with a mean value of 3.7 ± 2.0 mm, and 30 of the 44 involved stations (68%) contained foci of <4.0 mm diameter. This finding is supported by previous studies wherein the metastatic foci of false-negative lymph nodes by PET were small. Takamochi et al. [17] reported such foci to be 1–7.5 mm (mean, 3.4 mm; n = 12), while Nomori et al. [18] found that false-negative (n = 8) and true-positive (n = 28) lymph nodes on PET contained foci ranging from 0.5 to 9 mm (mean, 3 mm) and 4–18 mm (mean, 10 mm), respectively. No metastatic foci smaller than 4 mm were detected with PET.

The pattern of occult MLN metastasis shown in the present study was identical to the typical distribution in NSCLC patients reported by Naruke et al. [19] These authors reviewed 1815 patients who underwent systematic lymph node dissection, and examined which nodes had the highest likelihood of metastasis. Distribution of MLN metastasis from each lobe in their series was as follows: right upper lobe tumor, #3 (12.3%) and/or #4 (8%); right middle lobe tumor, #3 and/or #7 (16.4%); right lower lobe tumor, #7 (13.7%); left upper lobe tumor, #5 (12.3%) and/or #6 (6.7%); and left lower lobe tumor, #7 (11.9%). Based on our data, patients with tumors in the right upper or middle lobe are potential candidates for cervical mediastinoscopy because their possible metastatic mediastinal lymph nodes (#3, #4, #1, #2, #7) are easily accessible by cervical mediastinoscopy or endobronchial ultrasound-guided transbronchial needle aspiration. Skip metastasis was found in 10 patients (42%) in

Table 6
Distribution of size of metastatic foci.

	Size (mm)											Total
	<1.0	<2.0	<3.0	<4.0	<5.0	<6.0	<7.0	<8.0	<9.0	<10.0	≥10.0	
No.	1	5	12	12	5	3	2	1	0	3	0	44

the present series, which is a relatively high incidence compared to other reports for NSCLC patients (25–29%) [15]. In these 10 patients, it is speculated that metastatic MLNs were negative for integrated FDG-PET/CT and CT because they are the first nodal target of lymph from a primary tumor site and metastatic foci were still small.

Because of the small size of the metastatic foci, occult MLNs are considered difficult to be detected by endobronchial ultrasound-guided transbronchial needle aspiration. Therefore, we recommend cervical mediastinoscopy for the patients with tumors in the right upper or middle lobe who have risk factors for occult MLN metastasis.

5. Conclusions

The present study demonstrated that adenocarcinoma, tumors located in the upper or middle lobe, tumor size >3 cm, and SUV_{max} of primary tumor >4.0 g/ml are risk factors for occult MLN metastasis in patients with NSCLC diagnosed as clinical N0-1 by preoperative integrated FDG-PET/CT and CT. The metastatic foci of involved stations were small, with 68% of foci measuring <4.0 mm. The pattern of occult MLN metastasis discerned in the present study was typical for reported distribution of metastatic foci in NSCLC patients. Patients with tumors in the right upper or middle lobe are considered candidates for cervical mediastinoscopy because their metastatic mediastinal lymph nodes (#3, #4, #1, #2, #7) are easily accessible by these modalities.

Conflict of interest statement

None of the authors has any financial or other potential conflict of interest.

Acknowledgement

This study was partly supported by a grant from Osaka Anti-cancer Foundation.

References

- [1] Jemal A, Siegel R, Ward E, Murray T, Xu J, Thun MJ. Cancer statistics, 2007. *CA Cancer J Clin* 2007;57:43–66.
- [2] Roth JA, Atkinson EN, Fossella F, Komaki R, Bernadette RM, Putnam Jr JB, et al. Long-term follow-up of patients enrolled in a randomized trial comparing perioperative chemotherapy and surgery with surgery alone in resectable stage IIIA non-small-cell lung cancer. *Lung Cancer* 1998;21:1–6.
- [3] Rusch VW, Albain KS, Crowley JJ, Rice TW, Lonchyna V, McKenna Jr R, et al. Surgical resection of stage IIIA and stage IIIB non-small-cell lung cancer after concurrent induction chemoradiotherapy. A Southwest Oncology Group trial. *J Thorac Cardiovasc Surg* 1993;105:97–104, discussion 104–6.
- [4] Rosell R, Gomez-Codina J, Camps C, Maestre J, Padille J, Canto A, et al. A randomized trial comparing preoperative chemotherapy plus surgery with surgery alone in patients with non-small-cell lung cancer. *N Engl J Med* 1994;330:153–8.
- [5] Cerfolio RJ, Ojha B, Bryant AS, Raghuvver V, Mountz JM, Bartolucci AA. The accuracy of integrated PET-CT compared with dedicated PET alone for the staging of patients with nonsmall cell lung cancer. *Ann Thorac Surg* 2004;78:1017–23, discussion 1017–23.
- [6] Al-Sarraf N, Aziz R, Gately K, Lucey J, Wilson L, McGovern E, et al. Pattern and predictors of occult mediastinal lymph node involvement in non-small cell lung cancer patients with negative mediastinal uptake on positron emission tomography. *Eur J Cardiothorac Surg* 2008;33:104–9.
- [7] Lee PC, Port JL, Korst RJ, Liss Y, Meherally DN, Altorki NK. Risk factors for occult mediastinal metastases in clinical stage I non-small cell lung cancer. *Ann Thorac Surg* 2007;84:177–81.
- [8] Cerfolio RJ, Bryant AS, Ojha B, Bartolucci AA. The maximum standardized uptake values on positron emission tomography of a non-small cell lung cancer predict stage, recurrence, and survival. *J Thorac Cardiovasc Surg* 2005;130:151–9.
- [9] Mountain CF. Revisions in the International System for Staging Lung Cancer. *Chest* 1997;111:1710–7.
- [10] Naruke T, Suemasu K, Ishikawa S. Lymph node mapping and curability at various levels of metastasis in resected lung cancer. *J Thorac Cardiovasc Surg* 1978;76:832–9.
- [11] World Health Organization Histological Typing of Lung and Pleural Tumours. 4th ed. Lyon: IARC Press; 2004.
- [12] Cerfolio RJ, Bryant AS, Eloubeidi MA. Routine mediastinoscopy and esophageal ultrasound fine-needle aspiration in patients with non-small cell lung cancer who are clinically N2 negative: a prospective study. *Chest* 2006;130:1791–5.
- [13] Mori K, Yokoi K, Saito Y, Tominaga K, Miyazawa N. Diagnosis of mediastinal lymph node metastases in lung cancer. *Jpn J Clin Oncol* 1992;22:35–40.
- [14] Kamiyoshihara M, Kawashima O, Ishikawa S, Morishita Y. Mediastinal lymph node evaluation by computed tomographic scan in lung cancer. *J Cardiovasc Surg* 2001;42:119–24.
- [15] Shields TW. Pathology of carcinoma of the lung. In: Shields TW, LoCicero J, Reed CE, Feins RH, editors. *General Thoracic Surgery*, vol. II, 7th ed. New York: Lippincott Williams & Wilkins; 2009. p. 1311–40.
- [16] Downey RJ, Akhurst T, Gonen M, Vincent A, Bains MS, Larson S, et al. Preoperative F-18 fluorodeoxyglucose-positron emission tomography maximal standardized uptake value predicts survival after lung cancer resection. *J Clin Oncol* 2004;22:3255–60.
- [17] Takamochi K, Yoshida J, Murakami K, Niho S, Ishii G, Nishimura M, et al. Pitfalls in lymph node staging with positron emission tomography in non-small cell lung cancer patients. *Lung Cancer* 2005;47:235–42.
- [18] Nomori H, Watanabe K, Ohtsuka T, Naruke T, Suemasu K, Uno K. The size of metastatic foci and lymph nodes yielding false-negative and false-positive lymph node staging with positron emission tomography in patients with lung cancer. *J Thorac Cardiovasc Surg* 2004;127:1087–92.
- [19] Naruke T, Tsuchiya R, Kondo H, Nakayama H, Asamura H. Lymph node sampling in lung cancer: how should it be done? *Eur J Cardiothorac Surg* 1999;16(Suppl. 1):S17–24.



Alveolar space filling ratio as a favorable prognostic factor in small peripheral squamous cell carcinoma of the lung

Yukihiro Watanabe^a, Tomoyuki Yokose^{a,*}, Yuji Sakuma^d, Chikako Hasegawa^a, Haruhiro Saito^b, Kozo Yamada^b, Hiroyuki Ito^c, Masahiro Tsuboi^c, Haruhiko Nakayama^c, Yoichi Kameda^a

^a Pathology Division, Kanagawa Cancer Center Research Institute, 2-1-1, Nakao, Asahi-Ku, Yokohama, Kanagawa 241-0815, Japan

^b Respiratory Medicine Division, Kanagawa Cancer Center Research Institute, 2-1-1, Nakao, Asahi-Ku, Yokohama, Kanagawa 241-0815, Japan

^c Thoracic Surgery Division, Kanagawa Cancer Center Hospital, Kanagawa Cancer Center Research Institute, 2-1-1, Nakao, Asahi-Ku, Yokohama, Kanagawa 241-0815, Japan

^d Molecular Pathology and Genetics Division, Kanagawa Cancer Center Research Institute, 2-1-1, Nakao, Asahi-Ku, Yokohama, Kanagawa 241-0815, Japan

ARTICLE INFO

Article history:

Received 11 June 2010

Received in revised form

21 September 2010

Accepted 3 December 2010

Keywords:

Lung

Squamous cell carcinoma

Peripheral type

Alveolar space filling ratio

Prognostic factor

Microinvasive carcinoma

ABSTRACT

Introduction: Squamous cell carcinomas (SqCCs) of the lung can be divided into two types according to the location of primary site; one is central type and another is peripheral type. Many reports on the central type revealed the clinicopathological characteristics relating carcinogenesis, therapeutics and prognosis. On the other hand, those on the peripheral type are very a few and prognostic indicators of peripheral type have not been enough elucidated. The aim of this study was to clarify clinicopathological prognostic factors of small peripheral SqCCs of the lung 30 mm or less.

Materials and methods: We evaluated various 15 clinicopathological parameters in 81 patients with peripheral type SqCCs, which are defined as tumors located in or more peripheral from the third branching bronchus, measuring 30 mm or less in diameter.

Results: Univariate analyses were performed using the log rank test and multivariate analyses using logistic regression model. As a result, two factors had a statistically significant influence on outcome of the patients in the univariate analysis; no relapse was observed in the patients with the ratio of alveolar space filling (ASF) area to tumor area of 70% or more and the maximum diameter of invasive area measuring 10 mm or less in size ($P=0.0214$, $P=0.0373$, respectively). Meanwhile, multivariate analysis showed that the ASF ratio of 70% or more significantly affected the outcome of the patients ($P=0.0337$), however the maximum diameter of invasive area did not ($P=0.2136$). We could not show the unfavorable prognostic factor contributory to tumor relapse.

Conclusions: We have shown that the ASF ratio is a significantly favorable prognostic factor for small peripheral type. Especially the focally invasive tumors with ASF ratio of 70% or more might be classified as “a microinvasive carcinoma” of the peripheral SqCCs of the lung and tumors with ASF ratio 100% as noninvasive carcinoma.

© 2010 Elsevier Ireland Ltd. All rights reserved.

1. Introduction

Lung cancer is the leading cause of cancer death in the United States and other countries, including Japan. Squamous cell carcinoma (SqCC) occupies approximately 30% in all lung cancers [1]. About 70% of SqCC has been reported to arise from central portion of the lung, whereas the remaining 30% does from periphery [2]. According to a statistical report by Kodama in Japan, the peripheral type of SqCC goes on increasing around 50% of all SqCCs of the lung [3]. Many reports of the central type SqCC have revealed the pathological and clinical characteristics relating therapeutics

and prognosis, but those of the peripheral type have been very a few because the biological behavior has not been well evaluated [4–6].

Central type SqCC of the lung is generally believed to arise from bronchial dysplastic epithelium through the multiple steps of dysplasia–carcinoma sequence, which is the same well-known carcinogenic pathway as observed in uterine cervical neoplasia [7]. During this pathway, carcinoma in situ (CIS) is the earliest stage of the progression of central type SqCC and showed favorable prognosis. If the CIS lesion is localized, a non-surgical procedure like photodynamic therapy may be selected instead of operation. On the other hand, the carcinogenic pathway observed as dysplasia–CIS sequence in the central type has not been fully established in the peripheral SqCC of the lung, so that the standard treatment of peripheral SqCC in the early stage is confined to lobectomy at the present time.

* Corresponding author. Tel.: +81 45 391 5761; fax: +81 45 361 4692.

E-mail address: yokose-t@kcch.jp (T. Yokose).




Fig. 1. Peripheral squamous cell carcinoma (SqCC) consisting of two types of area; alveolar space filling (ASF) area and non-ASF area. (A) Histology of alveolar space filling (ASF) area. Hematoxylin and eosin (H&E). (B) Elastic fiber framework of ASF area. Elastica von Gieson (EVG). (C) Histology of non-ASF area. H&E. (D) Destruction of elastic fiber framework was observed. EVG.

Recently, several studies of peripheral SqCC have been presented. Funai et al. studied 109 cases of peripheral SqCC and indicated that those can be divided into three subtypes based on two distinctive structural patterns based on the tumor growth pattern and the condition of the elastic fiber framework (EFF): the alveolar space filling (ASF) type, the expanding type and the mixed type consisting of the mixture of the former two types [5]. Pure ASF type tumors, which showed no destruction of elastic fiber framework, had a favorable prognosis, showed 100% 5-year survival and thought to be classified as CIS of peripheral SqCCs. Meanwhile, Maeshima et al. studied 101 cases of peripheral SqCCs and proposed three types based on the minimal tumor nest (MTN) pattern: large nest consisting of more than 6 tumor cells, small nest consisting of 2–5 tumor cells, or single cell [6]. They considered tumors with single cell invasive component appeared to be highly malignant, whereas those composed only of large tumor nest or small tumor nest components seemed to have a relatively low potential for malignancy and recurrence, despite these two types also being invasive.

We consider that the clarification of clinicopathological prognostic factors in the peripheral SqCCs leads to the prediction of recurrence or survival and contribution to increase of therapeutic selections. The aim of this study was to identify favorable and unfavorable clinicopathological factors for small peripheral SqCCs of the lung.

2. Materials and methods

2.1. Patients

Between January 1993 and September 2008, surgical resection was performed in 364 cases of lung SqCCs at Kanagawa Cancer Center Hospital. The subjects of the study were consecutive 81 patients

with small peripheral SqCC of the lung measuring 30 mm or less in diameter. “Peripheral type” in this study was defined as tumors located in or more peripheral to the third branching bronchus.

The patients included 72 men and 9 women with a median age of 70.5 years (range, 47–84 years). We obtained the various clinical parameters from a medical record of each patient, including age, gender, smoking habit, exposure to asbestos, and limited or standard resection. The cases were classified according to the 7th edition of the Union International Against Cancer Staging System [8].

2.2. Pathologic review

All tumor slides stained with hematoxylin and eosin (H&E) and by the Elastica von Gieson method (EvG) for elastic fibers were reviewed by two of the authors (YW and TY) to confirm the histological differentiation, tumor stage, vessel invasion, lymphatic permeation, pleural involvement, presence or absence of usual interstitial pneumonia (UIP), the degree of lymphocytic inflammatory cell infiltration into the tumor nests, the size of necrotic area, the maximum diameter of invasive growth area which was characterized by the destruction of EFF and stromal cell reaction (Fig. 1), the proportion of ASF area to the entire tumor area (ASF ratio), and the minimal tumor nest (MTN) pattern. The degree of inflammation was defined as follows; mild lymphocytic tumor nest infiltration, the number of less than 10% of tumor cells per nest; moderate, 10 to less than 50%; severe, 50% or more. Definition of UIP met histological features of UIP pattern in the American Thoracic Society/European Respiratory Society international multidisciplinary consensus classification of idiopathic interstitial pneumonias [9]. ASF area showed tumor cells fill the alveoli and grow from one alveolus to another with no destruction of the elastic framework as if tumor cells penetrate the pores of Cohn and no desmoplastic

change in stroma which usually occurs in invasive lesion (Fig. 1) [5]. When a tumor had areas of tumor nests surrounded by thin fibrous septa without elastic fibers, the areas were excluded from ASF area. We determined the ASF ratio by visual examination using H&E and EvG staining specimens and 70% as a threshold (data not shown). A tumor nest was defined as either a nested group of tumor cells or a single tumor cell invading individually, similarly the MTN was defined as the smallest group of tumor cells observed in the primary tumor. The size of the MTN in each patient was classified as follows according to the report of Maeshima et al.: large nest (consisting of more than 6 tumor cells), small nest (consisting of 2–5 tumor cells) or single cell [6]. When discordant assessments arose, consensus was achieved after joint review of the specimen over a multiheaded microscope.

2.3. Statistical analysis

Log-rank test was used to select significant factors examined for discriminating the patients with a relapse from those without a relapse. To identify which factors had a significant influence on the patients' outcome, logistic regression models were used. The correlations between clinicopathological variables were evaluated by the chi-square test or Fisher's exact test, as appropriate. *P*-values <0.05 were considered significant.

3. Results

We reviewed 81 patients with SqCC of the peripheral lung measuring 30 mm or less in diameter. Clinicopathological characteristics are shown in Table 1. Peripheral SqCC patients showed male preponderance (89%) and high smoking habit (99%). In the present population, 70% took lobectomy, 80% diagnosed as T1a or T1b, 75% showed the tumor size over 20 mm, vessel invasion was more frequent than lymphatic permeation and lymph node metastasis was only 9%. UIP was observed in 9% of the patients. The recurrence rate for all patients was 19.7%. In the univariate analysis, only two factors had a statistically significant influence on the patients' outcome (Table 2). Neither the patients with ASF ratio 70% or more nor the maximum diameter of invasive area of 10 mm or less in size relapsed (ASF area, $P=0.0214$; maximum diameter of invasive area, $P=0.0373$). None of other factors; MTN size, lymph node metastasis, vessel invasion and lymphatic permeation showed significant difference in disease free survival (DFS) rate. Multivariate analysis showed that the ASF ratio 70% or more ($P=0.0337$), not the maximum diameter of invasive area ($P=0.2136$), significantly affected the outcome of the patients (Table 3). Fig. 2 shows disease-free survival curves for patients with ASF ratio 70% or more and ASF ratio less than 70%. Table 4 shows the clinicopathological characteristics of the tumors with ASF ratio 70% or more in comparison to those with ASF ratio less than 70%. The tumors with ASF ratio 70% or more showed female preference, less maximum diameter of tumor invasion size or vascular invasion, less single cell pattern, and less inflammation. Moreover, no patients with those tumors showed recurrence or death.

4. Discussion

In the present study, we evaluated several clinicopathological factors to reveal prognostic indications for peripheral lung SqCCs and found ASF ratio to be significant. The multivariate analysis showed that the patients having focally invasive tumors with ASF ratio 70% or more had no recurrence as same as those pure ASF type SqCCs (ASF ratio 100%).

The definition of ASF depends on the observation of alveolar EFF whose complete preservation is considered to represent noninva-

Table 1
Patients characteristics ($n=81$).

	Number of patients (%)
Total	81
Gender	
Male	72 (89%)
Female	9 (11%)
Age (years, range)	70.5 (47–84)
Smoking habit	
Non-smoker	1 (1%)
Past-smoker	36 (45%)
Current-smoker	44 (54%)
Surgery	
Lobectomy + LN	57 (70%)
Partial resection	24 (30%)
T factor	
T1a	22 (27%)
T1b	43 (53%)
T2	16 (20%)
Tumor size	
<20 mm	20 (25%)
20–30 mm	61 (75%)
Lymph node metastasis	
Negative	74 (91%)
Positive	7 (9%)
Vessel invasion	
Negative	55 (68%)
Positive	26 (32%)
Lymphatic permeation	
Negative	71 (87%)
Positive	10 (13%)
Pleural involvement	
P0	65 (80%)
P1–3	16 (20%)
Exposure to asbestos	
Negative	72 (88%)
Positive	9 (12%)
Tumor location	
Patients without UIP	
Upper lobe	39 (48%)
Middle and lower lobe	33 (41%)
Patients with UIP	
Upper lobe	2 (2%)
Middle and lower lobe	7 (9%)

SqCC, squamous cell carcinoma; UIP, usual interstitial pneumonia.

sive area. It has been well-known in lung adenocarcinoma that the preservation of alveolar structure [10–13] and alveolar EFF has significant association with prognosis [14–16]. Funai et al. considered pure ASF type tumors on peripheral SqCCs of the lung to be non-invasive cancers which were thought to be equivalent to tumors without EFF destruction on adenocarcinomas of the lung [5]. However, only 5% (5 cases of 109) of the patients had pure ASF type. That was only 1 case 1.2% in our study. The prognostic factors that are applicable to the majority of peripheral SqCCs have been desired. Since Funai et al. did not refer the proportion of ASF, we attempted to evaluate the ratio to clarify the significance as a prognostic factor. In the present study, nineteen patients (23.5%) had a tumor with ASF ratio 70% or more which was equivalent to a pure ASF type tumor of the Funai's study; we were able to select more cases with favorable prognosis in manner of detailed analysis of ASF ratio than Funai's study. In addition, the invasive tumors with ASF ratio 70% or more may appear to have low grade malignancy and no recurrence, therefore it could be said that they are micro-invasive carcinomas.

In 2009, Samuel and Yousem studied 62 peripheral SqCCs of lung focusing on the patterns of marginal or peripheral growth and rate of an alveolar filling (AF) component [17]. Comparing those tumors with 20% or more of their diameters composed of an AF pattern to those tumors of equivalent diameter without AF pattern, no differences in average survival were observed. In our advance investigation, we obtained the same result (data was not shown).

Table 2
Examination of prognostic factors on recurrence in peripheral squamous cell carcinoma of lung by univariate analysis.

Clinicopathological factors	Number of no recurrence	Number of recurrence	Univariate analysis (P value)
Total	65	16	
Gender			
Male	57	15	0.676
Female	8	1	
Brinkman index			
0–999	34	8	0.870
1000–	31	8	
Smoking habit			
Current smoker	8	36	0.765
Past smoker	8	28	
No smoker	0	1	
T factor			
T1a	17	4	0.792
T1b	36	8	
T2	12	4	
Lymph node metastasis			
Negative	74	16	0.229
Positive	7	0	
Vessel invasion			
Negative	44	11	0.701
Positive	21	5	
Lymphatic permeation			
Negative	55	16	0.146
Positive	10	0	
Necrotising area			
<50%	47	12	0.869
≥50%	18	4	
Pleural involvement			
P0	53	12	0.541
P1–3	12	4	
Size of minimal tumor nest			
Large nest	4	0	0.284
Small nest	15	2	
Single cell	46	14	
Differentiation			
Well	5	3	0.314
Moderate	27	7	
Poor	33	6	
Alveolar space filling			
≥70%	19	0	0.0214
<70%	46	16	
Inflammatory cell infiltration			
Mild	60	11	0.304
Moderate or severe	21	5	
Background lung			
With UIP	7	2	0.739
Without UIP	58	14	
Diameter of invasion area			
<10 mm	16	0	0.0373
≥10 mm	49	16	

The maximum diameter of invasive area (MDIA) of 10 mm or less in size had a statistically significant influence on the patients' outcome in univariate analysis but not in multivariate analysis. MDIA in peripheral SqCC had not been well evaluated, although the MDIA of 5 mm or less in adenocarcinoma was proved to be a good prog-

Table 3
Examination of prognostic factors on recurrence in peripheral squamous cell carcinoma of lung by multivariate analysis.

Parameters	P value
ASF ratio 70% or more	0.038
Invasive diameter 10mm or less	0.094
T factor	0.454
Age	0.428
Histological differentiation	0.493
Size of minimal tumor nest	0.826
Sex	0.936

Table 4
Correlation between clinicopathological features and peripheral squamous cell carcinoma of lung with ASF ratio 70% or more and less than 70%.

Factors	ASF ≥ 70% (n = 19)	70% > ASF (n = 62)	P value
Sex			0.046*
Male	14	58	
Female	5	4	0.702
Age (years)			
<70	8	21	
≤70	11	41	
Asbestos exposure			0.745
Absent	16	56	
Present	3	6	
Tumor size (mm)			0.221
≤20	8	15	
<20	11	47	
T			0.867
1a + 1b	16	49	
2	3	13	
N			0.894
0	18	56	
1 or 2	1	6	
ly			0.357
Negative	15	56	
Positive	4	6	
v			0.043*
Negative	17	38	
Positive	2	24	
PL			0.867
P0	16	49	
P1–3	3	13	
pm			0.528
pm0	19	61	
pm1	0	1	
Differentiation			0.226
Well	0	8	
Moderate or poorly	19	54	
Necrosis			0.696
Mild	15	44	
Moderate or severe	4	18	
Nesting			0.020*
Small or large	10	51	
Single cell	9	11	
Tumor invasion (mm)			0.00*
≤10	12	4	
<10	7	58	
Inflammation			0.040*
Mild	18	42	
Moderate or severe	1	20	
Interstitial pneumonia			0.178
Absent	19	53	
Present	0	9	
Recurrence			0.032*
Absent	19	46	
Present	0	16	
Death			0.032*
Live	19	46	
Died	0	16	

* Statistically significant factor.

nosis factor and thought to represent a microinvasive carcinoma [18]. We think that MDIA lost significance because ASF ratio 70% or more became a confounding factor to MDIA as shown in Table 4.

Inflammation is generally thought to be a positive prognostic factor in non-small cell carcinoma, but not in this case [19]. We have supposed that previous reports which regarded inflammation as a favorable factor conducted prognostification in patients with advanced cancers. But we compared probably minimally invasive cancers with overt invasive cancers; therefore analysis of inflammation might bring an opposite result.

In other studies, vascular invasion, lymphatic permeation, and perineural invasion have been reported to be important prognostic factors in peripheral SqCC. Saijo et al. reported that peripheral type SqCC had higher vascular invasion and higher pleural invasion and




Fig. 2. Disease-free survival curves for patients with ASF ratio 70% or more and ASF ratio less than 70%.

with less lymph node metastasis compared with central type SqCC [20]. In our study those all factors had not influence to patient's outcome, however, only vascular invasion was observed frequently in the tumors with ASF ratio less than 70%, significantly, and probably the study of Saijo et al. included many invasive tumors with ASF less than 70%.

We propose from the present data that peripheral invasive SqCC with ASF ratio 70% or more could be classified as microinvasive carcinoma and peripheral SqCC with ASF ratio 100% as noninvasive carcinoma. Anatomical lobectomy with mediastinal lymph node dissection is the present standard of care for such patients with stage I or II, and adjuvant chemotherapy may be performed in patients with stage IB or more advanced stage. In a similar manner to noninvasive adenocarcinoma of the lung, limited resection could be an acceptable treatment for a tumor with ASF ratio 100% and partial resection or simple lobectomy without lymph node dissection and/or adjuvant therapy for a tumor with ASF ratio 70% or more in the future. However, since it might be difficult to select candidate cases of favorable peripheral SqCC by the radiological procedure in the present, additional studies are needed before modification of the standard therapy by the basis of the proposed classification.

Conflict of interest statement

The authors report no conflicts of interest.

Acknowledgements

We thank Ms. Seki and Ms. Ueda for their technical assistance. This work was supported in part by the Kanagawa Cancer Research Fund and Kanagawa Prefectural Hospitals Cancer Research Fund.

References

- [1] Kakizoe T, Sugiura S, Segami K, Yamaguchi N, Shimizu H, Yamada T, et al. Cancer statistics in Japan 1999. Tokyo, Japan: Foundation of Promotion of Cancer Research; 1999. p. 12.
- [2] Travis WD, Brambilla E, Muller-Hermelink HK, Harris CC, editors. World Health Organization classification of tumors. Pathology and genetic of tumors of the lung, pleura, thymus and heart. Lyon: IARC Press; 2004. p. 26–30.
- [3] Furonaka O, Takeshima Y, Awaya H, Ishida H, Kohno N, Inai K. Aberrant methylation of *p14ARF*, *p15INK4b* and *p16INK4a* genes and location of the primary site in pulmonary squamous cell carcinoma. *Pathol Int* 2004;54: 549–55.
- [4] Sakurai H, Asamura H, Watanabe S, Suzuki K, Tsuchiya R. Clinicopathological features of peripheral squamous cell carcinoma of the lung. *Ann Thorac Surg* 2004;78:222–7.
- [5] Funai K, Yokose T, Ishii G, Araki K, Yoshida J, Nishimura M, et al. Clinicopathological characteristics of peripheral squamous cell carcinoma of the lung. *Am J Surg Pathol* 2003;27:978–84.
- [6] Maeshima A, Maeshima A, Asamura H, Matsuno Y. Histologic prognostic factors for small-sized squamous cell carcinomas of the peripheral lung. *Lung Cancer* 2006;52:53–8.
- [7] Carter D. Squamous cell carcinoma of the lung: an update. *Semin Diagn Pathol* 1985;2(4):226–34.
- [8] Sobin L, Gospodarowicz M, Wittekind C, editors. International union against cancer. TNM classification of malignant tumours. 7th ed. West Sussex, UK: John Wiley and Sons Ltd.; 2009.
- [9] American Thoracic Society, European Respiratory Society. American Thoracic Society/European Respiratory Society international multidisciplinary consensus classification of idiopathic interstitial pneumonias. General principles and recommendations. *Am J Respir Crit Care Med* 2002;165:277–304.
- [10] Kurokawa T, Matsuno Y, Noguchi M, Mizuno S, Shimotsato Y. Surgically curable early adenocarcinoma in the periphery of the lung. *Am J Surg Pathol* 1994;18(5):431–8.
- [11] Terasaki H, Niki T, Matsuno Y, Yamada T, Maeshima A, Asamura H, et al. Lung adenocarcinoma with mixed bronchioalveolar and invasive components. Clinicopathological features, subclassification by extent of invasive foci, and immunohistochemical characterization. *Am J Surg Pathol* 2003;27:937–51.
- [12] Fujita A, Kameda Y, Goya T. Clinicopathology of stromal invasion in lung adenocarcinoma. *Pathol Int* 2009;59:1–6.
- [13] Sakuma Y, Okamoto N, Saito H, Yamada K, Yokose T, Kiyohashi M, et al. A logistic regression predictive model and the outcome of patients with resected lung adenocarcinoma of 2 cm or less in size. *Lung Cancer* 2009;65: 85–90.
- [14] Noguchi M, Morikawa A, Kawasaki M, Matsuno Y, Yamada T, Hirohashi S, et al. Small adenocarcinoma of the lung. Histologic characteristics and prognosis. *Cancer* 1995;75(12):2844–52.
- [15] Yokose T, Suzuki K, Nagai K, Nishikawa Y, Sasaki S, Ochiai A, et al. Favorable and unfavorable morphological prognostic factors in peripheral adenocarcinoma of the lung 3 cm or less in diameter. *Lung Cancer* 2000;29:179–88.
- [16] Goto K, Yokose T, Kodama T, Nagai K, Nishikawa Y, Ando M, et al. Detection of early invasion on the basis of basement membrane destruction in small adenocarcinomas of the lung and its clinical implications. *Mod Pathol* 2001;5:274–84.
- [17] Samuel A, Yousem. Peripheral squamous cell carcinoma of lung: patterns of growth with particular focus on airspace filling. *Human Pathol* 2009;40: 861–7.
- [18] Sakurai H, Maeshima A, Watanabe S, Suzuki K, Tsuchiya R, Maeshima AM, et al. Grade of stromal invasion in small adenocarcinoma of the lung. Histopathological minimal invasion and prognosis. *Am J Surg Pathol* 2004;28:198–206.
- [19] Ruffini E, Asioli S, Filosso PL, Lyberis P, Bruna MC, Macri L, et al. Clinical significance of tumor-infiltrating lymphocytes in lung neoplasms. *Ann Thorac Surg* 2009;87:371–2.
- [20] Saijo T, Ishii G, Nagai K, Funai K, Nitadori J, Tsuta K, et al. Differences in clinicopathological and biological features between central-type and peripheral-type squamous cell carcinoma of the lung. *Lung Cancer* 2006;52:37–45.



CASE REPORT

Open Access

Stromal micropapillary pattern predominant lung adenocarcinoma - a report of two cases

Miki Ohe^{1,2,4*}, Tomoyuki Yokose¹, Yuji Sakuma³, Sachie Osanai¹, Chikako Hasegawa¹, Kota Washimi¹, Kimitoshi Nawa², Tetsukan Woo², Rurika Hamanaka², Haruhiko Nakayama², Yoichi Kameda¹, Kouzo Yamada² and Takeshi Isobe⁴

Abstract

Generally, adenocarcinomas with micropapillary pattern, featuring small papillary tufts lacking a central fibrovascular core, are thought to have poor prognosis. This pattern has been described in various organs. However, tumor cells with micropapillary pattern of lung adenocarcinoma are more often seen to float within alveolar spaces (aerogenous micropapillary pattern, AMP) than in fibrotic stroma like other organs (stromal micropapillary pattern, SMP) and SMP predominant lung adenocarcinoma (SMPPLA) has not been well described yet. We presented two cases of SMPPLA which were found in the last four years. Both the cases showed more than 50% of SMP in the tumor area. The majority of the stromal micropapillary clusters expressed MUC1 and epithelial membrane antigen along the outer surface of cell membrane. On the other hand, connective tissues surrounding stromal micropapillary clusters showed no reactivity for epithelial markers (thyroid transcription factor-1 and cytokeratin) or endothelial marker (D2-40 and CD34). It means clusters of SMP do not exist within air space or lymphatic or vessel lumens. The tumors with SMP often presented lymphatic permeation and vessel invasion, and intriguingly, one of the two cases showed metastasis to the mediastinal lymph node. Additionally, both the cases showed *EGFR* point mutations of exon 21. These results suggest that SMPPLA might be associated with poor prognosis and effective for *EGFR* tyrosine kinase inhibitors.

Keywords: lung adenocarcinoma, micropapillary subtype, stromal micropapillary pattern, aerogeneous micropapillary pattern

Background

A new lung adenocarcinoma classification has been proposed by the International Association for the Study of Lung Cancer, American Thoracic Society and European Respiratory Society (IASLC/ATS/ERS). In this classification, the micropapillary subtype of lung adenocarcinoma (MSLA) was recommended as a newly added subtype of lung adenocarcinoma to lepidic, acinar, papillary, and solid subtypes defined in the 2004 World Health Organization (WHO) classification [1,2]. Generally, the micropapillary pattern is defined as tumor cells growing in papillary tufts, which lack fibrovascular cores surrounded by lacunar spaces and has been reported to be associated with a high incidence of nodal metastasis and poor prognosis [3-6]. This pattern has been described in various organs such as

breast [7,8], urinary bladder [9,10], ovary [11,12], salivary gland [13], and is known to behave aggressively. In other organs than the lung, this pattern was observed mainly in stroma as invasive components (stromal micropapillary pattern: SMP) [7-19]; however in lung, MSLA is widely recognized as floating tumor cells within alveolar spaces (aerogenous micropapillary pattern: AMP) [3,4].

We examined whether SMP predominant subtypes were present in lung adenocarcinoma. During the period from February 2007 to December 2010, 559 patients with lung adenocarcinoma were consecutively treated by surgical resection at the Kanagawa Cancer Center, Kanagawa, Japan, and we found only two cases of SMP predominant lung adenocarcinoma (SMPPLA) (0.36%). We reported the cases of SMPPLA and attempted to describe the clinicopathological features.

* Correspondence: ohem@kcch.jp

¹Department of Pathology, Kanagawa Cancer Center, Kanagawa, Japan

Full list of author information is available at the end of the article