

Table 4 Clinicopathologic characteristics of adenoma lesions detected during primary and secondary examinations

Characteristic	Primary		Secondary		<i>p</i>
	NBI	WLI	WLI	NBI	
Adenoma lesions	306 (78.7)	310 (72.2)	83 (21.3)	119 (27.8)	0.03
Location					
Cecum	57 (90.5)	49 (89.1)	6 (9.5)	6 (10.9)	0.80
Ascending colon	120 (78.9)	129 (71.7)	32 (21.1)	51 (28.3)	0.13
Transverse colon	129 (74.1)	132 (68.0)	45 (25.9)	62 (32.0)	0.20
Morphology					
Polypoid	136 (89.5)	139 (78.1)	16 (10.5)	39 (21.9)	0.006
Ip	8 (88.9)	12 (100)	1 (11.1)	0	0.24
Isp	13 (100)	15 (83.3)	0	3 (16.7)	0.12
Is	115 (88.5)	112 (75.7)	15 (11.5)	36 (24.3)	0.006
Flat and depressed	170 (72.0)	171 (68.6)	67 (28.0)	80 (31.4)	0.42
IIa	167 (71.4)	167 (68.2)	67 (28.6)	78 (31.8)	0.45
IIa + IIc	3 (100)	3 (75.0)	0	1 (25.0)	0.35
IIc	0	1 (50.0)	0	1 (50.0)	–
Size (mm)					
1–4	148 (74.4)	154 (65.3)	51 (25.6)	82 (34.7)	0.04
5–9	108 (79.4)	109 (76.8)	28 (20.6)	33 (23.2)	0.59
≥10	50 (92.6)	47 (92.2)	4 (7.4)	4 (7.8)	0.93
Histopathological findings					
TSA	7 (77.8)	12 (85.7)	2 (22.2)	2 (14.3)	0.62
LGD	277 (77.7)	283 (70.9)	80 (22.3)	116 (29.1)	0.04
LGD with villous	2 (100)	0	0	0	–
HGD	12 (93.3)	7 (92.9)	1 (6.7)	1 (7.1)	0.72
HGD with villous	2 (100)	6 (100)	0	0	–
Invasive cancer	6 (100)	2 (100)	0	0	–

Data represent the number of lesions (%)

NBI narrow band imaging, WLI white light imaging, TSA traditional serrated adenoma, LGD adenoma with low-grade dysplasia, HGD adenoma with high-grade dysplasia

these problems are usually solved by using high-definition colonoscopy. In addition, the wider colorectal lumen than the esophageal lumen in NBI is considered to be a reason for the screen darkness. An NBI setting different from that used in esophageal observation by high-definition colonoscopy is, therefore, indispensable for polyp detection in the colon and rectum. Uraoka et al. [25, 26] have reported that the A-5 image setting of the surface structure enhancement function, together with the level 3 adaptive index of hemoglobin color enhancement function, seem to be the most suitable settings for the detection of colorectal adenomas. In accordance with these findings, we used only high-definition colonoscopy to compare WLI with NBI and we applied surface structure enhancement level A-5 and adaptive index of hemoglobin color enhancement level 3 for NBI.

Our study results did not show a significant difference between NBI and WLI in the primary outcome measure but results were significantly different in the secondary outcome measure. Specifically, we found no significant difference in the adenoma detection rates by primary NBI versus primary WLI. We consider these results reliable

because there was no significant difference in the bowel preparation results or total observation times between the groups. Only expert colonoscopists performed the procedures in this study; therefore, it is necessary to clarify the usefulness of NBI for adenoma detection by all colonoscopists, including novices, in the future. Further, the detection rates of adenoma lesions by primary NBI and WLI were 78.7 and 72.2 % when we considered the detection rate of adenoma lesions by both primary and secondary examinations to be 100 %. In other words, the adenoma miss rates by primary NBI and WLI were 21.3 and 27.8 % ($p = 0.03$). The higher miss rate of WLI is similar to that reported previously. [5–7] Furthermore, Kaltenbach et al. [20] reported that NBI did not improve the colorectal adenoma miss rate compared to WLI in a randomized controlled trial using tandem colonoscopy (NBI–WLI vs. WLI–WLI; 12.6 vs. 12.1 %, respectively). Their adenoma miss rate using WLI was lower than the adenoma miss rate of 10–30 % reported in other studies. However, when we compared our results with their findings, it was evident that our miss rate was high. The differences between the study of Kaltenbach et al. and our




Fig. 2 **a** A flat elevated lesion was detected in the transverse colon. The size of the lesion was 7 mm in diameter. The final histopathological diagnosis was adenoma with low-grade dysplasia (LGD). **a** The polyp was missed by WLI; **b** the same lesion was identified by

subsequent NBI. **c, d** A depressed lesion was detected in the transverse colon. The size of the lesion was 5 mm in diameter. The final histopathological diagnosis was LGD. **c** The polyp was missed by WLI; **d** the same lesion was identified by subsequent NBI

study are the use of LUCERA versus EXERA II and the single-center versus multicenter design. Furthermore, we believe that a difference in the resolution of NBI and WLI may have influenced the results, because we were able to detect small lesions with NBI.

In line with previous reports [16, 17], we found that significantly higher numbers of small lesions (<5 mm) and/or LGD lesions were detected by NBI than by WLI [16, 17]. Further, nearly all the adenoma lesions we detected were flat elevated or polypoid in shape, and two were depressed. Depressed lesions are considered to have a higher malignant potential than polypoid ones of similar size [31–34]. The superiority of NBI over WLI in the detection of depressed lesions was not proven in the present study; however, we believe NBI is a promising modality for detecting small neoplastic lesions. The advantage of NBI endoscopy is simply to get the NBI view when we use a one-touch electrical button and to avoid indigo carmine dye-spraying. In addition, we can diagnose a lesion at the same time as it is detected. In the colorectal region, NBI is useful for differentiating non-neoplastic from neoplastic

lesions, and magnifying NBI is effective for determining the depth of invasion in early neoplasms [35–37].

In the present study, we could not evaluate serrated lesions because the pathological diagnosis of serrated lesions (particularly, sessile serrated adenoma) is not yet unified among Japanese pathologists. The number of TSAs detected by primary NBI and WLI were 7 and 12, respectively, and those identified by secondary WLI and NBI were 2 and 2, respectively. The miss rates of primary NBI and WLI for these lesions were not significantly different (22.2 vs. 14.3 %; $p = 0.62$).

This study has several limitations. First, the procedures were conducted only in the right colon, because Uraoka et al. [17] reported higher adenoma detection rates with NBI in the right colon, and a higher adenoma miss rate has been reported in the right colon than in the left colon [5]. Further, because complete back-to-back colonoscopy is sometimes uncomfortable for patients without sedation, we defined the region from the cecum to the splenic flexure as the target area in our study. Another limitation is that WLI was used for colonoscopy insertion in both the study groups, which could have

influenced the NBI results if some lesions were identified during insertion. However, we used the same imaging condition and study design for tandem colonoscopy in both the groups. Moreover, the detected lesions were removed endoscopically using WLI, because of the darkness of the screen with NBI. We cannot entirely exclude the possibility that some switches of endoscopic treatment influenced the detection rates in both the groups. Endoscopic treatment is, however, usually focused on the small area in which the polyp is located and the examiner likely concentrates on the endoscopic treatment rather than on the detection of additional lesions. Another limitation is that both the NBI and WLI examinations were performed by the same endoscopist. There may be investigator bias. However, we believe this does not substantially influence the results, because this was a multicenter trial and the endoscopists performed procedures for both the NBI–WLI and WLI–NBI groups. Another limitation is that, in the distinction between neoplasia and non-neoplasia, NBI may be expected to have a small advantage. In the present study, expert colonoscopists examined the lesions; such experts are able to distinguish between neoplasia and non-neoplasia using WLI as well as NBI. Furthermore, because all detected non-neoplastic lesions were removed or biopsied, the difference between NBI and WLI with respect to the distinction between neoplasia and non-neoplasia would not have affected the overall adenoma detection rate. Therefore, we believe that the primary endpoint of the adenoma detection rate was not affected by this discrepancy. Finally, the current NBI systems have problems such as darkness and noise. Even if these systems are used with high-definition colonoscopy, the brightness of the screen is still not sufficient. Further, NBI system-related improvement is necessary for enhanced adenoma detection.

In conclusion, NBI does not have a higher adenoma detection rate during primary colonoscopy than WLI, but it has a lower adenoma miss rate in the proximal colon by tandem colonoscopy. NBI can be expected to represent a suitable modality for screening colonoscopy, because the miss rate is low.

Acknowledgments This paper was presented at the 18th United European Gastroenterology Week (UEGW), October 23–27, 2010, in Barcelona, Spain. This study was supported in part by Grant No. H21-009 from the Ministry of Health, Labour and Welfare of Japan.

Conflict of interest The authors declare that they have no conflict of interest.

References

1. Winawer SJ, Zauber AG, O'Brien MJ, Ho MN, Gottlieb L, Sternberg SS, et al. Randomized comparison of surveillance intervals after colonoscopic removal of newly diagnosed adenomatous polyps. The National Polyp Study Workgroup. *N Engl J Med.* 1993;328:901–6.
2. Winawer SJ, Zauber AG, Ho MN, O'Brien MJ, Gottlieb LS, Sternberg SS, et al. Prevention of colorectal cancer by colonoscopic polypectomy. The National Polyp Study Workgroup. *N Engl J Med.* 1993;329:1977–81.
3. Levin B, Lieberman DA, McFarland B, Andrews KS, Brooks D, Bond J, et al. Screening and surveillance for the early detection of colorectal cancer and adenomatous polyps, 2008: a joint guideline from the American Cancer Society, the US Multi-Society Task Force on Colorectal Cancer, and the American College of Radiology. *Gastroenterology.* 2008;134:1570–95.
4. Kaminski MF, Regula J, Kraszewska E, Polkowski M, Wojciechowska U, Didkowska J, et al. Quality indicators for colonoscopy and the risk of interval cancer. *N Engl J Med.* 2010;362:1795–803.
5. Rex DK, Cutler CS, Lemmel GT, Rahmani EY, Clark DW, Helper DJ, et al. Colonoscopic miss rates of adenomas determined by back-to-back colonoscopies. *Gastroenterology.* 1997;112:24–8.
6. Postic G, Lewin D, Bickerstaff C, Wallace MB. Colonoscopic miss rates determined by direct comparison of colonoscopy with colon resection specimens. *Am J Gastroenterol.* 2002;97:3182–5.
7. Heresbach D, Barrioz T, Lapalus MG, Coumaros D, Bauret P, Potier P, et al. Miss rate for colorectal neoplastic polyps: a prospective multicenter study of back-to-back video colonoscopies. *Endoscopy.* 2008;40:284–90.
8. Brooker JC, Saunders BP, Shah SG, Thapar CJ, Thomas HJ, Atkin WS, et al. Total colonic dye-spray increases the detection of diminutive adenomas during routine colonoscopy: a randomized controlled trial. *Gastrointest Endosc.* 2002;56:333–8.
9. Hurlstone DP, Cross SS, Slater R, Sanders DS, Brown S. Detecting diminutive colorectal lesions at colonoscopy: a randomised controlled trial of pan-colonic versus targeted chromoscopy. *Gut.* 2004;53:376–80.
10. Rex DK, Chadalawada V, Helper DJ. Wide angle colonoscopy with a prototype instrument: impact on miss rates and efficiency as determined by back-to-back colonoscopies. *Am J Gastroenterol.* 2003;98:2000–5.
11. Deenadayalu VP, Chadalawada V, Rex DK. 170 degrees wide-angle colonoscope: effect on efficiency and miss rates. *Am J Gastroenterol.* 2004;99:2138–42.
12. Triadafilopoulos G, Watts HD, Higgins J, Van Dam J. A novel retrograde-viewing auxiliary imaging device (Third Eye Retroscope) improves the detection of simulated polyps in anatomic models of the colon. *Gastrointest Endosc.* 2007;65:139–44.
13. DeMarco DC, Odstrcil E, Lara LF, Bass D, Herdman C, Kinney T, et al. Impact of experience with a retrograde-viewing device on adenoma detection rates and withdrawal times during colonoscopy: the Third Eye Retroscope study group. *Gastrointest Endosc.* 2010;71:542–50.
14. Matsushita M, Hajiro K, Okazaki K, Takakuwa H, Tominaga M. Efficacy of total colonoscopy with a transparent cap in comparison with colonoscopy without the cap. *Endoscopy.* 1998;30:444–7.
15. East JE, Suzuki N, Stavrinidis M, Guenther T, Thomas HJ, Saunders BP. Narrow band imaging for colonoscopic surveillance in hereditary non-polyposis colorectal cancer. *Gut.* 2008;57:65–70.
16. Inoue T, Murano M, Murano N, Kuramoto T, Kawakami K, Abe Y, et al. Comparative study of conventional colonoscopy and pan-colonic narrow-band imaging system in the detection of neoplastic colonic polyps: a randomized, controlled trial. *J Gastroenterol.* 2008;43:45–50.
17. Uraoka T, Saito Y, Matsuda T, Sano Y, Ikehara H, Mashimo Y, et al. Detectability of colorectal neoplastic lesions using a

- narrow-band imaging system: a pilot study. *J Gastroenterol Hepatol.* 2008;23:1810–5.
18. Rex DK, Helbig CC. High yields of small and flat adenomas with high-definition colonoscopes using either white light or narrow band imaging. *Gastroenterology.* 2007;133:42–7.
 19. Adler A, Pohl H, Papanikolaou IS, Abou-Rebyeh H, Schachschal G, Veltzke-Schlieker W, et al. A prospective randomised study on narrow-band imaging versus conventional colonoscopy for adenoma detection: does narrow-band imaging induce a learning effect? *Gut.* 2008;57:59–64.
 20. Kaltenbach T, Friedland S, Soetikno R. A randomised tandem colonoscopy trial of narrow band imaging versus white light examination to compare neoplasia miss rates. *Gut.* 2008;57:1406–12.
 21. Adler A, Aschenbeck J, Yenerim T, Mayr M, Aminimalai A, Drossel R, et al. Narrow-band versus white-light high definition television endoscopic imaging for screening colonoscopy: a prospective randomized trial. *Gastroenterology.* 2009;136:410–6.
 22. Gono K, Yamazaki K, Doguchi N, et al. Endoscopic observation of tissue by narrow band illumination. *Opt Rev.* 2003;10:211–5.
 23. Gono K, Obi T, Yamaguchi M, Ohyama N, Machida H, Sano Y, et al. Appearance of enhanced tissue features in narrow-band endoscopic imaging. *J Biomed Opt.* 2004;9:568–77.
 24. Muto M, Minashi K, Yano T, Saito Y, Oda I, Nonaka S, et al. Early detection of superficial squamous cell carcinoma in the head and neck region and esophagus by narrow band imaging: a multicenter randomized controlled trial. *J Clin Oncol.* 2010;28:1566–72.
 25. Uraoka T, Sano Y, Saito Y, et al. Narrow-band imaging for improving colorectal adenoma detection: appropriate system function settings are required. *Gut.* 2009;58:604–5.
 26. Uraoka T, Higashi R, Saito Y, et al. Impact of narrow-band imaging in screening colonoscopy. *Dig Endosc.* 2010;22(Suppl 1):S54–6.
 27. Chiu HM, Lin JT, Wang HP, Lee YC, Wu MS. The impact of colon preparation timing on colonoscopic detection of colorectal neoplasms—a prospective endoscopist-blinded randomized trial. *Am J Gastroenterol.* 2006;101:2719–25.
 28. East JE, Suzuki N, Arebi N, Bassett P, Saunders BP. Position changes improve visibility during colonoscopy withdrawal: a randomized, blinded, crossover trial. *Gastrointest Endosc.* 2007;65:263–9.
 29. The Paris endoscopic classification of superficial neoplastic lesions: esophagus, stomach, and colon: November 30 to December 1, 2002. *Gastrointest Endosc.* 2003;58:S3–43.
 30. Hamilton SR, Aaltonen LA, editors. World Health Organization classification of tumours: pathology and genetics of tumours of the digestive system. Lyon: IARC Press; 2000. p. 104–19.
 31. Kudo S, Kashida H, Tamura T, et al. Colonoscopic diagnosis and management of nonpolypoid early colorectal cancer. *World J Surg.* 2000;24:1081–90.
 32. Saitoh Y, Waxman I, West AB, Popnikolov NK, Gatalica Z, Watari J, et al. Prevalence and distinctive biologic features of flat colorectal adenomas in a North American population. *Gastroenterology.* 2001;120:1657–65.
 33. Tsuda S, Veress B, Toth E, Fork FT. Flat and depressed colorectal tumours in a southern Swedish population: a prospective chromoendoscopic and histopathological study. *Gut.* 2002;51:550–5.
 34. Soetikno R, Friedland S, Kaltenbach T, Chayama K, Tanaka S. Nonpolypoid (flat and depressed) colorectal neoplasms. *Gastroenterology* 2006;130:566–76 (quiz 588–9).
 35. Sano Y, Ikematsu H, Fu KI, Emura F, Katagiri A, Horimatsu T, et al. Meshed capillary vessels by use of narrow-band imaging for differential diagnosis of small colorectal polyps. *Gastrointest Endosc.* 2009;69:278–83.
 36. Chiu HM, Chang CY, Chen CC, Lee YC, Wu MS, Lin JT, et al. A prospective comparative study of narrow-band imaging, chromoendoscopy, and conventional colonoscopy in the diagnosis of colorectal neoplasia. *Gut.* 2007;56:373–9.
 37. Ikematsu H, Matsuda T, Emura F, Saito Y, Uraoka T, Fu KI, et al. Efficacy of capillary pattern type IIIA/IIIB by magnifying narrow band imaging for estimating depth of invasion of early colorectal neoplasms. *BMC Gastroenterol.* 2010;10:33.

The use of computed tomographic colonography in predicting the difficulty of endoscopic treatment for large protruding neoplasms

Taku Sakamoto · Mototaka Miyake ·
Takeshi Nakajima · Takahisa Matsuda ·
Hirokazu Taniguchi · Yutaka Saito · Gen Iinuma

Accepted: 1 December 2011 / Published online: 14 December 2011
© Springer-Verlag 2011

Dear Editor:

Submucosal fibrosis is considered a major hurdle in endoscopic submucosal dissection (ESD), which is an effective technique for treating colorectal neoplasms. Although it is relatively easy to predict the difficulty of endoscopic resection for laterally spreading tumors (LST), it is more difficult in the case of large protruding lesions (Paris type 0-Is, >20 mm in diameter) such as villous tumors, which are often eventually removed by multiple piecemeal resection. However, considering the high incidence of recurrent or residual tumors and the difficulty of precise histopathological evaluation, piecemeal endoscopic mucosal resection (EMR) should be avoided. Instead, when endoscopic resection is predicted to be difficult, laparoscopic assisted colectomy is a viable treatment option; therefore, the objective evaluation of fibrotic changes in the submucosal layer is important. Although this evaluation may be accomplished using endoscopic miniprobe ultrasonography (mEUS), which is effective in assessing the depth of invasion in both gastric cancer and colorectal neoplasms as well as that of submucosal fibrosis for flat or depressed lesions, its limited depth of penetration is a disadvantage in assessing protruding lesions. Moreover, the observation of lesions located on

the oral side of folds using mEUS is also considered difficult, and a different method may be more suitable for identifying these cases.

Computed tomographic colonography (CTC) is currently an established technique for colorectal imaging that allows the evaluation of both endoluminal and transluminal features and has good diagnostic performance in T staging of colorectal cancer. Using CTC, images can be reconstructed in almost any plane and can be used to create three-dimensional images while maintaining diagnostic image resolution and without influencing the location of the lesion. It can also be used to evaluate both lesion morphology and intratumoral features, including those of protruding lesions and lesions located on the oral side of folds. Moreover, several studies have shown that the degree of fibrosis contributes to an enhancement pattern in contrast-enhanced CTC (CE-CTC). Thus, CE-CTC may potentially be used for evaluating the shape and morphological changes of the bowel wall, including the degree of fibrotic changes in the submucosal layer. In this letter, we describe the use of CTC in the identification of patients with intramucosal protruded-type neoplasm predicted to present endoscopic difficulties.

All patients who had undergone preoperative colonoscopy and same day CE-CTC at our institution from January 2006 to December 2008 were identified through retrospective analysis. After pretreatment colonoscopy and standard preparation, patients underwent CE-CTC examination for staging using a 64-multidetector row CT scanner (Aquilion; Toshiba Medical Systems, Tokyo, Japan). The scan range was from the abdomen to the pelvis, with the following parameters: 120 kV; 200–400 mA with automatic exposure control; 64 rows×0.5 mm collimation; and helical pitch, 53 (pitch factor, 0.828). Anticholinergic drugs were injected intravenously immediately before each examination, and gas insufflation was performed via the anus with an

T. Sakamoto (✉) · T. Nakajima · T. Matsuda · Y. Saito
Endoscopy Division, National Cancer Center Hospital,
5-1-1 Tsukiji,
Chuo-ku, Tokyo 104-0045, Japan
e-mail: tasakamo@ncc.go.jp

M. Miyake · G. Iinuma
Department of Diagnostic Radiology, National Cancer Center Hospital,
Chuo-ku, Tokyo, Japan

H. Taniguchi
Pathology Division, National Cancer Center Hospital,
Chuo-ku, Tokyo, Japan

automated CO₂ insufflator (Protocol; E-Z-EM, New York). For TNM staging, all patients were administered a total of 150 mL of contrast medium [Omnipaque (300 mg/mL); Daiichi-Sankyo Pharmaceuticals, Tokyo, Japan] intravenously with an autoinjector at a rate of 3.0 mL/s. The scan delays were set at 50 s (early phase) and 90 s (delayed phase) after the injection of the contrast medium. CE-CTC images were acquired for each patient in prone and supine positions and reviewed by an experienced radiologist for identifying “bundle-like low-density areas (BLDA)” in the submucosa, which were characterized as a lower density area than that of protruding intramucosal neoplasms in the early phase.

Patients with protruding intramucosal neoplasms larger than 20 mm in diameter were included in our study, whereas those with LSTs as per the endoscopic data were excluded. The 11 patients that met these criteria were divided into two groups according to the outcome of the endoscopic treatment: 6 cases of patients were included in group A, which comprised the difficult resections that had a piecemeal resected specimen with more than 10 pieces and/or a procedure time longer than 180 min, while group B included the remaining 5 cases. En bloc resection was achieved in 36% cases (4 out of 11), and the median duration of the procedure time was 95 min (interquartile range, 80–200 min). In group A, the median size of the lesions was 52.5 mm (range, 40–60), with 83% lesions located in the colon and 17% in the rectum. In group B, the lesions had a median size of 40 mm (range, 30–50 mm), and 80% was located in the rectum. No significant differences in the size or location of lesions were observed between the two groups ($P=0.113$ and $P=0.067$, respectively); however, the duration of the procedure was significantly longer in group A than group B (median time, 200 vs. 70 min; $P=0.008$). The rate of positive findings for BLDAs was 83% (five out of six) in group A and 20% (one out of five) in group B. Histopathological examination showed that the presence of BLDA corresponded to the presence of submucosal fibrosis.

Regarding clinical outcomes, en bloc resection was achieved in only 36% of the cases in this study. Although ESD was planned in nine cases, the procedure was abandoned and converted to piecemeal EMR in 56% (five out of nine) of

these cases because of technical difficulties associated with submucosal fibrosis. As a previous Japanese multicenter study has reported that the en bloc resection rate of ESD for treatment of colorectal neoplasms was 88%, it appears that endoscopic treatment for large protruding lesions is more difficult than that for other colorectal neoplasms. Additionally, confounding factors other than submucosal fibrosis, such as location and lesion size, may have potentially contributed to the lack of success of endoscopic treatment in our study. In particular, group A included a greater number of colonic lesions; colonic lesion ESD is known to be technically more difficult to treat than rectal ESD. However, as there was no significant difference in lesion size between groups, submucosal fibrosis seemed to be the most important factor determining the difficulty of endoscopic treatment.

Our study has certain limitations. First, we reviewed only intramucosal neoplastic lesions that were treated endoscopically. Although the presence of submucosal fibrosis could be confirmed, precise correspondence between histological and CTC findings may be difficult to conclude in cases where patients require piecemeal resection. Comparing histological and CTC findings in surgically resected intramucosal neoplasms would enhance the accuracy and acceptance of preoperative BLDA assessment by CTC. Second, only a single experienced radiologist evaluated the presence of BLDA in our study, and our findings require validation by studies involving multiple radiologists. Third, our study was a pilot study with a small sample size; a prospective multicenter trial would provide further clarity regarding the utility of CTC in predicting difficulties in endoscopic resection. Despite these limitations, our study suggests that the presence of BLDA corresponds well with the presence of fibrosis in the submucosal layer and that CTC may be an effective tool for preoperatively estimating the difficulty of the endoscopic resection of large protruding tumors in the colon.

Financial disclosures No funding was provided for this study, and the authors have no relevant financial relationships to report.

GASTROENTEROLOGY

Time saving with narrow-band imaging for distinguishing between neoplastic and non-neoplastic small colorectal lesions

Taku Sakamoto, Takahisa Matsuda, Takaya Aoki, Takeshi Nakajima and Yutaka Saito

Endoscopy Division, National Cancer Center Hospital, Tokyo, Japan

Key words

colonoscopy, colorectal lesion, magnifying chromoendoscopy, narrow-band imaging.

Accepted for publication 30 June 2011.

Correspondence

Dr Taku Sakamoto, Endoscopy Division, National Cancer Center Hospital, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan. Email: tasakamo@ncc.go.jp

Abstract**Background and Aims:** For colonoscopic examinations, the narrow-band imaging (NBI) system is more convenient and timesaving than magnifying chromoendoscopy (MCE). However, the time-saving aspects of NBI techniques have not been assessed. The present study compared interpretation times between NBI and MCE techniques in distinguishing between neoplastic and non-neoplastic small colorectal lesions.**Methods:** Between January and March 2010, 693 consecutive patients who underwent colonoscopy at the National Cancer Center Hospital, Tokyo, Japan, were enrolled. When the first lesion was detected by conventional white-light observation, the patient was randomly assigned to undergo a sequence of NBI and MCE observations (group A: NBI-MCE, group B: MCE-NBI). The time to diagnosis with each modality (NBI, from changing to NBI until diagnosis; MCE, from the start of indigo carmine solution spraying until diagnosis) was recorded by an independent observer. The sensitivity, specificity, and diagnostic accuracy of the first modality used in each group (NBI or MCE) were assessed by referring to the histopathological data.**Results:** Seventy-one patients with 137 lesions were randomized to group A, and 80 patients with 163 lesions to group B. The median interpretation times were 12 s (interquartile range [IQR]: 7–19 s) in group A, and 17 s (IQR: 12–24 s) in group B, the difference being significant ($P < 0.001$). No significant differences were observed between NBI and MCE in terms of sensitivity, specificity, and diagnostic accuracy.**Conclusions:** NBI reduces the interpretation times for distinguishing between neoplastic and non-neoplastic small lesions during colonoscopies, without loss of diagnostic accuracy.**Introduction**

The usefulness of narrow-band imaging (NBI) systems with magnification, for differentiating between neoplastic and non-neoplastic lesions in colonoscopy, has been commonly reported.^{1–9} Prior to the development of NBI techniques, pit pattern analysis using magnifying chromoendoscopy (MCE) was considered to be the most reliable method, not only for differentiating between neoplastic and non-neoplastic lesions, but also for evaluating the invasion depth of early colorectal cancers.^{10–15} Most previous studies showed high diagnostic accuracies of pit pattern analysis with MCE (> 90%), especially in the differentiation of neoplastic lesions from non-neoplastic ones.^{10–12} Pit pattern analysis is, therefore, considered a gold standard for accurate endoscopic diagnosis of colorectal lesions. However, some studies have suggested that analysis of microcapillary vessel patterns of lesions using NBI with magnification shows almost the same diagnostic accuracy as pit pattern analysis.^{1–6} Moreover, NBI is reputed to be convenient and time saving compared to MCE, and thus the NBI system has

received increasing attention. Previous studies on the efficacy of NBI tended to emphasize only the merits of NBI's diagnostic capabilities, and did not precisely assess the duration of diagnostic interpretation times.^{1–9} The primary aim of this study was to compare the interpretation times between NBI and MCE in distinguishing between neoplastic and non-neoplastic lesions. Furthermore, we attempted to determine diagnostic accuracy and the levels of agreement between NBI and MCE techniques in the analysis of identical lesions.

Methods**Patients**

A total of 693 consecutive patients who underwent total colonoscopy at the National Cancer Center Hospital (NCCH), Tokyo, Japan, between January and March 2010, were considered eligible for enrolment in the study. Patients with familial adenomatous polyposis, inflammatory bowel disease, advanced colorectal




Figure 1 Flow chart of the colonoscopic diagnoses used in this study. MCV, microcapillary vessels; NBI, narrow-band imaging.

cancer, and poor colonoscopy preparation were excluded. The study was conducted prospectively, and the study protocol was approved by the NCCH institutional review board. Written, informed consent for diagnosis and treatment was obtained from all patients prior to the procedures. Lesions were detected in 151 patients and selected for study.

Colonoscopic examination

Patients prepared for colonoscopy by ingesting 2–3 L of polyethylene glycol-electrolyte solution on the morning of the examination day. Scopolamine butylbromide (10 mg) was administered intravenously to avoid bowel movement prior to examination in patients with no contraindication for this agent.

All examinations were performed using magnifying colonoscopes (CF-H260AZI or PCF-Q240ZI; Olympus Optical, Tokyo, Japan) and a standard videoendoscopic system (EVIS LUCERA system; Olympus Optical, Japan) with two light sources: one for the standard optical broadband filter, and the other for the NBI system (surface structure enhancement function, A-5 setting; and adaptive IHb color enhancement function, level 3 setting). During the procedure, whenever a lesion was detected by standard colonoscopy, analyses of capillary-vessel patterns with NBI magnification and analyses of pit patterns using MCE were performed. Patients were randomly assigned to first undergo observation with either NBI or MCE (group A: NBI–MCE, group B: MCE–NBI) (Fig. 1). For pit pattern analyses, indigo carmine (0.4%) was sprayed directly on the mucosal surface after washing with proteinase to remove any overlying mucous. The mucosal crypt patterns observed with chromoendoscopy were assessed using Kudo's classification.^{16–18} In the case of NBI, four different microvascular architecture patterns were identified according to Sano's classification.^{1,5,6,19}

After endoscopic evaluation, all detected lesions, unless contraindicated for endoscopic resection, were treated by biopsy, hot biopsy, snare polypectomy, endoscopic mucosal resection (EMR), or endoscopic submucosal dissection (ESD). All 11 endoscopists involved in the study had performed more than 200 colonoscopies with magnification, and had worked for more than 6 months at the NCCH. All endoscopists were familiar with NBI and MCE images, and received lectures throughout the course of the study during a weekly case conference meeting.

Assessments

The time to diagnosis with each modality (NBI, from changing to NBI until diagnosis; MCE, from starting the indigo carmine solution spraying until diagnosis) was recorded by an independent observer (Fig. 1). The sensitivity, specificity, and diagnostic accuracy of the first observation modality (NBI or MCE) in each group were assessed by referring to histopathological data.

Statistical analysis

Interpretation times of the two modalities were described by median values and interquartile ranges (IQR). Group medians were compared using the Wilcoxon rank sum test. Diagnostic performances for differentiating between neoplastic and non-neoplastic lesions were compared using the χ^2 -test. All *P*-values were determined by two-tailed tests, and values of *P* < 0.05 were considered statistically significant. All calculations were performed using STATA software, version 10.0 (StataCorp, College Station, TX, USA).

Table 1 Clinical characteristics of patients and lesions

	Group A (n = 137/71 patients)	Group B (n = 163/80 patients)	P-value [†]
Age (mean ± SD) years	65.6 ± 10.7	65.1 ± 8.6	0.373
Gender (male/female)	50/21	54/26	0.699
Median size (IQR)	5 (3–7)mm	5 (3–7)mm	0.887
No. cases by location (%)			0.765
Right colon	70 (51)	83 (51)	
Left colon	42 (38)	60 (37)	
Rectum	15 (11)	20 (12)	
No. cases by macroscopic type (%)			0.386
Is/lsp, lp	43 (31)	63 (38)	
Is + Ila	2 (2)	1 (1)	
Ila	92 (67)	98 (60)	
Ila + IIc	0 (0)	1 (1)	

[†]P-values represent the significance of the differences between the two groups. IQR, interquartile range; SD, standard deviation.

Table 2 Histopathological results

No. cases of neoplastic lesions (%)	
Tubular adenoma	216 (79)
Tubulo-villous adenoma	2 (1)
Well-differentiated adenocarcinoma	15 (6)
Sessile serrated adenoma	6 (2)
No. cases of non-neoplastic lesions (%)	
Hyperplastic polyp	28 (11)
Inflammatory polyp	3 (1)

Table 3 Diagnostic performances of magnifying chromoendoscopy (MCE) and narrow-band imaging (NBI), and the significance of the differences between them

	MCE	NBI	P-value
Sensitivity	93.7% (224/239)	94.1% (225/239)	0.863
Specificity	80.6% (25/31)	74.2% (23/31)	0.415
Accuracy	92.2% (249/270)	91.9% (248/270)	0.857

No. cases are in parentheses.

Results

Seventy-one patients with 137 lesions were randomized to group A, and 80 patients with 163 lesions to group B. The clinical features of the lesions detected in each group are presented in Table 1. There were no significant differences in age, sex, lesion size, location, and macroscopic type between the two groups.

In group A, the median interpretation times were 12 s (IQR: 7–19 s) at the first diagnosis by NBI (time A), and 17 s (IQR: 10–29.5 s) at the second diagnosis by MCE. In group B, the median interpretation times were 17 s (IQR: 12–24 s) at the first diagnosis by MCE (time B), and 11 s (IQR: 8–17 s) at the second diagnosis by NBI. Comparison of times A and B values indicated a significant difference in the interpretation time between NBI and MCE ($P < 0.001$).

Of the 270 colorectal lesions for which the histopathological findings were confirmed, 31 were hyperplastic or inflammatory polyps, six were sessile serrated adenomas (SSA), 218 were adenomas, and 15 were adenocarcinomas (Table 2). Interpretation of neoplastic lesions (adenoma, adenocarcinoma, and SSA) by NBI and MCE revealed sensitivities of 94.1% and 93.7%, respectively, specificities of 74.2% and 80.6% respectively, and diagnostic accuracies of 91.9% and 92.2%, respectively; none of these parameters showed significant differences between MCE and NBI (Table 3). Almost perfect intraobserver agreement was observed for the diagnoses of neoplastic lesions for each modality ($\kappa = 0.92$).

Discussion

The present study demonstrated that NBI can reduce the interpretation times for distinguishing between neoplastic and non-neoplastic lesions during colonoscopies. One of the advantages of using the NBI system is its relative simplicity, requiring only a single touch of a button to immediately change to the NBI mode from the conventional white-light observation mode. Moreover, the time required to differentiate between neoplastic and non-neoplastic lesions using the NBI system was statistically 4 or 5 s less than that for chromoendoscopic techniques. The extended interpretation time for MCE techniques was due to several factors, including: (i) the time required for spraying and suctioning indigo carmine solution; (ii) the time required to locate lesions in deep fold areas where fluid is easily retained, such as in the ascending colon; and (iii) the extra time required in cases where the patient has strong bowel movement or multiple lesions. The time saved using NBI was only 4 or 5 s per lesion, and the clinical benefit was rather small. However, the time saved would be particularly useful for the examination of patients with multiple lesions or with relatively poor preparation. In addition, time-saving modalities are important in cases where the lesions are located in curvy areas, such as the hepatic and splenic flexure, or when it is difficult to maneuver the colonoscope due to paradoxical movement.

Regarding the diagnostic performance for differentiating between neoplastic and non-neoplastic lesions, the NBI and MCE techniques showed almost the same values for sensitivity, specificity, and diagnostic accuracy, as determined in previous studies.^{1–6,10–12} Our data were obtained in a prospectively-designed study, and its applicability is limited to the roles of MCE and NBI in differentiating between neoplastic and non-neoplastic lesions.




Figure 2 Endoscopic images of a case of sessile serrated adenoma/polyp. Regular capillary vessel surrounding round crypts was identified with magnifying narrow-band imaging. Slight dilation of crypts or inhomogeneous pits was recognized with magnifying chromoendoscopy.

We should note that, despite its longer interpretation time, chromoendoscopy using indigo carmine dye spraying is not without value in endoscopic examinations. Treatment strategies for neoplastic lesions require detailed assessments of the configurations of lesions, for which MCE is well suited. For example, flat and depressed types of lesions, which characterize laterally-spreading tumors (LST), are likely precursors of advanced carcinoma; possible treatments are broadly divided into conventional EMR, piecemeal EMR, and ESD. In particular, LST non-granular lesions showing a very flat and smooth configuration with pseudopodia-like appearances larger than 20 mm in size are considered to be a definite indication for ESD because of the high risk of submucosal invasion;^{20–22} indigo carmine dye spraying is essential to clarify these features.

For diagnostic performance, the specificity of NBI is slightly lower than that of MCE, although the differences observed in this study were not statistically significant; this might be due to the presence of SSA cases in this study. Most SSA might be diagnosed as non-neoplastic lesions by NBI, but as neoplastic lesions by MCE. SSA are histologically characterized as follows: dilation of crypts, branching, presence of horizontal glands at the base, presence of mature mucinous cells at the base of crypts etc.^{23,24} Dilation of crypts or inhomogeneous pits reflecting histopathologically-evident atypia can be observed by MCE (Fig. 2). The shapes of crypts are similar to the hyperplastic polyps diagnosed on the basis of type II pit patterns (almost homogeneous stellar of papillary pits) by Kudo's classification; therefore, we can distinguish between SSA and hyperplastic polyps from the minor differences in pit pattern. However, some cases showed relatively dilated microcapillary vessels surrounding small crypts, rather than normal mucosa or typical small hyperplastic polyps characteristic of SSA. A clear definition suggesting the difference in capillary vessel pattern between SSA and hyperplastic polyps has not been provided yet, which resulted in non-neoplastic lesion diagnoses by NBI (Fig. 2). SSA thus need to be independently confirmed; the NBI findings for SSA might offer insights into a

subset of colorectal adenocarcinomas characteristic of neoplastic lesion, but more detailed study is needed to clarify the meaning of NBI findings.

There are several limitations to this study. First, this study was conducted at a single medical centre, where we usually apply Sano's classification to interpret microcapillary vessel patterns. However, other classification systems for NBI have been suggested in Japan, including the simultaneous evaluation of mucosal surface patterns and microcapillary vessels. In this approach, the NBI image settings might be slightly different than those in the present study, leading to slightly different results. Second, the endoscopists who performed the colonoscopies in this study were proficient in both NBI and MCE. Thus, the general efficacy of NBI requires revalidation studies, including those with general endoscopists. Third, all diagnoses were performed using magnifying endoscopy, but there might be some institutions where colonoscopies without magnification are performed in routine examinations. Therefore, the applicability of our results to endoscopic examinations without magnification is unclear.

In conclusion, NBI can reduce the interpretation time required for distinguishing between neoplastic and non-neoplastic lesions during colonoscopies without loss of diagnostic accuracy. Moreover, the differential diagnoses for a lesion by NBI are the same as those obtained by MCE. These results suggest that NBI can replace MCE as a diagnostic tool for assessing colorectal neoplasms.

References

- 1 Machida H, Sano Y, Yamamoto Y *et al.* Narrow band imaging for differential diagnosis of colorectal mucosal lesions: a pilot study. *Endoscopy* 2004; **36**: 1094–8.
- 2 Hirata M, Tanaka S, Oka S *et al.* Magnifying endoscopy with narrow band imaging for diagnosis of colorectal tumors. *Gastrointest. Endosc.* 2007; **65**: 988–95.

- 3 Chiu HM, Chang CY, Chen CC *et al.* A prospective comparative study of narrow-band imaging, chromoendoscopy, and conventional colonoscopy in the diagnosis of colorectal neoplasia. *Gut* 2007; **56**: 373–9.
- 4 Tischendorf JJ, Wasmuth HE, Koch A *et al.* Value of magnifying chromoendoscopy and narrow band imaging (NBI) in classifying colorectal polyps: a prospective controlled study. *Endoscopy* 2007; **39**: 1092–6.
- 5 Katagiri A, Fu KI, Sano Y *et al.* Narrow band imaging with magnifying colonoscopy as diagnostic tool for predicting histology of early colorectal neoplasia. *Aliment. Pharmacol. Ther.* 2008; **27**: 1269–74.
- 6 Sano Y, Ikematsu H, Fu KI *et al.* Meshed capillary vessels by use of narrow-band imaging for differential diagnosis of small colorectal polyps. *Gastrointest. Endosc.* 2009; **69**: 278–83.
- 7 Rastogi A, Bansal A, Wani S *et al.* Narrow-band imaging colonoscopy—a pilot feasibility study for the detection of polyps and correlation of surface patterns with polyp histologic diagnosis. *Gastrointest. Endosc.* 2008; **67**: 280–6.
- 8 Rastogi A, Pondugula K, Bansal A *et al.* Recognition of surface mucosal and vascular patterns of colon polyps by using narrow-band imaging: interobserver and intraobserver agreement and prediction of polyp histology. *Gastrointest. Endosc.* 2009; **69**: 716–22.
- 9 Tischendorf JJ, Schirin-Sokhan R, Streez K *et al.* Value of magnifying endoscopy in classifying colorectal polyps based on vascular pattern. *Endoscopy* 2010; **42**: 22–7.
- 10 Togashi K, Konishi F, Ishizuka T *et al.* Efficacy of magnifying endoscopy in the differential diagnosis of neoplastic and non-neoplastic polyps of the large bowel. *Dis. Colon. Rectum.* 1999; **42**: 1602–8.
- 11 Konishi K, Kaneko K, Kurahashi T *et al.* A comparison of magnifying and non magnifying colonoscopy for diagnosis of colorectal polyps: a prospective study. *Gastrointest. Endosc.* 2003; **57**: 48–53.
- 12 Fu KI, Sano Y, Kato S *et al.* Chromoendoscopy using indigo carmine dye spraying with magnifying observation is the most reliable method for differential diagnosis between non-neoplastic and neoplastic colorectal lesions: a prospective study. *Endoscopy* 2004; **36**: 1089–93.
- 13 Kanao H, Tanaka S, Oka S *et al.* Clinical significance of type V(I) pit pattern subclassification in determining the depth of invasion of colorectal neoplasms. *World J. Gastroenterol.* 2008; **14**: 211–17.
- 14 Onishi T, Tamura S, Kuratani Y *et al.* Evaluation of the depth score of type V pit patterns in crypt orifices of colorectal neoplastic lesions. *J. Gastroenterol.* 2008; **43**: 291–7.
- 15 Matsuda T, Fujii T, Saito Y *et al.* Efficacy of the invasive/non-invasive pattern by magnifying chromoendoscopy to estimate the depth of invasion of early colorectal neoplasms. *Am. J. Gastroenterol.* 2008; **103**: 2700–6.
- 16 Kudo S, Tamura S, Nakajima T *et al.* Diagnosis of colorectal tumorous lesions by magnifying endoscopy. *Gastrointest. Endosc.* 1996; **44**: 8–14.
- 17 Kudo S, Kashida H, Tamura T *et al.* Colonoscopic diagnosis and management of nonpolypoid early colorectal cancer. *World J. Surg.* 2000; **24**: 1081–90.
- 18 Kashida H, Kudo S. Magnifying colonoscopy, early colorectal cancer, and flat adenomas. In: Wayne JD, Rex DK, Williams CB, eds. *Colonoscopy: Principles and Practice*. Malden: Blackwell, 2003; 478–86.
- 19 Ikematsu H, Matsuda T, Emura F *et al.* Efficacy of capillary pattern type IIIA/IIIB by magnifying narrow band imaging for estimating depth of invasion of early colorectal neoplasms. *BMC Gastroenterol.* 2010; **10**: 33.
- 20 Uraoka T, Saito Y, Matsuda T *et al.* Endoscopic indications for endoscopic mucosal resection of laterally spreading tumours in the colorectum. *Gut* 2006; **55**: 1592–7.
- 21 Saito Y, Uraoka T, Matsuda T *et al.* Endoscopic treatment of large superficial colorectal tumors: a case series of 200 endoscopic submucosal dissections (with video). *Gastrointest. Endosc.* 2007; **66**: 966–73.
- 22 Saito Y, Fukuzawa M, Matsuda T *et al.* Clinical outcome of endoscopic submucosal dissection versus endoscopic mucosal resection of large colorectal tumors as determined by curative resection. *Surg. Endosc.* 2010; **24**: 343–52.
- 23 Yang S, Farraye FA, Mack C *et al.* BRAF and KRAS mutations in hyperplastic polyps and serrated adenomas of the colorectum: relationship to histology and CpG island methylation status. *Am. J. Surg. Pathol.* 2004; **28**: 1452–9.
- 24 Torlakovic EE, Gomez JD, Driman DK *et al.* Sessile serrated adenoma (SSA) vs traditional serrated adenoma (TSA). *Am. J. Surg. Pathol.* 2008; **32**: 21–9.

Research Article

Visualization of Laterally Spreading Colorectal Tumors by Using Image-Enhanced Endoscopy

Naoto Tamai,¹ Yutaka Saito,¹ Taku Sakamoto,¹ Takeshi Nakajima,¹ Takahisa Matsuda,¹ Namasivayam Vikneswaran,² and Hisao Tajiri³

¹Endoscopy Division, National Cancer Center Hospital, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan

²MBBS, Department of Gastroenterology and Hepatology, Singapore General Hospital, Singapore 769608

³Department of Gastroenterology and Hepatology, The Jikei University School of Medicine, 3-25-8 Nishi Shinbashi, Minato-ku, Tokyo 105-8461, Japan

Correspondence should be addressed to Yutaka Saito, ytsaito@ncc.go.jp

Received 20 September 2011; Accepted 15 October 2011

Academic Editor: Cesare Hassan

Copyright © 2012 Naoto Tamai et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Laterally spreading tumors may sometimes evade detection by colonoscopy. This study aimed to evaluate the use of image-enhanced endoscopy for visualizing laterally spreading tumors of the nongranular type. We reviewed consecutive patients with 47 non-granular-type laterally spreading tumors that had been examined using white-light imaging, autofluorescence imaging, narrow-band imaging, and chromoendoscopy with indigo carmine. The quality of visualization was evaluated using a 5-point scale by less- and more-experienced endoscopists. Autofluorescence imaging provided significantly better visualization than white-light imaging for both less-experienced and experienced endoscopists. On the other hand, no significant differences were observed between the quality of visualization provided by white-light imaging and narrow-band imaging for less-experienced endoscopists. Autofluorescence imaging provides high-quality visualization of non-granular-type laterally spreading tumors on still images. Multicenter trials should be conducted to confirm the usefulness of autofluorescence imaging in detecting laterally spreading colorectal tumors.

1. Introduction

Colorectal carcinoma is one of the most common cancers worldwide, and its prevalence is steadily increasing in Japan [1]. Colonoscopy is considered the gold standard for the detection of neoplastic lesions at risk of progression to colorectal carcinoma. However, according to the results of back-to-back colonoscopies by Rex et al., the miss rate for adenomas ≥ 1 cm was 6% [2]. Laterally spreading tumors (LSTs) constitute a subset of nonpolypoidal colonic lesions, which are characterized by lateral and circumferential extension along the colonic wall rather than vertical growth [3]. LSTs are further classified based on their macroscopic appearance. The granular type LST (LST-G) is defined by the presence of aggregates of even or uneven nodules on the surface, whereas the non-granular-type LST (LST-NG) has a smooth surface lacking the granulonodular formations [4, 5]. Owing to the flat shape of LSTs, the miss rate for these tumors might be higher than the 6% reported by Rex

et al. In addition, LSTs, particularly the NG type, have a higher potential for malignancy; nearly 30% of LST-NGs are associated with lymph follicular or multifocal submucosal invasion [6]. A reduction in the miss rate for LST-NG could therefore contribute to colorectal cancer prevention. Emerging data suggest that the use of image-enhanced endoscopy (IEE) such as autofluorescence imaging (AFI) and narrow-band imaging (NBI) may lead to improvements in polyp detection rates, although this notion remains controversial [7–15]. In our experience, we have encountered many LST-NG lesions that were better visualized by IEE than by white-light imaging (WL). The aim of this study was to evaluate the quality of visualization of LST-NG provided by IEE.

2. Methods


From September 2009 to April 2011, consecutive patients with LST-NG lesions resected by endoscopic submucosal

TABLE 1: Characteristics of lesions.

Number of lesions	47
Number of patients	45
Sex	
Male	31
Female	24
Age (years)	
Median	69
Range	50–80
Tumor size (mm)	
Median	30
Range	20–60
Tumor location	
Cecum	1
Colon	39
Rectum	7
Histopathology	
Adenoma	5
m-ca	24
sm superficial (sm1*)	11
sm deep (sm2-3)	7


*sm1 : sm < 1000 μ m.

dissection (ESD) in our institution were included in this study. The inclusion criteria for performing ESD on LST-NGs were as follows: (1) evidence of a noninvasive pattern [15–17] and (2) lesions larger than 20 mm that were difficult to resect enbloc by using conventional EMR [18]. First, endoscopic examinations were performed using the white-light mode of the AFI videoendoscope system to identify LST-NG lesions, once lesions were detected, the colonoscopist conducted AFI and NBI examinations by switching first to the AFI mode followed by the NBI mode, and finally lesions were examined by chromoendoscopy (CE) using the white-light mode. AFI colonoscopes (EVIS CF-FH260AZI; Olympus Medical Systems, Tokyo, Japan), light sources (EVIS CLV-260SL; Olympus Medical Systems), and video processors (EVLUCERA CV-260SL; Olympus Medical Systems) were used in this study. The AFI videoendoscope system is a novel illumination method that produces real-time pseudocolor images. Neoplastic lesions involve a thickening of the mucosal layer and increased hemoglobin so such lesions emit weaker autofluorescence compared to nonneoplastic lesions; therefore nonneoplastic lesion appears green, while neoplastic lesion has a magenta image [7]. The AFI system allowed for immediate switching from WL to AFI and NBI with a button on the control head of the endoscope. CE was performed using 0.4% indigo carmine. Images of the lesions from WL, NBI, AFI, and CE without magnification were captured and electronically archived in the electronic medical records of our hospital. The images were selected by an experienced endoscopist blinded to this study. The WL, NBI, AFI, and CE images for each lesion were downloaded. The images of all the lesions were randomly arranged, and a Microsoft PowerPoint presentation was created. These images did not contain any information to identify the



*Chi-square test

FIGURE 1: Visualization of LST-NG in group A.



*Fisher's exact test

FIGURE 2: Visualization of LST-NG in group B.

patient or the lesion. The PowerPoint presentations were sent to the respective raters for their independent evaluation. The images were assessed by 2 groups of endoscopists (A and B). Group A comprised 2 physicians with no previous experience in IEE, and group B comprised 2 endoscopists, each of whom had analyzed over 100 cases by using IEE. Each endoscopic image was assessed and given a global rating for visualization based on the ability to detect the lesion and the clarity of the tumor margins. The images were rated by the endoscopists on a 5-point scale as follows: 5, very well visualized; 4, well visualized; 3, moderately well visualized; 2, poorly visualized; 1, very poorly visualized. The ratings of the images were analyzed separately for groups A and B. For each group of raters, the quality of visualization of lesions that received a score of 4 or more from both the raters was classified as "good". The quality of visualization of lesions with a score below 4 was classified as "poor."

3. Statistical Analysis

Statistical analysis was performed using SPSS for Windows (SPSS, Release 6.0; SPSS Inc., Chicago, Ill, USA, 1993). Statistical significance was defined as a *P*-value less than 0.05.




FIGURE 3: LST-NG lesions categorized as “wellvisualized” using AFI. Location: Transverse colon. Size of the lesion: 45 mm. Macroscopic type: Ila (LST-NG). Pathological findings: well-differentiated adenocarcinoma, low-grade atypia, Pm.

4. Results

In all, 49 LST-NG lesions in 47 patients were included in this study. Two patients with lesions were excluded from this study, because the lesions were not observed in the same field in each of the 4 modalities. Finally, a total of 47 LST-NG lesions in 45 patients were evaluated (Table 1). Of the 47 lesions analyzed in group A, the quality of visualization was categorized as “good” for 6 lesions using AFI, 13 using NBI, and 25 using CE. AFI (36/47) provided significantly better visualization than WL (9/47) ($P < 0.001$). Similarly, there was a significant difference between the quality of visualization using CE (25/47) and WLI (9/47) ($P < 0.05$). There was no significant difference, however, between WLI (9/47) and NBI (25/47) (Figure 1). Regarding AFI visualization, there was no significant difference in the macroscopic subtype, tumor location, or underlying histology between well-visualized and poorly visualized lesions, but well-visualized lesions were larger than the poorly visualized lesions (Table 2).

In group B, the quality of visualization was assessed as “good” for 4 lesions by using WLI, 16 lesions by using AFI, 13 lesions by using NBI, and 16 lesions by using CE. There was

a significant difference in the frequency of well-visualized lesions between AFI (16/47) and WLI (4/47) ($P < 0.001$). Similarly, a significant difference in visualization quality was observed between CE (16/47) and WLI (16/47) ($P < 0.01$) and between NBI (13/47) and WLI (4/47) ($P < 0.05$) in group B (Figure 2). Regarding AFI, there was no significant difference in the macroscopic subtype, tumor location, or underlying histology between well-visualized and poorly visualized lesions. Well-visualized lesions were larger than the poorly visualized ones (Table 3).

5. Discussion

Based on the results of our study, AFI provides good-quality visualization of LST-NG lesions, not only for experienced endoscopists but for less-experienced endoscopists as well. The utility of AFI for the detection of colorectal tumors still remains controversial, with studies reporting mixed results [7–9, 15, 19]. In this study, 2 LST-NG lesions were determined to be well visualized by 4 endoscopists (Figures 3 and 4.). As Figures 1 and 2 show, we observed LST-NG lesions that were better visualized using AFI than the other methods. The relationship between visualization and




FIGURE 4: LST-NG lesions categorized as “wellvisualized” by using AFI. Location: lower rectum. Size of the lesion: 45 mm. Macroscopic type: IIa (LST-NG). Pathological findings: well and moderately differentiated adenocarcinoma, pSM (350 μ m).

detection is uncertain. However, better visualization may enable improved detection of LST lesions, especially those of the NG type, which have been shown to be difficult to detect with CE [4]. It is particularly important to improve the detection rate of LST-NGs, because they are more likely to harbor malignancy; nearly 30% of LSTs of the NG type involve lymph follicular or multifocal submucosal invasion [6]. Though LST-NG lesions are less prevalent than polypoidal lesions, their greater malignant potential necessitates reliable detection methods. This study suggests that AFI is superior to WLI for the detection of LST-NG lesions at least on still images. In the present study, there was no significant difference in the quality of visualization of LST-NGs between WLI and NBI for the less-experienced endoscopists.

We also evaluated LST-G lesions in the same fashion as for the LST-NGs. As shown in Figures 5 and 6, AFI also provided good-quality visualization of LST-G lesions for the less-experienced endoscopists, despite the lack of a significant difference in visualization quality between WLI and AFI for the experienced endoscopists. This result indicates that an advantage of AFI might be that it simplifies observations for less-experienced endoscopists. We also compared the backgrounds of the LST-NG lesions between those with good

TABLE 2: Backgrounds of the LST-NG lesion evaluated by AFI in group A.

	Quality of visualization		<i>P</i>
	Good	Poor	
Macroscopic type			
Flat elevated	32	9	0.30*
Flat or flat depressed	4	2	
Lesion size (mm)			
Median	25	35	<0.05**
Range	20–50	20–60	
Location			
Rectum	6	1	0.34*
Cecum or colon	30	10	
Pathological finding			
Adenoma	4	2	0.30*
Adenocarcinoma	32	9	

* Fisher's exact test.

** Mann-Whitney test.


versus poor visualization quality by using AFI. There were no significant differences between lesions that had good versus poor visualization quality with respect to macroscopic

TABLE 3: Characterization of LST-NG lesions by AFI in group B.

	Quality of visualization		P
	Good	Poor	
Macroscopic type			
Flat elevated	16	25	0.07*
Flat or flat depressed	0	6	
Lesion size (mm)			
Median	25	30	<0.05**
Range	20–45	20–60	
Location			
Rectum	2	5	0.32*
Cecum or colon	14	26	
Pathological finding			
Adenoma	4	2	0.08*
Adenocarcinoma	12	29	

* Fisher's exact test.

**Mann-Whitney test.



*Fisher's exact test

**Chi-square test


FIGURE 5: Visualization of LST-G in group A.

type, location, or pathological findings. However, the well-visualized lesions were larger than the poorly visualized lesions in groups A and B. To obtain a whole image of a large lesion, it is necessary to maintain sufficient distance between the tip of the scope and the lesion, which may affect the visibility of the lesion.

This study had several limitations. Only still images were evaluated, and it is uncertain if these findings can be applied to real-time video endoscopy. A relatively small sample precludes any multivariate analysis. Larger studies are needed to define the factors influencing the quality of visualization.

6. Conclusion

AFI provides good-quality visualization of LST-NG lesions on still images. However, to confirm the detectability of LST-NG lesions by using AFI, multicenter trials should be performed.



*Chi-square test

FIGURE 6: Visualization of LST-G in group B.

Conflict of Interests

All authors have no conflict of interests or financial ties to disclose.

References

- [1] T. Matsuda, T. Marugame, K. Kamo et al., "Cancer incidence and incidence rates in Japan in 2002: based on data from 11 population-based cancer registries," *Japanese Journal of Clinical Oncology*, vol. 38, no. 9, pp. 641–648, 2008.
- [2] D. K. Rex, C. S. Cutler, G. T. Lemmel et al., "Colonoscopic miss rates of adenomas determined by back-to-back colonoscopies," *Gastroenterology*, vol. 112, no. 1, pp. 24–28, 1997.
- [3] S. Kudo, "Endoscopic mucosal resection of flat and depressed types of early colorectal cancer," *Endoscopy*, vol. 25, no. 7, pp. 455–461, 1993.
- [4] S. E. Kudo, H. Kashida, T. Tamura et al., "Colonoscopic diagnosis and management of nonpolypoid early colorectal cancer," *World Journal of Surgery*, vol. 24, no. 9, pp. 1081–1090, 2000.
- [5] S. Kudo, R. Shimoda, and H. Kashida, "Laterally spreading tumor of colon: definition and history (in Japanese with English abstract)," *Stomach Intestine*, vol. 40, pp. 1721–1725, 2005.
- [6] T. Uraoka, Y. Saito, T. Matsuda et al., "Endoscopic indications for endoscopic mucosal resection of laterally spreading tumours in the colorectum," *Gut*, vol. 55, no. 11, pp. 1592–1597, 2006.
- [7] T. Matsuda, Y. Saito, K.-I. Fu et al., "Does autofluorescence imaging videoendoscopy system improve the colonoscopic polyp detection rate?—a pilot study," *The American Journal of Gastroenterology*, vol. 103, no. 8, pp. 1926–1932, 2008.
- [8] Y. Takeuchi, T. Inoue, N. Hanaoka et al., "Autofluorescence imaging with a transparent hood for detection of colorectal neoplasms: a prospective, randomized trial," *Gastrointestinal Endoscopy*, vol. 72, no. 5, pp. 1006–1013, 2010.
- [9] T. Kuiper, F. J. C. van den Broek, A. H. Naber et al., "Endoscopic trimodal imaging detects colonic neoplasia as well as standard video endoscopy," *Gastroenterology*, vol. 140, no. 7, pp. 1887–1894, 2011.

- [10] T. Uraoka, Y. Saito, T. Matsuda et al., "Detectability of colorectal neoplastic lesions using a narrow-band imaging system: a pilot study," *Journal of Gastroenterology and Hepatology*, vol. 23, no. 12, pp. 1810–1815, 2008.
- [11] A. Rastogi, D. S. Early, N. Gupta et al., "Randomized, controlled trial of standard-definition white-light, high-definition white-light, and narrow-band imaging colonoscopy for the detection of colon polyps and prediction of polyp histology," *Gastrointestinal Endoscopy*, vol. 74, no. 3, pp. 593–602, 2011.
- [12] T. Inoue, M. Murano, N. Murano et al., "Comparative study of conventional colonoscopy and pan-colonic narrow-band imaging system in the detection of neoplastic colonic polyps: a randomized, controlled trial," *Journal of Gastroenterology*, vol. 43, no. 1, pp. 45–50, 2008.
- [13] A. Rastogi, A. Bansal, S. Wani et al., "Narrow-band imaging colonoscopy—a pilot feasibility study for the detection of polyps and correlation of surface patterns with polyp histologic diagnosis," *Gastrointestinal Endoscopy*, vol. 67, no. 2, pp. 280–286, 2008.
- [14] A. Adler, H. Pohl, I. S. Papanikolaou et al., "A prospective randomised study on narrow-band imaging versus conventional colonoscopy for adenoma detection: does narrow-band imaging induce a learning effect?" *Gut*, vol. 57, no. 1, pp. 59–64, 2008.
- [15] H. Suzuki, Y. Saito, T. Matsuda, T. Nakajima, and T. Kikuchi, "Prospective case study on characterization of colorectal adenomas comparing AFI with NBI," *Diagnostic and Therapeutic Endoscopy*, vol. 2011, Article ID 963618, 6 pages, 2011.
- [16] T. Fujii, R. T. Hasegawa, Y. Saitoh et al., "Chromoscopy during colonoscopy," *Endoscopy*, vol. 33, no. 12, pp. 1036–1041, 2001.
- [17] Y. Saito, F. Emura, T. Matsuda et al., "Invasive pattern is an indication for surgical treatment," *Gut*, 2004, <http://gut.bmjournals.com/cgi/eletters/53/2/284>.
- [18] T. Matsuda, T. Fujii, Y. Saito et al., "Efficacy of the invasive/non-invasive pattern by magnifying chromoendoscopy to estimate the depth of invasion of early colorectal neoplasms," *The American Journal of Gastroenterology*, vol. 103, no. 11, pp. 2700–2706, 2008.
- [19] Y. Saito, T. Uraoka, Y. Yamaguchi et al., "A prospective, multicenter study of 1111 colorectal endoscopic submucosal dissections (with video)," *Gastrointestinal Endoscopy*, vol. 72, no. 6, pp. 1217–1225, 2010.

