

Miida T, Nishimura K, Okamura T, Hirayama S, Ohmura H, Yoshida H, Miyashita Y, Ai M, Tanaka A, Sumino H, Murakami M, Inoue I, Kayamori Y, Nakamura M, Nobori T, Miyazawa Y, Teramoto T, Yokoyama S.	A multicenter study on the precision and accuracy of homogeneous assays for LDL-cholesterol: comparison with a beta-quantification method using fresh serum obtained from non-diseased and diseased subjects.	Atherosclerosis	225(1)	208-15	2012
Toh R, Ishida T, Nishimura K, Nonaka H, Inoue Y, Kitagawa Y, Suematsu M, Miki T, Emoto N, Hirata K.	Comparison of medium-dose losartan/hydrochlorothiazide and maximal-dose angiotensin II receptor blockers in the treatment of Japanese patients with uncontrolled hypertension: the Kobe-CONNECT Study.	Hypertens Res	35(11)	1080-6	2012
Higashiyama A, Wakabayashi I, Kubota Y, Adachi Y, Hayashibe A, Nishimura K, Sugiyama D, Kadota A, Imano H, Miyamatsu N, Miyamoto Y, Okamura T.	Does high-sensitivity C-reactive protein or low-density lipoprotein cholesterol show a stronger relationship with the cardio-ankle vascular index in healthy community dwellers?: the KOBE study.	J Atheroscler Thromb	76(6)	1864-73	2012
Hasokawa M, Shinohara M, Tsugawa H, Bamba T, Fukusaki E, Nishiumi S, Nishimura K, Yoshida M, Ishida T, Hirata K.	Identification of biomarkers of stent restenosis with serum metabolomic profiling using gas chromatography/mass spectrometry.	Circ J	76(8)	1864-73	2012

