

- in 70-year-old men using the euglycaemic insulin clamp. *Diabetologia*, 2005; 48: 862-867
- 28) Wiberg B, Sundström J, Zethelius B, Lind L: Insulin sensitivity measured by the euglycaemic insulin clamp and proinsulin levels as predictors of stroke in elderly men. *Diabetologia*, 2009; 52: 90-96
- 29) DeFronzo RA, Ferrannini E, Simonson DC: Fasting hyperglycemia in non-insulin-dependent diabetes mellitus: contributions of excessive hepatic glucose production and impaired tissue glucose uptake. *Metabolism*, 1989; 38: 387-395
- 30) Sindelar DK, Chu CA, Venson P, Donahue EP, Neal DW, Cherrington AD: Basal hepatic glucose production is regulated by the portal vein insulin concentration. *Diabetes*, 1998; 47: 523-529
- 31) Abdul-Ghani MA, Jenkinson CP, Richardson DK, Tripathy D, DeFronzo RA: Insulin secretion and action in subjects with impaired fasting glucose and impaired glucose tolerance: results from the Veterans Administration Genetic Epidemiology Study. *Diabetes*, 2006; 55: 1430-1435
- 32) Martinez-Hervas S, Argente C, Garcia-Jodar J, Priego A, Real JT, Carratala A, Carmena R, Ascaso JF: Misclassification of subjects with insulin resistance and associated cardiovascular risk factors by homeostasis model assessment index: utility of a postprandial method based on oral glucose tolerance test. *Metabolism*, 2011; 60: 740-746
- 33) Festa A, D'Agostino R Jr, Howard G, Mykkanen L, Tracy RP, Haffner SM: Chronic subclinical inflammation as part of the insulin resistance syndrome: the Insulin Resistance Atherosclerosis Study (IRAS). *Circulation*, 2000; 102: 42-47
- 34) Steinberg HO, Chaker H, Leaming R, Johnson A, Brechtel G, Baron AD: Obesity/insulin resistance is associated with endothelial dysfunction: implications for the syndrome of insulin resistance. *J Clin Invest*, 1996; 97: 2601-2610
- 35) Meigs JB, Mittleman MA, Nathan DM, Tofler GH, Singer DE, Murphy-Sheehy PM, Lipinska I, D'Agostino RB, Wilson PW: Hyperinsulinemia, hyperglycemia, and impaired hemostasis: the Framingham Offspring Study. *JAMA*, 2000; 283: 221-228
- 36) Kaptoge S, Di Angelantonio E, Lowe G, Pepys MB, Thompson SG, Collins R, Danesh J: C-reactive protein concentration and risk of coronary heart disease, stroke, and mortality: an individual participant meta-analysis. *Lancet*, 2010; 375: 132-140

The mutual exacerbation of decreased kidney function and hypertension

Yoshihiro Kokubo

See original paper on page 505

Hypertension is the strongest risk factors for cardiovascular disease worldwide [1,2]. The total population-attributable fractions of higher blood pressure for cardiovascular disease have been estimated as approximately 50% in men and 30% in women [3]. When high blood pressure levels and other risk factors, such as diabetes mellitus [4] or chronic kidney disease (CKD) [5], are combined, the risk of cardiovascular disease becomes much higher. The prevention of hypertension is the best way to prevent cardiovascular disease.

Recently, CKD has become a major public health problem and a risk factor for stroke and coronary heart disease [6]. In end-stage renal disease, the cardiovascular disease mortality rate is more than 10 times as high as that in the general population [7]. In asymptomatic general populations or outpatients, a severely or moderately decreased glomerular filtration rate (GFR) has been shown by most studies to be an independent risk factor for stroke and coronary heart disease [6]. Some studies have shown CKD as an independent risk factor for cardiovascular disease in low-risk or general population [5,8,9]. It is extremely important for the prevention of stroke that we maintain patients' renal function.

Reduced renal function is associated with increased levels of inflammatory factors [10,11], abnormal apolipoprotein levels [10], elevated plasma homocysteine [10], enhanced coagulability [11], anemia, left ventricular hypertrophy, increased arterial calcification, endothelial dysfunction, and arterial stiffness [12]. These factors may contribute to elevated blood pressure.

In prospective cohort studies, hypertension has been shown to be a risk factor for end-stage renal disease in both men and women [13,14]. The Multiple Risk Factor Intervention Trial study showed that elevations of blood

pressure are a strong independent risk factor for end-stage renal disease [14]. Compared with optimal blood pressure, hypertension is a risk factor for end-stage renal disease, with adjusted relative risks (95% confidence intervals) for stages 1–4 hypertension of 3.1 (2.3–4.3), 6.0 (4.3–8.4), 11.2 (7.7–16.2), and 22.1 (14.2–34.3), respectively. Hypertensive patients (BP \geq 160/95 mm Hg) have a five-fold greater decline in GFR (2.7 ml/min per 1.73 m²/year), compared with patients with BP less than 140/90 mm Hg [15]. The Second Manifestations of Arterial Disease study shows that in the presence of albuminuria, higher blood pressure was associated with a greater GFR decrease ($\beta = 3.86$; 95% confidence intervals, 2.34–5.38 for hypertension presence) by a prospective vascular patients study [16]. In addition, the Systolic Hypertension in the Elderly Program study shows that the adjusted relative risk (95% confidence intervals) associated with the highest (\geq 175 mmHg) compared with the lowest quartile (158–163 mm Hg) of SBP was 2.44 (1.67–3.56) for decline in kidney function [17]. Higher blood pressure is a risk factor for decline in kidney function.

There is no original article on the association between decreasing GFR and incident hypertension in general population. In this issue of the *Journal of Hypertension*, Takase *et al.* [18] studied whether GFR can predict incident hypertension in a normotensive general population ($n = 7684$). During the follow-up period (30 624 person-years), hypertension developed in 2031 participants (66.3 per 1000 person-years). The multivariable-adjusted hazard ratio of incident hypertension in the highest tertile (4.4–76.1 ml/min per 1.73 m²) was 1.40 (95% confidence intervals 1.26–1.57) compared with the first tertile. A reduction in GFR of 10 ml/min per 1.73 m² was associated with an 11% increase in risk for incident hypertension. In addition, they investigated the impact of baseline GFR on yearly increases in SBP. The yearly increase in SBP significantly accelerated with a decreasing baseline GFR ($P < 0.01$). A reduction in GFR is a novel predictor of the onset of hypertension in a normotensive general population. Recently, the Suita Study has shown that CKD in the high-normal blood pressure category at the baseline survey was a risk factor for incident hypertension (multivariable-adjusted hazard ratio = 1.41) [19].

A higher level of blood pressure decreases renal function, and a decreased GFR raises blood pressure. In other

Journal of Hypertension 2012, 30:468–469

Department of Preventive Cardiology, National Cerebral and Cardiovascular Center, Suita, Japan

Correspondence to Yoshihiro Kokubo, MD, PhD, FAHA, FACC, FESC, FISH, Department of Preventive Cardiology, National Cerebral and Cardiovascular Center, 5-7-1, Fujishiro-dai, Suita, Osaka, 565-8565 Japan. Tel: +81 6 6833 5012 x2176; fax: +81 6 6833 5300; e-mail: ykokubo@hsp.ncvc.go.jp

J Hypertens 30:468–469 © 2012 Wolters Kluwer Health | Lippincott Williams & Wilkins.

DOI:10.1097/HJH.0b013e328350ade8

words, increases in blood pressure and the decreases of renal function exacerbate each other. In the Suita Study, the risk of cardiovascular disease was higher in CKD patients with normal and high-normal blood pressure than in non-CKD individuals in the same blood pressure categories. In order to prevent cardiovascular disease, control of both blood pressure and renal function are important.

ACKNOWLEDGEMENTS

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Kubo M, Hata J, Doi Y, Tanizaki Y, Iida M, Kiyohara Y. Secular trends in the incidence of and risk factors for ischemic stroke and its subtypes in Japanese population. *Circulation* 2008; 118:2672–2678.
2. Sjol A, Thomsen KK, Schroll M. Secular trends in blood pressure levels in Denmark 1964–1991. *Int J Epidemiol* 1998; 27:614–622.
3. Kokubo Y, Kamide K, Okamura T, Watanabe M, Higashiyama A, Kawanishi K, *et al*. Impact of high-normal blood pressure on the risk of cardiovascular disease in a Japanese urban cohort: the Suita study. *Hypertension* 2008; 52:652–659.
4. Kokubo Y, Okamura T, Watanabe M, Higashiyama A, Ono Y, Miyamoto Y, *et al*. The combined impact of blood pressure category and glucose abnormality on the incidence of cardiovascular diseases in a Japanese urban cohort: the Suita Study. *Hypertens Res* 2010; 33:1238–1243.
5. Kokubo Y, Nakamura S, Okamura T, Yoshimasa Y, Makino H, Watanabe M, *et al*. The relationship between blood pressure category and incidence of stroke and myocardial infarction in an urban Japanese population with and without chronic kidney disease: the Suita Study. *Stroke* 2009; 40:2674–2679.
6. Sarnak MJ, Levey AS, Schoolwerth AC, Coresh J, Cullerton B, Hamm LL, *et al*. Kidney disease as a risk factor for development of cardiovascular disease: a statement from the American Heart Association Councils on Kidney in Cardiovascular Disease, High Blood Pressure Research, Clinical Cardiology, and Epidemiology and Prevention. *Hypertension* 2003; 42:1050–1065.
7. USRDS: the United States Renal Data System. *Am J Kidney Dis* 2003; 42(6 Suppl 5):1–230.
8. Go AS, Chertow GM, Fan D, McCulloch CE, Hsu CY. Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med* 2004; 351:1296–1305.
9. Ninomiya T, Kiyohara Y, Kubo M, Tanizaki Y, Doi Y, Okubo K, *et al*. Chronic kidney disease and cardiovascular disease in a general Japanese population: the Hisayama Study. *Kidney Int* 2005; 68:228–236.
10. Muntner P, Hamm LL, Kusek JW, Chen J, Whelton PK, He J. The prevalence of nontraditional risk factors for coronary heart disease in patients with chronic kidney disease. *Ann Intern Med* 2004; 140:9–17.
11. Shlipak MG, Fried LF, Crump C, Bleyer AJ, Manolio TA, Tracy RP, *et al*. Elevations of inflammatory and procoagulant biomarkers in elderly persons with renal insufficiency. *Circulation* 2003; 107:87–92.
12. Feldman HI, Appel LJ, Chertow GM, Cifelli D, Cizman B, Daugirdas J, *et al*. The Chronic Renal Insufficiency Cohort (CRIC) Study: design and methods. *J Am Soc Nephrol* 2003; 14 (7 Suppl 2):S148–S153.
13. Tozawa M, Iseki K, Iseki C, Kinjo K, Ikemiya Y, Takishita S. Blood pressure predicts risk of developing end-stage renal disease in men and women. *Hypertension* 2003; 41:1341–1345.
14. Klag MJ, Whelton PK, Randall BL, Neaton JD, Brancati FL, Ford CE, *et al*. Blood pressure and end-stage renal disease in men. *N Engl J Med* 1996; 334:13–18.
15. Vupputuri S, Batuman V, Muntner P, Bazzano LA, Lefante JJ, Whelton PK, *et al*. Effect of blood pressure on early decline in kidney function among hypertensive men. *Hypertension* 2003; 42:1144–1149.
16. Vlek AL, van der Graaf Y, Braam B, Moll FL, Nathoe HM, Visseren FL. Blood pressure and decline in kidney function in patients with atherosclerotic vascular disease: a cohort study. *Am J Kidney Dis* 2009; 54:820–829.
17. Young JH, Klag MJ, Muntner P, Whyte JL, Pahor M, Coresh J. Blood pressure and decline in kidney function: findings from the Systolic Hypertension in the Elderly Program (SHEP). *J Am Soc Nephrol* 2002; 13:2776–2782.
18. Takase H, Dohi Y, Toriyama T, Okado T, Tanaka S, Sonoda H, Kimura G. Evaluation of risk for incident hypertension using glomerular filtration rate in the normotensive general population. *J Hypertens* 2012; 30:505–512.
19. Kokubo Y, Nakamura S, Watanabe M, Kamide K, Kawano Y, Kawanishi K, *et al*. Cardiovascular risk factors associated with incident hypertension according to blood pressure categories in nonhypertensive population in the Suita Study: an urban cohort study. *Hypertension* 2011; 58:e132.



Weight Reduction in Primary Care – Comprehensive Dietary Counseling and the Use of Healthy Delivered “Bento (Lunch Boxes)” –

Yoshihiro Kokubo, MD, PhD

A dramatically increasing prevalence of obesity has been observed worldwide in recent decades.^{1,2} Obesity is a risk factor for cardiovascular risk factors, such as hypertension, type 2 diabetes mellitus, and hyperlipidemia,^{3,4} and as a consequence, obese persons have an increased risk of atherosclerosis,⁵ stroke,⁶ ischemic heart disease,⁷ heart failure,⁸ and mortality.⁹ To prevent cardiovascular disease, obese persons must make lifestyle modifications to reduce their weight properly.

Article p 1335

The behavior targets for obesity prevention are focused on healthy diet and physical activity.¹⁰ The healthy diet consists of eating a lot of fruits, vegetables, whole grains, and nuts; limiting calories from added sugars, solid fats, and alcohol; and regulating energy intake rather than eating until the plate is empty. According to the Practical Guide – Identification, Evaluation, and Treatment of Overweight and Obesity in Adults, successful weight reduction requires paying particular attention to the following topics: choosing a balanced diet comprising fats, carbohydrates, and proteins; evaluating nutrition labels to determine caloric content and food composition; giving priority to low-calorie foods; avoiding high-calorie ingredients during cooking (eg, fats and oils); avoiding overconsumption of high-calorie foods (high-fat and high-carbohydrate foods); drinking an adequate amount of water; reducing portion sizes; and limiting alcohol consumption.¹¹ In general, healthcare providers, especially physicians, nurses, and/or dietitians, conduct counseling for the prevention of obesity. There are 3 worthwhile reasons for talking to obese persons about lifestyle modification: (1) participants will understand that a healthy lifestyle is important; (2) an interview about current lifestyle habits opens the door to modifying these habits; and (3) patients may be more responsive to lifestyle modifications when the advice comes from a healthcare provider.¹⁰

Recently, a randomized controlled trial, the Practice-based Opportunities for Weight Reduction (POWER) trial, was performed to determine the effectiveness of 2 behavioral weight loss interventions. The target population consisted of obese adults (≥ 21 years of age) who had one or more cardiovascular risk factors. The first intervention provided patients with weight-loss support remotely by telephone, a study-specific

Web site, and e-mail (remote support only group), and the other intervention provided in-person support during group and individual sessions, along with the 3 remote means of support (in-person support group). There was also a control group in which weight loss was self-directed (control group). The mean change in weight from baseline was significantly decreased by -4.6 kg and -5.1 kg in the group receiving remote support only and the in-person support group, respectively, compared with -0.8 kg in the control group.¹² Participants with obesity achieved and sustained clinically significant weight reduction over 1 year in the in-person support and in the remote support only group, where information was delivered remotely without face-to-face contact between participants and weight loss counselors for 2 years. Now that remote support coaching for weight-loss outcomes is similar to that of in-person visits, the use of mobile technologies to deliver behavioral weight-loss treatment appears to be useful in primary care. However, regardless of the amount of weight lost, it is challenging to maintain weight reduction for many years. Help with maintaining weight loss in the long term may be necessary.

There have been many previous clinical trials of calorie-restricted diets or formula food for weight loss, but there have been few original articles about the combination of delivered meals and dietary counseling in patients with hypertension and/or type 2 diabetes mellitus as an intervention study.^{13,14} Troyer et al conducted a 1-year randomized controlled trial with 298 persons, among whom 50% received 7 Dietary Approach to Stop Hypertension (DASH) meals per week for 1 year. The DASH meals were found to increase compliance with dietary recommendations among noncompliant older adults with cardiovascular disease.¹³ However, in that study, meals were not delivered daily to each individual's house. Individual meal delivery may be simpler and result in greater weight loss.

In a related work that appears in this issue of the Journal, Noda et al studied the effects of dietary counseling by registered dietitians and the use of delivered correctly calorie-controlled meals.¹⁵ They recruited 200 patients with hypertension and/or type 2 diabetes mellitus who were randomly divided into 2 groups, with and without dietary counseling, and consumption of an ordinary diet for 1 month. Each group was then subdivided into 2 groups, with and without dietary counseling, and received calorie-controlled “bento (lunch boxes)” for the

The opinions expressed in this article are not necessarily those of the editors or of the Japanese Circulation Society.

Received April 25, 2012; accepted April 25, 2012; released online May 17, 2012

Department of Preventive Cardiology, National Cerebral and Cardiovascular Center, Suita, Japan

Mailing address: Yoshihiro Kokubo, MD, PhD, FACC, FAHA, FESC, FJSH, Department of Preventive Cardiology, National Cerebral and Cardiovascular Center, 5-7-1 Fujishiro-dai, Suita 565-8565, Japan. E-mail: ykokubo@hsp.ncvc.go.jp

ISSN-1346-9843 doi:10.1253/circ.CJ-12-0554

All rights are reserved to the Japanese Circulation Society. For permissions, please e-mail: cj@j-circ.or.jp

following month. The combination of dietary counseling and delivered calorie-controlled meals was effective for weight reduction as well as alleviating cardiovascular risk factors. Comprehensive counseling that includes the delivery of healthy “bento” may be important for lifestyle modification. However, weight loss is mostly associated with weight regain and, as a result, may not be successful over a person’s lifetime. Further comprehensive cross-over intervention studies are required to support the long-term prevention of obesity, along with the use of a validated nutritional survey to test interventional studies of counseling by dietitians and delivering proper calorie-controlled meals.

References

1. Caterson ID, Hubbard V, Bray GA, Grunstein R, Hansen BC, Hong Y, et al. Prevention Conference VII: Obesity, a worldwide epidemic related to heart disease and stroke: Group III: Worldwide comorbidities of obesity. *Circulation* 2004; **110**: e476–e483.
2. Kumanyika SK, Obarzanek E, Stettler N, Bell R, Field AE, Fortmann SP, et al; American Heart Association Council on Epidemiology and Prevention, Interdisciplinary Committee for Prevention. Population-based prevention of obesity: The need for comprehensive promotion of healthful eating, physical activity, and energy balance: A scientific statement from the American Heart Association Council on Epidemiology and Prevention, interdisciplinary committee for prevention (formerly the expert panel on population and prevention science). *Circulation* 2008; **118**: 428–464.
3. Saito I. Epidemiological evidence of type 2 diabetes mellitus, metabolic syndrome, and cardiovascular disease in Japan. *Circ J* 2012; **76**: 1066–1073.
4. Kashihara H, Lee JS, Kawakubo K, Tamura M, Akabayashi A. Criteria of waist circumference according to computed tomography-measured visceral fat area and the clustering of cardiovascular risk factors. *Circ J* 2009; **73**: 1881–1886.
5. Lim S, Despres JP, Koh KK. Prevention of atherosclerosis in overweight/obese patients: In need of novel multi-targeted approaches. *Circ J* 2011; **75**: 1019–1027.
6. Furukawa Y, Kokubo Y, Okamura T, Watanabe M, Higashiyama A, Ono Y, et al. The relationship between waist circumference and the risk of stroke and myocardial infarction in a Japanese urban cohort: The Suita study. *Stroke* 2010; **41**: 550–553.
7. Cho E, Manson JE, Stampfer MJ, Solomon CG, Colditz GA, Speizer FE, et al. A prospective study of obesity and risk of coronary heart disease among diabetic women. *Diabetes Care* 2002; **25**: 1142–1148.
8. Komukai K, Minai K, Arase S, Ogawa T, Nakane T, Nagoshi T, et al. Impact of body mass index on clinical outcome in patients hospitalized with congestive heart failure. *Circ J* 2012; **76**: 145–151.
9. Tsugane S, Sasaki S, Tsubono Y. Under- and overweight impact on mortality among middle-aged Japanese men and women: A 10-yr follow-up of JPHC Study Cohort I. *Int J Obes Relat Disord* 2002; **26**: 529–537.
10. Cornier MA, Marshall JA, Hill JO, Maahs DM, Eckel RH. Prevention of overweight/obesity as a strategy to optimize cardiovascular health. *Circulation* 2011; **124**: 840–850.
11. National Institutes of Health, National Heart, Lung and Blood Institute, North American Association for the Study of Obesity. The Practical Guide: Identification, Evaluation, and Treatment of Overweight and Obesity in Adults. NIH Publication Number 00-4084. October 2000. Available at: http://www.nhlbi.nih.gov/guidelines/obesity/prctgd_c.pdf (accessed May 8, 2008).
12. Appel LJ, Clark JM, Yeh HC, Wang NY, Coughlin JW, Daumit G, et al. Comparative effectiveness of weight-loss interventions in clinical practice. *N Engl J Med* 2011; **365**: 1959–1968.
13. Troyer JL, Racine EF, Ngugi GW, McAuley WJ. The effect of home-delivered Dietary Approach to Stop Hypertension (DASH) meals on the diets of older adults with cardiovascular disease. *Am J Clin Nutr* 2010; **91**: 1204–1212.
14. Imai S, Kozai H, Matsuda M, Hasegawa G, Obayashi H, Togawa C, et al. Intervention with delivery of diabetic meals improves glycemic control in patients with type 2 diabetes mellitus. *J Clin Biochem Nutr* 2008; **42**: 59–63.
15. Noda K, Zhang B, Iwata A, Nishikawa H, Ogawa M, Ninomiya T, et al on behalf of the STYLIST Study Investigators. Lifestyle changes through the use of delivered meals and dietary counseling in a single-blind study: The STYLIST Study. *Circ J* 2012; **76**: 1335–1344.

Japanese 2011 guidelines for prevention and treatment of osteoporosis—executive summary

Hajime Orimo · Toshitaka Nakamura · Takayuki Hosoi · Masayuki Iki · Kazuhiro Uenishi · Naoto Endo · Hiroaki Ohta · Masataka Shiraki · Toshitsugu Sugimoto · Takao Suzuki · Satoshi Soen · Yoshiki Nishizawa · Hiroshi Hagino · Masao Fukunaga · Saeko Fujiwara

Received: 7 September 2012 / Accepted: 16 October 2012

© The Author(s) 2012. This article is published with open access at SpringerLink.com

Abstract

Introduction In 1998, the first Japanese practice guidelines on osteoporosis was published. It has been updated several times, with the most recent being the full-scale 2011 edition and its abridged edition. The present guidelines provide

information for the managements of primary osteoporosis in postmenopausal women and men over 50 years old, a summary of the evidence for the treatment of secondary osteoporosis, and a summary of the evidence for the prevention of osteoporosis in younger people.

A Report of the Committee for Developing Guidelines for Prevention and Treatment of Osteoporosis: Japan Osteoporosis Society, Japanese Society for Bone and Mineral Research, and Japan Osteoporosis Foundation

H. Orimo
Japan Osteoporosis Foundation,
Tokyo, Japan

T. Nakamura
Department of Orthopedic Surgery, School of Medicine,
University of Occupational and Environmental Health,
Kitakyushu, Fukuoka, Japan

T. Hosoi (✉)
Department of Clinical Research and Development, National
Center for Geriatrics and Gerontology,
35 Gengo, Morioka-machi,
Obu City, Aichi Prefecture 474-8511, Japan
e-mail: t-hosoi@ncgg.go.jp

M. Iki
Department of Public Health,
Kinki University Faculty of Medicine,
Osakasayama, Osaka, Japan

K. Uenishi
Laboratory of Physiological Nutrition, Kagawa Nutrition
University,
Sakato, Saitama, Japan

N. Endo
Division of Orthopedic Surgery, Department of Regenerative and
Transplant Medicine, Niigata University Graduate School of
Medical and Dental Sciences,
Niigata, Niigata, Japan

H. Ohta
Clinical Medical Research Center,
Women's Medical Center of Sanno Medical Center,
Tokyo, Japan

M. Shiraki
Research Institute and Practice for Involutional Diseases,
Azumino, Nagano, Japan

T. Sugimoto
Internal Medicine I, Shimane University Faculty of Medicine,
Shimane, Shimane, Japan

T. Suzuki
Research Institute, National Center for Geriatrics and Gerontology,
Obu, Aichi, Japan

S. Soen
Department of Orthopaedic Surgery and Rheumatology,
Nara Hospital, Kinki University School of Medicine,
Ikoma, Nara, Japan

Y. Nishizawa
Osaka City University,
Osaka, Osaka, Japan

H. Hagino
School of Health Science, Tottori University Faculty of Medicine,
Yonago, Tottori, Japan

Method The present Executive Summary is primarily based on the content of the 2011 Japanese abridged edition. One of the key changes is revision of the criteria for initiation of pharmacological treatment, along with an introduction of the fracture risk factors used in FRAX®. Key figures and tables were selected from the Japanese abridged edition and a reference list was added.

Result and conclusions The essential points of the Japanese practice guidelines on osteoporosis were translated into English for the first time. It is hoped that the content of the guidelines becomes known throughout the world.

Keywords Criteria for initiation of pharmacological treatment · Diagnosis of osteoporosis · Fracture risk assessment · Prevention of osteoporosis · Secondary osteoporosis · Treatment of osteoporosis

Preamble

In 1998, we published the “Guidelines for (Pharmacological) Treatment of Osteoporosis 1998” under the name of the Working Group for Developing Guidelines for Osteoporosis in the Osteoporosis Research Project supported by the Ministry of Health and Welfare (present-day Ministry of Health, Labor, and Welfare) of Japan. Although they were the first Japanese guidelines for the diagnosis and treatment of osteoporosis and also set a precedent for evidence-based practice guidelines in Japan, there were few effective therapeutic agents for osteoporosis available in Japan at that time. The 1998 edition was updated in 2002.

There has been tremendous change in the field of osteoporosis inside and outside Japan since that update. Addressing osteoporosis has become a more urgent issue also in Japan because of its fast-aging society. Therefore, we published the comprehensive “Guidelines for Prevention and Treatment of Osteoporosis 2006” under the name of the Committee for Developing Guidelines for Prevention and Treatment of Osteoporosis 2006, an ad hoc organization comprising the Japan Osteoporosis Society, Japanese Society for Bone and Mineral Research, and Japan Osteoporosis Foundation. Emphasizing prevention, covering secondary osteoporosis, presenting the criteria for initiation of pharmacological treatment, and grading the recommendation for each therapeutic agent, these guidelines were highly rated in the medical and clinical

arenas. Immediately thereafter we published an abridged edition to disseminate the content of the 2006 Guidelines to a greater number of doctors and healthcare professionals.

In late 2011, the 2006 Guidelines and its abridged edition were updated. Key changes are as follows: profile of the research progress on bone quality, revision of the criteria for initiation of pharmacological treatment (associated with the re-examination of the risk factors for fracture and introducing FRAX®), more detailed descriptions about secondary osteoporosis (including new information on the relationship between lifestyle-related diseases and fracture risk), evaluation of new therapeutic agents, and bone metabolic markers covered by public insurance. The present Executive Summary is primarily based on the content of the updated 2011 Japanese abridged edition. Only the most key figures and tables were selected from the Japanese abridged edition and a reference list was added. We hope this Executive Summary contributes to the advancement of medical care for osteoporosis in Asia and the world.

In developing the guidelines, a systematic literature search of MEDLINE, EMBSE, Cochrane Library, and PubMed was conducted. The treatment recommendations in these clinical guidelines were determined by the consensus of the committee. The draft guidelines were available for physician comments at the annual meetings of the Japan Osteoporosis Society in 2010 and 2011.

The funding for all costs to produce the guidelines and this position paper was obtained from the Japan Osteoporosis Society, Japanese Society for Bone and Mineral Research, and Japan Osteoporosis Foundation. All of the authors state they have no conflict of interest related to the guidelines or this position paper.

Definition, epidemiology, and etiology

Definition

The United States National Institutes of Health (NIH) Consensus Development Conference on Osteoporosis Prevention, Diagnosis, and Therapy held in 2000 proposed a new definition of osteoporosis as follows: Osteoporosis is defined as a skeletal disorder characterized by compromised bone strength predisposing a person to an increased risk of fractures. Further, it was stated that bone strength reflects the integration of two main features: bone mineral density (BMD), which accounts for almost 70 % of bone strength, and bone quality, which accounts for the remaining 30 %.

Risk factors for fractures vary among individuals, and include presence or absence of fragility fractures, family history, lifestyle factors, as well as BMD. Therefore, in clinical practice, the risk of fracture should be comprehensively evaluated based on these clinical risk factors for each individual.

M. Fukunaga
Kawasaki Medical School,
Kurashiki, Okayama, Japan

S. Fujiwara
Department of Clinical Studies, Radiation Effects Research
Foundation,
Hiroshima, Hiroshima, Japan

Recently, some algorithms have been developed to quantitatively estimate an individual's fracture risk by integrating multiple risk factors (see "Risk factors for fracture" for FRAX[®]).

Epidemiology

The estimated number of osteoporotic patients aged 40 or over in Japan is 12,800,000 (3,000,000 men and 9,800,000 women), based on the result of a survey of the prevalence of osteoporosis (diagnosed with BMD at the lumbar vertebrae or proximal femur) stratified by age in the general population (Fig. 1) [1] and the population structure stratified by age groups in 2005. Furthermore, the estimated annual incidence of osteoporosis, based on the BMD at the lumbar vertebrae in the population aged between 40 and 79 years, is 0.6 % in men and 2.3 % in women.

The estimated incidence of proximal femoral fractures due to osteoporosis in Japan was 148,100 (31,300 men and 116,800 women) in 2007 [2]. A follow-up study targeting a rural population revealed that the 10-year cumulative incidence of vertebral fractures was 5.1 and 14 % for men and women in their 60s, respectively, and 10.8 and 22.2 % among men and women in their 70s, respectively [3]. However, a long-term trend shows that a later year of birth is associated with a lower incidence of vertebral fractures.

The incidence of proximal femoral fractures was found to be higher in western Japan than in eastern Japan. As compared to reports from Western countries, the incidence of proximal femoral fractures is lower and that of vertebral fractures is similar or higher in Japan.

Etiology

From middle-age onward, BMD decreases and bone quality deteriorates with advancing age, resulting in loss of bone strength. Especially in women, BMD decreases sharply in

the perimenopausal period and for several years thereafter. In addition to this natural course, genetic factors, nutritional deficiency since childhood and puberty, lack of exercise, and unhealthy lifestyle also cause loss of bone strength. Primary osteoporosis is the clinical condition in which these factors have caused a significant loss of bone strength.

Bone remodeling consists of bone resorption by osteoclasts and bone formation by osteoblasts, a mechanism to maintain bone strength. If bone resorption increases with advancing age and menopause and exceeds the rate of bone formation, BMD will begin to decrease. Low BMD is caused by activation of osteoclasts due to estrogen deficiency associated with menopause, and by inadequate secondary mineralization, microarchitecture deterioration, and a decrease in capacity for absorbing calcium associated with advancing age, among other factors (Fig. 2).

Inadequate secondary mineralization and microarchitecture deterioration result in deterioration of bone quality, which is, however, also affected by the cell function of synthesizing bone matrix, conditions surrounding bone matrix (i.e., levels of oxidation and glycation), and levels of vitamins D and K. When oxidative stress and glycation increase in association with aging and lifestyle-related diseases, the non-enzymatic (nonphysiological) cross-links (see "Prevention of falls") increase between collagen molecules in the bone matrix, resulting in a loss of bone strength (Fig. 2).

Prognosis

Fractures associated with osteoporosis, in particular proximal femoral fractures, lead to impairment in mobility and vital functions and an increase in mortality. The relative risk of overall mortality is high in older women with a low BMD and vertebral deformity, and the greater the number of vertebral fractures, the higher the risk of mortality. Decreased BMD at the proximal femur increases the long-

Fig. 1 Estimated prevalence of osteoporosis in Japan. Osteoporosis was diagnosed from BMD at vertebrae L₂₋₄ (a) and proximal femur (b). Data from Yoshimura [1] (Copyright© 2009 Springer Science + Business Media BV)





Fig. 2 Factors causing deterioration of bone strength



term mortality risk, regardless of the presence or absence of vertebral fracture.

According to a survey on quality of life (QOL), patients with osteoporosis score lower on factors related to posture/body shape and falls/psychological in a self-assessment of QOL than persons in the general population who have undergone an osteoporosis screening.

Low BMD is strongly related to the Certification of Needed Long-Term Care for the public nursing-care insurance system in Japan. That is, osteoporosis or low BMD is one of the most significant factors for becoming fragile/immobilized or even becoming bedridden or institutionalized. Therefore, prevention of osteoporotic fractures is likely to prevent reduced mobility or immobilization.

Diagnosis

Diagnostic procedures

The procedures for diagnosis of osteoporosis are shown in Fig. 3 [4].

For the diagnosis of osteoporosis, a medical interview, physical examination, diagnostic imaging, and blood and urine examinations (including measurement of bone metabolic markers) should be conducted first. Then, bone assessment must be conducted with bone mass measurement and spinal radiography. Based on this information, diseases causing low bone mass or secondary osteoporosis should be excluded, and then an accurate diagnosis of primary osteoporosis should be made based on the diagnostic criteria (see "Diagnostic criteria for primary osteoporosis").

Information obtained in the diagnostic process about factors that could contribute to osteoporosis and the risk factors for fractures (e.g., family history, prevalent

fractures, and bone metabolic markers) should be used to evaluate the severity of osteoporosis and the fracture risk. This information will also be useful to provide guidance about lifestyle modification and to select the optimal therapeutic strategy.

Clinical presentation


In the absence of a fracture, osteoporosis is nearly asymptomatic. However, patients with osteoporosis are predisposed to the development of fractures due to loss of bone strength, and the occurrence of fractures will severely impair their QOL (Fig. 4). Osteoporotic fracture is also called fragile fracture.

Proximal femoral fractures directly lead to decreases in the activities of daily living (ADL) and can lead to patients being bedridden, resulting in poor prognosis.

The estimated prevalence of vertebral fractures in Japanese in their early 70s is 25 % and is 43 % in person over 80 years old. The occurrence of vertebral fractures often leads to subsequent vertebral fractures. Since a vertebral deformity persists after the fracture heals, accumulation of vertebral fractures in multiple sites causes kyphosis (round back). Progressive kyphosis leads to deterioration of QOL due to significantly limited ADL and lumbar backache, and can cause functional declines or disorders of the digestive, respiratory, and cardiac systems.

Some lifestyle-related diseases which cause atherosclerosis such as diabetes mellitus (DM), hypertension, dyslipidemia, and chronic kidney diseases (CKD) have attracted attention in relation to osteoporosis. In particular, DM and CKD predispose patients to osteoporosis, and increase their fracture risk (see "Prevention of falls"). The possibility of hidden osteoporosis always should be considered during medical care of patients with lifestyle-related diseases.

Fig. 3 Procedure for the diagnosis of osteoporosis. *YAM* young adult mean (20 to 44 years of age). Adapted from Orimo [4] (Copyright© 2001 Springer Science + Business Media BV)



Medical interview and physical examination


The objectives of the medical interview and physical examination are to assess the presence and symptoms of osteoporotic fractures, risk factors for osteoporosis and fractures, and to obtain information for the differential diagnosis.

Family history of proximal femoral fractures (in either or both parents), loss of height (4 cm or more relative to the height at 25 years of age), current smoking, and excessive alcohol consumption (3 units/day or more, 1 unit=8–10 g

ethanol) are particularly important risk factors for osteoporotic fractures. Therefore, taking a careful history including these factors is needed. History of glucocorticoids use, rheumatoid arthritis, and lifestyle-related diseases such as diabetes mellitus are important information for the differential diagnosis.

In regard to the physical findings, a rounded back, fewer than 20 teeth, and a value of less than -4 on the Female Osteoporosis Self-Assessment Tool for Asians are key factors that strongly suggest osteoporosis.

Fig. 4 Clinical presentation and prognosis of osteoporosis



Bone assessment

It is recommended that BMDs of the lumbar spine and/or proximal femur are measured by dual-energy X-ray absorptiometry (DXA). When there is a fracture or deformity in the lumbar vertebrae that increases the influence of an artifact on spine BMD, the data of lumbar spine should not be used. If the measurement at either of these sites is not successful (because of bilateral hip surgery, multiple fractures of the lumbar vertebra, severe vertebral deformity, or excessive obesity, etc.), another choice is forearm bone.

Microdensitometry has been developed in Japan to radiologically assess BMD, mainly of cortical bone in the second metacarpal.

The speed of sound and broadband ultrasound attenuation through bone are measured with quantitative ultrasound (QUS). This is a non-invasive measurement technique and may provide reliable information on bone quality along with the BMD. However, it is easily affected by measurement conditions, among other factors. The parameters used in QUS were standardized by the QUS Standardization Committee of the Japan Osteoporosis Society in 2010 [5].

Fracture evaluation

Radiography of the thoracic and lumbar vertebrae are essential for assessment of fracture, deformity, or change in the vertebrae, and for exclusion of other similar disorders that present with lower back pain, round back, or low bone mass. In the Japanese diagnostic criteria, the presence of fragility fractures alone confirms the diagnosis of osteoporosis (see "Diagnostic criteria for primary osteoporosis"). Since most of the prevalent fragility fractures, however, are vertebral fractures, usually without pain, radiography is fundamental for their proper diagnosis. Either semiquantitative assessment or quantitative morphometry is used. The lateral DXA images for vertebral fracture assessment can be used, but more clinical experience in Japan is needed to make a recommendation.

If used during the early period after a fracture has occurred (within 2 weeks), MRI provides a better diagnostic yield than plain radiography. MRI is helpful particularly for fresh vertebral fractures, because the height of the vertebral body often does not decrease in the early period. Since it is, however, impractical to diagnose all the cases with MRI, MRI is recommended when it is necessary to distinguish osteoporotic fractures including non-vertebral fractures from those caused by other diseases, or for a detailed examination regarding complicating diseases.

Bone metabolic markers

The increase of bone metabolic markers is a BMD-independent predictor of fractures, and bone metabolic

markers are one of the indices of fracture risk. There are two types of bone metabolic markers: bone resorption markers and bone formation markers. Examinations of blood or urine for these bone metabolic markers easily provide information on the bone metabolic state (Fig. 5) [6].

Bone metabolic markers are useful particularly for the following situations. (1) The patient has little understanding of the need for treatment. (2) The patient is scheduled to receive pharmacotherapy. (3) It is difficult to decide what drug to choose. (4) You want to adopt an appropriate treatment for the patient's pathological condition. Bone metabolic markers are also useful for evaluation of the response to treatment. Thus, it is recommended to measure them at the time of diagnosis if possible.

Among bone metabolic markers, undercarboxylated osteocalcin (ucOC) can be used as an index of vitamin K deficiency in the bones.

When the values of bone resorption markers are abnormally high, the presence of other metabolic bone diseases is suspected.

Differential diagnosis

The targets of differentiation from primary osteoporosis are secondary osteoporosis and other bone-related diseases. Secondary osteoporosis is caused by other diseases or treatments, but its clinical state can seem similar to that of primary osteoporosis, while other bone-related diseases display a clinical state that is different from that of primary osteoporosis. Some instances of secondary osteoporosis and other bone-related diseases are critical or require immediate medical attention. Further, most types of secondary osteoporosis require a therapeutic strategy different from that for primary osteoporosis, and the appropriate treatment of the causative diseases may lead to a dramatic improvement in secondary osteoporosis. Therefore, the differential diagnosis is an extremely important process, despite the prevalence of secondary osteoporosis being low. The probability of secondary osteoporosis is relatively high among premenopausal women and men.

Information for the differential diagnosis can be obtained in every step of the diagnostic process. In the medical interview, thorough medical and surgical histories are needed, including current medications. Radiography may be useful for exclusion of osteomalacia and bone metastases of malignant tumors. Various causative states of secondary osteoporosis may be suspected by the results of blood and urine examinations, for example, hypercalcemia, hypocalcemia, elevated alkaline phosphatase level, and proteinuria.

It is usually considered that patients who visit specialized medical institutes, such as university hospitals, are likely to have secondary osteoporosis due to endocrine diseases and others.




Fig. 5 Measurement of bone metabolic markers in drug treatment of osteoporosis. #1: in patients taking bisphosphonates, measure after stopping drug for at least 6 months, and in patients taking other osteoporosis drugs, measure after stopping drug for at least 1 month. #2: measure one

type each of a resorption marker and formation marker. #3: excluding eldecalcitol. #4: in patients expected to be on long-term bisphosphonate therapy, measure bone resorption markers and BAP or P1NP. Nishizawa [6] (Copyright© 2012 Springer Science + Business Media BV)

Diagnostic criteria for primary osteoporosis

After excluding both the presence of other diseases characterized by low bone mass and the possibility of secondary osteoporosis, primary osteoporosis should be diagnosed by a two-step approach: (1) presence or absence of fragility fractures and (2) BMD or assessment of osteopenia on spinal radiography (Fig. 6) [4].




Fig. 6 Diagnostic criteria for primary osteoporosis (updated in 2000). Primary osteoporosis is diagnosed according to these criteria in the absence of diseases causing low bone mass or secondary osteoporosis. #1: fragility fracture is a nontraumatic bone fracture that is caused by slight external force to a bone with low BMD (BMD less than 80 % of YAM). Sites of fracture include the spine, proximal femur, and the distal end of the radius. #2: BMD usually refers to lumbar BMD. However, when the measurement is inappropriate for reasons such as spinal deformity, the proximal femur BMD should be used. When measurement at those sites is difficult, BMD of the radius, second metacarpal bone, or calcaneus will be used. Revision of additional T-scores is under consideration. Adapted from Orimo [4] (Copyright © 2001 Springer Science + Business Media BV)

Primary osteoporosis is diagnosed on the presence of any fragility fractures (defined as a nontraumatic bone fracture caused by slight external force to a bone with low bone mass, which correlates to a BMD < 80 % of young adult mean (YAM) or radiographic osteopenia of the spine) at sites including spine, proximal femur, and the distal end of radius. If there is no fragility fracture, the BMD level is used to diagnose the patient as “normal”, “decreased bone mass”, or “osteoporosis”. Evaluation of osteopenia based on spinal radiography should be used as supplementary means, and quantitative bone densitometry is preferable for bone assessment.

The T-score to YAM of BMD, not the percentage, is used as diagnostic criteria internationally. A T-score of -1.5 represents a value of -1.5 standard deviation of the YAM and is approximately equivalent to 80 % of the YAM in Japan. A T-score of -2.5 is approximately equivalent to 70 % of the YAM. Internationally, the proximal femur is considered to be the standard measurement site for BMD.

Risk factors

Risk factors for fracture

Major risk factors for osteoporotic fractures are female gender, advanced age, low BMD, and prevalent fractures. In addition, many other factors affect fracture risk directly or indirectly. Although a poor intake of calcium increases fracture risk via low BMD, other risk factors for fractures such as age, prevalent fracture, family history of fractures, smoking, and drinking are independent of BMD. Low body weight also is a BMD-independent risk factor, but only for proximal femoral fractures.

The FRAX® (Fracture Risk Assessment Tool) was developed to estimate the 10-year probability of fractures in individual patients by the World Health Organization (WHO) in 2008 based on 11 risk factors identified from worldwide data in ten cohorts. FRAX is a convenient tool to easily identify a person at high risk for fractures, and therefore has been incorporated into the criteria for initiation of pharmacological treatment in the present guidelines (see "Criteria for initiation of pharmacological treatment").

Prevention

Primary prevention of osteoporosis

The most important measure for primary prevention of osteoporosis is education appropriate to each age group: in early life to acquire as high a peak bone mass (PBM) as possible, to maintain acquired PBM through exercise thereafter, and to minimize its decrease after menopause.

A study on the age-specific distribution of bone mass in Japanese women revealed that PBM is achieved at 18 years of age [7]. Thus, before age 18 is the most effective time for physicians to encourage young people to increase PBM to its maximal level. Guidance on maintenance of adequate weight, active intake of calcium, and weight-bearing exercise is effective.

For middle-aged and older persons, guidance on maintenance of adequate weight, aerobic exercises especially walking, and weight-bearing exercise is effective. Smoking cessation and limiting alcohol intake to less than 3 units/day (1 unit=8–10 g ethanol) is likely to decrease the fracture risk.

Prevention of falls

Most proximal femoral fractures in elderly people occur because of a fall. Risk factors for proximal femoral fractures are a past history of falls and the number of falls, and fall-related factors including generalized weakness, paralysis, muscular weakness, use of sleep-inducing drugs, and decreased vision.

Approaches to prevent falls include (1) exercise interventions (e.g., training to increase strength of muscle, balance, walking ability, and flexibility); (2) non-exercise interventions (e.g., instruction about medication, diet, and environment, along with education and guidance for behavior modification); and (3) multifactorial intervention (e.g., in addition to 1 and 2, an individualized approach based on the physical and mental functioning, environment, and medical assessment of a patient).

In elderly people, vitamin D deficiency increases the risk of falls, and administration of vitamin D can reduce the frequency of falls.

Wearing a hip protector is effective for the prevention of proximal femoral fractures; especially in high-risk groups in elderly care facilities.

Osteoporosis screening

Osteoporosis screening is spreading as a part of the Elderly Health Services (currently as a project under the Health Promotion Law) in Japan, and is performed every 5 years in women from 40 to 70 years old. The screening rate (the percentage of women who underwent osteoporosis screening against the entire target female population) was 4.6 % in 2005.

Osteoporosis screening for people of middle and older age is aimed at early detection of asymptomatic osteoporotic patients and persons at risk of osteoporosis to prevent future fractures. Persons at risk of osteoporosis should be given guidance on diet and exercises, and asymptomatic patients should be targets for early intervention (secondary prevention).

In screening, persons should be classified as either "Complete examination required", "Guidance required", or "No apparent abnormality" based on the results of the medical interview and bone mass measurement (Fig. 7) [8]. The criteria for requiring a complete examination is a bone mass of less than 80 % of YAM; this is different from the diagnostic criteria for osteoporosis (i.e., when BMD is less than 70 % of YAM in the absence of fragility fracture). In addition, bone mass measurement at the calcaneus (including QUS), which is not used to diagnose osteoporosis, is also permitted in the screening. The reason for these differences is that screening should identify the persons requiring the full diagnostic assessment for osteoporosis.

FRAX® will become suitable for osteoporosis screening after the cutoff values for fracture probability are established for complete examination and for guidance.


Treatment

Criteria for initiation of pharmacological treatment

The goals of osteoporosis treatment are prevention of fracture as a complication and maintenance of good skeletal health. Important strategies to reduce the fracture risk in osteoporotic patients are treatment with a bone resorption inhibitor or bone formation stimulant and guidance to establish a lifestyle that leads to maintenance and enhancement of bone strength and to avoid risk factors for fractures, such as a fall, that are independent of a decrease in bone strength.

The risk factors for fracture include low BMD, factors that contribute to a decrease in BMD, and deterioration of bone matrix, including lifestyle-related diseases. A prevalent fragility fracture is the most important among all these

Fig. 7 Criteria for osteoporosis screening. Risk factors for osteoporosis: non-modifiable factors: aging, sex (female), race, family history, late menarche, early menopause, and history of fracture. Modifiable factors: deficiency of calcium, vitamin D, and K; excess intake of phosphorus and salt; excessive calorie restriction; low BMI; lack of physical activity and sunlight; smoking; and excess intake of alcohol and coffee Orimo [8]



factors with the exception of low BMD. Family history of proximal femoral fractures significantly increases the fracture risk even in persons without a fragility fracture who have a "low bone mass" based on their BMD.

Based on this new knowledge about risk factors and the consideration about using FRAX® (see "Risk factors for fracture"), the criteria for initiating pharmacological treatment to prevent fragility fracture was established as shown in Fig. 8. In these criteria, FRAX® is used to consider whether or not to initiate pharmacological treatment in persons without a fragility fracture who have a low bone mass.

This is because persons with a fracture risk comparable to patients with osteoporosis possibly could be included in this group and need other measures to assess the magnitude of the fracture risk other than low BMD. Considering that the 10-year probability of major osteoporotic fractures in the patients receiving pharmacological treatment was observed around 15 % in Japanese clinical settings, we adopted 15 % as a treatment threshold for the persons with low bone mass. In the guidelines, FRAX® is not used in the first-line screening to determine the persons who need further examination such as bone densitometry. As stated earlier, the




Fig. 8 Criteria for initiation of pharmacological treatment. #1: this means proximal femoral fracture and/or vertebral fracture caused by slight external force after menopause in women and after age 50 in men. #2: this means distal forearm, proximal humerus, pelvis, lower leg and/or rib fracture caused by slight external force after menopause in women and after age 50 in men. #3: revision of additional T-scores is under consideration for some measurement sites. #4: this should be applied in persons

<75 years. Additionally, a lower cutoff value does not include all young persons in and around their 50s for whom pharmacological treatment is recommended based on the present diagnostic criteria. #5: as these criteria refer to primary osteoporosis, they should not be applied to persons whose FRAX® risk factors are "glucocorticoid", "rheumatoid arthritis", or "secondary osteoporosis". That is, these criteria should be applied only in persons who answer "No" to each of these items

cutoff value for the screening in Japan is being studied. The cutoff value of a 15 % 10-year probability is used for women and men younger than 75 years old, because almost all of the persons of this age group have a value above 15 % and thus its power as a cutoff value is too weak.

Evaluation of response to treatment

The optimal method for bone mass measurement to evaluate the therapeutic effect is DXA at the lumbar vertebrae on the anteroposterior direction, because it is sensitive enough to detect changes in bone mass. If the bone mass cannot be measured precisely at the lumbar vertebrae, measurement at the total hip is recommended. The timing of measurement should be determined based on the least significant change of each method.

The efficacy of drugs with significant effects on bone metabolism can be evaluated by measuring bone metabolic markers. It is beneficial to measure bone resorption markers at 3 to 6 months after the initiation of treatment and bone formation markers every 6 to 12 months. Attention should be paid to the minimum significant change of each marker.

Plain radiography is useful for detection of incident vertebral fractures after the initiation of treatment. CT, MRI, and bone scintigraphy are sometimes required for confirmation of minor fractures, incomplete fractures, and unapparent fractures, and for differentiation from other clinical conditions including tumors.

QOL assessment using the Japanese Osteoporosis Quality of Life Questionnaire (JOQOL) is useful also for evaluation of therapeutic effects.

Basic treatments (non-pharmacological treatment)

A daily intake of calcium (700 to 800 mg) is recommended to optimize the effect of pharmacological treatment. It has been reported that calcium derivatives and calcium supplements may increase the risk of cardiovascular diseases. However, dietary intake of the same amount of calcium has not been shown to increase cardiovascular risk. Moreover, those adverse findings were reported from outside Japan, where calcium intake, serum lipid levels, and BMI are different from those in Japan. At this time, calcium as a medicine or supplement should not exceed 500 mg per dose.

Vitamin D (recommended daily intake, 10 to 20 μ g) and vitamin K (250 to 300 μ g) are also essential, and they should be prescribed to be taken as a medicine when it is difficult for the patient to obtain a sufficient amount from dietary sources. Hyperhomocysteinemia due to vitamin deficiency (vitamins B₆, B₁₂, and folic acid) involved in homocysteine metabolism has been shown to be a BMD-independent risk factor for fracture. It is recommended to warn patients not to consume excessive amounts of phosphorus, salt content, caffeine, and alcohol.

It has been demonstrated that high-impact activities, resistance exercises, back muscle exercises, stretching exercises, aerobic exercises, walking, and balance training can increase BMD and prevent vertebral fractures and falls in patients with osteoporosis.

In terms of pain relief, few data from randomized controlled trials are available about the effects of various physical therapies, nerve blocks, and surgeries; however, the efficacy of some drugs has been demonstrated.

Pharmacological treatment

These Guidelines detail the effect of each therapeutic agent used in Japan on BMD and the risk of vertebral fracture, non-vertebral fracture, and proximal femoral fracture, based on evidence from Japan and abroad. Each recommendation is also graded (Table 1). In regard to some therapeutic agents, the effect on QOL is also described. Table 2 shows the prescription drugs covered by the public health insurance in Japan.

For the selection of therapeutic agents, the full range of drug-related information must be considered: the efficacy of each medicine on BMD, fracture risk, QOL including pain, bone metabolic markers, risk of fall, as well as safety, including effects other than those on bone metabolism per se and adverse effects. Further, the patient's clinical state must be considered.

The systematic review published by MacLean and colleagues indicated that bisphosphonates (alendronate and risedronate) are a first-line agent for patients at high risk of vertebral, non-vertebral, or proximal femoral fracture [9]. Parathyroid hormone derivatives are first-line agents for patients at high risk of vertebral or non-vertebral fracture. Selective estrogen receptor modulators (SERMs) are first-line agents for patients at high risk of vertebral fracture. Minodronic acid, a bisphosphonate developed in Japan, is expected to be used for the high-risk group for vertebral fracture. Eldecalcitol, an active vitamin D₃ derivative developed in Japan, is expected to be used for the high-risk group for vertebral or non-vertebral fracture. However, more data are required for these new agents.

Estrogen derivatives

A postmenopausal decrease in bone mass is caused by estrogen deficiency. Therefore, estrogen replacement has been considered to be an effective treatment option for osteoporosis since early times. Estrogen replacement is useful also for prevention and treatment of other diseases and symptoms caused by estrogen deficiency. Administration of estrogen to young amenorrheic women or relatively young postmenopausal women can prevent osteoporosis. Estrogen is also useful for treatment of osteoporosis in women with climacteric symptoms in relatively early stage of postmenopause. Conjugated estrogen, estradiol, and estriol are the approved estrogen derivatives in Japan.

Table 1 Grading of recommendation of therapeutic agents for osteoporosis in Japan

Therapeutic agent		BMD	Vertebral fracture	Non-vertebral fracture	Proximal femoral fracture
Calcium	Calcium L-aspartate hydrate	C	C	C	C
	Dibasic calcium phosphate hydrate	C	C	C	C
Estrogen	Estriol	C	C	C	C
	Conjugated estrogens ^a	A	A	A	A
	Estradiol	A	C	C	C
Active vitamin D ₃	Alfacalcidol	B	B	B	C
	Calcitriol	B	B	B	C
	Eldecacitol	A	A	B	C
Vitamin K ₂	Menatetrenone	B	B	B	C
Bisphosphonate	Etidronate disodium	A	B	C	C
	Alendronate sodium hydrate	A	A	A	A
	Sodium risedronate hydrate	A	A	A	A
	Minodronic acid hydrate	A	A	C	C
SERM	Raloxifene hydrochloride	A	A	B	C
	Bazedoxifene acetate	A	A	B	C
Calcitonin ^b	Elcatonin	B	B	C	C
	Calcitonin (Salmon)	B	B	C	C
PTH	Teriparatide (genetical recombination)	A	A	A	C
Other drugs	Ipriflavone	C	C	C	C
	Nandrolone decanoate	C	C	C	C

A strongly recommended to use, B recommended to use, C not enough evidence to recommend use, D recommended not to use

^a Administration of conjugated estrogen for osteoporosis is not covered by the public health insurance in Japan

^b Calcitonin has an analgesic effect, and reduces pain due to osteoporosis (grade A)

Table 2 Prescriptions of anti-osteoporotic agents covered by the public health insurance in Japan (as of September 2011)

Generic name	Launched	Prescription for osteoporosis
Calcium L-aspartate hydrate	1968	1.2 mg/day, p.o.
Dibasic calcium phosphate hydrate	1985	3 g/day, p.o.
Estriol	1969	1 mg/day, p.o.
Conjugated estrogens	1999	Not covered by the public insurance
Estradiol	2008	1 mg/day, p.o.
Alfacalcidol ^a	1981	0.5 or 1 µg/day, p.o. (adult)
Calcitriol	1986	0.5 µg/day, p.o.
Eldecacitol ^a	2011	0.75 or 0.5 µg/day, p.o.
Menatetrenone ^a	1995	45 mg/day, p.o.
Etidronate disodium	1990	200 or 400 mg/day, p.o. (intermittent)
Alendronate sodium hydrate	2001	5 mg/day or 35 mg/w, p.o.
Sodium risedronate hydrate	2002	2.5 mg/day or 17.5 mg/w, p.o.
Minodronic acid hydrate ^a	2009	1 mg/day or 50 mg/4w, p.o.
Raloxifene hydrochloride	2004	60 mg/day, p.o.
Bazedoxifene acetate	2010	20 mg/day, p.o.
Elcatonin ^a	1982	20 IU/w, i.m.
Calcitonin (Salmon)	1990	20 IU/w, i.m.
Teriparatide (genetical recombination)	2010	24 µg/day, s.c. (up to 24 months)
Ipriflavone	1988	200 mg/day, p.o.
Nandrolone decanoate	1984	25 or 50 mg/3 w, i.m.

Teriparatide acetate, a new drug developed in Japan, came to market in November 2011. Prescription is 56.5 µg/w, s.c., up to 72 weeks

^aAgents developed in Japan

Although conjugated estrogen increases BMD and prevents vertebral, non-vertebral, and proximal femoral fracture, it is not covered by the public health insurance in Japan for the treatment of osteoporosis.

Estradiol increases BMD, but there is little evidence that it prevents fractures.

There is almost no evidence about the effects of estriol.

Alfacalcidol and calcitriol (active vitamin D₃ derivatives)

Alfacalcidol and calcitriol are active vitamin D₃ derivatives. Alfacalcidol, developed in Japan, is a prodrug requiring hydroxylation in the liver for activation. Because these derivatives were approved for the treatment of osteoporosis in 1983 and 1989, respectively, there is insufficient large clinical trial data. However, several reports suggested these agents maintain lumbar BMD at a significantly higher level as compared to placebo, or reduce the risk of vertebral and non-vertebral fractures (not statistically significant; Fig. 9a) [10].

It has been reported also that vitamin D deficiency causes atrophy of the type II muscle fibers, and that vitamin D supplementation improves trunk imbalance. Active vitamin D₃ derivatives (alfacalcidol and calcitriol) reduce falls among the elderly (Fig. 9b) [11]. These active vitamin D₃ derivatives have been confirmed to be safe, even for long-term use, and they are recommended for the elderly (see "Combination therapy" for combination with bisphosphonate).

Eldecalcitol (active vitamin D₃ derivative)

Although the conventional active vitamin D₃ derivatives have been reported to be effective for preventing fractures, they have not been shown to increase BMD significantly. Various vitamin D₃ derivatives have been investigated; of these eldecalcitol was developed in Japan. Eldecalcitol showed superior efficacy to alfacalcidol to increase BMD (Fig. 10a) [12], while its effect on calcium absorption was nearly unchanged. Eldecalcitol may exert its actions by




Fig. 9 Meta-analyses on the efficacy of vitamin D. **a** Relative risk for vertebral fractures after treatment with vitamin D. *RR* relative risk, *CI* confidence interval. *Open rhombus* indicates using calcitriol and *closed rhombus* using alfacalcidol. Adapted from Papadimitropoulos

[10] (Copyright© 2002 The Endocrine Society). **b** Compared risk of falling between vitamin D-treated group and control group. *OR* odds ratio, *CI* confidence interval. Adapted from Bischoff-Ferrari [11] (Copyright© 2004 American Medical Association)




Fig. 10 Effect of eldecalcitol compared with alfacalcidol. **a** Change in lumbar BMD. Data are mean \pm SE, * $p < 0.001$ vs alfacalcidol group by Student *t* test (Matsumoto [12] (Copyright© 2011 Elsevier)). **b** Incidence of vertebral fracture. Hazard ratio (HR) is 0.74 and 95 % confidence interval (CI) is 0.56–0.97. Data from Matsumoto [12] (Copyright© 2011 Elsevier). **c** Incidence of non-vertebral fractures. HR for three major non-

vertebral fractures is 0.52 and 95 % CI is 0.29–0.93, $p = 0.031$. Three major non-vertebral sites mean humerus, wrist, and hip, i.e., the three sites of major non-vertebral fractures recognized as osteoporotic fractures in FRAX®. HR for wrist fractures is 0.29 and 95 % CI is 0.11–0.77, $p = 0.005$. Data from Matsumoto [12] (Copyright© 2011 Elsevier) and the website of Pharmaceuticals and Medical Devices Agency (in Japanese)

promoting calcium absorption from the small intestine, similar to the conventional active vitamin D₃ derivatives, and prevent bone resorption by inhibiting osteoclastic function.

In a comparative study of eldecalcitol and alfacalcidol, the incidence of vertebral fractures was found to be significantly lower in the eldecalcitol group (Fig. 10b) [12]. While there was no significant difference in the overall incidence of non-vertebral fractures between the eldecalcitol and alfacalcidol groups, there was a trend towards a greater decrease in the incidence of non-vertebral fractures at the three major sites (humerus, wrist, and hip) in the eldecalcitol group than in the alfacalcidol group (Fig. 10c) [12]. Of note, the incidence of wrist fractures was significantly reduced in the eldecalcitol group.

Clinical trials of eldecalcitol have been conducted in patients over a wide range of age and severity, and this agent can be used across the entire spectrum of patients with osteoporosis.

Menatetrenone (vitamin K₂ derivative)

In elderly women and patients with osteoporosis being treated with a bisphosphonate, insufficient intake of vitamin K is a BMD-independent risk factor for fractures. Menatetrenone, a vitamin K₂ derivative, promotes carboxylation of osteocalcin, and thereby it reduces the serum level of ucOC, an index of vitamin K deficiency.

Menatetrenone slightly increases lumbar BMD and reduces vertebral and non-vertebral fractures (Fig. 11)

[13]. Menatetrenone is considered to exert its fracture-reducing effect via a mechanism of action other than increasing BMD.

Etidronate (bisphosphonate)

Notably, for etidronate, a first-generation bisphosphonate, there is a small margin between its serum level for the onset of its inhibitory actions on bone resorption and the serum level for its inhibitory effects on bone formation. Close attention must be paid to its narrow safety range. Thus, a cyclical intermittent treatment strategy (200 to 400 mg/day once daily for 2 weeks, followed by a rest period of 10 to 12 weeks) is essential.

Because etidronate reduces bone resorption, it is effective particularly for high-turnover osteoporosis, and it maintains bone mass even in low-turnover osteoporosis. Etidronate reduces blood and urine levels of bone metabolic markers. Etidronate reduces incident vertebral fractures in patients who have vertebral fractures. There is no clear evidence about whether or not etidronate reduces non-vertebral fractures.

Alendronate (bisphosphonate)

Alendronate, a second-generation bisphosphonate, has a very wide safety range. Its inhibitory effect on bone resorption is exerted at a much smaller dose than the dose for its inhibitory effect on bone formation (approximately 1/6,000).




Fig. 11 Meta-analysis on the efficacy of menatetrenone on fractures. OR odds ratio, CI confidence interval (Cockayne [13] (Copyright© 2006 American Medical Association))

Many clinical trials and meta-analyses have shown that alendronate increases BMD, reduces fractures at the vertebra/non-vertebra, proximal femur, and distal end of the forearm; and improves the bone metabolic marker profile. Alendronate has been reported to reduce vertebral fracture and increase lumbar BMD also in men with osteoporosis.

In terms of QOL, a decrease in the duration of bed rest for low back pain, a decrease in the days of activity restriction, and improvement of arthralgia and pain-related QOL scores after treatment with alendronate have been reported (see "Combination therapy" for the combination with active vitamin D₃ derivatives).

A once-weekly dose of alendronate (35 mg), compared to a daily dose of alendronate (5 mg) was shown to have a similar effect on lumbar BMD and urinary levels of type I collagen cross-linked N-telopeptides (NTX); the incidence of adverse reactions and drug discontinuation was lower in the once-weekly group.

Risedronate (bisphosphonate)

Risedronate, a third-generation bisphosphonate, has a strong inhibitory effect on bone resorption.

Many clinical trials and meta-analyses have shown that risedronate increases BMD and reduces fractures at the vertebra/non-vertebra and proximal femur in postmenopausal women. Risedronate was reported to increase lumbar BMD also in men with osteoporosis. Large-scale clinical

trials in North America, Europe, and Australia have shown preventive effects with risedronate against incident vertebral fracture from the first year of treatment. In Japan, it was reported that risedronate improved scores for body pain, vitality, and social functioning in QOL assessment using the SF-36 scale.

Once-weekly risedronate (35 mg), compared to daily risedronate (5 mg), was shown to increase BMD at the femoral neck and trochanter to the same degree in a study in the USA. In a Japanese clinical trial, once-weekly risedronate (17.5 mg), compared to daily risedronate (2.5 mg), increased lumbar BMD to the same degree at 48 weeks.


Minodronic acid (bisphosphonate)

Minodronic acid is the only domestically developed bisphosphonate for osteoporosis, and the only bisphosphonate which has been investigated for its inhibitory effect on fracture in Japanese patients at doses approved in Japan. Minodronic acid has the strongest inhibitory effect on bone resorption among the bisphosphonates currently available in Japan.

The efficacy of minodronic acid on BMD at the lumbar spine and total hip is equivalent to alendronate (Fig. 12a) [14]. In addition, minodronic acid significantly increased BMD in patients who had a poor response to other bisphosphonates.

Minodronic acid reduced vertebral fracture risk by 59 % in Japanese patients with osteoporosis (Fig. 12b) [15], and no

Fig. 12 Effect of minodronic acid on BMD and vertebral fracture. **a** Percent change in lumbar spine and total hip BMD. *Solid line* is minodronic acid 1 mg ($n=134$) and *broken line* is alendronate 5 mg ($n=135$). Data are mean \pm SE (Hagino [14] (Copyright© 2009 Elsevier)). **b** Incidence of vertebral fracture. *Solid line* is minodronic acid ($n=339$) and *broken line* is placebo ($n=328$). Relative risk is 0.411 (95 % confidence interval 0.267–0.634) by Cox regression model. * $p<0.0001$ by log-rank test between the groups (Matsumoto [15] (Copyright© 2009 Springer Science + Business Media BV))



difference was observed in the effect between patients above and below 75 years of age. No clinical trial to determine the effect of minodronic acid on non-vertebral fracture or proximal femoral fracture has been conducted. The results of the ongoing Japanese Osteoporosis Intervention Trial (JOINT)-04 initiated in 2011 by the Adequate Treatment of Osteoporosis (A-TOP) Research Group (see “Combination therapy”) are greatly anticipated to answer these questions. Minodronic acid is available for daily use (1 mg) and once every 4 weeks (50 mg).

Raloxifene (SERM)

Raloxifene, a selective estrogen receptor modulator, binds to the estrogen receptor (ER) with an affinity equivalent to estrogen and induces a conformational change at the helix 12 in the C-terminal part of ER; this conformational change produced by raloxifene is different from that produced by estrogen. Thus, raloxifene has a tissue-selective pharmacological action: it shows estrogen-like effects on bone, but not on the breast or uterus.

The Multiple Outcomes of Raloxifene Evaluation, a large-scale randomized controlled trial with 7,705 patients in 25 countries, demonstrated that raloxifene increased BMD and reduced incident vertebral fractures, regardless of the presence or absence of prevalent vertebral fractures and even in subjects with low bone mass (osteopenia). Additionally, raloxifene significantly reduced the incidence of non-vertebral fractures in patients with severe vertebral fractures. In Japan, a 3-year post-marketing surveillance

demonstrated that the overall incidence of clinical fractures was as low as 1.2 %.

Many observational studies from Japan and abroad demonstrated the effect of raloxifene on QOL, including pain relief. A meta-analysis revealed that raloxifene decreases the overall mortality by 10 %.

Venous thromboembolism is one of the clinically important adverse events of SERMs. The incidence of venous thromboembolism in patients treated with raloxifene is 0.2 %, stated in the drug package insert, based on the results of a 3-year post-marketing surveillance conducted in 7,557 Japanese patients.

Bazedoxifene (SERM)

Bazedoxifene, a SERM, has an estrogen-like action selectively on bone metabolism and lipid metabolism, but not on the breast or uterus.

An international multi-center clinical trial demonstrated that bazedoxifene increases BMD and reduces vertebral fractures, similar to raloxifene. Although no overall reduction on non-vertebral fractures was observed with bazedoxifene, the incidence of non-vertebral fracture in postmenopausal women at a higher risk of fracture was significantly reduced by bazedoxifene as compared to placebo and raloxifene. Additionally, the higher the FRAX® score, the more effectively bazedoxifene reduced osteoporotic fractures. Bazedoxifene was also reported to improve the profile of bone metabolic markers. The effect of bazedoxifene on proximal femoral fracture has not been studied yet.