

each sex and age group are shown in Table 6.

#### REFERENCES

- 1) Food and Nutrition Board, Institute of Medicine. 2005. Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acid, Cholesterol, Protein, and Amino Acids, p 107–264. National Academies Press, Washington DC.
- 2) Brooks GA, Butte NF, Rand WM, Flatt JP, Caballero B. 2004. Chronicle of the Institute of Medicine physical activity recommendation: how a physical activity recommendation came to be among dietary recommendations. *Am J Clin Nutr* **79**(Suppl): S921–930.
- 3) Yanai R, Masuda T, Kitagawa S, Nagao N, Nagao M, Matsueda S. 2006. Relationship between overreporting or underreporting among young males and females and physical factors, psychosocial factors, and lifestyle habits. *Journal of Kawasaki Medical Welfare Society* **16**: 109–119.
- 4) Shimada M, Nishimuta M, Kodama N, Yoshitake Y. 2006. Existence in subject of low plasma triiodothyronine correlated with post-absorptive resting metabolism measurement of T3 is essential for determining standard basal metabolic rate. *Jpn J Phys Fitness Sports Med* **55**: 295–306 (in Japanese).
- 5) Taguchi M, Higuchi M, Oka J, Yoshiga C, Ishida Y, Matsushita M. 2001. Basal metabolic rate in Japanese female endurance athletes. *Jpn J Nutr Diet* **59**: 127–134 (in Japanese).
- 6) Usui C, Takahashi E, Gando Y, Sanada K, Oka J, Miyachi M, Tabata I, Higuchi M. 2007. Relationship between blood adipocytokines and resting energy expenditure in young and elderly women. *J Nutr Sci Vitaminol* **53**: 529–535.
- 7) Yamamura C, Tanaka S, Futami J, Oka J, Ishikawa Takata K, Kashiwazaki H. 2003. Activity diary method for predicting energy expenditure as evaluated by a whole-body indirect human calorimeter. *J Nutr Sci Vitaminol* **49**: 262–269.
- 8) Ganpule AA, Tanaka S, Ishikawa-Takata K, Tabata I. 2007. Interindividual variability in sleeping metabolic rate in Japanese subjects. *Eur J Clin Nutr* **61**: 1256–1261.
- 9) Hirose M. 1989. Study of basal metabolism of modern middle age Japanese. *Ehime Medical* **8**: 192–210 (in Japanese).
- 10) Hioki C, Arai M. 2007. Bofutsushosan use for obesity with IGT: search for scientific basis and development of effective therapy with Kampo medicine. *J Trad Med* **24**: 115–127.
- 11) Usui C, Oka J, Yamakawa J, Sasaki Y, Higuchi M. 2003. Basal metabolic rate and its determinants in post-menopausal women. *Jpn J Phys Fitness Sport Med* **52**: 189–198.
- 12) Yokozeki T. 1993. Basal metabolic rate and physical activity in the elderly. *J Jpn Soc Nutr Food Sci* **46**: 451–458 (in Japanese).
- 13) Yokozeki T. 1993. Basal metabolic rate and energy requirement of bed-ridden elderly women. *J Jpn Soc Nutr Food Sci* **46**: 459–466 (in Japanese).
- 14) Tahara Y. 1983. Seasonal variation of heat production by body composition in basal metabolic condition and cold exposure. *J Jpn Soc Nutr Food Sci* **36**: 255–263 (in Japanese).
- 15) Maeda T, Fukushima T, Ishibashi K, Higuchi S. 2007. Involvement of basal metabolic rate in determination of type of cold tolerance. *J Physiol Anthropol* **26**: 415–418.
- 16) Shetty PS, Henry CJ, Black AE, Prentice AM. 1996. Energy requirements of adults: an update on basal metabolic rates (BMRs) and physical activity levels (PALs). *Eur J Clin Nutr* **50**: S11–23.
- 17) Black AE, Coward WA, Cole TJ, Prentice AM. 1996. Human energy expenditure in affluent societies: an analysis of 574 doubly-labelled water measurements. *Eur J Clin Nutr* **50**: 72–92.
- 18) Schutz Y, Weinsier RL, Hunter GR. 2001. Assessment of free-living physical activity in humans: an overview of currently available and proposed new measures. *Obes Res* **9**: 368–379.
- 19) Ishikawa-Takata K, Tabata I, Sasaki S, Rafamantanantsoa HH, Okazaki H, Okubo H, Tanaka S, Yamamoto S, Shirota T, Uchida K, Murata M. 2008. Physical activity level in healthy free-living Japanese estimated by doubly-labelled water method and International Physical Activity Questionnaire. *Eur J Clin Nutr* **62**: 885–891.
- 20) Ebine N, Feng JY, Homma M, Saitoh S, Jones PJ. 2000. Total energy expenditure of elite synchronized swimmers measured by the doubly labeled water method. *Eur J Appl Physiol* **83**: 1–6.
- 21) Ebine N, Rafamantanantsoa HH, Nayuki Y, Yamanaka K, Tashima K, Ono T, Saitoh S, Jones PJ. 2002. Measurement of total energy expenditure by the doubly labeled water method in professional soccer players. *J Sports Sci* **20**: 391–397.
- 22) Ebine N, Shimada M, Tanaka H, Nishimuta M, Yoshitake Y, Saitoh S, Jones PJH. 2002. Comparative study of total energy expenditure in Japanese men using doubly labeled water method against activity record, heart rate monitoring, and accelerometer methods. *Jpn J Phys Fitness Sport Med* **51**: 151–163 (in Japanese).
- 23) Rafamantanantsoa HH, Ebine N, Yoshioka M, Higuchi H, Yoshitake Y, Tanaka H, Saitoh S, Jones PJ. 2002. Validation of three alternative methods to measure total energy expenditure against the doubly labeled water method for older Japanese men. *J Nutr Sci Vitaminol* **48**: 517–523.
- 24) Touno M, Rafamantanantsoa HH, Ebine N, Peng HY, Yoshitake Y, Tanaka H, Saitoh S. 2003. Measurement of total energy expenditure of firefighters on normal work shift. *Jpn J Phys Fitness Sport Med* **52**: 265–274 (in Japanese).
- 25) Rafamantanantsoa HH, Ebine N, Yoshioka M, Yoshitake Y, Tanaka H, Saitoh S. 2003. The effectiveness of three-day dietary records with advanced photo system camera for measuring energy intake in Japanese men as determined by doubly labeled water technique. *J Clin Biochem Nutr* **33**: 33–38.
- 26) Rafamantanantsoa HH, Ebine N, Yoshioka M, Yoshitake Y, Tanaka H, Saitoh S, Jones PJ. 2003. The role of exercise physical activity in varying the total energy expenditure in healthy Japanese men 30 to 69 years of age. *J Nutr Sci Vitaminol* **49**: 120–124.
- 27) Peng HY, Yoshitake Y, Saitoh S. 2004. Validity of methods to measure total energy expenditure of middle-aged women: a validation study against doubly labeled water method. *J Jpn Soc Study Obes* **10**: 163–172.
- 28) Peng HY, Shibata U, Yoshitake Y, Saito S, Omi N. 2005. Energy balance and nutritional status in middle-aged Japanese women with a long-term habit of exercise. *J Jpn Soc Nutr Food Sci* **58**: 329–335.

- 29) Peng HY, Saito S, Hikiyara Y, Ebine N, Yoshitake Y. 2005. Energy expenditure, body composition and maximal oxygen uptake in middle-aged Japanese women who have long-term habits of exercising. *Jpn J Phys Fitness Sport Med* **54**: 237–248 (in Japanese).
- 30) Hikiyara Y, Saitoh S, Yoshitake Y. 2005. Validity of methods to measure total energy expenditure of baseball players in Japanese high school. *J Jpn Soc Nutr Food Sci* **54**: 363–372 (in Japanese).
- 31) Adachi M, Sasayama K, Hikiyara Y, Okishima K, Mizuuchi H, Sunami Y, Shiomi M, Nishimuta M, Kikunaga S, Tanaka H, Saitoh S, Yoshitake Y. 2007. Assessing daily physical activity in elementary school students used by accelerometer: a validation study against doubly labeled water method. *Jpn J Phys Fitness Sports Med* **56**: 347–356 (in Japanese).
- 32) Yamamoto S, Ishikawa-Takata K, Bessyo K, Tanimoto M, Miyachi M, Tanaka S, Totani M, Tabata I. 2008. Basal metabolic rate and physical activity level in bodybuilders. *Jpn J Nutr* **66**: 195–200 (in Japanese).
- 33) Yamada Y, Yokoyama K, Noriyasu R, Osaki T, Adachi T, Itoi A, Naito Y, Morimoto T, Kimura M, Oda S. 2009. Light-intensity activities are important for estimating physical activity energy expenditure using uniaxial and triaxial accelerometers. *Eur J Appl Physiol* **105**: 141–152.
- 34) Baarends EM, Schols AM, Pannemans DL, Westerterp KR, Wouters EF. 1997. Total free living energy expenditure in patients with severe chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* **155**: 549–554.
- 35) Sawaya AL, Saltzman E, Fuss P, Young VR, Roberts SB. 1995. Dietary energy requirements of young and older women determined by using the doubly labeled water method. *Am J Clin Nutr* **62**: 338–344.
- 36) Rothenberg E, Bosaeus I, Lernfelt B, Landahl S, Steen B. 1998. Energy intake and expenditure: validation of a diet history by heart rate monitoring, activity diary and doubly labeled water. *Eur J Clin Nutr* **52**: 832–838.
- 37) Reilly JJ, Lord A, Bunker VW, Prentice AM, Coward WA, Thomas AJ, Briggs RS. 1993. Energy balance in healthy elderly women. *Br J Nutr* **69**: 21–27.
- 38) Bonnefoy M, Normand S, Pachiardi C, Lacour JR, Laville M, Kostka T. 2001. Simultaneous validation of ten physical activity questionnaires in older men: a doubly labeled water study. *J Am Geriatr Soc* **49**: 28–35.
- 39) Blanc S, Schoeller DA, Bauer D, Danielson ME, Tylavsky F, Simonsick EM, Harris TB, Kritchevsky SB, Everhart JE. 2004. Energy requirements in the eighth decade of life. *Am J Clin Nutr* **79**: 303–310.
- 40) Manini TM, Everhart JE, Patel KV, Schoeller DA, Colbert LH, Visser M, Tylavsky F, Bauer DC, Goodpaster BH, Harris TB. 2006. Daily activity energy expenditure and mortality among older adults. *JAMA* **296**: 171–179.
- 41) Rothenberg EM, Bosaeus IG, Steen BC. 2003. Energy expenditure at age 73 and 78—A five year follow-up. *Acta Diabetol* **40**: S134–138.
- 42) Fuller NJ, Sawyer MB, Coward WA, Paxton P, Elia M. 1996. Components of total energy expenditure in free-living elderly men (over 75 years of age): measurement, predictability and relationship to quality-of-life indices. *Br J Nutr* **75**: 161–173.
- 43) Rothenberg EM, Bosaeus IG, Westerterp KR, Steen BC. 2000. Resting energy expenditure, activity energy expenditure and total energy expenditure at age 91–96 years. *Br J Nutr* **84**: 319–324.
- 44) Wren RE, Blume H, Mazariegos M, Solomons N, Alvarez JO, Goran MI. 1997. Body composition, resting metabolic rate, and energy requirements of short- and normal-stature, low-income Guatemalan children. *Am J Clin Nutr* **66**: 406–412.
- 45) Fontvieille AM, Harper IT, Ferraro RT, Spraul M, Ravussin E. 1993. Daily energy expenditure by five-year-old children, measured by doubly labeled water. *J Pediatr* **123**: 200–207.
- 46) Bunt JC, Salbe AD, Harper IT, Hanson RL, Tataranni PA. 2003. Weight, adiposity, and physical activity as determinants of an insulin sensitivity index in Pima Indian children. *Diabetes Care* **26**: 2524–2530.
- 47) Franks PW, Ravussin E, Hanson RL, Harper IT, Allison DB, Knowler WC, Tataranni PA, Salbe AD. 2005. Habitual physical activity in children: the role of genes and the environment. *Am J Clin Nutr* **82**: 901–908.
- 48) Hoos MB, Plasqui G, Gerver WJ, Westerterp KR. 2003. Physical activity level measured by doubly labeled water and accelerometry in children. *Eur J Appl Physiol* **89**: 624–626.
- 49) Livingstone MB, Coward WA, Prentice AM, Davies PS, Strain JJ, McKenna PG, Mahoney CA, White JA, Stewart CM, Kerr MJ. 1992. Daily energy expenditure in free-living children: comparison of heart-rate monitoring with the doubly labeled water (2H2(18)O) method. *Am J Clin Nutr* **56**: 343–352.
- 50) Dugas LR, Ebersole K, Schoeller D, Yanovski JA, Barquera S, Rivera J, Durazo-Arzivu R, Luke A. 2008. Very low levels of energy expenditure among pre-adolescent Mexican-American girls. *Int J Pediatr Obes* **3**: 123–126.
- 51) Luke A, Roizen NJ, Sutton M, Schoeller DA. 1994. Energy expenditure in children with Down syndrome: correcting metabolic rate for movement. *J Pediatr* **125**: 829–838.
- 52) Ramirez-Marrero FA, Smith BA, Sherman WM, Kirby TE. 2005. Comparison of methods to estimate physical activity and energy expenditure in African American children. *Int J Sports Med* **26**: 363–371.
- 53) Treuth MS, Figueroa-Colon R, Hunter GR, Weinsier RL, Butte NF, Goran MI. 1998. Energy expenditure and physical fitness in overweight vs non-overweight prepubertal girls. *Int J Obes Relat Metab Disord* **22**: 440–447.
- 54) Treuth MS, Butte NF, Wong WW. 2000. Effects of familial predisposition to obesity on energy expenditure in multiethnic prepubertal girls. *Am J Clin Nutr* **71**: 893–900.
- 55) Maffei C, Pinelli L, Zaffanello M, Schena F, Iacumin P, Schutz Y. 1995. Daily energy expenditure in free-living conditions in obese and non-obese children: comparison of doubly labelled water (2H2(18)O) method and heart-rate monitoring. *Int J Obes Relat Metab Disord* **19**: 671–677.
- 56) Spadano JL, Bandini LG, Must A, Dallal GE, Dietz WH. 2005. Longitudinal changes in energy expenditure in girls from late childhood through midadolescence. *Am J Clin Nutr* **81**: 1102–1109.
- 57) Anderson SE, Bandini LG, Dietz WH, Must A. 2004. Relationship between temperament, nonresting energy expenditure, body composition, and physical activity in girls. *Int J Obes Relat Metab Disord* **28**: 300–306.
- 58) DeLany JP, Bray GA, Harsha DW, Volaufova J. 2006. Energy expenditure and substrate oxidation predict changes in body fat in children. *Am J Clin Nutr* **84**: 862–870.
- 59) DeLany JP, Bray GA, Harsha DW, Volaufova J. 2002. Energy expenditure in preadolescent African American

- and white boys and girls: the Baton Rouge Children's Study. *Am J Clin Nutr* **75**: 705–713.
- 60) Perks SM, Roemmich JN, Sandow-Pajewski M, Clark PA, Thomas E, Weltman A, Patrie J, Rogol AD. 2000. Alterations in growth and body composition during puberty. IV. Energy intake estimated by the youth-adolescent food-frequency questionnaire: validation by the doubly labeled water method. *Am J Clin Nutr* **72**: 1455–1460.
- 61) DeLany JP, Bray GA, Harsha DW, Volaufova J. 2004. Energy expenditure in African American and white boys and girls in a 2-y follow-up of the Baton Rouge Children's Study. *Am J Clin Nutr* **79**: 268–273.
- 62) Bandini LG, Schoeller DA, Dietz WH. 1990. Energy expenditure in obese and nonobese adolescents. *Pediatr Res* **27**: 198–203.
- 63) Bratteby LE, Sandhagen B, Fan H, Enghardt H, Samuelson G. 1998. Total energy expenditure and physical activity as assessed by the doubly labeled water method in Swedish adolescents in whom energy intake was underestimated by 7-d diet records. *Am J Clin Nutr* **67**: 905–911.
- 64) Arvidsson D, Slinde F, Hulthen L. 2005. Physical activity questionnaire for adolescents validated against doubly labeled water. *Eur J Clin Nutr* **59**: 376–383.
- 65) Slinde F, Arvidsson D, Sjöberg A, Rossander-Hulthén L. 2003. Minnesota leisure time activity questionnaire and doubly labeled water in adolescents. *Med Sci Sports Exerc* **35**: 1923–1928.
- 66) Ekelund U, Aman J, Yngve A, Renman C, Westerterp K, Sjöström M. 2002. Physical activity but not energy expenditure is reduced in obese adolescents: a case-control study. *Am J Clin Nutr* **76**: 935–941.
- 67) Butte NF, Wong WW, Hopkinson JM, Heinz CJ, Mehta NR, Smith EO. 2000. Energy requirements derived from total energy expenditure and energy deposition during the first 2 y of life. *Am J Clin Nutr* **72**: 1558–1569.
- 68) Tennefors C, Coward WA, Hernell O, Wright A, Forsum E. 2003. Total energy expenditure and physical activity level in healthy young Swedish children 9 or 14 months of age. *Eur J Clin Nutr* **57**: 647–653.
- 69) Davies PS, Gregory J, White A. 1995. Physical activity and body fatness in pre-school children. *Int J Obes Relat Metab Disord* **19**: 6–10.
- 70) Atkin LM, Davies PS. 2000. Diet composition and body composition in preschool children. *Am J Clin Nutr* **72**: 15–21.
- 71) Reilly JJ, Jackson DM, Montgomery C, Kelly LA, Slater C, Grant S, Paton JY. 2004. Total energy expenditure and physical activity in young Scottish children: mixed longitudinal study. *Lancet* **363**: 211–212.
- 72) Salbe AD, Weyer C, Harper I, Lindsay RS, Ravussin E, Tataranni PA. 2002. Assessing risk factors for obesity between childhood and adolescence: II. Energy metabolism and physical activity. *Pediatrics* **110**: 307–314.
- 73) Hernández-Triana M, Salazar G, Díaz E, Sánchez V, Basabe B, González S, Díaz ME. 2002. Total energy expenditure by the doubly-labeled water method in rural preschool children in Cuba. *Food Nutr Bull* **23**: 76–81.
- 74) Montgomery C, Reilly JJ, Jackson DM, Kelly LA, Slater C, Paton JY, Grant S. 2004. Relation between physical activity and energy expenditure in a representative sample of young children. *Am J Clin Nutr* **80**: 591–596.
- 75) Hoos MB, Gerver WJ, Kester AD, Westerterp KR. 2003. Physical activity levels in children and adolescents. *Int J Obes Relat Metab Disord* **27**: 605–609.
- 76) FAO. 2004. Human energy requirements. Report of a Joint FAO/WHO/UNU Expert Consultation. FAO Food and Nutrition Technical Report Series No. 1. FAO, Rome.
- 77) Butte NF, King JC. 2005. Energy requirements during pregnancy and lactation. *Public Health Nutr* **8**: 1010–1027.
- 78) Goldberg GR, Prentice AM, Coward WA, Davies HL, Murgatroyd PR, Sawyer MB, Ashford J, Black AE. 1991. Longitudinal assessment of the components of energy balance in well-nourished lactating women. *Am J Clin Nutr* **54**: 788–798.
- 79) Forsum E, Kabir N, Sadurskis A, Westerterp K. 1992. Total energy expenditure of healthy Swedish women during pregnancy and lactation. *Am J Clin Nutr* **56**: 334–342.
- 80) Goldberg GR, Prentice AM, Coward WA, Davies HL, Murgatroyd PR, Wensing C, Black AE, Harding M, Sawyer M. 1993. Longitudinal assessment of energy expenditure in pregnancy by the doubly labeled water method. *Am J Clin Nutr* **57**: 494–505.
- 81) Kopp-Hoolihan LE, van Loan MD, Wong WW, King JC. 1999. Longitudinal assessment of energy balance in well-nourished, pregnant women. *Am J Clin Nutr* **69**: 697–704.
- 82) Butte NF, Wong WW, Treuth MS, Ellis KJ, O'Brian Smith E. 2004. Energy requirements during pregnancy based on total energy expenditure and energy deposition. *Am J Clin Nutr* **79**: 1078–1087.
- 83) Takimoto H, Sugiyama T, Fukuoka H, Kato N, Yoshiike N. 2006. Maternal weight gain ranges for optimal fetal growth in Japanese women. *Int J Gynecol Obstet* **92**: 272–278.
- 84) Butte NF, Wong WW, Hopkinson JM. 2001. Energy requirements of lactating women derived from doubly labeled water and milk energy output. *J Nutr* **131**: 53–58.
- 85) Suzuki K, Sasaki A, Shinzawa K, Totani M. 2004. Milk intake by breast-fed infants before weaning. *Jpn J Nutr Diet* **62**: 369–372 (in Japanese).
- 86) Hirose J, Endo M, Nagao S, Mizushima K, Narita H, Shibata K. 2008. Amount of breast milk sucked by Japanese breast feeding infants. *J Jpn Soc Breastfeed Res* **2**: 23–28.
- 87) Yamawaki N, Yamada M, Kan-no T, Kojima T, Kaneko T, Yonekubo A. 2005. Macronutrient, mineral and trace element composition of breast milk from Japanese women. *J Trace Elements Med Biol* **19**: 171–181.
- 88) Tanaka S, Ohkawara K, Ishikawa-Takata K, Morita A, Watanabe S. 2008. Accuracy of predictive equations for basal metabolic rate and contribution of abdominal fat distribution to basal metabolic rate in obese Japanese people. *Anti-Aging Med* **5**: 17–21.
- 89) Takahashi E, Higuchi M, Hosokawa Y, Tabata I. 2007. Basal metabolic rate and body composition of Japanese young adult females. *Jpn J Nutr Diet* **65**: 241–247 (in Japanese).
- 90) Ainsworth BE, Haskell WL, Whitt MC, Irwin ML, Swartz AM, Strath SJ, O'Brien WL, Bassett DR Jr, Schmitz KH, Emplainscourt PO, Jacobs DR Jr, Leon AS. 2000. Compendium of physical activities: an update of activity codes and MET intensities. *Med Sci Sports Exerc* **32**: S498–516.
- 91) Ohkawara K, Tanaka S, Ishikawa-Takata K, Tabata I. 2008. Twenty-four-hour analysis of elevated energy expenditure after physical activity in a metabolic chamber: models of daily total energy expenditure. *Am J Clin Nutr* **87**: 1268–1276.

