

<http://www.cdc.gov/tularemia/resources/whotularemiamanual.pdf> are appropriate for Japanese tularemia patients. Some β -lactams might also be effective for some patients in Japan. These findings provide important information related to recommendations for post-exposure prophylaxis and therapy for tularemia patients in Japan.

Acknowledgements

This study was supported by a Health Science Research Grants from the Ministry of Health, Labour and Welfare of Japan (H22-Shinkou-Ippan-010 and H23-Shinkou-Wakate-022). N.S. was supported by a Honjo International Scholarship Foundation.

Conflict of interest

None to declare

References

1. Ellis, J., Oyston, P.C., Green, M., et al. (2002): Tularemia. Clin. Microbiol. Rev., 15, 631-646
2. Sjostedt, A. (2007): Tularemia: history, epidemiology, pathogen physiology, and clinical manifestations. Ann. NY Acad. Sci., 1105 1-29
3. Ohara, Y., Sato, T. and Homma, M. (1996): Epidemiological analysis of tularemia in Japan (yato-byo). FEMS Immunol. Med. Microbiol., 13, 185-189
4. Park, C.H., Nakanishi, A., Hatai, H., et al. (2009): Pathological and microbiological studies of Japanese Hare (*Lepus brachyurus angustidens*) naturally infected with *Francisella tularensis* subsp. *holarctica*. J. Vet. Med. Sci., 71, 1629-1635
5. Fujita, O., Uda, A., Hotta, A., et al. (2008): Genetic diversity of *Francisella tularensis* subspecies *holarctica* strains isolated in Japan. Microbiol. Immunol., 52, 270-276
6. Johansson, A., Urich, S.K., Chu, M.C., et al. (2002): In vitro susceptibility to quinolones of *Francisella tularensis* subspecies *tularensis*. Scand. J. Infect. Dis., 34, 327-330
7. Tomaso, H., Al Dahouk, S., Hofer, E., et al. (2005): Antimicrobial susceptibilities of Austrian *Francisella tularensis holarctica* biovar II strains. Int. J Antimicrob. Agents., 26, 279-284

8. Valade, E., Vaissaire, J., Merens, A., et al. (2008): Susceptibility of 71 French isolates of *Francisella tularensis* subsp. *holarctica* to eight antibiotics and accuracy of the Etest method. *J. Antimicrob. Chemother.*, 62, 208-210
9. Yesilyurt, M., Kilic, S., Celebi, B., et al. (2011): Antimicrobial susceptibilities of *Francisella tularensis* subsp. *holarctica* strains isolated from humans in the Central Anatolia region of Turkey. *J. Antimicrob. Chemother.*, 66, 2588-2592
10. Hotta, A., Uda, A., Fujita, O., et al. (2007): Preparation of monoclonal antibodies for detection and identification of *Francisella tularensis*. *Clin. Vaccine Immunol.*, 14, 81-84
11. Baker, C.N., Hollis, D.G. and Thornsberry, C. (1985): Antimicrobial susceptibility testing of *Francisella tularensis* with a modified Mueller-Hinton broth. *J. Clin. Microbiol.*, 22, 212-215
12. Markowitz, L.E., Hynes, N.A., De La Cruz, P., et al. (1985): Tick-borne tularemia. An outbreak of lymphadenopathy in children. *JAMA*, 254, 2922-2925
13. Pfeifer, Y., Cullik, A. and Witte, W. (2010): Resistance to cephalosporins and carbapenems in Gram-negative bacterial pathogens. *Int. J. Med. Microbiol.*, 300, 371-379

