

Table 1. Baseline Demographic and Disease Characteristics

Parameter	Value
N	10
Age, median years (range)	62 (52-70)
Male sex, n (%)	4 (40)
Japanese race, n (%)	10 (100)
Host <i>IL28B</i> genotype,* n (%)	
CC	2 (20)
CT	8 (80)
HCV genotype 1b, n (%)	10 (100)
HCV RNA, mean log ₁₀ IU/mL (SD)	6.8 (0.61)
ALT, mean U/L (SD)	60.6 (32.9)
Platelets × 10 ⁹ cells/mL, median (min, max)	150.5 (84.0, 166.0)
Total bilirubin, median mg/dL (min, max)	0.8 (0.6, 1.2)
Albumin, median g/dL (min, max)	3.9 (3.1, 4.2)
INR, median (min, max)	1.0 (1.0, 1.1)

*SNP rs12979860.

Abbreviation: *IL28B*, interleukin-28B; HCV, hepatitis C virus; SD, standard deviation; ALT, alanine aminotransferase; min, minimum; max, maximum; INR, international normalized ratio; SNP, single-nucleotide polymorphism.

were female and all were Japanese (Table 1). All enrolled patients were infected with genotype 1b, reflecting the predominance of this subtype in Japan, although the study protocol did not exclude patients with HCV genotype 1a.⁶ Two patients were *IL28B* genotype CC (SNP rs12979860) and 8 were CT. Nine patients completed 24 weeks of therapy; 1 patient discontinued at week 2 because of a grade 4 total bilirubin elevation (see below). Among the 9 patients treated for 24 weeks, asunaprevir was dosed at 600 mg twice-daily for 12-21 weeks before the dose was reduced to 200 mg twice-daily (Fig. 1).

Virologic Response. Serum HCV RNA levels decreased rapidly in all patients (Fig. 2); mean reductions from baseline were 4.4 log₁₀ IU/mL at week 1,




Fig. 1. ALT levels: individual patients. Serum ALT levels for the 9 patients who completed 24 weeks of treatment; the patient who discontinued at week 2 is not presented. Shaded area indicates the treatment period; arrows indicate the points at which the dose of asunaprevir was reduced from 600 to 200 mg twice-daily. Arrow and line colors are the same for each patient.




Fig. 2. HCV RNA levels: individual patients. Individual patient plasma HCV RNA levels during 24 weeks of treatment and through 24 weeks post-treatment (week 48) are shown. LLQ = 15 IU/mL.

5.3 log₁₀ IU/mL at week 2, and 5.8 log₁₀ IU/mL from week 4 through the end of treatment. At week 4, HCV RNA was undetectable (RVR) in 4 of 10 (40%) patients and below the assay LLQ in 9 of 10 (90%; Fig. 3). No patients qualified for discontinuation or addition of pegIFN/RBV. At week 8, HCV RNA was undetectable in 9 of 10 patients (all who remained on treatment) and remained undetectable through the end of treatment and follow-up. SVR₁₂, the primary endpoint, and SVR₂₄ were achieved by 90% of patients, including all 9 who completed 24 weeks of therapy. The patient who discontinued treatment at week 2 had low-level HCV RNA at discontinuation (1.8 log₁₀ IU/mL), but HCV RNA was undetectable at follow-up visits 2, 3, 4, 13, and 24 weeks after discontinuation.

Viral Breakthrough and Relapse. There was no viral breakthrough during treatment or relapse of HCV RNA post-treatment. Analysis of baseline samples revealed variants reported to confer minimal to low




Fig. 3. HCV RNA endpoints. Categorical HCV RNA endpoints are indicated for the 10 study patients. One patient discontinued at week 2 and was counted as a treatment failure at the time points shown. However, HCV RNA was undetectable in this patient at 2, 3, 13, and 24 weeks post-treatment.

Table 2. On-Treatment Adverse Events Occurring in ≥ 2 Patients

Event	Patients, n (%)
Diarrhea	7 (70)
Headache	4 (40)
ALT increased	3 (30)
AST increased	3 (30)
Lymphopenia	2 (20)
Abdominal discomfort	2 (20)
Malaise	2 (20)
Pyrexia	2 (20)
Nasopharyngitis	2 (20)
Lipase increased	2 (20)
Back pain	2 (20)

Abbreviations: ALT, alanine aminotransferase; AST, aspartate aminotransferase.

levels of resistance to daclatasvir.²² NS5A substitutions L28M and L31M were detected in 1 patient each, and Y93H was detected in 2 other patients. NS3 protease substitutions reported to confer resistance to telaprevir, boceprevir, and TMC-435 were detected²⁵; T54S was identified in 1 patient, and Q80L was identified in 3. In 1 patient, both NS3 protease substitutions (T54S and Q80L) and an NS5A substitution (Y93H) were detected. There was no consistent association between detection of these variants and virologic outcomes.

Safety. The most frequently reported adverse events were diarrhea and headache, all of which were mild (grade 1) (Table 2). The patient who discontinued (see below) experienced multiple grade 3 or 4 adverse events and laboratory abnormalities on treatment. In the other 9 patients, there were no grade 3 or 4 transaminase elevations or other grade 3 or 4 events, no clinically relevant changes in electrocardiogram parameters, and no lymphopenia of any severity. Two transient grade 1 ALT elevations were reported, and 1 grade 2 elevation that began at week 16 and persisted until the end of treatment, after which it normalized within 2 weeks (Fig. 1). There were no notable differences in ALT before and after asunaprevir dose reduction.

There were two serious adverse events. A 54-year-old male was hospitalized with grade 3 pyrexia and persistent diarrhea 11 days after initiating study treatment. Loxoprofen was initiated, and body temperature normalized and diarrhea improved after 4 days. The patient remained on study treatment. The second event concerned a 60-year-old woman with a history of ulcerative colitis who discontinued study treatment after 2 weeks because of a grade 4 bilirubin elevation with multiple complicating features. Five days before discontinuation, she presented with infectious gastro-

enteritis and was treated with cefotiam and was subsequently hospitalized with fever, vomiting, and diarrhea. Meropenem, human serum albumin, and furosemide were initiated. At discontinuation of study drugs, laboratory findings included total bilirubin of 7.7 mg/dL and grade 3 lymphopenia and serum phosphorus reduction; transaminases and alkaline phosphatase were within normal ranges. In the week after discontinuation, white cell and eosinophil counts became elevated; total bilirubin improved and transaminases remained normal. Two weeks after discontinuation, grade 4 ALT and aspartate-aminotransferase elevations and a grade 3 lipase elevation were reported. Six weeks after discontinuation, bilirubin and transaminase elevations were resolved and lipase improved to within $2\times$ ULN.

Discussion

This study assessed combination oral DAA therapy in a difficult-to-treat population with multiple adverse prognostic features, including HCV genotype 1b infection, primarily *IL28B* CT genotype, generally older age, and null response to previous Peg-IFN/RBV therapy.^{10,13,14} These patients represent a group with a significant need for new therapeutic options.

A DAA-only therapeutic strategy may be particularly appropriate for null responders, who have previously shown only marginal response to Peg-IFN/RBV.^{13,14} The combination of two highly potent DAAs cleared detectable virus rapidly in this study; HCV RNA was undetectable by week 8 in all 9 patients treated for 24 weeks. This outcome compares favorably with those observed when null responders received a combination of Peg-IFN/RBV and a single NS3 protease inhibitor, telaprevir or TMC435.^{15,26} In these studies, HCV RNA remained detectable in 36% to approximately 50% of patients after 12 weeks.

HCV RNA remained undetectable 12 (SVR₁₂) and 24 weeks (SVR₂₄) post-treatment in all patients who completed treatment. This contrasts with the poor results obtained with Peg-IFN/RBV retreatment and the reported 37% SVR rate of genotype 1b null responders who received Peg-IFN/RBV and telaprevir.^{10,13-15} Additional follow-up of patients from this study will assess whether SVR₂₄ is predictive of long-lasting viral clearance with this dual DAA therapy, as it is with Peg-IFN/RBV. It is interesting that HCV RNA was persistently undetectable post-treatment in the patient who discontinued after only 2 weeks of treatment. With early discontinuation data from only this single case, at present, the result must be considered an anomaly. The factors that contributed to viral

clearance are uncertain, although the patient's *IL28B* CC genotype suggests increased sensitivity to endogenous interferon²⁷; the possible influence of concurrent acute gastroenteritis or other complicating factors is unknown. However, coupled with the attainment of SVR₁₂ in all other patients, this outcome suggests that required duration of therapy, which is currently predicated on data from Peg-IFN-based regimens, may need reassessment for DAA-only regimens, and, possibly, that certain patient populations can be treated for very short durations.

The high SVR rate is consistent with limited data from a related U.S.-based study, in which 2 of 2 null responders with HCV genotype 1b and who were treated with daclatasvir and asunaprevir achieved SVR₂₄.²³ However, only 2 of 9 patients with genotype 1a achieved SVR₂₄ with the dual DAA regimen, compared with 9 of 10 patients who received both DAAs and Peg-IFN/RBV. These differences suggest that viral genotype can influence responses to DAA regimens that do not include Peg-IFN/RBV, and outcomes can be optimized with individualized therapy that considers viral genotype, among other factors. Because of the high SVR rate, the potential influence of other baseline and on-treatment parameters could not be assessed, other than to observe that unfavorable predictors of Peg-IFN/RBV response, such as older age and *IL28B* CT genotype,^{27,28} had no measureable impact on outcomes.

There was no viral breakthrough on treatment. In view of the rapid emergence of resistance in some studies of short-term DAA monotherapy,^{29,30} these findings support the concept that dual DAA therapy reduces the risk of viral breakthrough, in addition to increasing antiviral activity. Resistance analyses revealed that before treatment, some patients carried NS5A and NS3 polymorphisms predicted to reduce sensitivity to daclatasvir and some HCV protease inhibitors, respectively.^{22,25} There was no clear relationship between the presence of these polymorphisms and minor interpatient differences in the rate of early virologic response; however, further study in larger patient cohorts will help determine whether baseline polymorphisms can influence virologic response with this regimen.

The adverse event profile of the dual DAA regimen compares favorably with the more frequent and severe events reported with Peg-IFN/RBV, although patient numbers in this study were limited. The mild diarrhea experienced by several patients has been reported previously with asunaprevir and is common with other drugs of this class.^{15,18,24} Though a role

for daclatasvir and/or asunaprevir in the two serious adverse events could not be ruled out and the investigator considered these events drug related, multiple confounding factors existed. The case of pyrexia was consistent with a viral infection and resolved with treatment. In the case of hyperbilirubinemia that led to discontinuation, the time course of laboratory abnormalities and related events suggests a link to the use of cefotiam and meropenem for treatment of infectious gastroenteritis. Both of these agents have been associated with vomiting, diarrhea, and hyperbilirubinemia.^{31,32}

The asunaprevir dose was reduced during treatment because of transaminase elevations observed with 600 mg twice-daily in a concurrent study.²⁴ In this sentinel cohort, viral suppression was maintained in all patients after dose reduction, and no grade 3 or 4 transaminase elevations occurred during treatment at either dose of asunaprevir. One patient experienced grade 2 transaminase elevations that began at week 16 and persisted during treatment, despite asunaprevir dose reduction at week 19. Although these elevations were not severe, their rapid normalization post-treatment suggests a possible relationship to study treatment. None of the 9 patients treated for 24 weeks experienced transaminase elevations post-treatment. Although grade 4 transaminase elevations occurred 2 weeks post-treatment in the patient who discontinued, the timing of these events and multiple other complications suggest that they were not related directly to study treatment.

In conclusion, the combination of daclatasvir and asunaprevir achieved a high rate of SVR₂₄ in patients with HCV genotype 1b infections and previous null response to Peg-IFN/RBV. These results support the concept that HCV infection can be cured with two DAAs without Peg-IFN/RBV, even in difficult-to-treat populations that lack robust IFN responsiveness. Further research will assess the benefits of DAA combinations in larger, more diverse patient populations.

Acknowledgment: The authors thank the patients and their families, research staff at all participating sites, and Bristol-Myers Squibb Research and Development colleagues.

References

1. Armstrong GL, Wasley A, Simard EP, McQuillan GM, Kuhnert WL, Alter MJ. The prevalence of hepatitis C virus infection in the United States, 1999 through 2002. *Ann Intern Med* 2006;144:705-714.
2. World Health Organization. Global alert and response (GAR): hepatitis C. 2011 (January 26). Available at: <http://www.who.int/csr/disease/hepatitis/whocdscrlyo2003/en/index4.html>. Accessed on June 29, 2011.

3. Legrand-Abrevanel F, Colson P, Leguillou-Guillemette H, Alric L, Ravaux I, Lunel-Fabiani F, et al. Influence of the HCV subtype on the virological response to pegylated interferon and ribavirin therapy. *J Med Virol* 2009;81:2029-2035.
4. Nicot F, Alric L, Barange K, Metivier S, Dramard JM, Combis JM, et al. Influence of HCV genotype 1 subtypes on the virus response to PEG interferon alpha-2a plus ribavirin therapy. *J Med Virol* 2011;83:437-444.
5. Cornberg M, Razavi HA, Alberti A, Bernasconi E, Buti M, Cooper C, et al. A systematic review of hepatitis C virus epidemiology in Europe, Canada, and Israel. *Liver Int* 2011;31(Suppl 2):30-60.
6. Sievert W, Altraif I, Razavi HA, Abdo A, Ahmed EA, Alomair A, et al. A systematic review of hepatitis C virus epidemiology in Asia, Australia, and Egypt. *Liver Int* 2011;31(Suppl 2):61-80.
7. Negro F, Alberti A. The global health burden of hepatitis C virus infection. *Liver Int* 2011;31(Suppl 2):1-3.
8. Fried MW, Shiffman ML, Reddy KR, Smith C, Marinos G, Goncalves FL, Jr., et al. Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 2002;347:975-982.
9. McHutchison JG, Lawitz EJ, Shiffman ML, Muir AJ, Galler GW, McCone J, et al. Peginterferon alfa-2b or alfa-2a with ribavirin for treatment of hepatitis C infection. *N Engl J Med* 2009;361:580-593.
10. Ghany MG, Strader DB, Thomas DL, Seeff LB; American Association for the Study of Liver Diseases. Diagnosis, management, and treatment of hepatitis C: an update. *HEPATOLOGY* 2009;49:1335-1374.
11. Poordad F, McCone J Jr., Bacon BR, Bruno S, Manns MP, Sulkowski MS, et al. Boceprevir for untreated chronic HCV genotype 1 infection. *N Engl J Med* 2011;364:1195-1206.
12. Jacobson IM, McHutchison JG, Dusheiko G, Di Bisceglie AM, Reddy KR, Bzowej NH, et al. Telaprevir for previously untreated chronic hepatitis C virus infection. *N Engl J Med* 2011;364:2405-2416.
13. Poynard T, Colombo M, Bruix J, Schiff E, Terg R, Flamm S, et al. Peginterferon alfa-2b and ribavirin: effective in patients with hepatitis C who failed interferon alfa/ribavirin therapy. *Gastroenterology* 2009;136:1618-1628.
14. Jensen DM, Marcellin P, Freilich B, Andreone P, Di Bisceglie A, Brandao-Mello CE, et al. Re-treatment of patients with chronic hepatitis C who do not respond to peginterferon-alpha2b: a randomized trial. *Ann Intern Med* 2009;150:528-540.
15. Zeuzem S, Andreone P, Pol S, Lawitz E, Diago M, Roberts S, et al. Telaprevir for retreatment of HCV infection. *N Engl J Med* 2011;364:2417-2428.
16. Soriano V, Peters MG, Zeuzem S. New therapies for hepatitis C virus infection. *Clin Infect Dis* 2009;48:313-320.
17. Ohara E, Hiraga N, Imamura M, Iwao E, Kamiya N, Yamada I, et al. Elimination of hepatitis C virus by short term NS3-4A and NS5B inhibitor combination therapy in human hepatocyte chimeric mice. *J Hepatol* 2011;54:872-878.
18. Gane EJ, Roberts SK, Stedman CA, Angus PW, Ritchie B, Elston R, et al. Oral combination therapy with a nucleoside polymerase inhibitor (RG7128) and danoprevir for chronic hepatitis C genotype 1 infection (INFORM-1): a randomised, double-blind, placebo-controlled, dose-escalation trial. *Lancet* 2010;376:1467-1475.
19. Gao M, Nettles RE, Belema M, Snyder LB, Nguyen VN, Fridell RA, et al. Chemical genetics strategy identifies an HCV NS5A inhibitor with a potent clinical effect. *Nature* 2010;465:96-100.
20. McPhee F, Levesque PC, Li D, Zhu J, Friborg J, Sheaffer A, et al. Identification and preclinical profile of the novel HCV NS3 protease inhibitor BMS-650032 [abstract]. *J Hepatol* 2010;52(Suppl 1):S296.
21. Bifano M, Sevinsky H, Bedford BR, Coumbis J, Eley T, Huang SP, et al. Coadministration of BMS-790052 and BMS-650032 does not result in a clinically meaningful pharmacokinetic interaction in healthy subjects [abstract]. *HEPATOLOGY* 2010;52(Suppl):719A.
22. Fridell RA, Qiu D, Wang C, Valera L, Gao M. Resistance analysis of the hepatitis C virus NS5A inhibitor BMS-790052 in an in vitro replication system. *Antimicrob Agents Chemother* 2010;54:3641-3650.
23. Lok A, Gardiner D, Lawitz E, Martorell C, Everson G, Ghalib R, et al. Quadruple therapy with BMS-790052, BMS-650032, and peg-IFN/RBV for 24 weeks results in 100% SVR12 in HCV genotype 1 null responders [abstract]. *J Hepatol* 2011;54:S536.
24. Bronowicki JP, Pol S, Thuluvath PJ, Larrey D, Martorell CT, Rustgi VK, et al. BMS-650032, an NS3 inhibitor, in combination with peginterferon alfa-2a and ribavirin in treatment-naive subjects with genotype 1 chronic hepatitis C infection [abstract]. *J Hepatol* 2011;54:S472.
25. Romano KP, Ali A, Royer WE, Schiffer CA. Drug resistance against HCV NS3/4A inhibitors is defined by the balance of substrate recognition versus inhibitor binding. *Proc Natl Acad Sci U S A* 2010;107:20986-20991.
26. Zeuzem S, Foster GR, Fried MW, Hezode C, Hirschfield GM, Nikitin I, et al. The ASPIRE trial: TMC435 in treatment-experienced patients with genotype-1 HCV infection who have failed previous pegIFN/RBV treatment [abstract]. *J Hepatol* 2011;54:S546.
27. Thompson AJ, Muir AJ, Sulkowski MS, Ge D, Fellay J, Shianna KV, et al. Interleukin-28B polymorphism improves viral kinetics and is the strongest pretreatment predictor of sustained virologic response in genotype 1 hepatitis C virus. *Gastroenterology* 2010;139:120-129.
28. Manns MP, McHutchison JG, Gordon SC, Rustgi VK, Shiffman M, Reindollar R, et al. Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomised trial. *Lancet* 2001;358:958-965.
29. Sarrazin C, Kieffer TL, Bartels D, Hanzelka B, Muh U, Welker M, et al. Dynamic hepatitis C virus genotypic and phenotypic changes in patients treated with the protease inhibitor telaprevir. *Gastroenterology* 2007;132:1767-1777.
30. Susser S, Welsch C, Wang Y, Zettler M, Domingues FS, Karey U, et al. Characterization of resistance to the protease inhibitor boceprevir in hepatitis C virus-infected patients. *HEPATOLOGY* 2009;50:1709-1718.
31. AstraZeneca. Merrem (meropenem) IV prescribing information. 2010. Available at: <http://www1.astrazeneca-us.com/pi/MerremIV.pdf>. Accessed on June 29, 2011.
32. Imada A, Hirai S. Cefotiam hexetil. *Int J Antimicrob Agents* 1995;5:85-99.

Amino Acid Substitution in HCV Core/NS5A Region and Genetic Variation Near *IL28B* Gene Affect Treatment Efficacy to Interferon plus Ribavirin Combination Therapy

Norio Akuta^a Fumitaka Suzuki^a Miharuru Hirakawa^a Yusuke Kawamura^a
Hitomi Sezaki^a Yoshiyuki Suzuki^a Tetsuya Hosaka^a Masahiro Kobayashi^a
Mariko Kobayashi^b Satoshi Saitoh^a Yasuji Arase^a Kenji Ikeda^a
Kazuaki Chayama^c Yusuke Nakamura^d Hiromitsu Kumada^a

^aDepartment of Hepatology, and ^bLiver Research Laboratory, Toranomon Hospital, Tokyo, ^cDepartment of Medical and Molecular Science, Division of Frontier Medical Science, Programs for Biomedical Research, Graduate School of Biomedical Science, Hiroshima University, Hiroshima, and ^dLaboratory of Molecular Medicine, Human Genome Center, Institute of Medical Science, University of Tokyo, Tokyo, Japan

Key Words

Hepatitis C virus · Interferon · Ribavirin · Core region · NS5A region · ISDR · IRRDR · *IL28B*

Abstract

Objective: To evaluate predictive factors of treatment efficacy to interferon (IFN)/ribavirin in patients infected with HCV genotype 1b (HCV-1b). **Methods:** This study investigated pretreatment predictors, including viral- (aa substitutions in core aa 70/91 and NS5A-ISDR/IRRDR) and host-related factors (genetic variation near *IL28B* gene), to 48-week IFN/ribavirin in 490 Japanese adults infected with HCV-1b. **Results:** The proportion of patients who showed end-of-treatment response (ETR), sustained virological response (SVR), and SVR after ETR was 76, 54, and 76%, respectively. There was a significant positive correlation between the number of aa substitutions in ISDR and those in IRRDR. Concerning the substitution of core aa 91, the number of aa substitutions in ISDR/IRRDR of patients with Leu91 was significantly higher

than that of patients with Met91. Furthermore, levels of viremia were influenced by aa substitutions in core aa 91 and ISDR/IRRDR. By multivariate analysis, rs8099917 genotype was an important predictor of ETR and SVR. With regard to viral factors, core aa 70/91 was an important predictor of ETR, and SVR after ETR. ISDR was an important predictor of SVR, and SVR after ETR. **Conclusion:** aa substitution in core/NS5A region and genetic variation near *IL28B* were important predictors of treatment efficacy to IFN/ribavirin.

Copyright © 2011 S. Karger AG, Basel

Introduction

Treatment of chronic hepatitis C virus (HCV) infection with interferon (IFN) combined with ribavirin carries potential serious side effects and is costly, especially when used long enough to achieve a high sustained virological response (SVR) in patients infected with HCV genotype 1b (HCV-1b) and high viral loads. For these rea-

KARGER

Fax +41 61 306 12 34
E-Mail karger@karger.ch
www.karger.com

© 2011 S. Karger AG, Basel
0300-5526/12/0553-0231\$38.00/0

Accessible online at:
www.karger.com/int

Norio Akuta, MD
Department of Hepatology, Toranomon Hospital
2-2-2 Toranomon, Minato-ku
Tokyo 105-0001 (Japan)
Tel. +81 44 877 5111, E-Mail akuta-gi@umin.ac.jp

sons, those patients who do not achieve SVR need to be identified, so as to free them of unnecessary side effects and reduce costs, preferably before the start of the combination therapy.

Viral- and host-related factors are useful as predictors of treatment efficacy to 48-week IFN/ribavirin combination therapy. With regard to viral factors, amino acid (aa) substitutions at position 70 and/or 91 in the core region of HCV-1b are pretreatment predictors of virological response to combination therapy [1–4], and also affect clinical outcome, including hepatocarcinogenesis [5, 6]. Furthermore, the NS5A region of HCV-1b, including IFN-sensitivity-determining region (ISDR) [7, 8] and IFN/ribavirin resistance-determining region (IRRD) [9, 10], are also useful as pretreatment predictors of virological response to combination therapy [11, 12]. With regard to host factors, genetic variations near *IL28B* gene (rs8099917, rs12979860) on chromosome 19, which encodes IFN- λ -3, are pretreatment predictors of virological response to combination therapy in individuals infected with HCV-1 [13–16], and also affect clinical outcome, including spontaneous clearance of HCV [17]. A recent report identified genetic variation near *IL28B* gene and aa substitution of the core region as predictors of SVR to triple therapy of telaprevir/pegylated (PEG)-IFN/ribavirin in Japanese patients infected with HCV-1b [18]. However, to our knowledge, there are no previous reports of IFN/ribavirin combination therapy based on multivariate analysis to investigate pretreatment predictors, including all of aa substitutions in core aa 70/91 and NS5A-ISDR/IRRD, and genetic variation near *IL28B* gene.

The aim of the present study was to investigate predictive factors of treatment efficacy, including viral- (aa substitutions in core aa 70/91 and NS5A-ISDR/IRRD) and host-related factors (genetic variation near *IL28B* gene), to 48-week IFN/ribavirin in Japanese adults infected with HCV-1b.

Patients and Methods

Study Population

A total of 1,249 HCV-1b-infected Japanese adult patients were consecutively recruited into the study protocol of combination therapy with IFN (PEG-IFN α -2b or IFN α -2b) plus ribavirin between December 2001 and January 2009 at Toranomon Hospital, Tokyo, Japan. Among these, 490 patients, who could complete a total of 48 weeks of combination therapy, were enrolled in this retrospective study, and fulfilled the following criteria: (1) negativity for hepatitis B surface antigen (HBsAg) in serum; (2) HCV-1b only confirmed by sequence analysis; (3) HCV-RNA levels of ≥ 5.0 log IU/ml determined by the COBAS TaqMan HCV test

(Roche Diagnostics, Tokyo, Japan) within the preceding 2 months of enrolment; (4) no hepatocellular carcinoma; (5) body weight >40 kg; (6) lack of coinfection with human immunodeficiency virus; (7) no previous treatment with antiviral or immunosuppressive agents within the preceding 3 months of enrolment; (8) none was an alcoholic; lifetime cumulative alcohol intake was <500 kg; (9) none had other forms of liver diseases, such as hemochromatosis, Wilson disease, primary biliary cirrhosis, alcoholic liver disease, or autoimmune liver disease, and (10) none of the females was pregnant or breastfeeding.

The study protocol was in compliance with the Good Clinical Practice Guidelines and the 1975 Declaration of Helsinki, and was approved by the institutional review board. Each patient gave their informed consent before participating in this trial.

The treatment efficacy was evaluated in terms of HCV-RNA negativity at the end of treatment (end-of-treatment response (ETR)) and 24 weeks after the completion of therapy (SVR), based on the COBAS TaqMan HCV test (Roche Diagnostics). SVR in patients who achieved ETR was defined as SVR after ETR. ETR, SVR, and SVR after ETR could be evaluated in 487 (99%), 448 (91%), and 321 (66%) of 490 patients, respectively.

422 (86%) patients received PEG-IFN α -2b at a median dose of 1.4 μ g/kg (range 0.7–1.9) subcutaneously each week plus oral ribavirin at a median dose of 11.1 mg/kg (range 3.7–15.1) daily for 48 weeks. The remaining 68 (14%) patients received 6 million units of IFN α -2b intramuscularly each day for 48 weeks (daily for the initial 2 weeks, followed by three times per week for 46 weeks), and oral ribavirin at a median dose of 11.3 mg/kg (range 6.8–13.4) daily for 48 weeks.

Table 1 summarizes the profiles and laboratory data of the 490 patients at the commencement of treatment. They included 310 males and 180 females aged 20–75 years (median 54).

Measurement of HCV RNA

The antiviral effects of treatment on HCV were assessed by measuring plasma HCV-RNA levels. In this study, HCV-RNA levels were evaluated at least once every month before, during, and after therapy. HCV-RNA concentrations were determined using the COBAS TaqMan HCV test (Roche Diagnostics). The linear dynamic range of the assay was 1.2–7.8 log IU/ml, and the undetectable samples were defined as negative.

Detection of aa Substitutions in Core, and NS5A Regions of HCV-1b

With the use of HCV-J (accession No. D90208) as a reference [19], the sequence of 1–191 aa in the core protein of HCV-1b was determined and then compared with the consensus sequence constructed on the previous study to detect substitutions at aa 70 of arginine (Arg70) or glutamine/histidine (Gln70/His70) and aa 91 of leucine (Leu91) or methionine (Met91) [1]. The sequence of 2,209–2,248 aa in the NS5A of HCV-1b (ISDR) reported by Enomoto et al. [7, 8] was determined, and the number of aa substitutions in ISDR was defined as wild-type (WT) (0, 1) or non-wild-type (non-WT) (≥ 2) in comparison with HCV-J. Furthermore, the sequence of 2,334–2,379 aa in the NS5A of HCV-1b (IRRD) reported by El-Shamy et al. [9, 10] was determined and then compared with the consensus sequence constructed on the previous study. In the present study, aa substitutions of the core region and NS5A-ISDR/IRRD of HCV-1b were analyzed by direct sequencing [10, 18].

Genetic Variation near *IL28B* Gene

Samples for genome-wide association survey were genotyped using the Illumina HumanHap610-Quad Genotyping BeadChip. Genotyping data were subjected to quality control before the data analysis. Genotyping for replication and fine mapping was performed by use of Invader assay, TaqMan assay, or direct sequencing as described previously [20, 21].

In this study, genetic variations near *IL28B* gene (rs8099917), reported as the pretreatment predictors of treatment efficacy in Japanese patients [14, 18], were investigated.

Statistical Analysis

Non-parametric tests (Mann-Whitney U test, χ^2 test and Fisher's exact probability test) were used to compare the characteristics of the groups. Correlation analysis was evaluated by the Spearman rank correlation test. Uni- and multivariate logistic regression analyses were used to determine those factors that significantly contributed to ETR, SVR, and SVR after ETR. The odds ratios (OR) and 95% confidence intervals (95% CI) were also calculated. All *p* values <0.05 by the two-tailed test were considered significant. Variables that achieved statistical significance (*p* < 0.05) on univariate analysis were entered into multiple logistic regression analysis to identify significant independent predictive factors. Each variable was transformed into categorical data consisting of two simple ordinal numbers for uni- and multivariate analyses. Potential predictive factors associated with ETR, SVR, and SVR after ETR included the following variables: sex, age, history of blood transfusion, familial history of liver disease, body mass index, aspartate aminotransferase (AST), alanine aminotransferase (ALT), albumin, γ -glutamyl transpeptidase (GGT), leukocyte count, hemoglobin, platelet count, level of viremia, α -fetoprotein, total cholesterol, high-density lipoprotein cholesterol, low-density lipoprotein cholesterol, triglycerides, uric acid, ribavirin dose/body weight, genetic variation near *IL28B* gene, and aa substitution in the core region, and NS5A-ISDR/IRRDR. Statistical analyses were performed using SPSS software (SPSS Inc., Chicago, Ill., USA).

Results

Response to Therapy

ETR was achieved by 372 of 487 (76%) patients, SVR by 244 of 448 (54%), and SVR after ETR by 244 of 321 (76%).

Number of aa Substitutions in NS5A-ISDR and NS5A-IRRDR

As a whole, 0, 1, and ≥ 2 aa substitutions in ISDR were found in 56% (227 of 406), 23% (95 of 406), and 21% (84 of 406) of patients, respectively. Thus, the percentage of patients with ≤ 1 aa substitution in ISDR (WT) was 79% (322 of 406). Furthermore, ≤ 3 , 4–5, and ≥ 6 aa substitutions in IRRDR were found in 36% (73 of 200), 34% (67 of 200), and 30% (60 of 200) of patients, respectively (fig. 1).

Table 1. Patient profile and laboratory data at commencement of the 48-week combination therapy of IFN + ribavirin in 490 patients infected with HCV-1b

<i>Demographic data</i>	
Number of patients	490
Male/female	310/180
Age, years	54 (20–75)
History of blood transfusion	169 (34%)
Family history of liver disease	96 (20%)
Body mass index, kg/m ²	22.6 (15.7–34.7)
<i>Laboratory data</i>	
Level of viremia, log IU/ml	6.4 (2.2–7.7)
Serum AST, IU/l	50 (16–296)
Serum ALT, IU/l	67 (12–836)
Serum albumin, g/dl	3.9 (3.1–4.7)
GGT, IU/l	44 (10–592)
Leukocyte count, n/mm ³	4,700 (1,200–10,900)
Hemoglobin, g/dl	14.4 (10.6–18.1)
Platelet count, $\times 10^4$ /mm ³	16.7 (6.4–37.5)
α -Fetoprotein, μ g/l	5 (1–459)
Total cholesterol, mg/dl	170 (96–284)
High-density lipoprotein cholesterol, mg/dl	46 (13–95)
Low-density lipoprotein cholesterol, mg/dl	100 (32–190)
Triglycerides, mg/dl	90 (33–416)
Uric acid, mg/dl	5.5 (2.3–9.4)
<i>Treatment</i>	
PEG-IFN α -2b/IFN α -2b	422/68
Ribavirin dose, mg/kg	11.2 (3.7–15.1)
<i>aa substitutions in the HCV-1b</i>	
Core aa 70, arginine/glutamine (histidine)	266/151
Core aa 91, leucine/methionine	246/169
ISDR of NS5A, 0/1/ ≥ 2	227/95/84
IRRDR of NS5A, $\leq 3/4-5/\geq 6$	73/67/60
<i>Genetic variation near IL28B gene</i>	
rs8099917 genotype, TT/TG/GG	150/65/4

Data represent number of patients with percentages in parentheses, or median (range) values.

The correlation between ISDR and IRRDR was analyzed. There was a significant positive correlation between the number of aa substitutions in ISDR and those in IRRDR (*r* = 0.308, *p* < 0.001) (fig. 2).

aa Substitutions in the Core Region and NS5A-ISDR/IRRDR

Concerning the substitution of core aa 70, the number of aa substitutions in ISDR of 256 patients with Arg70 (median 0) was not significantly different from that of 146 patients with Gln70 (His70) (median 0) (fig. 3a). Fur-




Fig. 1. The number of aa substitutions in NS5A-ISDR and NS5A-IRRDR. The percentage of patients with ≤ 1 aa substitution in ISDR (WT) was 79%.




Fig. 2. Correlation between NS5A-ISDR and NS5A-IRRDR. There was a significant positive correlation between the number of aa substitutions in ISDR and that in IRRDR ($r = 0.308$, $p < 0.001$).

thermore, the number of aa substitutions in IRRDR of 123 patients with Arg70 (median 5) was also not significantly different from that of 77 patients with Gln70 (His70) (median 4) (fig. 3b).

Concerning the substitution of core aa 91, the number of aa substitutions in ISDR of 240 patients with Leu91 (median 1) was significantly higher than that of 161 patients with Met91 (median 0) ($p < 0.001$) (fig. 3c). Furthermore, the number of aa substitutions in IRRDR of 111 patients with Leu91 (median 5) was significantly higher than that of 89 patients with Met91 (median 3) ($p < 0.001$) (fig. 3d).

Viremia Level and aa Substitutions in Core Region/ISDR/IRRDR

Concerning the number of substitutions in ISDR, viremia levels of 321 patients with WT (median 6.5) were significantly higher than those of 84 patients with non-WT (median 5.7) ($p < 0.001$) (fig. 4a).

Concerning the number of substitutions in IRRDR, viremia levels of 140 patients with ≤ 5 substitutions (median 6.4) were significantly higher than those of 60 patients with ≥ 6 (median 6.1) ($p = 0.027$) (fig. 4b).

Concerning the substitution of core aa 70, viremia levels of 265 patients with Arg70 (median 6.4) were not significantly different from those of 151 patients with Gln70 (His70) (median 6.3) (fig. 4c).

Concerning the substitution of core aa 91, viremia levels of 169 patients with Met91 (median 6.5) were significantly higher than those of 245 patients with Leu91 (median 6.2) ($p = 0.028$) (fig. 4d).

Thus, levels of viremia were influenced by aa substitutions in core aa 91 and ISDR/IRRDR.

Treatment Response according to the Number of aa Substitutions in IRRDR

Concerning the number of aa substitutions in IRRDR, a significantly higher proportion of patients with ≥ 4 aa substitutions (58%) showed SVR compared to patients with ≤ 3 (42%) ($p = 0.039$). In contrast, the SVR rate was not significantly different between patients with ≤ 4 (49%) and those with ≥ 5 (57%) aa substitutions. Likewise, the SVR rate was not significantly different between patients with ≤ 5 (51%) and those with ≥ 6 (55%) aa substitutions (fig. 5a).

The ETR rate was not significantly different between patients with ≤ 3 (74%) and those with ≥ 4 (82%) aa substitutions, nor between patients with ≤ 4 (76%) and those with ≥ 5 (83%). Likewise, the ETR rate was not significantly different between those with ≤ 5 (79%) and those with ≥ 6 (80%) aa substitutions (fig. 5b).

The SVR rate after ETR was not significantly different between patients with ≤ 3 (61%) and those with ≥ 4 (74%) aa substitutions, nor between patients with ≤ 4 (67%) and those with ≥ 5 (72%). Likewise, they were not significantly different between patients with ≤ 5 (67%) and those with ≥ 6 (75%) aa substitutions (fig. 5c).

Thus, it was useful as predictor of SVR to categorize into two groups of ≤ 4 and ≥ 5 aa substitutions by univariate analysis. However, the ETR and SVR after ETR rates were not significantly different according to the number of aa substitutions in IRRDR.




Fig. 3. aa substitutions in the core region and NS5A-ISDR/IRRDR. **a, b** Concerning the substitution of core aa 70, the number of aa substitutions in ISDR/IRRDR of patients with Arg70 was not significantly different from that of patients with Gln70 (His70). **c, d** Concerning the substitution of core aa 91, the number of aa substitutions in ISDR/IRRDR of patients with Leu91 was significantly higher than that of patients with Met91 ($p < 0.001$).

Predictors of SVR as Determined by Uni- and Multivariate Analyses

Univariate analysis identified 15 parameters that correlate with SVR: gender (male sex; $p < 0.001$), age (< 55 years; $p < 0.001$), ribavirin dose (≥ 11.0 mg/kg; $p = 0.006$), AST (< 58 IU/l; $p = 0.039$), leukocyte count ($\geq 4,500/\text{mm}^3$; $p = 0.043$), hemoglobin (≥ 14.0 g/dl; $p = 0.001$), platelet count ($\geq 15.0 \times 10^4/\text{mm}^3$; $p < 0.001$), GGT (< 50 IU/l; $p = 0.028$), uric acid (≥ 5.5 mg/dl; $p = 0.005$), level of viremia (< 6.0 log IU/ml; $p < 0.001$), α -fetoprotein (< 10 $\mu\text{g/l}$; $p < 0.001$), genetic variation in rs8099917 (genotype TT; $p < 0.001$), substitution of aa 70 (Arg70; $p < 0.001$), the number of aa substitutions in ISDR (non-WT; $p < 0.001$) and IRRDR (≥ 4 ; $p = 0.039$). Figure 6 shows the SVR rate according to aa substitution in the core/NS5A region and genetic variation near *IL28B* by univariate analysis.

Multivariate analysis that included the above variables identified 3 parameters that independently influenced

SVR: genetic variation in rs8099917 (genotype TT; $p < 0.001$), gender (male sex; $p < 0.001$), and the number of aa substitutions in ISDR (non-WT; $p = 0.027$) (table 2).

Predictors of ETR as Determined by Uni- and Multivariate Analyses

Univariate analysis identified 14 parameters that correlated with ETR: gender (male sex; $p = 0.001$), age (< 55 years; $p = 0.004$), AST (< 39 IU/l; $p = 0.027$), hemoglobin (≥ 14.0 g/dl; $p = 0.035$), platelet count ($\geq 15.0 \times 10^4/\text{mm}^3$; $p < 0.001$), albumin (≥ 3.9 g/dl; $p = 0.014$), GGT (< 50 IU/l; $p < 0.001$), uric acid (≥ 5.5 mg/dl; $p = 0.003$), level of viremia (< 6.0 log IU/ml; $p = 0.001$), low-density lipoprotein cholesterol (≥ 85 mg/dl; $p = 0.004$), α -fetoprotein (< 10 $\mu\text{g/l}$; $p < 0.001$), genetic variation in rs8099917 (genotype TT; $p < 0.001$), substitution of aa 70 (Arg70; $p < 0.001$), and the number of aa substitutions in ISDR (non-WT; $p = 0.021$). Figure 7 shows the ETR rate according to aa




Fig. 4. Viremia level and aa substitutions in core region/ISDR/IRRDR. **a** Concerning the number of substitutions in ISDR, viremia levels of patients with WT were significantly higher than those of patients with non-WT ($p < 0.001$). **b** Concerning the number of substitutions in IRRDR, viremia levels of patients with ≤ 5 aa substitutions were significantly higher levels than those of patients with ≥ 6 ($p = 0.027$). **c** Concerning the substitution of

core aa 70, viremia levels of patients with Arg70 were not significantly different from those of patients with Gln70 (His70). **d** Concerning the substitution of core aa 91, viremia levels of patients with Met91 were significantly higher than those of patients with Leu91 ($p = 0.028$). Thus, levels of viremia might be influenced by aa substitutions in core aa 91 and ISDR/IRRDR.




Fig. 5. Treatment response according to the number of aa substitutions in NS5A-IRRDR. **a** A significantly higher proportion of patients with ≥ 4 (58%) aa substitutions showed SVR compared to patients with ≤ 3 (42%) ($p = 0.039$), and it was useful as predictor

of SVR to categorize into two groups of ≤ 4 and ≥ 5 aa substitutions by univariate analysis. **b**, **c** ETR and SVR after ETR rates were not significantly different according to the number of aa substitutions in IRRDR.




Fig. 6. SVR rate according to aa substitution in core/NS5A region and genetic variation near *IL28B* by univariate analysis.




Fig. 7. ETR rate according to aa substitution in core/NS5A region and genetic variation near *IL28B* by univariate analysis.

Table 2. Factors associated with SVR to 48-week IFN + ribavirin combination therapy in patients infected with HCV-1b, identified by multivariate analysis

Factor	Category	OR (95% CI)	p
rs8099917 genotype	1: TG+GG	1	
	2: TT	16.7 (4.54–61.3)	<0.001
Gender	1: Female	1	
	2: Male	10.5 (3.47–32.3)	<0.001
ISDR of NS5A	1: WT	1	
	2: Non-WT	5.68 (1.22–26.3)	0.027

Only variables that achieved statistical significance ($p < 0.05$) on multivariate logistic regression are shown.

Table 3. Factors associated with ETR response to 48-week IFN + ribavirin combination therapy in patients infected with HCV-1b, identified by multivariate analysis

Factor	Category	OR (95% CI)	p
rs8099917 genotype	1: TG+GG	1	
	2: TT	18.2 (6.29–52.6)	<0.001
Level of viremia log IU/ml	1: ≥6.0	1	
	2: <6.0	9.20 (2.59–32.6)	0.001
Core aa 70	1: Gln70 (His70)	1	
	2: Arg70	4.68 (1.65–13.3)	0.004
Serum albumin g/dl	1: <3.9	1	
	2: ≥3.9	3.08 (1.11–8.47)	0.030

Only variables that achieved statistical significance ($p < 0.05$) on multivariate logistic regression are shown.




Fig. 8. SVR after ETR rate according to aa substitution in core/NS5A region and genetic variation near *IL28B* by univariate analysis.

substitution in the core/NS5A region and genetic variation near *IL28B* by univariate analysis.

Multivariate analysis that included the above variables identified 4 parameters that independently influenced ETR: genetic variation in rs8099917 (genotype TT; $p < 0.001$), level of viremia ($< 6.0 \log \text{ IU/ml}$; $p = 0.001$), substitution of aa 70 (Arg70; $p = 0.004$), and albumin ($\geq 3.9 \text{ g/dl}$; $p = 0.030$) (table 3).

Predictors of SVR after ETR as Determined by Uni- and Multivariate Analyses

Univariate analysis identified 11 parameters that influenced SVR after ETR: gender (male sex; $p < 0.001$), age (< 55 years; $p < 0.001$), ribavirin dose ($\geq 11.0 \text{ mg/kg}$; $p = 0.025$), leukocyte count ($\geq 4,500/\text{mm}^3$; $p = 0.033$), hemoglobin ($\geq 14.0 \text{ g/dl}$; $p = 0.025$), platelet count ($\geq 15.0 \times 10^4/\text{mm}^3$; $p = 0.001$), level of viremia ($< 6.0 \log \text{ IU/ml}$; $p = 0.020$), total cholesterol ($< 170 \text{ mg/dl}$; $p = 0.017$), α -fetoprotein ($< 10 \mu\text{g/l}$; $p = 0.004$), substitution of aa 70 and 91 (Arg70 and/or Leu91; $p = 0.044$), and the number of aa substitutions in ISDR (non-WT; $p = 0.001$). Figure 8 shows the SVR after ETR rate according to aa substitution in the core/NS5A region and genetic variation near *IL28B* by univariate analysis.

Multivariate analysis that included the above variables identified 6 parameters that independently influenced the SVR after ETR: gender (male sex; $p < 0.001$), ribavirin dose ($\geq 11.0 \text{ mg/kg}$; $p = 0.002$), the number of aa substitutions in ISDR (non-WT; $p = 0.012$), substitution of aa 70 and 91 (Arg70 and/or Leu91; $p = 0.023$), platelet count ($\geq 15.0 \times 10^4/\text{mm}^3$; $p = 0.033$), and α -fetoprotein ($< 10 \mu\text{g/l}$; $p = 0.042$) (table 4).

Comparison of Factors Associated with Treatment Efficacy Identified by Multivariate Analysis

Table 5 shows the variables that achieved statistical significance on multivariate logistic regression for each evaluation of treatment efficacy. Rs8099917 genotype was an important predictor of ETR and SVR. With regard to viral factors, core region was an important predictor of ETR, and SVR after ETR. ISDR was an important predictor of SVR, and SVR after ETR. Level of viremia was an important predictor of ETR. Thus, genetic variation near *IL28B* and viral factors (core region, ISDR, and level of viremia) were important predictors of treatment efficacy. Furthermore, gender, α -fetoprotein, albumin, and platelet count were also identified as other important predictors of treatment efficacy, in addition to genetic variation near *IL28B* and viral factors.

Discussion

Using multivariate analysis, the present study identified viral- (aa substitutions in core aa 70/91 and NS5A-ISDR/IRRDR) and host-related factors (genetic variation near *IL28B* gene) that influenced treatment efficacy to 48-week IFN/ribavirin combination therapy, which is in agreement with recent findings [22, 23]. Identification of these viral and host factors before the start of IFN/ribavirin combination therapy should help to select better therapeutic regimens, including triple therapy of telaprevir/PEG-IFN/ribavirin [24–26], for those patients who are less likely to achieve SVR.

According to the number of substitutions in ISDR, a previous report showed that levels of viremia were sig-

Table 4. Factors associated with SVR in patients who achieved ETR response to 48-week IFN + ribavirin combination therapy in patients infected with HCV-1b, identified by multivariate analysis

Factor	Category	OR (95% CI)	p
Gender	1: Female	1	<0.001
	2: Male	4.27 (2.15–8.55)	
Ribavirin dose, mg/kg	1: <11.0	1	0.002
	2: ≥11.0	2.95 (1.48–5.86)	
ISDR of NS5A	1: WT	1	0.012
	2: Non-WT	4.00 (1.35–11.8)	
Core aa 70 and 91	1: Gln70 (His70) and Met91	1	0.023
	2: Arg70 and/or Leu91	2.96 (1.16–7.52)	
Platelet count × 10 ⁴ /mm ³	1: <15.0	1	0.033
	2: ≥15.0	2.19 (1.07–4.50)	
α-Fetoprotein μg/l	1: ≥10	1	0.042
	2: <10	2.66 (1.04–6.80)	

Only variables that achieved statistical significance (p < 0.05) on multivariate logistic regression are shown.

Table 5. Comparison of factors associated with efficacy of 48-week IFN + ribavirin combination therapy in patients infected with HCV-1b, identified by multivariate analysis

Factor	ETR response (at 48 weeks)	SVR after ETR response	SVR
<i>IL28B</i>	rs8099917 p < 0.001, 18.2 (6.29–52.6) ^a		rs8099917 p < 0.001, 16.7 (4.54–61.3) ^a
Virus	Core aa 70 p = 0.004, 4.68 (1.65–13.3) ^a	Core aa 70 and 91 p = 0.023, 2.96 (1.16–7.52) ^a	
	Level of viremia p = 0.001, 9.20 (2.59–32.6) ^a	ISDR p = 0.012, 4.00 (1.35–11.8) ^a	ISDR p = 0.027, 5.68 (1.22–26.3) ^a
Others	Albumin p = 0.030, 3.08 (1.11–8.47) ^a	α-Fetoprotein p = 0.042, 2.66 (1.04–6.80) ^a	
		Platelet count p = 0.033, 2.19 (1.07–4.50) ^a	
		Gender p < 0.001, 4.27 (2.15–8.55) ^a	Gender p < 0.001, 10.5 (3.47–32.3) ^a
		Ribavirin dose p = 0.002, 2.95 (1.48–5.86) ^a	

Only variables that achieved statistical significance (p < 0.05) on multivariate logistic regression are shown.
^a OR (95% CI).

nificantly lower in patients with non-WT of ISDR than in those with WT [8]. The present study indicated that substitution of IRRDR and core aa 91, in addition to substitution of ISDR, also significantly influenced levels of viremia. Furthermore, there was a significant positive correlation between the number of aa substitutions in

ISDR and those in IRRDR, and the number of aa substitutions in ISDR/IRRDR of patients with Leu91 was significantly higher than that of patients with Met91. To our knowledge, this is the first report of the relationship between viremia levels and aa substitutions in core region/ISDR/IRRDR. This result might be interpreted to mean

that core aa 91/ISDR/IRRDR might be associated with viremia levels involved in resistance to combination therapy. Further studies that examine the functional impact of aa substitutions to combination therapy should be conducted to confirm the above finding.

The present results showed that α -fetoprotein, albumin, platelet count, and gender were predictors of virological response to IFN/ribavirin combination therapy. Previous data indicated that absence of advanced liver fibrosis was a positive predictor of SVR to IFN monotherapy and IFN/ribavirin combination therapy [2, 3, 13, 27–29], and that advanced liver fibrosis was usually associated with higher levels of α -fetoprotein, and lower levels of albumin and platelet count [1, 3, 30–32]. Furthermore, gender is also a predictor of treatment response to IFN/ribavirin combination therapy [2, 3, 14]. In the present study based on a large number of patients, histopathological changes in the liver and gender were identified as independent predictors of virological response, in addition to genetic variation near *IL28B* and viral factors (core region, ISDR, and level of viremia).

In a previous study, multivariate analysis identified core region, gender, and stage of liver fibrosis as parameters that independently influenced the SVR of patients who achieved early virological response, but ISDR was not entered into uni- and multivariate analysis [3]. To our knowledge, the present study based on multivariate analysis is the first report to identify ISDR as pretreatment

predictor of SVR after ETR to combination therapy. Interestingly, ISDR was not a predictor of ETR, but was a significant predictor of SVR to combination therapy. Thus, the underlying mechanisms of failure to develop SVR in those patients who achieve HCV-RNA negativity remain unclear. Further studies that examine the impact of aa substitutions of ISDR to combination therapy should be conducted to confirm the above finding.

One limitation of the present study was that aa substitutions in areas other than the core region and NS5A-ISDR/IRRDR of the HCV genome were not examined. Other limitations were differences in host factors including race [24, 33, 34] and differences in viral factors, such as the distribution of HCV-1a or -1b, and geographic diversities of HCV-1b [35]. Further large-scale prospective studies are necessary to investigate whether the present results relate to the efficacy of 48-week IFN/ribavirin combination therapy, and further understanding of the complex interaction between virus- and host-related factors should facilitate the development of more effective therapeutic regimens.

Acknowledgement

This study was supported in part by a Grant-in-Aid from the Ministry of Health, Labor and Welfare, Japan.

References

- 1 Akuta N, Suzuki F, Sezaki H, Suzuki Y, Hosaka T, Someya T, Kobayashi M, Saitoh S, Watahiki S, Sato J, Matsuda M, Kobayashi M, Arase Y, Ikeda K, Kumada H: Association of amino acid substitution pattern in core protein of hepatitis C virus genotype 1b high viral load and non-virological response to interferon-ribavirin combination therapy. *Intervirology* 2005;48:372–380.
- 2 Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Arase Y, Ikeda K, Kumada H: Predictive factors of early and sustained responses to peginterferon plus ribavirin combination therapy in Japanese patients infected with hepatitis C virus genotype 1b: amino acid substitutions in the core region and low-density lipoprotein cholesterol levels. *J Hepatol* 2007;46:403–410.
- 3 Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Arase Y, Ikeda K, Kumada H: Predictors of viral kinetics to peginterferon plus ribavirin combination therapy in Japanese patients infected with hepatitis C virus genotype 1b. *J Med Virol* 2007;79:1686–1695.
- 4 Donlin MJ, Cannon NA, Yao E, Li J, Wahed A, Taylor MW, Belle SH, Di Bisceglie AM, Aurora R, Tavis JE: Pretreatment sequence diversity differences in the full-length hepatitis C virus open reading frame correlate with early response to therapy. *J Virol* 2007; 81:8211–8224.
- 5 Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Arase Y, Ikeda K, Kumada H: Amino acid substitutions in the hepatitis C virus core region are the important predictor of hepatocarcinogenesis. *Hepatology* 2007; 46:1357–1364.
- 6 Fishman SL, Factor SH, Balestrieri C, Fan X, Dibisceglie AM, Desai SM, Benson G, Branch AD: Mutations in the hepatitis C virus core gene are associated with advanced liver disease and hepatocellular carcinoma. *Clin Cancer Res* 2009;15:3205–3213.
- 7 Enomoto N, Sakuma I, Asahina Y, Kurosaki M, Murakami T, Yamamoto C, Izumi N, Marumo F, Sato C: Comparison of full-length sequences of interferon sensitive and resistant hepatitis C virus 1b. Sensitivity to interferon is conferred by amino acid substitutions in the NS5A region. *J Clin Invest* 1995;96:224–230.
- 8 Enomoto N, Sakuma I, Asahina Y, Kurosaki M, Murakami T, Yamamoto C, Ogura Y, Izumi N, Marumo F, Sato C: Mutations in the nonstructural protein 5A gene and response to interferon in patients with chronic hepatitis C virus 1b infection. *N Engl J Med* 1996; 334:77–81.
- 9 El-Shamy A, Sasayama M, Nagano-Fujii M, Sasase N, Imoto S, Kim SR, Hotta H: Prediction of efficient virological response to pegylated interferon/ribavirin combination therapy by NS5A sequences of hepatitis C virus and anti-NS5A antibodies in pre-treatment sera. *Microbiol Immunol* 2007;51:471–482.

- 10 El-Shamy A, Nagano-Fujii M, Sasase N, Imoto S, Kim SR, Hotta H: Sequence variation in hepatitis C virus nonstructural protein 5A predicts clinical outcome of pegylated interferon/ribavirin combination therapy. *Hepatology* 2008;48:38–47.
- 11 Shirakawa H, Matsumoto A, Joshita S, Komatsu M, Tanaka N, Umemura T, Ichijo T, Yoshizawa K, Kiyosawa K, Tanaka E: Nagano Interferon Treatment Research Group: Pre-treatment prediction of virological response to peginterferon plus ribavirin therapy in chronic hepatitis C patients using viral and host factors. *Hepatology* 2008;48:1753–1760.
- 12 Mori N, Imamura M, Kawakami Y, Saneto H, Kawaoka T, Takaki S, Aikata H, Takahashi S, Chayama K: Hiroshima Liver Study Group: Randomized trial of high-dose interferon- α -2b combined with ribavirin in patients with chronic hepatitis C: Correlation between amino acid substitutions in the core/NS5A region and virological response to interferon therapy. *J Med Virol* 2009;81:640–649.
- 13 Ge D, Fellay J, Thompson AJ, Simon JS, Shianna KV, Urban TJ, Heinzen EL, Qiu P, Bertelsen AH, Muir AJ, Sulkowski M, McHutchison JG, Goldstein DB: Genetic variation in *IL28B* predicts hepatitis C treatment-induced viral clearance. *Nature* 2009;461:399–401.
- 14 Tanaka Y, Nishida N, Sugiyama M, Kurosaki M, Matsuura K, Sakamoto N, Nakagawa M, Korenaga M, Hino K, Hige S, Ito Y, Mita E, Tanaka E, Mochida S, Murawaki Y, Honda M, Sakai A, Hiasa Y, Nishiguchi S, Koike A, Sakaïda I, Imamura M, Ito K, Yano K, Masaki N, Sugauchi F, Izumi N, Tokunaga K, Mizokami M: Genome-wide association of *IL28B* with response to pegylated interferon- α and ribavirin therapy for chronic hepatitis C. *Nat Genet* 2009;41:1105–1109.
- 15 Suppiah V, Moldovan M, Ahlenstiel G, Berg T, Weltman M, Abate ML, Bassendine M, Spengler U, Dore GJ, Powell E, Riordan S, Sheridan D, Smedile A, Fragomeli V, Müller T, Bahlo M, Stewart GJ, Booth DR, George J: *IL28B* is associated with response to chronic hepatitis C interferon- α and ribavirin therapy. *Nat Genet* 2009;41:1100–1104.
- 16 Rauch A, Kutalik Z, Descombes P, Cai T, Di Iulio J, Mueller T, Bochud M, Battegay M, Bernasconi E, Borovicka J, Colombo S, Cerny A, Dufour JF, Furrer H, Günthard HF, Heim M, Hirschel B, Malinverni R, Moradpour D, Müllhaupt B, Witteck A, Beckmann JS, Berg T, Bergmann S, Negro F, Telenti A, Bochud PY: Swiss Hepatitis C Cohort Study; Swiss HIV Cohort Study: Genetic variation in *IL28B* is associated with chronic hepatitis C and treatment failure: a genome-wide association study. *Gastroenterology* 2010;138:1338–1345.
- 17 Thomas DL, Thio CL, Martin MP, Qi Y, Ge D, O’Huigin C, Kidd J, Kidd K, Khakoo SI, Alexander G, Goedert JJ, Kirk GD, Donfield SM, Rosen HR, Tobler LH, Busch MP, McHutchison JG, Goldstein DB, Carrington M: Genetic variation in *IL28B* and spontaneous clearance of hepatitis C virus. *Nature* 2009;461:798–801.
- 18 Akuta N, Suzuki F, Hirakawa M, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Saitoh S, Arase Y, Ikeda K, Chayama K, Nakamura Y, Kumada H: Amino acid substitution in HCV core region and genetic variation near the interleukin-28B gene predict viral response to telaprevir with peginterferon and ribavirin. *Hepatology* 2010;52:421–429.
- 19 Kato N, Hijikata M, Ootsuyama Y, Nakagawa M, Ohkoshi S, Sugimura T, Shimotohno K: Molecular cloning of the human hepatitis C virus genome from Japanese patients with non-A, non-B hepatitis. *Proc Natl Acad Sci USA* 1990;87:9524–9528.
- 20 Ohnishi Y, Tanaka T, Ozaki K, Yamada R, Suzuki H, Nakamura Y: A high-throughput SNP typing system for genome-wide association studies. *J Hum Genet* 2001;46:471–477.
- 21 Suzuki A, Yamada R, Chang X, Tokunaga S, Sawada T, Suzuki M, Nagasaki M, Nakayama-Hamada M, Kawaida R, Ono M, Ohtsuki M, Furukawa H, Yoshino S, Yukioka M, Tohma S, Matsubara T, Wakitani S, Teshima R, Nishioka Y, Sekine A, Iida A, Takahashi A, Tsunoda T, Nakamura Y, Yamamoto K: Functional haplotypes of *PAD14*, encoding citrullinating enzyme peptidylarginine deiminase 4, are associated with rheumatoid arthritis. *Nat Genet* 2003;34:395–402.
- 22 Kurosaki M, Tanaka Y, Nishida N, Sakamoto N, Enomoto N, Honda M, Sugiyama M, Matsuura K, Sugauchi F, Asahina Y, Nakagawa M, Watanabe M, Sakamoto M, Maekawa S, Sakai A, Kaneko S, Ito K, Masaki N, Tokunaga K, Izumi N, Mizokami M: Pre-treatment prediction of response to pegylated-interferon plus ribavirin for chronic hepatitis C using genetic polymorphism in *IL28B* and viral factors. *J Hepatol* 2011;54:439–448.
- 23 Hayes CN, Kobayashi M, Akuta N, Suzuki F, Kumada H, Abe H, Miki D, Imamura M, Ochi H, Kamatani N, Nakamura Y, Chayama K: HCV substitutions and *IL28B* polymorphisms on outcome of peg-interferon plus ribavirin combination therapy. *Gut* 2011;60:261–267.
- 24 McHutchison JG, Everson GT, Gordon SC, Jacobson IM, Sulkowski M, Kauffman R, McNair L, Alam J, Muir AJ: PROVE1 Study Team: Telaprevir with peginterferon and ribavirin for chronic HCV genotype 1 infection. *N Engl J Med* 2009;360:1827–1838.
- 25 McHutchison JG, Manns MP, Muir AJ, Terrault NA, Jacobson IM, Afdhal NH, Heathcote EJ, Zeuzem S, Reesink HW, Garg J, Bsharat M, George S, Kauffman RS, Adda N, Di Bisceglie AM: PROVE3 Study Team: Telaprevir for previously treated chronic HCV infection. *N Engl J Med* 2010;362:1292–1303.
- 26 Hézode C, Forestier N, Dusheiko G, Ferenci P, Pol S, Goeser T, Bronowicki JP, Bourlière M, Gharakhanian S, Bengtsson L, McNair L, George S, Kieffer T, Kwong A, Kauffman RS, Alam J, Pawlotsky JM, Zeuzem S: PROVE2 Study Team: Telaprevir and peginterferon with or without ribavirin for chronic HCV infection. *N Engl J Med* 2009;360:1839–1850.
- 27 Jouet P, Roudot-Thoraval F, Dhumeaux D, Metreau JM: Comparative efficacy of interferon alfa in cirrhotic and noncirrhotic patients with non-A, non-B, C hepatitis. *Gastroenterology* 1994;106:686–690.
- 28 Poynard T, McHutchinson J, Goodman Z, Ling MH, Albrecht J: Is an ‘a la carte’ combination interferon alfa-2b plus ribavirin regimen possible for the first-line treatment in patients with chronic hepatitis C? The AL-GOVIROC Group. *Hepatology* 2000;31:211–218.
- 29 Bruno S, Camma C, Di Marco V, Rumi M, Vinci M, Camozzi M, Rebusci C, Di Bona D, Colombo M, Craxi A, Mondelli MU, Pinzello G: Peginterferon alfa-2b plus ribavirin for naïve patients with genotype 1 chronic hepatitis C: a randomized controlled trial. *J Hepatol* 2004;41:474–481.
- 30 Bayati N, Silverman AL, Gordon SC: Serum α -fetoprotein levels and liver histology in patients with chronic hepatitis C. *Am J Gastroenterol* 1998;93:2452–2456.
- 31 Chu CW, Hwang SJ, Luo JC, Lai CR, Tsay SH, Li CP, Wu JC, Chang FY, Lee SD: Clinical, virological, and pathologic significance of elevated serum α -fetoprotein levels in patients with chronic hepatitis C. *J Clin Gastroenterol* 2001;32:240–244.
- 32 Hu KQ, Kyulo NL, Lim N, Elhazim B, Hillebrand DJ, Bock T: Clinical significance of elevated α -fetoprotein in patients with chronic hepatitis C, but not hepatocellular carcinoma. *Am J Gastroenterol* 2004;99:860–865.
- 33 McHutchinson JG, Poynard T, Pianko S, Gordon SC, Reid AE, Dienstag J, Morgan T, Yao R, Albrecht J: The impact of interferon plus ribavirin on response to therapy in black patients with chronic hepatitis C. The International Hepatitis Interventional Therapy Group. *Gastroenterology* 2000;119:1317–1323.
- 34 Kaplan DE, Sugimoto K, Ikeda F, Stadanlick J, Valiga M, Shetty K, Reddy KR, Chang KM: T-cell response relative to genotype and ethnicity during antiviral therapy for chronic hepatitis C. *Hepatology* 2005;41:1365–1375.
- 35 Nakano I, Fukuda Y, Katano Y, Nakano S, Kumada T, Hayakawa T: Why is the interferon sensitivity-determining region (ISDR) system useful in Japan? *J Hepatol* 1999;30:1014–1022.



Complicated Relationships of Amino Acid Substitution in Hepatitis C Virus Core Region and *IL28B* Genotype Influencing Hepatocarcinogenesis

Norio Akuta,¹ Fumitaka Suzuki,¹ Yuya Seko,¹ Yusuke Kawamura,¹ Hitomi Sezaki,¹ Yoshiyuki Suzuki,¹ Tetsuya Hosaka,¹ Masahiro Kobayashi,¹ Tasuku Hara,¹ Mariko Kobayashi,² Satoshi Saitoh,¹ Yasuji Arase,¹ Kenji Ikeda,¹ and Hiromitsu Kumada¹

The impact of amino acid (aa) 70 substitution in the core region on hepatocarcinogenesis and survival for liver-related death in patients of hepatitis C virus (HCV) genotype 1b (HCV-1b), who had not received antiviral therapy, is unknown. The relationships among aa 70 substitution, *IL28B* genotype, and hepatocarcinogenesis are also not clear. A total of 1,181 consecutive HCV-infected patients, who had not received antiviral therapy, were included in a follow-up study to determine predictive factors of hepatocarcinogenesis and survival for liver-related death. The cumulative hepatocarcinogenesis rates in HCV-1b of Gln70(His70) (glutamine (histidine) at aa 70) were significantly higher than those in HCV-1b of Arg70 (arginine at aa 70) and HCV-2a/2b. The cumulative survival rates for liver-related death in HCV-1b of Gln70(His70) were significantly lower than those in HCV-1b of Arg70 and HCV-2a/2b. Multivariate analysis identified gender (male), age (≥ 60 years), albumin (< 3.9 g/dL), platelet count ($< 15.0 \times 10^4/\text{mm}^3$), aspartate aminotransferase (≥ 67 IU/L), and HCV subgroup (HCV-1b of Gln70(His70)) as determinants of both hepatocarcinogenesis and survival rates for liver-related death. In HCV-1b patients, the cumulative change rates from Arg70 to Gln70(His70) by direct sequencing were significantly higher than those from Gln70(His70) to Arg70. In patients of Arg70 at the initial visit, the cumulative change rates from Arg70 to Gln70(His70) in *IL28B* rs8099917 non-TT genotype were significantly higher than those in the TT genotype. **Conclusion:** Substitution of aa 70 in the core region of HCV-1b is an important predictor of hepatocarcinogenesis and survival for liver-related death in HCV patients who had not received antiviral therapy. The *IL28B* genotype might partly affect changes over time of dominant amino acid in core aa 70 of HCV-1b. (HEPATOLOGY 2012;56:2134-2141)

Hepatitis C virus (HCV) usually causes chronic infection that can result in chronic hepatitis, liver cirrhosis, and hepatocellular carcinoma (HCC).^{1,2} At present, treatments based on interferon (IFN), in combination with ribavirin, are the mainstay for combating HCV infection. In Japan, HCV genotype 1b (HCV-1b) and high viral loads account for more than 70% of HCV infections, making it difficult to treat patients with chronic hepatitis C.³

Despite numerous lines of epidemiologic evidence connecting HCV infection and the development of

HCC, it remains controversial whether HCV itself plays a direct role or an indirect role in the pathogenesis of HCC.⁴ It has become evident that HCV core region has oncogenic potential through the use of transgenic mice, but the clinical impact of the core region on hepatocarcinogenesis is still unclear.⁵ Previous reports indicated that amino acid (aa) substitutions at position 70 in the HCV core region of patients infected with HCV-1b are pretreatment predictors of poor virological response to pegylated IFN (PEG-IFN)/ribavirin combination therapy and triple therapy

Abbreviations: aa, amino acid; ALT, alanine aminotransferase; AST, aspartate aminotransferase; HCC, hepatocellular carcinoma; HCV, hepatitis C virus; PEG/IFN, pegylated interferon.

From the ¹Department of Hepatology, Toranomon Hospital, and Okinaka Memorial Institute for Medical Research, Tokyo, Japan; ²Liver Research Laboratory, Toranomon Hospital, Tokyo, Japan.

Received May 1, 2012; Revised May 30, 2012; accepted June 17, 2012.

Supported in part by Grants-in-Aid for scientific research and development from the Ministry of Health, Labor and Welfare and Ministry of Education Culture Sports Science and Technology, Government of Japan.

of telaprevir/PEG-IFN/ribavirin,⁶⁻⁹ and also affects hepatocarcinogenesis.¹⁰⁻¹³ These reports support the findings of oncogenic potential by core region from the clinical aspect. However, its impact on hepatocarcinogenesis and survival for liver-related death in patients of HCV-1b who had not received antiviral therapy is still unknown.

The *IL28B* genotype is a poor predictor of virological response to PEG-IFN/ribavirin combination therapy and triple therapy of telaprevir/PEG-IFN/ribavirin^{9,14-17} and is reported to be associated with HCC, although its impact on HCC is controversial.¹⁸⁻²¹ Furthermore, treatment-resistant substitution of core aa 70 (glutamine/histidine at aa 70 (Gln70/His70)), which might affect hepatocarcinogenesis, was significantly more frequent in patients with treatment-resistant genotype (non-TT) than -sensitive genotype (TT) at *IL28B* rs8099917.²¹⁻²³ Thus, the significant linkage between substitution of aa 70 and *IL28B* genotype had been shown, but it is not clarified whether the existence of a complex interaction between the virus and host might affect hepatocarcinogenesis.

The present study included 1,181 consecutive HCV-infected patients who had not received antiviral therapy. The aims of the study were: (1) To evaluate the impact of aa substitutions in the core region of HCV-1b on hepatocarcinogenesis and survival for liver-related death; and (2) To investigate the association of *IL28B* genotype and time-dependent aa changes in the core region of HCV-1b.

Patients and Methods

Patients. Among 2,799 consecutive HCV-infected patients in whom antiviral therapy (IFN monotherapy or IFN plus ribavirin combination therapy) was not induced between December 1962 and November 2010 at Toranomon Hospital, 1,181 were selected in the present study based on the following criteria. (1) Positive for anti-HCV (third-generation enzyme immunoassay, Chiron, Emerville, CA) and positive for HCV RNA (nested polymerase chain reaction [PCR]), at the initial visit. (2) Patients without HCC at the initial visit. (3) Patients infected with single genotype of

Table 1. Profiles and laboratory data at the initial visit of 1,181 patients infected with HCV, who had not received antiviral therapy

Demographic data	
Number of patients	1,181
Sex (male/female)	608/573
Age (years)*	60 (20-93)
History of blood transfusion	526 (49.2%)
Family history of liver disease	201 (20.3%)
Lifetime cumulative alcohol intake (>500 kg)	110 (10.8%)
Laboratory data*	
Total bilirubin (mg/dl)	0.7(0.1-20.0)
Aspartate aminotransferase (IU/l)	71 (13-1,052)
Alanine aminotransferase (IU/l)	88 (4-1,210)
Albumin (g/dl)	4.1 (1.0-5.5)
Hemoglobin (g/dl)	14.0 (7.8-18.0)
Platelet count ($\times 10^4/\text{mm}^3$)	15.3 (2.6-52.9)
HCV genotype (1b / 2a or 2b)	750/431
Levels of viremia (high viral load)	757 (74.4%)
Amino acid substitutions in the HCV genotype 1b	
Core aa 70 (arginine / glutamine (histidine))	431/319
Core aa 91 (leucine / methionine)	482/268

Data are number and percentages of patients, except those denoted by *, which represent the median (range) values.

HCV-1b, 2a, or 2b. (4) In HCV-1b, patients analyzed aa substitutions of the core region by direct sequencing, one or more times from the initial visit. (5) Patients negative for hepatitis B surface antigen (radioimmunoassay, Dainabot, Tokyo, Japan). (6) Patients free of coinfection with human immunodeficiency virus. (7) Patients free of other types of chronic liver disease, including hemochromatosis, Wilson's disease, primary biliary cirrhosis, alcoholic liver disease, autoimmune liver disease, inherited liver disease including alpha-1 antitrypsin deficiency, and hepatic venous outflow block. (8) Patients who consented to the study.

Table 1 summarizes the profiles and laboratory data at the initial visit of 1,181 patients infected with HCV who had not received antiviral therapy. They did not receive antiviral therapy based on concerns about adverse effects, lack of time for treatment, physician recommendation based on the appearance of depression and cardiopulmonary disease, lower levels of aspartate aminotransferase (AST) / alanine aminotransferase (ALT), or elderly patients. They included 608 males and 573 females, aged 20 to 93 years (median, 60 years). The median follow-up time from the initial visit until death or until the last visit was 9.0 years (range, 0.0-37.7

Address reprint requests to: Norio Akuta, M.D., Department of Hepatology, Toranomon Hospital, 2-2-2 Toranomon, Minato-ku, Tokyo, 105-0001, Japan. E-mail: akuta-gi@umin.ac.jp; fax: +81-44-860-1623.

Copyright © 2012 by the American Association for the Study of Liver Diseases.

View this article online at wileyonlinelibrary.com.

DOI 10.1002/hep.25949

Potential conflict of interest: Norio Akuta has received speakers' bureau from MSD K.K., and holds a right to get some loyalty from SRL. Inc. Hiromitsu Kumada has received speakers' bureau from MSD K.K., Mitsubishi Tanabe Pharma, Dainippon Sumitomo Pharma, Bristol-Myers Squibb, and holds a right to get some loyalty from SRL. Inc. Fumitaka Suzuki has received speakers' bureau from Bristol-Myers Squibb. The other authors have nothing to disclose.

years). The study protocol was approved by the Human Ethics Review Committee of the institution.

Laboratory Investigations. Blood samples were frozen at -80°C within 4 hours of collection and were not thawed until used for testing. Anti-HCV, HCV RNA, HCV genotype, and aa substitutions of the HCV-1b core region were assayed using stored frozen sera. HCV genotype was determined by PCR using a mixed primer set derived from nucleotide sequences of the NS5 region.²⁴ HCV RNA quantitative analysis was measured by branched DNA assay v. 2.0 (Chiron), AMPLICOR GT HCV Monitor v. 2.0 using the 10-fold dilution method (Roche Molecular Systems, Pleasanton, CA), or COBAS TaqMan HCV test (Roche Diagnostics, Tokyo, Japan). High viral load of viremia levels was defined as branched DNA assay ≥ 1.0 Meq/mL, AMPLICOR GT HCV Monitor $\geq 100 \times 10^3$ IU/mL, or COBAS TaqMan HCV test ≥ 5.0 log IU/mL. Low viral load was defined as branched DNA assay < 1.0 Meq/mL, AMPLICOR GT HCV Monitor $< 100 \times 10^3$ IU/mL, or COBAS TaqMan HCV test < 5.0 log IU/mL.

Detection of Amino Acid Substitutions in Core Regions of HCV-1b. In the present study, aa substitutions of the core region of HCV-1b were analyzed by direct sequencing. HCV RNA was extracted from serum samples and reverse transcribed with random primer and MMLV reverse transcriptase (Takara Syuzo, Tokyo, Japan). Nucleic acids of the core region were amplified by nested PCR using the following primers. The first-round PCR was performed with CE1 (sense, 5'-GTC TGC GGA ACC GGT GAG TA-3', nucleotides: 134-153) and CE2 (antisense, 5'-GAC GTG GCG TCG TAT TGT CG-3', nucleotides: 1096-1115) primers, and the second-round PCR with CC9 (sense, 5'-ACT GCT AGC CGA GTA GTG TT-3', nucleotides: 234-253) and CE6 (antisense, 5'-GGA GCA GTC GTT CGT GAC AT-3', nucleotides: 934-953) primers. All samples were initially denatured at 95°C for 2 minutes. The 35 cycles of amplification were set as follows: denaturation for 30 seconds at 95°C , annealing of primers for 30 seconds at 55°C , and extension for 1 minute at 72°C with an additional 7 minutes for extension. Then $1 \mu\text{L}$ of the first PCR product was transferred to the second PCR reaction. Other conditions for the second PCR were the same as the first PCR, except that the second PCR primers were used instead of the first PCR primers. The amplified PCR products were purified by the QIA quick PCR purification kit (Qiagen, Tokyo, Japan) after agarose gel electrophoresis and then used for direct sequencing. Dideoxynucleotide termination sequencing

was performed with the Big Dye Deoxy Terminator Cycle Sequencing kit (Perkin-Elmer, Tokyo, Japan).

With the use of HCV-J (accession no. D90208) as a reference,²⁵ the dominant sequence of 1-191 aa in the core protein of HCV-1b was determined by direct sequencing and then compared with the consensus sequence constructed on 50 clinical samples to detect substitutions at aa 70 of arginine (Arg70) or glutamine/histidine (Gln70/His70) and aa 91 of leucine (Leu91) or methionine (Met91).⁶ Especially, patients were classified into three HCV subgroups according to HCV genotype in combination with aa substitutions in HCV-1b core region (HCV-1b of Arg70, HCV-1b of Gln70(His70), and HCV-2a/2b).

Determination of IL28B Genotype. IL28B rs8099917 was genotyped by the Invader assay, TaqMan assay, or direct sequencing, as described.^{26,27}

Follow-Up and Diagnosis of HCC. Follow-up of patients was made on a monthly to trimonthly basis after the initial visit. Imaging diagnosis was made one or more times per year with ultrasonography, computed tomography, or magnetic resonance imaging. During this time, liver-related death, which included HCC, cholangiocellular carcinoma, liver failure, or esophageal variceal bleeding, was also evaluated.

Statistical Analysis. The cumulative rates of hepatocarcinogenesis, survival for liver-related death, and amino acid changes in the core region were calculated using the Kaplan-Meier technique; differences between the curves were tested using the log-rank test. Statistical analyses of hepatocarcinogenesis, survival, and amino acid changes, according to groups, were calculated using the period from the initial visit. Stepwise Cox regression analysis was used to determine independent predictive factors that were associated with hepatocarcinogenesis and survival for liver-related death. The hazard ratio (HR) and 95% confidence interval (95% CI) was also calculated. Potential predictive factors associated with hepatocarcinogenesis and survival for liver-related death included the variables: sex, age, history of blood transfusion, family history of liver disease, lifetime cumulative alcohol intake, total bilirubin, AST, ALT, albumin, hemoglobin, platelet count, levels of viremia, and HCV subgroup according to HCV genotype in combination with aa substitution in core region. Variables that achieved statistical significance ($P < 0.05$) on univariate analysis were tested by multivariate Cox proportional hazard model to identify significant independent factors. Statistical comparisons were performed using the SPSS software (Chicago, IL). $P < 0.05$ by the two-tailed test were considered significant.




Fig. 1. Cumulative hepatocarcinogenesis rates according to HCV genotype in combination with amino acid substitutions in core region of HCV-1b. The rates were significantly different among the three HCV subgroups ($P < 0.001$; log-rank test). Especially, the rates in HCV-1b of Gln70(His70) were significantly higher than those in HCV-1b of Arg70 ($P = 0.028$; log-rank test) and HCV-2a/2b ($P < 0.001$; log-rank test), and the rates in HCV-1b of Arg70 were also significantly higher than those in HCV-2a/2b ($P < 0.001$; log-rank test).

Results

Hepatocarcinogenesis Rates and Survival Rates for Liver-Related Death in Patients Infected With HCV Who Had Not Received Antiviral Therapy. During the follow-up, 413 patients (35.0%) developed HCC. The cumulative hepatocarcinogenesis rates were 16.3, 34.3, 48.3, 58.7, and 69.1% at the end of 5, 10, 15, 20, and 25 years, respectively. The median interval between the initial visit and detection of HCC was 6.2 years (range, 0.1-31.7 years).

During the follow-up period, 243 patients (20.6%) died due to liver-related causes, and 97 of 243 (90.5%) developed HCC. The cumulative survival rates for liver-related death were 96.2, 84.8, 68.9, 55.0, and 46.1% at the end of 5, 10, 15, 20, and 25 years, respectively. The median interval between the initial visit and liver-related death was 10.1 years (range, 0.4-35.8 years).

Hepatocarcinogenesis Rates and Survival Rates for Liver-Related Death According to HCV Genotype in Combination with Amino Acid Substitutions in Core Region of HCV-1b. During the follow-up, 163 patients (51.3%), 175 (41.2%), and 75 (17.6%) developed HCC in HCV-1b of Gln70(His70), HCV-1b of Arg70, and HCV-2a/2b, respectively. In HCV-1b of Gln70(His70), HCV-1b of Arg70, and HCV-2a/2b, the cumulative hepatocarcinogenesis rates were 21.7, 19.3, 8.0% at the end of 5 years; 44.4, 39.4, 18.2% at the end of 10 years; 60.4, 52.7, 29.1% at the end of

15 years; 71.6, 60.3, 43.1% at the end of 20 years; and 87.1, 69.8, 46.9% at the end of 25 years, respectively. The rates were significantly different among the three HCV subgroups ($P < 0.001$) (Fig. 1). Especially, the rates in HCV-1b of Gln70(His70) were significantly higher than those in HCV-1b of Arg70 ($P = 0.028$) and HCV-2a/2b ($P < 0.001$), and the rates in HCV-1b of Arg70 were also significantly higher than those in HCV-2a/2b ($P < 0.001$).

During the follow-up, 104 patients (34.4%), 97 (23.4%), and 42 (10.0%) died due to liver-related causes in HCV-1b of Gln70(His70), HCV-1b of Arg70, and HCV-2a/2b, respectively. In HCV-1b of Gln70(His70), HCV-1b of Arg70, and HCV-2a/2b, the cumulative survival rates for liver-related death were 95.2, 95.4, 97.9% at the end of 5 years; 77.7, 83.3, 93.9% at the end of 10 years; 58.4, 68.4, 81.2% at the end of 15 years; 39.3, 58.4, 69.0% at the end of 20 years; and 33.8, 47.5, 59.5% at the end of 25 years, respectively. The rates were significantly different among the three HCV subgroups ($P < 0.001$) (Fig. 2). Especially, the rates in HCV-1b of Gln70(His70) were significantly lower than those in HCV-1b of Arg70 ($P = 0.016$) and HCV-2a/2b ($P < 0.001$), and the rates in HCV-1b of Arg70 were also significantly lower than those in HCV-2a/2b ($P < 0.001$).

Predictive Factors Associated with Hepatocarcinogenesis and Survival for Liver-Related Death in Patients Infected with HCV Who Had Not Received Antiviral Therapy. The data for the whole population




Fig. 2. Cumulative survival rates for liver-related death according to HCV genotype in combination with amino acid substitutions in the core region of HCV-1b. The rates were significantly different among the three HCV subgroups ($P < 0.001$; log-rank test). Especially, the rates in HCV-1b of Gln70(His70) were significantly lower than those in HCV-1b of Arg70 ($P = 0.016$; log-rank test) and HCV-2a/2b ($P < 0.001$; log-rank test), and the rates in HCV-1b of Arg70 were also significantly lower than those in HCV-2a/2b ($P < 0.001$; log-rank test).

Table 2. Factors associated with hepatocarcinogenesis in patients infected with HCV, who had not received antiviral therapy, identified by multivariate analysis

[Factors]	[Category]	Hazard ratio (95% confidence interval)	P
Gender	1: female	1	
	2: male	1.78 (1.44-2.21)	<0.001
Age (years)	1: <60	1	
	2: ≥60	1.68 (1.35-2.09)	<0.001
Albumin (g/dl)	1: ≥3.9	1	
	2: <3.9	1.94 (1.55-2.42)	<0.001
Platelet count (× 10 ⁴ /mm ³)	1: ≥15.0	1	
	2: <15.0	2.89 (2.25-3.72)	<0.001
Aspartate aminotransferase (IU/l)	1: <67	1	
	2: ≥67	1.92 (1.47-2.52)	<0.001
HCV subgroup	1: HCV-2a/2b	1	
	2: HCV-1b with Arg70	1.91 (1.42-2.55)	<0.001
	3: HCV-1b with Gln70(His70)	1.94 (1.45-2.61)	<0.001

Cox proportional hazard model

sample were analyzed to determine those factors that could predict hepatocarcinogenesis and survival for liver-related death.

Univariate analysis identified eight parameters that significantly correlated with hepatocarcinogenesis. These included gender (male; $P < 0.001$), age (≥ 60 years; $P < 0.001$), total bilirubin (≥ 1.2 mg/dL; $P < 0.001$), AST (≥ 67 IU/L; $P < 0.001$), ALT (≥ 85 IU/L; $P < 0.001$), platelet count ($< 15.0 \times 10^4/\text{mm}^3$; $P < 0.001$), albumin (< 3.9 g/dL; $P < 0.001$), and lifetime cumulative alcohol intake (≥ 500 kg; $P = 0.025$). Furthermore, the rates in HCV-1b of Gln70(His70) were significantly higher than those in HCV-1b of Arg70 ($P = 0.028$) and HCV-2a/2b ($P < 0.001$). These factors were entered into multivariate analysis, which then identified six parameters that significantly influenced hepatocarcinogenesis independently: gender (male; HR 1.78, $P < 0.001$), age (≥ 60 years; HR 1.68, $P < 0.001$), albumin (< 3.9 g/dL; HR 1.94, $P < 0.001$), platelet count ($< 15.0 \times 10^4/\text{mm}^3$; HR 2.89, $P < 0.001$), AST (≥ 67 IU/L; HR 1.92, $P < 0.001$), and HCV subgroup (HCV-1b of Gln70(His70); HR 1.94, $P = 0.001$) (Table 2).

Univariate analysis identified seven parameters that significantly correlated with survival for liver-related death. These included gender (male; $P < 0.001$), age (≥ 60 years; $P < 0.001$), total bilirubin (≥ 1.2 mg/dL; $P < 0.001$), AST (≥ 67 IU/L; $P < 0.001$), ALT (≥ 85 IU/L; $P < 0.001$), platelet count ($< 15.0 \times 10^4/\text{mm}^3$; $P < 0.001$), and albumin (< 3.9 g/dL; $P < 0.001$). Furthermore, the rates in HCV-1b of Gln70(His70)

were significantly lower than those in HCV-1b of Arg70 ($P = 0.016$) and HCV-2a/2b ($P < 0.001$). These factors were entered into multivariate analysis, which then identified six parameters that significantly influenced survival for liver-related death independently: gender (male; HR 1.91, $P < 0.001$), age (≥ 60 years; HR 1.61, $P = 0.001$), albumin (< 3.9 g/dL; HR 2.49, $P < 0.001$), platelet count ($< 15.0 \times 10^4/\text{mm}^3$; HR 3.69, $P < 0.001$), AST (≥ 67 IU/L; HR 4.16, $P < 0.001$), and HCV subgroup (HCV-1b of Gln70(His70); HR 2.16, $P < 0.001$) (Table 3).

IL28B Genotype and Time-Dependent Amino Acid Changes in Core Region of HCV-1b. Among 1,181 patients, 359 could be evaluated for changes over time of dominant amino acid by direct sequencing in core aa 70 of HCV-1b. Furthermore, among 359 patients, 142 could also be analyzed for the relationship between *IL28B* rs8099917 genotype and time-dependent changes of core aa 70.

In 199 patients of Arg70 at the initial visit, 34 patients (17.1%) changed from Arg70 to Gln70(His70) during the follow-up. Inversely, in 160 patients of Gln70(His70) at the initial visit, eight patients (5.0%) changed from Gln70(His70) to Arg70 during the follow-up. In change from Arg70 to Gln70(His70), and change from Gln70(His70) to Arg70, the cumulative change rates were 3.0, 0% at the end of 5 years; 16.8, 5.8% at the end of 10 years; 27.4, 11.5% at the end of 15 years; and 38.9, 16.7% at the end of 20 years, respectively. The cumulative change rates from Arg70 to Gln70(His70) were significantly higher than those from Gln70(His70) to Arg70 ($P = 0.002$).

In 78 patients of Arg70 and TT genotype at the initial visit, nine (11.5%) changed from Arg70 to Gln70(His70) during the follow-up. In 11 patients of Arg70 and non-TT genotype at the initial visit, seven (63.6%) changed from Arg70 to Gln70(His70) during the follow-up. In TT and non-TT genotype, the cumulative change rates were 0, 9.1% at the end of 5 years; 3.2, 65.4% at the end of 10 years; 14.8, 65.4% at the end of 15 years; and 29.0, 65.4% at the end of 20 years, respectively. The cumulative change rates in non-TT genotype were significantly higher than those in TT genotype ($P < 0.001$) (Fig. 3A).

In 30 patients of Gln70(His70) and TT genotype at the initial visit, three patients (10.0%) changed from Gln70(His70) to Arg70 during the follow-up. In 23 patients of Gln70(His70) and non-TT genotype at the initial visit, no patients changed from Gln70(His70) to Arg70 during the follow-up. In TT and non-TT genotype, the cumulative change rates were 0, 0% at