


15. Valsecchi M, Mauri L, Casellato R, Prioni S, Loberto N, et al. (2007) Ceramide and sphingomyelin species of fibroblasts and neurons in culture. *J Lipid Res* 48: 417–424.
16. Watanabe T, Sudoh M, Miyagishi M, Akashi H, Arai M, et al. (2006) Intracellular-diced dsRNA has enhanced efficacy for silencing HCV RNA and overcomes variation in the viral genotype. *Gene Ther* 13: 883–892.
17. Fujita T, Inoue K, Yamamoto S, Ikumoto T, Sasaki S, et al. (1994) Fungal inhibition of sphingomyelin synthesis reduces atherogenesis in *Isaria sinclairii* metabolite. *J Antibiot (Tokyo)* 47: 208–215.
18. Miyake Y, Kozutsumi Y, Nakamura S, Fujita T, Kawasaki T (1995) Serine palmitoyltransferase is the primary target of a sphingosine-like immunosuppressant, ISP-1/myriocin. *Biochem Biophys Res Commun* 211: 396–403.
19. Park TS, Panek RL, Mueller SB, Hanselman JC, Rosebury WS, et al. (2004) Inhibition of sphingomyelin synthesis reduces atherogenesis in apolipoprotein E-knockout mice. *Circulation* 110: 3465–3471.
20. Miyanari Y, Hijikata M, Yamaji M, Hosaka M, Takahashi H, et al. (2003) Hepatitis C virus non-structural proteins in the probable membranous compartment function in viral genome replication. *J Biol Chem* 278: 50301–50308.
21. Diamond DL, Jacobs JM, Paepfer B, Proll SC, Gritsenko MA, et al. (2007) Proteomic profiling of human liver biopsies: hepatitis C virus-induced fibrosis and mitochondrial dysfunction. *Hepatology* 46: 649–657.
22. Tardif KD, Mori K, Siddiqui A (2002) Hepatitis C virus subgenomic replicons induce endoplasmic reticulum stress activating an intracellular signaling pathway. *J Virol* 76: 7453–7459.
23. Pettus BJ, Chalfant CE, Hannun YA (2002) Ceramide in apoptosis: an overview and current perspectives. *Biochim Biophys Acta* 1585: 114–125.
24. Tepper AD, Ruurs P, Wiedmer T, Sims PJ, Borst J, et al. (2000) Sphingomyelin hydrolysis to ceramide during the execution phase of apoptosis results from phospholipid scrambling and alters cell-surface morphology. *J Cell Biol* 150: 155–164.
25. Liu YY, Han TY, Giuliano AE, Hansen N, Cabot MC (2000) Uncoupling ceramide glycosylation by transfection of glucosylceramide synthase antisense reverses adriamycin resistance. *J Biol Chem* 275: 7138–7143.
26. Taguchi Y, Kondo T, Watanabe M, Miyaji M, Umehara H, et al. (2004) Interleukin-2-induced survival of natural killer (NK) cells involving phosphatidylinositol-3 kinase-dependent reduction of ceramide through acid sphingomyelinase, sphingomyelin synthase, and glucosylceramide synthase. *Blood* 104: 3285–3293.
27. Diamond DL, Syder AJ, Jacobs JM, Sorensen CM, Walters KA, et al. (2010) Temporal proteome and lipidome profiles reveal hepatitis C virus-associated reprogramming of hepatocellular metabolism and bioenergetics. *PLoS Pathog* 6: e1000719.
28. Huitema K, van den Dikkenberg J, Brouwers JF, Holthuis JC (2004) Identification of a family of animal sphingomyelin synthases. *EMBO J* 23: 33–44.
29. Merrill AH, Jr., Schmelz EM, Dillehay DL, Spiegel S, Shayman JA, et al. (1997) Sphingolipids—the enigmatic lipid class: biochemistry, physiology, and pathophysiology. *Toxicol Appl Pharmacol* 142: 208–225.
30. Huwiler A, Kolter T, Pfeilschifter J, Sandhoff K (2000) Physiology and pathophysiology of sphingolipid metabolism and signaling. *Biochim Biophys Acta* 1485: 63–99.
31. Hannun YA, Luberto C, Argraves KM (2001) Enzymes of sphingolipid metabolism: from modular to integrative signaling. *Biochemistry* 40: 4893–4903.
32. van Genderen IL, Brandimarti R, Torrisi MR, Campadelli G, van Meer G (1994) The phospholipid composition of extracellular herpes simplex virions differs from that of host cell nuclei. *Virology* 200: 831–836.
33. Brugger B, Glass B, Haberkant P, Leibrecht I, Wieland FT, et al. (2006) The HIV lipidome: a raft with an unusual composition. *Proc Natl Acad Sci U S A* 103: 2641–2646.
34. Aizaki H, Morikawa K, Fukasawa M, Hara H, Inoue Y, et al. (2008) Critical role of virion-associated cholesterol and sphingolipid in hepatitis C virus infection. *J Virol* 82: 5715–5724.
35. Syed GH, Amako Y, Siddiqui A (2010) Hepatitis C virus hijacks host lipid metabolism. *Trends Endocrinol Metab* 21: 33–40.
36. Berger KL, Cooper JD, Heaton NS, Yoon R, Oakland TE, et al. (2009) Roles for endocytic trafficking and phosphatidylinositol 4-kinase III alpha in hepatitis C virus replication. *Proc Natl Acad Sci U S A* 106: 7577–7582.
37. Borawski J, Troke P, Puyang X, Gibaja V, Zhao S, et al. (2009) Class III phosphatidylinositol 4-kinase alpha and beta are novel host factor regulators of hepatitis C virus replication. *J Virol* 83: 10058–10074.
38. Tai AW, Benita Y, Peng LF, Kim SS, Sakamoto N, et al. (2009) A functional genomic screen identifies cellular cofactors of hepatitis C virus replication. *Cell Host Microbe* 5: 298–307.
39. Vaillancourt FH, Pilote L, Cartier M, Lippens J, Liuzzi M, et al. (2009) Identification of a lipid kinase as a host factor involved in hepatitis C virus RNA replication. *Virology* 387: 5–10.
40. Amemiya F, Maekawa S, Itakura Y, Kanayama A, Matsui A, et al. (2008) Targeting lipid metabolism in the treatment of hepatitis C virus infection. *J Infect Dis* 197: 361–370.
41. Ishitsuka R, Sato SB, Kobayashi T (2005) Imaging lipid rafts. *J Biochem* 137: 249–254.
42. Ramstedt B, Slotte JP (2002) Membrane properties of sphingomyelins. *FEBS Lett* 531: 33–37.
43. He Q, Suzuki H, Sharma N, Sharma RP (2006) Ceramide synthase inhibition by fumonisin B1 treatment activates sphingolipid-metabolizing systems in mouse liver. *Toxicol Sci* 94: 388–397.
44. Bligh EG, Dyer WJ (1959) A rapid method of total lipid extraction and purification. *Can J Biochem Physiol* 37: 911–917.
45. Inoue K, Umehara T, Ruegg UT, Yasui F, Watanabe T, et al. (2007) Evaluation of a cyclophilin inhibitor in hepatitis C virus-infected chimeric mice in vivo. *Hepatology* 45: 921–928.
46. Kashiwakuma T, Hasegawa A, Kajita T, Takata A, Mori H, et al. (1996) Detection of hepatitis C virus specific core protein in serum of patients by a sensitive fluorescence enzyme immunoassay (FEIA). *J Immunol Methods* 190: 79–89.
47. Ikeda K, Shimizu T, Taguchi R (2008) Targeted analysis of ganglioside and sulfatide molecular species by LC/ESI-MS/MS with theoretically expanded multiple reaction monitoring. *J Lipid Res* 49: 2678–2689.
48. Taguchi R, Nishijima M, Shimizu T (2007) Basic analytical systems for lipidomics by mass spectrometry in Japan. *Methods Enzymol* 432: 185–211.
49. Ikeda K, Oike Y, Shimizu T, Taguchi R (2009) Global analysis of triacylglycerols including oxidized molecular species by reverse-phase high resolution LC/ESI-QTOF MS/MS. *J Chromatogr B Analyt Technol Biomed Life Sci* 877: 2639–2647.
50. Ikeda K, Mutoh M, Teraoka N, Nakanishi H, Wakabayashi K, et al. (2011) Increase of oxidant-related triglycerides and phosphatidylcholines in serum and small intestinal mucosa during development of intestinal polyp formation in Min mice. *Cancer Sci* 102: 79–87.



Impairment of interferon regulatory factor-3 activation by hepatitis C virus core protein basic amino acid region 1

Kazuaki Inoue^{a,b}, Kyoko Tsukiyama-Kohara^{a,c,d}, Chiho Matsuda^a, Mitsutoshi Yoneyama^e, Takashi Fujita^f, Shusuke Kuge^g, Makoto Yoshiba^b, Michinori Kohara^{a,*}

^a Department of Microbiology and Cell Biology, The Tokyo Metropolitan Institute of Medical Science, 2-1-6 Kamikitazawa, Setagaya-ku, Tokyo 156-0057, Japan

^b Division of Gastroenterology, Showa University Fujigaoka Hospital, 1-30 Aoba-ku, Fujigaoka, Yokohama 227-8501, Japan

^c Department of Experimental Phylaxiology, Faculty of Life Sciences, Kumamoto University, 1-1-1 Honjo, Kumamoto City, Kumamoto 860-8556, Japan

^d Joint Faculty of Veterinary Medicine, Kagoshima University, 1-21-24 Korimoto, Kagoshima City, Kagoshima 890-0065, Japan

^e Medical Microbiology Research Center, Chiba University, 1-8-1 Inohana Chuo-Ku, Chiba-City, Chiba 260-8673, Japan

^f Laboratory of Molecular Genetics, Institute for Virus Research, Kyoto University, Kyoto 606-8507, Japan

^g Laboratory of Microbiology, Tohoku Pharmaceutical University, Komatsushima, Aoba-ku, Sendai 981-8558, Japan

ARTICLE INFO

Article history:

Received 17 October 2012

Available online 30 October 2012

Keywords:

Hepatitis C virus

IRF-3

Core protein

Type I interferon signaling

Basic amino acid region 1

Interferon beta

ABSTRACT

Interferon regulatory factor-3 (IRF-3), a key transcriptional factor in the type I interferon system, is frequently impaired by hepatitis C virus (HCV), in order to establish persistent infection. However, the exact mechanism by which the virus establishes persistent infection has not been fully understood yet. The present study aimed to investigate the effects of various HCV proteins on IRF-3 activation, and elucidate the underlying mechanisms. To achieve this, full-length HCV and HCV subgenomic constructs corresponding to structural and each of the nonstructural proteins were transiently transfected into HepG2 cells. IFN- β induction, plaque formation, and IRF-3 dimerization were elicited by Newcastle disease virus (NDV) infection. The expressions of IRF-3 homodimer and its monomer, Ser386-phosphorylated IRF-3, and HCV core protein were detected by immunofluorescence and western blotting. IFN- β mRNA expression was quantified by real-time PCR (RT-PCR), and IRF-3 activity was measured by the levels of IRF-3 dimerization and phosphorylation, induced by NDV infection or polyriboinosinic:polyribocytidylic acid [poly(I:C)]. Switching of the expression of the complete HCV genome as well as the core proteins, E1, E2, and NS2, suppressed IFN- β mRNA levels and IRF-3 dimerization, induced by NDV infection. Our study revealed a crucial region of the HCV core protein, basic amino acid region 1 (BR1), to inhibit IRF-3 dimerization as well as its phosphorylation induced by NDV infection and poly (I:C), thus interfering with IRF-3 activation. Therefore, our study suggests that rescue of the IRF-3 pathway impairment may be an effective treatment for HCV infection.

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

Hepatitis C virus (HCV), a flavivirus comprising a positive-sense, single-stranded RNA (ssRNA) of approximately 9.6 kb [1], causes persistent disease in infected individuals, possibly leading to chronic liver injury [2]. Despite the approximately 170 million individuals worldwide suffering from HCV infection that ranges from chronic hepatitis to hepatocellular carcinoma (HCC) [3,4], the exact mechanism by which the virus establishes persistent infection is not fully resolved.

The innate immune system is activated immediately upon infection as the first line of host defense against invading pathogens, with type I interferon (IFN) signaling being the crucial step

in the antiviral response [5]. The IFN system is, therefore, a prime target of HCV and other viruses in order to establish persistent infections [6], wherein the disruption of the type I IFN-activation pathway forms the most efficient strategy for HCV. Studies on HCV IFN-interference mechanisms have revealed that the HCV proteins NS5A and E2 selectively inhibit the double-stranded RNA-activated protein kinase (PKR) [7,8], an IFN-inducible antiviral molecule that controls transcription and translation [6]. IFN- β , a crucial molecule in type I IFN signaling, is regulated by several cellular factors associated with the activation of interferon regulatory factor-3 (IRF-3), leading to its rapid induction following viral infection [9,10]. However, IFN- β induction is impaired in HCV-infected cells, thus resulting in the disruption of IFN downstream signaling cascade [11].

IRF-3, a key constitutively expressed transcriptional factor localized in the cytoplasm in its inactive form [9], is activated upon

* Corresponding author. Fax: +81 3 5316 3137.

E-mail address: kohara-mc@igakuken.or.jp (M. Kohara).

phosphorylation, whereby it translocates to the nucleus to act as a transcriptional factor for positive regulatory domain (PRD) I of the IFN- β promoter. IRF-3 activation also induces phosphorylation of Ser385 and Ser386 or the serine/threonine (Ser/Thr) cluster between amino acids (aa) 396 and 405 (located at the C-terminus of IRF-3), a step that is essential for dimerization and nuclear translocation.

A previous study by Foy et al. showed that the NS3/4A serine protease derived from a subgenomic replicon participates in the suppression of the cellular pathway that activates IRF-3 [12]. The NS3/4A protein of HCV disrupts signaling of the double-stranded RNA (dsRNA) receptors, retinoic acid-inducible gene-I (RIG-I), and Toll-like receptor 3 (TLR3) by inducing proteolysis of interferon promoter stimulator-1 (IPS-1) [13–15] and Toll/interleukin-1 receptor (TIR) domain-containing adaptor protein inducing IFN- β (TRIF) [16], as well as by suppressing the downstream activation of IFN- β [13].

In view of the above observations and the emerging data on the role of HCV in regulating the IRF-3 pathway by additional

mechanisms, we aimed to investigate the effects of various HCV proteins on IRF-3 activation, and further elucidate the underlying novel mechanisms.

2. Material and methods

2.1. Transient expression of the HCV core proteins E1, E2, and NS3-4A

HepG2 cells were transfected to express E1, E2, or NS3-4A HCV core protein under the control of EF promoter (Invitrogen). The HCV core expression vectors were derived from HCR6 (genotype 1b), HCR24-12K (genotype 2a), or HCR24-12Q (genotype 2a). The E1, E2, and NS3 clones derived from HCR6 contained either the full-length cDNAs encoding the core protein or 1 of the 3 different deletions (deletion mutants), each of which lacked 1 of the 3 basic amino acid regions (BR), BR1 deletion (aa 4–14), BR2 deletion (aa 37–44), and BR3 deletion (aa 57–72) [17]. HepG2 cells were transfected with 4 μ g of the core cDNA (amino acids 1–191), E1 (amino




Fig. 1. (A) Structures of the conditional expression vectors for HCV RNAs and proteins. The cDNA clones that displayed highest level of homology to the consensus sequences among the 3 clones were used to construct *HCR6-Rz* (nt 1–9611). *HCR6-Fse* clone harbored a termination codon introduced at nucleotide 3606 and truncated *HCR6-Age* clone lacked nucleotides 155–1967. They were flanked with ribozyme (Rz) and hepatitis D virus ribozyme (HDV-Rz) sequences under the control of the CAG promoter in the Cre/loxP switching expression cassette, which consisted of the neomycin resistance gene, as a stuffer region flanked by the loxP sequence. (B) IFN- β mRNA levels in the cell lines *HCR6-Rz*, *HCR6-Fse*, and *HCR6-Age* before and after the expression of the HCV protein. The results are expressed as copy numbers per microgram of total RNA, as quantified by RT-PCR. (C) Efficiency of NDV plaque formation. Plaque assays were performed on Vero cells for NDV infectivity in *HCR6-Rz*, *HCR6-Fse*, or *HCR6-Age* before and after the expression of the HCV protein. The plaque numbers were counted 3 days after NDV inoculation. The ratio indicates the plaque numbers after the expression of the HCV genome divided by the plaque numbers before the expression of the HCV genome. (D) Suppression of IFN- β mRNA induction by HCV expression, 18 days after NDV inoculation. Day 0, before the expression of the HCV genome; Day 48, after the expression of the HCV genome. The results are expressed relative to the levels on Day 0 (100%) in each of the 3 HCV-expressing systems. p.i.h., post-inoculation hour.

acids 192–383), E2 (amino acids 384–809), NS3-4A (amino acids 1027–1711), or the core regions lacking the BR, in 35-mm dishes, by using Lipofectamine 2000 (Invitrogen) at 37 °C for 6 h. The medium was subsequently replaced with normal culture medium, and the cells were harvested after 48 h.

2.2. Newcastle disease virus (NDV) infection and addition of polyriboinosinic:polyribocytidylic acid [poly(I:C)]

NDV (Miyadera strain) was propagated from swabs by using the embryonated egg culture method, as described in the Supplementary methods.

Poly(I:C) (20 µg/well; GE Healthcare) was added to HepG2 cells in 35-mm dishes, 48 h after transfection with the core expression vector derived from HCR6 (genotype 1b).

2.3. Native PAGE of IRF-3 and phosphorylated IRF-3

Cells were lysed in 30 µL of lysis buffer (50 mM Tris-HCl [pH 8.0], 1% NP-40, 150 mM NaCl, 100 µg/mL leupeptin, 1 mM PMSF, 5 mM Na₃VO₄), mixed vigorously, and centrifuged at 15,000 rpm for 10 min; the supernatant was isolated. Total protein samples (10 µg) were electrophoresed on a 7.5% native PAGE gel (Bio-Rad Laboratories) and transferred onto a PVDF membrane. IRF-3 homodimer and its monomer and Ser386-phosphorylated IRF-3 were detected by western blotting using polyclonal rabbit antibodies, anti-human IRF-3 (1:1000) [18] and, anti-human Ser386-phosphorylated IRF-3 (1:10,000), respectively. Detection was achieved by enhanced chemiluminescence (ECL; Amersham, UK) according to the manufacturer's instructions. The rabbit anti-human IRF-3 and anti-human Ser386-phosphorylated IRF-3 antibodies were described previously [19].

2.4. Quantification of IFN-β gene expression

IFN-β mRNA expression was quantified by using real-time PCR (RT-PCR) as described previously [20] and Supplementary methods using the following primers and probes: sense (5'-CCATCTATGAGATGCTCCAGAA-3'), antisense (5'-TTTCTCCAGGACTGTCTTCA-GA-3') and probe (5'-AGCACTGGCTGGAATGAGACTATTGTTG-3').

3. Results

3.1. Induction of IFN and IRF-3 by HCV-Rz

To evaluate the effect of HCV gene persistent expression, cell lines expressing the HCV genomes in a Cre/loxP expression system [20] were established by transfecting the full-genome HCV (*HCR6-Rz*); core, E1, E2, and NS2 (*HCR6-Fse*); and E2~NS5b (*HCR6-Age*) (Fig. 1A) clones into HepG2 cells. Of the 3 HCV expression systems, only *HCR6-Rz* transiently induced endogenous IFN-β expression (Fig. 1B). No endogenous IFN-β was detected by RT-PCR on Day 0 or Day 48 (long culture; LC) [21] in any of the 3 systems (Fig. 1B).

3.2. Effect of HCV expression on NDV infection and IFN-β induction

Further, we examined whether the persistent expression of HCV genome influenced the plaque formation activity of NDV in *HCR6-Rz*-, *HCR6-Fse*-, and *HCR6-Age*-expressing cell lines. Prior to the expression of HCV proteins, the plaque numbers were similar across all the cell lines (Fig. 1C). Expression of HCV proteins for more than 48 days [21] increased plaque numbers in the *HCR6-Rz*- and *HCR6-Fse*-expressing cells from 7- to 8-fold (Fig. 1C); however, plaque numbers in the *HCR6-Age*-expressing cells remained constant. These findings thus indicate that HCV structural proteins

interfered with the induction of IFN-β mRNA, after NDV infection (18 h post-infection). Therefore, we measured IFN-β mRNA levels in the 3 cell lines, *HCR6-Rz*, *HCR6-Fse*, and *HCR6-Age*, by RT-PCR before (Day 0) and after (Day 48) inoculation with NDV (Fig. 1D). The mRNA expression of IFN-β was not observed prior to NDV infection in any of the 3 cell lines (Day 0). Notably, after 18 h of NDV inoculation and prior to the expression of various HCV proteins (Day 0), the levels of IFN-β mRNA transcription were similar among *HCR6-Rz*-, *HCR6-Fse*-, and *HCR6-Age*-expressing cell lines (Fig. 1D). Following HCV protein expression, the induced IFN-β mRNA expression was reduced to 20% in both *HCR6-Rz*- and *HCR6-Fse*-expressing cell lines. However, IFN-β mRNA expression remained constant in the *HCR6-Age*-expressing cell line.




Fig. 2. (A) HCV inhibition of IRF-3 nuclear translocation. IRF-3 is a constitutively expressed transcriptional factor that localizes in the cytoplasm in a diffuse manner, when inactive. After NDV inoculation and prior to the expression of the HCV genome, IRF-3 translocated to the nucleus but was retained at a perinuclear site in the *HCR6-Rz*-expressing cells (Day 48). IRF-3 colocalized with the HCV core protein. (B–D) Suppression of IRF-3 dimerization by HCV expression. The dimeric and monomeric forms of IRF-3 were detected by western blotting on native PAGE gels. The influence of (B) *HCR6-Rz* expression, (C) *HCR6-Fse* expression, and (D) *HCR6-Age* expression on IRF-3 dimerization is shown. NDV was used to induce IRF-3 dimerization.




Fig. 3. (A) Structures of the HCR6 core, E1, and E2 expression vectors encoding the HCV core (aa 1–191), E1 (amino acids 192–383), and E2 (amino acids 384–809) proteins, respectively, under the control of the EF1 promoter. (B) Left panel, IRF-3 dimerization induced by NDV before transfection with the expression vectors; right panel, IRF-3 dimerization induced by NDV after transfection with the expression vectors. (C) IFN- β mRNA induction levels at 12 and 18 h after NDV inoculation into HepG2 cells transfected with the vector alone, core, E1, or E2 proteins. The results are expressed relative to the induction levels of IFN- β in HepG2 cells transfected with the vector alone (100%), for post-inoculation, each time. (D) IRF-3 dimerization on administration of 10 mg/mL poly(I:C) before the expression (Day 0), and at 2, 4, 8, and 12 h after the expression of the HCV core (pEF-Core) or vector plasmid DNA. (E) Effects of the expression of vector (lane 1), R6 core (lane 2), R24 core (lane 3), R24 core (Q12K) (lane 4), R6-NS3-4A (lane 5), N1-NS3-4A (lane 6), and mock control (lane 7) on IRF-3 dimerization (upper column a) and phosphorylation of serine residue at amino acid 386 in IRF-3 (lower column b), after infection with NDV for 18 h. F. Effects of the expressions of R6-C, R24-C, R24-C(Q12K), and R6-NS3 on IFN- β induction, 18 h after NDV inoculation. The IFN- β mRNA levels were assayed by RT-PCR. The results are expressed relative to the induction levels of IFN- β in HepG2 cells transfected with the vector alone (100%).

3.3. Effect of HCV on IRF-3 localization, nuclear translocation, and dimerization by NDV

The effect of HCV expression on cellular localization of IRF-3 was analyzed in HCR6-Rz-expressing cells infected with NDV

before (Day 0) and after LC (Fig. 2A). Prior to NDV infection, IRF-3 was detected in the cytoplasm by immunofluorescence. Notably, after 18 h of NDV inoculation and prior to HCR6-Rz protein expression, when IFN- β induction and IRF-3 dimerization were maximal, a substantial amount of IRF-3 translocated to the nucleus.

However, this nuclear translocation was suppressed in the presence of HCV proteins (Fig. 2A) and resulted in the co-localization of the HCV core protein with IRF-3 at perinuclear sites (Fig. 2A, superimposed image of IRF-3 and core protein immunostaining).

To elucidate the mechanism underlying the suppression of IFN- β mRNA in *HCR6-Rz-* and *HCR6-Fse*-expressing cells, we examined the effect of HCV expression on IRF-3 dimerization after NDV infection (Fig. 2B). Interestingly, the levels of IRF-3 dimerization peaked at 12–18 h after NDV infection in the 3 cell lines lacking HCV expression (Day 0; Fig. 2). However, in the *HCR6-Rz-* and *HCR6-Fse*-expressing cell lines, IRF-3 dimerization was found to be significantly reduced, (Fig. 2B and C) when compared to that in the *HCR6-Age*-expressing cells (Fig. 2D).

3.4. Identification of the HCV genome region responsible for the inhibition of IRF-3 dimerization and IFN- β induction

To identify the HCV genome region responsible for suppression of IRF-3 dimerization, HepG2 cells were transfected to express the HCV core regions derived from HCR6, E1, or E2 (genotype 1b; Fig. 3A). Protein expression was confirmed by western blotting (data not shown). The HCV core protein suppressed IRF-3 dimerization, but E1 and E2 expressions had no effect on the dimerization (Fig. 3B). Expression of E1, E2, or the vector alone did not alter the levels of IFN- β mRNA induced by NDV infection in HepG2 cells (Fig. 3C), but significantly reduced IFN- β mRNA levels at both 12 and 18 h after infection (Fig. 3C).

3.5. Effect of HCV core protein expression on IRF-3 dimerization through TLR3

Among the synthetic dsRNAs, poly(I:C) is a potent inducer of IFN- β through TLR3. Accordingly, HepG2 cells transfected with poly(I:C) and the vector control (pEF1-vector) showed IRF-3 dimerization (Fig. 3D). In contrast, IRF-3 dimerization was suppressed in HepG2 cells expressing the HCV core protein albeit the induction of IFN- β mRNA following poly(I:C) expression (data not shown).

3.6. Effect of the HCV core protein NS3 and core proteins derived from genotype 2a on IRF-3 dimerization, compared to proteins derived from genotype 1b

Further, we investigated whether the HCV core protein NS3 and core proteins derived from other genotypes exerted the same effects on IRF-3 after 18 h of NDV infection. The core proteins derived from genotypes 1b (R6) and 2a (R24-12Q and R24-12K) suppressed IRF-3 dimerization in cells infected with NDV (Fig. 3E(a)). In contrast, IRF-3 dimerization remained unaltered in the presence of R6 clone NS3 protein, but was suppressed by the N clone NS3 protein. Thus, NS3-4A protein of R6 clone suppressed IRF-3 dimerization to a relatively lesser extent compared to that of the N1 strain. Similar results were obtained for the phosphorylation at Ser386 in IRF-3 (Fig. 3E(b)). The IFN- β mRNA transcription was quantified in HepG2 cells by RT-PCR after transfection with these expression vectors (Fig. 3F).

3.7. Identification of the HCV core region responsible for suppressing IRF-3 dimerization

We sought to identify the region of the HCV core protein responsible for suppressing IRF-3 dimerization. Expression vectors encoding the entire HCV core or the core region lacking 1 of the 3 basic amino acid regions (BR) that influenced nuclear translocation [17] were transfected into HepG2 cells, and the effects on IRF-3 dimerization were examined (Fig. 4A). Protein




Fig. 4. (A) Structures of the HCR6 core, E1, and E2 expression vectors carrying the complete core, BR1 deletion (aa 4–14), BR2 deletion (aa 37–44), and BR3 deletion (aa 57–72). (B) Western blotting to confirm the expression of the mutated core proteins. (C) Effects of the expression of various mutated core proteins on IRF-3 dimerization and IRF-3 phosphorylation at Ser386, 18 h after NDV inoculation. (D) Effects of the expression of each type of core protein region on IFN- β mRNA synthesis, 18 h after NDV inoculation. The results are expressed relative to the induction levels of IFN- β in HepG2 cells transfected with the vector alone (100%). IFN- β mRNA levels were assayed by RT-PCR.

expression of the core and the deletion mutants (BR1, BR2, and BR3) was confirmed by western blotting (Fig. 4B). IRF-3 dimerization, phosphorylation at Ser386 of IRF-3, and induction of IFN- β mRNA were suppressed in HepG2 cells expressing the entire core, a deletion of BR2, or a deletion of BR3 (Fig. 4C), but not in cells expressing the BR1-deleted HCV core regions (Fig. 4C and D).

4. Discussion

The present study indicates that the HCV core protein inhibits IRF-3 dimerization, IRF-3 phosphorylation at Ser386, and IFN- β induction. In addition, our study showed that the effect of the core protein derived from genotype 1b was similar to that of the core protein derived from genotype 2a, indicating that the inhibitory effect of the core protein might be effective in several genotypes of HCV. These findings are corroborated by a previous study by Foy et al. [12] who showed that HCV NS3/4A disrupts virus-associated-kinase-mediated IRF-3 activation, which further results in the suppression of IRF-3 phosphorylation, nuclear translocation, and IRF-3-dependent ISRE/PRDI activation. These findings indicate that attenuation of the IFN system was achieved through NS3/4A proteins via the interference of IRF-3 activation, thus strengthening our results, which show the potential of HCV core protein to interfere with IRF-3 activation in promoting persistent infection.

Furthermore, the present study showed that the N-terminal region of the core protein and BR-1 domain in particular are responsible for inactivating IRF-3. The N-terminal region (amino acids 1–59) of the HCV core protein has been identified as the binding region for a DEAD box protein (DDX3) [22]. Human DDX3, a putative RNA helicase, is a member of the highly conserved DEAD box subclass that includes the expression of murine PL10, *Xenopus* An3, and yeast Ded 1 proteins. Recently, expression of DDX3 was found to enhance IFN- β promoter induction by TBK1/IKK ϵ , whereas silencing of DDX3 inhibited IFN- β promoter and virus- or dsRNA-induced IRF-3 activation [23]. It was shown that Vaccinia virus K7 protein also binds to DDX3 and inhibits pattern recognition receptor-induced IFN- β induction by preventing TBK1/IKK ϵ -mediated IFN- β induction via impaired TBK1/IKK ϵ -induced activation of IRF-3 [23]. A previous study by Oshiumi et al. showed that DDX3 C-terminal region (amino acids 622–662) directly binds to the IFN-beta promoter stimulator-1 (IPS-1) CARD-like domain [24] as well as the N-terminal HCV core protein [36]. The present study demonstrated that the expression of the core protein decreased the levels of DDX3 expression (data not shown). This is in agreement with the result of a previous study, which showed that DDX3 is downregulated in HCV-associated hepatocellular carcinoma (HCC) and silencing of DDX3 accelerates cell growth [25]. Collectively, these findings suggest that DDX3 may be the target of the core protein for inhibiting IRF-3 activation.

In conclusion, our study revealed a crucial region of the HCV core protein, basic amino acid region 1, to interfere with IRF-3 activation and thereby inhibit the IFN signaling cascades. Therefore, the inhibitory effects that result in the IRF-3 pathway impairment could be rescued by deleting the basic region 1 of core protein, thus suggesting that it might be an effective treatment for HCV infection. Future studies involving DDX3 modification by the HCV core protein may be interesting to explore the cell growth-dysregulation mechanisms.

Acknowledgments

This study was supported in part by a grant from the Ministry of Education, Culture, Sports, Science and Technology of Japan; a grant from the Ministry of Health, Labour and Welfare of Japan; and the Program for Promotion of Fundamental Studies in Health Sciences of the National Institute of Biomedical Innovation of Japan; and the Cooperative Research Project on Clinical and

Epidemiological Studies of Emerging and Re-emerging Infectious Diseases.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.bbrc.2012.10.079>.

References

- [1] N. Kato, M. Hijikata, Y. Ootsuyama, et al., Molecular cloning of the human hepatitis C virus genome from Japanese patients with non-A, non-B hepatitis, *Proc. Natl. Acad. Sci. USA* 87 (1990) 9524–9528.
- [2] L.B. Seeff, F.B. Hollinger, H.J. Alter, et al., Long-term mortality and morbidity of transfusion-associated non-A, non-B, and type C hepatitis: a National Heart, Lung, and Blood Institute collaborative study, *Hepatology* 33 (2001) 455–463.
- [3] H.B. El-Serag, Hepatocellular carcinoma and hepatitis C in the United States, *Hepatology* 36 (2002) S74–S83.
- [4] D. Thomas, The hepatitis C viruses, in: C. Hagedorn, C. Rice (Eds.), *Hepatitis C Epidemiology*, Springer-Verlag, Berlin, 1999, pp. 25–42.
- [5] G.R. Foster, Interferons in host defense, *Semin. Liver Dis.* 17 (1997) 287–295.
- [6] R. Kaufman, Translation control of genome expression, in: N. Sonenberg, J. Hershey, M. Matthews (Eds.), *The double strand RNA-activated protein kinase PKR*, Cold Spring Harbor, New York, 2000, pp. 503–527.
- [7] M.J. Gale Jr., M.J. Korth, N.M. Tang, et al., Evidence that hepatitis C virus resistance to interferon is mediated through repression of the PKR protein kinase by the nonstructural 5A protein, *Virology* 230 (1997) 217–227.
- [8] D.R. Taylor, S.T. Shi, P.R. Romano, et al., Inhibition of the interferon-inducible protein kinase PKR by HCV E2 protein, *Science* 285 (1999) 107–110.
- [9] M. Yoneyama, W. Suhara, Y. Fukuhara, et al., Direct triggering of the type I interferon system by virus infection: activation of a transcription factor complex containing IRF-3 and CBP/p300, *Embo. J.* 17 (1998) 1087–1095.
- [10] J. Hiscott, Triggering the innate antiviral response through IRF-3 activation, *J. Biol. Chem.* 282 (2007) 15325–15329.
- [11] E.F. Meurs, A. Breiman, The interferon inducing pathways and the hepatitis C virus, *World J. Gastroenterol.* 13 (2007) 2446–2454.
- [12] E. Foy, K. Li, C. Wang, et al., Regulation of interferon regulatory factor-3 by the hepatitis C virus serine protease, *Science* 300 (2003) 1145–1148.
- [13] E. Meylan, J. Curran, K. Hofmann, et al., Cardif is an adaptor protein in the RIG-I antiviral pathway and is targeted by hepatitis C virus, *Nature* 437 (2005) 1167–1172.
- [14] X.D. Li, L. Sun, R.B. Seth, et al., Hepatitis C virus protease NS3/4A cleaves mitochondrial antiviral signaling protein off the mitochondria to evade innate immunity, *Proc. Natl. Acad. Sci. USA* 102 (2005) 17717–17722.
- [15] R. Lin, J. Lacoste, P. Nakhaei, et al., Dissociation of a MAVS/IPS-1/VISA/Cardif-IKKeppilon molecular complex from the mitochondrial outer membrane by hepatitis C virus NS3-4A proteolytic cleavage, *J. Virol.* 80 (2006) 6072–6083.
- [16] K. Li, E. Foy, J.C. Ferreón, et al., Immune evasion by hepatitis C virus NS3/4A protease-mediated cleavage of the Toll-like receptor 3 adaptor protein TRIF, *Proc. Natl. Acad. Sci. USA* 102 (2005) 2992–2997.
- [17] R. Suzuki, Y. Matsuura, T. Suzuki, et al., Nuclear localization of the truncated hepatitis C virus core protein with its hydrophobic C terminus deleted, *J. Gen. Virol.* 76 (Pt 1) (1995) 53–61.
- [18] W. Suhara, M. Yoneyama, I. Kitabayashi, et al., Direct involvement of CREB-binding protein/p300 in sequence-specific DNA binding of virus-activated interferon regulatory factor-3 holocomplex, *J. Biol. Chem.* 277 (2002) 22304–22313.
- [19] M. Mori, M. Yoneyama, T. Ito, et al., Identification of Ser-386 of interferon regulatory factor 3 as critical target for inducible phosphorylation that determines activation, *J. Biol. Chem.* 279 (2004) 9698–9702.
- [20] K. Tsukiyama-Kohara, S. Tone, I. Maruyama, et al., Activation of the CKI-CDK-Rb-E2F pathway in full genome hepatitis C virus-expressing cells, *J. Biol. Chem.* 279 (2004) 14531–14541.
- [21] T. Nishimura, M. Kohara, K. Izumi, et al., Hepatitis C virus impairs p53 via persistent overexpression of 3beta-hydroxysterol Delta24-reductase, *J. Biol. Chem.* 284 (2009) 36442–36452.
- [22] A.M. Owsianka, A.H. Patel, Hepatitis C virus core protein interacts with a human DEAD box protein DDX3, *Virology* 257 (1999) 330–340.
- [23] M. Schroder, M. Baran, A.G. Bowie, Viral targeting of DEAD box protein 3 reveals its role in TBK1/IKKeppilon-mediated IRF activation, *EMBO J.* 27 (2008) 2147–2157.
- [24] H. Oshiumi, K. Sakai, M. Matsumoto, T. Seya, DEAD/H BOX 3 (DDX3) helicase binds the RIG-I adaptor IPS-1 to up-regulate IFN-beta-inducing potential, *Eur. J. Immunol.* 40 (4) (2010) 940–948.
- [25] P.C. Chang, C.W. Chi, G.Y. Chau, et al., DDX3, a DEAD box RNA helicase, is deregulated in hepatitis virus-associated hepatocellular carcinoma and is involved in cell growth control, *Oncogene* 25 (2006) 1991–2003.

Immunization with a Recombinant Vaccinia Virus That Encodes Nonstructural Proteins of the Hepatitis C Virus Suppresses Viral Protein Levels in Mouse Liver

Satoshi Sekiguchi¹, Kiminori Kimura², Tomoko Chiyo¹, Takahiro Ohtsuki¹, Yoshimi Tobita¹, Yuko Tokunaga¹, Fumihiko Yasui¹, Kyoko Tsukiyama-Kohara³, Takaji Wakita⁴, Toshiyuki Tanaka⁵, Masayuki Miyasaka⁶, Kyosuke Mizuno⁷, Yukiko Hayashi⁸, Tsunekazu Hishima⁸, Kouji Matsushima⁹, Michinori Kohara^{1*}

1 Department of Microbiology and Cell Biology, Tokyo Metropolitan Institute of Medical Science, Setagaya-ku, Tokyo, Japan, **2** Division of Hepatology, Tokyo Metropolitan Komagome Hospital, Bunkyo-ku, Tokyo, Japan, **3** Transboundary Animal Diseases Center, Joint Faculty of Veterinary Medicine, Kagoshima University, Korimoto, Kagoshima, Japan, **4** Department of Virology II, National Institute of Infectious Diseases, Shinjuku-ku, Tokyo, Japan, **5** Laboratory of Immunobiology, Department of Pharmacy, School of Pharmacy, Hyogo University of Health Sciences, Chuo-ku, Kobe, Japan, **6** Laboratory of Immunodynamics, Department of Microbiology and Immunology, Osaka University Graduate School of Medicine, Suita, Osaka, Japan, **7** Chemo-Sero-Therapeutic Research Institute, Okubo, Kumamoto, Japan, **8** Department of Pathology, Tokyo Metropolitan Komagome Hospital, Bunkyo-ku, Tokyo, Japan, **9** Department of Molecular Preventive Medicine, School of Medicine, University of Tokyo, Bunkyo-ku, Tokyo, Japan

Abstract

Chronic hepatitis C, which is caused by infection with the hepatitis C virus (HCV), is a global health problem. Using a mouse model of hepatitis C, we examined the therapeutic effects of a recombinant vaccinia virus (rVV) that encodes an HCV protein. We generated immunocompetent mice that each expressed multiple HCV proteins via a *Cre/loxP* switching system and established several distinct attenuated rVV strains. The HCV core protein was expressed consistently in the liver after polyinosinic acid-polycytidylic acid injection, and these mice showed chronic hepatitis C-related pathological findings (hepatocyte abnormalities, accumulation of glycogen, steatosis), liver fibrosis, and hepatocellular carcinoma. Immunization with one rVV strain (rVV-N25), which encoded nonstructural HCV proteins, suppressed serum inflammatory cytokine levels and alleviated the symptoms of pathological chronic hepatitis C within 7 days after injection. Furthermore, HCV protein levels in liver tissue also decreased in a CD4 and CD8 T-cell-dependent manner. Consistent with these results, we showed that rVV-N25 immunization induced a robust CD8 T-cell immune response that was specific to the HCV nonstructural protein 2. We also demonstrated that the onset of chronic hepatitis in CN2-29^(+/-)/MxCre^(+/-) mice was mainly attributable to inflammatory cytokines, (tumor necrosis factor) TNF- α and (interleukin) IL-6. Thus, our generated mice model should be useful for further investigation of the immunological processes associated with persistent expression of HCV proteins because these mice had not developed immune tolerance to the HCV antigen. In addition, we propose that rVV-N25 could be developed as an effective therapeutic vaccine.

Citation: Sekiguchi S, Kimura K, Chiyo T, Ohtsuki T, Tobita Y, et al. (2012) Immunization with a Recombinant Vaccinia Virus That Encodes Nonstructural Proteins of the Hepatitis C Virus Suppresses Viral Protein Levels in Mouse Liver. PLoS ONE 7(12): e51656. doi:10.1371/journal.pone.0051656

Editor: Naglaa H. Shoukry, University of Montreal, Canada

Received: March 13, 2012; **Accepted:** November 5, 2012; **Published:** December 17, 2012

Copyright: © 2012 Sekiguchi et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was supported by grants from the Ministry of Education, Culture, Sports, Science, and Technology of Japan; the Program for Promotion of Fundamental Studies in Health Sciences of the Pharmaceuticals and Medical Devices Agency of Japan; and the Ministry of Health, Labor, and Welfare of Japan. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: kohara-mc@igakuken.or.jp

Introduction

Hepatitis C virus (HCV) is a major public health problem; approximately 170 million people are infected with HCV worldwide [1]. HCV causes persistent infections that can lead to chronic liver diseases such as chronic hepatitis, liver cirrhosis, and hepatocellular carcinoma (HCC) [2]. Antiviral drugs are not highly effective in individuals with a chronic infection; furthermore, an effective vaccine against HCV has not been developed. A convenient animal model of HCV infection will greatly facilitate the development of an effective HCV vaccine.

Transgenic mice that express HCV proteins have been generated to study HCV expression [3,4]; however, in each of

these cases, the relevant transgene is expressed during embryonic development; therefore, the transgenic mice become immunotolerant to the transgenic products, and consequently, the adult mice are not useful for investigations of the pathogenesis of chronic hepatitis C. To address this problem, we developed a system that can drive conditional expression of an HCV transgene; our system involves the *Cre/loxP* system and a recombinant adenovirus capable of expressing Cre recombinase [5,6]. Concerns have been expressed that an adenovirus and transient expression of HCV proteins could induce immune responses [5] and, therefore, obscure any evidence of the effect of the host immune responses on chronic liver pathology. Therefore, here, we used a *Cre/loxP* switching system to generate an immunocompetent mouse model

of HCV protein expression; with this system, we could study the host immune responses against HCV proteins.

Folgori et al. (2006) reported effective vaccination of chimpanzees with an adenoviral vector and plasmid DNA encoding the HCV nonstructural region. This technique protected the liver tissues from acute hepatitis, which results when whole animals are challenged with virus [7]. However, this vaccine has not yet been shown to be effective against chronic HCV infection.

Here, we aimed to address how HCV expression causes chronic liver diseases and to provide new options for HCV vaccine development. Using LC16m8, a highly attenuated strain of vaccinia virus (VV), we generated three recombinant vaccinia viruses (rVVs) that each encoded one of three different HCV proteins and found that one recombinant virus (rVV-N25), which encoded nonstructural HCV proteins, resolved pathological chronic hepatitis C symptoms in the liver. We also found that immunization with rVV-N25 suppressed HCV core protein levels in the livers of transgenic mice; moreover, this suppression was mediated by CD4 and CD8 T cells, as has been previously reported [8].

Results

Generation of a Model of Persistent HCV Protein Expression

To produce adult mice that express an HCV transgene, we bred CN2-29 transgenic mice, which carry an HCV transgene, [5,6,9] with Mx1-Cre transgenic mice [10], which express Cre recombinase in response to interferon (IFN)- α or a chemical inducer of IFN- α , poly(I:C) (Figure 1A). Following poly(I:C) injection, the HCV transgene was rearranged, and HCV sequences were expressed in the livers of F1 progeny (CN2-29^(+/-)/MxCre^(+/-) mice) within 7 days after poly(I:C) injection (Figure 1B).

To evaluate the characteristic features of these CN2-29^(+/-)/MxCre^(+/-) mice, we analyzed serum alanine aminotransferase (ALT) and liver HCV core protein levels after poly(I:C) injection. As illustrated in Figure 1C, serum ALT levels increased and reached a peak at 24 h after the first poly(I:C) injection; this elevation appeared to be a direct result of the poly(I:C) treatment, which causes liver injury [11]. After this peak, serum ALT levels dropped continuously until day 4, and then ALT levels began to increase, as did HCV core protein levels. Thereafter, the HCV core protein was expressed consistently for at least 600 days.

Histological analysis showed HCV core protein expression in most hepatocytes of the transgenic mice; these mice showed evidence of lymphocytic infiltration that was caused by the HCV core proteins (Figure 1D and E). These observations, in addition to the modified histology activity index (HAI) scores, indicated that expression of HCV proteins caused chronic hepatitis in the CN2-29^(+/-)/MxCre^(+/-) mice because a weak, though persistent, immune response followed an initial bout of acute hepatitis (Figure S1). Moreover, we observed a number of other pathological changes in these mice – including swelling of hepatocytes, abnormal architecture of liver-cell cords, abnormal accumulation of glycogen, steatosis, fibrosis, and HCC (Figures 1E and F, Table S1). Steatosis was mild in the younger mice (day 21) and became increasingly severe over time (days 120 and 180; Figure S2). Importantly, none of the pathological changes were observed in the CN2-29^(+/-)/MxCre^(-/-) mice after poly(I:C) injection (Figure 1F).

Recombinant Vaccinia Virus Immunization in HCV Transgenic Mice

To determine whether activation of the host immune response caused the reduction with HCV protein levels in the livers of CN2-29^(+/-)/MxCre^(+/-) mice, we used a highly attenuated VV strain, LC16m8, to generate three rVVs [12]. Each rVV encoded a different HCV protein; rVV-CN2 encoded mainly structural proteins, rVV-N25 encoded nonstructural proteins, and rVV-CN5 encoded the entire HCV protein region (Figure 2A). Because rVVs can express a variety of proteins and induce strong and long-term immunity, they have been evaluated as potential prophylactic vaccines [13].

We used western blots to confirm that each HCV protein was expressed in cell lines. Each of seven proteins – the core, E1, E2, NS3-4A, NS4B, NS5A, and NS5B – was recognized and labeled by a separate cognate antibody directed (Figure S3). To induce effective immune responses against HCV proteins in transgenic mice, we injected an rVV-HCV (rVV-CN2, rVV-CN5, or rVV-N25) or LC16m8 (as the control) intradermally into CN2-29^(+/-)/MxCre^(+/-) mice 90 days after poly(I:C) injection (Figure 2B). Analysis of liver sections 7 days after immunization with rVV-N25 revealed dramatic improvement in a variety of pathological findings associated with chronic hepatitis – including piecemeal necrosis, hepatocyte swelling, abnormal architecture of liver-cell cords, abnormal accumulation of glycogen, and steatosis (Figures 2C–E). Collectively, these results demonstrated that only the rVV-N25 treatment resulted in histological changes indicative of improvement in the chronic hepatitis suffered by the transgenic mice.

To determine whether rVV-N25 treatment induced the same effect in other strains of HCV transgenic mice, we analyzed RzCN5-15^(+/-)/MxCre^(+/-) mice, which express all HCV proteins; in these mice, chronic hepatitis was resolved within 28 days of immunization with rVV-N25. Taken together, these findings indicated that rVV-N25 had a dramatic therapeutic effect on both types of HCV transgenic mice (Figure S4).

Treatment with rVV-N25 Reduced the HCV Core Protein Levels in the Livers

To assess in detail the effects of rVV-HCV immunization on HCV protein clearance from the livers of CN2-29^(+/-)/MxCre^(+/-) mice, we monitored the levels of HCV core protein in liver samples via ELISA. We found that within 28 days after immunization the HCV core protein levels were significantly lower in livers of rVV-N25-treated mice than in those of control mice (Figure 3A). Immunohistochemical analysis indicated that, within 28 days after immunization, levels of HCV core protein were substantially lower in the livers of CN2-29^(+/-)/MxCre^(+/-) mice than in those of control mice (Figure 3B). Importantly, neither resolution of chronic hepatitis nor reduction in the HCV protein levels was observed in the mice treated with LC16m8, rVV-CN2, or rVV-CN5. These results indicated that HCV non-structural proteins might be important for effects of therapeutic vaccines. In contrast, rVV-CN5 which encoded HCV structural and non-structural proteins did not show any significant effects. These results indicated that HCV structural proteins might have inhibited the therapeutic effects of the non-structural proteins. Therefore, it may be important to exclude the HCV structural proteins (aa 1–541) as antigenic proteins when developing therapeutic vaccines against chronic hepatitis C.

In addition, we measured serum ALT levels in CN2-29^(+/-)/MxCre^(+/-) mice from all four treatment groups 28 days after rVV-HCV immunization. Serum ALT levels were not significant-




Figure 1. Pathogenesis in immunocompetent mice with persistent HCV expression. (A) Structure of CN2-29^(+/-)/MxCre^(+/-) and the Cre-mediated activation of the transgene unit. R6CN2 HCV cDNA was cloned downstream of the CAG promoter, neomycin-resistant gene (*neo*), and poly A (pA) signal flanked by two *loxP* sequences. This cDNA contains the core, E1, E2, and NS2 regions. (B) Cre-mediated genomic DNA recombination. After poly(I:C) injection, genomic DNA was extracted from liver tissues and analyzed by quantitative RTD-PCR for Cre-mediated transgenic recombination. The transgene was almost fully recombined in transgenic mouse livers 7 days after the injection. In all cases, n = 3 mice per group. (C) HCV core protein expression was sustained for at least 600 days after poly(I:C) injection. (D) Immunohistochemical analysis revealed that most hepatocytes expressed the HCV core protein within 6 days after injection. (E) Liver sections from CN2-29^(+/-)/MxCre^(+/-) mice after the poly(I:C) injection. Infiltrating lymphocytes (arrows) were observed on days 6 and 180; Hepatocellular carcinoma (HCC) was observed on day 360. In contrast, these pathological changes were not observed in CN2-29^(+/-)/MxCre^(-/-) mice after the injection. The inset image shows abnormal mitosis in a tumor cell. (F) Hepatocyte swelling and abnormal architecture of liver-cell cords (silver staining), as well as abnormal glycogen accumulation (PAS staining) were observed on day 90 in CN2-29^(+/-)/MxCre^(+/-) mice. We observed steatosis (oil-red-O staining) on day 180 and, subsequently, fibrosis (Azan staining) on day 480. The scale bars indicate 50 μ m. doi:10.1371/journal.pone.0051656.g001




Figure 2. Effects of rVV-HCV treatment on the CN2-29^(+/-)/MxCre^(+/-) mice. (A) HCV gene structure in the CN2-29^(+/-)/MxCre^(+/-) mice and recombinant vaccinia viruses (rVV-HCV). MxCre/CN2-29 cDNA contains the core, E1, E2, and NS2 regions. The rVV-CN2 cDNA contains the core, E1, E2, and NS2 regions. The rVV-N25 cDNA contains the NS2, NS3, NS4A, NS4B, NS5A, and NS5B regions. The rVV-CN5 cDNA contains the entire HCV region. (B) Four groups of CN2-29^(+/-)/MxCre^(+/-) mice were inoculated intradermally with rVV-CN2, rVV-N25, rVV-CN5, or LC16m8 90 days after the poly(I:C) injection. Blood, liver, and spleen tissue samples were collected 7 and 28 days after the inoculation. (C) Liver sections from the four groups of CN2-29^(+/-)/MxCre^(+/-) mice 7 days after the inoculation. The sections were stained with H&E, silver, oil-red-O, or PAS. The scale bars indicate 50 μ m. (D) Histological evaluation of piecemeal necrosis in the four groups of CN2-29^(+/-)/MxCre^(+/-) mice 7 days after inoculation. (E) Histological evaluation of steatosis in the four groups of CN2-29^(+/-)/MxCre^(+/-) mice 7 days after inoculation. Significant relationships are indicated by a P-value. doi:10.1371/journal.pone.0051656.g002




Figure 3. Effects of HCV core protein expression on the livers of CN2-29^(+/−)/MxCre^(+/−) mice inoculated with rVV-HCV. (A) Expression of the HCV core protein in the four treatment groups of CN2-29^(+/−)/MxCre^(+/−) mice 28 days after the inoculation. Significant relationships are indicated by a P-value. (B) H&E staining and immunohistochemical analysis for HCV core protein in the LC16m8-, rVV-CN2-, rVV-CN5-, or rVV-N25-treated CN2-29^(+/−)/MxCre^(+/−) mice 28 days after the inoculation. Liver sections were stained with the anti-core monoclonal antibody. The scale bars indicate 50 μm. (C) Effects of HCV core protein expression on serum ALT levels in the four treatment groups of CN2-29^(+/−)/MxCre^(+/−) mice 28 days after immunization. (D) Cre-mediated genomic DNA recombination in the four treatment groups 28 days after immunization. (E) Expression of HCV mRNA in the LC16m8- or rVV-N25-treated CN2-29^(+/−)/MxCre^(+/−) mice 28 days after immunization. In all cases, n=6 mice per group. doi:10.1371/journal.pone.0051656.g003

ly different in the rVV-N25-treated mice and control mice (Figure 3C); this finding indicated that rVV-N25 treatment did not cause liver injury and that the antiviral effect was independent of hepatocyte destruction.

We hypothesized that the reduction in the levels of HCV core protein in rVV-HCV-treated mice was not caused by cytolytic elimination of hepatocytes that expressed HCV proteins. To investigate this hypothesis, we conducted an RTD-PCR analysis of genomic DNA from liver samples of CN2-29^(+/−)/MxCre^(+/−) mice. The recombined transgene was similar in rVV-N25-treated and control mice 28 days after immunization (Figure 3D). We also measured the expression of HCV mRNA in LC16m8-treated CN2-29^(+/−)/MxCre^(+/−) mice with that in rVV-N25-treated CN2-29^(+/−)/MxCre^(+/−) mice 28 days after immunization; the HCV mRNA levels did not differ between rVV-N25-treated CN2-29^(+/−)/MxCre^(+/−) and control mice (Figure 3E). These results indicated that rVV-N25-induced suppression of HCV core protein expression could be controlled at a posttranscriptional level.

Role of CD4 and CD8 T cells in rVV-N25-treated Mice

Viral clearance is usually associated with CD4 and CD8 T-cell activity that is regulated by cytolytic or noncytolytic antiviral mechanism [14]. To determine whether CD4 or CD8 T-cell activity was required for the reduction in HCV core protein levels

in the livers of transgenic mice, we analyzed the core protein levels in CN2-29^(+/−)/MxCre^(+/−) mice immunized with rVV-N25 in the absence of CD4 or CD8 T cells (Figure 4A). As expected, the mice lacking CD4 or CD8 T cells failed to show a reduction in HCV core protein levels (Figure 4B).

However, in mice lacking either CD4 or CD8 T-cells, the pathological changes associated with chronic hepatitis were resolved following rVV-N25 immunization, and the steatosis score of rVV-N25-treated mice was significantly lower than that of control mice (Figures 4C–E). These results indicated that CD4 and CD8 T cells were not responsible for the rVV-N25-induced amelioration of histological findings and that other inflammatory cell types may play an as-yet-undefined role in the resolution of the pathological changes in these mice.

rVV-N25 Immunization Induced an NS2-specific Activated CD8 T cells Response

Because we found that HCV protein reduction in the liver required CD8 T cells, we tested whether HCV-specific CD8 T cells were present in splenocytes 28 days after immunization. To determine the functional reactivity of HCV-specific CD8⁺ T cells, we performed a CD107a mobilization assay and intracellular IFN-γ staining. CN2-29 transgenic mice expressed the HCV structural protein and the NS2 region. However, rVV-N25 comprised only




Figure 4. Role of CD4 and CD8 T cells in rVV-N25-treated mice. (A) Schematic diagram depicts depletion of CD4 and CD8 T cells via treatment with monoclonal antibodies. (B) Comparison of HCV core protein expression in control, CD4-depleted, and CD8-depleted mice 28 days after immunization with LC16m8 or rVV-N25. (C, D) Histological analysis of liver samples from CD4-depleted or CD8-depleted CN2-29^(+/-)/MxCre^(+/-) mice

28 days after immunization with LC16m8 or rVV-N25. The scale bars indicate 100 μm (C) and 50 μm (D). (E) Histological evaluation of steatosis in liver samples from CD4-depleted or CD8-depleted CN2-29^(+/-)/MxCre^(+/-) mice 28 days after immunization with LC16m8 or rVV-N25. Significant relationships are indicated by a P-value. doi:10.1371/journal.pone.0051656.g004

a HCV nonstructural protein. Thus, we focused on the role of the NS2 region as the target for CD8 T cells and generated EL-4 cell lines that expressed the NS2 antigen or the CN2 antigen.

Isolated splenocytes from immunized mice were co-cultured with EL-4CN2 or EL-4NS2 cell lines for 2 weeks and analyzed.

Cytolytic cell activation can be measured using CD107a, a marker of degranulation [15]. The ratio of CD8⁺CD107a⁺ cells to all CD8 T cells significantly increased in rVV-N25-treated splenocytes after co-culture with EL-4CN2 or EL-4NS2 ($P < 0.05$), whereas splenocytes that had been treated with any other rVV were not detected (Figure 5A, B and C). These results indicated that rVV-N25 treatment increased the frequency of HCV NS2-specific activated CD8 T cells. Consistent with these results, the ratio of CD8⁺IFN- γ ⁺ cells to all CD8 T cells for rVV-N25-treated mice was also significantly higher than that for mice treated with any other rVV ($P < 0.05$). Taken together, these findings indicated that rVV-N25 induced an effective CD8 T-cell immune response and that NS2 is an important epitope for CD8 T cells.

rVV-N25 Immunization Suppressed Inflammatory Cytokines Production

To determine whether rVV-N25 treatment affected inflammatory cytokine production, we measured serum levels of inflammatory cytokines after rVV immunization. The serum levels of these inflammatory cytokines increased in the CN2-29^(+/-)/MxCre^(+/-) mice (Figure 6A, Figure S5). Immunization with rVV-N25 affected serum levels of inflammatory cytokines in CN2-29^(+/-)/MxCre^(+/-) mice and caused a return to the cytokine levels observed in wild-type untreated mice (Figure 6A). In wild-type mice, the cytokine levels remained unchanged after immunization (Figure 6A). These results indicated that inflammatory cytokines were responsible for liver pathogenesis in the transgenic mice.

To test the hypothesis that inflammatory cytokines were responsible for liver pathogenesis in CN2-29^(+/-)/MxCre^(+/-) mice, we administered transgenic mouse serum intravenously into nontransgenic mice. We observed the development of chronic hepatitis in the nontransgenic mice within 7 days after the serum transfer (Figures 6B and C). This finding was consistent with the




Figure 5. Immunization with rVV-N25 induced CD8 T-cell degranulation, a marker for cytotoxicity, and IFN- γ production. (A) The numbers represent the percentage of CD107a positive cells and negative cells (left two columns) and IFN- γ -positive cells and negative cells (right two columns). (B, C) The ratio of CD8⁺IFN- γ ⁺ cells to all CD8 T cells for rVV-N25-treated mice was significantly higher than that for mice treated with any other rVV. Splenocytes (4×10^6 per well) were cultured with EL-4CN2 or EL-4NS2 cell lines in RPMI 1640 complete medium including 3% T-STIMTM with ConA for 2 weeks. Harvested cells were incubated for 4 h with EL-4, EL-4CN2, or EL-4NS2 in combination with PE-labeled anti-CD107a mAb and monensin in RPMI 1640 complete medium with 50 IU/mL IL-2, according to the manufacturer's instruction. After incubation, cell suspensions were washed with PBS, and the cells were further stained with APC-labeled anti-IFN- γ mAb and Pacific blue-labeled anti-CD8 mAb. Harvested cells were stained with anti-CD107a-PE, anti-IFN- γ -APC, or anti-CD8-Pacific blue. Results that are representative of three independent experiments are shown. Significant relationships are indicated by P-value. doi:10.1371/journal.pone.0051656.g005




Figure 6. Immunization with rVV-N25 suppresses serum inflammatory cytokine levels. (A) Daily cytokine levels in the serum of CN2-29^(+/-)/MxCre^(+/-) mice during the week following immunization with LC16m8, rVV-CN2, rVV-N25, or rVV-CN5. Values represent means \pm SD (n = 3) and reflect the concentrations relative to those measured on day 0. The broken lines indicate the baseline data from wild-type mice. In all cases, n = 6 mice per group. (B) Liver sections from CN2-29^(+/-)/MxCre^(+/-) and CN2-29^(+/-)/MxCre^(-/-) mice. (C) Histology activity index (HAI) scores of liver samples taken from CN2-29^(+/-)/MxCre^(+/-) or CN2-29^(+/-)/MxCre^(-/-) mice. (D) Liver sections from CN2-29^(+/-)/MxCre^(+/-) mice in which TNF- α was neutralized and the IL-6 receptor was blocked. The scale bars indicate 50 μ m. (E) HAI scores of liver samples taken from CN2-29^(+/-)/MxCre^(+/-) in which TNF- α was neutralized and the IL-6 receptor was blocked. Tg and non-Tg indicate CN2-29^(+/-)/MxCre^(+/-) and CN2-29^(+/-)/MxCre^(-/-), respectively. (F) Macrophages were the main producers of TNF- α and IL-6 in CN2-29^(+/-)/MxCre^(+/-) mice following poly(I:C) injection. (G) Immunization with rVV-N25 reduced the number of macrophages in liver samples from CN2-29^(+/-)/MxCre^(+/-) mice and suppressed TNF- α and IL-6 production from macrophages (Figure 6G). Significant relationships are indicated by a P-value. doi:10.1371/journal.pone.0051656.g006

hypothesis that inflammatory mediators played a key role in inducing hepatitis. Furthermore, to investigate whether TNF- α and IL-6 played particularly critical roles in the pathogenesis of chronic hepatitis in the transgenic mice, we neutralized TNF- α and blocked the IL-6 receptor in the livers of these mice. As expected, chronic hepatitis did not develop in these mice. (Figure 6D and E).

Next, to determine which cell population(s) produced TNF- α , IL-6, or both during continuous HCV expression in CN2-29^(+/-)/MxCre^(+/-) mice, we isolated intrahepatic lymphocytes (IHLs) and labeled the macrophages (the F4/80⁺ cells) with anti-TNF- α and anti-IL-6 antibodies using an intracellular cytokine detection method. Macrophages in CN2-29^(+/-)/MxCre^(-/-) mice produced small amounts of TNF- α and IL-6, while those in CN2-29^(+/-)/MxCre^(+/-) mice produced much larger amounts of these cytokines (Figure 6F).

Finally, we evaluated whether rVV-N25 treatment affected the number of macrophages, cytokine production by macrophages, or both; specifically, we isolated IHLs from CN2-29^(+/-)/MxCre^(+/-) mice 7 days after immunization with rVV-N25 or with LC16m8. The percentage of macrophages (CD11b⁺F4/80⁺) among IHLs and IL-6 production from these macrophages were significantly lower in rVV-N25-treated mice than in control mice (Figure 6G). Though the percentage of TNF- α -producing macrophages was not significantly different in rVV-N25-treated and control mice (P = 0.099), rVV-N25 treatment appeared to suppress these macrophages. These results demonstrated that rVV-N25 had a suppressive effect on activated macrophages, and they indicated that this suppression ameliorated the histological indicators of chronic hepatitis.

Discussion

Various HCV transgenic mouse models have been developed and used to examine immune response to HCV expression and the effects of pathogenic HCV protein on hepatocytes [4,16,17]. However, these transgenic mice develop tolerance to the HCV protein; therefore, examining immune response to HCV protein has been difficult.

To overcome the problem of immune tolerance in mouse models of HCV expression, we developed an HCV model in mice that relies on conditional expression of the core, E1, E2, and NS2 proteins and the Cre/loxP switching system [5,6]; we showed that the injection of an Ad-Cre vector enhanced the frequency of HCV-specific activated CD8 T cells in the liver of these mice and caused liver injury. However, the Ad-Cre adenovirus vector alone causes acute hepatitis in wild-type mice. Nevertheless, the transgenic model was useful for evaluating interactions between the host immune system and viral protein (serum ALT level over 2,000 IU/L) [5]; HCV core protein levels were reduced and expression of this protein was transient (about 2 weeks). Therefore, this Ad-Cre-dependent model cannot be used to effectively investigate immune responses to chronic HCV hepatitis.

Here, we used poly (I:C)-induced expression of Cre recombinase to generate HCV transgenic mice in order to study the effect of HCV protein and confirmed that these mice developed chronic active hepatitis—including steatosis, lipid deposition, and hepatocellular carcinoma. These pathological findings in the transgenic mice were very similar to those in humans with chronic hepatitis C; therefore, this mouse model of HCV may be useful for analyzing the immune response to chronic hepatitis. However, experimental results obtained with this mouse model may not directly translate to clinical findings from patients with HCV infection because the expression of HCV proteins was not liver specific in these mice. Furthermore, poly(I:C) injection can activate innate immune responses and, consequently, might induce temporary liver injury [18]. Additionally, poly(I:C) injection has an adjuvant effect; specifically, it stimulates TLR3 signaling [19].

To evaluate whether poly(I:C) injection caused hepatitis in CN2-29^(+/-)/MxCre^(-/-) mice, we examined serum ALT levels and liver histology following poly(I:C) injection. We found that, following poly(I:C) injection, serum ALT levels in CN2-29^(+/-)/MxCre^(-/-) mice increased, reached a peak one day after injection, declined from day 1 to day 6, and were not elevated thereafter; this time-course indicated that poly(I:C) injection alone did not induce continuous liver injury (figure S6). Based on these findings, we believe that the effects of poly(I:C) injection in these mice did not confound our analysis of chronic hepatitis.

Immunization with rVV-N25 suppressed HCV protein levels in the liver, and this suppression was associated with ameliorated pathological chronic hepatitis findings (see Figure 3). Importantly, rVV-N25 treatment did not cause liver injury based on the serum ALT levels; therefore, this treatment was unlikely to have cytopathic effects on infected hepatocytes. These findings provided strong evidence that rVV-N25 treatment effectively halted the progression of chronic hepatitis. Immunization with plasmid DNA or with recombinant vaccinia virus can effectively induce cellular and humoral immune responses and exert a protective effect against challenge with HCV infection [20,21]. However, findings from these previous studies revealed HCV immunization of both uninfected, naïve animals and immune-tolerant animals induced a HCV-specific immune response. In the model describe here; the animals were immune competent for HCV; therefore, our findings provided further important evidence that rVV-N25 was effective in the treatment of chronic hepatitis.

In addition, we demonstrated that rVV-N25 treatment in the absence of CD4 and CD8 T cells had no effect on HCV clearance. This important observation indicated that rVV-N25-induced HCV clearance was mediated by CD4 and CD8 T cells. Many studies have shown that spontaneous viral clearance during acute HCV infection is characterized by a vigorous, broadly reactive CD4 and CD8 T-cell response. [8,22] HCV clearance and hepatocellular cytotoxicity are both mediated by CD8 antigen-specific (cytotoxic T lymphocyte) CTLs [23]. Consistent with these observations, rVV-N25 treatment effectively induced the accumulation of NS2-specific CD8 T cells, which express high levels of

CD107a and IFN- γ in the spleen. Notably, even with rVV-N25 immunization, the frequency of activated CD8 T cells was very low, and a minimum of 2-weeks incubation was required to distinguish the difference between rVV treatments. Even if a small population of specific CD8+ T cells played a relevant role in the reduction of core protein, it is difficult to assert that the only NS2-specific CD8+ T cells were important to this reduction. However, based on the results presented in Figure 4B, we are able to conclude that at least CD8+ and/or CD4+ T cells were important to the reduction in HCV core protein. Therefore, to elucidate the mechanism of HCV protein clearance, further investigation of not only the other T cell epitopes but also other immunocompetent cells is required.

Interestingly, rVV-N25 treatment—but not the rVV-CN2 or rVV-CN5 treatment—efficiently induced a HCV-specific activated CD8 T cells response; this difference in efficacy could have one or more possible causes. The HCV structural proteins (core, E1, and E2 proteins) in the rVV-CN construct may cause the difference; Saito et al. reported that injection with plasmid constructs encoding the core protein induced a specific CTL response in BALB/c mice [24]. Reportedly, CTL activity against core or envelope protein is completely absent from transgenic mice immunized with a plasmid encoding the HCV structural proteins, but core-specific CTL activity is present in transgenic mice that were immunized with a plasmid encoding the HCV core [21]. In contrast, when recombinant vaccinia virus expressing different regions of the HCV polyprotein were injected into BALB/c mice, only the HCV core protein markedly suppressed vaccinia-specific CTL responses [25]. Thus, the HCV core protein may have an immunomodulatory function [26]. Based on these reports and our results, we hypothesize that the causes underlying the effectiveness of rVV-N25 treatment were as follows: 1) this rVV construct included the core and envelope proteins and 2) the core protein had an immune-suppressive effect on CTL induction. Therefore, we suggest that exclusion of the core and envelope antigen as immunogen is one important factor in HCV vaccine design.

Interestingly, immunization with rVV-N25 rapidly suppressed the inflammatory response; however, immunization with either of the other rVVs did not (see Figure 6A). This result indicated that rVV-N25 may modulate inflammation via innate immunity, as well as via acquired immunity. Reportedly, Toll-like receptor (TLR)-dependent recognition pathways play a role in the recognition of poxviruses [27]. TLR2 and TLR9 have also been implicated in the recognition of the vaccinia virus [28,29]. These findings indicate that TLR on dendritic cells may modulate the immunosuppressive effect of rVV-N25 in our model of HCV infection; however, further examination of this hypothesis is required. The finding that pathological symptoms in the HCV transgenic mice were completely blocked by intravenous injection of TNF- α and IL-6 neutralizing antibodies indicated that the progression of chronic hepatitis depended on inflammatory cytokines in serum, rather than the HCV protein levels in hepatocytes. Lymphocytes, macrophages, hepatocytes, and adipocytes each produce TNF- α and IL-6 [30,31], and HCV-infected patients have elevated levels of TNF- α and IL-6 [32,33]. Both cytokines also contribute to the maintenance of hepatosteatosis in mice fed a high-fat diet [34], and production of TNF- α and IL-6 is elevated in obese mice due to the low grade inflammatory response that is caused by lipid accumulation [35]. These findings indicate that both cytokines are responsible for HCV-triggered hepatosteatosis, and anti-cytokine neutralization is a potential treatment for chronic hepatitis if antiviral therapy is not successful.

The reduction of macrophages in number might be due to the induction of apoptosis by vaccinia virus *in vitro* infection as

previously reported [36]. To understand the mechanisms responsible for the reduction of the number of macrophage, we performed another experiment to confirm whether the macrophages were infected with vaccinia virus inoculation. However, based on PCR analyses; vaccinia virus DNA was not present in liver tissue that contained macrophages (Figure S7). Furthermore, apoptosis of macrophages was not detected in liver samples (Data not shown). Based on these results, it is unlikely that the reduction in the number of macrophages was due to apoptosis induced by vaccinia virus infection. Although rVV-N25 reduced the number of macrophage, precise mechanism is still unknown. Further examination to elucidate the mechanism is required.

In conclusion, our findings demonstrated that rVV-N25 is a promising candidate for an HCV vaccine therapy. Additionally, the findings of this study indicate that rVV-N25 immunization can be used for prevention of HCV infection and as an antiviral therapy against ongoing HCV infection.

Materials and Methods

Ethics Statement

All animal care and experimental procedures were performed according to the guidelines established by the Tokyo Metropolitan Institute of Medical Science Subcommittee on Laboratory Animal Care; these guidelines conform to the Fundamental Guidelines for Proper Conduct of Animal Experiment and Related Activities in Academic Research Institutions under the jurisdiction of the Ministry of Education, Culture, Sports, Science and Technology, Japan, 2006. All protocols were approved by the Committee on the Ethics of Animal Experiments of the Tokyo Metropolitan Institute of Medical Science (Permit Number: 11–078). All efforts were made to minimize the suffering of the animals.

Animals

R6CN2 HCV cDNA (nt 294–3435) [37] and full genomic HCV cDNA (nt 1–9611) [38,39] were cloned from a blood sample taken from a patient (#R6) with chronic active hepatitis (Text S1). The infectious titer of this blood sample has been previously reported [40]. R6CN2HCV and R6CN5HCV transgenic mice were bred with Mx1-Cre transgenic mice (purchased from Jackson Laboratory) to produce R6CN2HCV-MxCre and R6CN5HCV-MxCre transgenic mice, which were designated CN2-29^(+/-)/MxCre^(+/-) and RzCN5-15^(+/-)/MxCre^(+/-) mice, respectively. Cre expression in the livers of these mice was induced by intraperitoneal injection of polyinosinic acid–polycytidylic acid [poly(I:C)] (GE Healthcare UK Ltd., Buckinghamshire, England); 300 μ L of a poly(I:C) solution (1 mg/mL in phosphate-buffered saline [PBS]) was injected three times at 48-h intervals. All animal care and experimental procedures were performed according to the guidelines established by the Tokyo Metropolitan Institute of Medical Science Subcommittee on Laboratory Animal Care.

Histology and Immunohistochemical Staining

Tissue samples were fixed in 4% paraformaldehyde in PBS, embedded in paraffin, sectioned (4- μ m thickness), and stained with hematoxylin and eosin (H&E). Staining with periodic acid–Schiff stain, Azan stain, silver, or Oil-red-O was also performed to visualize glycogen degeneration, fibrillization, reticular fiber degeneration, or lipid degeneration, respectively.

For immunohistochemical staining, unfixed frozen liver sections were fixed in 4% paraformaldehyde for 10 min and then incubated with blocking buffer (1% bovine serum albumin in PBS) for 30 min at room temperature. Subsequently, the sections were incubated with biotinylated mouse anti-HCV core mono-

clonal antibody (5E3) for 2 h at room temperature. After being washed with PBS, the sections were incubated with streptavidin–Alexa Fluor 488 (Invitrogen). The nuclei were stained with 4',6-diamidino-2-phenylindole (DAPI). Fluorescence was observed using a confocal laser microscope (Laser scanning microscope 510, Carl Zeiss).

Generation of rVVs

The pBR322-based plasmid vector pBMSF7C contained the ATI/p7.5 hybrid promoter within the hemagglutinin gene region of the vaccinia virus, which was reconstructed from the pSFJ1-10 plasmid and pBM vector [41,42]. Separate full-length cDNAs encoding either the HCV structural protein, nonstructural protein, or all HCV proteins were cloned from HCV R6 strain (genotype 1b) RNA by RT-PCR. Each cDNA was inserted into a separate pBMSF7C vector downstream of the pBMSF7C ATI/p7.5 hybrid promoter; the final designation of each recombinant plasmid was pBMSF7C-CN2, pBMSF7C-N25, or pBMSF7C-CN5 (Figure 2). They were then transfected into primary rabbit kidney cells infected with LC16m8 (multiplicity of infection = 10). The virus–cell mixture was harvested 24 h after the initial transfection by scrapping; the mixture was then frozen at -80°C until use. The hemagglutinin-negative recombinant viruses were cloned as previously described [42] and named rVV-CN2, rVV-N25, or rVV-CN5. Insertion of the HCV protein genes into the LC16m8 genome was confirmed by direct PCR, and expression of each protein from the recombinant viruses was confirmed by western blot analysis. The titers of rVV-CN2, rVV-N25, and rVV-CN5 were determined using a standard plaque assay and RK13 cells.

Statistical Analysis

Data are shown as mean \pm SD. Data were analyzed using the nonparametric Mann–Whitney or Kruskal–Wallis tests or ANOVA as appropriate; GraphPad Prism 5 for Macintosh (GraphPad) was used for all analyses. P values <0.05 were considered statistically significant.

Supporting Information

Figure S1 HAI score of liver samples taken from CN2-29^(+/-)/MxCre^(+/-) mice.
(EPS)

Figure S2 Lipid degeneration in samples of liver taken from CN2-29^(+/-)/MxCre^(+/-) mice.

References

- Lauer GM, Walker BD (2001) Hepatitis C virus infection. *N Engl J Med* 345: 41–52.
- Alter MJ (1995) Epidemiology of hepatitis C in the West. *Semin Liver Dis* 15: 5–14.
- Kawamura T, Furusaka A, Koziel MJ, Chung RT, Wang TC, et al. (1997) Transgenic expression of hepatitis C virus structural proteins in the mouse. *Hepatology* 25: 1014–1021.
- Moriya K, Fujie H, Shintani Y, Yotsuyanagi H, Tsutsumi T, et al. (1998) The core protein of hepatitis C virus induces hepatocellular carcinoma in transgenic mice. *Nat Med* 4: 1065–1067.
- Wakita T, Katsume A, Kato J, Taya C, Yonekawa H, et al. (2000) Possible role of cytotoxic T cells in acute liver injury in hepatitis C virus cDNA transgenic mice mediated by Cre/loxP system. *J Med Virol* 62: 308–317.
- Wakita T, Taya C, Katsume A, Kato J, Yonekawa H, et al. (1998) Efficient conditional transgene expression in hepatitis C virus cDNA transgenic mice mediated by the Cre/loxP system. *J Biol Chem* 273: 9001–9006.
- Folgori A, Capone S, Ruggeri L, Meola A, Sporeno E, et al. (2006) A T-cell HCV vaccine eliciting effective immunity against heterologous virus challenge in chimpanzees. *Nat Med* 12: 190–197.
- Chisari FV, Ferrari C (1995) Hepatitis B virus immunopathology. *Springer Semin Immunopathol* 17: 261–281.
- Machida K, Tsukiyama-Kohara K, Seike E, Tone S, Shibasaki F, et al. (2001) Inhibition of cytochrome c release in Fas-mediated signaling pathway in transgenic mice induced to express hepatitis C viral proteins. *J Biol Chem* 276: 12140–12146.
- Kuhn R, Schwenk F, Aguet M, Rajewsky K (1995) Inducible gene targeting in mice. *Science* 269: 1427–1429.
- Li K, Chen Z, Kato N, Gale M Jr, Lemon SM (2005) Distinct poly(I:C) and virus-activated signaling pathways leading to interferon-beta production in hepatocytes. *J Biol Chem* 280: 16739–16747.
- Sugimoto M, Yamanouchi K (1994) Characteristics of an attenuated vaccinia virus strain, LC16m0, and its recombinant virus vaccines. *Vaccine* 12: 675–681.
- Youn JW, Hu YW, Tricoche N, Pfahler W, Shata MT, et al. (2008) Evidence for protection against chronic hepatitis C virus infection in chimpanzees by immunization with replicating recombinant vaccinia virus. *J Virol* 82: 10896–10905.
- Guidotti LG, Rochford R, Chung J, Shapiro M, Purcell R, et al. (1999) Viral clearance without destruction of infected cells during acute HBV infection. *Science* 284: 825–829.
- Burkett MW, Shafer-Weaver KA, Strobl S, Baseler M, Malyguine A (2005) A novel flow cytometric assay for evaluating cell-mediated cytotoxicity. *J Immunother* 28: 396–402.

(EPS)

Figure S3 HCV protein expression after infection of LC16m8, rVV-CN2, rVV-N25, or rVV-CN5 into HepG2 cells.

(EPS)

Figure S4 Effects of treatment with rVV-N25 in RzCN5-15^(+/-)/MxCre^(+/-) mice.

(EPS)

Figure S5 Daily cytokine profiles of the serum from CN2-29^(+/-)/MxCre^(+/-) mice during the week following inoculation with LC16m8, rVV-CN2, rVV-N25, or rVV-CN5.

(EPS)

Figure S6 The immune response following poly(I:C) injection in the acute phase.

(EPS)

Figure S7 Detection of vaccinia virus DNA in the skin, liver, and spleen after inoculation with attenuated vaccinia virus (Lister strain) or highly attenuated vaccinia virus (LC16m8 strain).

(EPS)

Table S1 Incidence of hepatocellular carcinoma in male and female transgenic mice at 360, 480, and 600 days after poly(I:C) injection.

(EPS)

Text S1 Supporting information including material and methods, and references.

(DOCX)

Acknowledgments

We thank Dr. Fukashi Murai for supporting this study. We also thank Dr. Keiji Tanaka for providing the MxCre mice, Dr. Shigeo Koyasu for providing the GK1.5 (anti-CD4) and 53–6.72 (anti-CD8) monoclonal antibodies, and Dr Takashi Tokuhisa for helpful discussions.

Author Contributions

Performed the experiments: SS KK TC Y. Tobita TO FY Y. Tokunaga. Analyzed the data: SS KK TC MK. Contributed reagents/materials/analysis tools: KT-K TW TT MM K. Mizuno YH TH K. Matsushima. Wrote the paper: SS KK MK. Study concept and design: MK.

16. Pasquinelli C, Shoenberger JM, Chung J, Chang KM, Guidotti LG, et al. (1997) Hepatitis C virus core and E2 protein expression in transgenic mice. *Hepatology* 25: 719–727.
17. Lerat H, Honda M, Beard MR, Loesch K, Sun J, et al. (2002) Steatosis and liver cancer in transgenic mice expressing the structural and nonstructural proteins of hepatitis C virus. *Gastroenterology* 122: 352–365.
18. Lang KS, Georgiev P, Recher M, Navarini AA, Bergthaler A, et al. (2006) Immunoprivileged status of the liver is controlled by Toll-like receptor 3 signaling. *The Journal of clinical investigation* 116: 2456–2463.
19. Jasani B, Navabi H, Adams M (2009) Ampligen: a potential toll-like 3 receptor adjuvant for immunotherapy of cancer. *Vaccine* 27: 3401–3404.
20. Elmowalid GA, Qjao M, Jeong SH, Borg BB, Baumert TF, et al. (2007) Immunization with hepatitis C virus-like particles results in control of hepatitis C virus infection in chimpanzees. *Proc Natl Acad Sci U S A* 104: 8427–8432.
21. Sato J, Murata K, Lechmann M, Manickan E, Zhang Z, et al. (2001) Genetic immunization of wild-type and hepatitis C virus transgenic mice reveals a hierarchy of cellular immune response and tolerance induction against hepatitis C virus structural proteins. *J Virol* 75: 12121–12127.
22. Crispe IN (2009) The liver as a lymphoid organ. *Annu Rev Immunol* 27: 147–163.
23. Chisari FV (2005) Unscrambling hepatitis C virus-host interactions. *Nature* 436: 930–932.
24. Saito T, Sherman GJ, Kurokohchi K, Guo ZP, Donets M, et al. (1997) Plasmid DNA-based immunization for hepatitis C virus structural proteins: immune responses in mice. *Gastroenterology* 112: 1321–1330.
25. Large MK, Kittlesen DJ, Hahn YS (1999) Suppression of host immune response by the core protein of hepatitis C virus: possible implications for hepatitis C virus persistence. *Journal of immunology* 162: 931–938.
26. Dustin LB, Rice CM (2007) Flying under the radar: the immunobiology of hepatitis C. *Annu Rev Immunol* 25: 71–99.
27. Bowie A, Kiss-Toth E, Symons JA, Smith GL, Dower SK, et al. (2000) A46R and A52R from vaccinia virus are antagonists of host IL-1 and toll-like receptor signaling. *Proc Natl Acad Sci U S A* 97: 10162–10167.
28. Zhu J, Martinez J, Huang X, Yang Y (2007) Innate immunity against vaccinia virus is mediated by TLR2 and requires TLR-independent production of IFN- β . *Blood* 109: 619–625.
29. Samuelsson C, Hausmann J, Lauterbach H, Schmidt M, Akira S, et al. (2008) Survival of lethal poxvirus infection in mice depends on TLR9, and therapeutic vaccination provides protection. *J Clin Invest* 118: 1776–1784.
30. Sheikh MY, Choi J, Qadri I, Friedman JE, Sanyal AJ (2008) Hepatitis C virus infection: molecular pathways to metabolic syndrome. *Hepatology* 47: 2127–2133.
31. Tilg H, Moschen AR, Kaser A, Pines A, Dotan I (2008) Gut, inflammation and osteoporosis: basic and clinical concepts. *Gut* 57: 684–694.
32. Malaguamera M, Di Fazio I, Laurino A, Ferlito L, Romano M, et al. (1997) Serum interleukin 6 concentrations in chronic hepatitis C patients before and after interferon-alpha treatment. *Int J Clin Pharmacol Ther* 35: 385–388.
33. Larrea E, Garcia N, Qjan C, Civeira MP, Prieto J (1996) Tumor necrosis factor alpha gene expression and the response to interferon in chronic hepatitis C. *Hepatology* 23: 210–217.
34. Park EJ, Lee JH, Yu GY, He G, Ali SR, et al. (2010) Dietary and genetic obesity promote liver inflammation and tumorigenesis by enhancing IL-6 and TNF expression. *Cell* 140: 197–208.
35. Gregor MF, Hotamisligil GS (2011) Inflammatory mechanisms in obesity. *Annu Rev Immunol* 29: 415–445.
36. Humlova Z, Vokurka M, Esteban M, Melkova Z (2002) Vaccinia virus induces apoptosis of infected macrophages. *The Journal of general virology* 83: 2821–2832.
37. Choo QL, Kuo G, Weiner AJ, Overby LR, Bradley DW, et al. (1989) Isolation of a cDNA clone derived from a blood-borne non-A, non-B viral hepatitis genome. *Science* 244: 359–362.
38. Tsukiyama-Kohara K, Tone S, Maruyama I, Inoue K, Katsume A, et al. (2004) Activation of the CKI-CDK-Rb-E2F pathway in full genome hepatitis C virus-expressing cells. *J Biol Chem* 279: 14531–14541.
39. Nishimura T, Kohara M, Izumi K, Kasama Y, Hirata Y, et al. (2009) Hepatitis C virus impairs p53 via persistent overexpression of 3 β -hydroxysterol Delta24-reductase. *J Biol Chem* 284: 36442–36452.
40. Shimizu YK, Purcell RH, Yoshikura H (1993) Correlation between the infectivity of hepatitis C virus in vivo and its infectivity in vitro. *Proc Natl Acad Sci U S A* 90: 6037–6041.
41. Yasui F, Kai C, Kitabatake M, Inoue S, Yoneda M, et al. (2008) Prior immunization with severe acute respiratory syndrome (SARS)-associated coronavirus (SARS-CoV) nucleocapsid protein causes severe pneumonia in mice infected with SARS-CoV. *J Immunol* 181: 6337–6348.
42. Kitabatake M, Inoue S, Yasui F, Yokochi S, Arai M, et al. (2007) SARS-CoV spike protein-expressing recombinant vaccinia virus efficiently induces neutralizing antibodies in rabbits pre-immunized with vaccinia virus. *Vaccine* 25: 630–637.



ELSEVIER

Ribosomal production and *in vitro* selection of natural product-like peptidomimetics: The FIT and RaPID systems

Christopher J Hipolito and Hiroaki Suga

Bioactive natural product peptides have diverse architectures such as non-standard sidechains and a macrocyclic backbone bearing modifications. *In vitro* translation of peptides bearing these features would provide the research community with a diverse collection of natural product peptide-like molecules with a potential for drug development. The ordinary *in vitro* translation system, however, is not amenable to the incorporation of non-proteinogenic amino acids or genetic encoding of macrocyclic backbones. To circumvent this problem, flexible tRNA-acylation ribozymes (flexizymes) were combined with a custom-made reconstituted translation system to produce the flexible *in vitro* translation (FIT) system. The FIT system was integrated with mRNA display to devise an *in vitro* selection technique, referred to as the random non-standard peptide integrated discovery (RaPID) system. It has recently yielded an *N*-methylated macrocyclic peptide having high affinity ($K_d = 0.60$ nM) for its target protein, E6AP.

Address

Department of Chemistry, Graduate School of Science, The University of Tokyo, Tokyo 113-0033, Japan

Corresponding author: Suga, Hiroaki (hsuga@chem.s.u-tokyo.ac.jp)

Current Opinion in Chemical Biology 2012, 16:196–203

This review comes from a themed issue on
Biocatalysis and Biotransformation
Edited by Jon S Thorson and Ben Shen

Available online 6th March 2012

1367-5931/\$ – see front matter

© 2012 Elsevier Ltd. All rights reserved.

DOI 10.1016/j.cbpa.2012.02.014

Introduction

Bioactive natural product peptides are biosynthesized by either ribosomal translation machinery or nonribosomal peptide synthetases (NRPSs) and derivatized by post-translational modification enzymes or tailoring NRPS enzymes, respectively [1]. Unlike ordinary peptides, these ‘non-standard’ peptides are chemically diversified with D-amino acids, unique sidechains, *N*-methylation, and macrocyclic backbones (Figure 1). In the case of NRPSs, this diversity is largely attributed to the multidomain NRPSs’ ability to activate and modify amino acids before condensation. Domains are grouped into modules, and each module is in general responsible for the activation, modification and condensation of a single amino acid. Nature arranges these modules into an assembly line so that an

exact peptide sequence can be produced in an NRPS-templated manner. The idea of producing custom non-standard peptides or their *de novo* libraries has attracted many researchers and driven efforts toward engineering NRPSs, but manipulating the modular NRPS components to create tailored non-standard peptides has thus far met limited success owing to their mechanistic and functional complexities. Presumably, directed evolution of NRPSs, similar to the recent work reported by Fischbach *et al.* [2] as well as Evans *et al.* [3], may lead to a solution of this herculean challenge in order to achieve the goal of constructing *de novo* libraries in the future.

Biosynthetic pathways involving post-translational modification enzymes are less complex than those involving NRPSs. In fact, genetic and mechanistic studies on such enzymes have increased our knowledge of essential recognition elements in substrates, enabling us to design non-natural substrates. Moreover, recent studies using synthetic peptides as well as coupling with *in vivo* incorporation of an artificial amino acid has allowed for the expansion of substrate repertoires [4–7]. Thus, continuing such efforts will lead us to the goal of synthesis of *de novo* libraries and, ultimately, discovering bioactive compounds with new chemical scaffolds against therapeutic targets of interest.

As a completely different technology, recent advances in artificial manipulation of the genetic code have shown the potential to produce natural product-like non-standard peptides by using translation machinery (*vide infra*). In this technology, unlike the above two biosynthetic processes, the sequences of such non-standard peptides are encoded in messenger RNA (mRNA) or its parental DNA. This feature facilitates the construction of their *de novo* libraries with high sequence complexities and, moreover, enables us to devise a one-pot screening (or selection) system to discover active non-standard peptides. In this review, we shall focus on discussing this emerging new technology.

Genetic code reprogramming as a distinct alternative approach

Genetic code reprogramming, which allows for a variety of non-standard amino acids to be incorporated by using sense codon suppressions, is a technique that was introduced within the last decade [8,9]. Genetic code reprogramming in general requires the use of a reconstituted *in vitro* translation system [10]. For instance, Forster *et al.* demonstrated in 2003 that reconstitution of only limited members of *Escherichia coli* translation factors (initiation