

Table 1. Cont.

OK-99-10	75	<i>Theonella</i> sp.	Porifera	EtOAc	Shimoji Island
<u>OK-99-12</u>	53	<i>Isis hippuris</i>	Cnidaria	EtOAc	Shimoji Island
OK-99-13	68	<i>Acanthella</i> sp.	Porifera	EtOAc	Shimoji Island
OK-99-15	64	<i>Phyllospongia</i> sp.	Porifera	EtOAc	Shimoji Island
<u>OK-99-17</u>	59	<i>Petrosia</i> sp.	Porifera	EtOAc	Shimoji Island
OK-99-18	80	<i>Fasciospongia rimosa</i>	Porifera	EtOAc	Shimoji Island
OK-99-20	77	<i>Echinoclathria</i> sp.	Porifera	EtOAc	Shimoji Island
OK-99-21	68	<i>Strongylophora</i> sp.	Porifera	EtOAc	Shimoji Island
OK-99-23	74	<i>Dysidea herbacea</i>	Porifera	EtOAc	Shimoji Island
<u>OK-99-26</u>	55	<i>Dysidea</i> cf. <i>arenaria</i>	Porifera	EtOAc	Shimoji Island
OK-99-28	123	<i>Plakortis</i> sp.	Porifera	EtOAc	Shimoji Island
OK-99-31	118	<i>Spongia</i> sp.	Porifera	EtOAc	Okinawa Island
OK-99-34	119	<i>Theonella swinhoei</i>	Porifera	EtOAc	Okinawa Island
OK-99-35	108	<i>Petrosia</i> sp.	Porifera	EtOAc	Okinawa Island
OK-99-36	90	<i>Acanthella</i> sp.	Porifera	EtOAc	Okinawa Island
OK-99-37	102	<i>Luffariella</i> sp.	Porifera	EtOAc	Okinawa Island
OK-99-41	62	<i>Dysidea</i> cf. <i>arenaria</i>	Porifera	EtOAc	Okinawa Island
OK-99-43	85	<i>Xestospongia</i> sp.	Porifera	EtOAc	Okinawa Island
OK-99-44	61	<i>Dysidea arenaria</i>	Porifera	EtOAc	Okinawa Island
OK-99-47	108	<i>Dysidea</i> cf. <i>arenaria</i>	Porifera	EtOAc	Okinawa Island
OK-99-49	90	<i>Petrosia</i> sp.	Porifera	EtOAc	Chibishi
OK-99-51	69	<i>Isis hippuris</i>	Cnidaria	EtOAc	Chibishi
OK-99-52	78	<i>Petrosia</i> sp.	Porifera	EtOAc	Kuro Island
OK-99-55	65	<i>Acanthella</i> sp.	Porifera	EtOAc	Kuro Island
OK-99-57	84	<i>Theonella swinhoei</i>	Porifera	EtOAc	Kuro Island
OK-99-63	117	<i>Epipolasis kushimotoensis</i>	Porifera	EtOAc	Kuro Island
OK-99-64	98	<i>Xestospongia</i> sp.	Porifera	EtOAc	Kuro Island
SG1-1-2	77	<i>Comanthus gisleni</i>	Echinodermata	MeOH	Kume Island
SG1-2-2	112	<i>Stephanometra indica</i>	Echinodermata	MeOH	Kume Island
<u>SG1-5-2</u>	33	<i>Comantella</i> sp. cf. <i>maculata</i>	Echinodermata	MeOH	Kume Island
SG1-9-2	57	<i>Phanogenia gracilis</i>	Echinodermata	MeOH	Kume Island
<u>SG1-12-2</u>	39	<i>Comanthus parvicirrus</i>	Echinodermata	MeOH	Kume Island
SG1-14-2	117	<i>Comaster schlegelii</i>	Echinodermata	MeOH	Kume Island
<u>SG1-15-2</u>	26	Colobometridae sp.	Echinodermata	MeOH	Kume Island
SG1-16-2	66	<i>Cenometra bella</i>	Echinodermata	MeOH	Kume Island
SG1-19-2	78	<i>Comaster nobilis</i>	Echinodermata	MeOH	Kume Island
<u>SG1-21-2</u>	32	<i>Oxycomanthus</i> sp.	Echinodermata	MeOH	Kume Island
<u>SG1-23-1</u>	-3	<i>Alloeocomatella polycladia</i>	Echinodermata	EtOAc	Kume Island

Table 1. Cont.

<u>SG1-24-1</u>	24	<i>Comanthus</i> sp.	Echinodermata	EtOAc	Kume Island
<u>SG1-26-2</u>	51	<i>Oxycomanthus benetti</i>	Echinodermata	MeOH	Kume Island
<u>SG1-28-2</u>	38	<i>Lamprometra palmata</i>	Echinodermata	MeOH	Kume Island
<u>SG1-30-1</u>	25	<i>Colobometra perspinosa</i>	Echinodermata	EtOAc	Kume Island
<u>SG1-31-1</u>	26	<i>Comanthus</i> sp.	Echinodermata	EtOAc	Kume Island
<u>SG1-33-1</u>	32	<i>Basilometra boschmai</i>	Echinodermata	EtOAc	Kume Island
SG3-1	82	<i>Stereonephthya</i> sp.	Cnidaria	EtOAc	Tokashiki Island
SG3-4	73	<i>Dysidea</i> cf. <i>arenaria</i>	Porifera	EtOAc	Tokashiki Island
SG3-6	74	<i>Stylotella</i> sp.	Porifera	EtOAc	Tokashiki Island
SG3-10	139	<i>Epipolasis</i> sp.	Porifera	EtOAc	Tokashiki Island
SG3-11	97	<i>Nephtea</i> sp.	Cnidaria	EtOAc	Tokashiki Island
SG3-21	106	<i>Myrmekioderma</i> sp.	Porifera	EtOAc	Tokashiki Island
SG3-25	111	<i>Pseudoceratina purpurea</i>	Porifera	EtOAc	Tokashiki Island
SG3-26	95	<i>Leucetta</i> sp.	Porifera	EtOAc	Tokashiki Island
SG3-28	65	<i>Lyngbya</i> sp.	Cyanobacteria	EtOAc	Tokashiki Island
SG3-29	61	<i>Dysidea</i> sp.	Porifera	EtOAc	Tokashiki Island

Total number of marine organisms: 61; Marine organisms that strongly inhibit NS3 helicase activity (<30%) (boldface and underlined): 5; Extracts of organisms that exhibit intermediate inhibition of NS3 helicase activity (30%–60%) (underlined): 11; EtOAc: Ethyl acetate; MeOH: Methanol.

Figure 1. Schematic representation of the PET assay system for unwinding activity of HCV NS3 helicase. The fluorescent dye (BODIPY FL) is attached to the cytosine at the 5'-end of the fluorescent strand and quenched by the guanine base at the 3'-end of the complementary strand via photoinduced electron transfer. When the helicase unwinds the double-strand RNA substrate, the fluorescence of the dye emits bright light upon the release of the dye from the guanine base. The capture strand, which is complementary to the complementary strand, prevents the reannealing of the unwound duplex.






Figure 2. *Alloeocomatella polycladia* belongs to a class of feather star (Echinodermata, Crinoidea). The ethyl acetate fraction prepared from the marine organism was designated SG1-23-1 in this study.



Figure 3. Effect of SG1-23-1 on the unwinding activity of NS3 helicase. **(A)** NS3 helicase activity was measured by PET assay. The reactions were carried out in the absence or presence of SG1-23-1. Helicase activity in the absence of SG1-23-1 was defined as 100% helicase activity. Each value represents the mean of three independent reactions. Error bars indicate standard deviation. The data represent three independent experiments. **(B)** The unwinding activity of NS3 helicase was measured by RNA unwinding assay using radioisotope-labeled RNA. The heat-denatured single-strand RNA (26-mer) and the partial duplex RNA substrate were applied to lanes 1 and 2, respectively. The duplex RNA was reacted with NS3 (300 nM) in the presence of SG1-23-1 (lanes 4 to 9, 16 to 500 µg/mL). The resulting samples were subjected to native polyacrylamide gel electrophoresis.

A**B**


2.2. Effect of SG1-23-1 on HCV NS3 ATPase and RNA Binding Activities

The unwinding ability of HCV helicase is dependent on ATP binding, ATP hydrolysis, and RNA binding [8,9]. We examined the effect of SG1-23-1 on the ATPase activity of NS3 helicase. The ratio of free phosphate ($^{32}\text{P-Pi}$) in ATP ($^{32}\text{P-ATP}$) was measured in the presence of SG1-23-1. The reaction

was carried out between 16 and 500 μg of SG1-23-1 per milliliter. ATPase activity was slightly increased at 16 μg SG1-23-1 per milliliter and slightly decreased at 500 μg SG1-23-1 per milliliter (Figure 4A). However, the helicase activity was decreased to less than 10% in the presence of 50 μg of SG1-23-1 per milliliter (Figure 3A,B). Next, we examined the effect of SG1-23-1 on the binding of NS3 helicase to single-strand RNA (ssRNA). A gel-mobility shift assay was employed to estimate the binding activity of NS3 to 21 mer of ssRNA. The binding of NS3 to ssRNA was inhibited with SG1-23-1 in a dose-dependent manner (Figure 4B). These results suggest that SG1-23-1 contains the compound that inhibits RNA binding to NS3 helicase.

Figure 4. Effect of SG1-23-1 on ATPase and RNA-binding activities of NS3 helicase.

(A) The reaction mixtures were incubated with $[\gamma\text{-}^{32}\text{P}]$ ATP as described in Materials and Methods. The reaction mixtures were subjected to thin-layer chromatography. The start positions and migrated positions of ATP and free phosphoric acid are indicated as “Origin”, “ $^{32}\text{P}\text{-ATP}$ ”, and “ $^{32}\text{P}\text{-Pi}$ ”, respectively, on the left side of this figure. The data represent three independent experiments. (B) Gel mobility shift assay for RNA-binding activity of NS3 helicase. The reaction was carried out at the indicated concentration of SG1-23-1. The reaction mixture was subjected to gel mobility shift assay. The data represent three independent experiments.



2.3. Effect of SG1-23-1 on HCV RNA Replication in HCV 1b Replicon Cells

We investigated the effect of SG1-23-1 on both viral replication and growth of the replicon cell lines. The cell lines possess viral subgenomic RNAs derived from three genotype 1b strains (strains N [33], Con1 [34], and O [35]) or a full genomic RNA derived from the O strain [35]. Each cell line was treated with various concentrations of SG1-23-1. The treated cells were harvested 72 h post-treatment. Treatment with SG1-23-1 suppressed HCV RNA replications of all cell lines in a dose-dependent manner irrespectively of full- and sub-genome replicons; it exhibited no effect below 25 $\mu\text{g}/\text{mL}$ and

little effect on cellular viability at the highest concentration, 50 $\mu\text{g/mL}$ (Figure 5C,D). Both HCV NS3 and NS5A were decreased at the protein level in a dose-dependent manner, corresponding to the viral replication, but beta-actin was not changed in the cell line harboring subgenome replicon RNA of the Con1 strain (Figure 5E).

The inhibitory effect of SG1-23-1 on HCV replication is summarized in Table 2. The inhibitory effects on the HCV replication of the subgenome replicon derived from Con1, O, and N strains were 22.9 ± 0.4 , 19.9 ± 1.8 , and 44.2 ± 1.5 $\mu\text{g/mL}$, respectively, as EC_{50} ; and 48.1 ± 1.5 , 48.5 ± 0.3 , and >50 $\mu\text{g/mL}$, respectively, as EC_{90} . Treatment with SG1-23-1 inhibited the replication of the subgenome replicon of the O strain (EC_{50} : 19.9 ± 1.8 $\mu\text{g/mL}$; EC_{90} : 48.5 ± 0.3 $\mu\text{g/mL}$) at a more potent level than the replication of the full genomic replicon of the O strain (EC_{50} : 39.5 ± 0.8 $\mu\text{g/mL}$; EC_{90} : >50 $\mu\text{g/mL}$). When luciferase of firefly or *Renilla* was expressed under the control of the EF promoter, neither showed a significant change in activity in the presence of SG1-23-1 (Figure 5F). The replicon RNA of HCV is composed of the 5'-UTR of HCV, indicator genes (luciferase and drug-resistant genes), encephalomyocarditis virus (EMCV) IRES, the viral genes encoding complete or nonstructural proteins, and the 3'-UTR of HCV in that order [33–35]. The replicon RNA replicated autonomously in several HCV replication-permissive cell lines derived from several hepatoma cell lines. Nonstructural proteins in replicon cells were polycistronically translated through EMCV IRES. The cap-dependent translated mRNA, including *Renilla* luciferase, EMCV IRES, and the firefly luciferase/neomycin-resistant gene in that order, was constructed to examine the effect of the extract on EMCV-IRES-dependent translation (Figure 5G). When the expression of the mRNA was transcribed by an EF promoter of the transfected plasmid in the presence of SG1-23-1, the ratio of firefly luciferase activity to *Renilla* luciferase activity was not changed, suggesting that treatment with SG1-23-1 exhibited no effect on EMCV-IRES-dependent translation (Figure 5H). Thus, the inhibitory effect of SG1-23-1 on the luciferase activity must correspond to the replication efficiency of the replicon RNA but not to the inhibition of luciferase activity or the inhibition of EMCV-IRES-dependent translation. The inhibitory effect of the extract on the viral replication is similar to that of the extract on the helicase activity with respect to the values of IC_{50} and EC_{50} (Figure 3A and Table 2). These results suggest that treatment with SG1-23-1 inhibits HCV replication in a manner similar to that of the inhibitory effect on NS3 helicase activity.

Figure 5. Effect of SG1-23-1 on viral replication in replicon cell lines. (A–D) Huh7 Lunet/Con1 LUN Sb #26 (A), Huh7 rep Feo (B), Huh7#94/ORN3-5B#24 (C), and OR6 (D) cell lines were incubated in medium containing various concentrations of SG1-23-1. Luciferase and cytotoxicity assays were carried out as described in Materials and Methods. Error bars indicate standard deviation. The data represent three independent experiments. (E) Protein extract was prepared from Huh7 Lunet/Con1 LUN Sb #26 cells treated for 72 h with an indicated concentration of SG1-23-1 and then was subjected to Western blotting using antibodies to NS3, NS5A, and beta-actin. (F) Huh7 cell line transfected with pEF Fluc IN vector or pEF Rluc IN was established in the presence of G418. Both cell lines were incubated without (control) and with 50 $\mu\text{g/mL}$ SG1-23-1. Firefly or *Renilla* luciferase activity was measured 72 h post-treatment. Luciferase activity was normalized with protein concentration. Error bars indicate standard deviation. The data represent three

independent experiments. (G) Schematic structure of the plasmid, pEF RLuc EMCV IRES Feo. The bicistronic gene is transcribed under the control of elongation factor 1 α (EF1 α) promoter. The upstream cistron encoding *Renilla* luciferase (RLuc) is translated by a cap-dependent mechanism. The downstream cistron encodes the fusion protein (Feo), which consists of the firefly luciferase (FLuc) and neomycin phosphotransferase (Neo^r), and is translated under the control of the EMCV IRES. (H) Huh7 cell line transfected with pEF RLuc EMCV IRES Feo was established in the presence of G418. The cells were incubated for 72 h without (control) and with 50 μ g/mL of SG1-23-1. Firefly or *Renilla* luciferase activity was measured by the method described in Materials and Methods and was normalized by the protein concentration. F/R: Relative ratio of Firefly luciferase activity to *Renilla* luciferase activity. F/R is presented as a percentage of the control condition. Error bars indicate standard deviation. The data represent three independent experiments.








Table 2. Anti-HCV activity of SG1-23-1 in different replicon cell lines of genotype 1b.


Replicon Cell Line	Virus Strain (Genotype 1b)	EC ₅₀ ^a (µg/mL)	EC ₉₀ ^b (µg/mL)	CC ₅₀ ^c (µg/mL)
<u>Subgenome</u>				
Huh7 Lunet/ Con1 LUN Sb #26	Con1	22.9 ± 0.4	48.1 ± 1.5	>50
Huh7 rep Feo	N	44.2 ± 1.5	>50	>50
Hu7#94/ORN3-5B#24	O	19.9 ± 1.8	48.8 ± 0.3	>50
<u>Full genome</u>				
OR6	O	39.5 ± 0.8	>50	>50

All data represent means ± standard deviation for three independent experiments; ^a Fifty percent effective concentration based on the inhibition of HCV replication; ^b Ninety percent effective concentration based on the inhibition of HCV replication; ^c Fifty percent cytotoxicity concentration based on the reduction of cell viability.

2.4. Effect of SG1-23-1 on the Interferon (IFN) Signaling Pathway

It has been reported that the HCV replication in cultured cells is potently inhibited by interferon (IFN) [36,37]. We examined whether or not treatment with SG1-23-1 induces interferon from replicon cells. The replicon cells were treated with various concentrations of interferon-alpha 2b or 50 µg of SG1-23-1 per milliliter. The treated cells were harvested at 72 h post-treatment. The interferon-inducible genes, MxA and 2',5'-OAS, were induced with IFN-alpha 2b but not with SG1-23-1 (Figure 6). These results suggest that the inhibitory effect of SG1-23-1 on the replication of the HCV replicon is independent of the IFN signaling pathway.

Figure 6. Effect of SG1-23-1 on interferon signaling pathway. Huh7 Lunet/Con1 LUN Sb #26 cells were treated without (lane 1) or with 1, 10, or 100 U/mL IFNα-2b (lanes 2–4), and 50 µg/mL SG1-23-1 (lane 5) for 48 h. The mRNAs of MxA, 2',5'-OAS, and glyceraldehyde-3-phosphate dehydrogenase (GAPDH) as an internal control were detected by reverse-transcription polymerase chain reaction (RT-PCR). Error bars indicate standard deviation. The data represent three independent experiments.



Treatment with SG1-23-1 suppressed the helicase activity of NS3 in a dose-dependent manner and exhibited an IC_{50} of 11.7 $\mu\text{g/mL}$. Interestingly, treatment with SG1-23-1 inhibited the RNA binding activity of the helicase but not the ATPase activity of NS3. Treatment with SG1-23-1 inhibited the luciferase activity corresponding to the HCV replication in the replicon cell lines, but not the enzymatic activity of luciferase or the translational activity of EMCV IRES, suggesting that treatment with SG1-23-1 decreases HCV replication. Figure 4 shows that the viral proteins NS3 and NS5A in replicon cells were decreased by treatment with SG1-23-1, supporting the notion that SG1-23-1 inhibits HCV replication but not the enzymatic activity of luciferase. The inhibition of cell growth would not contribute to the inhibition of HCV replication by SG1-23-1 (Figure 3 and Table 2). Treatment with SG1-23-1 did not induce the interferon-stimulated genes in the replicon cell lines (Figure 6), suggesting that inhibition of HCV replication by treatment with SG1-23-1 is not due to interferon induction or interferon signaling. The extract SG1-23-1 inhibited the HCV replicon with an EC_{50} of 22 to 44 $\mu\text{g/mL}$, which is similar to the value of IC_{50} . These results suggest that the anti-HCV compound(s) included in *A. polycladia* can suppress viral replication by inhibiting NS3 helicase activity.

3. Experimental Section

3.1. Preparation of Extracts from Marine Organisms

All marine organisms used in this study were collected by hand during scuba diving off Shimoji, Okinawa, Chibishi, Kuro, Kume, and Tokashiki Islands in Okinawa Prefecture, Japan. In the case of OK-99-tagged extract, a specimen was soaked in ethanol. The ethanol-soluble fraction was concentrated, and the resulting aqueous material was suspended in ethyl acetate (EtOAc). The organic fraction was used for screening.

Each specimen from Kume was soaked in ethanol. The ethanol-soluble fraction was concentrated. The resulting material was suspended in EtOAc. The EtOAc-soluble fraction was used for screening and tagged with SG1 and the last digit of “1”. The water layer was concentrated to dryness and suspended in methanol (MeOH). The MeOH-soluble fraction was used for screening and tagged with SG1 and the last digit of “2”.

Each specimen from Tokashiki was extracted three times with acetone. After removal of acetone from the solution, the residual material was suspended in EtOAc. The EtOAc-soluble fraction was used for screening and tagged with SG3.

All samples were dried and then solubilized in dimethyl sulfoxide (DMSO) before testing.

3.2. High-Throughput Screening of NS3 Helicase Inhibitors

A continuous fluorescence assay based on photoinduced electron transfer (PET) was described previously [38] and was slightly modified with regard to the reaction mixture. A schematic diagram of the PET assay for HCV NS3 helicase activity is shown in Figure 1. The double-strand RNA was prepared as a substrate by annealing, at a 1:2 molar ratio, a 5' BODIPY FL-labeled 37-mer (5'-CUAUUACCUCCACCCUCAUAACCUUUUUUUUUUUUUUUU-3') to a 23-mer (GGUUAUGAG GGUGGAGGUAAUAG). When unwound by HCV NS3 helicase, the unlabeled ssRNA was captured by a DNA capture strand (5'-CTATTACCTCCACCCTCATAACC-3'). A fluorescent-dye-labeled

oligonucleotide was purchased from J-Bio 21 Corporation. BODIPY FL was attached to the 5'-end via an aminohexylphosphate linker with a six-carbon spacer. Unlabeled oligonucleotides were purchased from Japan Bio Services Co., Ltd. The PET NS3 helicase assay was carried out in 22 μ L of 25 mM MOPS-NaOH (pH 6.5) containing 3 mM MgCl₂, 2 mM dithiothreitol (DTT), 4 U RNasin, 50 nM of the double-strand RNA described above, 100 nM DNA capture strand, 5 mM ATP, and the extract (25 μ g/mL) and 240 nM HCV NS3 helicase. The reaction was started by the addition of HCV NS3 helicase. The reaction mixture was incubated at 37 °C for 30 min. The fluorescence intensity was recorded every 5 s until 5 min post-reaction, and then every 30 s between 5 and 30 min post-reaction by using a LightCycler 1.5 (Roche Diagnostics, Tokyo, Japan). The initial reaction velocity was calculated and represented as NS3 helicase activity.

3.3. ATPase Assay

NS3 ATPase activity was determined by the method of Gallinari *et al.* [39] with slight modifications. The reaction was carried out at 37 °C for 10 min in 10 μ L of the reaction mixture containing 25 mM MOPS-NaOH (pH 7.0), 1 mM DTT, 5 mM MgCl₂, 5 mM CaCl₂, 1 mM [γ -³²P] ATP (Muromachi, Tokyo, Japan), 300 nM NS3, and 0.1 μ g poly (U) per microliter and an indicated concentration of SG1-23-1, and then was terminated by the addition of 15 microliters of 10 mM EDTA. Two microliters of the reaction mixture were spotted onto a polyethyleneimine cellulose sheet (Merck, Darmstadt, Germany) and then developed in 0.75 M LiCl/1 M formic acid solution at room temperature for 20 min. The sheet was air-dried completely and then exposed to an image plate. Radioactive bands were visualized with an Image Reader FLA-9000 and quantified by Multi Gauge V 3.11 software (version 3.11; Fujifilm: Tokyo, Japan, 2008).

3.4. RNA Helicase Assay

NS3 RNA helicase assay was carried out by the method of Gallinari *et al.* [39] with slight modifications. The substrate for annealing two complementary RNA oligonucleotides, 5'-AGAGAGAGAGGUUGAGAGAGAGAGAGUUUGAGAGAGAGAG-3' (40-mer, template strand) and 5'-CAAACUCUCUCUCUCAACAAAAAA-3' (26-mer, release strand) was purchased from Shanghai GenePharma Co., Ltd. The release strand was labeled at the 5'-end with [γ -³²P] ATP (Muromachi, Tokyo, Japan) using the T4 polynucleotide kinase (Toyobo, Osaka, Japan) at 37 °C for 60 min and then purified by phenol chloroform extraction. The template and the labeled release strands were annealed at a molar ratio of 3:1 (template: release), denatured at 80 °C for 5 min, and slowly renatured at 23 °C for 30 min in an annealing buffer consisting of 20 mM Tris-HCl (pH 8), 0.5 M NaCl, and 1 mM EDTA. The partial duplex RNA substrate was purified on a G-50 micro column (GE Healthcare, Uppsala, Sweden) and stored at -20 °C in H₂O containing 0.25 U of RNasin Plus (Promega, Madison, WI, USA) per microliter.

SG1-23-1 extract was added at various concentrations to a helicase reaction mixture consisting of 25 mM MOPS-NaOH (pH 7.0), 2.5 mM DTT, 2.5 U of RNasin Plus (Promega), 100 μ g of BSA per milliliter, and 3 mM MgCl₂. The mixture was supplemented with 300 nM NS3 protein and 5 fM ³²P-labeled partial duplex RNA substrate. It was then preincubated at 23 °C for 15 min. After adding ATP at a final concentration of 5 mM, the reaction mixture (20 μ L) was incubated at 37 °C for 30 min

and stopped by adding 5 μ L helicase termination buffer consisting of 0.1 M Tris-HCl (pH 7.5), 20 mM EDTA, 0.5% SDS, 0.1% Nonidet P-40, 0.1% bromophenol blue, 0.1% xylene cyanol, and 25% glycerol. The terminated reaction mixture was subjected to native TBE 10% polyacrylamide gel electrophoresis. The radioactive RNAs in the gel were visualized with an Image Reader FLA-9000 (Fujifilm) and quantified by Multi Gauge V 3.11 software.

3.5. RNA Binding Assay

RNA binding to NS3 helicase was analyzed by gel mobility shift assay [40]. First, let-7 single-strand RNA (5'-UGAGGUAGUAGGUUGUAUAGU-3') was incubated with [γ - 32 P] ATP (Muromachi, Tokyo, Japan) and T4 polynucleotide kinase (Toyobo) at 37 °C for 60 min for labeling at the 5'-end of the single-strand RNA. The reaction mixture was subjected to phenol chloroform extraction for purification of labeled RNA. The reaction was carried out at room temperature for 15 min in 20 μ L of the mixture consisting of 30 mM Tris-HCl (pH 7.5), 100 mM NaCl, 2 mM MgCl₂, 1 mM DTT, 1 unit of RNasin Plus (Promega) per microliter, 300 nM NS3, 5 fmol let-7-labeled ssRNA, and an indicated concentration of SG1-23-1. The reaction was stopped by adding an equal volume of dye solution consisting of 0.025% bromophenol blue, 10% glycerol, and 0.5 \times Tris/borate/EDTA (TBE). The resulting mixture was subjected to native 6% polyacrylamide gel electrophoresis (acrylamide: bis acrylamide = 19:1). The radioactive RNA was visualized with the Image Reader FLA-9000 and quantified by Multi Gauge V 3.11 software.

3.6. Cell Lines

The following Huh-7-derived cell lines used in this study were maintained in Dulbecco's modified Eagle's medium containing 10% fetal calf serum and 0.5 mg/mL G418: The Lunet/Con1 LUN Sb #26 cell line, which harbors the subgenomic replicon RNA of the Con1 strain (genotype 1b) [34]; the Huh7/ORN3-5B #24 cell line, which harbors the subgenomic replicon RNA of the O strain (genotype 1b) [35]; the Huh7 Rep Feo cell line, which harbors the subgenomic replicon RNA of the N strain (genotype 1b) [33]; and the OR6 cell line, which harbors the full genomic RNA of the O strain (genotype 1b) [35].

3.7. Determination of Luciferase Activity in HCV Replicon Cells

HCV replicon cells were seeded at 2×10^4 cells per well in a 48-well plate 24 h before treatment. The extract SG1-23-1 was added to the culture medium at various concentrations. The treated cells were harvested 72 h post-treatment and lysed in cell culture lysis reagent (Promega) or *Renilla* luciferase assay lysis buffer (Promega). Luciferase activity in the harvested cells was estimated with a luciferase assay system (Promega) or a *Renilla* luciferase assay system (Promega). The resulting luminescence was detected by the Luminescencer-JNR AB-2100 (ATTO, Tokyo, Japan) and corresponded to the expression level of the HCV replicon.

3.8. Determination of Cytotoxicity in HCV Replicon Cells

HCV replicon cells were seeded at a density of 1×10^4 cells per well in a 96-well plate and incubated at 37 °C for 24 h. The extract fraction of the sample code SG1-23-1 was added to the culture medium at various concentrations. These cells were treated with an indicated concentration of the extract fraction and then were harvested 72 h post-treatment. Cell viability was measured by dimethylthiazol carboxymethoxy-phenylsulfophenyl tetrazolium (MTS) assay using a CellTiter 96 aqueous one-solution cell proliferation assay kit (Promega).

3.9. Effects on Activities of Luciferase and Internal Ribosome Entry Site (IRES)

The plasmid pEF Fluc IN and pEF Rluc EMCV IRES Feo were described previously [41]. The firefly luciferase gene was replaced with the *Renilla* luciferase gene in the plasmid pEF Fluc IN. The resulting plasmid was designated as pEF RlucIN in this study. The Huh7 cells were transfected with the pEF Fluc IN, pEF Rluc IN, or pEF Rluc EMCV IRES Feo and then were established in a medium containing 0.25 mg/mL G418 as described previously [41]. These cell lines were seeded at 2×10^4 cells per well in a 48-well plate 24 h before treatment, treated with 50 µg/mL extract SG1-23-1, and then harvested at 72 h post-treatment. Activities of firefly and *Renilla* luciferases in pEF Rluc EMCV IRES Feo were measured with the dual luciferase reporter assay system (Promega). Total protein concentration was measured using the BCA Protein Assay Reagent Kit (Thermo Scientific, Rockford, IL, USA) to normalize luciferase activity.

3.10. Western Blotting

The cells were lysed in lysis buffer containing Cell Culture Lysis Reagent (Promega). The cell lysate was subjected to SDS-10% polyacrylamide gel (SDS-PAGE). The proteins in the gel were transferred onto a polyvinylidene fluoride (PVDF) membrane. The resulting membrane was incubated with the primary antibodies at 4 °C overnight and then was washed three times with PBS containing 0.02% Tween 20 (PBS-T). The resulting membrane was reacted with a horseradish peroxidase-labeled anti-IgG antibody at room temperature for 2 h and then was washed three times with PBS-T. The reacted proteins were visualized with ImmunoStar LD (Wako Pure Chemical, Osaka, Japan). The antibodies to NS3 (Abcam, Cambridge, UK), NS5A (ViroGen, Watertown, MA, USA) and beta-actin were purchased from New England Biolabs (Beverly, MA, USA) and were used as the primary antibodies in this study.

3.11. Reverse-Transcription Polymerase Chain Reaction (RT-PCR)

The previously described method of RT-PCR [41] was slightly modified, as described below. Total RNA was isolated from cultured cells with the RNAqueous-4PCR kit (Ambion, Austin, TX, USA) and then was reverse-transcribed with a Superscript III reverse transcriptase (Invitrogen, Carlsbad, CA, USA). The transcribed mRNA was amplified with PCR using AmpliTaq Gold DNA polymerase (Applied Biosystems, Foster City, CA, USA) and an appropriate primer pair. Primer sequences targeting the genes encoding 2',5'-oligoadenylate synthetase (2',5'-OAS), myxovirus resistance protein A (MxA), and glyceraldehyde-3-phosphate dehydrogenase (GAPDH) were described previously [41].

4. Conclusions

In conclusion, we showed that the ethyl acetate extract from *Alloeocomatella polycladia* significantly inhibits HCV replication by suppressing viral helicase activity. The purification of an inhibitory compound from the extract of *Alloeocomatella polycladia* will be required in order to improve the efficacy of chemical modification of the compound(s).

Acknowledgments

We thank R. Bartenschlager for kindly providing cell lines and plasmids, and H. Kasai and I. Katoh for their helpful comments and discussions. This work was supported in part by grants-in-aid from the Ministry of Health, Labor (H22-kanen-004, 006 and 009), and Welfare and from the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

References

1. Baldo, V.; Baldovin, T.; Trivello, R.; Floreani, A. Epidemiology of HCV infection. *Curr. Pharm. Des.* **2008**, *14*, 1646–1654.
2. Seeff, L.B. Natural history of chronic hepatitis C. *Hepatology* **2002**, *36*, S35–S46.
3. Moriishi, K.; Matsuura, Y. Host factors involved in the replication of hepatitis C virus. *Rev. Med. Virol.* **2007**, *17*, 343–354.
4. Tsukiyama-Kohara, K.; Iizuka, N.; Kohara, M.; Nomoto, A. Internal ribosome entry site within hepatitis C virus RNA. *J. Virol.* **1992**, *66*, 1476–1483.
5. Bartenschlager, R.; Ahlborn-Laake, L.; Mous, J.; Jacobsen, H. Nonstructural protein 3 of the hepatitis C virus encodes a serine-type proteinase required for cleavage at the NS3/4 and NS4/5 junctions. *J. Virol.* **1993**, *67*, 3835–3844.
6. Kim, D.W.; Gwack, Y.; Han, J.H.; Choe, J. C-terminal domain of the hepatitis C virus NS3 protein contains an RNA helicase activity. *Biochem. Biophys. Res. Commun.* **1995**, *215*, 160–166.
7. Failla, C.; Tomei, L.; de Francesco, R. Both NS3 and NS4A are required for proteolytic processing of hepatitis C virus nonstructural proteins. *J. Virol.* **1994**, *68*, 3753–3760.
8. Belon, C.A.; Frick, D.N. Helicase inhibitors as specifically targeted antiviral therapy for hepatitis C. *Future Virol.* **2009**, *4*, 277–293.
9. Frick, D.N. The hepatitis C virus NS3 protein: a model RNA helicase and potential drug target. *Curr. Issues Mol. Biol.* **2007**, *9*, 1–20.
10. Kwong, A.D.; Rao, B.G.; Jeang, K.T. Viral and cellular RNA helicases as antiviral targets. *Nat. Rev. Drug Discov.* **2005**, *4*, 845–853.
11. Manns, M.P.; Wedemeyer, H.; Cornberg, M. Treating viral hepatitis C: efficacy, side effects, and complications. *Gut* **2006**, *55*, 1350–1359.
12. McHutchison, J.G.; Everson, G.T.; Gordon, S.C.; Jacobson, I.M.; Sulkowski, M.; Kauffman, R.; McNair, L.; Alam, J.; Muir, A.J.; Team, P.S. Telaprevir with peginterferon and ribavirin for chronic HCV genotype 1 infection. *N. Engl. J. Med.* **2009**, *360*, 1827–1838.

13. Zeuzem, S.; Hultcrantz, R.; Bourliere, M.; Goeser, T.; Marcellin, P.; Sanchez-Tapias, J.; Sarrazin, C.; Harvey, J.; Brass, C.; Albrecht, J. Peginterferon alfa-2b plus ribavirin for treatment of chronic hepatitis C in previously untreated patients infected with HCV genotypes 2 or 3. *J. Hepatol.* **2004**, *40*, 993–999.
14. Asselah, T.; Marcellin, P. New direct-acting antivirals' combination for the treatment of chronic hepatitis C. *Liver Int.* **2011**, *31 Suppl 1*, 68–77.
15. Jazwinski, A.B.; Muir, A.J. Direct-acting antiviral medications for chronic hepatitis C virus infection. *Gastroenterol. Hepatol. (NY)* **2011**, *7*, 154–162.
16. Lange, C.M.; Sarrazin, C.; Zeuzem, S. Review article: specifically targeted anti-viral therapy for hepatitis C—A new era in therapy. *Aliment. Pharmacol. Ther.* **2010**, *32*, 14–28.
17. Hofmann, W.P.; Zeuzem, S. A new standard of care for the treatment of chronic HCV infection. *Nat. Rev. Gastroenterol. Hepatol.* **2011**, *8*, 257–264.
18. Kwong, A.D.; Kauffman, R.S.; Hurter, P.; Mueller, P. Discovery and development of telaprevir: An NS3-4A protease inhibitor for treating genotype 1 chronic hepatitis C virus. *Nat. Biotechnol.* **2011**, *29*, 993–1003.
19. Kieffer, T.L.; Kwong, A.D.; Picchio, G.R. Viral resistance to specifically targeted antiviral therapies for hepatitis C (STAT-Cs). *J. Antimicrob. Chemother.* **2010**, *65*, 202–212.
20. Thompson, A.J.; McHutchison, J.G. Antiviral resistance and specifically targeted therapy for HCV (STAT-C). *J. Viral. Hepat.* **2009**, *16*, 377–387.
21. Belon, C.A.; High, Y.D.; Lin, T.I.; Pauwels, F.; Frick, D.N. Mechanism and specificity of a symmetrical benzimidazolephenylcarboxamide helicase inhibitor. *Biochemistry* **2010**, *49*, 1822–1832.
22. Maga, G.; Gemma, S.; Fattorusso, C.; Locatelli, G.A.; Butini, S.; Persico, M.; Kukreja, G.; Romano, M.P.; Chiasserini, L.; Savini, L.; *et al.* Specific targeting of hepatitis C virus NS3 RNA helicase. Discovery of the potent and selective competitive nucleotide-mimicking inhibitor QU663. *Biochemistry* **2005**, *44*, 9637–9644.
23. Chin, Y.W.; Balunas, M.J.; Chai, H.B.; Kinghorn, A.D. Drug discovery from natural sources. *AAPS J.* **2006**, *8*, E239–E253.
24. Koehn, F.E.; Carter, G.T. The evolving role of natural products in drug discovery. *Nat. Rev. Drug Discov.* **2005**, *4*, 206–220.
25. Li, J.W.; Vederas, J.C. Drug discovery and natural products: End of an era or an endless frontier? *Science* **2009**, *325*, 161–165.
26. Ahmed-Belkacem, A.; Ahnou, N.; Barbotte, L.; Wychowski, C.; Pallier, C.; Brillet, R.; Pohl, R.T.; Pawlotsky, J.M. Silibinin and related compounds are direct inhibitors of hepatitis C virus RNA-dependent RNA polymerase. *Gastroenterology* **2010**, *138*, 1112–1122.
27. Wagoner, J.; Negash, A.; Kane, O.J.; Martinez, L.E.; Nahmias, Y.; Bourne, N.; Owen, D.M.; Grove, J.; Brimacombe, C.; McKeating, J.A.; *et al.* Multiple effects of silymarin on the hepatitis C virus lifecycle. *Hepatology* **2010**, *51*, 1912–1921.
28. Ciesek, S.; von Hahn, T.; Colpitts, C.C.; Schang, L.M.; Friesland, M.; Steinmann, J.; Manns, M.P.; Ott, M.; Wedemeyer, H.; Meuleman, P.; *et al.* The green tea polyphenol, epigallocatechin-3-gallate, inhibits hepatitis C virus entry. *Hepatology* **2011**, *54*, 1947–1955.

29. Takeshita, M.; Ishida, Y.; Akamatsu, E.; Ohmori, Y.; Sudoh, M.; Uto, H.; Tsubouchi, H.; Kataoka, H. Proanthocyanidin from blueberry leaves suppresses expression of subgenomic hepatitis C virus RNA. *J. Biol. Chem.* **2009**, *284*, 21165–21176.
30. Donia, M.; Hamann, M.T. Marine natural products and their potential applications as anti-infective agents. *Lancet Infect. Dis.* **2003**, *3*, 338–348.
31. Molinski, T.F.; Dalisay, D.S.; Lievens, S.L.; Saludes, J.P. Drug development from marine natural products. *Nat. Rev. Drug Discov.* **2009**, *8*, 69–85.
32. Mayer, A.M.; Glaser, K.B.; Cuevas, C.; Jacobs, R.S.; Kem, W.; Little, R.D.; McIntosh, J.M.; Newman, D.J.; Potts, B.C.; Shuster, D.E. The odyssey of marine pharmaceuticals: A current pipeline perspective. *Trends Pharmacol. Sci.* **2010**, *31*, 255–265.
33. Yokota, T.; Sakamoto, N.; Enomoto, N.; Tanabe, Y.; Miyagishi, M.; Maekawa, S.; Yi, L.; Kurosaki, M.; Taira, K.; Watanabe, M.; *et al.* Inhibition of intracellular hepatitis C virus replication by synthetic and vector-derived small interfering RNAs. *EMBO Rep.* **2003**, *4*, 602–608.
34. Frese, M.; Barth, K.; Kaul, A.; Lohmann, V.; Schwarzle, V.; Bartenschlager, R. Hepatitis C virus RNA replication is resistant to tumour necrosis factor- α . *J. Virol.* **2003**, *84*, 1253–1259.
35. Ikeda, M.; Abe, K.; Dansako, H.; Nakamura, T.; Naka, K.; Kato, N. Efficient replication of a full-length hepatitis C virus genome, strain O, in cell culture, and development of a luciferase reporter system. *Biochem. Biophys. Res. Commun.* **2005**, *329*, 1350–1359.
36. Blight, K.J.; Kolykhalov, A.A.; Rice, C.M. Efficient initiation of HCV RNA replication in cell culture. *Science* **2000**, *290*, 1972–1974.
37. Guo, J.T.; Bichko, V.V.; Seeger, C. Effect of alpha interferon on the hepatitis C virus replicon. *J. Virol.* **2001**, *75*, 8516–8523.
38. Tani, H.; Akimitsu, N.; Fujita, O.; Matsuda, Y.; Miyata, R.; Tsuneda, S.; Igarashi, M.; Sekiguchi, Y.; Noda, N. High-throughput screening assay of hepatitis C virus helicase inhibitors using fluorescence-quenching phenomenon. *Biochem. Biophys. Res. Commun.* **2009**, *379*, 1054–1059.
39. Gallinari, P.; Brennan, D.; Nardi, C.; Brunetti, M.; Tomei, L.; Steinkuhler, C.; De Francesco, R. Multiple enzymatic activities associated with recombinant NS3 protein of hepatitis C virus. *J. Virol.* **1998**, *72*, 6758–6769.
40. Huang, Y.; Liu, Z.R. The ATPase, RNA unwinding, and RNA binding activities of recombinant p68 RNA helicase. *J. Biol. Chem.* **2002**, *277*, 12810–12815.
41. Jin, H.; Yamashita, A.; Maekawa, S.; Yang, P.; He, L.; Takayanagi, S.; Wakita, T.; Sakamoto, N.; Enomoto, N.; Ito, M. Griseofulvin, an oral antifungal agent, suppresses hepatitis C virus replication *in vitro*. *Hepatol. Res.* **2008**, *38*, 909–918.

© 2012 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).

