

play an important role in prevention of both early and metastatic cancer, the role of NK cell activity in controlling breast cancer is still controversial and few studies have addressed whether enhancing this activity is of clinical benefit to breast cancer patients.

A variety of mechanisms are involved in controlling neoplastic cells by NK cells, one of which is the direct release of cytolytic granules that contain perforin, granzymes, and granzyme B by exocytosis to kill target cells (i.e., the granule exocytosis pathway) [3, 4]. Most mature human NK cells in peripheral blood (PB) constitutively express granzyme B and perforin, and have basal cytotoxicity against NK-sensitive targets. Cytokine exposure with interleukin (IL)-2 or IL-15 is known to increase the baseline granzyme B and perforin abundance and cytotoxic activity of NK cells, and also converts basal NK cytotoxicity to lymphokine-activated killing. Another mechanism is mediated by death-inducing ligands such as Fas ligand (FasL) and tumor necrosis factor (TNF)-related apoptosis-inducing ligand (TRAIL). Fas, a TNF family protein, is expressed on breast cancer cell membranes [5, 6], suggesting that activation of the Fas/FasL pathway induces apoptosis mediated by caspase activation. An additional mechanism is involved when HER2-overexpressing breast cancer cells are targeted because differential levels of HER2 expression in normal versus HER2-overexpressing tumor cells, together with the clear involvement of HER2 in tumor progression, make HER2 an ideal target for therapeutic approaches. NK cell-mediated antibody-dependent cell-mediated cytotoxicity (ADCC) is thought to contribute to the therapeutic effects of monoclonal antibodies (mAbs) specifically directed against the extracellular domain of HER2 (trastuzumab).

NK cells are abundant in the liver in contrast to their relatively small percentage in the peripheral lymphatics and other lymphatic organs in rodents [7–9] and humans [10]; however, the underlying reason for the anatomically biased distribution of NK cells has not been elucidated. In addition, liver NK cells have been shown to mediate higher cytotoxic activity against tumor cells than PB NK cells do in rodents [7–9, 11]. However, these functional differences between PB and liver NK cells have not been extensively investigated in humans because of the limited availability of appropriate samples. In this study, we extracted NK cells from allograft liver perfusates in clinical liver transplantation and examined the quantitative and qualitative cytotoxic functions of those liver NK cells targeting various breast cancer cell lines in comparison with PB NK cells. Through the experiments, we attempted to define whether PB NK cells can recognize and kill breast cancer cells, and whether liver resident NK cells can hinder metastasis of breast cancer to the liver, to assess the potential therapeutic use of NK cells, i.e., by either activation of endogenous NK

cells or adoptive transfer of in vitro-activated autologous NK cells. As the therapeutic efficacy of endogenous or exogenous NK cells likely depends on their migration and accumulation at tumor metastasis sites, we further analyzed the expression of receptors and ligands for chemokines secreted from breast cancer cells on PB and liver NK cells.

## Materials and methods

### Isolation of liver and PB lymphocytes

Liver mononuclear cells (LMNCs) were obtained from liver perfusates in clinical living donor liver transplantation as previously described [10]. LMNCs were isolated by gradient centrifugation with Separate-L (Muto Pure Chemicals Co., Ltd, Tokyo, Japan). PB mononuclear cells (PBMCs) were also isolated by gradient centrifugation with Separate-L from heparinized PB from healthy volunteers and liver transplant donors. LMNCs and PBMCs were suspended in DMEM (Gibco, Grand Island, NY) supplemented with 10% heat-inactivated fetal calf serum (FCS) (Sanko Chemical Co., Tokyo, Japan), 25 mmol/l HEPES buffer (Gibco), 50  $\mu$ mol/l 2-mercaptoethanol (Katayama Chemical Co., Osaka, Japan), 50 U/ml penicillin, and 50  $\mu$ g/ml streptomycin (Gibco) (10% DMEM). The ethics committee at Hiroshima University Hospital approved this study.

### Cell culture

LMNCs and PBMCs were cultured with human recombinant IL-2 (100 Japanese reference U/ml; Takeda, Tokyo, Japan) in DMEM at 37°C in a 5% CO<sub>2</sub> incubator. Cells were harvested for further analyses after 5 days in culture. Cell viability was assessed by the dye-exclusion test.

### Isolation of NK cells

LMNCs and PBMCs were separated into a CD3<sup>-</sup>CD56<sup>+</sup> NK cell fraction and a non-NK cell fraction (T cells, NKT cells, B cells, and monocytes/macrophages) by magnetic cell sorting (Miltenyi Biotec, Bergisch Gladbach, Germany), using the human NK cell isolation kit (Miltenyi Biotec) according to the manufacturer's instructions. The purity of isolated fractions was assessed by FCM, and only preparations with purities >90% were used for functional studies.

### Cell lines

The human breast cancer cell lines were obtained as follows: MDA-MB-231 and MDA-MB-468 were from ATCC

(Manassas, VA), MDA-MB-453 was from Riken Cell Bank (Saitama, Japan), and MCF-7 was kindly provided by Dr. A. Kikuchi (Osaka University, Suita, Japan). MDA-MB-231, MDA-MB-453, and MDA-MB468 cells were cultured in Leibovitz's L-15 medium (Invitrogen) supplemented with 10% FCS at 37°C in a 0% CO<sub>2</sub> incubator. MCF-7 cells were cultured in DMEM supplemented with 10% FCS at 37°C in a 5% CO<sub>2</sub> incubator. The phenotypic features of those cell lines were well documented [12, 13].

#### Flow cytometric analyses

Flow cytometric analyses were performed using a FACSCalibur<sup>®</sup> cytometer (BD Biosciences, San Jose, CA). The following mAbs were used for surface staining of the lymphocytes: anti-CD3-fluorescein isothiocyanate (FITC), HIT3a; anti-CD56-phycoerythrin (PE), B159; anti-NKG2D-allophycocyanin (APC), 1D11; anti-CD183-APC, CXCR3 (1C6/CXR3); and anti-FasL-biotin, NOK-1. The biotinylated mAb was visualized using APC-streptavidin (all mAbs from BD Biosciences, Franklin Lakes, NJ).

To analyze activating receptors or TRAIL expression on NK cells, LMNCs and PBMCs were stained with anti-CD3-FITC, anti-CD56-APC, anti-TRAIL-PE, RIK-2; anti-NKp30-PE, P30-15; anti-NKp44-PE, P44-8.1; and anti-NKp46-PE, 9E2/NKp46; mAbs (all mAbs from BD Biosciences). To analyze inhibitory receptors on NK cells, LMNCs and PBMCs were stained with anti-CD3-APC, SP34-2; anti-CD56-PE, anti-CD-158a-FITC, HP-3E4; anti-CD-158b-FITC, CH-L; or anti-CD94-FITC, HP-3D9; mAbs (all mAbs from BD Biosciences). To analyze the expression of CD94/NKG2A or CD94/NKG2C on NK cells, LMNCs and PBMCs were stained with anti-CD3-PerCP, SK7; (BD Biosciences), anti-CD56-APC, anti-CD94-FITC, and anti-NKG2A-PE, z199; (Beckman Coulter), or anti-NKG2C-PE, 134591; (R&D Systems, Minneapolis, MN) mAbs. To analyze TRAIL receptors on breast cancer cell lines, they were stained with biotin-conjugated anti-TRAIL-R1/DR4, DJR1; anti-TRAIL-R2/DR5, DJR2-4; anti-TRAIL-R3/decoy receptor (DcR) 1, DJR3; or anti-TRAIL-R4/DcR2, DJR4-1; mAbs (all mAbs from eBioscience). To analyze HER2 expression on breast cancer cell lines, breast cancer cells were stained with PE-conjugated anti-HER2 mAb, Neu 24.7; (Becton-Dickinson). All biotinylated mAbs were visualized with APC-streptavidin (BD Biosciences). Dead cells were excluded from analysis by light-scatter and propidium iodide staining.

#### Immunohistochemistry

Surgically resected breast specimens were obtained from breast cancer patients who had undergone curative tumor resection at Hiroshima University Hospital. Breast specimens that had been pathologically proven to be normal


(i.e., excluding mastopathic and mastitic tissues) were used as normal controls. Each tumor section (4- $\mu$ m thickness) was deparaffinized and subjected to antigen retrieval by incubation in 10 mM citrate buffer (sodium citrate, pH 6.0) at 99°C for 25 min. To block non-specific antibody binding, sections were incubated with blocking solution (PBS containing Sanglorpor I.V., 1 mg/ml; CSL Behring AG, Bern, Switzerland) for 20 min. Sections were then incubated overnight at 4°C in blocking solution in the presence of the first antibody (biotin-conjugated anti-TRAIL-R1/DR4, anti-TRAIL-R2/DR5, anti-TRAIL-R3/DcR1, and anti-TRAIL-R4/DcR2) (5  $\mu$ g/ml) (all mAbs from eBioscience) or biotin-conjugated mouse IgG1,  $\kappa$  isotype control (P3) at the same concentration (eBioscience). Sections were washed twice in PBS, and primary antibody binding sites were visualized using the Dako EnVision Kit (Dako, Copenhagen, Denmark) according to the manufacturer's instructions. Sections were faintly counterstained with Harris' hematoxylin and mounted with glycerol gelatin.

#### Cytotoxicity assay

The <sup>51</sup>Cr-labeled breast cancer cells were incubated in a total volume of 200  $\mu$ l with effector cells in L-15 medium in round-bottomed 96-well microtiter plates (BD Falcon) for 4 h. IL-2-stimulated NK cells were used as effectors at an effector-to-target (E:T) ratio of 1.25:1–10:1. To evaluate the ADCC of trastuzumab, the assay was performed in the presence of 0.1  $\mu$ g/ml trastuzumab (kindly provided by Genentech Inc., San Francisco, CA). Target cells were incubated either in culture medium alone to determine spontaneous <sup>51</sup>Cr release or in a mixture of the culture medium and 2% Nonidet P-40 (Nakalai Tesque Inc., Kyoto, Japan) to determine the maximum <sup>51</sup>Cr release for controls. The radioactivity of the cell-free supernatants was measured in a gamma counter. The percentage of specific <sup>51</sup>Cr release was calculated as % cytotoxicity = [(cpm of experimental release – cpm of spontaneous release)/(cpm of maximum release – cpm of spontaneous release)]  $\times$  100. In some experiments, the <sup>51</sup>Cr-labeled breast cancer cells were incubated with effector cells either for 4 or 18 h in the presence of 10  $\mu$ g/ml of anti-TRAIL (N2B2) mAb, 10  $\mu$ g/ml of anti-FasL (MFL3) mAb (both from BD Pharmingen), and/or 50 nmol/l of concanamycin A (CMA) (Wako Chemicals, Osaka, Japan), which inhibits perforin-mediated cytotoxicity [14].

#### Cytometric bead assay

Chemokine production in the cell culture supernatants was analyzed by the cytometric bead array (CBA) using Human MIG (CXCL9), Human IP-10 (CXCL10), and Human I-TAC (CXCL11) Flex Sets (BD Bioscience), according to



**Fig. 1** The proportions of CD56<sup>+</sup>CD3<sup>-</sup> NK and CD56<sup>+</sup>CD3<sup>+</sup> NKT cells in LMNCs extracted from liver perfusates were significantly higher than those in PBMCs. **a** Flow cytometric (FCM) analyses of freshly isolated LMNCs obtained from liver perfusates and PBMCs from the corresponding donors were analyzed after staining with mAbs against CD3 and CD56. Lymphocytes were gated by forward scatter and side scatter. FCM profiles are representative of six and five independent experiments (using PBMCs and LMNCs, respectively). *a* CD3<sup>-</sup>CD56<sup>bright</sup> NK cell subset, *b* CD3<sup>-</sup>CD56<sup>dim</sup> NK cell subset, *c* CD3<sup>+</sup>CD56<sup>+</sup> (NKT) cell subset. **b** Percentages of CD3<sup>-</sup>CD56<sup>bright</sup> NK, CD3<sup>-</sup>CD56<sup>dim</sup> NK, and CD3<sup>+</sup>CD56<sup>+</sup> NKT cells among mononuclear cells (mean  $\pm$  SEM, PBMCs; *n* = 6, LMNCs; *n* = 5). **c** Histograms showing the log fluorescence intensities obtained for


TRAIL staining after gating CD3<sup>-</sup>CD56<sup>bright</sup> NK and CD3<sup>-</sup>CD56<sup>dim</sup> NK cell subsets of LMNCs and PBMCs from the corresponding donors. Data are shown as overlays with comparison against isotype controls. Histogram profiles are representative of independent experiments (PBMCs; *n* = 6, LMNCs; *n* = 5). **d** Mean fluorescence intensities (MFI) of TRAIL staining on NK cells freshly isolated from LMNCs and PBMCs (CD56<sup>bright</sup> NK cells; *open column*, CD56<sup>dim</sup> NK cells; *closed column*). Data represent mean  $\pm$  SEM (PBMCs; *n* = 6 LMNCs; *n* = 5). Statistical analyses were performed using the paired Student's *t* test (*\*P* < 0.05). *NK* natural killer, *TRAIL* TNF-related apoptosis-inducing ligand, *LMNC* liver mononuclear cells, *PBMC* peripheral blood mononuclear cells, *mAb* monoclonal antibody

the manufacturer's instructions, for the production of CXCL9, CXCL10, and CXCL11, respectively.

#### Cell migration assay

Migration assays were performed in transwell culture inserts (BD Falcon) of 6.4-mm diameter and 3- $\mu$ m pore

filters. MDA-MB231 and MDA-MB468 cells ( $1 \times 10^6$ /well) were cultured in the lower chamber of a 24-well plate (BD Falcon) in 0.5 ml L-15 medium. After 2 days, IL-2-activated NK cells derived from PBMCs and LMNCs in 0.2 ml L-15 medium were added to the upper chamber ( $1 \times 10^6$ /well), and cells were allowed to migrate for 2 h.



**Fig. 2** Phenotypic analyses of human PB and liver NK cells. **a** Flow cytometric (FCM) analyses of PBMCs and LMNCs cultivated with IL-2 for 5 days were performed after staining with mAbs against CD3 and CD56. FCM profiles are representative of five and six independent experiments (using PBMCs and LMNCs, respectively). Percentages of NK and NKT cells are indicated (mean  $\pm$  SEM, PBMC;  $n = 5$ , LMNC;  $n = 6$ ). **b** Expression of cytotoxic effector molecules on NK cell subsets among LMNCs or PBMCs freshly isolated or cultivated with or without IL-2 was analyzed. Expression of TRAIL

and FasL on electronically gated CD3<sup>-</sup>CD56<sup>+</sup> NK cells was analyzed by FCM. Numbers above the lesion marker line indicate the percentages of cells expressing TRAIL and FasL, and numbers below the line indicate the median fluorescence intensity of expression of whole NK cells (mean  $\pm$  SEM,  $n = 4$  each). PBMCs were obtained from the corresponding LMNC donor. Histogram profiles are representative of independent experiments. Dotted lines represent negative control staining with isotype-matched mAbs. \* $P < 0.05$  PB NK cells versus liver NK cells

### Enzyme-linked immunosorbent assay

The supernatants from the cell migration assay and coculture of MDA-MB231 or MDA-MB468 with IFN- $\gamma$  for 2 or 6 h were used. IFN- $\gamma$  and CXCL10 levels in the cell culture supernatants were determined by an enzyme-linked immunosorbent assay (ELISA) with the Quantikine kit (R&D Systems), according to the manufacturer instructions. Absorbance was measured at 492 nm on a microplate reader (MTP-300; CORONA Electric, Ibaraki, Japan).

### Statistical analysis

Data are presented as mean  $\pm$  SEM. The statistical differences of the results were analyzed by the 2-tailed, paired *t* test and Mann–Whitney *U* test, using Excel. *P* values of  $<0.05$  were considered statistically significant.

## Results

### Phenotypic properties of human NK cells


NK cells are abundant in the liver in contrast to their relatively small percentage in the peripheral lymphatics and other lymphatic organs in humans [10]. While NK cells in circulating lymphocytes have been phenotypically and functionally defined, those that reside in the liver remain to be characterized. We phenotypically analyzed the LMNCs that were extracted from the perfusates of allograft livers during liver transplantation surgery. The proportions of CD56<sup>+</sup>CD3<sup>-</sup> NK and CD56<sup>+</sup>CD3<sup>+</sup> NKT cells in the LMNCs extracted from liver perfusates were significantly higher than those in the PBMCs (Fig. 1a). Although this non-destructive method might allow some extent of contamination with circulating mononuclear cells, these data were consistent with previous reports using the enzymatic dissociation method [15]. Among CD56<sup>+</sup>CD3<sup>-</sup> NK cells, CD56<sup>bright</sup> cells, which constitutively expressed TRAIL, were abundant in LMNCs but were almost undetectable in PBMCs (Fig. 1b–d). On effector molecule analyses, the expression of TRAIL was significantly upregulated in both liver and PB NK cells after cultivation with IL-2 for 5 days. Both the proportion of the TRAIL<sup>+</sup> fraction and the staining intensity of liver NK cells were significantly higher than those of PB NK cells (Fig. 2a). Neither PB nor liver NK cells expressed FasL even after IL-2 stimulation (Fig. 2b). We further analyzed the C-type lectin-like receptors CD94, NKG2A, and NKG2C and killer cell immunoglobulin-like receptors (KIR) such as CD158a and CD158b (Fig. 3). CD94 recognizes the non-classical MHC class Ib molecule HLA-E, whereas KIRs are MHC class I-restricted molecules that recognize HLA-A, HLA-B, HLA-C, and HLA-G


**Fig. 3** IL-2 stimulation maintained the expression of C-type lectin-like receptors and killer cell immunoglobulin-like receptors (KIR) in both human PB and liver NK cells. Expression of the C-type lectin-like receptors CD94, NKG2A and NKG2C, and KIR such as CD158a and CD158b on NK cell subsets among LMNCs or PBMCs freshly isolated or cultivated with or without IL-2 was analyzed. Expression on electronically gated CD3<sup>-</sup>CD56<sup>+</sup> NK cells was analyzed by FCM. Numbers above the lesion marker line indicate the percentages of cells expressing each molecule, and numbers below the line indicate the median fluorescence intensity of expression of whole NK cells (mean  $\pm$  SEM, *n* = 4 each). PBMCs were obtained from the corresponding LMNC donor. Histogram profiles are representative of independent experiments. Dotted lines represent negative control staining with isotype-matched mAbs. \**P* < 0.05 PB NK cells versus liver NK cells

molecules. All freshly isolated PB and liver NK cells expressed CD94, and cell subpopulations expressed CD158a/CD158b (Fig. 3). No statistically significant differences were observed in the expression of CD94, NKG2A, NKG2C, CD158a, and CD158b between PB and liver NK cells. IL-2 stimulation maintained the expression of these molecules in both liver and PB NK cells, indicating that these cells retain their ability to protect self-MHC class I-expressing cells from NK cell-mediated death. On the analyses of the cytotoxicity-associated receptors, including NKp30, NKp44, NKp46, and NKG2D, no statistically significant differences were found between PB and liver NK cells even after IL-2 stimulation (Fig. 4). Although liver NK cells tended to express higher levels of NKp44 and NKp46 than did PB NK cells, the differences did not reach statistical significance.

### Breast cancer cells express the death-inducing receptor

Susceptibility to TRAIL-induced apoptosis may be related to the expression levels of multiple receptors on target cells. TRAIL binds to at least four receptors: two of these are death-inducing receptors (TRAIL-R1/DR4 and TRAIL-R2/DR5) containing cytoplasmic death domains and signal apoptosis, whereas the other two are death-inhibitory receptors (TRAIL-R3/DcR1 and TRAIL-R4/DcR2) that lack a functional death domain and do not mediate apoptosis; all have similar affinities for TRAIL and the latter two may act as decoys [16, 17]. The susceptibility to TRAIL-induced apoptosis is related to the expression levels of those receptors in tumor cells. We investigated the expression patterns of TRAIL-DR and TRAIL-DcR in both normal mammary gland and breast cancer tissue samples. Ductal cells in normal mammary gland tissues expressed TRAIL-DR4 together with TRAIL-DcR1 (Fig. 5a). Breast cancer cells showed a much higher expression of TRAIL-DR4 than did normal mammary gland cells, but little TRAIL-DcR1, regardless of the HER2 type. Similar to the clinical breast cancer tissues, all the tested breast cancer cell lines expressed high TRAIL-DR4 together with TRAIL-DR5, but no TRAIL-DcR1 and TRAIL-DcR2, regardless of their HER2 status (Fig. 5b).





**Fig. 4** IL-2 stimulation led to elevated expression of NKp30, NKp44, and NKG2D in both human PB and liver NK cells. Expression of the cytotoxicity-associated receptors, including NKp30, NKp44, NKp46, and NKG2D, in NK cell subsets among LMNCs or PBMCs freshly isolated or cultivated with or without IL-2 was analyzed. Expression in electronically gated CD3<sup>+</sup>CD56<sup>+</sup> NK cells was analyzed by FCM. Numbers above the lesion marker line indicate the percentages of

cells expressing each molecule, and numbers below the line indicate the median fluorescence intensity of expression of whole NK cells (mean ± SEM, n = 4 each). PBMCs were obtained from the corresponding LMNC donor. Histogram profiles are representative of independent experiments. Dotted lines represent negative control staining with isotype-matched mAbs.

### IL-2-stimulated NK cells showed significant cytotoxicity against breast cancer cells

Cytotoxicity assays of NK cells isolated from LMNCs and PBMCs as effectors and various breast cancer cell lines as targets were performed. Cells were stimulated by 5-day culture with IL-2 before use in the cytotoxicity assays. Liver NK cells showed more vigorous cytotoxicity against all tested cell lines (MDA-MB231, MDA-MB453, MDA-MB468, and MCF-7) compared with PB NK cells (Fig. 6). Addition of trastuzumab enhanced the cytotoxicity of both liver and PB NK cells toward MDA-MB231, MDA-MB453, and MCF-7, which express HER2. Although MDA-MB468 has been reported to be a triple-negative breast cancer cell line [12], it seemed to express dim HER2 on phenotypic analysis of the breast cancer cell lines in this study (Fig. 5b). This may explain why trastuzumab did not promote the cytotoxicity of PB NK cells but somewhat enhanced the cytotoxicity of liver NK cells toward MDA-MB468. Nevertheless, these observations suggest the involvement of HER2/trastuzumab-mediated ADCC. Despite the strong cytotoxicity exhibited by IL-2-stimulated donor liver NK cells, their cytotoxicities toward one-haplotype identical allogeneic and autologous lymphoblasts were negligible (data not shown).

### TRAIL and perforin are involved in the cytotoxicity of NK cells against breast cancer cells

To determine the contribution of TRAIL to the cytotoxicity of NK cells against breast cancer cells, the effect of a neutralizing anti-TRAIL mAb was examined in a cytotoxicity assay with PB and liver NK cells as effectors and MDA-MB231 cells as the target. Both PB and liver NK cell-induced cytotoxicity was inhibited partially by the anti-TRAIL mAb alone and more profoundly by the combination of the anti-TRAIL mAb and CMA, indicating that TRAIL and perforin are involved in NK cell-mediated cytotoxicity (Fig. 7). Remarkable levels of inhibition of NK cell-induced cytotoxicity were observed with anti-TRAIL mAb at 18-h culture when compared with those at 4 h. This finding is consistent with the results of the previous study with a mouse model demonstrating that death receptor-mediated NK cell kill needs longer incubation times than perforin-mediated NK cell kill [18].

NK cells were preferentially drawn by chemokines secreted from breast cancer cells, presumably through the CXCL10/CXCR3 axis



The distribution of NK cells is known to be associated with their expression of receptors and ligands for chemokines secreted from infectious or neoplastic sites [19–21]. We

found that NK cells freshly isolated from PBMCs and LMNCs highly expressed CXCR3, which binds to the chemokines CXCL9, CXCL10, and CXCL11 secreted by breast cancer cells. IL-2 activation increased the levels of CXCR3 expression on both NK cell types (Fig. 8a, b). We further investigated the secretion activities of these various chemokines from the breast cancer cell lines. Significant levels of CXCL10 were detected in the culture supernatants of three of four breast cancer cell lines: MDA-MB231, MDA-MB453, and MDA-MB468 (Fig. 9a). PB or liver NK cells activated with IL-2 were cultured in the upper compartment of transwell tissue culture plates in the presence or in the absence of MDA-MB231 or MDA-MB468 cells in the lower compartment for 2 h. The migration of PB and liver NK cells through the membrane was markedly promoted by the presence of tumor cells in the lower compartment, suggesting that NK cells are preferentially drawn by chemokines secreted from tumor cells (Fig. 9b). Significant levels of CXCL10 were detected only in the culture supernatants in the lower compartment with MDA-MB231 (Fig. 9c), suggesting that the CXCL10/CXCR3 axis plays an important role in the accumulation of NK cells in tumor sites. The MDA-MB231 cell line produced a lot more CXCL10 when PB NK cells were added (Fig. 9a, c), suggesting that soluble factors secreted from NK cells promoted the production of CXCL10 from this cell line. Taken together with the fact that CXCL10 is an IFN- $\gamma$ -inducible protein [22] and that IL-2 augments the active production of IFN- $\gamma$  from NK cells, we could assume that IFN- $\gamma$  secreted from NK cells promotes CXCL10 production from the breast cancer cell line, which in turn accelerates the migration of CXCR3-expressing NK cells into the tumor site. Consistent with this hypothesis, the CXCL10 levels were well correlated with IFN- $\gamma$  levels in the culture supernatants of the cell migration assay (Fig. 9d). In addition, we directly confirmed that IFN- $\gamma$  promoted the production of CXCL10 from MDA-MB231 and MDA-MB468 in a dose-dependent manner (Fig. 9e, f).

### Discussion

Human NK cells can be divided into the CD56<sup>bright</sup> and CD56<sup>dim</sup> subsets. These subsets have different phenotypic expression and may have different functions, although the direct functional significance of the expression levels of the CD56 antigen remains unknown. We previously demonstrated that CD56<sup>bright</sup> NK cells, which constitutively express low levels of TRAIL, are abundant in the liver [10]. CD56<sup>bright</sup> NK cells also constitutively express the high-affinity heterotrimeric IL-2R (IL-2R $\alpha\beta\gamma$ ) [23, 24]; hence, this subset has a high proliferative response to IL-2 and expand and survive through the upregulation of bcl-2 in






**Fig. 5** Breast cancer cells express death-inducing TRAIL-DR4 but lack death-inhibitory TRAIL-DcR1 and TRAIL-DcR2. **a** Immunohistochemical expression of TRAIL-DcR1, TRAIL-DcR2, TRAIL-DR4, and TRAIL-DR5 in normal breast tissue and tumor sites of HER2 (+)- and HER2 (-)-type breast cancer tissues. Immunopathological findings are representative of three individual samples in each breast cancer category. Magnification  $\times 200$ . Scale bar 100  $\mu\text{m}$ . **b** Surface expression of HER2 and TRAIL receptors on the surface of MDA-MB231, MDA-MB453, MDA-MB468, and MCF-7 was analyzed by FCM. Dotted lines represent negative control staining with isotype-matched mAbs. Numbers indicate the mean fluorescence intensity (MFI) of cells that stained positively for HER2 and TRAIL receptors. TRAIL, TNF-related apoptosis-inducing ligand; FCM flow cytometric, mAb monoclonal antibody, TNF tumor necrosis factor


2 in vitro [23, 26]. In this study, CD56<sup>bright</sup> NK cells exclusively survived and significantly upregulated TRAIL expression after in vitro cultivation of both PBMCs and LMNCs with IL-2 (Fig. 2). IL-2 stimulation also increased the surface expression of inhibitory receptors such as the KIR, including CD158a/158b and C-type lectin-like receptors (the CD94/NKG2 complex). CD94, which is expressed on essentially all NK cells, uses HLA-E expression as a sensor for the overall HLA class I level of a cell. In contrast, individual KIR family members are expressed on certain NK cell subsets and exhibit finer specificity for HLA class I allotypes and can distinguish between groups of HLA-A, HLA-B, and HLA-C allotypes. Ligation of such KIRs/CD94 to HLA class I molecules on self cells results in inhibition of NK cell cytotoxic activity, as originally predicted by the “missing-self” hypothesis [1, 27]. This regulation ensures

vitro in response to IL-2 [25, 26]. In contrast, resting CD56<sup>dim</sup> NK cells, which express IL-2R $\beta\gamma$  only, show almost no proliferation in response to even high doses of IL-

**Fig. 6** Liver NK cells showed more vigorous cytotoxicity against breast cancer cell lines compared with PB NK cells. Cytotoxic activities of NK cells isolated from IL-2-stimulated PBMCs and LMNCs with or without trastuzumab against target cells (MDA-MB231, MDA-MB453, MDA-MB468, and MCF-7) were analyzed by the <sup>51</sup>Cr release assay. NK cells were isolated from PBMCs and LMNCs after stimulation with IL-2 for 5 days by magnetic sorting (purity > 90%). Data represent the mean  $\pm$  SEM of values from triplicate samples and represent four similar experiments



**Fig. 7** NK cell-induced cytotoxicity was inhibited partially by the anti-TRAIL mAb alone and more profoundly by the combination of anti-TRAIL mAb and CMA. Isolated PB and liver NK cell populations were used as effector cells (*E*) in assays of cytotoxicity against the target (*T*) MDA-MB231 human breast cancer cell line. Cytotoxicity assays were performed at an *E/T* ratio of 10:1, 5:1, or 2.5:1 in the presence or in the absence of anti-TRAIL (N2B2) mAb (10 µg/ml) and/or concanamycin A (CMA) (50 nmol/l). Data are the average ± SEM values from triplicate samples and represent four similar experiments, in which five different donor individuals were used (the results of four other experiments are shown in Supplementary Figures 1 and 2). Error bars not shown appear within the data point. *NK* natural killer, *TRAIL* tumor necrosis factor-related apoptosis-inducing ligand, *mAb* monoclonal antibody, *FasL* Fas ligand, *E/T* effector-to-target, *CMA* concanamycin A




that cells expressing none, altered, or reduced MHC-I molecules, such as malignant or virus-infected cells, are eliminated by NK cells. The modulated expression of KIRs/CD94 by IL-2 is likely associated with the changed cytotoxic target-discriminating ability of NK cells upon their exposure to IL-2.

The significantly upregulated TRAIL expression on the IL-2-stimulated NK cells implies that they have the ability to target cancer cells expressing death-inducing receptors. TRAIL is a member of the TNF superfamily, which includes TNF and FasL [28]. The expression of TNF and FasL leads to damage of normal tissues in addition to their proapoptotic

effect on transformed cells [29, 30], limiting their clinical applications. Conversely, TRAIL selectively induces apoptosis in transformed cells but not in most normal cells [28, 31, 32], making it a promising candidate for tumor therapy. However, intravenous delivery of recombinant TRAIL has met with problems, including a short pharmacokinetic half-life [32], necessitating frequent and high doses to produce the desired effect. The use of TRAIL-expressing NK cells as a delivery vector might promise both targeted and prolonged delivery of this death ligand.

TRAIL binds DR4 and DR5, leading to the formation of the death-inducing signaling complex and the



**Fig. 8** CXC chemokine receptor 3 (CXCR3) expression was significantly upregulated on both liver and PB NK cells after cultivation with IL-2. **a** Histograms representing the log fluorescence intensities for CXCR3 expression on freshly isolated and IL-2-stimulated liver and PB NK cells. Dotted lines represent negative control staining with isotype-matched mAbs. Numbers (mean ± SEM) indicate the percentages of cells in each group that were positive for CXCR3 expression (PBMCs;  $n = 6$ , LMNCs;  $n = 7$ ). Histogram profiles are representative of independent experiments. **b** Numbers indicate the mean fluorescence intensity (MFI) of cells that stained positively for CXCR3 on liver and PB NK cells (fresh NK cells open column, activated NK cells closed column). Data represent mean ± SEM (PBMCs;  $n = 6$ , LMNCs;  $n = 7$ ). Statistical analyses were performed using the paired Student's  $t$  test (\*\* $P < 0.01$ ). NK natural killer, LMNC liver mononuclear cell, PBMC peripheral blood mononuclear cell, mAb monoclonal antibody


Fas-associated protein with death domain. In turn, these complexes recruit caspase-8 (or caspase-10), which plays an important role in apoptosis induction either by direct activation of downstream effector caspases (caspase-3, caspase-6, and caspase-7) or by cleaving apoptotic molecules (Bcl-2 and Bcl-xL), resulting in further activation of the caspase-9 complex [33]. In this study, breast cancer cells of clinical

samples showed much higher expression of TRAIL-DR4 than normal mammary glands but exhibited little TRAIL-DcR1, regardless of HER2 type. Similarly, all the tested breast cancer cell lines expressed TRAIL-DR4 but not TRAIL-DcR1 and TRAIL-DcR2, regardless of their HER2-status, suggesting that they are susceptible to TRAIL-induced apoptosis.

We tested various breast cancer cell lines to evaluate their susceptibility to NK cell-mediated cytotoxicity. Notably, liver NK cells showed more vigorous cytotoxicity against all the tested cell lines than did PB NK cells (Fig. 6), although the underlying mechanism remains unclear. The contribution of TRAIL to NK cell cytotoxicity was determined using the neutralizing anti-TRAIL mAb (Fig. 7). Trastuzumab addition remarkably enhanced the cytotoxicity of both NK cell types toward HER2-overexpressing breast cancer cell lines, indicating that HER2/trastuzumab-mediated ADCC was involved. As ADCC requires the activation and engagement of the CD16 Fc $\gamma$ R on NK cells by Ab-coated targets, CD56<sup>dim</sup> NK cells, which highly express CD16 (Fc $\gamma$  receptor III), are generally thought to exhibit greater levels of ADCC than do the CD56<sup>bright</sup> subset [34]. On the other hand, the majority of CD56<sup>bright</sup> NK cells expanded after activation with IL-2 expressed CD16 and efficiently mediated ADCC [20], explaining the HER2/trastuzumab-mediated ADCC observed in this study.


The cytotoxic ability of NK cells against cancer cells presumably requires contact between NK cells and their target cells. In general, NK cells are detected infrequently in tumors and their presence in the infiltrate consistently correlates with a good prognosis and increased patient survival [35, 36]. Chemokines acting on CXCR3 and CX3CR1 are considered major determinants of NK cell infiltration. CX3CR1 expression in gastric adenocarcinoma samples directly correlates with the number of NK cells infiltrating the tumor, and patients with higher CX3CL1 levels had a significantly better prognosis than patients with low CX3CL1 levels [37]. Similarly, our in vitro demonstration that the CXCL10/CXCR3 axis plays a role in the attraction between activated NK cells and breast cancer cells suggests that this chemokine system recruits NK cells to cancer cell sites and elicits antitumoral responses. In addition, we proposed a novel mechanistic paradigm in which IFN- $\gamma$  secreted from NK cells promotes the production of CXCL10 from breast cancer cells, which in turn further accelerates the migration of CXCR3-expressing NK cells into the tumor site (Fig. 10).

Given the efficacy of NK cells to selectively eliminate abnormal cells, a variety of approaches have been taken to selectively augment NK cell response to tumors [38, 39]. Several therapeutic cytokines primarily act through NK cells (e.g., IL-2, IL-12, IL-15, and IFNs), and many studies have shown that activation of NK cell differentiation and



**Fig. 9** NK cells were preferentially drawn by chemokines secreted from breast cancer cells, presumably through the CXCL10/CXCR3 axis. **a** Levels of various chemokines (CXCL9, CXCL10, and CXCL11) in the culture supernatants of breast cancer cell lines (MDA-MB231, MDA-MB453, MDA-MB468, and MCF-7) were analyzed using CBA Flex Sets. Supernatants were collected after 2 days of cultivation. **b** Migration assays were performed in transwell culture inserts with 3-μm pore filters. MDA-MB231 and MDA-MB468 cell lines were cultured in the lower chamber of the plate for 2 days and IL-2-activated NK cells from PBMCs and LMNCs were added to the upper chamber. After 2 h, the migrated NK cells were counted. Results are presented as mean migrated cell numbers ± SEM (*n* = 3). **c** Levels of chemokine in the medium of lower chambers in the migration assays described above were

measured by CBA assay. Results are the average ± SEM values from triplicate samples and represent three individual experiments. **d** Levels of IFN-γ in the medium of lower chambers in the migration assays described above were measured by ELISA. The results are the average ± SEM values from triplicate samples and represent three individual experiments. **e** MDA-MB231 cells were cultured with various doses of IFN-γ for 2 and 6 h, and the levels of CXCL10 in the medium were measured by CBA assay. Results are the average ± SEM values from triplicate samples and represent three individual experiments. **f** MDA-MB468 cells were cultured with various doses of IFN-γ for 2 and 6 h, and the levels of CXCL10 in the medium were measured by CBA assay. Results are the average ± SEM values from triplicate samples and represent three individual experiments



**Fig. 10** Mechanistic paradigm of interaction between NK and breast cancer cells.  $\text{IFN-}\gamma$  secreted from NK cells promotes the production of CXCL10 from breast cancer cells, which in turn further accelerates migration of CXCR3-expressing NK cells into the tumor site. Migrated NK cells kill breast cancer cells by either of the two major mechanisms that require direct contact between NK cells and target

cells. In the first, cytoplasmic granule toxins, perforin, and granzymes are secreted by exocytosis and together induce apoptosis of the target cell. The second mechanism involves the engagement of death receptors on target cells by expressing of their cognate ligands (TRAIL) on NK cells, resulting in apoptosis of the target cells

function leads to more efficient elimination of tumor growth [40–44]. Despite these promising advances, the systemic administration of cytokines, such as IL-2, which non-specifically activate a broad range of immune cell types, is associated with significant toxicity [40, 45]. Recent animal experiments have demonstrated the ability of adoptive transfer of NK cells to mount a therapeutic antitumor response [46, 47], and translational clinical research suggests that NK cells are useful for controlling human malignancy [48–50]. Our results have proven that PB NK cells can kill breast cancer cells and liver NK cells can hinder metastasis of breast cancer to the liver, which suggests the potential therapeutic use of NK cells, i.e., by either activation of endogenous NK cells or adoptive transfer of in vitro-activated autologous NK cells. Although liver NK cells displayed higher cytotoxicity than PB NK cells, no clinically applicable method for obtaining liver NK cells from patients with breast cancer has yet been established. Alternatively, locally infusing IL-2 into the liver through the portal vein likely activates endogenous liver NK cells, which in turn might infiltrate or accumulate to the tumor site probably through the CXCL10/CXCR3 axis.

**Acknowledgments** We thank Drs. Kohei Ishiyama, and Masahiro Ohira for their advice and encouragement and Drs. Doshali Marlen, Yuka Igarashi and Nabin Basnet, and Ms. Yuko Ishida and Ms. Midori Kiyokawa for their expert technical assistance. This work was

supported by a Grant-in-Aid for Scientific Research (A) from the Japan Society for the Promotion of Science and a Grant-in-Aid for the Research on Hepatitis and BSE from the Japanese Ministry of Health, Labour and Welfare.

**Conflict of interest** None.

## References

1. Trinchieri G (1989) Biology of natural killer cells. *Adv Immunol* 47:187–376
2. Robertson MJ, Ritz J (1990) Biology and clinical relevance of human natural killer cells. *Blood* 76(12):2421–2438
3. Cullen SP, Martin SJ (2008) Mechanisms of granule-dependent killing. *Cell Death Differ* 15(2):251–262
4. Pardo J, Aguilo JI, Anel A, Martin P, Joeckel L, Borner C et al (2009) The biology of cytotoxic cell granule exocytosis pathway: granzymes have evolved to induce cell death and inflammation. *Microbes Infect* 11(4):452–459
5. Sjöström-Mattsson J, Von Boguslawski K, Bengtsson NO, Mjåland I, Salmenkivi K, Blomqvist C (2009) The expression of p53, bcl-2, bax, fas and fasL in the primary tumour and lymph node metastases of breast cancer. *Acta Oncol* 48(8):1137–1143
6. Lahiry L, Saha B, Chakraborty J, Adhikary A, Hossain DM et al (2010) Theaflavins target Fas/caspase-8 and Akt/pBad pathways to induce apoptosis in p53-mutated human breast cancer cells. *Carcinogenesis* 31(2):259–268
7. Takahashi M, Ogasawara K, Takeda K, Hashimoto W, Sakihara H, Kumagai K et al (1996) LPS induces NK1.1+ alpha beta T

- cells with potent cytotoxicity in the liver of mice via production of IL-12 from Kupffer cells. *J Immunol* 156(7):2436–2442
8. Crispe IN, Mehal WZ (1996) Strange brew: T cells in the liver. *Immunol Today* 17(11):522–525
  9. Seki S, Habu Y, Kawamura T, Takeda K, Dobashi H, Ohkawa T et al (2000) The liver as a crucial organ in the first line of host defense: the roles of Kupffer cells, natural killer (NK) cells and NK1.1 Ag+ T cells in T helper 1 immune responses. *Immunol Rev* 174:35–46
  10. Ishiyama K, Ohdan H, Mitsuta H, Arihiro K, Asahara T (2006) Difference in cytotoxicity against hepatocellular carcinoma between liver and periphery natural killer cells in humans. *Hepatology* 43(2):362–372
  11. Ochi M, Ohdan H, Mitsuta H, Onoe T, Tokita D, Hara H et al (2004) Liver NK cells expressing TRAIL are toxic against self hepatocytes in mice. *Hepatology* 39(5):1321–1331
  12. Rahman M, Davis SR, Pumphrey JG, Bao J, Nau MM, Meltzer PS et al (2009) TRAIL induces apoptosis in triple-negative breast cancer cells with a mesenchymal phenotype. *Breast Cancer Res Treat* 113(2):217–230
  13. Dastjerdi K, Tabar GH, Dehghani H, Haghparast A (2011) Generation of an enriched pool of DNA aptamers for an HER2-overexpressing cell line selected by cell SELEX. *Biotechnol Appl Biochem* 58(4):226–230
  14. Kataoka T, Shinohara N, Takayama H, Takaku K, Kondo S, Yonehara S et al (1996) Concanamycin A, a powerful tool for characterization and estimation of contribution of perforin- and Fas-based lytic pathways in cell-mediated cytotoxicity. *J Immunol* 156(10):3678–3686
  15. Kawarabayashi N, Seki S, Hatsuse K, Ohkawa T, Koike Y, Aihara T et al (2000) Decrease of CD56(+)T cells and natural killer cells in cirrhotic livers with hepatitis C may be involved in their susceptibility to hepatocellular carcinoma. *Hepatology* 32(5):962–969
  16. Pan G, O'Rourke K, Chinnaiyan AM, Gentz R, Ebner R, Ni J et al (1997) The receptor for the cytotoxic ligand TRAIL. *Science* 276(5309):111–113
  17. Sheridan JP, Marsters SA, Pitti RM, Gurney A, Skubatch M, Baldwin D et al (1997) Control of TRAIL-induced apoptosis by a family of signaling and decoy receptors. *Science* 277(5327):818–821
  18. Wallin RP, Screpanti V, Michaelsson J, Grandien A, Ljunggren HG (2003) Regulation of perforin-independent NK cell-mediated cytotoxicity. *Eur J Immunol* 33(10):2727–2735
  19. Salazar-Mather TP, Orange JS, Biron CA (1998) Early murine cytomegalovirus (MCMV) infection induces liver natural killer (NK) cell inflammation and protection through macrophage inflammatory protein 1alpha (MIP-1alpha)-dependent pathways. *J Exp Med* 187(1):1–14
  20. Guimond MJ, Wang B, Croy BA (1998) Engraftment of bone marrow from severe combined immunodeficient (SCID) mice reverses the reproductive deficits in natural killer cell-deficient tg epsilon 26 mice. *J Exp Med* 187(2):217–223
  21. Hedrick JA, Saylor V, Figueroa D, Mizoue L, Xu Y, Menon S et al (1997) Lymphotoxin is produced by NK cells and attracts both NK cells and T cells in vivo. *J Immunol* 158(4):1533–1540
  22. Groom JR, Luster AD (2011) CXCR3 ligands: redundant, collaborative and antagonistic functions. *Immunol Cell Biol* 89(2):207–215
  23. Caligiuri MA, Zmuidzinis A, Manley TJ, Levine H, Smith KA, Ritz J (1990) Functional consequences of interleukin 2 receptor expression on resting human lymphocytes. Identification of a novel natural killer cell subset with high affinity receptors. *J Exp Med* 171(5):1509–1526
  24. Nagler A, Lanier LL, Phillips JH (1990) Constitutive expression of high affinity interleukin 2 receptors on human CD16-natural killer cells in vivo. *J Exp Med* 171(5):1527–1533
  25. Caligiuri MA, Murray C, Robertson MJ, Wang E, Cochran K, Cameron C et al (1993) Selective modulation of human natural killer cells in vivo after prolonged infusion of low dose recombinant interleukin 2. *J Clin Invest* 91(1):123–132
  26. Baume DM, Robertson MJ, Levine H, Manley TJ, Schow PW, Ritz J (1992) Differential responses to interleukin 2 define functionally distinct subsets of human natural killer cells. *Eur J Immunol* 22(1):1–6
  27. Karre K, Ljunggren HG, Piontek G, Kiessling R (1986) Selective rejection of H-2-deficient lymphoma variants suggests alternative immune defence strategy. *Nature* 319(6055):675–678
  28. Wiley SR, Schooley K, Smolak PJ, Din WS, Huang CP, Nicholl JK et al (1995) Identification and characterization of a new member of the TNF family that induces apoptosis. *Immunity* 3(6):673–682
  29. Ogasawara J, Watanabe-Fukunaga R, Adachi M, Matsuzawa A, Kasugai T, Kitamura Y et al (1993) Lethal effect of the anti-Fas antibody in mice. *Nature* 364(6440):806–809
  30. Nagata S (1997) Apoptosis by death factor. *Cell* 88(3):355–365
  31. Ashkenazi A, Pai RC, Fong S, Leung S, Lawrence DA, Marsters SA et al (1999) Safety and antitumor activity of recombinant soluble Apo2 ligand. *J Clin Invest* 104(2):155–162
  32. Walczak H, Miller RE, Ariail K, Gliniak B, Griffith T, Kubin M et al (1999) Tumor necrosis factor-related apoptosis-inducing ligand in vivo. *Nat Med* 5(2):157–163
  33. Herr I, Debatin KM (2001) Cellular stress response and apoptosis in cancer therapy. *Blood* 98(9):2603–2614
  34. Nagler A, Lanier LL, Cwirla S, Phillips JH (1989) Comparative studies of human FcR3-positive and negative natural killer cells. *J Immunol* 143(10):3183–3191
  35. Ishigami S, Natsugome S, Tokuda K, Nakajo A, Xiangming Che, Iwashige H et al (2000) Prognostic value of intratumoral natural killer cells in gastric carcinoma. *Cancer* 88(3):577–583
  36. Villegas FR, Santiago C, Villarrubia VG, Jimenez R, Chillon MJ, Jareno J et al (2002) Prognostic significance of tumor infiltrating natural killer cells subset CD57 in patients with squamous cell lung cancer. *Lung Cancer* 35(1):23–28
  37. Hyakudomi M, Matsubara T, Hyakudomi R, Yamamoto T, Kinugasa S, Yamanoi A et al (2008) Increased expression of fractalkine is correlated with a better prognosis and an increased number of both CD8+ T cells and natural killer cells in gastric adenocarcinoma. *Ann Surg Oncol* 15(6):1775–1782
  38. Miller JS, Soignier Y, Panoskaltis-Mortari A, McNearney SA, Yun GH, Fautsch SK et al (2005) Successful adoptive transfer and in vivo expansion of human haploidentical NK cells in patients with cancer. *Blood* 105(8):3051–3057
  39. Klingemann HG, Martinson J (2004) Ex vivo expansion of natural killer cells for clinical applications. *Cytotherapy* 6(1):15–22
  40. Rosenberg SA, Lotze MT, Muul LM, Chang AE, Avis FP, Leitman S et al (1987) A progress report on the treatment of 157 patients with advanced cancer using lymphokine-activated killer cells and interleukin-2 or high-dose interleukin-2 alone. *N Engl J Med* 316(15):889–897
  41. Brunda MJ, Luistro L, Warriar RR, Wright RB, Hubbard BR, Murphy M et al (1993) Antitumor and antimetastatic activity of interleukin 12 against murine tumors. *J Exp Med* 178(4):1223–1230
  42. Takeda K, Hayakawa Y, Smyth ML, Kayagaki N, Yamaguchi N, Kakuta S et al (2001) Involvement of tumor necrosis factor-related apoptosis-inducing ligand in surveillance of tumor metastasis by liver natural killer cells. *Nat Med* 1:94–100
  43. Kennedy MK, Glaccum M, Brown SN, Butz EA, Viney JL, Embers M et al (2000) Reversible defects in natural killer and memory CD8 T cell lineages in interleukin 15-deficient mice. *J Exp Med* 191(5):771–780

44. Smyth MJ, Cretney E, Takeda K, Wilttrout RH, Sedger LM, Kayagaki N et al (2001) Tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) contributes to interferon gamma-dependent natural killer cell protection from tumor metastasis. *J Exp Med* 193(6):661–670
45. Talmadge JE, Phillips H, Schindler J, Tribble H, Pennigton R (1987) Systematic preclinical study on the therapeutic properties of recombinant human interleukin 2 for the treatment of metastatic disease. *Cancer Res* 47(21):5725–5732
46. Helms MW, Prescher JA, Cao YA, Schaffert S, Contag CH (2010) IL-12 enhances efficacy and shortens enrichment time in cytokine-induced killer cell immunotherapy. *Cancer Immunol Immunother* 59(9):1325–1334
47. Pegram HJ, Haynes NM, Smyth MJ, Kershaw MH, Darcy PK (2010) Characterizing the anti-tumor function of adoptively transferred NK cells in vivo. *Cancer Immunol Immunother* 59(8):1235–1246
48. Iliopoulou EG, Kountourakis P, Karamouzis MV, Doufexis D, Ardavanis A, Baxevanis CN et al (2010) A phase I trial of adoptive transfer of allogeneic natural killer cells in patients with advanced non-small cell lung cancer. *Cancer Immunol Immunother* 59(12):1781–1789
49. Milani V, Stangl S, Issels R, Gehrman M, Wagner B, Hube K (2009) Anti-tumor activity of patient-derived NK cells after cell-based immunotherapy—a case report. *J Transl Med* 7:50
50. Knorr DA, Kaufman DS (2010) Pluripotent stem cell-derived natural killer cells for cancer therapy. *Transl Res* 156(3):147–154



## Prevention of recurrence after curative treatment for hepatocellular carcinoma

Tsuyoshi Kobayashi · Kohei Ishiyama · Hideki Ohdan

Received: 16 August 2012 / Accepted: 26 October 2012  
© Springer Japan 2012

**Abstract** Hepatocellular carcinoma often recurs even after curative treatment. In addition to its high frequency of metastasis, hepatocellular carcinoma recurrence is characterized by multicentric carcinogenesis arising in the liver damaged by viral infection with the hepatitis B or hepatitis C virus. This is considered to complicate the initial treatment and recurrence prevention strategy for hepatocellular carcinoma, and accordingly, there is no established adjuvant therapy to prevent recurrence. Preventive adjuvant therapy should be administered to high-risk patients, and should be optimized based on individual risk factors. This review will summarize the current status and future prospects of preventive therapy for the recurrence of hepatocellular carcinoma after curative treatment. Although transcatheter arterial embolization/chemoembolization prior to curative treatment can induce tumor necrosis in some patients, several studies have failed to show any improvement in survival. Postoperative interferon therapy may contribute to prolonging the survival in specific groups of patients. No established adjuvant therapy against advanced hepatocellular carcinoma that prevents metastasis has been established so far. Novel treatment strategies incorporating molecular and immunological mechanisms are expected in the future.

**Keywords** Hepatocellular carcinoma · Adjuvant therapy · Recurrence

---

T. Kobayashi (✉) · K. Ishiyama · H. Ohdan  
Department of Gastroenterological and Transplant Surgery,  
Applied Life Sciences, Institute of Biomedical and Health  
Sciences, Hiroshima University, 1-2-3 Kasumi,  
Minami-ku, Hiroshima 734-8551, Japan  
e-mail: tsukoba@hiroshima-u.ac.jp

### Introduction

Hepatocellular carcinoma (HCC) often recurs even after curative treatment. In addition to its high frequency of metastasis, HCC recurrence is characterized by multicentric carcinogenesis arising in the liver damaged by infections with hepatitis B virus (HBV) and hepatitis C virus (HCV). This is considered to complicate the initial treatment and recurrence prevention strategies for HCC.

According to the Clinical Practice Guidelines for Hepatocellular Carcinoma—The Japan Society of Hepatology 2009 update, the choice of treatment for HCC is based on three factors: the degree of liver damage, the number of tumors and the tumor diameter [1]. Treatments include liver resection, local ablation therapy, transcatheter arterial embolization/chemoembolization (TAE/TACE), hepatic arterial infusion chemotherapy, liver transplantation and palliative care. Among these treatments, liver resection, local ablation therapy and liver transplantation are considered to be curative treatments. However, HCC frequently recurs even after curative treatment, and there is no established adjuvant therapy to prevent recurrence. Such preventive adjuvant therapy should be administered to high-risk patients, and the administration of such treatments should be optimized based on individual risk factors. This review will summarize the current status and future prospects of preventive therapy for HCC recurrence after curative treatment.

Risk factors for the recurrence of hepatocellular carcinoma

The risk factors for HCC recurrence after curative treatment include the tumor stage, vascular invasion, number of tumors, tumor size, capsular formation and liver function [1, 2].

Among these factors, the stage, vascular invasion, number of tumors, tumor size and capsular formation are considered to be related to metastasis. Liver function, however, is likely to be primarily related to the multicentric nature of carcinogenesis. The strategy selected for adjuvant therapy should be based on an individual's risk factors; patients at a high risk of metastasis are good candidates for metastasis-preventing therapies. In high-risk patients with multicentric carcinogenesis, such as those with HCV- or HBV-positive HCC, preventive therapy should include aggressive antiviral therapy.

The vascular invasion and the degree of tumor differentiation play important roles in tumor recurrence after liver transplantation in patients with HCC [2]. Liver transplantation simultaneously treats both the background liver disease and liver tumors; therefore, factors related to the background liver disease are not likely to influence recurrence after transplantation. Cases of early recurrence after transplantation are often caused by the progression of pre-existing micro-metastases, or the implantation of tumor cells circulating in the peripheral blood. Therefore, an effective preventive strategy should focus on providing antitumor effects against these remaining cells. In addition, after liver transplantation, patients must be treated with immunosuppressive agents. Accordingly, it is important to take preventive measures against the reactivation of hepatitis viruses in hepatitis virus-infected patients.

#### Adjuvant therapy prior to curative treatment

Transcatheter arterial embolization/chemoembolization (TAE/TACE) is a procedure wherein embolic material is introduced into the hepatic artery with or without an antitumor agent. This treatment has been administered to patients with unresectable HCC. Two randomized controlled trials

(RCT) demonstrated that TAE/TACE improved the antitumor effect and survival rate compared with conservative treatment in patients with unresectable HCC [3, 4]. TAE/TACE is recommended for patients with advanced hepatocellular carcinoma, which is inoperable, and who are not candidates for local ablation therapy in the Clinical Practice Guidelines for Hepatocellular Carcinoma—The Japan Society of Hepatology 2009 update [1]. TAE/TACE is also administered as a preoperative chemotherapy for resectable HCC prior to hepatic resection or local ablative therapy. More recently, it has been administered prior to liver transplantation.

#### TAE/TACE prior to hepatic resection

A number of studies have reported inconsistent results on the effect of TAE/TACE prior to hepatic resection. Some reports suggest that TAE/TACE improves the prognosis, while others do not. To date, three RCTs have been reported, and none has shown an improvement in the recurrence-free survival, which was the primary endpoint (Table 1) [5–7]. Although the trials differed in terms of their patient inclusion criteria and embolization methods, the results have been uniformly negative.

#### TAE/TACE prior to local ablative therapy

The Clinical Practice Guidelines for Hepatocellular Carcinoma—The Japan Society of Hepatology 2009 update recommend that good candidates for local ablation therapy are patients with liver function graded Child–Pugh class A or B, and three or fewer tumors measuring 3 cm or less in diameter [1]. Conventionally, percutaneous ethanol injection (PEI) has been administered; more recently, radiofrequency ablation (RFA) has become popular. However, the

**Table 1** Randomized clinical trials about TAE/TACE before curative treatment for hepatocellular carcinoma

References	Cases	Inclusion criteria	Regimen	Treatment	5-year survival (vs. control)	5-year disease-free survival (vs. control)	Result
Wu et al. [5]	52	Larger than 10 cm	TACE (EPI)	RES	40 vs. 50 % (3 years)	32 vs. 60 % (3 years)	Not effective
Yamasaki et al. [6]	97	Solitary, 2 to 5 cm	TAE	RES	62.7 vs. 61.7 %	39.1 vs. 31.1 %	Not effective
Koda et al. [8]	52	Smaller than 3 cm	TACE (EPI)	PEI	40.4 vs. 37.7 %	19.3 vs. 80.1 % <sup>a</sup> (3 years)	Effective (recurrence)
Akamatsu et al. [9]	42	Uninodular	TAE	PEI, RFA	82.4 vs. 82.2 % (3 years)	33.8 vs. 34.3 % (3 years)	Not effective
Zhou et al. [7]	108	Larger than 5 cm	TACE (CDDP, MMC, FU)	RES	30.7 vs. 21.1 %	12.8 vs. 8.9 %	Not effective

TAE transcatheter arterial embolization, TACE transcatheter arterial chemo-embolization, EPI epirubicin, CDDP cisplatin, MMC mitomycin C, FU fluorouracil, RES resection, PEI percutaneous ethanol injection, RFA radiofrequency ablation

<sup>a</sup> Recurrence rates

rate of local recurrence after these procedures is high. The use of local ablative therapy preceded by TAE/TACE has been compared to local ablative therapy alone in two RCTs (Table 1) [8, 9]. One RCT compared combination therapy with PEI alone, and the other compared combination therapy with PEI or RFA alone. The results of these studies showed that combination therapy significantly reduced the local recurrence compared with PEI alone, but not with the RFA alone [9]. Neither of the studies found an improvement in the survival rate; therefore, this approach is not recommended in the various treatment guidelines [1, 2, 10].

#### TAE/TACE prior to liver transplantation

It is not yet clear whether preoperative preventive therapy in patients with HCC improves their prognosis after liver transplantation [11]. Several studies have investigated whether preoperative TACE reduces the recurrence of HCC after liver transplantation, but these were retrospective studies, and no RCT has been reported. Decaens et al. [12] reported no significant difference in the 5-year survival rates (59.4 vs. 59.3 %) and 5-year disease-free survival rates (69.3 vs. 64.1 %) between patients treated with preoperative TACE ( $n = 100$ ) and those without adjuvant therapy ( $n = 100$ ) in a multicenter case-control study. Although other studies have shown a favorable prognosis after transplantation in patients who responded well to TACE, and have demonstrated the efficacy of preoperative TACE in reducing the dropout rate while patients were waiting for liver transplantation, there is no evidence demonstrating that TACE improves the overall or recurrence-free survival [13–15].

#### Adjuvant therapy after curative treatment

##### Antiviral therapy in patients with viral hepatitis after curative treatment

Virus eradication by interferon therapy is effective against chronic hepatitis C and compensated cirrhosis type C. Because patients with HCC frequently have viral hepatitis, interferon therapy is administered after curative treatment to eradicate the virus and to repress inflammation. Eight RCTs have investigated the effectiveness of interferon therapy after curative treatment of HCC (Table 2) [16–23]. Shiratori et al. administered PEI to 74 HCV-positive HCC patients with three or fewer tumors; 49 of these patients were subsequently treated with interferon. Interferon therapy did not change the rate of recurrence, but significantly improved the survival [18]. On the other hand, Mazzaferro et al. [21] performed hepatic resection in 150 HCV-positive patients with HCC; 76 of these patients were subsequently treated with interferon. The 5-year recurrence-free survival rate in those treated with interferon (24.3 %) was not significantly different from those not treated with interferon (5.8 %). In summary, one of the RCTs found interferon to be effective in reducing recurrence and improving survival after the curative treatment for HCC, while the other showed a limited effect of interferon in a selected subgroup. However, several meta-analyses of multiple studies have shown that interferon therapy reduced the recurrence and prolonged the survival rate [24–29].

Although interferon was used alone in these studies, the current standard of care for treating chronic hepatitis C is the use of pegylated-interferon in combination with

**Table 2** Randomized clinical trials about adjuvant interferon therapy after curative treatment for hepatocellular carcinoma

References	Cases	Inclusion criteria	IFN	Treatment	5-year survival (vs. control)	5-year recurrence rates (vs. control)	Result
Ikeda et al. [16]	20	HCV	IFN beta	RES, PEI	ND	0 vs. 100 % (2 years)	Effective (recurrence)
Kubo et al. [17]	30	HCV	IFN alpha	RES	ND	ND	Effective (survival)
Shiratori et al. [18]	74	HCV	IFN alpha	PEI	68 vs. 48 %	80 vs. 92 %	Effective (survival)
Lin et al. [19]	30	HBV, HCV	IFN alpha	PAI	ND	47 vs. 90 % (4 years)	Effective (recurrence)
Sun et al. [20]	236	HBV	IFN alpha	RES	63.8 vs. 38.8 m <sup>a</sup>	31.2 vs. 17.7 m <sup>b</sup>	Effective (survival)
Mazzaferro et al. [21]	150	HCV	IFN alpha	RES	63.6 vs. 52.4 %	24.3 vs. 5.8 % <sup>c</sup>	Not effective
Lo et al. [22]	80	HBV	IFN alpha	RES	79 vs. 61 %	ND	Not effective
Chen et al. [23]	268	HBV, HCV	IFN alpha	RES	75.4 vs. 72.5 %	42.7 vs. 45.5 % <sup>c</sup>	Not effective

HCV hepatitis C virus, HBV hepatitis B virus, IFN interferon, RES resection, PEI percutaneous ethanol injection, PAI percutaneous acetic acid injection, ND not described

<sup>a</sup> Median survival time

<sup>b</sup> Median disease-free survival time

<sup>c</sup> Recurrence-free survival rates

ribavirin. This combination therapy has demonstrated a higher rate of sustained virological response (SVR) than interferon monotherapy. Some studies have also evaluated its effectiveness as an adjuvant therapy after curative treatment for HCC [30, 31]. Our recent report showed that postoperative administration of pegylated-interferon plus ribavirin in patients with HCV-positive HCC resulted in a 5-year survival rate of 91.7 %, which was significantly higher than that of the historical control group (50.6 %) [31]. In addition, two recent RCTs have investigated telaprevir, an HCV genotype 1 protease inhibitor. The ADVANCE trial investigated the effect of adding telaprevir to the pegylated-interferon plus ribavirin combination therapy (PR group) in 1,088 untreated genotype 1 hepatitis C patients [32]. Telaprevir was used with the combination therapy for either 12 weeks (T12PR) or 8 weeks (T8PR). The SVR was 44 % in the PR group, compared with 75 and 69 % in the T12PR and T8PR groups, respectively. This demonstrates a significant additive effect of the combination therapy. The REALIZE trial investigated the effect of adding telaprevir to pegylated-interferon plus ribavirin combination therapy (PR group) in 663 treated genotype 1 hepatitis C patients [33]. The telaprevir combination therapy was used either for 12 weeks (T12PR) or for 12 weeks after an initial 4 weeks of PR and was followed by 32 weeks of PR (lead-in T12PR48). The SVR was 17 % in the PR group as compared with 64 and 66 % in the T12PR and lead-in T12PR48 groups, respectively, demonstrating a significant improvement. Based on these positive outcomes, telaprevir combination therapy is likely to become a standard therapy used for HCV genotype 1 hepatitis. Although telaprevir is not recommended for elderly patients, patients with thrombocytopenia or with low hemoglobin, this drug is expected to be useful as an adjuvant therapy for selected patients after curative treatment for HCC.

Nucleoside analogues are effective against hepatitis B. The results of an RCT demonstrated that lamivudine suppressed carcinogenesis arising from hepatitis B [34]. Nucleoside analogues suppress the replication of HBV, repress inflammation and reverse liver fibrosis. Although nucleoside analogues are expected to be used as an adjuvant therapy in patients with hepatitis B virus, no prospective randomized studies have demonstrated its efficacy. It may be difficult to perform an RCT with or without nucleoside analogues after curative treatment for HCC, because these drugs are already recommended for patients with HBV, especially patients with a high viral load.

#### *Adjuvant chemotherapy after curative treatment*

HCC is generally insensitive to anti-cancer drugs, and the response rate to systemic chemotherapy against

unresectable advanced HCC is less than 20 %. In many cases, HCC develops on a background of chronic liver disease; accordingly, worsening liver function can lead to insufficient dosing or a deteriorated prognosis. Until the introduction of sorafenib, there was no standard therapy with proven efficacy for unresectable HCC [35]. Hepatic arterial infusion chemotherapy is considered to result in a high local concentration and have fewer adverse systemic effects, because the systemic concentration of the anti-cancer drug is reduced. A number of small RCTs have investigated the use of various adjuvant chemotherapies for reducing the rates of metastasis and recurrence after curative treatment of HCC (Table 3) [36–47]. Oral, intravenous and intrahepatic arterial administration routes have all been investigated individually or in combination. Although two previous RCTs have reported that neither 1-hexylcarbonyl-5-fluorouracil nor uracil-tegafur could reduce the recurrence of HCC, a recent small RCT reported that oral administration of capecitabine reduced the recurrence rate [36–38]. There is currently no evidence that intravenous chemotherapy is effective. The most frequent route of administration is intrahepatic arterial administration, the efficacy of which was demonstrated in a meta-analysis [48]. However, the included studies were single center experiences, with a small number of patients, and incorporated various treatment regimens. As such, there is no established evidence for a single treatment protocol. Still, some RCTs showed reduced recurrence of HCC with portal vein tumors and improved survival in a population with limited inclusion criteria [39, 45–47]. Because HCC is considered to progress via the portal vein, and patients with complicating portal vein tumors have an increased risk of metastasis and recurrence, hepatic arterial infusion chemotherapy is considered to be important in these patients. Therefore, multicenter studies appropriately designed to target patients with a high risk of metastasis are needed for further investigations.

Sorafenib is a molecule that selectively suppresses receptor tyrosine kinases, including the VEGF receptor and PDGF receptor, as well as the serine/threonine kinase Raf in the MAP kinase cascade [49, 50]. In 2008, the SHARP trial, a large multicenter trial conducted primarily in the US and Europe demonstrated that sorafenib significantly prolonged the survival in patients with advanced hepatocellular carcinoma [35]. Sorafenib is the first agent proven to improve the survival in HCC patients in a large phase III trial. A recent, large clinical trial investigated the efficacy of sorafenib for preventing recurrence after curative therapy for HCC; the results are forthcoming (STORM trial).

Other studies have investigated various adjuvant therapies. In a randomized study, Takayama et al. [51] demonstrated that adoptive immunotherapy after hepatic resection significantly reduced the recurrence rate in 150