

39. Schulz O, Fichlmair A, Rehwinkel J, Rogers NC, Scheuner D, et al. (2010) Protein kinase R contributes to immunity against specific viruses by regulating interferon mRNA integrity. *Cell Host Microbe* 7: 354–361.
40. Min JY, Li S, Sen GC, Krug RM (2007) A site on the influenza A virus NS1 protein mediates both inhibition of PKR activation and temporal regulation of viral RNA synthesis. *Virology* 363: 236–243.
41. Khaperskyy DA, Hatchette TF, McCormick C (2011) Influenza A virus inhibits cytoplasmic stress granule formation. *FASEB J*.
42. Guo Z, Chen LM, Zeng H, Gomez JA, Plowden J, et al. (2007) NS1 protein of influenza A virus inhibits the function of intracytoplasmic pathogen sensor, RIG-I. *Am J Respir Cell Mol Biol* 36: 263–269.
43. Mibayashi M, Martinez-Sobrido L, Loo YM, Cardenas WB, Gale M Jr, et al. (2007) Inhibition of retinoic acid-inducible gene 1-mediated induction of beta interferon by the NS1 protein of influenza A virus. *J Virol* 81: 514–524.
44. Gack MU, Albrecht RA, Urano T, Inn KS, Huang IC, et al. (2009) Influenza A virus NS1 targets the ubiquitin ligase TRIM25 to evade recognition by the host viral RNA sensor RIG-I. *Cell Host Microbe* 5: 439–449.
45. Li W, Li Y, Kedersha N, Anderson P, Emara M, et al. (2002) Cell proteins TIA-1 and TIAR interact with the 3' stem-loop of the West Nile virus complementary minus-strand RNA and facilitate virus replication. *J Virol* 76: 11989–12000.
46. Iseni F, Garcin D, Nishio M, Kedersha N, Anderson P, et al. (2002) Sendai virus trailer RNA binds TIAR, a cellular protein involved in virus-induced apoptosis. *EMBO J* 21: 5141–5150.
47. Simpson-Holley M, Kedersha N, Dower K, Rubins KH, Anderson P, et al. (2010) Formation of antiviral cytoplasmic granules during orthopoxvirus infection. *J Virol* 85: 1581–1593.
48. Yoneyama M, Suhara W, Fukuhara Y, Fukuda M, Nishida E, et al. (1998) Direct triggering of the type I interferon system by virus infection: activation of a transcription factor complex containing IRF-3 and CBP/p300. *EMBO J* 17: 1087–1095.
49. Daly C, Reich NC (1993) Double-stranded RNA activates novel factors that bind to the interferon-stimulated response element. *Mol Cell Biol* 13: 3756–3764.
50. Iwamura T, Yoneyama M, Yamaguchi K, Suhara W, Mori W, et al. (2001) Induction of IRF-3/-7 kinase and NF-kappaB in response to double-stranded RNA and virus infection: common and unique pathways. *Genes Cells* 6: 375–388.
51. Momose F, Kikuchi Y, Komase K, Morikawa Y (2007) Visualization of microtubule-mediated transport of influenza viral progeny ribonucleoprotein. *Microbes Infect* 9: 1422–1433.
52. Onoguchi K, Yoneyama M, Takemura A, Akira S, Taniguchi T, et al. (2007) Viral infections activate types I and III interferon genes through a common mechanism. *J Biol Chem* 282: 7576–7581.
53. Brummelkamp TR, Bernards R, Agami R (2002) A system for stable expression of short interfering RNAs in mammalian cells. *Science* 296: 550–553.
54. Naldini L, Blomer U, Gally P, Ory D, Mulligan R, et al. (1996) In vivo gene delivery and stable transduction of nondividing cells by a lentiviral vector. *Science* 272: 263–267.
55. Zufferey R, Nagy D, Mandel RJ, Naldini L, Trono D (1997) Multiply attenuated lentiviral vector achieves efficient gene delivery in vivo. *Nat Biotechnol* 15: 871–875.
56. Bridge AJ, Pebernard S, Ducraux A, Nicolauz AL, Iggo R (2003) Induction of an interferon response by RNAi vectors in mammalian cells. *Nat Genet* 34: 263–264.
57. Jo S, Kawaguchi A, Takizawa N, Morikawa Y, Momose F, et al. (2010) Involvement of vesicular trafficking system in membrane targeting of the progeny influenza virus genome. *Microbes Infect* 12: 1079–1084.

Development of a drug assay system with hepatitis C virus genome derived from a patient with acute hepatitis C

Kyoko Mori · Youki Ueda · Yasuo Ariumi ·
Hiromichi Dansako · Masanori Ikeda ·
Nobuyuki Kato

Received: 5 October 2011 / Accepted: 1 January 2012 / Published online: 18 January 2012
© Springer Science+Business Media, LLC 2012

Abstract We developed a new cell culture drug assay system (AH1R), in which genome-length hepatitis C virus (HCV) RNA (AH1 strain of genotype 1b derived from a patient with acute hepatitis C) efficiently replicates. By comparing the AH1R system with the OR6 assay system that we developed previously (O strain of genotype 1b derived from an HCV-positive blood donor), we demonstrated that the anti-HCV profiles of reagents including interferon- γ and cyclosporine A significantly differed between these assay systems. Furthermore, we found unexpectedly that rolipram, an anti-inflammatory drug, showed anti-HCV activity in the AH1R assay but not in the OR6 assay, suggesting that the anti-HCV activity of rolipram differs depending on the HCV strain. Taken together, these results suggest that the AH1R assay system is useful for the objective evaluation of anti-HCV reagents and for the discovery of different classes of anti-HCV reagents.

Keywords HCV · Acute hepatitis C · Anti-HCV drug assay system · Anti-HCV activity of rolipram

Introduction

Hepatitis C virus (HCV) infection frequently causes chronic hepatitis, which progresses to liver cirrhosis and hepatocellular carcinoma. HCV is an enveloped virus with a positive single-stranded 9.6 kb RNA genome, which

encodes a large polyprotein precursor of approximately 3,000 amino acid (aa) residues [1, 2]. This polyprotein is cleaved by a combination of the host and viral proteases into at least 10 proteins in the following order: Core, envelope 1 (E1), E2, p7, non-structural 2 (NS2), NS3, NS4A, NS4B, NS5A, and NS5B [1].

Human hepatoma HuH-7 cell culture-based HCV replicon systems derived from a number of HCV strains have been widely used for various studies on HCV RNA replication [3, 4] since the first replicon system (based on the Con1 strain of genotype 1b) was developed in 1999 [5]. Genome-length HCV RNA replication systems (see Fig. 2 for details) derived from a limited number of HCV strains (H77, N, Con1, O, and JFH-1) are also sometimes used for such studies, as they are more useful than the replicon systems lacking the structural region of HCV, although the production of infectious HCV from the genome-length HCV RNA has not been demonstrated to date [3, 4]. Furthermore, these RNA replication systems have been improved enough to be suitable for the screening of anti-HCV reagents by the introduction of reporter genes such as luciferase [3, 4, 6]. We also developed an HuH-7-derived cell culture assay system (OR6) in which genome-length HCV RNA (O strain of genotype 1b derived from an HCV-positive blood donor) encoding renilla luciferase (RL) efficiently replicates [7]. Such reporter assay systems could save time and facilitate the mass screening of anti-HCV reagents, since the values of luciferase correlated well with the level of HCV RNA after treatment with anti-HCV reagents. Furthermore, OR6 assay system became more useful as a drug assay system than the HCV subgenomic replicon-based reporter assay systems developed to date [3, 4], because the older systems lack the Core-NS2 regions containing structural proteins likely to be involved in the events that take place in the HCV-infected human liver.

K. Mori · Y. Ueda · Y. Ariumi · H. Dansako ·
M. Ikeda · N. Kato (✉)
Department of Tumor Virology, Okayama University Graduate
School of Medicine, Dentistry, and Pharmaceutical Sciences,
2-5-1 Shikata-cho, Okayama 700-8558, Japan
e-mail: nkato@md.okayama-u.ac.jp

Indeed, by the screening of preexisting drugs using the OR6 assay system, we have identified mizoribine [8], statins [9], hydroxyurea [10], and teprenone [11] as new anti-HCV drug candidates, indicating that the OR6 assay system is useful for the discovery of anti-HCV reagents.

On the other hand, we previously established for the first time an HuH-7-derived cell line (AH1) that harbors genome-length HCV RNA (AH1 strain of genotype 1b) derived from a patient with acute hepatitis C [12]. In that study, we noticed different anti-HCV profiles of interferon (IFN)- γ or cyclosporine A (CsA) between AH1 and O cells supporting genome-length HCV RNA (O strain) replication [7]. From these results, we supposed that the diverse effects of IFN- γ or CsA were attributable to the difference in HCV strains [12].

To test this assumption in detail, we first developed an AH1 strain-derived assay system (AH1R) corresponding to the OR6 assay system, and then performed a comparative analysis using AH1R and OR6 assay systems. In this article, we report that the difference in HCV strains causes the diverse effects of anti-HCV reagents, and we found unexpectedly by AH1R assay that rolipram, an anti-inflammatory drug, is an anti-HCV drug candidate.

Materials and methods

Reagents

IFN- α , IFN- γ , and CsA were purchased from Sigma-Aldrich (St. Louis, MO). Rolipram was purchased from Wako Pure Chemical Industries (Osaka, Japan).

Plasmid construction

The plasmid pAH1RN/C-5B/PL,LS,TA,(VA)₃ was constructed from pAH1 N/C-5B/PL,LS,TA,(VA)₃ encoding genome-length HCV RNA clone 2 (See Fig. 2) obtained from AH1 cells [12], by introducing a fragment of the RL gene from pORN/C-5B into the *AscI* site before the neomycin phosphotransferase (*Neo*^R) gene as previously described [7].

RNA synthesis

The plasmid pAH1RN/C-5B/PL,LS,TA,(VA)₃ DNA was linearized by *XbaI*, and used for RNA synthesis with T7 MEGAscript (Ambion, Austin TX) as previously described [7].

Cell cultures

AH1R and OR6 cells supporting genome-length HCV RNAs were cultured in Dulbecco's modified Eagle's

medium (DMEM) supplemented with 10% fetal bovine serum (FBS) and 0.3 mg/mL of G418 (Geneticin; Invitrogen, Carlsbad, CA). AH1c-cured cells, which were created by eliminating HCV RNA from AH1 cells [12] by IFN- γ treatment, were also cultured in DMEM supplemented with 10% FBS.

RNA transfection and selection of G418-resistant cells

Genome-length HCV (AH1RN/C-5B/PL,LS,TA,(VA)₃) RNA synthesized *in vitro* was transfected into AH1c cells by electroporation, and the cells were selected in the presence of G418 (0.3 mg/mL) for 3 weeks as described previously [13].

RL assay for anti-HCV reagents

To monitor the effects of anti-HCV reagents, RL assay was performed as described previously [14]. Briefly, the cells were plated onto 24-well plates (2×10^4 cells per well) in triplicate and cultured with the medium in the absence of G418 for 24 h. The cells were then treated with each reagent at several concentrations for 72 h. After treatment, the cells were subjected to a luciferase assay using the RL assay system (Promega, Madison, WI). From the assay results, the 50% effective concentration (EC₅₀) of each reagent was determined.

Quantification of HCV RNA

Quantitative reverse transcription-polymerase chain reaction (RT-PCR) analysis for HCV RNA was performed using a real-time LightCycler PCR (Roche Applied Science, Indianapolis, IN, USA) as described previously [7]. The experiments were done in triplicate.

IFN- α treatment to evaluate the assay systems

To monitor the anti-HCV effect of IFN- α on AH1R cells, 2×10^4 cells and 5×10^5 cells were plated onto 24-well plates (for luciferase assay) and 10 cm plates (for quantitative RT-PCR assay) in triplicate, respectively, and cultured for 24 h. The cells were then treated with IFN- α at final concentrations of 0, 1, 10, and 100 IU/mL for 24 h, and subjected to luciferase and quantitative RT-PCR assays as described above.

Western blot analysis

The preparation of cell lysates, sodium dodecyl sulfate-polyacrylamide gel electrophoresis, and immunoblotting analysis with a PVDF membrane were performed as described previously [13]. The antibodies used in this study were those against HCV Core (CP11 monoclonal antibody;

Institute of Immunology, Tokyo), NS5B, and E2 (generous gifts from Dr. M. Kohara, Tokyo Metropolitan Institute of Medical Science, Japan). Anti- β -actin antibody (AC-15; Sigma, St. Louis, MO, USA) was used as a control for the amount of protein loaded per lane. Immunocomplexes were detected with the Renaissance enhanced chemiluminescence assay (Perkin-Elmer Life Sciences, Boston, MA).

WST-1 cell proliferation assay

The cells were plated onto 96-well plates (1×10^3 cells per well) in triplicate and then treated with rolipram at several concentrations for 72 h. After treatment, the cells were subjected to the WST-1 cell proliferation assay (Takara Bio, Otsu, Japan) according to the manufacturer's protocol. From the assay results, the 50% cytotoxic concentration (CC_{50}) of rolipram was estimated. The selective index (SI) value of rolipram was also estimated by dividing the CC_{50} value by the EC_{50} value.

RT-PCR and sequencing

To amplify the genome-length HCV RNA, RT-PCR was performed separately in two fragments as described previously [7, 15]. Briefly, one fragment covered from 5'-untranslated region to NS3, with a final product of approximately 6.2 kb, and the other fragment covered from NS2 to NS5B, with a final product of approximately 6.1 kb. These fragments overlapped at the NS2 and NS3 regions and were used for sequence analysis of the HCV open reading frame (ORF) after cloning into pBR322MC. PrimScript (Takara Bio) and KOD-plus DNA polymerase (Toyobo, Osaka, Japan) were used for RT and PCR, respectively. The nucleotide sequences of each of the three independent clones obtained were determined using the Big Dye terminator cycle sequencing kit on an ABI PRISM 310 genetic analyzer (Applied Biosystems, Foster City, CA, USA).

Statistical analysis

Differences between AH1R and OR6 cell lines were tested using Student's *t* test. *P* values <0.05 were considered statistically significant.

Results

Development of a luciferase reporter assay system that facilitates the quantitative monitoring of genome-length HCV-AH1 RNA replication

To develop an HCV AH1 strain-derived assay system corresponding to the OR6 assay system [7], a genome-length HCV RNA encoding RL (AH1RN/C-5B/PL,LS,TA,(VA)₃)

was transfected into AH1c cells. Following 3 weeks of culturing in the presence of G418, more than 10 colonies were obtained, and then 8 colonies (#2, #3, #4, #5, #6, #8, #13, and #14) were successfully proliferated. We initially selected colonies #2, #3, and #14 because they had high levels of RL activity ($>4 \times 10^6$ U/ 1.6×10^5 cells) (Fig. 1a). However, RT-PCR and the sequencing analyses revealed that the genome-length HCV-AH1 RNAs obtained from these colonies each had an approximately 1 kb deletion in the E2 region (data not shown). In this regard, we previously observed similar phenomenon and described the difficulty of the development of a luciferase reporter assay system using the genome-length HCV RNA of more than 12 kb [7], suggesting that the NS5B polymerase possesses the limited elongation ability (probably up to a total length of 12 kb). Indeed, in that study, we could overcome this obstacle by the selection of the colony harboring a complete genome-length HCV RNA among the obtained G418-resistant colonies [7]. Therefore, we next carried out the selection among the other colonies. Fortunately, we found that colony #4, showing a rather high level of RL activity (2×10^6 U/ 1.6×10^5 cells), possessed a complete genome-length HCV-AH1 RNA without any deleted forms, although most of the other colonies possessed some amounts of a deleted form in addition to a complete genome-length HCV-AH1 RNA (data not shown). We demonstrated that the HCV RNA sequence was not integrated into the genomic DNA in colony #4 (data not shown). From these results, we finally selected colony #4, and it was thereafter referred to as AH1R and used for the following studies.

We first demonstrated that AH1R cells expressed sufficient levels of HCV proteins (Core, E2, and NS5B) by Western blot analysis for the evaluation of anti-HCV reagents, and the expression levels were almost equivalent to those in OR6 cells (Fig. 1b). In this analysis, we confirmed that the size of the E2 protein in AH1R cells was 7 kDa larger than that in OR6 cells (Fig. 1b), as observed previously [12]. This result indicates that AH1R cells express AH1 strain-derived E2 protein possessing two extra N-glycosylation sites [12]. We next demonstrated good correlations between the levels of RL activity and HCV RNA in AH1R cells (Fig. 1c), as we previously demonstrated in OR6 cells treated with IFN- α for 24 h [7]. These correlations indicate that AH1R cells were as useful as OR6 cells as a luciferase assay system.

Aa substitutions detected in genome-length HCV RNA in AH1R cells

To examine whether or not genome-length HCV RNA in AH1R cells possesses additional conserved mutations such as adaptive mutations, we performed a sequence analysis of HCV RNA in AH1R cells. The results (Fig. 2) revealed that




Fig. 1 Characterization of AH1R cells harboring genome-length HCV RNA. **a** Selection of G418-resistant cell clones. The levels of HCV RNA in G418-resistant cells were monitored by RL assay. **b** Western blot analysis. AH1c, AH1R, and OR6 cells were used for the comparison. Core, E2, and NS5B were detected by Western blot analysis. β -actin was used as a control for the amount of protein loaded per lane. **c** RL activity is correlated with HCV RNA level.

The AH1R cells were treated with IFN- α (0, 1, 10, and 100 IU/mL) for 24 h, and then a luciferase reporter assay (right panel) and quantitative RT-PCR (left panel) were performed. The relative luciferase activity (RLU) (%) or HCV RNA (%) calculated at each point, when the level of luciferase activity or HCV RNA in non-treated cells was assigned to be 100%, is presented here

two additional mutations accompanying aa substitutions (W860R (NS2) and A1218E (NS3)) were detected commonly among the three independent clones sequenced, suggesting that these additional mutations are required for the efficient replication or stability of genome-length HCV RNA. The P1115L (NS3), L1262S (NS3), V1897A (NS4B), and V2360A (NS5A) mutations derived from the sAH1 replicon [12] were conserved in AH1R cell-derived clones. However, AH1-clone-2-specific mutations (T1338A and V1880A) were almost reverted to the consensus sequences of AH1 RNA [12] except for V1880A in AH1R clone 2 (Fig. 2). In addition, the Q63R (Core) mutation was observed in two of three clones (Fig. 2).

Comparison between the AH1R and OR6 assay systems regarding the sensitivities to IFN- α , IFN- γ , and CsA

Using quantitative RT-PCR analysis, we previously examined the anti-HCV activities of IFN- α , IFN- γ , and CsA in AH1 and O cells, and noticed different anti-HCV profiles of IFN- γ and CsA between AH1 and O cells [12]. In that study, AH1 cells seemed to be more sensitive than the O cells to CsA (significant difference was observed

when 0.063, 0.12, or 0.25 μ g/mL of CsA was used). Conversely, AH1 cells seemed to be less sensitive than the O cells to IFN- γ (significant difference was observed when 1 or 10 IU/mL of IFN- γ was used). However, we were not able to determine precisely the EC₅₀ values of these reagents, because of the unevenness of the data obtained by RT-PCR.

After developing the AH1R assay system in this study, we determined the EC₅₀ values of IFN- α , IFN- γ , and CsA using the AH1R assay and compared the values with those obtained by the OR6 assay. The results revealed that AH1R assay was more sensitive than OR6 assay to IFN- α (EC₅₀; 0.31 IU/mL for AH1R, 0.45 IU/mL for OR6) (Fig. 3a) and CsA (EC₅₀; 0.11 μ g/mL for AH1R, 0.42 μ g/mL for OR6) (Fig. 3b), and that the OR6 assay was more sensitive than the AH1R assay to IFN- γ (EC₅₀; 0.69 IU/mL for AH1R, 0.28 IU/mL for OR6) (Fig. 3c). Regarding these anti-HCV reagents, the anti-HCV activities observed between the AH1R and OR6 assays differed significantly in all of the concentrations examined (Fig. 3). In addition, regarding these anti-HCV reagents, cell growth was not suppressed within the concentrations used. Regarding IFN- γ and CsA, the present results clearly support those of our previous





Fig. 2 Aa substitutions detected in intracellular AH1R genome-length HCV RNA. The *upper portion* shows schematic gene organization of genome-length HCV RNA encoding the *RL* gene developed in this study. Genome-length HCV RNA consists of 2 cistrons. In the first cistron, *RL* is translated as a fusion protein with *Neo^R* by HCV-IRES, and in the second cistron, all of HCV proteins (C-NS5B) are translated by encephalomyocarditis virus (EMCV)-IRES introduced in the region upstream of C-NS5B regions. Genome-length HCV RNA-replicating cells possess the G418-resistant phenotype because *Neo^R* is produced by the efficient replication of genome-length HCV RNA. Therefore, when genome-length HCV RNA is excluded from the cells or when its level is decreased, the cells are killed in the presence of G418. In this system, anti-HCV activity is able to evaluate the value of the reporter (*RL* activity) instead of the quantification of HCV RNA or HCV proteins. In addition, it has been known that the infectious HCV is not produced from this RNA replication system [3, 4, 6]. Core to NS5B regions of three independent clones (*AH1R clones 1–3*) sequenced are presented. W860R and A1218E conserved substitutions are indicated by *asterisks*. Q63R substitutions detected in two of three clones are each indicated by a *small dot*. Core to NS5B regions of *AH1 clone 2*, used to establish the AH1R cell line, are also presented. AH1-specific conserved substitutions and *AH1-clone-2*-specific substitutions are indicated by *open circles* and *black circles*, respectively

study [12]. Therefore, we suggest that the diverse effects of these anti-HCV reagents are due to the difference in HCV strains, although we are not able to completely exclude the possibility that AH1R cells are compromised cells causing the different responses against anti-HCV reagents. In summary, the previous and present findings suggest that the AH1R assay system is also useful for the evaluation of anti-HCV reagents as an independent assay system.

Fig. 3 The diverse effects of anti-HCV reagents on AH1R and OR6 assay systems. AH1R and OR6 cells were treated with anti-HCV reagents for 72 h, and then the *RL* assay was performed as described in Fig. 1c. **a** Effect of IFN- α . **b** Effect of CsA. **c** Effect of IFN- γ



Anti-HCV activity of rolipram was clearly observed in the AH1R assay, but not in the OR6 assay

From the above findings, we supposed that the anti-HCV reagents reported to date might show diverse effects between the drug assay systems derived from the different HCV strains. To test this assumption, we used the AH1R and OR6 assay systems to evaluate the anti-HCV activity of more than 10 pre-existing drugs (6-Azauridine, bisindolyl maleimide 1, carvedilol, cehalotaxine, clemizole, 2'-deoxy-5-fluorouridine, esomeprazole, guanazole, hemin, homoharringtonine, methotrexate, nitazoxanide, resveratrol, rolipram, silibinin A, Y27632, etc.), which other groups had evaluated using an assay system derived from the Con1 strain (genotype 1b) or JFH-1 strain (genotype 2a). The results revealed that most of these reagents in the AH1R assay showed similar levels of anti-HCV activities compared with those in the OR6 assay or those of the previous studies (data not shown). However, we found that only rolipram, a selective phosphodiesterase 4 (PDE4) inhibitor [16] that is used as an anti-inflammatory drug, showed moderate anti-HCV activity (EC_{50} 31 μ M; CC_{50} > 200 μ M; SI > 6) in the AH1R assay, but no such activity in the OR6 assay (upper panel in Fig. 4a). This remarkable difference was confirmed by Western blot analysis (lower panel in Fig. 4a). It is unlikely that rolipram's anti-HCV activity is due to the inhibition of exogenous *RL*, *Neo^R* or encephalomyocarditis virus internal ribosomal entry site (EMCV-IRES), all of which are encoded in the genome-length HCV RNA, because the AH1R and OR6 assay systems possess the same structure of genome-length HCV RNA except for HCV ORF. To demonstrate that rolipram's anti-HCV activity is not due to the clonal specificity of the cells or the specificity of genome-length HCV RNA, we examined the anti-HCV activity of rolipram using the monoclonal HCV replicon RNA-replicating cells (sAH1 cells for AH1 strain [12], and sO cells for O strain [13]). The results




Fig. 4 Anti-HCV activity of rolipram. **a** Rolipram sensitivities on genome-length HCV RNA replication in AH1R and OR6 assay systems. AH1R and OR6 cells were treated with rolipram for 72 h, followed by RL assay (black circle with linear line in the upper panels) and WST-1 assay (black triangle with broken line in the upper panels). The relative value (%) calculated at each point, when the level in non-treated cells was assigned to 100%, is presented here. Western blot analysis of the treated cells for the HCV Core was also

performed (lower panels). **b** Rolipram sensitivities on HCV replicon RNA replication in sAH1 and sO cells. sAH1 and sO cells were treated with rolipram for 72 h, and extracted total RNAs were subjected to quantitative RT-PCR for HCV 5' untranslated region as described previously [7]. The HCV RNA (%) calculated at each point, when the level of HCV RNA in non-treated cells was assigned to be 100%, is presented here

revealed by quantitative RT-PCR that rolipram showed moderate anti-HCV activity (EC₅₀ 66 μM) in sAH1 cells, but no such activity in sO cells (Fig. 4b). Anti-HCV activity of rolipram in sAH1 cells was a little weaker than that in AH1R cells (Fig. 4b). The similar phenomenon that the anti-HCV activity in genome-length HCV RNA-based reporter assay is stronger than that in HCV subgenomic replicon-based reporter assay was observed regarding other anti-HCV reagents in our previous studies [14, 17, 18]. This result suggests that the anti-HCV activity of rolipram is not either a clone-specific or genome-length HCV RNA-specific phenomenon. In our previous studies also [14, 18], we demonstrated that anti-HCV activities of several reagents including ribavirin and statins were not due to the clonal specificity of the cells. On the other hand, it was recently reported that rolipram did not show anti-HCV activity in the JFH-1 strain-derived assay [19]. Taken together, the previous and present results suggest that rolipram's anti-HCV activity differs depending on the HCV strain. In summary, rolipram was identified as a new anti-HCV candidate using the AH1R assay system.

Discussion

In the present study, we developed for the first time a drug assay system (AH1R), derived from the HCV-AH1 strain (from a patient with acute hepatitis C), in which HCV-AH1

RNA is efficiently replicated. Using this system, we found that rolipram, an anti-inflammatory drug, had potential anti-HCV activity. This potential had not been detected by preexisting assay systems such as OR6, in which HCV-O RNA was derived from an HCV-positive blood donor. Since an HCV replicon harboring the sAH1 cell line, the parent of the AH1R cell line, was obtained from OR6-cured cells [12], the divergence in rolipram's effects between AH1R and OR6 cells is probably attributable to the difference in HCV strains rather than to the difference in cell clones. Indeed, rolipram's anti-HCV activity was not observed in another ORL8 assay system (O strain), which was recently developed using a new hepatoma Li23 cell line (data not shown) [15]. Therefore, we propose that multiple assay systems derived from different HCV strains are required for the discovery of anti-HCV reagents such as rolipram or for the objective evaluation of anti-HCV activity.

Comparative evaluation analysis of anti-HCV activities of IFN-α, IFN-γ, and CsA using AH1-strain-derived AH1R and O-strain-derived OR6 assay systems demonstrated that each of these anti-HCV reagents showed significantly diverse antiviral effects between the two systems. Regarding IFN-γ and CsA, the present results obtained using a luciferase reporter assay fully supported our previous findings [12] using quantitative RT-PCR analysis. However, in the present analysis, we noticed that IFN-α also showed significantly diverse effects (especially at less than 1 IU/mL) between the AH1R and OR6 assays.

The differences in IFN- α sensitivity may be attributable to the difference in aa sequences in the IFN sensitivity-determining region (ISDR; aa 2209–2248 in the HCV-1b genotype), in which aa substitutions correlate well with IFN sensitivity in patients with chronic hepatitis C [20], because the AH1 strain possesses three aa substitutions (T2217A, H2218R, and A2224 V) in ISDR, whereas the O strain possesses no aa substitutions. However, no report has demonstrated the correlation between IFN sensitivity and the substitution numbers in ISDR using the cell culture-based HCV RNA replication system.

Alternatively, Akuta et al. [21] reported that aa substitutions at position 70 and/or position 91 in the HCV Core region of patients infected with the HCV-1b genotype are pretreatment predictors of null virological response (NVR) to pegylated IFN/ribavirin combination therapy. In particular, substitutions of arginine (R) by glutamine (Q) at position 70, and/or leucine (L) by methionine (M) at position 91, were common in NVR. The patients with position-70 substitutions often showed little or no decrease in HCV RNA levels during the early phase of IFN- α treatment [21]. Regarding this point, it is interesting that position 70 in the AH1 strain is R (wild type) and that in the O strain is Q (mutant type), whereas position 91 is L (wild type) in both strains. Therefore, wild-type R in position 70 of the AH1 strain may contribute to the high sensitivity to IFN- α in the AH1R assay. Regarding positions 70 and 91 of the HCV Core, it is noteworthy that, among all of the HCV strains used thus far to develop HCV replicon systems, only the AH1 strain possesses double wild-type aa (data not shown). Therefore, the AH1R assay system may be useful for further study of sensitivity to IFN/ribavirin treatment.

The anti-HCV activity of rolipram, which is currently used as an anti-inflammatory drug, is interesting, although its anti-HCV mechanism is unclear. As a selective PDE4 inhibitor [16], rolipram may attenuate fibroblast activities that can lead to fibrosis and may be particularly effective in the presence of transforming growth factor (TGF)- β 1-induced fibroblast stimulation [22]. On the other hand, HCV enhances hepatic fibrosis progression through the generation of reactive oxygen species and the induction of TGF- β 1 [23]. Taken together, the previous and present results suggest that rolipram may inhibit both HCV RNA replication and HCV-enhanced hepatic fibrosis. However, it is unclear that rolipram shows anti-HCV activity against the majority of HCV strains, because rolipram has been effective for AH1 strain, but not for O strain. Although rolipram's anti-HCV activity would be HCV-strain-specific, it is not clear which HCV strain is the major type regarding the sensitivity to rolipram. Since developed assay systems using genome-length HCV RNA-replicating cells are limited to several HCV strains including O and AH1

strains to date, further analysis using the assay systems of other HCV strains will be needed to clarify this point.

In this study, we demonstrated that the AH1R assay system, which was for the first time developed using an HCV strain derived from a patient with acute hepatitis C, showed different sensitivities against anti-HCV reagents in comparison with assay systems in current use, such as OR6 assay. Therefore, AH1R assay system would be useful for various HCV studies including the evaluation of anti-HCV reagents and the identification of antiviral targets.

Acknowledgment This study was supported by grants-in-aid for research on hepatitis from the Ministry of Health, Labor, and Welfare of Japan. K. M. was supported by a Research Fellowship for Young Scientists from the Japan Society for the Promotion of Science.

References

1. N. Kato, *Acta Med. Okayama* **55**, 133–159 (2001)
2. N. Kato, M. Hijikata, Y. Ootsuyama, M. Nakagawa, S. Ohkoshi, T. Sugimura, K. Shimotohno, *Proc. Natl. Acad. Sci. USA* **87**, 9524–9528 (1990)
3. R. Bartenschlager, S. Sparacio, *Virus Res.* **127**, 195–207 (2007)
4. D. Moradpour, F. Penin, C.M. Rice, *Nat. Rev. Microbiol.* **5**, 453–463 (2007)
5. V. Lohmann, F. Korner, J. Koch, U. Herian, L. Theilmann, R. Bartenschlager, *Science* **285**, 110–113 (1999)
6. M. Ikeda, N. Kato, *Adv. Drug Deliv. Rev.* **59**, 1277–1289 (2007)
7. M. Ikeda, K. Abe, H. Dansako, T. Nakamura, K. Naka, N. Kato, *Biochem. Biophys. Res. Commun.* **329**, 1350–1359 (2005)
8. K. Naka, M. Ikeda, K. Abe, H. Dansako, N. Kato, *Biochem. Biophys. Res. Commun.* **330**, 871–879 (2005)
9. M. Ikeda, K. Abe, M. Yamada, H. Dansako, K. Naka, N. Kato, *Hepatology* **44**, 117–125 (2006)
10. A. Nozaki, M. Morimoto, M. Kondo, T. Oshima, K. Numata, S. Fujisawa, T. Kaneko, E. Miyajima, S. Morita, K. Mori, M. Ikeda, N. Kato, K. Tanaka, *Arch. Virol.* **155**, 601–605 (2010)
11. M. Ikeda, Y. Kawai, K. Mori, M. Yano, K. Abe, G. Nishimura, H. Dansako, Y. Ariumi, T. Wakita, K. Yamamoto, N. Kato, *Liver Int.* **31**, 871–880 (2011)
12. K. Mori, K. Abe, H. Dansako, Y. Ariumi, M. Ikeda, N. Kato, *Biochem. Biophys. Res. Commun.* **371**, 104–109 (2008)
13. N. Kato, K. Sugiyama, K. Namba, H. Dansako, T. Nakamura, M. Takami, K. Naka, A. Nozaki, K. Shimotohno, *Biochem. Biophys. Res. Commun.* **306**, 756–766 (2003)
14. K. Mori, M. Ikeda, Y. Ariumi, H. Dansako, T. Wakita, N. Kato, *Virus Res.* **157**, 61–70 (2011)
15. N. Kato, K. Mori, K. Abe, H. Dansako, M. Kuroki, Y. Ariumi, T. Wakita, M. Ikeda, *Virus Res.* **146**, 41–50 (2009)
16. S.J. MacKenzie, M.D. Houslay, *Biochem. J.* **347**, 571–578 (2000)
17. M. Yano, M. Ikeda, K. Abe, H. Dansako, S. Ohkoshi, Y. Aoyagi, N. Kato, *Antimicrob. Agents Chemother.* **51**, 2016–2027 (2007)
18. G. Nishimura, M. Ikeda, K. Mori, T. Nakazawa, Y. Ariumi, H. Dansako, N. Kato, *Antiviral Res.* **82**, 42–50 (2009)
19. P. Gastaminza, C. Whitten-Baue, F.V. Chisari, *Proc. Natl. Acad. Sci. USA* **107**, 291–296 (2010)
20. N. Enomoto, I. Sakuma, Y. Asahina, M. Kurosaki, T. Murakami, C. Yamamoto, Y. Ogura, N. Izumi, F. Marumo, C. Sato, *N. Engl. J. Med.* **334**, 77–81 (1996)

21. N. Akuta, F. Suzuki, Y. Kawamura, H. Yatsuji, H. Sezaki, Y. Suzuki, T. Hosaka, M. Kobayashi, M. Kobayashi, Y. Arase, K. Ikeda, H. Kumada, *J. Med. Virol.* **79**, 1686–1695 (2007)
22. S. Togo, X. Liu, X. Wang, *Am. J. Physiol. Lung Cell. Mol. Physiol.* **296**, L959–L969 (2009)
23. W. Lin, W.L. Tsai, R.X. Shao, G. Wu, L.F. Peng, L.L. Barlow, W.J. Chung, L. Zhang, H. Zhao, J.Y. Jang, R.T. Chung, *Gastroenterology* **138**, 2509–2518 (2010)

Development of hepatitis C virus production reporter-assay systems using two different hepatoma cell lines

Midori Takeda,¹ Masanori Ikeda,¹ Yasuo Ariumi,¹† Takaji Wakita² and Nobuyuki Kato¹

Correspondence

Masanori Ikeda
maikeda@md.okayama-u.ac.jp

¹Department of Tumor Virology, Okayama University, Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama 700-8544, Japan

²Department of Virology II, National Institute of Infectious Disease, Tokyo 162-8640, Japan

A hepatitis C virus (HCV) infection system was developed previously using the HCV JFH-1 strain (genotype 2a) and HuH-7 cells, and this cell culture is so far the only robust production system for HCV. In patients with chronic hepatitis C, the virological effects of pegylated interferon and ribavirin therapy differ depending on the HCV strain and the genetic background of the host. Recently, we reported the hepatoma-derived Li23 cell line, in which the JFH-1 life cycle is reproduced at a level almost equal to that in HuH-7-derived RSc cells. To monitor the HCV life cycle more easily, we here developed JFH-1 reporter-assay systems using both HuH-7- and Li23-derived cell lines. To identify any genetic mutations by long-term cell culture, HCV RNAs in HuH-7 cells were amplified 130 days after infection and subjected to sequence analysis to find adaptive mutation(s) for robust virus replication. We identified two mutations, H2505Q and V2995L, in the NS5B region. V2995L but not H2505Q enhanced JFH-1 RNA replication. However, we found that H2505Q but not V2995L enhanced HCV RNA replication of strain O (genotype 1b). We also selected highly permissive D7 cells by serial subcloning of Li23 cells. The expression levels of claudin-1 and Niemann–Pick C1-like 1 in D7 cells are higher than those in parental Li23 cells. In this study, we developed HCV JFH-1 reporter-assay systems using two distinct hepatoma cell lines, HuH-7 and Li23. The mutations in NS5B resulted in different effects on strains O and JFH-1 HCV RNA replication.

Received 21 December 2011

Accepted 26 March 2012

INTRODUCTION

Hepatitis C virus (HCV) infection frequently causes chronic hepatitis and leads to liver cirrhosis and hepatocellular carcinoma. Elimination of HCV by antiviral reagents seems to be the most efficient therapy to prevent fatality.

HCV belongs to the family *Flaviviridae* and contains a positive ssRNA genome of 9.6 kb. The HCV genome encodes a single polyprotein precursor of approximately 3000 aa, which is cleaved by host and viral proteases into at least 10 proteins in the following order: Core, envelope 1 (E1), E2, p7, non-structural 2 (NS2), NS3, NS4A, NS4B, NS5A and NS5B (Kato, 2001; Kato *et al.*, 1990; Tanaka *et al.*, 1996).

Evaluation of anti-HCV reagents was difficult before the development of the HCV replicon system (Lohmann *et al.*, 1999). The HCV replicon system enabled investigation of anti-HCV reagents and the cellular factors involved in HCV RNA replication. Following introduction of the replicon system, genome-length HCV RNA-replication systems and reporter-assay systems were also developed (Ikeda *et al.*, 2002, 2005; Lohmann *et al.*, 2001; Pietschmann *et al.*, 2002). In 2005, an HCV infection system was developed using the genotype 2a JFH-1 strain (Lindenbach *et al.*, 2005; Wakita *et al.*, 2005; Zhong *et al.*, 2005). The JFH-1 infection system has been used to study not only viral RNA replication, but also virus infection and release. This HCV cell-culture system was developed using the human hepatoma cell line HuH-7 and, thus far, HuH-7 is the only cell line to exhibit robust HCV production. Therefore, we intended to test the susceptibility of various other cell lines to HCV RNA replication. We reported previously that the hepatoma cell line Li23 supports robust HCV RNA replication and is also susceptible to authentic JFH-1 infection (Kato *et al.*, 2009). Microarray analysis

†Present address: Center for AIDS Research, Kumamoto University, Kumamoto 860-0811, Japan.

Three supplementary figures are available with the online version of this paper.

revealed that HuH-7 and Li23 cells exhibited distinct gene-expression profiles (Mori *et al.*, 2010). For example, we identified three genes (New York oesophageal squamous cell carcinoma 1, β -defensin-1 and galectin-3) showing Li23-specific expression. Using HuH-7 and Li23 cells in combination with HCV strain O (genotype 1b), we developed drug-assay systems (OR6 and ORL8, respectively) by introducing the *Renilla* luciferase (RL) gene (Ikeda *et al.*, 2005; Kato *et al.*, 2009). We found and reported that the sensitivities to anti-HCV reagents were different between the HuH-7 and Li23 assay systems; for example, the Li23 assay system was 10 times more sensitive to ribavirin than the HuH-7 assay system (Mori *et al.*, 2011). Methotrexate showed very strong anti-HCV activity in the Li23 assay system, although it showed very weak anti-HCV activity in the HuH-7 assay system (Ueda *et al.*, 2011). These results encouraged us to develop a JFH-1 reporter-assay system using HuH-7 and Li23 cells. This JFH-1 reporter-assay system not only facilitated monitoring of virus infection and release steps, but also provided us with new information that could be missed in these steps when using only a HuH-7 assay system. However, increasing the size of the viral genome by introducing exogenous genes [RL and the encephalomyocarditis virus internal ribosomal site (EMCV-IRES)] reduced the efficiency of HCV RNA replication. To overcome this issue, we tried to improve the efficiency of HCV RNA replication by introducing adaptive mutations and by subcloning the parental cells.

Here, we developed JFH-1 HCV production reporter-assay systems in HuH-7- and Li23-derived cells using adaptive mutations and subcloned cells, which monitor the life cycle of HCV with luciferase activity. We also tested the effect of the mutations in NS5B from the JFH-1 strain on RNA replication of the specific genotype 1b O strain.

RESULTS

HCV mutations caused by long-term cell culture

The efficiency of HCV RNA replication depends on viral genetic mutations, host cells and viral genome size. For development of the HCV reporter-assay system, use of a longer viral genome reduced the efficiency of virus replication. To compensate for this issue, we tried to introduce adaptive mutations into the JFH-1 genome. We examined the viral sequences of JFH-1 130 days after infection of HuH-7-derived RSc cells. We performed RT-PCR for three parts of the viral genome: Core to NS2, NS3 to NS5A, and NS5B to 3'X. These three parts were separated by the *AgeI*, *SpeI*, *BsrGI* and *XbaI* sites on the viral genome. We introduced PCR products into the cloning vector and three independent clones were subjected to sequencing analysis.

In the Core to NS2 region between the *AgeI* and *SpeI* sites (designated AS), there were eight common mutations with

amino acid substitutions: lysine to glutamate at aa 78 (K78E) in Core, P251L and A351D in E1, V402A, I414T and K715N in E2, Y771C in p7, and D962G in NS2 (Fig. 1a). In the NS3 to NS5A region between *SpeI* and *BsrGI* sites (designated SB), there were eight common mutations with amino acid substitutions: V1460I and M1611T in NS3, and I2270T, Q2307R, S2363L, M2392T, S2426A and C2441S in NS5A (Fig. 1b). In the NS5B to 3'X region between the *BsrGI* and *XbaI* sites (designated BX), there was only one common mutation with an amino acid substitution, V2995L in NS5B (Fig. 1c). The determined sequences were studied further to enhance HCV RNA replication in the JFH-1 reporter assay.

Effect of genetic mutations on HCV RNA replication

To monitor the virus life cycle more easily, we constructed dicistronic JFH-1 with a reporter gene, pJR/C-5B. The first cistron contained the RL gene and was translated by the HCV-IRES. The second cistron contained the JFH-1 ORF and was translated by the EMCV-IRES. This construct facilitated monitoring of all steps of the virus life cycle by quantification of RL activity. However, the use of a longer viral genome resulted in lower replication efficiency. We tested the effect on HCV RNA replication of amino acid substitution caused during long-term cell culture.

The amino acid substitution clusters from three independent clones in Core to NS2 (AS-1, AS-2, AS-3) were introduced into pJR/C-5B. *In vitro*-transcribed HCV RNA was introduced into HuH-7-derived RSc cells, and RL activities were monitored 24, 48 and 72 h after electroporation (Fig. 2a). AS-3 exhibited higher replication efficiency than the wild type (WT). However, the replication efficiency of AS-2 was almost equal to that of the WT, and AS-1 exhibited lower replication efficiency than the WT. AS-3 possessed the highest replication efficiency among the tested JFH-1 mutants: at 72 h, the luciferase value of this clone was approximately 100 times that at 24 h.

The three pJR/C-5B constructs with mutations in NS3 to NS5A (SB-2, SB-3 and SB-4) were transcribed and introduced into RSc cells to compare the efficiency of HCV RNA replication (Fig. 2b). Unexpectedly, RL activity was not increased over 72 h after electroporation and exhibited a pattern similar to that of JFH-1 without the GDD motif. This result indicates that the mutation in NS3 to NS5A exhibited a negative effect on HCV RNA replication.

Finally, we tested the effect of the mutations in the NS5B region on HCV RNA replication. BX-2 contains two mutations with amino acid substitution (H2505Q and V2995L) and BX-7 contains only V2995L (Fig. 2c). JFH-1 with mutation(s) of BX-2 or BX-7 exhibited strong enhancement of HCV RNA replication. These results indicate that V2995L works as a strong replication-enhancing mutation (REM) in JFH-1 HCV RNA replication.




Fig. 1. Genetic mutations causing amino acid substitutions during long-term HCV infection. RT-PCR was performed for HCV RNAs from HuH-7 cells 130 days after JFH-1 infection. PCR products were subcloned into the pBluescript II plasmid. Three clones of (a) the Core to NS2 region between the *Agel* and *SpeI* sites (AS), (b) the NS3 to NS5A region between the *SpeI* and *BsrGI* sites (SB) and (c) the NS5B to 3'X region between the *BsrGI* and *XbaI* sites (BX) were subjected to sequencing analysis. ▼ and ▽ represent conservative and non-conservative amino acid substitutions, respectively.

Mutations in NS5B enhanced HCV RNA replication differently in genotypes 1b and 2a

V2995L in NS5B is a common substitution, occurring in three clones, and H2505Q is conserved in two clones (BX-2 and BX-10). We examined the corresponding amino acids at positions 2995 and 2505 in genotype 1b replication-competent HCV strains O, 1B-4 and KAH5 (Fig. 3a) (Nishimura *et al.*, 2009). The histidine at aa 2505 in JFH-1 is conserved in O, 1B-4 and KAH5 at the corresponding position, aa 2482. The valine at aa 2995 in JFH-1 is an

alanine in O, 1B-4 and KAH5 at the corresponding position, aa 2972 (Fig. 3a). It is not clear whether the adaptive mutation found in genotype 2a is effective in genotype 1b HCV. Therefore, we investigated the effect of V2995L and/or H2505Q substitution on genotype 1b HCV RNA replication. We introduced substitutions V2995L and/or H2505Q into the subgenomic replicon, pOR/3-5B (HCV-O). In contrast to the case of JFH-1, H2505Q but not V2995L enhanced HCV-O RNA replication (Fig. 3b). These results indicate that the mutations in NS5B derived from JFH-1 functioned differently in genotype 1b HCV-O RNA replication.




Fig. 2. Effect of amino acid substitutions on HCV RNA replication. (a) The Core to NS2 region; (b) the NS3 to NS5A region; (c) the NS5B to 3'X region. Amino acid substitutions were introduced into pJR/C5B and *in vitro*-synthesized RNAs were electroporated into HuH-7-derived RSc cells. RL activity was determined 24, 48 and 72 h after electroporation. dGDD, Negative control without the GDD motif; WT, wild type. * $P < 0.05$; ** $P < 0.01$.

HCV infection in HuH-7- and Li23-derived cell lines

As well as viral genetic mutations, the choice of host cells is important for the efficiency of HCV RNA replication. Cured cells in which HCV RNAs were eliminated by IFN- α , such as HuH-7.5, HuH-7.5.1 and our RSc cells, exhibit




Fig. 3. Effect of amino acid substitutions in NS5B on genotype 1b and 2a HCV RNA replication. (a) Alignment of amino acids at positions 2505 (JFH-1) and 2482 (genotype 1b) and around the adjacent region (upper panel). Alignment of amino acids at positions 2995 (JFH-1) and 2972 (genotype 1b) and around the adjacent region (lower panel). The HCV strains O, KAH5 and 1B-4 belong to genotype 1b. (b) H2505Q and/or V2995L were introduced into the HCV-O subgenomic replicon (pOR/3-5B), and transcribed RNAs were electroporated into RSc cells. RL activities were tested 24, 48 and 72 h after infection. dGDD, Negative control without the GDD motif; WT, wild type. ** $P < 0.01$.

higher replication efficiency than their parental HuH-7 cells (Ariumi *et al.*, 2007; Blight *et al.*, 2002; Zhong *et al.*, 2005). Therefore, we examined whether subcloned Li23 cells might enhance HCV RNA replication. We performed serial subcloning of Li23 cells from Li23-derived ORL8c cells by the limiting-dilution method (Fig. 4a). ORL8c cells are a cured cell line in which genome-length HCV RNAs were eliminated by interferon (IFN) treatment (Kato *et al.*, 2009). The subclonal Li23-derived cell lines were selected from among 50–100 independent single cells in 96-well plates by three-round limiting dilution from ORL8c cells (Fig. S1a, available in JGV Online). First, L8c15 cells were selected from their parental ORL8c cells by limiting dilution. Then, C22 cells were selected from their parental L8c15 cells by limiting dilution. Finally, D7 cells were selected from C22 cells by limiting dilution (Fig. S1b). Together, these steps resulted in the

selection of three subclonal cell lines that respectively exhibited the strongest replication efficiency in each round of selection. The lineages of the selected cell lines after three rounds of subcloning were designated L8c15, C22 and D7 cells, respectively.

We tested the subcloned cells for their HCV infectivities in comparison with those of HuH-7 and HuH-7-derived RSc cells. We reported previously that RSc cells could strongly support HCV replication and production (Kato *et al.*, 2009). Li23 and its derived ORL8c, L8c15, C22 and D7 cell lines were infected using the supernatant from RSc cells replicating JR/C-5B with BX-2 mutations at an m.o.i. of 0.2 (Fig. 4b, c). RL activities were determined 24, 48, 72 and 96

h after infection and f.f.u. ml⁻¹ were determined 48 h after infection. The efficiency of HCV infectivity was highest in D7 cells, followed in order by C22, L8c15 and Li23 cells. HCV RNA replication in D7 cells was almost equal to that in RSc cells. These results suggest that the subcloned cell lines exhibit higher susceptibility to HCV infection than their parental cells.

Next, we further characterized the susceptibility of D7 cells to HCV infection in comparison with RSc cells, because D7 cells exhibited the highest susceptibility to HCV infection among the Li23-derived cell lines. D7 cells also exhibited the highest production and release of Core into the supernatant among the parental C22-derived subclonal




Fig. 4. HCV infection in HuH-7- and Li23-derived cell lines. (a) History of the selection of subclonal Li23-derived cell lines. (b) HuH-7, HuH-7-derived RSc, and Li23-derived ORL8c, L8c15, C22 and D7 cells were inoculated with supernatant from RSc cells replicating JR/C5B/BX-2. ***P* < 0.01. (c) f.f.u. ml⁻¹ values were determined 48 h after infection of HuH-7- and Li23-derived cells with HCV using the supernatant from RSc cells replicating JR/C5B/BX-2. (d) f.f.u. ml⁻¹ values were determined 48 h after infection of RSc or D7 cells with HCV using the supernatant from RSc cells replicating JR/C5B/AS-3 or JR/C5B/BX-2. Supernatant from authentic JFH-1-replicating RSc cells was used as a positive control. (e) Core expression levels in RSc or D7 cells were determined 1, 2, 3 and 4 days after infection with JFH-1 with BX-2 mutations. Lanes: 1 and 6, mock-infected cells; 2 and 7, cells 1 day after infection; 3 and 8, cells 2 days after infection; 4 and 9, cells 3 days after infection; 5 and 10, cells 4 days after infection; 11 and 12, OR6c and OR6 cells, respectively; 13 and 14, ORL8c and ORL8 cells, respectively. OR6 and ORL8 were used as positive controls; OR6c and OR8c were used as negative controls. β-Actin was used as a control for the amount of protein loaded per lane.




Fig. 5. Expression levels of HCV receptors in HuH-7- and Li23-derived cells. Quantitative RT-PCR was performed for CD81, SR-BI, CLDN1, OCLN and NPC1L1 as described in Methods. Relative expression levels of mRNA are shown, when the expression level of each receptor in HuH-7 was assigned to be 1. GAPDH was used as an internal control. Experiments were done in triplicate.

cells (Fig. S1b). The susceptibility of the HCV reporter-assay system to HCV infection was examined using HuH-7- and Li23-derived cells. Supernatants from RSc cells replicating JR/C-5B with AS-3 or BX-2 mutations were used as inocula. The supernatant from authentic JFH-1-replicating RSc cells was used as a positive control. RSc and D7 cells were inoculated with each HCV-containing supernatant and f.f.u. ml⁻¹ were determined 48 h after infection. As shown in Fig. 4(d), the values of f.f.u. ml⁻¹ for AS-3 were 2.5×10^4 and 1.0×10^4 in RSc and D7 cells, respectively; those for BX-2 were 3.1×10^4 and 1.8×10^4 in RSc and D7 cells, respectively; and those for authentic JFH-1 were 2.9×10^5 and 1.2×10^5 in RSc and D7 cells, respectively. These results indicate that the infectivities of these three inocula were almost equal in RSc and D7 cells.

Next we examined Core expression after infection of RSc and D7 cells with HCV, as D7 cells exhibited the highest infectivity among the Li23-derived cell lines (Fig. 4e). Core was detected 2, 3 and 4 days after infection of the supernatant from RSc cells infected by JR/C-5B with BX-2. Although Core expression in D7 cells was slightly weaker than that in RSc cells, the signal of Core in HCV-infected D7 cells was equal to that in stable ORL8 cells. These results suggest that the JFH-1 reporter-assay system in Li23 cells is useful not only for the RL assay, but also for Core expression.

Expression of HCV receptors in parental and subclonal hepatoma cell lines

We tested expression of the HCV receptors CD81, scavenger receptor class B member I (SR-BI), claudin-1 (CLDN1) and occludin (OCLN). We also examined the expression of the recently reported HCV entry factor Niemann-Pick C1-like 1 (NPC1L1) (Sainz *et al.*, 2012). Expression levels of CD81 in Li23 and its subclonal cells were higher than those in HuH-7 and RSc cells (Fig. 5a). Although expression of CD81 in Li23-derived cell lines was lower than that in parental Li23 cells, interestingly the expression levels of CD81 increased during the rounds of selection. There is no difference in the expression of SR-BI among the cell lines tested (Fig. 5b). The expression of CLDN1 in Li23-derived cells was higher than that in parental Li23 cells (Fig. 5c). Expression levels of OCLN in Li23 and its subclonal cells were higher than those in HuH-7 and RSc cells (Fig. 5d). Finally, the expression of NPC1L1 in Li23-derived cell lines was higher than that in parental Li23 cells (Fig. 5e). It is noteworthy that the expression level of NPC1L1 in RSc cells was approximately 2 log₁₀ lower than that in parental HuH-7 cells. Taken together, these results indicate that the expression levels of CLDN1 and NPC1L1 in Li23-derived cells were higher than those in parental Li23 cells.

Life cycle of the HCV reporter-assay system in Li23-derived cells

We investigated whether D7 cells produce infectious HCV. First, D7 cells were inoculated using the supernatant from RSc cells replicating JR/C5B with BX-2, and the supernatant was stored at 17 days after infection. Then, the supernatant derived from the D7 cells was used as an inoculum for reinfection of naïve D7 cells. RL activities were determined 2, 6, 10 and 14 days after reinfection (Fig. 6). RL activity was increased after reinfection in D7 cells and reached a plateau 10 days after reinfection. These data indicate that the JFH-1 reporter-assay system is also useful for monitoring the HCV life cycle in Li23-derived cell lines.

DISCUSSION

In this study, we developed an HCV production reporter-assay system using two distinct hepatoma cell lines, HuH-7 and Li23. Robust HCV RNA replication and virus production were achieved by the introduction of REMs into the structural region or the NS5B region. These REMs were obtained from JFH-1-infected long-term-cultured cells. The two REMs in NS5B (H2505Q and V2995L substitutions) derived from JFH-1 had different effects on replication of genotype 1b HCV-O RNA and genotype 2a JFH-1 RNA. Furthermore, the subcloned Li23-derived D7 cells produced by serial limiting dilution supported this HCV production reporter-assay system.

Several groups have reported JFH-1 reporter virus systems (Koutsoudakis *et al.*, 2006; Marcello *et al.*, 2006; Pietschmann *et al.*, 2002; Wakita *et al.*, 2005). However, robust reporter virus production was limited within the study using HuH-7-derived cells. Therefore, we attempted to develop a JFH-1 reporter virus assay system using our previously reported line of Li23 cells (Kato *et al.*, 2009).




Fig. 6. HCV life cycle in Li23-derived D7 cells. D7 cells were inoculated with the supernatant from D7 cells after infection with JFH-1 with BX-2 mutants. RL activities were tested 2, 6, 10 and 14 days after infection.

The introduction of RL and EMCV-IRES genes into the HCV gene lengthened the genome of HCV by approximately 1.9 kb and led to a reduction in the efficiency of HCV RNA replication. To overcome this disadvantage, we adopted the following strategies: (i) introduce the REMs; (ii) select cloned Li23-derived cells with a highly permissive host condition by the serial limiting-dilution method. For the first purpose, we performed sequence analyses for HCV RNA from JFH-1-infected RSc cells. Mutations in the region from Core to NS2 or NS5B enhanced HCV RNA replication. However, combination of mutations from two different regions reduced HCV RNA replication (Fig. S2). The reason for this may be that these two mutation clusters were obtained from distinct RT-PCR-amplified clones and they were not necessarily located on the same viral genome. It has been reported that the combination of REMs exhibited an antagonistic effect on HCV RNA replication (Lohmann *et al.*, 2001). For the second purpose, we selected highly permissive Li23-derived clonal cells by the limiting-dilution method. We obtained three Li23-derived subclonal cell lines, L8c15, C22 and D7, in order from parental Li23-derived ORL8c cells. The efficiency of infectivity was highest in D7 cells, followed in order by C22, L8c15 and Li23 cells. D7 cells were highly permissive for infection of HCV with NS5B mutations.

As shown in Fig. 3(a), the histidine at aa 2505 in JFH-1 was conserved in the replication-competent O, 1B-4 and KAH5 strains at the corresponding position, aa 2482. The valine at aa 2995 in JFH-1 was alanine in strains O, 1B-4 and KAH5 at the corresponding position, aa 2972. The REMs in genotype 1b HCV were usually obtained by selection with neomycin after HCV RNA electroporation. Pietschmann *et al.* (2009) reported that REMs impaired infectious virus production. Most REMs are located in the NS3 and NS5A regions (Abe *et al.*, 2007; Blight *et al.*, 2002; Lohmann *et al.*, 2001; Pietschmann *et al.*, 2002). NS5A is a key molecule for virus production, and REMs affect the phosphorylation status of NS5A and the interaction with Core (Kato *et al.*, 2008; Masaki *et al.*, 2008; Tellinghuisen *et al.*, 2008). In contrast, our REMs in NS5B were obtained in JFH-1-infected long-term cell culture without drug selection. Taking this information into account, we considered that H2505Q in NS5B might not interfere with genotype 1b virus production. We attempted to apply this REM from genotype 2a to genotype 1b and found that H2505Q enhanced replication of the genotype 1b HCV-O replicon. We are currently investigating whether our NS5B REM could enhance genotype 1b HCV production. As for the substitution at aa 2995 in JFH-1 (aa 2972 in genotype 1b), we should be careful in interpretation, because the backgrounds at this position are different between genotypes 2 and 1. Analysis of an HCV database (<http://s2as02.genes.nig.ac.jp/>) revealed that the consensus amino acids at position 2995 in genotype 2 and at 2972 in genotype 1 were valine and alanine, respectively. Furthermore, alanine and valine are not found at position 2995 in genotype 2 or at 2972 in genotype 1, respectively. These observations

indicate that amino acid substitution between alanine and valine at these positions may be lethal for HCV of both genotypes. The amino acid at position 2995 in genotype 2 (2972 in genotype 1) is just upstream of a *cis*-acting replication element in NS5B. Therefore, the nucleotide at this position may affect the HCV RNA replication. To clarify this issue, further study will be needed.

A comparative study using HuH-7- and Li23-based JFH-1 reporter-assay systems would be expected to reveal new information on virus entry and release steps, because the backgrounds of these cells are different. Our recent study of these cells revealed the difference in sensitivities to anti-HCV reagents including ribavirin and methotrexate (Mori *et al.*, 2011; Ueda *et al.*, 2011). Furthermore, the IL28B genotype was different between HuH-7 and Li23 cells. The IL28B genotype (rs8099917) of HuH-7 cells renders them resistant to pegylated IFN and ribavirin, and Li23 cells are sensitive to pegylated IFN and ribavirin (M. Ikeda and N. Kato, unpublished data).

Recently, it was reported that stable expression of miR122 enhanced JFH-1 HCV production in Hep3B and HepG2 (Kambara *et al.*, 2012; Narbus *et al.*, 2011). It is noteworthy that the expression of miR122 in Li23-derived cells was almost the same as that in HuH-7 cells (Fig. S3). High-level expression of miR122 in Li23 cells may be one of the reasons that Li23 cells can support HCV production as robust as that in HuH-7 cells among the hepatocyte-derived cell lines. Interestingly, the expression levels of miR122 are higher in ORL8c, L8c15 and D7 cells, but not in C22 cells, than those in parental Li23 cells (Fig. S3). This result suggests that the expression level of miR122 may partly contribute to the fitness of HCV replication and production.

So far, we have only little information regarding the mechanism by which subclonal cells support HCV replication and production more efficiently than the parental cells. In this study, we found that the expression levels of CLDN1 and NPC1L1 in Li23-derived subclonal cells were higher than those in the parental Li23 cells. These results suggest that a high expression level of these entry factors in Li23-derived subclonal cells may contribute to enhanced virus entry. In the course of the experiment to determine the expression levels of NPC1L1 in HuH-7- and Li23-derived cell lines, we found that RSc cells expressed a very low level of NPC1L1 compared with the parental HuH-7 cells. Possible mechanisms for this are: (i) very low-level expression of NPC1L1 is sufficient for HCV entry; (ii) an unknown entry factor compensates for NPC1L1 in the entry step in RSc cells. Further study will be needed to clarify this issue.

In summary, we have developed JFH-1 reporter-assay systems using HuH-7-derived RSc and Li23-derived D7 cells. Expression levels of CLDN1 and NPC1L1 were higher than those in the parental Li23 cells. We found different effects of REMs (V2995L and H2505Q) in NS5B on virus RNA replication in genotype 2a and 1b HCV strains. These findings will become useful tools for the study of the life cycle of HCV.

METHODS

Cell cultures. RSc and ORL8c cells were derived from the cell lines HuH-7 and Li23, respectively, as described previously (Kato *et al.*, 2009). L8c15, D7 and C22 cells were selected from ORL8c, L8c15 and C22 cells, respectively, by limiting dilution. HuH-7 and RSc cells were cultured in Dulbecco's modified Eagle's medium (DMEM; Life Technologies) supplemented with 10% FBS (Life Technologies). Li23-derived cell lines were maintained in F12 medium (Life Technologies) and DMEM (1:1 in volume) supplemented with 1% FBS and epidermal growth factor (50 ng ml⁻¹; PeproTech, Inc.) as described previously (Kato *et al.*, 2009).

RT-PCR and sequencing analysis. RSc cells were infected with cell-culture-grown HCV (HCVcc) and cultured for 130 days. Total RNAs from these cells were prepared using an RNeasy extraction kit (Qiagen). These RNA samples were used for RT-PCR in order to amplify the Core to NS2 (4.0 kb), NS3 to NS5A (3.6 kb) and NS5B to 3'X (1.9 kb) regions. Reverse transcription was performed with an oligo(dA)₂₃ primer. The following primer pairs were employed: to amplify the Core to NS2 region, JFH-1/*AgeI* (5'-CCCAAGCTTACCGGTGAGTACACCGGAATTGC-3') and JFH-1/*SpeI* (5'-TGCCA-TGTGCCTTGGATAGGTACG-3'); for the NS3 to NS5A region, JFH-1/*SpeI* (5'-CCCAAGGGGTACAAAGTACTAGTGC-3') and JFH-1/*BsrGI* (5'-CCCAAGCTTTACTTTTTAGCCCTCTGTGAGGC-3'); for the NS5B to 3'X region, JFH-1/*BsrGI* (5'-CCGCTCGAGACCC-TTTGAGTAACTCGCTGTTGC-3') and JFH-1/*XbaI* (5'-GCTCTA-GACATGATCTGCAGAGAGACCAGTTAC-3'). SuperScript III reverse transcriptase (Invitrogen) and KOD-plus DNA polymerase (TOYOBO) were used for reverse transcription and PCR, respectively. PCR products were ligated into pBluescript II (Fermentas) and three independent clones were subjected to sequencing analysis.

Plasmid construction. pJR/C-5B plasmid is a dicistronic HCV JFH-1 construct. The RL gene and HCV ORF were introduced into the first and second cistrons, respectively. To construct this plasmid, we fused the JFH-1 5'UTR with the RL gene by overlap PCR, and the PCR products were ligated into pFGR-JFH-1 (GenBank accession no. AB237837) at the *AgeI* and *PmeI* sites. For the first PCR, the primer pair 5'-GCGCCTAGCCATGGCGTTAGTATG-3' (J5dC) and 5'-AAGCCATGGCCGGCCCTGGGCGACGGTTGGTGTCTTTTGG-3' (J5dCR) was employed to amplify the 5'UTR, and the primer pair 5'-AACCGTCGCCAGGCGCCGATGGCTTCCAAGGTGTACG-ACCCC-3' (JRL) and 5'-TCGAAATCTCGTGATGGCAGGTTGG-3' (JRLR) was employed to amplify the RL region. These first PCR products were used in the second PCR as the templates. For the second PCR, the primer pair J5dC and JRLR was employed to amplify the 5'UTR and RL. KOD-plus DNA polymerase was used for PCR.

The H2505H and/or A2995L mutations were introduced into the HCV-O replicon by QuikChange mutagenesis (Stratagene) as described previously (Ikeda *et al.*, 2002).

Luciferase reporter assay. For the luciferase assay, approximately 1.0–1.5 × 10⁴ HCV-harboring cells were plated onto 24-well plates in triplicate and were cultured for 24–96 h after electroporation or infection, as described previously (Ikeda *et al.*, 2005). The cells were harvested with *Renilla* lysis reagent (Promega) and subjected to RL assay according to the manufacturer's protocol.

Western blot analysis. Preparation of cell lysates, SDS-PAGE and immunoblotting were performed as described previously (Kato *et al.*, 2003). The antibodies used in this study were Core (CP11; Institute of Immunology, Tokyo, Japan) and β -actin (AC-15; Sigma) antibodies. Immunocomplexes were detected with a Renaissance enhanced chemiluminescence assay (PerkinElmer Life Science).

HCV infection and determination of f.f.u. To determine f.f.u. ml⁻¹, 6 × 10³ cells were plated onto a 96-well plate 24 h before infection. The supernatant of HCV RNA-replicating cells was diluted serially and was used as an inoculum. Forty-eight hours after infection, the cells were fixed and Core was stained with anti-Core antibody and HRP-conjugated mouse anti-IgG antibody. Then, the expression of Core was visualized with a DAB substrate kit (DAKO). Culture supernatants and cells were collected for quantification of Core by ELISA (Mitsubishi Kagaku Bio-Clinical Laboratories).

Quantitative RT-PCR analysis. Quantitative RT-PCR analysis for HCV receptors was performed using real-time LightCycle PCR (Roche Diagnostics) as described previously [Ikeda *et al.*, 2005]. The primer pairs for CD81, SR-BI, CLDN1 and OCLN were reported previously [Nakamura *et al.*, 2011]. The primer pair NPC1L1 (5'-AGATCTTCTTCTCCGCCTCCA-3') and NPC1L1R (5'-TGCCAG-AGCCGGGTTAAC-3') was used for NPC1L1.

Statistical analysis. Luciferase activities were compared statistically between the various treatment groups using Student's *t*-test. *P*-values of <0.05 were considered statistically significant. The mean ± SD was determined from at least three independent experiments.

ACKNOWLEDGEMENTS

The authors would like to thank Masayo Takemoto, Takashi Nakamura and Keiko Takeshita for their technical assistance. This work was supported by Grants-in-Aid for Research on Hepatitis from the Ministry of Health, Labour, and Welfare of Japan.

REFERENCES

- Abe, K., Ikeda, M., Dansako, H., Naka, K. & Kato, N. (2007). Cell culture-adaptive NS3 mutations required for the robust replication of genome-length hepatitis C virus RNA. *Virus Res* 125, 88–97.
- Ariumi, Y., Kuroki, M., Abe, K., Dansako, H., Ikeda, M., Wakita, T. & Kato, N. (2007). DDX3 DEAD-box RNA helicase is required for hepatitis C virus RNA replication. *J Virol* 81, 13922–13926.
- Blight, K. J., McKeating, J. A. & Rice, C. M. (2002). Highly permissive cell lines for subgenomic and genomic hepatitis C virus RNA replication. *J Virol* 76, 13001–13014.
- Ikeda, M., Yi, M., Li, K. & Lemon, S. M. (2002). Selectable subgenomic and genome-length dicistronic RNAs derived from an infectious molecular clone of the HCV-N strain of hepatitis C virus replicate efficiently in cultured Huh7 cells. *J Virol* 76, 2997–3006.
- Ikeda, M., Abe, K., Dansako, H., Nakamura, T., Naka, K. & Kato, N. (2005). Efficient replication of a full-length hepatitis C virus genome, strain O, in cell culture, and development of a luciferase reporter system. *Biochem Biophys Res Commun* 329, 1350–1359.
- Kambara, H., Fukuhara, T., Shiokawa, M., Ono, C., Ohara, Y., Kamitani, W. & Matsuura, Y. (2012). Establishment of a novel permissive cell line for the propagation of hepatitis C virus by expression of microRNA miR122. *J Virol* 86, 1382–1393.
- Kato, N. (2001). Molecular virology of hepatitis C virus. *Acta Med Okayama* 55, 133–159.
- Kato, N., Hijikata, M., Ootsuyama, Y., Nakagawa, M., Ohkoshi, S., Sugimura, T. & Shimotohno, K. (1990). Molecular cloning of the human hepatitis C virus genome from Japanese patients with non-A, non-B hepatitis. *Proc Natl Acad Sci U S A* 87, 9524–9528.
- Kato, N., Sugiyama, K., Namba, K., Dansako, H., Nakamura, T., Takami, M., Naka, K., Nozaki, A. & Shimotohno, K. (2003). Establishment of a hepatitis C virus subgenomic replicon derived from human hepatocytes infected in vitro. *Biochem Biophys Res Commun* 306, 756–766.
- Kato, T., Choi, Y., Elmowalid, G., Sapp, R. K., Barth, H., Furusaka, A., Mishi, S., Wakita, T., Krawczynski, K. & Liang, T. J. (2008). Hepatitis C virus JFH-1 strain infection in chimpanzees is associated with low pathogenicity and emergence of an adaptive mutation. *Hepatology* 48, 732–740.
- Kato, N., Mori, K., Abe, K., Dansako, H., Kuroki, M., Ariumi, Y., Wakita, T. & Ikeda, M. (2009). Efficient replication systems for hepatitis C virus using a new human hepatoma cell line. *Virus Res* 146, 41–50.
- Koutsoudakis, G., Kaul, A., Steinmann, E., Kallis, S., Lohmann, V., Pietschmann, T. & Bartenschlager, R. (2006). Characterization of the early steps of hepatitis C virus infection by using luciferase reporter viruses. *J Virol* 80, 5308–5320.
- Lindenbach, B. D., Evans, M. J., Syder, A. J., Wölk, B., Tellinghuisen, T. L., Liu, C. C., Maruyama, T., Hynes, R. O., Burton, D. R. & other authors (2005). Complete replication of hepatitis C virus in cell culture. *Science* 309, 623–626.
- Lohmann, V., Körner, F., Koch, J., Herian, U., Theilmann, L. & Bartenschlager, R. (1999). Replication of subgenomic hepatitis C virus RNAs in a hepatoma cell line. *Science* 285, 110–113.
- Lohmann, V., Körner, F., Dobierzewska, A. & Bartenschlager, R. (2001). Mutations in hepatitis C virus RNAs conferring cell culture adaptation. *J Virol* 75, 1437–1449.
- Marcello, T., Grakoui, A., Barba-Spaeth, G., Machlin, E. S., Kotenko, S. V., MacDonald, M. R. & Rice, C. M. (2006). Interferons α and λ inhibit hepatitis C virus replication with distinct signal transduction and gene regulation kinetics. *Gastroenterology* 131, 1887–1898.
- Masaki, T., Suzuki, R., Murakami, K., Aizaki, H., Ishii, K., Murayama, A., Date, T., Matsuura, Y., Miyamura, T. & other authors (2008). Interaction of hepatitis C virus nonstructural protein 5A with core protein is critical for the production of infectious virus particles. *J Virol* 82, 7964–7976.
- Mori, K., Ikeda, M., Ariumi, Y. & Kato, N. (2010). Gene expression profile of Li23, a new human hepatoma cell line that enables robust hepatitis C virus replication: comparison with HuH-7 and other hepatic cell lines. *Hepatol Res* 40, 1248–1253.
- Mori, K., Ikeda, M., Ariumi, Y., Dansako, H., Wakita, T. & Kato, N. (2011). Mechanism of action of ribavirin in a novel hepatitis C virus replication cell system. *Virus Res* 157, 61–70.
- Narbus, C. M., Israelow, B., Sourisseau, M., Michta, M. L., Hopcraft, S. E., Zeiner, G. M. & Evans, M. J. (2011). HepG2 cells expressing microRNA miR-122 support the entire hepatitis C virus life cycle. *J Virol* 85, 12087–12092.
- Nakamura, M., Fujino, T., Yada, R., Aoyagi, Y., Yasutake, K., Kohjima, M., Fukuizumi, K., Yoshimoto, T., Harada, N. & other authors (2011). Expression profiles of genes associated with viral entry in HCV-infected human liver. *J Med Virol* 83, 921–927.
- Nishimura, G., Ikeda, M., Mori, K., Nakazawa, T., Ariumi, Y., Dansako, H. & Kato, N. (2009). Replicons from genotype 1b HCV-positive sera exhibit diverse sensitivities to anti-HCV reagents. *Antiviral Res* 82, 42–50.
- Pietschmann, T., Lohmann, V., Kaul, A., Krieger, N., Rinck, G., Rutter, G., Strand, D. & Bartenschlager, R. (2002). Persistent and transient replication of full-length hepatitis C virus genomes in cell culture. *J Virol* 76, 4008–4021.
- Pietschmann, T., Zayas, M., Meuleman, P., Long, G., Appel, N., Koutsoudakis, G., Kallis, S., Leroux-Roels, G., Lohmann, V. & Bartenschlager, R. (2009). Production of infectious genotype 1b virus particles in cell culture and impairment by replication enhancing mutations. *PLoS Pathog* 5, e1000475.
- Sainz, B., Jr, Barretto, N., Martin, D. N., Hiraga, N., Imamura, M., Hussain, S., Marsh, K. A., Yu, X., Chayama, K. & other authors (2012).

Identification of the Niemann–Pick C1-like 1 cholesterol absorption receptor as a new hepatitis C virus entry factor. *Nat Med* 18, 281–285.

Tanaka, T., Kato, N., Cho, M. J., Sugiyama, K. & Shimotohno, K. (1996). Structure of the 3' terminus of the hepatitis C virus genome. *J Virol* 70, 3307–3312.

Tellinghuisen, T. L., Foss, K. L. & Treadaway, J. (2008). Regulation of hepatitis C virion production via phosphorylation of the NS5A protein. *PLoS Pathog* 4, e1000032.

Ueda, Y., Mori, K., Ariumi, Y., Ikeda, M. & Kato, N. (2011). Plural assay systems derived from different cell lines and hepatitis C virus strains

are required for the objective evaluation of anti-hepatitis C virus reagents. *Biochem Biophys Res Commun* 409, 663–668.

Wakita, T., Pietschmann, T., Kato, T., Date, T., Miyamoto, M., Zhao, Z., Murthy, K., Habermann, A., Kräusslich, H. G. & other authors (2005). Production of infectious hepatitis C virus in tissue culture from a cloned viral genome. *Nat Med* 11, 791–796.

Zhong, J., Gastaminza, P., Cheng, G., Kapadia, S., Kato, T., Burton, D. R., Wieland, S. F., Uprichard, S. L., Wakita, T. & Chisari, F. V. (2005). Robust hepatitis C virus infection in vitro. *Proc Natl Acad Sci U S A* 102, 9294–9299.



Identification of host genes showing differential expression profiles with cell-based long-term replication of hepatitis C virus RNA

Hiroe Sejima, Kyoko Mori, Yasuo Ariumi¹, Masanori Ikeda, Nobuyuki Kato*

Department of Tumor Virology, Okayama University Graduate School of Medicine, Dentistry, and Pharmaceutical Sciences, 2-5-1, Shikata-cho, Okayama 700-8558, Japan

ARTICLE INFO

Article history:

Received 5 February 2012

Received in revised form 18 April 2012

Accepted 19 April 2012

Available online 1 May 2012

Keywords:

HCV

HCV RNA replication system

Li23 cells

Long-term RNA replication

Upregulated host genes

Downregulated host genes

ABSTRACT

Persistent hepatitis C virus (HCV) infection frequently causes hepatocellular carcinoma. However, the mechanisms of HCV-associated hepatocarcinogenesis and disease progression are unclear. Although the human hepatoma cell line, HuH-7, has been widely used as the only cell culture system for robust HCV replication, we recently developed new human hepatoma Li23 cell line-derived OL, OL8, OL11, and OL14 cells, in which genome-length HCV RNA (O strain of genotype 1b) efficiently replicates. OL, OL8, OL11, and OL14 cells were cultured for more than 2 years. We prepared cured cells from OL8 and OL11 cells by interferon- γ treatment. The cured cells were also cultured for more than 2 years. cDNA microarray and RT-PCR analyses were performed using total RNAs prepared from these cells. We first selected several hundred highly or moderately expressed probes, the expression levels of which were upregulated or downregulated at ratios of more than 2 or less than 0.5 in each set of compared cells (e.g., parent OL8 cells versus OL8 cells cultured for 2 years). From among these probes, we next selected those whose expression levels commonly changed during a 2-year culture of genome-length HCV RNA-replicating cells, but which did not change during a 2-year culture period in cured cells. We further examined the expression levels of the selected candidate genes by RT-PCR analysis using additional specimens from the cells cultured for 3.5 years. Reproducibility of the RT-PCR analysis using specimens from recultured cells was also confirmed. Finally, we identified 5 upregulated genes and 4 downregulated genes, the expression levels of which were irreversibly altered during 3.5-year replication of HCV RNA. These genes may play roles in the optimization of the environment in HCV RNA replication, or may play key roles in the progression of HCV-associated hepatic diseases.

© 2012 Elsevier B.V. All rights reserved.

1. Introduction

Hepatitis C virus (HCV) is a causative agent of chronic hepatitis, which progresses to liver cirrhosis and hepatocellular carcinoma (HCC) (Choo et al., 1989; Saito et al., 1990; Thomas, 2000). However,

the mechanisms of HCV-associated hepatocarcinogenesis and disease progression are still unclear. HCV is an enveloped virus with a positive single-stranded 9.6 kb RNA genome, which encodes a large polyprotein precursor of approximately 3000 amino acid residues. This polyprotein is cleaved by a combination of the host and viral proteases into at least 10 proteins in the following order: Core, envelope 1 (E1), E2, p7, nonstructural protein 2 (NS2), NS3, NS4A, NS4B, NS5A, and NS5B (Hijikata et al., 1991, 1993; Kato et al., 1990).

The initial development of a cell culture-based replicon system (Lohmann et al., 1999) and a genome-length HCV RNA-replication system (Ikeda et al., 2002) using genotype 1b strains enabled the rapid progression of investigations into the mechanisms underlying HCV replication (Bartenschlager, 2005; Lindenbach and Rice, 2005). Furthermore, these RNA replication systems have been improved such that they have become suitable for the screening of anti-HCV reagents by the introduction of reporter genes such as luciferase (Ikeda et al., 2005; Krieger et al., 2001). Moreover, in 2005, an efficient virus production system using the JFH1 genotype 2a strain was developed using human hepatoma cell line HuH-7-derived cells (Wakita et al., 2005). However, to date, HuH-7-derived cells are used as the only cell culture

Abbreviations: HCV, hepatitis C virus; HCC, hepatocellular carcinoma; E1, envelope 1; EGF, epidermal growth factor; RT-PCR, reverse transcription-polymerase chain reaction; IFN, interferon; ACSM3, acyl-CoA synthetase medium-chain family member 3; ANGPT1, angiopoietin 1; CDKN2C, cyclin-dependent kinase inhibitor 2C; PLA1A, phospholipase A1 member A; SEL1L3, Sel-1 suppressor of lin-12-like 3; SLC39A4, solute carrier family 39 member 4; TBC1D4, TBC1 domain family member 4; WISP3, WNT1 inducible signaling pathway protein 3; ANXA1, annexin A1; AREG, amphiregulin; BASP1, brain abundant, membrane attached signal protein 1; CIDEC, cell death activator CIDE-3; CPB2, carboxypeptidase B2; HSPA6, heat-shock 70 kDa protein B'; PI3, peptidase inhibitor 3; SLC1A3, solute carrier family 1 member 3; THSD4, thrombospondin type-1 domain-containing protein 4; ICAM-1, intercellular adhesion molecule-1; ALXR, ANXA1 receptor.

* Corresponding author. Tel.: +81 86 235 7385; fax: +81 86 235 7392.

E-mail address: nkato@md.okayama-u.ac.jp (N. Kato).

¹ Present address: Center For AIDS Research, Kumamoto University, Kumamoto 860-0811, Japan.