

WT mice (1.5-fold), even though the serum ALT level in the ND-fed KO mice was modestly higher than that in the ND-fed WT mice.

We next conducted histological analyses of liver sections from the mice. H&E staining revealed that the livers from KO mice fed the HFD possessed not only steatotic areas but also scattered inflammatory foci composed of gathering nonparenchymal cells (Fig. 2b). Although inflammatory foci were also observed in the livers from KO mice fed the ND, a larger number of foci could clearly be seen in KO mice fed the HFD than in KO mice fed the ND (Fig. 2c). In contrast, WT mice fed the HFD, as well as those given the ND, showed few inflammatory foci. Taken together, these results indicated that the HFD augmented liver inflammation in KO mice but not in WT mice.

The HFD enhanced hepatic inflammation-related gene expression in the absence of iNKT cells

To understand the underlying mechanisms of the hepatic inflammation induced by the HFD in the absence of iNKT cells, we first examined the levels of several cytokines and chemokines in the livers from mice at week 5 after they had been started on the diets. Real-time RT-PCR analyses revealed that the messenger RNA expression of tumor necrosis factor (TNF)- α , IFN- γ , IL-10, chemokine (C-C motif) ligand (CCL) 2 and 4, and chemokine (C-X-C motif) ligand (CXCL) 9 and 10 were remarkably upregulated by the HFD, compared with the ND, in KO but not in WT mice (Fig. 3), although these values in KO mice fed the ND were modestly higher than those in WT mice fed the ND. In contrast, the messenger RNA expression of IL-4, -5, and -13 did not show any detectable levels in the livers from both WT and KO mice fed either the ND or HFD.

The HFD altered the proportions of subpopulations in liver mononuclear cells from KO mice, but not in those from WT mice

We next examined the phenotype of mononuclear cells in livers from mice at week 5 of feeding. Flow cytometric analyses demonstrated that the proportion of CD4+ TCR β + CD4 T cells was lower in KO mice fed the ND than in WT mice fed the ND (Fig. 4a), which might have resulted from a lack of iNKT cells partly composed of CD4+ cells [5]. The proportion of CD49b+ TCR β - NK cells or CD8+ TCR β + CD8 T cells was higher in KO mice fed the ND than in WT mice fed the ND. The HFD did not lead to any significant changes in the proportion of hepatic CD4 T, CD8 T, NK, or iNKT cells in WT mice. In contrast, the HFD induced significant increases in the proportion of hepatic CD4 T cells and CD8 T cells, but not of NK cells, in KO mice. We then examined the distribution of these cells




Fig. 3 Inflammatory cytokine and chemokine gene expression in the liver. Liver tissues were obtained from wild-type (WT) and α 18-deficient (KO) mice fed either a normal diet (ND) or a high-fat diet (HFD) for 5 weeks. Liver RNA levels of the indicated genes and beta-actin as a control were analyzed using real-time reverse transcription polymerase chain reaction (RT-PCR). Data are shown as the fold increase of HFD-fed WT, ND-fed KO, or HFD-fed KO compared with ND-fed WT mice, with means \pm SEM from five mice in each group. Data are representative of more than four independent experiments. TNF- α Tumor necrosis factor alpha, IFN- γ interferon gamma, IL interleukin, CCL chemokine (C-C motif) ligand, CXCL chemokine (C-X-C motif) ligand. §, not detected. * P < 0.05 versus KO fed ND

in the liver. Immunohistochemical examination revealed that CD4+ cells and CD8+ cells formed foci surrounding hepatocytes in the livers of KO mice (Fig. 4b), which partly corresponded to the inflammatory foci observed in the




Fig. 4 Involvement of CD4 T and/or CD8 T cells in liver inflammation. Livers were obtained from wild-type (WT) and $\alpha 18$ -deficient (KO) mice fed either a normal diet (ND) or a high-fat diet (HFD) for 5 weeks. **a** Prepared mononuclear cells from the livers were stained with cell markers indicated in “Materials and methods”. Proportions of the indicated cell population were analyzed by flow cytometry. Data shown are means \pm SEM from five mice in each group. Data are representative of more than four independent experiments. §, not detected. * $P < 0.05$ versus KO fed ND. **b** Liver sections were analyzed by immunohistochemical staining for CD4- or CD8-positive cells. Representative images are shown ($\times 200$)

H&E-stained liver sections of KO mice. WT mice fed either the ND or HFD did not display such foci consisting of stained cells in the livers. Collectively, these results suggested that CD4 T and/or CD8 T cells played a role in the HFD-enhanced liver inflammation in KO mice.

The HFD led to the development of liver fibrosis in the absence of iNKT cells

Persistent hepatic inflammation causes fibrotic changes in the liver [25]. To investigate whether inflammation with steatosis due to the HFD in KO mice would induce fibrosis in the liver, we fed the ND or HFD to WT and KO mice for a longer period of 15 weeks. H&E staining and Oil-red-O staining showed that KO mice fed the HFD possessed the inflammatory foci, together with lipid retention in the liver, at week 15, as well as showing these findings at week 5,




Fig. 5 Liver fibrosis following inflammation in the absence of iNKT cells. Livers were obtained from wild-type (WT) and $\alpha 18$ -deficient (KO) mice fed either a normal diet (ND) or a high-fat diet (HFD) for 15 weeks. **a** Liver tissues were stained with Sirius-Red to assess liver fibrosis. Representative images are shown ($\times 200$). **b** The stained areas were evaluated in five different fields per section. Data shown are means \pm SEM from five mice in each group. Data are representative of more than two independent experiments. * $P < 0.05$ versus KO fed ND

and that WT mice fed the HFD showed lipid retention, but few inflammatory foci, in the liver at week 15 (data not shown). Sirius-Red staining revealed clear fibrosis in the livers from KO mice fed the HFD and also in the livers from KO mice fed the ND, but to a much lesser extent (Fig. 5a). In contrast, the staining showed no obvious fibrosis in the livers from WT mice fed the HFD or in those given the ND. Quantitative analyses to evaluate the stained areas also showed that the HFD-fed KO mice possessed significantly greater areas of hepatic fibrosis than the ND-fed KO mice, while WT mice fed either the ND or HFD had few fibrotic areas (Fig. 5b). Taken together, these results indicated that the HFD led to the development of liver fibrosis accompanied by steatohepatitis in KO mice.

Discussion

An increasing amount of evidence suggests that iNKT cells play a role in immune responses in the liver [12], although

the exact implication of that role is controversial. iNKT cells, for instance, have been reported to play a critical role in animal models of liver injury induced by concanavalin A, α -galactosylceramide, or salmonella infection [26–28], suggesting a proinflammatory role of these cells. On the other hand, iNKT cells have been very recently implicated in the suppression of liver damage in a mouse model of cholestasis [29], suggesting an anti-inflammatory role of these cells. The present study, using iNKT cell-deficient mice fed an HFD, demonstrated that the HFD led to the development of steatohepatitis with fibrosis in the absence of iNKT cells, while the HFD led to steatosis but not steatohepatitis in the presence of these cells. This suggests that iNKT cells play a critical role in suppressing the development of inflammation and fibrosis in the steatotic liver.

Our real-time RT-PCR analyses demonstrated that CCL2, CCL4, CXCL9, and CXCL10 were remarkably upregulated by the HFD in KO mice but not in WT mice (Fig. 3). CCL2 or CCL4 has the ability to attract predominantly Th1 cells via chemokine (C–C motif) receptor 2 or 5, respectively. CXCL9 and CXCL10 also attract predominantly Th1 cells via chemokine (C–X–C motif) receptor 3 [30, 31]. Indeed, Th1 cytokines such as TNF- α and IFN- γ were remarkably upregulated by the HFD in KO mice but not in WT mice. Although IL-10, which is one of the anti-inflammatory cytokines, was also upregulated by the HFD in KO mice but not in WT mice, the upregulation of IL-10 may have counteracted the upregulation of the proinflammatory Th1 cytokines TNF- α and IFN- γ . Our flow cytometric analyses and immunohistochemical analyses showed that the proportions of CD4 T and CD8 T cells were increased (Fig. 4a) and that these cells also accumulated to form foci (Fig. 4b) in the livers of KO mice fed the HFD. Bigorgne et al. [32] reported that HFD-induced obesity in leptin-deficient ob/ob mice rendered hepatic mononuclear cells, particularly CD4 T and CD8 T cells, sensitive to chemokines such as CXCL12 and CXCL13, which attract T cells, suggesting an important role of chemokines in liver inflammation with steatosis. Although the sources of the chemokines upregulated in our model were not clear, these chemokines presumably play an important role in the infiltration of proinflammatory cells in the liver of the KO mice fed the HFD. iNKT cells suppress the production of these chemokines directly or indirectly; thus, they may prevent steatohepatitis induced by an HFD.

The liver can be anatomically exposed to gut-derived contents, such as food antigens and bacterial products, via the portal vein [33, 34]. Once these entities flow into the liver, they can activate a variety of cells in the liver, which may be associated with certain types of liver disease [33, 34]. Gut-derived food-antigens can activate T cells [33] and gut-derived bacterial products can stimulate all resident

cells in the liver, such as hepatocytes, Kupffer cells, stellate cells, and dendritic cells, via toll-like receptors [33–36]. Moreover, fat itself, particularly saturated fatty acids, stimulates an immune response in the liver [37, 38]. On the other hand, the liver is an immune-tolerogenic organ, in which immune-suppressive cells may play a critical role to keep this organ immunologically silent [33]. The present study demonstrated that liver inflammation was greatly exacerbated—where CD4 T and/or CD8 T cells infiltrated to form foci surrounding damaged hepatocytes—by an HFD in the absence of iNKT cells. This suggests a suppressive role of iNKT cells in the development of liver inflammation with steatosis. Thus, iNKT cells may play an important role in keeping the liver immunologically silent, and the absence of iNKT cells together with steatosis may elicit a break of hepatic immune tolerance, resulting in the activation of CD4 T and/or CD8 T cells to provoke liver inflammation. Consistent with this speculation is the observation that the absence of iNKT cells, even without steatosis, caused modest liver inflammation.

In conclusion, iNKT cells suppress liver inflammation progressing to fibrosis that is exacerbated by HFD-induced steatosis, thus contributing to the maintenance of immune homeostasis in the liver. This study has shed some light on iNKT cells as immunoregulatory cells and their key role in the pathogenesis of NAFLD.

Acknowledgments The authors thank Drs. Masaru Taniguchi and Ken-ichiro Seino for providing α 18-deficient BALB/c mice. This work was supported by a Grant-in-Aid for Scientific Research (to T. Takehara) and Global Centers of Excellence Program (to T. Miyagi) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

1. Angulo P. Nonalcoholic fatty liver disease. *N Engl J Med*. 2002;346:1221–31.
2. Clark JM, Brancati FL, Diehl AM. Nonalcoholic fatty liver disease. *Gastroenterology*. 2002;122:1649–57.
3. Maher JJ, Leon P, Ryan JC. Beyond insulin resistance: Innate immunity in nonalcoholic steatohepatitis. *Hepatology*. 2008;48:670–8.
4. Tilg H, Diehl AM. Cytokines in alcoholic and nonalcoholic steatohepatitis. *N Engl J Med*. 2000;343:1467–76.
5. Bendelac A, Savage PB, Teyton L. The biology of NKT cells. *Annu Rev Immunol*. 2007;25:297–336.
6. Godfrey DI, Kronenberg M. Going both ways: immune regulation via CD1d-dependent NKT cells. *J Clin Invest*. 2004;114:1379–88.
7. Trobonjaca Z, Leithäuser F, Möller P, Schirmbeck R, Reimann J. Activating immunity in the liver. I. Liver dendritic cells (but not hepatocytes) are potent activators of IFN-gamma release by liver NKT cells. *J Immunol*. 2001;167:1413–22.
8. Winau F, Hegasy G, Weiskirchen R, Weber S, Cassan C, Sieling PA, et al. Ito cells are liver-resident antigen-presenting cells for activating T cell responses. *Immunity*. 2007;26:117–29.

9. Matsuda JL, Mallevaey T, Scott-Browne J, Gapin L. CD1d-restricted iNKT cells, the 'Swiss-Army knife' of the immune system. *Curr Opin Immunol*. 2008;20:358–68.
10. Matsuda JL, Naidenko OV, Gapin L, Nakayama T, Taniguchi M, Wang CR, et al. Tracking the response of natural killer T cells to a glycolipid antigen using CD1d tetramers. *J Exp Med*. 2000;192:741–54.
11. Miyagi T, Takehara T, Tatsumi T, Kanto T, Suzuki T, Jinushi M, et al. CD1d-mediated stimulation of natural killer T cells selectively activates hepatic natural killer cells to eliminate experimentally disseminated hepatoma cells in murine liver. *Int J Cancer*. 2003;106:81–9.
12. Exley MA, Koziel MJ. To be or not to be NKT: natural killer T cells in the liver. *Hepatology*. 2004;40:1033–40.
13. Norris S, Doherty DG, Collins C, McEntee G, Traynor O, Hegarty JE, et al. Natural T cells in the human liver: cytotoxic lymphocytes with dual T cell and natural killer cell phenotype and function are phenotypically heterogeneous and include Valpha24-JalphaQ and gammadelta T cell receptor bearing cells. *Hum Immunol*. 1999;60:20–31.
14. Guebre-Xabier M, Yang S, Lin HZ, Schwenk R, Krzych U, Diehl AM. Altered hepatic lymphocyte subpopulations in obesity-related murine fatty livers: potential mechanism for sensitization to liver damage. *Hepatology*. 2000;31:633–40.
15. Li ZP, Soloski MJ, Diehl AM. Dietary factors alter hepatic innate immune system in mice with nonalcoholic fatty liver disease. *Hepatology*. 2005;42:880–5.
16. Ma X, Hua J, Li Z. Probiotics improve high fat diet-induced hepatic steatosis and insulin resistance by increasing hepatic NKT cells. *J Hepatol*. 2008;49:821–30.
17. Miyazaki Y, Iwabuchi K, Iwata D, Miyazaki A, Kon Y, Niino M, et al. Effect of high fat diet on NKT cell function and NKT cell-mediated regulation of Th1 responses. *Scand J Immunol*. 2008;67:230–7.
18. Xu CF, Yu CH, Li YM, Xu L, Du J, Shen Z. Association of the frequency of peripheral natural killer T cells with nonalcoholic fatty liver disease. *World J Gastroenterol*. 2007;13:4504–8.
19. Tajiri K, Shimizu Y, Tsuneyama K, Sugiyama T. Role of liver-infiltrating CD3+ CD56+ natural killer T cells in the pathogenesis of nonalcoholic fatty liver disease. *Eur J Gastroenterol Hepatol*. 2009;21:673–80.
20. Elinav E, Pappo O, Sklair-Levy M, Margalit M, Shibolet O, Gomori M, et al. Adoptive transfer of regulatory NKT lymphocytes ameliorates non-alcoholic steatohepatitis and glucose intolerance in ob/ob mice and is associated with intrahepatic CD8 trapping. *J Pathol*. 2006;209:121–8.
21. Cui J, Shin T, Kawano T, Sato H, Kondo E, Taura I, et al. Requirement for Valpha14 NKT cells in IL-12-mediated rejection of tumors. *Science*. 1997;278:1623–6.
22. Harada M, Magara-Koyanagi K, Watarai H, Nagata Y, Ishii Y, Kojo S, et al. IL-21-induced Bepsilon cell apoptosis mediated by natural killer T cells suppresses IgE responses. *J Exp Med*. 2006;203:2929–37.
23. Miyagi T, Takehara T, Tatsumi T, Suzuki T, Jinushi M, Kanazawa Y, et al. Concanavalin A injection activates intrahepatic innate immune cells to provoke an antitumor effect in murine liver. *Hepatology*. 2004;40:1190–6.
24. Kamada Y, Matsumoto H, Tamura S, Fukushima J, Kiso S, Fukui K, et al. Hypoadiponectinemia accelerates hepatic tumor formation in a nonalcoholic steatohepatitis mouse model. *J Hepatol*. 2007;47:556–64.
25. Friedman SL. Mechanisms of hepatic fibrogenesis. *Gastroenterology*. 2008;134:1655–69.
26. Ishigami M, Nishimura H, Naiki Y, Yoshioka K, Kawano T, Tanaka Y, et al. The roles of intrahepatic Valpha14(+) NK1.1(+) T cells for liver injury induced by Salmonella infection in mice. *Hepatology*. 1999;29:1799–808.
27. Kaneko Y, Harada M, Kawano T, Yamashita M, Shibata Y, Gejyo F, et al. Augmentation of Valpha14 NKT cell-mediated cytotoxicity by interleukin 4 in an autocrine mechanism resulting in the development of concanavalin A-induced hepatitis. *J Exp Med*. 2000;191:105–14.
28. Osman Y, Kawamura T, Naito T, Takeda K, Van Kaer L, Okumura K, et al. Activation of hepatic NKT cells and subsequent liver injury following administration of alpha-galactosylceramide. *Eur J Immunol*. 2000;30:1919–28.
29. Wintermeyer P, Cheng CW, Gehring S, Hoffman BL, Holub M, Brossay L, et al. Invariant natural killer T cells suppress the neutrophil inflammatory response in a mouse model of cholestatic liver damage. *Gastroenterology*. 2009;136:1048–59.
30. Mantovani A, Sica A, Sozzani S, Allavena P, Vecchi A, Locati M. The chemokine system in diverse forms of macrophage activation and polarization. *Trends Immunol*. 2004;25:677–86.
31. Moser B, Wolf M, Walz A, Loetscher P. Chemokines: multiple levels of leukocyte migration control. *Trends Immunol*. 2004;25:75–84.
32. Bigorgne AE, Bouchet-Delbos L, Naveau S, Dagher I, Prévot S, Durand-Gasselin I, et al. Obesity-induced lymphocyte hyperresponsiveness to chemokines: a new mechanism of fatty liver inflammation in obese mice. *Gastroenterology*. 2008;134:1459–69.
33. Crispe IN. Hepatic T cells and liver tolerance. *Nat Rev Immunol*. 2003;3:51–62.
34. Gao B, Jeong WI, Tian Z. Liver: an organ with predominant innate immunity. *Hepatology*. 2008;47:729–36.
35. Paik YH, Schwabe RF, Bataller R, Russo MP, Jobin C, Brenner DA. Toll-like receptor 4 mediates inflammatory signaling by bacterial lipopolysaccharide in human hepatic stellate cells. *Hepatology*. 2003;37:1043–55.
36. Seki E, De Minicis S, Osterreicher CH, Kluwe J, Osawa Y, Brenner DA, et al. TLR4 enhances TGF-beta signaling and hepatic fibrosis. *Nat Med*. 2007;13:1324–32.
37. Lee JY, Hwang DH. The modulation of inflammatory gene expression by lipids: mediation through Toll-like receptors. *Mol Cells*. 2006;21:174–85.
38. Lee JY, Zhao L, Hwang DH. Modulation of pattern recognition receptor-mediated inflammation and risk of chronic diseases by dietary fatty acids. *Nutr Rev*. 2010;68:38–61.

Altered interferon- α -signaling in natural killer cells from patients with chronic hepatitis C virus infection[☆]

Takuya Miyagi^{1,2}, Tetsuo Takehara¹, Kumiko Nishio¹, Satoshi Shimizu¹, Keisuke Kohga¹, Wei Li¹, Tomohide Tatsumi¹, Naoki Hiramatsu¹, Tatsuya Kanto¹, Norio Hayashi^{1,2,*}

¹Department of Gastroenterology and Hepatology, Osaka University Graduate School of Medicine, Suita, Osaka, Japan; ²Global Centers of Excellence Program, Frontier Biomedical Science Underlying Organelle Network Biology, Osaka University, Suita, Osaka, Japan

Background & Aims: Natural killer (NK) cells play an important role in the immune response against virus infection. Interferon (IFN)- α , an essential component in therapy against hepatitis C virus (HCV) infection, regulates NK cell function. However, it remains obscure how chronic HCV infection (CHC) modifies intracellular IFN- α signaling in NK cells. We investigated IFN- α signaling in NK cells in patients with CHC.

Methods: Peripheral blood mononuclear cells were obtained from patients with CHC and healthy subjects (HS) as controls.

Results: The expression level of signal transducer and activator of transcription (STAT) 1, a key molecule of IFN- α signaling, was clearly higher in NK cells from the CHC patients than in those from HS. The phosphorylation level of STAT1 with IFN- α stimulation was significantly greater in NK cells from the CHC patients than in those from the HS, while that of STAT4 was significantly less. These phosphorylation levels of STAT1 and STAT4 positively and negatively correlated with the STAT1 level in NK cells, respectively. The IFN- α induced messenger RNA level of the suppressor of cytokine signaling 1, which is a downstream gene of phosphorylated-STAT1, was clearly greater in NK cells from the CHC patients than in those from the HS, while that of IFN- γ , which is a downstream gene of phosphorylated-STAT4, was clearly lower.

Conclusions: These results indicate altered IFN- α signaling in NK cells in CHC patients, suggesting that this alteration is associated with the persistence of HCV infection and resistance to IFN- α therapy.

Keywords: Natural killer cells; Interferon; Hepatitis C virus; Signal transducer and activator of transcription.

Received 28 October 2009; received in revised form 24 March 2010; accepted 28 March 2010; available online 25 May 2010

* Funding: The authors who have taken part in this study declared that they do not have anything to disclose regarding funding or conflict of interest with respect to this manuscript.

Corresponding author. Address: Department of Gastroenterology and Hepatology, Osaka University Graduate School of Medicine, 2-2 Yamada-oka, Suita, Osaka 565-0871, Japan. Tel.: +81 6 6879 3621; fax: +81 6 6879 3629.

E-mail address: hayashin@gh.med.osaka-u.ac.jp (N. Hayashi).

Abbreviations: NK, natural killer; IFN, interferon; STAT, signal transducer and activator of transcription; HCV, hepatitis C virus; CHC, chronic hepatitis C virus infection; HS, healthy subject; SOCS, suppressor of cytokine signaling; mRNA, messenger RNA; PBMC, peripheral blood mononuclear cell; IL, interleukin; pSTAT, phosphorylated-signal transducer and activator of transcription; RT-PCR, reverse transcription polymerase chain reaction; NKT, natural killer T; ISG, interferon stimulated gene.

© 2010 European Association for the Study of the Liver. Published by Elsevier B.V. All rights reserved.

Introduction

Natural killer (NK) cells play an important role in innate immune responses against a variety of viral infections by directly killing infected cells with cytotoxic molecules such as perforin and granzyme [1]. The cells also have a great ability to secrete a key cytokine, interferon (IFN)- γ , which activates subsequent adaptive immune responses as well as inhibits viral replication [1,2]. Another major component in innate immune responses during viral infections is IFN- α , which is the most abundant cytokine released during viral infections [3]. In addition to its anti-viral effects, IFN- α activates NK cells to induce IFN- γ production via activation of the signal transducer and activator of transcription (STAT) 4, as well as its cytotoxic ability via activation of STAT1 [4–7].

Hepatitis C virus (HCV) causes persistent infection in more than 70% of infected patients. Whereas some of the patients show a carrier-like state, most develop chronic liver diseases, including chronic hepatitis, liver cirrhosis, and hepatocellular carcinoma, which is why HCV infection is a worldwide health problem [8]. The administration of IFN- α is a well-established anti-viral therapy for HCV infection. More than 90% of patients with acute HCV infection respond to IFN- α based therapy, while only around 50% of patients with chronic HCV infection (CHC) do [9–12], suggesting a mechanism by which persistent HCV infection leads to resistance to IFN- α based therapy. NK cell number has been demonstrated to decrease in patients with CHC, while it is controversial whether NK cell functions are impaired in patients with CHC [13–15]. It thus remains unclear whether perturbation of NK cells is involved in the persistence of CHC as well as resistance to the therapy [13–15].

In the present study, we investigated how chronic HCV infection modifies intracellular IFN- α signaling in NK cells. The expression level of total STAT1, a key molecule of IFN- α signaling, was clearly higher in NK cells from patients with CHC than in those from healthy subjects (HS). Phosphorylation of STAT1, resulting in induction of the suppressor of cytokine signaling (SOCS) 1 messenger RNA (mRNA) expression in response to IFN- α was clearly greater in NK cells from the CHC patients than



in those from the HS. On the other hand, the phosphorylation of STAT4, resulting in induction of IFN- γ mRNA expression in response to IFN- α , was clearly less in NK cells from the CHC patients than in those from the HS. NK cell degranulation was significantly enhanced in response to IFN- α in the HS, but not in the CHC patients. These findings suggest altered IFN- α signaling in NK cells in patients with CHC. The alteration of IFN- α signaling might be associated with the persistence of chronic HCV infection and resistance to IFN- α therapy.

Materials and methods

Subjects

Twenty-six patients with CHC (HCV RNA genotype 1) and 26 healthy volunteers were enrolled in this study. The profile of these subjects is shown in the Supplementary data and Supplementary Table 1. The study was approved by the ethical committee of Osaka University Hospital.

Isolation of peripheral blood mononuclear cell (PBMC) populations

PBMCs were isolated from fresh heparinized peripheral blood by Ficoll-Hypaque density gradient centrifugation as described [16].

In vitro stimulation of cells

Prepared cells were unstimulated or stimulated with either natural human IFN- α , recombinant human IFN- γ or recombinant human interleukin (IL)-12. The details are provided in the Supplementary data.

Cell lysates and Western blot analysis

Prepared cells were lysed as described [17]. A 25 μ g sample of protein was separated on 10% SDS polyacrylamide gels and transferred onto PVDF membrane. Monoclonal anti-STAT1 antibody (1/Stat1) was purchased from BD Biosciences (San Jose, CA, USA). Polyclonal anti- β -actin antibody from Abcam (Cambridge, MA, USA) was used as the loading control. Detection of immunolabeled proteins was performed as described [17].

Flow cytometric analysis

The staining of prepared cells was performed as described [16,18]. Briefly, cells were stained with fluorescein isothiocyanate-conjugated anti-CD3 (UCHT1) and biotin-conjugated anti-CD56 antibody (B159), fixed and permeabilized with Cytofix/Cytoperm (BD Biosciences) and cold pure methanol, and then stained with phycoerythrin-conjugated anti-phosphorylated-STAT (pSTAT) 4 (pY693) (38/p-Stat4) and Alexa Fluor[®] 647-conjugated anti-pSTAT1 (pY701) antibody (4a), or phycoerythrin-conjugated anti-STAT1 (1/Stat1) antibody alone, or the corresponding isotype control, followed by staining with peridinin chlorophyll protein-conjugated streptavidin (BD Biosciences). All antibodies were purchased from BD Biosciences. The stained cells were analyzed with a FACScan (Becton Dickinson, Mountain View, CA, USA), and the data were processed using the FlowJo program (Tree Star Inc., Ashland, OR, USA).

NK cell enrichment from PBMCs

To obtain pure populations of CD56⁺ CD3⁻ NK cells from PBMCs, NK cells were negatively isolated by magnetic cell sorting with a human NK cell isolation kit (Miltenyi Biotec, Gladbach, Germany). The purity of the isolated population was confirmed using FACS analysis and was more than 90%.

RNA isolation and analysis

Total RNA isolation and the real-time reverse transcription polymerase chain reaction (RT-PCR) analysis are presented in detail in the Supplementary data, Supplementary Fig. 1, and Supplementary Table 2.

NK cell degranulation assay

NK cell degranulation was assessed as described [19], with minor modifications. Details and representative data are provided in the Supplementary data and Supplementary Fig. 2.

Statistical analysis

The statistical significance of differences between the patient and control groups or that of changes due to IFN- α stimulation in the NK cell degranulation assay was determined by applying unpaired or paired Student's *t*-test, respectively. Correlations were assessed using the Pearson product-moment correlation coefficient. The statistical significance was defined as $p < 0.05$.

Results

NK cells from CHC patients showed a higher level of STAT1 expression than those from HS

Murine NK cells have been reported to exhibit a lower level of STAT1 expression than non-NK cells [17,18]. Also, the increase of STAT1 expression level in murine NK cells has been observed in viral infection [18]. We examined whether similar findings could be observed for human NK cells. Western blot analyses revealed that human NK cells from representative HS have a clearly lower level of STAT1 expression than non-NK cells (Fig. 1A). We then examined whether chronic HCV infection affected the STAT1 expression level in NK cells. It was clearly higher in NK cells from the CHC patients than in those from the HS (Fig. 1B).

We next examined STAT1 expression level within individual cells by flow cytometry. Consistent with the results of Western blot analyses, flow cytometric analyses demonstrated that NK cells from a representative healthy subject had a lower level of STAT1 expression than non-NK cells such as T cells or natural killer T (NKT) cells (Fig. 2A). NK cells from a representative CHC patient displayed a clearly higher level of STAT1 expression than




Fig. 1. STAT1 expression in NK cells from patients with chronic HCV infection. STAT1 protein levels were evaluated by Western blot analyses with β -actin measurement as a loading control. PBMCs were obtained from patients with chronic HCV infection (CHC) and healthy subjects (HS). NK cells and non-NK cells were purified from those cells. (A) STAT1 protein levels in NK cells and non-NK cells from two representative HS are shown. (B) STAT1 protein levels in NK cells from three other representative HS and three representative patients are shown.

Research Article




Fig. 2. STAT1 expression level in NK, T or NKT cells from patients with chronic HCV infection. STAT1 protein level was evaluated by flow cytometry, electronically gating on CD56⁺ CD3⁻ NK cells, CD56⁻ CD3⁺ T cells, and CD56⁺ CD3⁺ NKT cells. PBMCs were derived from patients with chronic HCV infection (CHC) and healthy subjects (HS). (A) Representative histograms from a patient and an HS are shown. Dotted lines show staining with the isotype control. Solid lines with shaded areas show staining with the antibody. Numbers are percentages of positive cells (positive cell rate) determined based on the isotype control staining. (B) Comparison of STAT1 expression level between the patients with CHC ($n = 11$) and HS ($n = 11$) are shown as positive cell rates. Each circle represents data for an individual. Horizontal bars represent means. * $p < 0.005$ vs. HS.

those from the healthy subject. Fig. 2B summarizes the profile of the intracellular STAT1 expression level of NK, T, and NKT cells. The intracellular STAT1 expression level of NK cells from the CHC patients was significantly higher than that from the HS, while that of T cells or NKT cells did not show any significant difference.

Altered activation of STAT1/4 occurred in response to IFN- α in NK cells from CHC patients

Activation of STAT1/4 in response to IFN- α in murine NK cells has been reported to shift from pSTAT4 dominant to pSTAT1 dominant as the intracellular STAT1 expression level increases [18]. This led us to examine the activation of STAT1/4 in response to IFN- α in human NK cells using samples from CHC patients and HS. Although IFN- α can phosphorylate both STAT1 and STAT4, IFN- γ can phosphorylate STAT1 and IL-12 can phosphorylate STAT4 in NK cells [1,7]. We examined the phosphorylation level of STAT1/4 in response to IFN- α , compared with IFN- γ or IL-12, in NK cells from the subjects by flow cytometry. It was found that IFN- α phosphorylated STAT1 more strongly in NK cells from the representative CHC patient than in those from the representative

HS, while IFN- γ did not (Fig. 3A). On the other hand, IFN- α phosphorylated STAT4 more weakly in NK cells from the CHC patient than in those from the HS, while IL-12 did not. Fig. 3B summarizes the profile of STAT1/4 phosphorylation level in NK cells in response to IFN- α , IFN- γ or IL-12. IFN- α , but not IFN- γ , phosphorylated STAT1 significantly more strongly in NK cells from the CHC patients than in those from the HS. IFN- α , but not IL-12, phosphorylated STAT4 significantly more weakly in NK cells from the CHC patients than in those from the HS. These results suggested altered signaling of IFN- α , but not of IFN- γ or of IL-12, in NK cells of patients with CHC.

We then examined the relationship between STAT1 expression level and STAT1/4 phosphorylation level in response to IFN- α in NK cells from the CHC patients. The phosphorylation level of STAT1 in response to IFN- α correlated significantly and positively with the STAT1 expression level in NK cells ($R^2 = 0.67$, $p < 0.003$), while that of STAT4 correlated significantly and negatively ($R^2 = 0.49$, $p < 0.02$) (Fig. 4). These results suggested that, as in murine NK cells, the activation of STAT1/4 in response to IFN- α in human NK cells shifts from a preference for pSTAT4 to one for pSTAT1 as intracellular STAT1 expression level increases.

Altered induction of interferon stimulated gene (ISG) expression occurred in response to IFN- α in NK cells

We examined how the altered activation of STAT1/4 in response to IFN- α in NK cells affected the induction of downstream gene expression. Real-time RT-PCR analyses revealed that the induction level of SOCS1 mRNA expression with IFN- α stimulation, which is a downstream gene of pSTAT1, was clearly greater in NK cells isolated from the CHC patients than in those from the HS, and that the induction level of IFN- γ mRNA expression with IFN- α stimulation, which is a downstream gene of pSTAT4, was clearly lower (Fig. 5). On the other hand, the mRNA induction level of perforin or granzyme B, which is a cytotoxic molecule induced by IFN- α via pSTAT1, was not greater but modestly lower, suggesting negative regulation by the large induction of SOCS1, which is a negative regulator of the pSTAT1 pathway [20].

Altered activation of NK cells occurred in response to IFN- α

To examine how the altered IFN- α signaling in NK cells affected the activation of NK cells in response to IFN- α , we evaluated the NK cell degranulation ability in response to IFN- α . NK cell degranulation assay showed that CD107a expression, as a marker of degranulation, in the presence of K562 cells was significantly up-regulated in response to IFN- α in NK cells from HS, but not in those from CHC patients (Fig. 6), suggesting altered NK cell activation in response to IFN- α in CHC patients.

Discussion

Both IFN- α and IFN- γ have been observed in sera from patients with CHC [21,22]. Their production may be induced by a host response to HCV within the liver, which would make these IFNs detectable in the systemic circulation of patients with CHC. The present study has shown that NK cells from patients with CHC display higher levels of STAT1 expression compared to those from HS (Fig. 1B and Fig. 2). STAT1 itself is one of the ISGs, whose




Fig. 3. Altered activation of STAT1/4 in response to IFN- α , but not to IFN- γ or IL-12 in NK cells from patients with chronic HCV infection. pSTAT1 and pSTAT4 protein levels were evaluated by flow cytometry with isotype control staining. PBMCs were derived from patients with chronic HCV infection (CHC) and healthy subjects (HS). (A) Prepared PBMCs were unstimulated or stimulated with natural IFN- α , IFN- γ or IL-12 for 90 min *in vitro*, and then collected. pSTAT1 and pSTAT4 protein levels were evaluated by flow cytometry, electronically gating on CD56⁺ CD3⁻ NK cells. Representative histograms of a patient and an HS are shown. Dotted lines show staining of stimulated cells with isotype control. Thick lines show staining of unstimulated cells with the antibody. Thin lines with shaded areas show staining of stimulated cells with the antibody. Positive cell rates were determined based on the staining with isotype controls. Numbers are increased positive cell rates which were determined by subtracting the positive cell rate of unstimulated cells from those of stimulated cells. (B) Comparison of pSTAT1/4 level in response to IFN- α , IFN- γ or IL-12 between the patients with CHC ($n = 11$) and the HS ($n = 11$) are shown as increased pSTAT1/4 positive cell rate. Each circle represents individual data. Horizontal bars represent means. * $p < 0.001$ vs. HS.

expression is up-regulated by IFN- α or IFN- γ [23,24]. It is thus possible that the higher level of STAT1 in NK cells from the CHC patients was up-regulated by IFN- α and/or IFN- γ induced by a chronic host response to HCV.

Our real-time RT-PCR analyses showed that the induction level of SOCS1 mRNA in response to IFN- α was significantly greater in NK cells from the CHC patients than in those from the HS (Fig. 5). This finding is consistent with the observation that the phosphorylation level of STAT1 in response to IFN- α

was significantly stronger in NK cells from the CHC patients than in those from the HS (Fig. 3), because SOCS1 is a downstream gene of pSTAT1 [20]. On the other hand, SOCS1 is an inducible negative regulator which inhibits further activation of the pSTAT1 pathway [20]. It is therefore possible that this greater up-regulation of SOCS1 owing to the greater level of STAT1 phosphorylation in response to IFN- α finally results in a weaker response to IFN- α , that is, a weaker induction of ISGs via the pSTAT1 pathway. Indeed, an increase of the STAT1 level,

Research Article




Fig. 4. Correlation between STAT1 level and pSTAT1/4 level in response to IFN- α in NK cells from patients with chronic HCV infection. The relationship was statistically analyzed between intracellular STAT1 level in NK cells and increased pSTAT1/4 level in response to IFN- α in NK cells from patients with chronic HCV infection. Each circle represents individual data. The lines represent regression lines.

which is itself a downstream gene of pSTAT1, by IFN- α based therapy, correlated negatively in vivo with the basal STAT1 expression level before therapy (Miyagi et al. unpublished data). The higher basal STAT1 expression causes a greater level of STAT1 phosphorylation in response to IFN- α , which is followed by a greater level of SOCS1 induction and then might result in a lower increase of STAT1 level. Since perforin and granzyme B are also ISGs from the pSTAT1 pathway [6,25], it

is reasonable that the mRNA induction of perforin and granzyme B in response to IFN- α in NK cells from the CHC patients was not significantly greater but modestly lower than those from the HS (Fig. 5). Indeed, the enhancement of degranulation by IFN- α , that is, the increase of CD107a expression, in the presence of K562 cells in response to IFN- α , was significant in NK cells from the HS, but not in those from the CHC patients (Fig. 6).




Fig. 5. Induction of ISGs in NK cells in response to IFN- α in vitro. NK cells were purified from PBMCs derived from patients with chronic HCV infection (CHC) and healthy subjects (HS). Isolated NK cells were untreated or treated with natural IFN- α in vitro for 3 h, and then collected. The collected cell RNA level of the indicated genes and β -actin as a control were analyzed by real-time RT-PCR. Data are shown as the fold increase of treated cells compared with untreated cells, with means \pm standard error of the mean from five subjects in each group. * $p < 0.05$ vs. HS.




Fig. 6. Increase of NK cell degranulation by IFN- α stimulation. PBMCs derived from patients with chronic HCV infection (CHC) and healthy subjects (HS) were treated with or without IFN- α in the presence of K562 cells. The CD107a expression of NK cells were evaluated by flow cytometry, electronically gating on CD56 $^{+}$ CD3 $^{-}$ NK cells. Data are shown as the frequency of CD107a-positive NK cells treated with or without IFN- α . Each circle represents individual data. * $p < 0.05$.

Patients who were clear of HCV due to IFN- α based therapy exhibited a significant increase in NK cell numbers and activity in the peripheral blood as well as in the liver compared to those who were not able to become clear of the virus [26–28]. IFN- γ produced from NK cells have been reported to suppress HCV replication in vitro [29]. IFN- γ from NK cells is also considered to activate the subsequent adaptive immune response during virus infection as well as inhibit viral replication [1,2,7]. These findings suggest the involvement of NK cells with IFN- γ in the response to IFN- α based therapy. The present study showed that the level of STAT4 phosphorylation in response to IFN- α in NK cells correlated negatively with the intracellular STAT1 level in NK cells (Fig. 4), and that IFN- γ induction, which is one of the downstream genes of pSTAT4, was clearly weaker in NK cells from the patients with CHC than in those from the HS (Fig. 5). The lower activation in CHC patients of the pSTAT4-to-IFN- γ pathway in response to IFN- α in NK cells compared to those from the HS might be one of the mechanisms which make CHC patients resistant to IFN- α based therapy.

Ahlenstiel et al. have recently reported that chronic exposure to HCV-induced IFN- α caused NK cells to become functionally polarized towards a cytotoxic phenotype, but that lacked an increase in IFN- γ production [30]. Moreover, Oliviero et al. showed that NK cells from CHC patients were of a predominantly activating phenotype, and that these phenotypic changes were associated with enhanced cytotoxic activity and defective IFN- γ production [31]. These reports would be associated with our finding that NK cells from the CHC patients displayed a high level of STAT1 expression. Cytotoxic molecules such as perforin and granzyme, as well as STAT1, belong to the ISGs [6,25]. A high level of STAT1 in NK cells in the CHC patients might correspond to a high level of cytotoxic molecules in NK cells resulting in enhanced cytotoxic activity at the basal level. Indeed, the CD107a expression at basal level in NK cells from the CHC patients seemed to be modestly higher than that in NK cells from the HS (Fig. 6), although NK cells from the CHC patients did not significantly increase the CD107a expression in response to IFN- α . On the other hand, our present study showed that STAT1 expression level in NK cells correlated negatively with the activation of STAT4 to produce IFN- γ in response to IFN- α in NK cells. A high level of STAT1 in NK cells would also cause defective IFN- γ production in the NK cells of patients with CHC.

Recent studies have revealed that the higher level of ISGs in hepatocytes as well as in PBMCs before IFN- α based therapy is associated with resistance to this therapy [32–33]. The present study demonstrated that NK cells from CHC patients displayed higher levels of STAT1, which is one of the ISGs, compared with those from HS (Fig. 2). Those who had a higher STAT1 level in NK cells showed less STAT4 phosphorylation and more STAT1 phosphorylation in response to IFN- α , resulting in less IFN- γ induction and more SOCS1 induction. Thus, it is possible that those who have a higher STAT1 level in NK cells would display less response to IFN- α based therapy. Our preliminary data with a small number of patients treated with IFN- α based therapy revealed a tendency for those who had a higher STAT1 level in NK cells to not respond well to the therapy in the context of HCV clearance by week 8 after initiation of therapy (Supplementary Fig. 3).

In the present study, we found that the expression level of total STAT1, a key molecule of IFN- α signaling, was clearly higher

in NK cells from the patients with CHC than in those from the HS. The phosphorylation levels of STAT1 and STAT4 with IFN- α stimulation were altered in NK cells from the CHC patients, compared with those from the HS. The induction of IFN- γ mRNA expression with IFN- α stimulation, which is one of the downstream genes of pSTAT4, was clearly weaker in isolated NK cells from the CHC patients than in those from the HS. The induction of SOCS1 mRNA expression, which is one of the downstream genes of pSTAT1, was clearly stronger. The enhancement of NK cell degranulation, the increase of CD107a expression, in response to IFN- α was significant in the HS, but not in the CHC patients. These results indicate that IFN- α signaling in NK cells is altered in CHC patients, suggesting that this alteration of IFN- α signaling is associated with the persistence of chronic HCV infection and resistance to IFN- α therapy. The basal total STAT1 level in NK cells might enable the prediction of the outcome of IFN- α therapy against HCV infection, and thus could serve as a molecular target for more effective IFN- α based therapy.

Conflict of interests

The authors who have taken part in this study declare that they do not have anything to disclose regarding funding or conflict of interest with respect to this manuscript.

Acknowledgements

This work was supported by Grants-in-aid for Scientific Research (to T. Takehara and T. Miyagi) and Global Centers of Excellence Program (to T. Miyagi) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.jhep.2010.03.018.

References

- [1] Biron CA, Nguyen KB, Pien GC, Cousens LP, Salazar-Mather TP. Natural killer cells in antiviral defense: function and regulation by innate cytokines. *Annu Rev Immunol* 1999;17:189–220.
- [2] Farrar MA, Schreiber RD. The molecular cell biology of interferon-gamma and its receptor. *Annu Rev Immunol* 1993;11:571–611.
- [3] García-Sastre A, Biron CA. Type 1 interferons and the virus–host relationship: a lesson in détente. *Science* 2006;312:879–882.
- [4] Cho SS, Bacon CM, Sudarshan C, Rees RC, Finbloom D, Pine R, et al. Activation of STAT4 by IL-12 and IFN-alpha: evidence for the involvement of ligand-induced tyrosine and serine phosphorylation. *J Immunol* 1996;157:4781–4789.
- [5] Matikainen S, Paananen A, Miettinen M, Kurimoto M, Timonen T, Julkunen I, et al. IFN-alpha and IL-18 synergistically enhance IFN-gamma production in human NK cells: differential regulation of Stat4 activation and IFN-gamma gene expression by IFN-alpha and IL-12. *Eur J Immunol* 2001;31:2236–2245.
- [6] Liang S, Wei H, Sun R, Tian Z. IFNalpha regulates NK cell cytotoxicity through STAT1 pathway. *Cytokine* 2003;23:190–199.
- [7] Lee SH, Miyagi T, Biron CA. Keeping NK cells in highly regulated antiviral warfare. *Trends Immunol* 2007;28:252–259.
- [8] Liang TJ, Rehermann B, Seeff LB, Hoofnagle JH. Pathogenesis, natural history, treatment, and prevention of hepatitis C. *Ann Intern Med* 2000;132:296–305.

Research Article

- [9] Kamal SM, Fouly AE, Kamel RR, Hockenjos B, Al Tawil A, Khalifa KE, et al. Peginterferon alfa-2b therapy in acute hepatitis C: impact of onset of therapy on sustained virologic response. *Gastroenterology* 2006;130:632–638.
- [10] Manns MP, McHutchison JG, Gordon SC, Rustgi VK, Shiffman M, Reindollar R, et al. Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomised trial. *Lancet* 2001;358:958–965.
- [11] Fried MW, Shiffman ML, Reddy KR, Smith C, Marinos G, Gonçales FLJ, et al. Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 2002;347:975–982.
- [12] Santantonio T, Fasano M, Sinisi E, Guastadisegni A, Casalino C, Mazzola M, et al. Efficacy of a 24-week course of PEG-interferon alpha-2b monotherapy in patients with acute hepatitis C after failure of spontaneous clearance. *J Hepatol* 2005;42:329–333.
- [13] Takehara T, Hayashi N. Natural killer cells in hepatitis C virus infection: from innate immunity to adaptive immunity. *Clin Gastroenterol Hepatol* 2005;3:S78–S81.
- [14] Golden-Mason L, Rosen HR. Natural killer cells: primary target for hepatitis C virus immune evasion strategies? *Liver Transpl* 2006;12:363–372.
- [15] Szabo G, Chang S, Dolganiuc A. Altered innate immunity in chronic hepatitis C infection: cause or effect? *Hepatology* 2007;46:1279–1290.
- [16] Jinushi M, Takehara T, Tatsumi T, Kanto T, Miyagi T, Suzuki T, et al. Negative regulation of NK cell activities by inhibitory receptor CD94/NKG2A leads to altered NK cell-induced modulation of dendritic cell functions in chronic hepatitis C virus infection. *J Immunol* 2004;173:6072–6081.
- [17] Takehara T, Uemura A, Tatsumi T, Suzuki T, Kimura R, Shiotani A, et al. Natural killer cell-mediated ablation of metastatic liver tumors by hydrodynamic injection of IFNalpha gene to mice. *Int J Cancer* 2007;120:1252–1260.
- [18] Miyagi T, Gil MP, Wang X, Louten J, Chu WM, Biron CA. High basal STAT4 balanced by STAT1 induction to control type 1 interferon effects in natural killer cells. *J Exp Med* 2007;204:2383–2396.
- [19] Alter G, Malenfant JM, Altfeld M. CD107a as a functional marker for the identification of natural killer cell activity. *J Immunol Methods* 2004;294:15–22.
- [20] Alexander WS, Hilton DJ. The role of suppressors of cytokine signaling (SOCS) proteins in regulation of the immune response. *Annu Rev Immunol* 2004;22:503–529.
- [21] Pfeffer LM, Madey MA, Riely CA, Fleckenstein JF. The induction of type I interferon production in hepatitis C-infected patients. *J Interferon Cytokine Res* 2009;29:299–306.
- [22] Dolganiuc A, Norkina O, Kodys K, Catalano D, Bakis G, Marshall C, et al. Viral and host factors induce macrophage activation and loss of toll-like receptor tolerance in chronic HCV infection. *Gastroenterology* 2007;133:1627–1636.
- [23] Der SD, Zhou A, Williams BR, Silverman RH. Identification of genes differentially regulated by interferon alpha, beta, or gamma using oligonucleotide arrays. *Proc Natl Acad Sci USA* 1998;95:15623–15628.
- [24] Ji X, Cheung R, Cooper S, Li Q, Greenberg HB, He XS. Interferon alfa regulated gene expression in patients initiating interferon treatment for chronic hepatitis C. *Hepatology* 2003;37:610–621.
- [25] Zimmerer JM, Lesinski GB, Kondadasula SV, Karpa VI, Lehman A, Raychaudhury A, et al. IFN-alpha-induced signal transduction, gene expression, and antitumor activity of immune effector cells are negatively regulated by suppressor of cytokine signaling proteins. *J Immunol* 2007;178:4832–4845.
- [26] Bonavita MS, Franco A, Paroli M, Santillo I, Benvenuto R, De Petrillo G, et al. Normalization of depressed natural killer activity after interferon-alpha therapy is associated with a low frequency of relapse in patients with chronic hepatitis C. *Int J Tissue React* 1993;15:11–16.
- [27] Appasamy R, Bryant J, Hassanein T, Van Thiel DH, Whiteside TL. Effects of therapy with interferon-alpha on peripheral blood lymphocyte subsets and NK activity in patients with chronic hepatitis C. *Clin Immunol Immunopathol* 1994;73:350–357.
- [28] Yamagiwa S, Matsuda Y, Ichida T, Honda Y, Takamura M, Sugahara S, et al. Sustained response to interferon-alpha plus ribavirin therapy for chronic hepatitis C is closely associated with increased dynamism of intrahepatic natural killer and natural killer T cells. *Hepatol Res* 2008;38:664–672.
- [29] Wang SH, Huang CX, Ye L, Wang X, Song L, Wang YJ, et al. Natural killer cells suppress full cycle HCV infection of human hepatocytes. *J Viral Hepat* 2008;15:855–864.
- [30] Ahlenstiel G, Titerence RH, Koh C, Edlich B, Feld JJ, Rotman Y, et al. Natural killer cells are polarized toward cytotoxicity in chronic hepatitis C in an interferon-alfa-dependent manner. *Gastroenterology* 2010;138:325–335.
- [31] Oliviero B, Varchetta S, Paudice E, Michelone G, Zaramella M, Mavilio D, et al. Natural killer cell functional dichotomy in chronic hepatitis B and chronic hepatitis C virus infections. *Gastroenterology* 2009;137:1151–1160.
- [32] Feld JJ, Nanda S, Huang Y, Chen W, Cam M, Pusek SN, et al. Hepatic gene expression during treatment with peginterferon and ribavirin: identifying molecular pathways for treatment response. *Hepatology* 2007;46:1548–1563.
- [33] Sarasin-Filipowicz M, Oakeley EJ, Duong FH, Christen V, Terracciano L, Filipowicz W, et al. Interferon signaling and treatment outcome in chronic hepatitis C. *Proc Natl Acad Sci USA* 2008;105:7034–7039.

I κ B ζ is essential for natural killer cell activation in response to IL-12 and IL-18

Tohru Miyake^{a,b,c}, Takashi Satoh^{a,b}, Hiroki Kato^{a,b}, Kazufumi Matsushita^{a,b}, Yutaro Kumagai^{a,b}, Alexis Vandenberg^d, Tohru Tani^e, Tatsushi Muta^e, Shizuo Akira^{a,b,1}, and Osamu Takeuchi^{a,b}

^aLaboratory of Host Defense, ^dLaboratory of Systems Immunology, WPI Immunology Frontier Research Center, and ^bResearch Institute for Microbial Diseases, Osaka University, Osaka 565-0871, Japan; ^cDepartment of Surgery, Shiga University of Medical Science, Seta-tsukinowacho, Shiga 520-2192, Japan; and ^eLaboratory of Cell Recognition and Response, Graduate School of Life Sciences, Tohoku University, Sendai 980-8578, Japan

Contributed by Shizuo Akira, September 1, 2010 (sent for review August 20, 2010)

I κ B ζ , encoded by *Nfkbiz*, is a nuclear I κ B-like protein harboring ankyrin repeats. I κ B ζ has been shown to regulate IL-6 production in macrophages and Th17 development in T cells. However, the role of I κ B ζ in natural killer (NK) cells has not been understood. In the present study, we found that the expression of I κ B ζ was rapidly induced in response to IL-18 in NK cells, but not in T cells. Analysis of *Nfkbiz*^{-/-} mice revealed that I κ B ζ was essential for the production of IFN- γ production and cytotoxic activity in NK cells in response to IL-12 and/or IL-18 stimulation. IL-12/IL-18-mediated gene induction was profoundly impaired in *Nfkbiz*^{-/-} NK cells. Whereas the phosphorylation of STAT4 was normally induced by IL-12 stimulation, STAT4 was not recruited to the *Ifng* gene regions in *Nfkbiz*^{-/-} NK cells. Acetylation of histone 3 K9 on *Ifng* regions was also abrogated in *Nfkbiz*^{-/-} NK cells. I κ B ζ was recruited on the proximal promoter region of the *Ifng* gene, and overexpression of I κ B ζ together with IL-12 activated the *Ifng* promoter. Furthermore, *Nfkbiz*^{-/-} mice were highly susceptible to mouse MCMV infection. Taken together, these results demonstrate that I κ B ζ is essential for the activation of NK cells and antiviral host defense responses.

IFN- γ | transcription | virus infection | STAT4

Antiviral host defense is mediated by coordinated action of multiple cell types, such as innate immune cells, T cells, and natural killer (NK) cells (1, 2). In response to virus infection, NK cells induce apoptosis of infected cells by expressing perforins and granzymes and produce high amounts of IFN- γ (3–5). Various cytokines are important for activating NK cells to produce IFN- γ and increasing their cytotoxic activity. Type I IFNs produced from innate immune cells as well as cytokines such as IL-2, IL-12, IL-15, IL-18, and IL-21 are critical for NK-cell activation. Furthermore, NK-cell receptors harboring immunoreceptor tyrosine-based activation motifs (ITAMs) or those interacting with ITAM-containing adaptor proteins induce cytolytic activity and IFN- γ production (4). For instance, antibody-coated target cells are recognized by Fc γ RIII, which mediates antibody-dependent cellular cytotoxicity and IFN- γ production.

The expression of the *Ifng* gene is one of the hallmarks of NK-cell activation. Because *Ifng* is also highly expressed in T helper 1 (Th1) cells, the mechanisms of the *Ifng* promoter activation has been extensively studied in Th1 cells and NK cells (6, 7). Stimulation of NK cells with IL-12 and IL-18 leads to production of IFN- γ as well as elevation of cytotoxic activity (8, 9). It has been shown that sequence elements conserved among species (also known as conserved noncoding sequences, CNSs) present up to 100 kb from the transcription start site (TSS) of *Ifng* are involved in controlling the expression of IFN- γ (7, 10–12). Various transcription factors are recruited to the CNS of *Ifng*. IL-12 induces phosphorylation and nuclear translocation of STAT4, which is recruited to the cis-regulatory region of *Ifng* (7, 13, 14). IFN- γ -inducible T-box transcription factor (T-bet) is critical for inducing *Ifng* by controlling histone acetylation in Th1 cells (11, 15, 16). T-bet is also required for the development of NK cells (17). In T cells, T-bet-dependent chromatin remodeling of the *Ifng*

locus induces recruitment of the NF- κ B p65 subunit to *Ifng* cis-elements (12).

IL-18 is structurally related to IL-1 β and known to work together with IL-12 to activate Th1 cells to produce IFN- γ (18). However, IL-18 does not induce Th1 development by itself, unlike IL-12. Recent reports show that IL-18 is involved in the differentiation of Th2 cells and Th17 cells in murine experimental models, in addition to Th1 (18, 19). Although IL-18 is reported to activate AP-1 via MAP kinases to express *Ifng* (20), the mechanism of how IL-18 potentiates IFN- γ production in NK cells is yet to be clarified.

I κ B ζ , also known as INAP or MAIL, is a nuclear factor belonging to the Bcl-3 family, which contains a nuclear localization domain in the N terminus and C-terminal ankyrin repeats (21). I κ B ζ is encoded by the *Nfkbiz* gene, and the expression of *Nfkbiz* is rapidly induced in response to various Toll-like receptor (TLR)/IL-1 receptor (IL-1R) stimuli in macrophages (22). The expressed I κ B ζ interacts with NF- κ B p50 subunit and positively regulates expression of a set of genes including *Il6*, *Il12b*, *Csf2*, *Defb2*, *Lcn2*, and so on (22, 23). In addition to NF- κ B binding sequences, flanking C/EBP-binding sites are also important for I κ B ζ -mediated control of transcription (23). Furthermore, a recent study shows that I κ B ζ is required for chromatin remodeling downstream of TLR stimuli (24). Recently, it was shown that T cells lacking I κ B ζ showed severe defects in development of Th17 cells, and I κ B ζ regulates IL-17 expression by cooperating with *Rorgt* (25). On the other hand, a report showed that overexpression of I κ B ζ induced *Ifng* in a cell line, although the mechanism was not understood (26). Nevertheless, it is unclear whether I κ B ζ plays any role in the activation of NK cells.

In the present study, we found that I κ B ζ was required for the activation of NK cells in response to IL-12 and IL-18. IL-12/IL-18-mediated gene expression including *Ifng* was profoundly impaired in *Nfkbiz*^{-/-} NK cells. I κ B ζ was recruited to the proximal promoter region of *Ifng* and able to transactivate the *Ifng* together with IL-12. Furthermore, *Nfkbiz*^{-/-} mice were highly susceptible to infection with mouse CMV (MCMV), a model virus known to be cleared by NK cells. Taken together, I κ B ζ plays an essential role in the activation of NK cells by triggering dynamic transcription factor recruitment and chromatin remodeling.

Results

I κ B ζ Expression Is Induced in Response to IL-18 in NK Cells. To investigate whether I κ B ζ plays a role in controlling cells other than macrophages, we first examined the expression levels of I κ B ζ in

Author contributions: T. Miyake, S.A., and O.T. designed research; T. Miyake, T.S., H.K., K.M., and Y.K. performed research; T. Muta contributed new reagents/analytic tools; T. Miyake, A.V., T.T., S.A., and O.T. analyzed data; and T. Miyake, S.A., and O.T. wrote the paper.

The authors declare no conflict of interest.

¹To whom correspondence should be addressed. E-mail: sakira@biken.osaka-u.ac.jp.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1012977107/-DCSupplemental.

NK cells and T cells. We prepared splenic NK, and naïve T cells. Then, T cells were cultured in the presence of either IL-12, anti-IL-4 Ab, and anti-CD3 to develop Th1 cells or of IL-4, anti-IL-12, and anti-CD3 to develop Th2 cells. Quantitative-PCR (Q-PCR) analysis using RNAs prepared from these cells revealed that NK cells expressed the *IκBζ* gene higher than naïve T cells (Fig. 1A). Polarization to Th1 or Th2 cells did not increase the expression of *Nfkbiz*. The expression of *IκBζ* was highly induced in response to IL-12 and IL-18 stimulation in NK cells (Fig. 1A). Next, we examined kinetics of the *IκBζ* expression in NK cells in response to IL-12, IL-18, or both. *IκBζ* expression was induced at 1 h after stimulation with IL-18 alone or after costimulation with IL-12 and IL-18, and gradually decreased by 8 h after stimulation (Fig. 1B). In contrast, IL-12 alone slowly increased *Nfkbiz* about twofold at 8 h after stimulation. On the other hand, *Ifng* expression was increased in response to both IL-12 and IL-18, but not to either cytokine alone, and the expression peaked at 4 h after stimulation in NK cells (Fig. 1B). These results demonstrate that *IκBζ* expression is higher in NK cells than in T cells, and that this is an early induced gene in response to IL-18 stimulation in NK cells.

***IκBζ* Is Critical for Activation of NK Cells in Response to IL-12 and IL-18.** The expression of *IκBζ* in NK cells prompted us to investigate a functional role of *IκBζ* in NK cells by using *IκBζ*-deficient (*Nfkbiz*^{-/-}) mice. Stimulation of *Nfkbiz*^{+/-} splenic NK cells with IL-12, or costimulation with IL-12 and IL-18, induced production of IFN-γ (Fig. 2A). In contrast, IFN-γ production was severely impaired in *Nfkbiz*^{-/-} NK cells in response to IL-12, and IL-12 together with IL-18. These results indicated that *IκBζ* is indispensable for IFN-γ production in response to IL-12 and IL-18. Production of IFN-γ in response to culture with immobilized IgG1 was comparable in wild-type NK cells (Fig. 2A), indicating that signaling pathways from FcγRIII to induce *Ifng* expression is not affected in the absence of *IκBζ*. We then analyzed cytotoxic activity of NK cells to IL-12 and IL-18 stimulation by a standard ⁵¹Cr release assay against YAC1 target cells. Cytotoxic activity of NK cells stimulated with IL-12 alone or costimulated with IL-12 and IL-18 was reduced in *Nfkbiz*^{-/-} mice (Fig. 2B). IL-18 alone failed to increase cytotoxic activity even in wild-type NK cells




Fig. 1. Expression of *IκBζ* in NK cells in response to IL-12 and IL-18 stimulation. (A) The expression of *IκBζ* in naïve T, Th1, Th2 cells, and NK cells in response to IL-12 and IL-18. (B) NK cells were stimulated with IL-12 and/or IL-18 for the indicated periods, and the expression levels of *IκBζ* and IFN-γ were determined by Q-PCR analysis. Data represent means ± SD of triplicates. Similar results were obtained in three independent experiments.




Fig. 2. Role of *IκBζ* in the activation of NK cells in response to IL-12 and IL-18 stimulation. (A) Splenic NK cells from *Nfkbiz*^{+/-} and *Nfkbiz*^{-/-} mice were stimulated with 1 ng/mL IL-12, 10 ng/mL IL-18, and a combination of IL-12 and IL-18 or the cells were cultured on the plates coated with IgG1 for 24 h. The concentrations of IFN-γ in the culture supernatant were measured by ELISA. Data represent means ± SD. Similar results were obtained in three independent experiments. (B) IL-12-induced splenic NK cells were cultivated with IL-12 together with or without IL-18 for 4 h. NK-cell cytotoxicities against YAC-1 cells were measured by a ⁵¹Cr release assay at different E:T ratios. The percentages of specific cell lysis are indicated. Data represent the mean ± SD of duplicate measurements. The data shown are representative of two independent experiments. (C) Splenic NK cells from *Nfkbiz*^{+/-} and *Nfkbiz*^{-/-} mice were stimulated with IL-12 and IL-18 for 24 h, and total RNA was subjected to microarray analysis using Affymetrix mouse genome 430 2.0 microarray chips. The data were normalized by robust multichip analysis (RMA), and genes up-regulated more than twofold in wild-type NK cells after stimulation were selected. The genes were hierarchically clustered as follows: The expression values for each gene were standardized as average = 0 and SD = 1. Complete linkage method was then applied using absolute value of Pearson correlation as distance. The resulted heatmap, dendrogram, and a color key are shown. A blue line in the color key indicates the frequency distribution colors in the heatmap. For the gene lists shown in this figure, see Dataset S1.

(data not shown). These results demonstrate that *IκBζ* is critical for activation of NK cells in response to IL-12 and IL-18.

***IκBζ* Is Dispensable for NK-Cell Development.** We next analyzed the differentiation of splenic NK cells in *Nfkbiz*^{-/-} mice. It is known that NK-cell progenitors initially express CD122 (IL15R),

CD127 (IL7R), and c-kit (27). Upon maturation, NK cells down-regulate CD127 and c-kit. The DX5⁺CD3⁻ NK-cell population in spleen was comparable between *Nfkbiz*^{+/-} and *Nfkbiz*^{-/-} mice (Fig. S1). Furthermore, CD122 was highly expressed, and c-kit and CD127 expressions were comparably suppressed both in wild-type and *Nfkbiz*^{-/-} splenic NK cells (Fig. S1). Collectively, IκBζ is dispensable for controlling differentiation of NK cells.

Profound Impairment in Gene Expression in *Nfkbiz*^{-/-} NK Cells in Response to IL-12 and IL-18 Stimulation. Next we examined the role of IκBζ in IL-12 and IL-18-mediated gene expression in NK cells. Wild-type and *Nfkbiz*^{-/-} NK cells were stimulated with IL-12 and IL-18 for 24 h, and gene expression profile was assessed by microarray analysis. In response to stimulation with IL-12 and IL-18, 96 genes were up-regulated more than twofold in wild-type NK cells. We then hierarchically clustered genes induced by IL-12 and IL-18 and found that they are roughly classified into those completely (58 genes) or partially (38 genes) dependent on IκBζ (Fig. 2C) (Dataset S1). Genes completely dependent on IκBζ include *Ifng*, *Il22*, *Ccr5*, and *Il18r1*. These data demonstrate that IκBζ is critical for IL-12 and IL-18-mediated gene expression in NK cells.

IκBζ Promotes STAT4 Accessibility in IFN-γ Genes in Response to IL-12 and IL-18 in NK Cells. IL-12 activates Janus kinases (JAKs) that phosphorylate STAT4 (14). STAT4 is localized in the cytoplasm in the resting cells, and phosphorylated STAT4 translocates to the nucleus for the transcriptional activation. Treatment with IL-12 greatly increased NK cells stained with anti-phospho-STAT4 (p-STAT4), and the levels of p-STAT4 positive cells were comparable between *Nfkbiz*^{+/-} and *Nfkbiz*^{-/-} mice (Fig. 3A). Furthermore, nuclear translocation of STAT4 was comparably induced in *Nfkbiz*^{+/-} and *Nfkbiz*^{-/-} NK cells in response to stimulation with IL-12 and IL-18 (Fig. 3B), indicating that IκBζ was dispensable for phosphorylation and nuclear translocation of STAT4. We then examined recruitment of STAT4 to *Ifng* CNSs by chromatin immunoprecipitation (ChIP) coupled with Q-PCR (ChIP-Q-PCR) analysis. We found that STAT4 was widely recruited to CNSs (-33 kb, -22 kb, -6 kb, intron 1a, +10 kb, +20 kb, and +30 kb from the TSS) of *Ifng* in response to IL-12 and IL-18 stimulation in wild-type NK cells (Fig. 4A). In contrast, STAT4 recruitment to the *Ifng* conserved elements was

severely impaired in *Nfkbiz*^{-/-} NK cells through the CNS (Fig. 4A). These results suggested that IκBζ promotes STAT4 accessibility to the CNS of *Ifng* gene in stimulated NK cells.

IκBζ Is Required for Change in Histone 3 Lysine 9 Acetylation in Response to IL-12 and IL-18 in NK Cells. It has been shown that histones of the *Ifng* loci were hyperacetylated even in the absence of stimulation in NK cells, compared with T cells (10). We performed ChIP analysis with anti-acetyl histone 3 lysine 9 (H3K9) antibody to assess H3K9 acetylation in *Ifng* CNS. The analysis revealed that intron regions of *Ifng* were hyperacetylated even without stimulation in wild-type and *Nfkbiz*^{-/-} NK cells (Fig. 4B). Whereas H3K9 acetylation levels at the -6 kb region of *Ifng* were up-regulated in response to IL-12 and IL-18 in wild-type NK cells, *Nfkbiz*^{-/-} NK cells failed to induce H3K9 acetylation (Fig. 4B). These results suggest that IκBζ is required for change in H3K9 acetylation on *Ifng* loci in response to IL-12 and IL-18 stimulation.

IκBζ Is Recruited to the *Ifng* Proximal Promoter Region. To examine the recruitment of IκBζ to the *Ifng* promoter, we examined ChIP analysis using anti-IκBζ antibody. In contrast to STAT4 recruitment or H3K9 acetylation, IκBζ was not recruited to the -6 kb region of *Ifng* locus in NK cells in response to IL-12 and IL-18 (Fig. 5A). In contrast, IκBζ was recruited to the proximal promoter region (-0.2 kb from the TSS) of *Ifng* (Fig. 5A). IκBζ failed to associate with other CNSs such as -22 kb and intron 1a (data not shown). On the other hand, STAT4 was not recruited to the proximal *Ifng* promoter (data not shown). To investigate whether IκBζ directly regulates *Ifng* through binding to the




Fig. 3. Nuclear translocation of STAT4 in NK cells in the absence of IκBζ. (A) Splenic NK cells were stimulated with IL-12 for 30 min. Phosphorylation of STAT4 was then determined by intracellular staining of cells with anti-phospho-STAT4 Ab followed by flow cytometry. (B) NK cells were stimulated with IL-12 and IL-18 for 1 h. Cytoplasmic (CP) and nuclear (NE) proteins were prepared and immunoblotted using anti-STAT4 Ab and anti-JNK as a control.




Fig. 4. Requirement of IκBζ in the recruitment of STAT4 and acetylation of H3K9 in NK cells in response to IL-12 and IL-18. Chromatin was isolated from wild-type and *Nfkbiz*^{-/-} NK cells stimulated with IL-12 and IL-18 for 4 h; recruitment of STAT4 (A) and acetylation of H3K9 (B) to indicated *Ifng* gene regions was determined by ChIP-QPCR analysis. The data are representative of two independent experiments.




Fig. 5. Regulation of *Ifng* proximal promoter region by IκBζ in NK cells. (A) Chromatin was isolated from splenic NK cells stimulated with IL-12 and IL-18 for 4 h, recruitment of IκBζ to indicated *Ifng* gene regions was determined by ChIP-QPCR analysis. (B) A 3.6-kb fragment of the human *Ifng* promoter luciferase reporter construct was transfected to EL4 cells with increasing amounts of IκBζ construct. The luciferase activity was measured 18 h after transfection. (C) EL4 cells were cotransfected with the *Ifng* promoter reporter construct and IκBζ, followed by stimulation with IL-12 or IL-18. The luciferase activity was measured 18 h after stimulation.

proximal promoter region, we expressed a reporter construct with the human *Ifng* promoter region (-3.6 kb to +70 k) linked to the luciferase gene, together with IκBζ in EL4 cells. As shown in Fig. 5B, overexpression of IκBζ enhanced luciferase reporter activity in a dose-dependent manner. Furthermore, stimulation with IL-12, but not IL-18, synergistically activated the *Ifng* promoter with overexpression of IκBζ (Fig. 5C), suggesting that IκBζ expression facilitates IL-12-mediated *Ifng* promoter activation. These observations suggest that the recruitment of IκBζ to the proximal promoter region is responsible for the transcriptional activation of *Ifng*.

Essential Role of IκBζ in Host Defense Against MCMV Infection. It is known that NK cells play an important role in host defense against MCMV infection (1, 5, 28). MCMV-induced activation of innate immune cells such as dendritic cells (DCs) induces the production of type I IFNs as well as IL-12, leading to activation of NK cells to produce IFN-γ. To examine the role of IκBζ in activating NK cells in vivo in response to virus infection, we intraperitoneally inoculated MCMV into wild-type and *Nfkbiz*^{-/-} mice. MCMV-induced production of IL-12p40, but not IFN-α, in the sera was partially impaired in *Nfkbiz*^{-/-} mice compared with wild-type mice (Fig. 6A), suggesting that IκBζ plays a role in expressing IL-12p40 in innate immune cells in response to MCMV. In contrast, the production of IFN-γ in the sera was completely abrogated in *Nfkbiz*^{-/-} mice (Fig. 6A). Furthermore, flow cytometric analysis revealed that the expression of IFN-γ in splenic NK cells was severely impaired in MCMV-infected *Nfkbiz*^{-/-} mice (Fig. 6B), although NK-cell cytotoxicity was not altered in *Nfkbiz*^{-/-} NK cells (data not shown). Although all wild-type mice and 80% of the *Nfkbiz*^{+/-} mice survived, *Nfkbiz*^{-/-} mice were highly susceptible and more than 80% of the *Nfkbiz*^{-/-} mice died within 6 d after MCMV infection (Fig. 6C). Taken together, IκBζ is essential for host defense against MCMV infection by inducing IFN-γ production from NK cells.

Discussion

In the present study, we examined the role of IκBζ in NK cells and found that this nuclear protein is essential for inducing a set of genes including *Ifng* in response to IL-12 and IL-18. IκBζ was rapidly induced in response to IL-18 in NK cells, although the




Fig. 6. Essential role of IκBζ in host defense against MCMV infection. (A) Wild-type and *Nfkbiz*^{-/-} mice (n = 3–5) were intraperitoneally infected with 1 × 10⁵ pfu MCMV, and the concentrations of IFN-α, IFN-γ, and IL-12p40 in the sera 36 h after infection were measured by ELISA. (B) The expression of IFN-γ in splenic NK cells prepared from *Nfkbiz*^{+/-} and *Nfkbiz*^{-/-} mice 36 h after MCMV infection was determined by intracellular staining of cells with anti-IFN-γ Ab followed by flow cytometry. (C) *Nfkbiz*^{+/+} (n = 5), *Nfkbiz*^{+/-} (n = 5), and *Nfkbiz*^{-/-} (n = 6) mice were intraperitoneally infected with MCMV, and the survival of mice was monitored daily. The data are representative of two independent infection experiments.

expression level was kept low in Th1 cells. IL-18 is a cytokine that is able to activate NK cells together with IL-12, although the precise mechanisms of the action of IL-18 has not been understood well (18). IL-18R triggers intracellular signaling pathways leading to the activation of NF-κB and MAP kinases (29). We have previously shown that IκBζ is a TLR/IL-1R-inducible gene in macrophages (22). Given that IL-18R and TLR/IL-1R share MyD88 as an adaptor molecule for the signaling, it is not surprising that IL-18 also induces IκBζ expression. IL-18 is known to augment IFN-γ production, when costimulated with IL-12.

IL-12 induces phosphorylation and nuclear translocation of STAT4, which is recruited to cis-regulatory regions of several genes including *Ifng* (6, 7). It has been shown that sequence elements conserved among species present up to 100 kb from the transcription start site of *Ifng* are involved in controlling the expression of *Ifng* (12). These distal elements may loop toward the *Ifng* gene for regulating its transcription (6, 7). Whereas the H3K9 acetylation as well as recruitment of STAT4 to distal conserved regions of *Ifng* was abrogated in *Nfkbiz*^{-/-} NK cells, IκBζ was recruited to a proximal promoter region of *Ifng*, but not to distal sequences. These results imply that initial recruitment of

I κ B ζ to the proximal region is required for further chromatin remodelling and efficient STAT4 binding. A report showed that I κ B ζ inhibits STAT3-mediated signaling by direct interaction (30). However, given that STAT4 and I κ B ζ are recruited to different conserved elements on the *Irfng* promoter, it seems reasonable to assume that I κ B ζ and STAT4 do not control the *Irfng* expression by physically interacting with each other.

I κ B ζ interacts with NF- κ Bp50 in macrophages, and the ankyrin repeats of this protein were required for synergistically increasing transcriptional activity (22). Furthermore, I κ B ζ is reported to be involved in histone modifications on the *Lcn2* promoter in LPS-stimulated macrophages (24). A previous report shows that NF- κ Bp50-deficient mice showed increased, but not decreased, IFN- γ production in response to IL-12 and IL-18 stimulation in NK cells (31). Further, no clear NF- κ B binding sequence was found in the proximal promoter region of *Irfng*. Thus, it is unlikely that NF- κ Bp50-I κ B ζ interaction plays a pivotal role in controlling IFN- γ production in NK cells. In Th17 cells, I κ B ζ was suggested to directly control IL-17 expression by recruiting to the *Il17* promoter regions (25). However, clear I κ B ζ responsive elements were not found in the proximal region of *Irfng*. Thus, it is possible that I κ B ζ is directly recruited to *Irfng* promoter, although further studies are required to address the mechanism of *Irfng* transactivation by I κ B ζ .

In contrast to the critical role of I κ B ζ in NK-cell activation in response to IL-12 and IL-18, *Nfkbiz*^{-/-} Th1 cells developed in the presence of IL-12 and IL-18 did not show a defect in producing IFN- γ in response to antigen-receptor stimulation (25). In contrast, I κ B ζ is involved in the development of Th-17 cells (25). We found that IL-18 stimulation failed to up-regulate I κ B ζ expression in T cells. Although NK cells produced IFN- γ by stimulation with IL-12 and IL-18, Th1 cells require T-cell receptor (TCR) stimulation in addition to culture with these cytokines. In addition to the difference in I κ B ζ expression levels in NK cells and T cells, differential requirement of antigen-receptor signaling in producing IFN- γ may explain the role of I κ B ζ in NK cells but not in Th1 cells.

Although I κ B ζ was essential for IFN- γ production in response to IL-12 and IL-18 stimulation, *Nfkbiz*^{-/-} NK cells produced IFN- γ normally in response to culture with IgG1, which is known to activate NK cells via Fc γ RIIIa. In contrast to IL-12 and IL-18, Fc γ RIII activates the intracellular signaling via recruiting ITAM-motif containing adaptors activating Syk tyrosine kinase and ZAP-70 (32). Subsequently, phospholipase C (PLC)- γ is activated to induce Ca²⁺ influx, MAP kinase activation, and NF- κ B activation. The Fc γ RIII and TCR largely share the signaling molecules, and future studies are required for identifying the mechanisms of how activating Fc γ RIII and TCR induce production of IFN- γ independent of I κ B ζ .

NK cells are important for the elimination of MCMV (1, 5). Type I IFNs and IL-12 produced from DC subsets such as plasmacytoid DC and CD8 α ⁺ DC are important for the NK-cell cytotoxic activity and production of IFN- γ , respectively (3, 4). Whereas MCMV-mediated production of IFN- α in the sera was normal, IL-12p40 levels were decreased in *Nfkbiz*^{-/-} mice. Because I κ B ζ is required for the production of IL-6 and IL-12p40 in response to TLR ligands in macrophages (22), impaired production of IL-12p40 in *Nfkbiz*^{-/-} mice is likely due to the defect in TLR-mediated MCMV recognition. Although type I IFNs are known to activate NK cells in response to virus infection, it has been shown that IFN- γ production to MCMV infection was not decreased even in the absence of type I IFN production or IFN β signaling. On the other hand, we did not find a defect in MCMV-induced NK-cell cytotoxicity in *Nfkbiz*^{-/-} mice. Because NK-cell cytotoxicity to MCMV infection was reported to be IL-12-independent, normal activation of NK-cell cytotoxicity to MCMV infection in *Nfkbiz*^{-/-} mice suggests that I κ B ζ is not involved in the type I IFN signaling.

In summary, our data demonstrate that I κ B ζ is essential for NK-cell activation in response to IL-12 and IL-18 stimulation by inducing various genes including IFN- γ . Furthermore, induction of I κ B ζ is one of the important host defense mechanisms against virus infection via NK cells. Given that I κ B ζ looks to control different set of genes in a cell-type specific manner, further studies will clarify the mechanisms of I κ B ζ in controlling gene expression.

Materials and Methods

Mice, Cells, and Reagents. *Nfkbiz*^{-/-} (22) and littermate control mice were used at 6–16 wk of age. Mouse experiments were conducted in accordance with institutional guidelines of animal care and use committees. Splenic NK cells were enriched via MACS separation with DX5 MicroBeads (Miltenyi Biotec). YAC-1 cells were as described previously (33). EL4 cells were obtained from ATCC (TIB-39). IL-12 and IL-2 were purchased from R&D Systems. Biotinylated rabbit anti-mouse IgG1 was purchased from Zymed. ELISA kits for IFN- γ and IL-12p40 were purchased from R&D Systems. IL-18 and ELISA kit for IFN α were purchased from PBL.

Q-PCR. Total RNA was isolated using TRIzol (Invitrogen) and reverse transcription was performed with ReverTra Ace (Toyobo) according to the manufacturer's instructions. For Q-PCR, cDNA fragments were amplified by real-time PCR Master Mix (Toyobo), Fluorescence from the taqman probe for each cytokine was detected by a 7500 real-time PCR system (Applied Biosystems). To determine the relative induction of cytokine mRNA in response to various stimuli, the mRNA expression level of each gene was normalized to the expression level of 18S rRNA.

Cytotoxicity Assay. Splenic NK cells were cultured with IL-2 (2,000 U/mL) for 7 d. IL-2-induced NK cells were stimulated with IL-12 (0.1 ng/mL) and IL-18 (1 ng/mL) for 4 h. The ⁵¹Cr-labeled YAC-1 cells (1 \times 10⁴ cells/well) were cultured with indicated numbers of NK cells. After 4-h incubation, the supernatant was harvested and the ⁵¹Cr radioactivity of supernatants were measured as described previously (33).

Intracellular Staining of STAT4. Antibodies for flow cytometry were purchased from BD or eBioscience. Splenocytes were stimulated with IL-12 (2 ng/mL) for 30 min and were fixed immediately with PhosFlow Lyse/Fix buffer (BD) for 10 min. Cells were washed once with PBS and stained with DX5-PE and CD3e-APC. Cells were then permeabilized by BD PhosFlow Perm Buffer III and incubated for 25 min on ice. Cells were washed with BD Pharmingen Stain Buffer and stained with Phospho-STAT4-FITC at room temperature for 30 min in the dark. Cells were washed and acquired on a FACS Calibur flow cytometer (BD) and analyzed using FlowJo (Tree Star).

Immunoblot Analysis. NK cells were lysed in a buffer containing 0.1% Nonidet-P40, 10 mM KCl, 10 mM Hepes-KOH (pH 7.8), 0.1 mM EDTA and a protease inhibitor mixture (Roche). Cell lysates were centrifuged at 5,000 rpm for 5 min, and the supernatant was harvested as cytoplasmic proteins. Pellets were lysed in a buffer containing 5 mM MgCl₂, 420 mM KCl, 50 mM Hepes-KOH (pH 7.8), 0.1 mM EDTA and 20% glycerol for 30 min on ice. The lysates were centrifuged at 18,500 \times g for 15 min, and the supernatant was harvested as nuclear proteins. Cell lysates were dissolved by SDS/PAGE and transferred onto a polyvinylidene difluoride membrane. The membrane was blotted with the specific antibodies to indicated proteins.

Chromatin Immunoprecipitation (ChIP). ChIP was performed by using a chip assay kit (Millipore). NK cells were stimulated with IL-12 (20 ng/mL) and IL-18 (100 ng/mL) for 4 h and were fixed with 1% formaldehyde for 10 min or 10 mM DMA for 20 min. Cells were then washed and resuspended in SDS buffer. Lysates were sonicated to obtain DNA fragments with a peak in size between 200 and 1,000 bp. Lysates precleared with protein A agarose/salmon sperm DNA were incubated with antibodies against STAT4 (Santa Cruz) or H3K9 (Abcam) and I κ B ζ (kindly provided by Drs. Kayama and Takeda, Osaka University, Suita, Japan), and immunoprecipitated at 4 $^{\circ}$ C overnight. The immune complexes were absorbed with beads. Beads were washed with low salt buffer, high salt buffer, LiCl wash buffer, and in TE buffer. Immune complexes were extracted with elution buffer (1% SDS 100 mM NaHCO₃, 10 mM DTT) and cross links were reversed by incubation overnight at 65 $^{\circ}$ C. After proteinase K treatment for 1 h, DNA was then purified via ethanol precipitation. The purified DNA was used in qPCR to assess the presence of target sequences. Quantitative RT-PCR

was performed with Thunderbird qPCR Mix (Toyobo) in an Applied Biosystem 7300. Primers used for amplifying *Irfng* loci were shown previously (10, 34).

Reporter Assay. EL4 cells were transfected with either an empty vector or the $\kappa B\zeta$ vector, human *Irfng* reporter plasmid containing the firefly luciferase gene (1 μ g) (Addgene plasmid 17598) (35) and 50 ng of pRL-TK (Promega) by using DMRIE-C reagent (Invitrogen). EL4 cells were incubated in the presence of IL-12 (2 ng/mL) and IL-18 (10 ng/mL) for 18 h after transfection. Luciferase activities were measured with a Dual-Luciferase reporter assay system (Promega).

Microarray. Splenic NK cells were stimulated with IL-12 (1 ng/mL) and IL-18 (10 ng/mL) for 24 h. Then, total RNA was extracted with TRIzol (Invitrogen Life Technologies) and further purified using an RNeasy kit (Qiagen). Biotin-labeled cDNA was synthesized from 100 ng of total RNA using the Ovation Biotin RNA amplification and labeling system (Nugen) according to the

manufacturer's protocol. Hybridization, staining, washing, and scanning of Affymetrix mouse Genome 430 2.0 microarray chips were conducted according to the manufacturer's instructions. Data analysis was performed using R.

ACKNOWLEDGMENTS. We thank Dr. M. Ueda at Kyoto Prefectural University of Medicine (Kyoto, Japan) for providing *Nfkbiz*^{-/-} mice, Dr. M. Dalod at Université de la Méditerranée (Marseille, France) for providing MCMV, and Drs. H. Kayama and K. Takeda at Osaka University (Suita, Japan) for providing anti- $\kappa B\zeta$ antibody. We thank all of our colleagues in our laboratory, E. Kamada for secretarial assistance, and Y. Fujiwara, M. Kumagai, and R. Abe for technical assistance. This research was supported by Funding Program for World-Leading Innovative R&D on Science and Technology of the Japan Society for the Promotion of Science, the Special Coordination Funds of the Japanese Ministry of Education, Culture, Sports, Science and Technology, grants from the Ministry of Health, Labor and Welfare in Japan, the Global Center of Excellence Program, and the National Institutes of Health Grant P01 AI070167.

- Beutler B, et al. (2007) Genetic analysis of resistance to viral infection. *Nat Rev Immunol* 7:753–766.
- Takeuchi O, Akira S (2010) Pattern recognition receptors and inflammation. *Cell* 140: 805–820.
- Biron CA, Nguyen KB, Pien GC, Cousens LP, Salazar-Mather TP (1999) Natural killer cells in antiviral defense: function and regulation by innate cytokines. *Annu Rev Immunol* 17:189–220.
- Lanier LL (2005) NK cell recognition. *Annu Rev Immunol* 23:225–274.
- Lanier LL (2008) Evolutionary struggles between NK cells and viruses. *Nat Rev Immunol* 8:259–268.
- Wilson CB, Rowell E, Sekimata M (2009) Epigenetic control of T-helper-cell differentiation. *Nat Rev Immunol* 9:91–105.
- Aune TM, Collins PL, Chang S (2009) Epigenetics and T helper 1 differentiation. *Immunology* 126:299–305.
- Okamura H, Kashiwamura S, Tsutsui H, Yoshimoto T, Nakanishi K (1998) Regulation of interferon-gamma production by IL-12 and IL-18. *Curr Opin Immunol* 10:259–264.
- Takeda K, et al. (1998) Defective NK cell activity and Th1 response in IL-18-deficient mice. *Immunity* 8:383–390.
- Chang S, Aune TM (2005) Histone hyperacetylated domains across the *Irfng* gene region in natural killer cells and T cells. *Proc Natl Acad Sci USA* 102:17095–17100.
- Hatton RD, et al. (2006) A distal conserved sequence element controls *Irfng* gene expression by T cells and NK cells. *Immunity* 25:717–729.
- Balasubramani A, et al. (2010) Modular utilization of distal cis-regulatory elements controls *Irfng* gene expression in T cells activated by distinct stimuli. *Immunity* 33: 35–47.
- Thierfelder WE, et al. (1996) Requirement for Stat4 in interleukin-12-mediated responses of natural killer and T cells. *Nature* 382:171–174.
- Levy DE, Darnell JE, Jr (2002) Stats: Transcriptional control and biological impact. *Nat Rev Mol Cell Biol* 3:651–662.
- Szabo SJ, et al. (2000) A novel transcription factor, T-bet, directs Th1 lineage commitment. *Cell* 100:655–669.
- Avni O, et al. (2002) T(H) cell differentiation is accompanied by dynamic changes in histone acetylation of cytokine genes. *Nat Immunol* 3:643–651.
- Townsend MJ, et al. (2004) T-bet regulates the terminal maturation and homeostasis of NK and Valpha14i NKT cells. *Immunity* 20:477–494.
- Nakanishi K, Yoshimoto T, Tsutsui H, Okamura H (2001) Interleukin-18 regulates both Th1 and Th2 responses. *Annu Rev Immunol* 19:423–474.
- Weaver CT, Harrington LE, Mangan PR, Gavrieli M, Murphy KM (2006) Th17: An effector CD4 T cell lineage with regulatory T cell ties. *Immunity* 24:677–688.
- Nakahira M, et al. (2002) Synergy of IL-12 and IL-18 for IFN-gamma gene expression: IL-12-induced STAT4 contributes to IFN-gamma promoter activation by up-regulating the binding activity of IL-18-induced activator protein 1. *J Immunol* 168:1146–1153.
- Yamazaki S, Muta T, Takeshige K (2001) A novel IkappaB protein, IkappaB-zeta, induced by proinflammatory stimuli, negatively regulates nuclear factor-kappaB in the nuclei. *J Biol Chem* 276:27657–27662.
- Yamamoto M, et al. (2004) Regulation of Toll/IL-1-receptor-mediated gene expression by the inducible nuclear protein IkappaBzeta. *Nature* 430:218–222.
- Matsuo S, Yamazaki S, Takeshige K, Muta T (2007) Crucial roles of binding sites for NF-kappaB and C/EBPs in IkappaB-zeta-mediated transcriptional activation. *Biochem J* 405:605–615.
- Kayama H, et al. (2008) Class-specific regulation of pro-inflammatory genes by MyD88 pathways and IkappaBzeta. *J Biol Chem* 283:12468–12477.
- Okamoto K, et al. (2010) IkappaBzeta regulates T(H)17 development by cooperating with ROR nuclear receptors. *Nature* 464:1381–1385.
- Raices RM, et al. (2009) A novel role for IkappaBzeta in the regulation of IFNgamma production. *PLoS ONE* 4:e6776.
- Di Santo JP (2006) Natural killer cell developmental pathways: A question of balance. *Annu Rev Immunol* 24:257–286.
- Vidal SM, Lanier LL (2006) NK cell recognition of mouse cytomegalovirus-infected cells. *Curr Top Microbiol Immunol* 298:183–206.
- Hoshino K, et al. (1999) Cutting edge: Generation of IL-18 receptor-deficient mice: Evidence for IL-1 receptor-related protein as an essential IL-18 binding receptor. *J Immunol* 162:5041–5044.
- Wu Z, et al. (2009) Nuclear protein IkappaB-zeta inhibits the activity of STAT3. *Biochem Biophys Res Commun* 387:348–352.
- Tato CM, et al. (2006) Opposing roles of NF-kappaB family members in the regulation of NK cell proliferation and production of IFN-gamma. *Int Immunol* 18:505–513.
- Nimmerjahn F, Ravetch JV (2008) Fcgamma receptors as regulators of immune responses. *Nat Rev Immunol* 8:34–47.
- Miyake T, et al. (2009) Poly I:C-induced activation of NK cells by CD8 alpha+ dendritic cells via the IPS-1 and TRIF-dependent pathways. *J Immunol* 183:2522–2528.
- Bandyopadhyay S, Qui HZ, Adler AJ (2009) In vitro and in vivo differentiated effector CD8 T cells display divergent histone acetylation patterns within the *Irfng* locus. *Immunol Lett* 122:214–218.
- Gonsky R, et al. (2000) Mucosa-specific targets for regulation of IFN-gamma expression: Lamina propria T cells use different cis-elements than peripheral blood T cells to regulate transactivation of IFN-gamma expression. *J Immunol* 164:1399–1407.

The *Jmjd3-Irf4* axis regulates M2 macrophage polarization and host responses against helminth infection

Takashi Satoh^{1,2,8}, Osamu Takeuchi^{1,2,8}, Alexis Vandenbon³, Koubun Yasuda⁴, Yoshiaki Tanaka⁵, Yutaro Kumagai^{1,2}, Tohru Miyake^{1,2}, Kazufumi Matsushita^{1,2}, Toshihiko Okazaki¹, Tatsuya Saitoh^{1,2}, Kiri Honma⁶, Toshifumi Matsuyama⁶, Katsuyuki Yui⁶, Tohru Tsujimura⁷, Daron M Standley³, Kenji Nakanishi⁴, Kenta Nakai⁶ & Shizuo Akira^{1,2}

Polarization of macrophages to M1 or M2 cells is important for mounting responses against bacterial and helminth infections, respectively. Jumonji domain containing-3 (*Jmjd3*), a histone 3 Lys27 (H3K27) demethylase, has been implicated in the activation of macrophages. Here we show that *Jmjd3* is essential for M2 macrophage polarization in response to helminth infection and chitin, though *Jmjd3* is dispensable for M1 responses. Furthermore, *Jmjd3* (also known as *Kdm6b*) is essential for proper bone marrow macrophage differentiation, and this function depends on demethylase activity of *Jmjd3*. *Jmjd3* deficiency affected trimethylation of H3K27 in only a limited number of genes. Among them, we identified *Irf4* as encoding a key transcription factor that controls M2 macrophage polarization. Collectively, these results show that *Jmjd3*-mediated H3K27 demethylation is crucial for regulating M2 macrophage development leading to anti-helminth host responses.

Innate immune cells such as macrophages sense the presence of microbial infection through pattern-recognition receptors and mount anti-microbial responses^{1–3}. The Toll-like receptor (TLR) family is one of the best-characterized PRR families recognizing various pathogens such as bacteria, viruses, protozoa and fungi. TLRs are crucial in evoking innate immune responses to infection by various pathogens, leading to production of inflammatory mediators, including proinflammatory cytokines, chemokines and interferons.

Macrophages are functionally polarized into M1 and M2 cells in response to infection with microorganisms and host mediators^{4,5}. M1 macrophages produce large amounts of nitric oxide by expressing inducible nitric oxide synthase (iNOS) and tumor necrosis factor (TNF), and are essential for clearing bacterial, viral and fungal infections⁶. Other macrophages, called alternatively activated macrophages or M2 macrophages, have an important role in responses to parasite infection, tissue remodeling, angiogenesis and tumor progression. M2 macrophages are characterized by their high expression of arginase-1 (*Arg1*), chitinase-like Ym1 (*Chi3l3*), found in inflammatory zone-1 (*Fizz1*, also called *Retnla*) mannose receptor (*Mrc1* encoding MR, also known as CD206), and chemokines such as CCL17 and CCL24 (refs. 7–11). The PRR system responsible for the recognition of helminth infection and M2 polarization has yet to be identified.

Cytokines and growth factors have been implicated in the reprogramming of M1 and M2 macrophages. Whereas interferon- γ (IFN- γ), produced by activated T cells and TLR ligands, induces M1 macrophage generation, stimulation of macrophages with interleukin-4 (IL-4) or IL-13 induces M2-type macrophages^{4,5}. In addition, immune complexes, IL-10 and glucocorticoid or secosteroid hormones are also known to generate M2 macrophages. Among growth factors, treatment of bone marrow cells with granulocyte-macrophage colony-stimulating factor (GM-CSF) and macrophage colony-stimulating factor (M-CSF) leads to generation of M1 and M2 cells, respectively^{12–15}.

TLRs trigger intracellular signaling pathways, inducing activation of a set of transcription factors, such as NF- κ B, AP-1, C/EBP β , PU.1 and interferon-regulatory factors (IRFs)^{1,16}. These transcription factors cooperatively upregulate the expression of multiple genes such as proinflammatory cytokines, leading to M1 macrophage polarization. Proteins induced by TLR signaling are known to modulate inflammatory responses. I κ B ζ , an I κ B family member, positively regulates certain genes such as *Il6* by interacting with NF- κ Bp50 or inducing histone modification^{17,18}, whereas ATF3 negatively regulates expression¹⁹. In contrast, transcription factors such as STAT6 and peroxisome proliferator-activated receptor- γ (PPAR γ) are involved in polarization of M2 macrophages^{5,20}.

¹Laboratory of Host Defense, World Premier Initiative Immunology Frontier Research Center, Osaka University, Osaka, Japan. ²Research Institute for Microbial Diseases, Osaka University, Osaka, Japan. ³Laboratory of Systems Immunology, World Premier International Immunology Frontier Research Center, Osaka University, Osaka, Japan. ⁴Department of Immunology and Medical Zoology, Hyogo College of Medicine, Hyogo, Japan. ⁵Laboratory of Functional Analysis *In Silico*, Human Genome Center, Institute of Medical Science, University of Tokyo, Tokyo, Japan. ⁶Department of Molecular Microbiology and Immunology, Graduate School of Biomedical Sciences, Nagasaki University, Nagasaki, Japan. ⁷Department of Pathology, Hyogo College of Medicine, Hyogo, Japan. ⁸These authors contributed equally to this work. Correspondence should be addressed to S.A. (sakira@biken.osaka-u.ac.jp).

Received 19 March; accepted 22 July; published online 22 August 2010; doi:10.1038/ni.1920






Figure 1 *Jmjd3* is dispensable for induction of M1 macrophage polarization. (a–d) ELISAs showing production of TNF (a,c) and IL-6 (b,d) in resident PECs (a,b) or thioglycollate-elicited PECs (c,d) from wild-type (WT) or *Jmjd3*^{-/-} bone marrow chimeras. Med, control medium; ND, not determined. (e) CD11b and F4/80 expression in PECs harvested from WT or *Jmjd3*^{-/-} chimeras either untreated (resident Mφ), 3 d after treatment with thioglycollate (TEM), or 24 or 48 h after intraperitoneal infection with *L. monocytogenes*. Boxes and numbers indicate percentage of F4/80⁺CD11b⁺ cells in total PECs. (f) Numbers of F4/80⁺CD11b⁺ macrophages in PECs harvested from WT or *Jmjd3*^{-/-} chimeras either untreated, 3 d after thioglycollate treatment or 24 or 48 h after intraperitoneal infection with *L. monocytogenes*. (g) ELISAs showing concentrations of TNF, IL-6, IFN-γ and IL-12p40 in the sera of WT or *Jmjd3*^{-/-} chimeras infected with *L. monocytogenes*. (h) Quantitative PCR analysis showing expression of TNF, IL-6, IL-12p40 and iNOS mRNAs (relative to 18S rRNA) in PECs in mice infected with *L. monocytogenes*. Results are representative of four (c–f) or two (a,b,g,h) independent experiments (error bars indicate s.d.).

In addition to regulation by transcription factors, epigenetic regulation is essential for controlling proper gene expression¹⁶. Histone modifications, particularly at the N-terminal tails, and dynamic chromatin remodeling have been shown to be important for controlling sets of genes. In the case of histone modification, trimethylation of H3K4 is associated with active gene transcription, whereas trimethylation of H3K9, H3K27 and H4K20 is linked to silencing of gene expression^{16,21,22}. The methylation of H3K27 is mediated by the Polycomb repressive complex-2 (PRC2), composed of Ezh2, Suz12 and Eed (ref. 23). Proteins harboring a Jumonji-C (JmjC) domain, such as Jmj3, Utx and Uty, are known to act as H3K27 demethylases, catalyzing the conversion of H3K27me3 (trimethylated) to H3K27me1 (monomethylated)^{24–26}.

It has been reported that the expression of *Jmjd3* is induced in macrophages by TLR stimuli in an NF-κB-dependent fashion²⁶. *Jmjd3* has also been identified as an early TLR-inducible gene in mouse macrophages by microarray analysis²⁷. H3K27 trimethylation is implicated in the silencing of gene expression, and it has been shown that Jmj3 is recruited to transcription start sites (TSSs) that have abundant RNA polymerase II and H3K4me3 (ref. 28). Jmj3 is reported to fine-tune macrophage activation by controlling *Bmp2* and *Hox* expression²⁶. Furthermore, Jmj3 has been linked to the control of development through the regulation of *Hox*, and to oncogenesis through promotion of the expression of *Ink4a*^{29,30}. Here we report the role of Jmj3 *in vivo* in controlling M2 macrophage polarization and identify *Irf4* as a Jmj3 target gene crucial for the regulation of macrophages.

RESULTS

Generation of *Jmjd3*^{-/-} mice

To investigate the functional roles of Jmj3 in immune responses *in vivo*, we generated *Jmjd3*^{-/-} mice (Supplementary Fig. 1a,b). Reverse-transcription PCR analysis showed that the expression

of *Jmjd3* was abrogated in *Jmjd3*^{-/-} mouse embryonic fibroblasts (Supplementary Fig. 1c). *Jmjd3*^{-/-} mice died perinatally, and adult *Jmjd3*^{-/-} mice were not obtained (Supplementary Fig. 2a). Histological examination revealed that alveolar cell walls were thickened with tissues, and almost no air space was observed in lungs of *Jmjd3*^{-/-} neonates (Supplementary Fig. 2b), suggesting that the postnatal lethal phenotype of *Jmjd3*^{-/-} mice is due to premature development of lung tissues. To analyze the role of *Jmjd3* in hematopoietic cells, we obtained fetal liver cells from *Jmjd3*^{-/-} E15.5 embryos and established bone marrow-chimeric mice. Flow cytometry revealed that populations of T cells, B cells, conventional dendritic cells, plasmacytoid dendritic cells, natural killer cells, neutrophils, F4/80⁺CD11b⁺ macrophages and Ly6Ch⁺CD11b⁺ inflammatory monocytes in the spleen were similar between wild-type and *Jmjd3*^{-/-} chimeras (Supplementary Fig. 3). Proliferative responses of splenic T and B cells to mitogens and antigen receptor stimuli were also not altered (Supplementary Fig. 4). Furthermore, *Jmjd3*^{-/-} splenic T cells differentiated into either type 1 or type 2 helper T cells (T_H1 and T_H2) produced normal amounts of IFN-γ or IL-4, respectively (Supplementary Fig. 5). These results indicate that T cells lacking *Jmjd3* are capable of differentiating into T_H1 and T_H2 cells in response to cytokines.

Jmjd3 is dispensable for M1 macrophages

As *Jmjd3* is a TLR-inducible gene in macrophages, we first examined cytokine production of peritoneally resident and thioglycollate-elicited peritoneal exudate cells (PECs) in response to TLR ligands including lipopeptide (Pam₃CSK₄, a TLR2 ligand), lipopolysaccharide (LPS, a TLR4 ligand), imidazoquinoline analog (R-848, a TLR7 ligand) and oligonucleotide with a CpG motif (CpG-DNA, a TLR9 ligand). Production of TNF and IL-6 in response to TLR ligands did not differ between wild-type and *Jmjd3*^{-/-} tissue-resident macrophages and thioglycollate-elicited PECs (Fig. 1a–d). PECs elicited




Figure 2 Crucial role of *Jmjd3* in M2 macrophage polarization in response to chitin administration. (a,b) Expression of CD11b and either F4/80 (a) or Siglec-F (b) in PECs harvested from wild-type (WT) or *Jmjd3*^{-/-} chimeric mice 2 d after peritoneal injection with chitin. Quadrants and numbers in a indicate percentage of cells in each gate (F4/80⁺CD11b⁺, F4/80⁺CD11b⁻, F4/80⁻CD11b⁺ and F4/80⁻CD11b⁻); circles and numbers in b indicate percentage of Siglec-F⁺ eosinophils in total PECs. (c,d) Numbers of F4/80⁺CD11b⁺ macrophages (c) or Siglec-F⁺ eosinophils (d) in PECs harvested from WT or *Jmjd3*^{-/-} chimeric mice 2 d after peritoneal injection with chitin. (e) Cell types within PEC population from chitin-treated mice, stained with Diff-Quick in cytospin centrifuges. Mφ, macrophages; EO, eosinophils. (f) Surface MR expression on peritoneal F4/80⁺CD11b⁺ macrophages from chitin-treated mice. Graph shows MR expression on macrophages on horizontal axis. (g) Quantitative PCR showing expression of Arg1, Ym1, Fizz1, MR and iNOS mRNAs (relative to 18S rRNA). Total RNA was prepared from PECs 48 h after administration of chitin. **P* < 0.05; ***P* < 0.01 (two-tailed Student's *t*-test). Results are representative of five (a–d), two (e) or three (f,g) independent experiments (error bars indicate s.d.).

by peptone treatment of wild-type or *Jmjd3*^{-/-} chimeras also produced similar amounts of TNF and IL-6 in response to TLR ligand stimulation (Supplementary Fig. 6a). Flow cytometry revealed that the proportions of F4/80⁺CD11b⁺Gr1⁻ cells among resident and thioglycollate-elicited PECs did not differ between wild-type and *Jmjd3*^{-/-} chimeras (Fig. 1e and Supplementary Fig. 7). The total number of peritoneally resident macrophages, thioglycollate-elicited macrophages and peptone-elicited macrophages was not changed in *Jmjd3*^{-/-} chimeras (Fig. 1f and Supplementary Fig. 6b). We then examined the role of *Jmjd3* in M1 macrophage polarization in response to *Listeria monocytogenes* infection. When *L. monocytogenes* was inoculated intraperitoneally, production of proinflammatory cytokines in the sera was similar in wild-type and *Jmjd3*^{-/-} chimeras (Fig. 1g). F4/80⁺CD11b⁺Gr1⁺ macrophages, F4/80^{int}CD11b⁺Gr1^{hi} neutrophils, and F4/80⁻CD11b^{int}B220⁺ and F4/80⁻CD11b⁻B220⁺ B cells were examined in PECs prepared from *L. monocytogenes*-infected mice (Supplementary Fig. 8). The number of macrophages recruited to the peritoneal cavity was similar in wild-type and *Jmjd3*^{-/-} chimeras (Fig. 1e,f). Furthermore, the expression of genes encoding TNF, IL-6, IL-12p40 and iNOS in PECs were similarly upregulated (Fig. 1h). Collectively, these data suggest that *Jmjd3* is not involved in the generation and recruitment of M1 macrophages in response to inflammatory reagents and bacterial infection *in vivo*.

Jmjd3 is needed for M2 polarization to chitin

Chitin is a polymerized sugar and a structural component of helminths, arthropods and fungi³¹. It has been shown that chitin administration recruits macrophages with the M2 phenotype to the site of administration, which are important for subsequent

recruitment of eosinophils^{32,33}. Indeed, we found that intraperitoneal administration of chitin recruited F4/80⁺CD11b⁺ macrophages (Fig. 2a), Siglec-F⁺CCR3⁺CD4⁻ eosinophils and CD11b^{int}B220⁺ B cells to the peritoneal cavity after 48 h in wild-type mice (Fig. 2b and Supplementary Fig. 9). Whereas the number of chitin-elicited F4/80⁺CD11b⁺ macrophages was similar between wild-type and *Jmjd3*^{-/-} chimeras (Fig. 2c), the recruitment of eosinophils was severely impaired in *Jmjd3*^{-/-} chimeras (Fig. 2b–e). Furthermore, the expression of MR on chitin-elicited macrophages was severely impaired in *Jmjd3*^{-/-} chimeras (Fig. 2f). Chitin-elicited macrophages, but not eosinophils or B cells, expressed high levels of mRNA encoding Arg1, Ym1, Fizz1 and MR, the hallmarks of M2 macrophages (Supplementary Fig. 10). The expression of genes encoding Arg1, Ym1, Fizz1 and MR was considerably lower in chitin-induced PECs obtained from *Jmjd3*^{-/-} chimeras compared with wild-type controls, whereas expression of the gene encoding iNOS, associated with M1 macrophages, was not altered (Fig. 2g). Of note, the numbers of eosinophils circulating in the blood were similar in wild-type and *Jmjd3*^{-/-} chimeric mice, suggesting that eosinophil development was not impaired by *Jmjd3* deficiency (data not shown). Together, these results suggest *Jmjd3* is necessary for M2 macrophage polarization in response to chitin administration.

Role of *Jmjd3* in helminth infection

Next we investigated whether *Jmjd3* contributes to the responses against helminth infection *in vivo*. We used the *Nippostrongylus brasiliensis* infection model, which induces a strong type 2 immune response in the lung. Bronchoalveolar lavage (BAL) staining 5 and 13 d after infection revealed that macrophages were similarly recruited to the lung in wild-type and *Jmjd3*^{-/-} bone marrow chimeras (Fig. 3a,b).

