

Prevalence of Hepatitis C Virus Genotype 1a in Japan and Correlation of Mutations in the NS5A Region and Single-Nucleotide Polymorphism of Interleukin-28B With the Response to Combination Therapy With Pegylated-Interferon-Alpha 2b and Ribavirin

Kazuhiko Hayashi,¹ Yoshiaki Katano,^{1*} Teiji Kuzuya,¹ Yoshihiko Tachi,¹ Takashi Honda,¹ Masatoshi Ishigami,¹ Akihiro Itoh,¹ Yoshiki Hirooka,¹ Tetsuya Ishikawa,¹ Isao Nakano,¹ Fumihiro Urano,² Kentaro Yoshioka,³ Hidenori Toyoda,⁴ Takashi Kumada,⁴ and Hidemi Goto¹

¹Department of Gastroenterology, Nagoya University Graduate School of Medicine, Showa-ku, Nagoya, Japan

²Department of Gastroenterology, Toyohashi Municipal Hospital, Toyohashi, Japan

³Division of Liver and Biliary Diseases, Department of Internal Medicine, Fujita Health University, Kutsukake-cho, Toyoake, Japan

⁴Department of Gastroenterology, Ogaki Municipal Hospital, Ogaki, Japan

Hepatitis C virus (HCV) genotype 1a is rare in Japanese patients and the clinical characteristics of this genotype remain unclear. The interferon (IFN) sensitivity-determining region (ISDR) and single-nucleotide polymorphisms (SNPs) of interleukin-28B (IL28B) among patients with HCV genotype 1b are associated with IFN response, but associations among patients with genotype 1a are largely unknown. This study investigated the clinical characteristics of genotype 1a and examined whether genomic heterogeneity of the ISDR and SNPs of IL28B among patients with HCV genotype 1a affects response to combination therapy with pegylated-IFN- α 2b and ribavirin. Subjects comprised 977 patients infected with HCV genotype 1, including 574 men and 412 women (mean age, 55.2 ± 10.6 years). HCV was genotyped by direct sequencing of the 5'-untranslated region and/or core regions and confirmed by direct sequencing of the NS5A region. HCV genotypes 1a ($n = 32$) and 1b ($n = 945$) were detected. Twenty-three (71.9%) of the 32 patients with genotype 1a were patients with hemophilia who had received imported clotting factors. Prevalence of genotype 1a after excluding patients with hemophilia was thus 0.9%. Of the 23 patients with genotype 1a who completed IFN therapy, 11 (47.8%) were defined as achieving sustained virological response. Factors related to sustained virological response by univariate analysis were IL28B and ISDR. In conclusion,

HCV genotype 1a is rare in Japan. The presence of IL28B genotype TT, and more than two mutations, in the ISDR are associated with a good response to IFN therapy in patients with HCV genotype 1a. *J. Med. Virol.* **84:438–444, 2012.** © 2012 Wiley Periodicals, Inc.

KEY WORDS: hepatitis C virus; genotype 1a; NS5A; IL 28B; interferon

INTRODUCTION

Hepatitis C virus (HCV) is a member of the Flaviviridae family and causes chronic hepatitis that can develop into cirrhosis and hepatocellular carcinoma [Seeff, 2002]. HCV infection is a significant global health problem, affecting 170 million individuals worldwide. HCV can be divided into six genotypes and several subtypes according to genomic heterogeneity [Simmonds et al., 2005]. Each genotype shows a unique distribution and clinical characteristics such

All authors have nothing to disclose.

*Correspondence to: Yoshiaki Katano, MD, PhD, Department of Gastroenterology, Nagoya University Graduate School of Medicine, 65 Tsurumai-cho, Showa-ku, Nagoya 466-8550, Japan. E-mail: ykatano@med.nagoya-u.ac.jp

Accepted 23 November 2011

DOI 10.1002/jmv.23207

Published online in Wiley Online Library (wileyonlinelibrary.com).

as interferon (IFN) responsiveness [Ghany et al., 2009]. HCV genotypes 1b, 2a, and 2b are the major types encountered in Japan [Enomoto et al., 1990; Hayashi et al., 2003]. Genotype 1a is common worldwide, but is rare in Japan except among individuals with hemophilia who have received imported clotting factors [Fujimura et al., 1996; Otagiri et al., 2002; Hayashi et al., 2003]. The prevalence and clinical characteristics, including IFN responsiveness, of Japanese patients with HCV genotype 1a are unclear. HCV NS5A protein reportedly includes a domain associated with IFN response. This domain, located in the NS5A region of HCV genotype 1b, is closely associated with response to IFN therapy and is known as the IFN sensitivity-determining region (ISDR) [Enomoto et al., 1996]. IFN acts to inhibit viral replication by inducing double-stranded RNA-dependent protein kinase (PKR). The ISDR is located at the 5' end of the PKR-binding domain and is inhibited by PKR in vitro [Gale et al., 1998]. ISDR heterogeneity of genotype 1b is thus an important factor that may affect response to IFN [Enomoto et al., 1996; Nakano et al., 1999; Pascu et al., 2004; Hayashi et al., 2011a]. Several studies have reported a relationship between ISDR and IFN responsiveness among patients with HCV genotype 1a [Hofgärtner et al., 1997; Zeuzem et al., 1997; Kumthip et al., 2011; Yahoo et al., 2011]. However, this remains controversial for genotype 1a, and the utility of ISDR sequences for predicting IFN responsiveness has not been investigated for HCV genotype 1a in Japan due to the rarity of this genotype. Both genetic heterogeneity of the HCV genome and host genetics contribute to IFN responsiveness. Several genome-wide association studies have thus been performed to clarify host factors associated with IFN responsiveness, revealing that interleukin-28B (IL28B) polymorphisms are strongly associated with response to IFN therapy [Ge et al., 2009; Suppiah et al., 2009; Tanaka et al., 2009; Thomas et al., 2009]. Combined use of the single-nucleotide polymorphisms (SNPs) of IL28B and amino acid substitutions in the core region and ISDR could thus improve the prediction of response to IFN in patients with HCV genotype 1b [Akuta et al., 2011; Hayashi et al., 2011b; Kurosaki et al., 2011]. However, the effects of a combined evaluation of the SNPs of IL28B and amino acid substitutions in the ISDR in patients with HCV genotype 1a on IFN response are unclear. The aim of the present study was to determine whether genomic heterogeneity of the ISDR and SNPs of IL28B among patients with HCV genotype 1a affect response to combination therapy with pegylated-IFN- α 2b and ribavirin.

PATIENTS AND METHODS

A total of 977 patients (569 men, 408 women) with chronic hepatitis C genotype 1 and high viral load (<100 KIU/ml) who were treated at Nagoya University Hospital and affiliated hospitals were enrolled in

this study. Mean age of patients was 55.1 ± 12.2 years (range: 18–75 years). None of the patients had a history of chronic alcohol abuse, autoimmune disease, or metabolic disease. Patients with active intravenous drug use and immigrants were excluded from this study. The core region (aa 30–110) and ISDR (aa 2,209–2,248) of HCV were examined by direct sequencing. SNPs of IL28B (rs8099917) were identified using a real-time polymerase chain reaction (PCR) system. Patients received subcutaneous injections of pegylated-IFN- α 2b (1.5 μ g/kg) once each week along with oral ribavirin (600 mg/day for patients <60 kg, 800 mg/day for 60–80 kg, 1,000 mg/day for >80 kg) for 48 weeks. Patients who became negative for HCV-RNA between 16 and 36 weeks after initiating IFN treatment had the IFN treatment extended to 72 weeks, in accordance with Japanese guidelines [Kumada et al., 2010]. HCV-RNA levels in serum samples were examined at 12 weeks, at the end of IFN therapy, and at 6 months after the end of treatment. Serum was stored at -80°C for virological examination at pretreatment. Early virological response was defined as HCV-negative status at 12 weeks. Patients who were persistently negative for serum HCV-RNA at 24 weeks after withdrawal of IFN treatment were considered to show sustained virological response. Written informed consent was obtained from each patient, and the study protocol conformed to the ethical guidelines of the 1975 Declaration of Helsinki.

Virological Analysis

HCV-RNA quantitative viremia load was determined by PCR. HCV was genotyped by direct sequencing of the 5'-untranslated region and/or core regions as described previously and confirmed by direct sequencing of the NS5A region [Otagiri et al., 2002; Dal Pero et al., 2007; Hayashi et al., 2011a]. Genotypes were classified according to the nomenclature proposed by Simmonds et al. [2005]. Direct sequencing of the core and NS5A-ISDR regions was performed as reported previously [Dal Pero et al., 2007; Hayashi et al., 2011a]. In brief, RNA was extracted from 140 μ l of serum using a commercial kit (QIAamp Viral RNA Kit; Qiagen, Valencia, CA) and dissolved in 50 μ l of diethylpyrocarbonate-treated water. RNA (10 ng) was used for reverse transcription with oligos and random hexamer primers with a commercial kit (iScript cDNA Synthesis Kit; Bio-Rad, Hercules, CA). The HCV core region and NS5A-ISDR were amplified by nested PCR. In brief, each 50- μ l PCR reaction mixture contained 100 nM of each primer, 1 ng of template cDNA, 5 μ l of GeneAmp 10 \times PCR buffer, 2 μ l of dNTPs, and 1.25 U of AmpliTaq Gold (Applied Biosystems, Foster City, CA). Primers for the core region were: sense, 5'-GGGAGGTCTCGTAGACCGTGCAC-CATG-3' and antisense, 5'-GAGMGGKATRTACCC-CATGAGRTCGGC-3'. Primers for the NS5A-ISDR were: sense, 5'-GCCTGGAGCCCTTGATGTC-3' and

TABLE I. Clinical Characteristic of Patients With HCV Genotype 1a

	N = 32
Age (y.o.)	36.4 ± 2.2
Sex: male/female	28/4
AST (IU/L)	48.8 ± 33.6
ALT (IU/L)	64.6 ± 57.8
Platelet (10 ⁴ /μl)	18.8 ± 6.0
HCV RNA level (KIU/ml)	2607.4 ± 3072.2
Source (clotting factor/BTF/unknown)	23/2/7

AST, aspartate aminotransferase; ALT, alanine aminotransferase; HCV, hepatitis C virus.

antisense, 5'-CTGCGTGAAGTGGTGAATAC-3'. Amplification conditions consisted of 10 min at 94°C, followed by 40 cycles of 94°C for 10 sec, 55°C for 30 sec, and 72°C for 30 sec in a thermal cycler (GeneAmp PCR System 9700; Applied Biosystems). The second PCR was performed using the same reaction buffer with the first-round PCR product as template, and the following sets of primers: for the core region, sense primer 5'-AGACCGTGCACCATGAGCAC-3' and antisense 5'-TACGCCGGGGGTCAKTRGGGCCCA-3'; and for the NS5A-ISDR, sense 5'-TGTTTCCCCACGCACTAC-3' and antisense 5'-TGATGGGCAGTTTT-TGTTCTTC-3'. PCR products were separated by electrophoresis on 2% agarose gels, stained with ethidium bromide, and visualized under ultraviolet light. PCR products were then purified and sequenced with the second-round PCR primers using a dye terminator sequencing kit (BigDye Terminator v1.1 Cycle Sequencing Kit; Applied Biosystems) and an ABI 310 DNA Sequencer (Applied Biosystems).

Genotyping Analysis

Detection of SNPs for IL28B (rs8099917) was conducted using a real-time PCR system. In brief, genomic DNA was extracted from 150 μl of whole blood with a commercial kit (QIAamp DNA Blood mini Kit; Qiagen) and dissolved in 50 μl of diethylpyrocarbonate-treated water. DNA (10 ng) was used for PCR and genotyping of IL28B SNP (rs8099917) was performed by TaqMan allelic discrimination (ABI-Prism 7300 SDS software; Applied Biosystems) with TaqMan SNP Genotyping Assays provided by Applied Biosystems (C_11710096_10).

Statistical Analysis

Data are expressed as mean ± standard deviation (SD). The paired *t*-test was used to analyze differences in variables. A value of *P* < 0.05 was considered statistically significant. Statview 5.0 software (SAS Institute, Cary, NC) was used for all analyses.

RESULTS

Thirty-two of the 977 patients (3.3%) were infected by genotype 1a. Clinical characteristics of patients with genotype 1a are summarized in Table I. Twenty-three cases involved patients with hemophilia who had received imported clotting factors. The prevalence of genotype 1a after excluding patients with hemophilia was 0.9%. A comparison of clinical characteristics according to hemophilia status is shown in Table II. No significant differences were apparent among the two groups. Differences in clinical characteristics between genotypes 1a and 1b are shown in Table III. Males were more frequent among patients with genotype 1a (87.5%) than among those with genotype 1b (57.2%), as the majority of patients with genotype 1a were young male patients with hemophilia. Sequence alignments of the core region at codons 71 and 90 showed arginine and cysteine, respectively, in all patients. The HCV core region of genotype 1a was thus well-conserved, with no significant mutations at codons 71 or 90. This is not similar to previous findings for genotype 1b [Akuta et al., 2005, 2011; Hayashi et al., 2011a,b; Kurosaki et al., 2011]. Alignment of the amino acid sequence for NS5A-ISDR is shown in Figure 1. The sequence of the HCV-1 strain was defined as the consensus sequence of genotype 1a, and the number of mutations to the chosen consensus sequence in ISDR was used to analyze the ISDR system. Sequences of the HCV-1 strain and HCV-1 strain with only one amino acid substitution were defined as wild-type, while ISDR sequences with more than two amino acid substitutions were defined as mutant-type. Twenty-seven strains were defined as wild-type and 5 strains were defined as mutant-type. IL28B genotypes could be obtained for 25 patients, and IL28B alleles were TT (n = 14) and TG (n = 11). Twenty-three patients received pegylated-IFN-α2b plus ribavirin therapy. Twenty patients were treated for 48 weeks, and 1 patient was treated for 72 weeks. Two patients were withdrawn at 24 weeks due to a

TABLE II. Clinical Characteristic According to Hemophilia

	Patients with hemophilia (N = 23)	Patients without hemophilia (N = 9)	<i>P</i> -value
Age (y.o.)	37.1 ± 9.2	37.1 ± 16.3	0.9966
Sex: male/female	22/1	6/3	0.0572
AST (IU/L)	51.2 ± 34.8	41.9 ± 30.9	0.5072
ALT (IU/L)	68.2 ± 55.8	54.0 ± 66.1	0.5566
Platelet (10 ⁴ /μl)	18.4 ± 6.8	19.8 ± 3.0	0.5602
HCV levels (KIU/ml)	2599.6 ± 3108.0	2630.0 ± 3176.5	0.9812

AST, aspartate aminotransferase; ALT, alanine aminotransferase; HCV, hepatitis C virus.

TABLE III. Clinical Characteristic According to Genotypes

	Genotype 1a (N = 32)	Genotype 1b (N = 945)	P-value
Age (y.o.)	36.4 ± 2.2	55.9 ± 11.6	0.0001
Sex: male/female	28/4	546/408	0.0004
Patients with hemophilia	23	4	0.0001
AST (IU/L)	48.8 ± 33.6	59.9 ± 45.0	0.1745
ALT (IU/L)	64.6 ± 57.8	64.6 ± 57.8	0.9894
Platelet (10 ⁴ /μl)	18.8 ± 6.0	17.2 ± 6.0	0.0918
HCV levels (KIU/ml)	2607.4 ± 3072.2	2011.5 ± 1453.8	0.0642

AST, aspartate aminotransferase; ALT, alanine aminotransferase; PLT, platelet count; HCV, hepatitis C virus.

lack of response to IFN therapy. Frequency of early virological response, characterized by undetectable HCV at 12 weeks, was 30.4% (7/23). Virological response rate at the end of treatment was 47.8% (11/23). Finally, 11 of 23 patients (47.8%) achieved sustained virological response. Clinical characteristics were compared between patients who achieved sustained virological response and patients who did not (Table IV), revealing significant differences in two factors on univariate analysis: IL28B and ISDR.

DISCUSSION

The present study investigated 977 patients with genotype 1 using direct sequencing of core and NS5A regions, revealing that genotype 1a is rare (3.3%) in

Japan. Of the 33 patients with genotype 1a, 23 (71.9%) were patients with hemophilia, confirming that the majority of cases with genotype 1a involve patients with hemophilia who have received imported clotting factors, as previously reported [Fujimura et al., 1996; Otagiri et al., 2002; Hayashi et al., 2003]. Analysis after excluding patients with hemophilia revealed the prevalence of genotype 1a in Japan was 0.9% (9/954). Recently, the distributions of HBV genotypes have been changing in Japan due to international exchange [Hayashi et al., 2007; Matsuura et al., 2009]. However, prevalences of HCV genotypes have remained stable because of the different modes of infection involved. The present study revealed that 11 (47.8%) of 23 patients achieved sustained virological response. The IFN responsiveness of HCV genotype 1a in Japanese patients was reported in 1999 from Okinawa, a far southern island in Japan [Sakugawa et al., 1997]. That study reported that the rate of sustained virological response tended to be higher in patients with genotype 1a than in those with genotype 1b, but no significant differences were identified because of the small number of patients with genotype 1a. Low virological response rates in both genotypes 1a and 1b were confirmed in the present Japanese patients, as in Caucasian patients [Manns et al., 2001; McHutchison et al., 2009]. No significant differences in sustained virological response rate were seen between genotypes 1a and 1b. Discriminating between genotypes 1a and 1b thus seems to have little clinical relevance in terms of IFN responsiveness. Viral factors associated with sustained virological response, including HCV genotype, have been studied most frequently and mutations in the core and NS5A regions of HCV genotype 1b have been associated with response to IFN therapy [Akuta et al., 2005, 2010, 2011; Okanoue et al., 2009; Nakagawa et al., 2010; Toyoda et al., 2010; Hayashi et al., 2011a; Hayes et al., 2011; Kumthip et al., 2011; Kurosaki et al., 2011]. These viral factors could improve prediction of sustained virological response for genotype 1a, as in 1b. Amino acid substitutions at positions 70 and 91 of the HCV core region in genotype 1b have been related to IFN responsiveness, liver steatosis, hepatic oxidative stress, insulin resistance, and carcinogenesis [Akuta et al., 2005, 2007, 2009; Tachi et al., 2010]. These substitutions may have substantial impacts on




Fig. 1. Alignment of the amino acid sequence for the NS5A-ISDR. In the sequence alignment, dashes indicate amino acids identical to consensus sequence HCV1. Sequences of the HCV1 strain and HCV1 strains with one-nucleotide substitutions were defined as wild-type ISDR, and all other strains were defined as mutant-type ISDR. ISDR, interferon sensitivity-determining region.

TABLE IV. Univariate Analysis: Factors Predictive of Sustained Virologic Response

Factors	Sustained virologic response (n = 11)	Non-sustained virologic response (n = 12)	P-value
Age (y.o.)	37.9 ± 10.9	39.8 ± 11.3	0.6958
Gender: male/female	10/1	10/2	0.9999
ALT (IU/L)	78.2 ± 50.8	62.6 ± 68.1	0.5435
AST (IU/L)	51.4.4 ± 29.2	48.8 ± 40.4	0.8616
PLT ($\times 10^3/\text{mm}^3$)	19.0 ± 5.4	19.3 ± 5.7	0.8870
HCV RNA level (KIU/ml)	1323.1 ± 1077.3	2567.0 ± 2940.8	0.2481
ISDR: wild/mutant	7/4	12/0	0.0373
IL28B:TT/TG	9/1	4/8	0.0115

AST, aspartate aminotransferase; ALT, alanine aminotransferase; PLT, platelet count; HCV, hepatitis C virus; ISDR, interferon sensitivity-determining region; IL28B, interleukin 28B.

the pathogenesis of HCV genotype 1a infection. However, the HCV core region of genotype 1a is well-conserved and no significant mutations were seen in the core region, which is associated with IFN responsiveness. Several reports have also found that the HCV core region, including positions 70 and 91, of HCV genotype 1a is highly conserved [Alestig et al., 2011; Kumthip et al., 2011]. Mutations in the core region of genotype 1a would be rare, so this region might be unsuitable for routine clinical use, unlike in genotype 1b. However, the number of patients in this study was small, and large studies including from other countries are needed to clarify these issues. The ISDR in the NS5A region of HCV genotype 1b is closely associated with response to IFN therapy. ISDR mutations of genotype 1b are well known to be more important in predicting sustained virological response in Japanese patients than European patients [Hofgärtner et al., 1997; Zeuzem et al., 1997; Nakano et al., 1999; Pascu et al., 2004; Hayashi et al., 2011a]. European studies have failed to detect the specific amino acid substitutions in ISDR of genotype 1a associated with IFN responsiveness [Hofgärtner et al., 1997; Zeuzem et al., 1997]. In this study, sustained virological response was achieved in 36.8% of patients with wild-type ISDR and 100% of patients with mutant-type ($P = 0.0373$). The present analysis showed a close relationship between ISDR of genotype 1a and sustained virological response, as in genotype 1b. Recent investigations in Thailand and Iran have failed to identify the usefulness of ISDR for HCV genotype 1a in predicting sustained virological response [Kumthip et al., 2011; Yahoo et al., 2011]. The high virological response rate and low prevalence of patients with mutations in the ISDR do not favor the use of ISDR analysis in predicting IFN responsiveness [Herion and Hoofnagle, 1997; Yokozaki et al., 2011]. Rates of sustained virological response among these studies were much higher than those in the present study (68.4% and 75% vs. 47.8%). The mean number of mutations in patients who achieved sustained virological response in the studies by Kumthip et al. [2011] and Yahoo et al. [2011], and the present group were 1.4, 1.4, and 1.6, respectively. Differences in sustained virological response and the number of mutations to the ISDR might underpin this discrepancy in the evaluation of ISDR. Although the sample size in

the present study was small, the results indicate that ISDR represents a strong indicator of progression to sustained virological response for patients with HCV genotype 1a. Amino acid substitutions in the ISDR of genotype 1a thus also play an important role in predicting sustained virological response in Japanese patients compared to patients from other countries. IL28B polymorphisms such as host genetics, as well as mutations in the HCV genome, contribute to IFN treatment outcomes. Rates of sustained virological response in patients in this study with TT and TG were 69.2% and 11.1%, respectively. The TG allele of the IL28B genotype was significantly associated with poor response to IFN therapy ($P = 0.0115$). SNPs of IL28B would regulate the expression of IFN-stimulated genes and affect IFN responsiveness. IL28B and ISDR thus exert independent effects on IFN responsiveness and both host and viral factors impacting IFN responsiveness would improve the prediction of sustained virological response. Several studies have thus reported that both the SNP of IL28B and mutations in the ISDR were associated with sustained virological response in patients with HCV genotype 1b [Akuta et al., 2011; Hayashi et al., 2011b; Kurosaki et al., 2011]. In the present study of HCV genotype 1a, among the 9 patients who had simultaneously the TG allele for IL28B and wild-type ISDR, only 1 achieved sustained virological response (11.1%). The best-sustained virological response was achieved in patients with mutant-type ISDR and the T allele (100%). The combination of SNPs for IL28B and mutations in ISDR may thus predict response to IFN therapy in patients with HCV genotype 1a as well as genotype 1b. Given the small sample size in this investigation, larger cohorts are needed to confirm the present results. Furthermore, infection with genotype 1a in Japanese patients is rare, making large-scale studies difficult to perform.

In conclusion, the prevalence of HCV genotype 1a is rare in Japan and the majority of cases involve patients with hemophilia. The TG genotype of IL28B is associated with poor response, while mutant-type ISDR is associated with good response to combination therapy with pegylated-IFN- α 2b and ribavirin in patients with HCV genotype 1a. Combined use of both IL28B and ISDR could improve the prediction of IFN response.

REFERENCES

- Akuta N, Suzuki F, Sezaki H, Suzuki Y, Hosaka T, Someya T, Kobayashi M, Saitoh S, Watahiki S, Sato J, Matsuda M, Kobayashi M, Arase Y, Ikeda K, Kumada H. 2005. Association of amino acid substitution pattern in core protein of hepatitis C virus genotype 1b high viral load and non-virological response to interferon-ribavirin combination therapy. *Intervirology* 48:372–380.
- Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Arase Y, Ikeda K, Kumada H. 2007. Amino acid substitutions in the hepatitis C virus core region are the important predictor of hepatocarcinogenesis. *Hepatology* 46:1357–1364.
- Akuta N, Suzuki F, Hirakawa M, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Saitoh S, Arase Y, Ikeda K, Kumada H. 2009. Amino acid substitutions in the hepatitis C virus core region of genotype 1b are the important predictor of severe insulin resistance in patients without cirrhosis and diabetes mellitus. *J Med Virol* 81:1032–1039.
- Akuta N, Suzuki F, Hirakawa M, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Saitoh S, Arase Y, Ikeda K, Chayama K, Nakamura Y, Kumada H. 2010. Amino acid substitution in hepatitis C virus core region and genetic variation near the interleukin 28B gene predict viral response to telaprevir with peginterferon and ribavirin. *Hepatology* 52:421–429.
- Akuta N, Suzuki F, Hirakawa M, Kawamura Y, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Saitoh S, Arase Y, Ikeda K, Chayama K, Nakamura Y, Kumada H. 2011. Amino acid substitution in HCV core/NS5A region and genetic variation near IL28B gene affect treatment efficacy to interferon plus ribavirin combination therapy. *Intervirology* (in press).
- Alestig E, Arnholm B, Eilard A, Lagging M, Nilsson S, Norström G, Wahlberg T, Wejstål R, Westin J, Lindh M. 2011. Core mutations, IL28B polymorphisms and response to peginterferon/ribavirin treatment in Swedish patients with hepatitis C virus genotype 1 infection. *BMC Infect Dis* 12:124.
- Dal Pero F, Tang KH, Gerotto M, Bortoletto G, Paulon E, Herrmann E, Zeuzem S, Alberti A, Naoum NV. 2007. Impact of NS5A sequences of hepatitis C virus genotype 1a on early viral kinetics during treatment with peginterferon-alpha 2a plus ribavirin. *J Infect Dis* 196:998–1005.
- Enomoto N, Takada A, Nakao T, Date T. 1990. There are two major types of hepatitis C virus in Japan. *Biochem Biophys Res Commun* 170:1021–1025.
- Enomoto N, Sakuma I, Asahina Y, Kurosaki M, Murakami T, Yamamoto C, Ogura Y, Izumi N, Marumo F, Sato C. 1996. Mutations in the nonstructural protein 5A gene and response to interferon in patients with chronic hepatitis C virus 1b infection. *N Engl J Med* 334:77–81.
- Fujimura Y, Ishimoto S, Shimoyama T, Narita N, Kuze Y, Yoshioka A, Fukui H, Tanaka T, Tsuda F, Okamoto H, Miyakawa Y, Mayumi M. 1996. Genotypes and multiple infections with hepatitis C virus in patients with haemophilia A in Japan. *J Viral Hepat* 3:79–84.
- Gale M, Jr., Blakely CM, Kwieciszewski B, Tan SL, Dossett M, Tang NM, Korth MJ, Polyak SJ, Gretch DR, Katze MG. 1998. Control of PKR protein kinase by hepatitis C virus nonstructural 5A protein: Molecular mechanisms of kinase regulation. *Mol Cell Biol* 18:5208–5218.
- Ge D, Fellay J, Thompson AJ, Simon JS, Shianna KV, Urban TJ, Heinzen EL, Qiu P, Bertelsen AH, Muir AJ, Sulkowski M, McHutchison JG, Goldstein DB. 2009. Genetic variation in IL28B predicts hepatitis C treatment-induced viral clearance. *Nature* 461:399–401.
- Ghany MG, Strader DB, Thomas DL, Seeff LB, American Association for the Study of Liver Diseases. 2009. Diagnosis, management, and treatment of hepatitis C: An update. *Hepatology* 49:1335–1374.
- Hayashi K, Fukuda Y, Nakano I, Katano Y, Toyoda H, Yokozaki S, Hayakawa T, Morita K, Nishimura D, Kato K, Urano F, Takamatsu J. 2003. Prevalence and characterization of hepatitis C virus genotype 4 in Japanese hepatitis C carriers. *Hepatitis Res* 25:409–414.
- Hayashi K, Katano Y, Takeda Y, Honda T, Ishigami M, Itoh A, Hirooka Y, Nakano I, Yano M, Goto H, Yoshioka K, Toyoda H, Kumada T. 2007. Comparison of hepatitis B virus subgenotypes in patients with acute and chronic hepatitis B and absence of lamivudine-resistant strains in acute hepatitis B in Japan. *J Med Virol* 79:366–373.
- Hayashi K, Katano Y, Ishigami M, Itoh A, Hirooka Y, Nakano I, Urano F, Yoshioka K, Toyoda H, Kumada T, Goto H. 2011a. Mutations in the core and NS5A region of hepatitis C virus genotype 1b and correlation with response to pegylated-interferon-alpha 2b and ribavirin combination therapy. *J Viral Hepat* 18:280–286.
- Hayashi K, Katano Y, Honda T, Ishigami M, Itoh A, Hirooka Y, Ishikawa T, Nakano I, Yoshioka K, Toyoda H, Kumada T, Goto H. 2011b. Association of interleukin 28B and mutations in the core and NS5A region of hepatitis C virus with response to peg-interferon and ribavirin therapy. *Liver Int* 9:1359–1365.
- Hayes CN, Kobayashi M, Akuta N, Suzuki F, Kumada H, Abe H, Miki D, Imamura M, Ochi I, Kamatani N, Nakamura Y, Chayama K. 2011. HCV substitutions and IL28B polymorphisms on outcome of peg-interferon plus ribavirin combination therapy. *Gut* 60:261–267.
- Herion D, Hoofnagle JH. 1997. The interferon sensitivity determining region: All hepatitis C virus isolates are not the same. *Hepatology* 25:769–770.
- Hofgärtner WT, Polyak SJ, Sullivan DG, Carithers RL, Jr., Gretch DR. 1997. Mutations in the NS5A gene of hepatitis C virus in North American patients infected with HCV genotype 1a or 1b. *J Med Virol* 53:118–126.
- Kumada H, Okanoue T, Onji M, Moriwaki H, Izumi N, Tanaka E, Chayama K, Sakisaka S, Takehara T, Oketani M, Suzuki F, Toyota J, Nomura H, Yoshioka K, Seike M, Yotsuyanagi H, Ueno Y, The Study Group for the Standardization of Treatment of Viral Hepatitis Including Cirrhosis, Ministry of Health, Labour and Welfare of Japan. 2010. Guidelines for the treatment of chronic hepatitis and cirrhosis due to hepatitis C virus infection for the fiscal year 2008 in Japan. *Hepatology* 40:8–13.
- Kumthip K, Pantip C, Chusri P, Thongsawat S, O'Brien A, Nelson KE, Maneekarn N. 2011. Correlation between mutations in the core and NS5A genes of hepatitis C virus genotypes 1a, 1b, 3a, 3b, 6f and the response to pegylated interferon and ribavirin combination therapy. *J Viral Hepat* 18:e117–e125.
- Kurosaki M, Tanaka Y, Nishida N, Sakamoto N, Enomoto N, Honda M, Sugiyama M, Matsuura K, Sugauchi F, Asahina Y, Nakagawa M, Watanabe M, Sakamoto M, Maekawa S, Sakai A, Kaneko S, Ito K, Masaki N, Tokunaga K, Izumi N, Mizokami M. 2011. Pre-treatment prediction of response to pegylated-interferon plus ribavirin for chronic hepatitis C using genetic polymorphism in IL28B and viral factors. *J Hepatol* 54:439–448.
- Manns MP, McHutchison JG, Gordon SC, Rustgi VK, Shiffman M, Reindollar R, Goodman ZD, Koury K, Ling M, Albrecht JK. 2001. Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment of chronic hepatitis C: A randomised trial. *Lancet* 358:958–965.
- Matsuura K, Tanaka Y, Hige S, Yamada G, Murawaki Y, Komatsu M, Kuramitsu T, Kawata S, Tanaka E, Izumi N, Okuse C, Kakumu S, Okanoue T, Hino K, Hiasa Y, Sata M, Maeshiro T, Sugauchi F, Nojiri S, Joh T, Miyakawa Y, Mizokami M. 2009. Distribution of hepatitis B virus genotypes among patients with chronic infection in Japan shifting toward an increase of genotype A. *J Clin Microbiol* 47:1476–1483.
- McHutchison JG, Lawitz EJ, Shiffman ML, Muir AJ, Galler GW, McCone J, Nyberg LM, Lee WM, Ghalib RH, Schiff ER, Galati JS, Bacon BR, Davis MN, Mukhopadhyay P, Koury K, Noviello S, Pedicone LD, Brass CA, Albrecht JK, Sulkowski MS, IDEAL Study Team. 2009. Peginterferon alfa-2b or alfa-2a with ribavirin for treatment of hepatitis C infection. *N Engl J Med* 361:580–593.
- Nakagawa M, Sakamoto N, Ueyama M, Mogushi K, Nagaie S, Itsui Y, Azuma S, Kakinuma S, Tanaka H, Enomoto N, Watanabe M. 2010. Mutations in the interferon sensitivity determining region and virological response to combination therapy with pegylated-interferon alpha 2b plus ribavirin in patients with chronic hepatitis C-1b infection. *J Gastroenterol* 45:656–665.
- Nakano I, Fukuda Y, Katano Y, Nakano S, Kumada T, Hayakawa T. 1999. Why is the interferon sensitivity-determining region (ISDR) system useful in Japan? *J Hepatol* 30:1014–1022.
- Okanoue T, Itoh Y, Hashimoto H, Yasui K, Minami M, Takehara T, Tanaka E, Onji M, Toyota J, Chayama K, Yoshioka K, Izumi N, Akuta N, Kumada H. 2009. Predictive values of amino acid sequences of the core and NS5A regions in antiviral therapy for

- hepatitis C: A Japanese multi-center study. *J Gastroenterol* 44:952–963.
- Otagiri H, Fukuda Y, Nakano I, Katano Y, Toyoda H, Yokozaki S, Hayashi K, Hayakawa T, Fukuda Y, Kinoshita M, Takamatsu J. 2002. Evaluation of a new assay for hepatitis C virus genotyping and viral load determination in patients with chronic hepatitis C. *J Virol Methods* 103:137–143.
- Pascu M, Martus P, Höhne M, Wiedenmann B, Hopf U, Schreiber E, Berg T. 2004. Sustained virological response in hepatitis C virus type 1b infected patients is predicted by the number of mutations within the NS5A-ISDR: A meta-analysis focused on geographical differences. *Gut* 53:1345–1351.
- Sakugawa H, Nakasone H, Kinjo F, Saito A, Keida Y, Kikuchi K, Oyadomari Y, Ishihara M, Nakasone K, Yogi S, Kinjo Y, Taira M. 1997. Clinical features of patients with chronic liver disease associated with hepatitis C virus genotype 1a/I in Okinawa, Japan. *J Gastroenterol Hepatol* 12:176–181.
- Seeff LB. 2002. Natural history of chronic hepatitis C. *Hepatology* 36:S35–S46.
- Simmonds P, Bukh J, Combet C, Deléage G, Enomoto N, Feinstone S, Halfon P, Inchauspé G, Kuiken C, Maertens G, Mizokami M, Murphy DG, Okamoto H, Pawlotsky JM, Penin F, Sablon E, Shin-I T, Stuyver LJ, Thiel HJ, Viazov S, Weiner AJ, Widell A. 2005. Consensus proposals for a unified system of nomenclature of hepatitis C virus genotypes. *Hepatology* 42:962–973.
- Suppiah V, Moldovan M, Ahlenstiel G, Berg T, Weltman M, Abate ML, Bassendine M, Spengler U, Dore GJ, Powell E, Riordan S, Sheridan D, Smedile A, Fragomeli V, Müller T, Bahlo M, Stewart GJ, Booth DR, George J. 2009. IL28B is associated with response to chronic hepatitis C interferon-alpha and ribavirin therapy. *Nat Genet* 41:1100–1104.
- Tachi Y, Katano Y, Honda T, Hayashi K, Ishigami M, Itoh A, Hirooka Y, Nakano I, Samejima Y, Goto H. 2010. Impact of amino acid substitutions in the hepatitis C virus genotype 1b core region on liver steatosis and hepatic oxidative stress in patients with chronic hepatitis C. *Liver Int* 30:554–559.
- Tanaka Y, Nishida N, Sugiyama M, Kurosaki M, Matsuura K, Sakamoto N, Nakagawa M, Korenaga M, Hino K, Hige S, Ito Y, Mita E, Tanaka E, Mochida S, Murawaki Y, Honda M, Sakai A, Hiasa Y, Nishiguchi S, Koike A, Sakaida I, Imamura M, Ito K, Yano K, Masaki N, Sugauchi F, Izumi N, Tokunaga K, Mizokami M. 2009. Genome-wide association of IL28B with response to pegylated interferon-alpha and ribavirin therapy for chronic hepatitis C. *Nat Genet* 41:1105–1109.
- Thomas DL, Thio CL, Martin MP, Qi Y, Ge D, O'Huigin C, Kidd J, Kidd K, Khakoo SI, Alexander G, Goedert JJ, Kirk GD, Donfield SM, Rosen HR, Tobler LH, Busch MP, McHutchison JG, Goldstein DE, Carrington M. 2009. Genetic variation in IL28B and spontaneous clearance of hepatitis C virus. *Nature* 461:798–801.
- Toyoda H, Kumada T, Tada T, Arakawa T, Hayashi K, Honda T, Katano Y, Goto H. 2010. Association between HCV amino acid substitutions and outcome of peginterferon and ribavirin combination therapy in HCV genotype 1b and high viral load. *J Gastroenterol Hepatol* 25:1072–1078.
- Yahoo N, Sabahi F, Shahzamani K, Malboobi MA, Jabbari H, Sharifi H, Mousavi-Fard SH, Merat S. 2011. Mutations in the E2 and NS5A regions in patients infected with hepatitis C virus genotype 1a and their correlation with response to treatment. *J Med Virol* 83:1332–1337.
- Yokozaki S, Katano Y, Hayashi K, Ishigami M, Itoh A, Hirooka Y, Nakano I, Goto H. 2011. Mutations in two PKR-binding domains in chronic hepatitis C of genotype 3a and correlation with viral loads and interferon responsiveness. *J Med Virol* 83:1727–1732.
- Zeuzem S, Lee JH, Roth WK. 1997. Mutations in the nonstructural 5A gene of European hepatitis C virus isolates and response to interferon alfa. *Hepatology* 25:740–744.

Favorable association between genetic polymorphisms near the *IL28B* gene and hepatic steatosis: Direct or indirect?

To the Editor,

We read with great interest the article by Tillmann *et al.* [1] that investigated the association between *IL28B* polymorphisms and hepatic steatosis, both of which are associated with a response to combination therapy with peginterferon (PEG-IFN) and ribavirin. It will provide new insight into the role of *IL28B* polymorphisms on the resistance to combination therapy against hepatitis C virus (HCV) infection.

Previous studies have reported the influence of amino acid substitutions at residue 70 of the HCV core region (from arginine to glutamine or histidine) on the resistance to combination therapy with PEG-IFN and ribavirin in patients infected with HCV genotype 1b [2–4]. In addition, prior studies have identified an association between amino acid substitutions at HCV core 70 and hepatic steatosis [5,6]. The percentage of patients with the mutant amino acid at residue 70 of the HCV core region increases with the progression of chronic hepatitis, suggesting that the mutation of the amino acid at residue 70 occurs during the natural course of chronic HCV infection [7]. Several recent studies have reported a lower prevalence of mutant amino acids at HCV core 70 in patients who have the *IL28B* polymorphism that is associated with a favorable response to combination therapy with PEG-IFN and ribavirin (i.e., CC genotype of rs12979860 and TT genotype of rs8099917) than in patients who have an unfavorable genotype [8,9]. These reports suggest that the mutation frequency of the HCV core 70 amino acid may differ according to the genetic polymorphism near the *IL28B* gene.

We analyzed polymorphisms of rs8099917 that corresponded to those of rs12979860, the rate of which is more than 99% of individuals of Japanese ethnicity [10], amino acid substitutions at HCV core 70, and hepatic steatosis based on biopsy specimens, which were obtained just prior to the start of the therapy and evaluated with the same criteria used by Tillmann *et al.*, in our 122 Japanese Mongolian patients infected with HCV genotype 1b. We found higher likelihoods of sustained virologic response in patients with the TT genotype of rs8099917, in patients with arginine at residue 70 of the HCV core region, and in patients without steatosis. We did not find significant association between *IL28B* polymorphisms and hepatic steatosis (absence of steatosis: TT genotype, 66 out of 85 (77.6%) vs. TG/GG genotype, 22 out of 37 (59.5%), $p = 0.0658$). We found significant associations between *IL28B* polymorphisms and the amino acid at residue 70 of the HCV core region (patients with arginine at HCV core 70: TT genotype, 71 out of 85 (83.5%) vs. TG/GG genotype, 14 out of 37 (37.8%), $p < 0.0001$) and also between the amino acid at HCV core 70 and hepatic steatosis (absence of steatosis: arginine at HCV core 70, 73 of 85 (85.9%) vs. glutamine/histidine at HCV core 70, 15 of 37 (40.5%), $p < 0.0001$). These associations may indicate that the polymorphisms near the *IL28B* gene may influence the mutation of the amino acid at residue 70 of the HCV core region, and that the amino acid mutation at HCV

core 70 may influence hepatic steatosis over the course of chronic HCV infection.

There are differences between our Japanese Mongolian population and the population studied by Tillmann *et al.* including ethnicity, the rate of correspondence between rs12979860 and rs8099917, the frequency of the mutation at residue 70 of the HCV core region, and the rate of hepatic steatosis. Furthermore, the number of patients infected with HCV genotype 1b in the Tillmann *et al.* cohort is unknown. Despite these facts, it would be interesting if they were to investigate the association of amino acid substitutions at residue 70 of the HCV core both with *IL28B* polymorphisms and with hepatic steatosis.

Conflict of interest

The authors who have taken part in this study declared that they do not have anything to disclose regarding funding or conflict of interest with respect to this manuscript.

References

- [1] Tillmann HL, Patel K, Muir AJ, Guy CD, Li JH, Lao XQ, et al. Beneficial *IL28B* genotype associated with lower frequency of hepatic steatosis in patients with chronic hepatitis C. *J Hepatol* 2011;55:1195–1200.
- [2] Akuta N, Suzuki F, Sezaki H, Suzuki Y, Hosaka T, Someya T, et al. Association of amino acid substitution pattern in core protein of hepatitis C virus genotype 1b high viral load and non-virological response to interferon-ribavirin combination therapy. *Intervirology* 2005;48:372–380.
- [3] Donlin MJ, Cannon NA, Yao E, Li J, Wahed A, Taylor MW, et al. Pretreatment sequence diversity differences in the full-length hepatitis C virus open reading frame correlate with early response to therapy. *J Virol* 2007;81:8211–8224.
- [4] Hayes NC, Kobayashi M, Akuta N, Suzuki F, Kumada H, Abe H, et al. HCV substitutions and *IL28B* polymorphisms on outcome of peg-interferon plus ribavirin combination therapy. *Gut* 2011;60:261–267.
- [5] Tachi Y, Katano Y, Honda T, Hayashi K, Ishigami M, Itoh A, et al. Impact of amino acid substitutions in the hepatitis C virus genotype 1b core region on liver steatosis and hepatic oxidative stress in patients with chronic hepatitis C. *Liver Int* 2010;30:554–559.
- [6] Sumida Y, Kanemasa K, Hara T, Inada Y, Sakai K, Imai S, et al. Impact of amino acid substitutions in hepatitis C virus genotype 1b core region on liver steatosis and glucose tolerance in non-cirrhotic patients without overt diabetes. *J Gastroenterol Hepatol* 2011;26:836–842.
- [7] Kobayashi M, Akuta N, Suzuki F, Hosaka T, Sezaki H, Kobayashi M, et al. Influence of amino-acid polymorphism in the core protein on progression of liver disease in patients infected with hepatitis C virus genotype 1b. *J Med Virol* 2010;82:41–48.
- [8] Abe H, Ochi H, Maekawa T, Hayes CN, Tsuge M, Miki D, et al. Common variation of *IL28* affects gamma-GTP levels and inflammation of the liver in chronically infected hepatitis C virus patients. *J Hepatol* 2010;53:439–443.
- [9] Akuta N, Suzuki F, Hirakawa M, Kawamura Y, Yatsuji H, Sezaki H, et al. Amino acid substitution in hepatitis C virus core region and genetic variation near the interleukin 28B gene predict viral response to telaprevir with peginterferon and ribavirin. *Hepatology* 2010;52:421–429.



[10] Tanaka Y, Nishida N, Sugiyama M, Tokunaga K, Mizokami M. λ -Interferons and the single nucleotide polymorphisms: a milestone to tailor-made therapy for chronic hepatitis C. *Hepatol Res* 2010;40:449-460.

Hidegori Toyoda*

Takashi Kumada

Department of Gastroenterology, Ogaki Municipal Hospital,
4-86 Minaminokawa, Ogaki, Gifu 503-8502, Japan

* Corresponding author. Tel.: +81 584 81 3341;

fax: +81 584 75 5715.

E-mail address: hmtoyoda@spice.ocn.ne.jp (H. Toyoda).

Reply to: “Favorable association between genetic polymorphisms near the *IL28B* gene and hepatic steatosis: Direct or indirect?”

To the Editor:

We thank Drs. Toyoda and Kumada for raising the additional point that HCV amino acid substitutions have also been demonstrated to influence steatosis in the setting of HCV infection. In their study of 122 patients, 85 of whom had a beneficial *IL28B* genotype, Toyoda and Kumada found a trend for steatosis to be associated with *IL28B* polymorphism, as only 22% of the patients with beneficial genotype have steatosis compared to 40% of patients with the less beneficial genotype. Thus, their *r*, though not significant, is in line with our study, where we likewise found a 25% and 27% higher rate of steatosis in patients with the less beneficial *IL28B* (“non-C/C” for rs12979860 or “non-T/T” for rs8999017) genotype in two different cohorts of 145 and 180 patients, respectively. Similar to our and Toyoda and Kumada’s results, Cai *et al.* found an association between the beneficial *IL28B* genotype and lower steatosis frequency [1]. However, a three center study by Trépo *et al.* failed to find a relevant association between *IL28B* genotype and steatosis, according to their statement [2]. This latter paper, however, did not show the data, and therefore it cannot be assessed whether the association was absent or only not significant. Toyoda and Kumada’s study showed a similar trend for steatosis with *IL28B*, whereby *IL28B* is associated with different mutations in the core region, the HCV core mutation clearly shows a higher association with steatosis.

In our article, we indicate that response to treatment in relation to steatosis seems unlikely to be explained by *IL28B* alone, and though not specifically mentioned, *IL28B* is likely not solely responsible for the association with steatosis. We had a small cohort of 54 non-genotype 1 patients of whom 19 were genotype 3 and 35 were genotype 2. Despite the fact that steatosis tended to be higher in “non-C/C” patients (4/20 [20%] vs. 7/15 [46%] in genotype 2 patients and 4/8 [50%] vs. 7/11 [63%] in genotype 3 patients; Table 1) this was not significant. However, the trend was similar across genotypes. Furthermore, we have data on genotype 1a and 1b in 60 and 75 patients from the fibrosis study, respectively. In concordance with the overall results, steatosis was less frequently present in C/C genotype patients with both HCV genotype 1a and 1b (Table 1).

A possible explanation lies in the virus itself as the authors correctly point out with a focus on HCV’s core protein. Unfortunately, we do not have the core antigen sequence of our patients. A previous work by Jhaveri *et al.* suggested a role for amino acids 182 and 186 of the core protein, linking steatosis *in vitro* to steatosis [3], but certainly amino acid 70 seems to be especially relevant in genotype 1b infection. There is also some evidence indicating that not all differences can be explained by viral core sequence variation [4].

Table 1. Frequency of steatosis.

		Presence of steatosis	
1A	Non-CC	19/43 (44.2%)	<i>p</i> = 0.019
	CC	2/17 (11.8%)	
1B	Non-CC	28/56 (50%)	<i>p</i> = 0.034
	CC	4/19 (21%)	
2	Non-CC	7/15 (46%)	n.s.
	CC	4/20 (20%)	
3	Non-CC	7/11 (63%)	n.s.
	CC	4/8 (50%)	

n.s., not significant.

Conflict of interest

The authors declared that they do not have anything to disclose regarding funding of conflict of interest with respect to this manuscript.

References

- [1] Cai T, Dufour JF, Muellhaupt B, Gerlach T, Heim M, Moradpour D, et al. on behalf of the Swiss Hepatitis C Cohort Study Group. Viral genotype-specific role of PNPLA3, PPARG, MBOAT2, and IL28B in hepatitis C virus-associated steatosis. *J Hepatol* 2011;55:529-535.
- [2] Trépo E, Pradat P, Potthoff A, Momozawa Y, Quertinmont E, Gustot T, et al. Impact of patatin-like phospholipase-3 (rs738409 C>G) polymorphism on fibrosis progression and steatosis in chronic hepatitis C. *Hepatology* 2011;54:60-69. doi:10.1002/hep.2435.
- [3] Jhaveri R, McHutchison J, Patel K, Qiang G, Diehl AM. Specific polymorphisms in hepatitis C virus genotype 3 core protein associated with intracellular lipid accumulation. *J Infect Dis* 2008;197:283-291.
- [4] Piodi A, Chouteau P, Lerat H, Hézode C, Pawlotsky JM. Morphological changes in intracellular lipid droplets induced by different hepatitis C virus genotype core sequences and relationship with steatosis. *Hepatology* 2008;48:16-27.

Hans Ludger Tillmann*

Jeanette J. McCarthy

Duke University, Durham, NC, United States

* Corresponding author.

E-mail addresses: hans.tillmann@duke.edu, hans.tillmann@medizin.uni-leipzig.de (H.L. Tillmann).

Short Communication

Reference change values for *lens culinaris* agglutinin-reactive α -fetoprotein and des- γ -carboxy prothrombin in patients with chronic hepatitis C**Futoshi Kanke^{1,*}, Takashi Kumada²,
Hidenori Toyoda² and Shinji Satomura¹**¹Diagnostic Division, Wako Pure Chemical Industries Ltd.,
Osaka, Japan²Department of Gastroenterology, Ogaki Municipal
Hospital, Ogaki, Japan**Abstract**

Background: *Lens culinaris* agglutinin-reactive α -fetoprotein (AFP-L3) and des- γ -carboxy prothrombin (DCP) have been routinely used as serological tumor markers of hepatocellular carcinoma (HCC) for surveillance. The aims of this study were: (i) to determine the biological variation of AFP-L3 and DCP in patients with chronic hepatitis C; and (ii) to calculate the reference change values (RCVs) of AFP-L3 and DCP.

Methods: Ten patients with cirrhosis due to hepatitis C virus (HCV) infection and without HCC were enrolled in the study. Serum samples were collected at 14-day intervals, and 10 samples in total were obtained for each patient. AFP-L3 and DCP levels were measured by microchip capillary electrophoresis and liquid-phase binding assay. Intra-individual (CV_I) and inter-individual (CV_G) biological variations and RCVs were estimated from the data generated.

Results: The CV_I was 29.0% for AFP-L3 and 24.6% for DCP, and CV_G was 63.5% for AFP-L3 and 40.4% for DCP. The RCVs for AFP-L3 and DCP were 68.3% and 58.5%, respectively.

Conclusions: Increases in values for AFP-L3 and DCP within 68.3% and 58.5% may be biological variations. Clinician should take these variations into consideration for the management of patients with HCV infection under surveillance of HCC.

Keywords: α -fetoprotein (AFP-L3); biological variation; des- γ -carboxy prothrombin (DCP); hepatocellular carcinoma; reference change value (RCV).

Hepatocellular carcinoma (HCC) is one of the most prevalent cancers. It is the sixth most common cancer in the world, and the third most common cause of cancer-related death (1). Chronic liver disease caused by hepatitis C viruses (HCV) is an important risk factor for the development of HCC (2). α -Fetoprotein (AFP) (3, 4), the percentage of *lens culinaris* agglutinin-reactive fraction of AFP to total AFP (AFP-L3) (5, 6), and des- γ -carboxy prothrombin (DCP) (7) have been routinely used as serological tumor markers of HCC for surveillance, diagnosis, or the prediction of outcome in clinical practice. Whereas established cut-off with fixed values of AFP, AFP-L3, and DCP are usually used to make clinical decisions, fluctuation of these values are often observed in patients without HCC. Previous studies reported that the reference change value (RCV) for AFP was useful for dynamic monitoring of AFP in patients with liver disease (8). However, clinically minimal significant changes on sequential measurements of AFP-L3 and DCP have never been investigated. The aims of this study were: (i) to determine the biological variation of AFP-L3 and DCP in patients with chronic hepatitis C including cirrhosis; and (ii) to calculate the RCVs of AFP-L3 and DCP as the basis for clinically minimal significant change value.

Ten patients (five males and five females) having cirrhosis with HCV but without HCC were enrolled into the study between April and September 2008. The age of the patients (mean \pm SD) were 69.6 \pm 5.1 years (range, 62–76 years). The platelet counts were 92.2 \pm 27.4 \times 1000/mL. All patients were positive for HCV RNA as determined by a commercial assay (HCV COBAS AmpliPrep/COBAS TaqMan System; Roche Molecular Systems, Pleasanton, CA, USA). With respect to liver function, eight patients had Child-Pugh class A and the remaining two patients had Child-Pugh class B liver function. The absence of HCC was confirmed in all patients by ultrasonography and Gd-EOB-DTPA enhanced magnetic resonance imaging that is most sensitive modality to detect hepatic tumors (9) at the time of entry in order to avoid inclusion of patients with occult HCC. In addition, these patients were followed for 2 years after serum sampling for the lack of the development of HCC. Serum samples were collected at 14-day intervals following a standard protocol in order to minimize pre-analytical variations, and 10 samples in total were obtained from each patient. Samples were collected following a standardized protocol in order to minimize

*Corresponding author: Futoshi Kanke, Diagnostics Division, Wako Pure Chemical Industries, Ltd., 1-2, Doshomachi 3, Chuo-ku, Osaka 540-8605, Japan

Fax: +81 6 62031917, E-mail: kanke.futoshi@wako-chem.co.jp

Received May 9, 2011; accepted January 27, 2012

pre-analytical variations. All serum samples were stored frozen at -80°C until measurement. The study protocol was in compliance with the Helsinki Declaration and was approved by the Ethics Committee of Ogaki Municipal Hospital. Written informed consent was obtained from each patient.

AFP-L3%, DCP, and AFP levels were measured by a microchip-based capillary electrophoresis and liquid-phase binding assay on the μTAS Wako i30 autoanalyzer (Wako Pure Chemical Industries, Ltd., Japan) (10). Because we purposed to compare biological variation of AFP-L3, which is expressed as a percentage of total AFP in a current use, with that of AFP-L3 concentration (expressed in units of ng/mL), AFP-L3 concentration (ng/mL) was calculated using the formula $\text{total AFP} \times (\text{AFP-L3}/100)$. Albumin (ALB), total bilirubin (T-Bil), aspartate aminotransferase (AST), and alanine aminotransferase (ALT) were measured on a general clinical chemistry analyzer as liver function analytes. ALB was measured by bromocresol purple method (L-Type WAKO ALB-BCP; Wako Pure Chemical Industries, Ltd.), T-Bil was measured by vanadate oxidation method (Total Bilirubin E-HA; Wako Pure Chemical Industries, Ltd.), and AST and ALT were measured by Japan Society Clinical Chemistry transferable method (L-Type AST.J2 and ALT.J2; Wako Pure Chemical Industries, Ltd.). Conversion factors from conventional units to SI units for ALB (g/dL \rightarrow g/L), T-Bil (mg/dL \rightarrow $\mu\text{mol/L}$), AST/ALT (U/L \rightarrow $\mu\text{kat/L}$), AFP/AFP-L3 (ng/mL \rightarrow $\mu\text{g/L}$) and DCP (mAU/mL \rightarrow $\mu\text{g/L}$) are 10, 17.1, 0.017, 1 and 0.012, respectively. Each sample was assayed in duplicates. Measurements were performed by the same analyst. To calculate analytical imprecision (CV_A), controls (AFP: 18.8 ng/mL, AFP-L3: 9.7%, DCP: 99 mAU/mL, ALB: 3.0 g/dL, T-BIL: 0.6 mg/dL, AST: 28 U/L, ALT: 25 U/L) were measured in duplicate with two runs per day over 21 days.

Data was analyzed by calculating intra- (CV_I), and inter-individual (CV_O) biological variations according to Fraser et al. (11, 12). We also calculated the index of individuality (IOI) using an equation of $\text{IOI} = \text{CV}_I / \text{CV}_O$ according to Harris et al. (11, 13). When the IOI for a particular test is ≤ 0.6 , conventional population-based reference intervals and cut-off values are of limited value for the classification of some unusual results for a particular individual. When the IOI is ≥ 1.4 , the variation of a particular individual will fit within the assay limits as those of the population and, therefore, population-derived reference value could be applied. The critical differences at 5% significance level (one-tail test) for AFP-L3%, AFP-L3 absolute concentration, DCP, AFP, ALB, T-Bil, AST, and ALT (including analytical variation) were calculated as $2.33 \times (\text{CV}_I^2 + \text{CV}_A^2)^{1/2}$. All statistical analyses were performed with JMP6 statistical software (SAS Institute Japan, Japan).

The median and range of AFP-L3%, AFP-L3 concentration, DCP, and AFP in each of 10 patients are shown in Figure 1. Although all of these values did not show a normal distribution by Shapiro-Wilk test, we did not use logarithmic transformation for these values. It was noted that the median value was below the recommended cut-off of 10% of AFP-L3% and 40 mAU/mL of DCP, respectively. Although the mean values of ALB, T-Bil, and ALT were within the respective reference ranges, the mean value of AST slightly exceeded the reference range (Table 1). Although we observed some cases in which the range was greater in comparison to other cases (ex. patient #2 for AFP and AFP-L3 concentration, patients #8 and #10 for AFP-L3%, and patients #2 and #6 for DCP), these were not outliers (11). Patient #2 showed greater ranges of AFP, AFP-L3 concentration and DCP, but the changes in these values in serial measurement were not spiky, which reportedly indicates the development of HCC in case of AFP (14), and the development of HCC was no




Figure 1 The median and range of AFP (A), AFP-L3% (B), AFP-L3 concentration (C), and DCP (D) for each patient.

Table 1 The mean, range, SD, CVA, CV_I, CV_G, RCV and IoI for AFP, AFP-L3%, AFP-L3 concentration, DCP albumin, total bilirubin, AST, and ALT.

Marker	Mean	Range	SD	CVA, %	CV _I , %	CV _G , %	RCV, %	IoI
AFP, ng/mL	12.771	71.712	15.3	2.1	52.1	113.1	121.8	0.461
AFP-L3%, %	4.45	8.21	3.0	3.8	29.0	63.5	68.3	0.457
AFP-L3, ng/mL	0.714	4.076	0.9	4.9	42.4	122.0	99.6	0.348
DCP, mAU/mL	18.0	47.2	8.3	4.7	24.6	40.4	58.5	0.609
Albumin, g/dL	3.97	1.95	0.5	0.8	3.7	12.5	8.9	0.300
Total bilirubin, mg/dL	0.81	1.40	0.3	2.4	17.5	32.3	41.2	0.540
AST, U/L	51.4	142.6	22.3	5.5	26.7	36.1	63.5	0.740
ALT, U/L	35.6	106.1	17.8	3.4	36.4	36.3	85.3	1.000

CV_A, analytical imprecision; CV_I, intra-individual biological variation; IoI, index of individuality; CV_G, inter-individual biological variation; RCV, reference change value.

observed under surveillance more than 2 years after study. Results of calculations for CV_A, CV_I, CV_G, RCV, and IOI are summarized in Table 1. The CV_I was 29.0% and CV_G was 63.5% for AFP-L3%, 42.4% and 122.0% for AFP-L3 concentration, 24.6% and 40.4% for DCP, and 52.1% and 113.1% for AFP. The CV_I and CV_G for AFP-L3% were smaller than those of AFP-L3 concentration. This finding indicated that AFP-L3 that was expressed as a percentage of total AFP as a current use is more appropriate than AFP-L3 concentration as a tumor marker for HCC, since the effects of biological variance on clinical decision-making can be reduced when AFP-L3 was expressed as a percentage. The IOI values for AFP-L3%, AFP-L3 concentration, DCP, and AFP were 0.457, 0.348, 0.609, and 0.461, respectively. The value was under 0.6 in case of AFP-L3%, AFP concentration, and AFP, and was far from 1.4 in case of DCP.

Measurement of tumor markers for HCC, AFP, AFP-L3, and DCP, are recommended for patients with high-risk of developing HCC in Japan (15), which plays an important role in HCC surveillance. Fixed cut-off values are used for AFP, AFP-L3, and DCP for clinical decision-making. However, our findings demonstrated that the cut-off values of AFP-L3 and DCP might be of limited clinical utility as an indicator of developing HCC, since the IOI scores were <0.6 in our study; the range of the values for an individual subject spans only a small part of the reference interval for a population. One previous study investigated CV_I, CV_G, and IOI score for AFP (8), with which the data for AFP in the present study was compatible, indicating limited clinical utility of the cut-off value for AFP of 20 ng/mL and 200 ng/mL.

Similar finding are reported for other tumor markers. Carcinoembryonic antigen (CEA) and CA 19-9 also have low IOI scores. It was determined that conventional or population-based reference intervals for CEA and CA19-9 are of limited utilities in interpretation of the assay results, since the IOI for these markers were <0.6 (16, 17). The use of RCVs would improve the clinical utility of these markers. Like these tumor markers, utilization of the RCVs for AFP-L3, DCP, and AFP might improve the accuracy for the detection of HCC under surveillance.

This study includes several limitations. The potential bias may be present in patient selections due to a study performed

at a single site, and the number of patients analyzed was small. Larger clinical studies are needed to further evaluate and confirm the RCVs for AFP-L3 and DCP. In addition, all patients analyzed were infected with HCV, and these analyses should be applied for patients with HBV infection or those without hepatitis viral infection.

In conclusion, conventional population-based cut-off values for AFP-L3 and DCP may be of limited value for indicating HCC development in HCV-infected patients at high-risk for HCC under surveillance, due to the effects of significant biological variations of these markers over population based cut-off selection. Analytical and intra-individual component of the biological variations would accounted for the significant changes in sequential marker concentrations, and these should be determined and taken into considerations in interpretation of the assay results. In the management of patients with chronic HCV infection, which is a major cause of HCC development in Japan, clinicians should take these values of biological variation of AFP-L3 and DCP (68.3% and 58.5%) into consideration when evaluating the changes of these biomarkers in serial measurements. Further clinical evaluation is needed to address whether the RCVs for AFP-L3, and DCP would be useful for HCC detection.

Conflict of interest statement

Authors' conflict of interest disclosure: The authors stated that there are no conflicts of interest regarding the publication of this article. The employment status of F. Kanke and S. Satomura at Wako Pure Chemical Industries Ltd. played no role in the study design; in the collection, analysis, and interpretation of data; in the writing of the report; or in the decision to submit the report for publication.

Research funding: None declared.

Employment or leadership: F. Kanke and S. Satomura are employees at Wako Pure Chemical Industries Ltd.

Honorarium: None declared.

References

1. Parkin D, Bray F, Ferlay J, Pisani P. Global cancer statistics, 2002. *CA Cancer J Clin* 2002;55:74–108.

2. Gomaa A, Khan SA, Toledano MB, Waked I, Taylor-Robinson SD. Hepatocellular carcinoma: epidemiology, risk factors and pathogenesis. *World J Gastroenterol* 2008;14:4300–8.
3. Abelev GI. Production of embryonal serum alpha-globulin by hepatomas: review of experimental and clinical data. *Cancer Res* 1968;28:1344–50.
4. Bisceglie AM, Sterling RK, Chung RT, Everhart JE, Dienstag JL, Bonkovsky HL, et al. Serum alpha-fetoprotein levels in patients with advanced hepatitis C: results from the HALT-C Trial. *J Hepatol* 2005;43:434–41.
5. Taketa K, Endo Y, Sekiya C, Tanikawa K, Koji T, Taga H, et al. A collaborative study for the evaluation of lectin-reactive alpha-fetoproteins in early detection of hepatocellular carcinoma. *Cancer Res* 1993;53:5419–23.
6. Tamura Y, Igarashi M, Kawai H, Suda T, Satomura S, Aoyagi Y. Clinical advantage of highly sensitive on-chip immunoassay for fucosylated fraction of alpha-fetoprotein in patients with hepatocellular carcinoma. *Dig Dis Sci* 2010;55:3576–83.
7. Yano Y, Yamashita F, Kuwaki K, Fukumori K, Kato O, Yamamoto H, et al. Clinical features of hepatitis C virus-related hepatocellular carcinoma and their association with alpha-fetoprotein and protein induced by vitamin K absence or antagonist-II. *Liver Int* 2006;26:789–95.
8. Trapé J, Botargues J, Porta F, Ricós C, Badal J, Salinas R, et al. Reference change value for alpha-fetoprotein and its application in early detection of hepatocellular carcinoma in patients with hepatic disease. *Clin Chem* 2003;49:1209–11.
9. Kagebayashi C, Yamaguchi I, Akinaga A, Kitano H, Yokoyama K, Satomura M, et al. Automated immunoassay system for AFP-L3% using on-chip electrokinetic reaction and separation by affinity electrophoresis. *Anal Biochem* 2009;388:306–11.
10. Kim SH, Kim SH, Lee J, Kim MJ, Jeon YH, Park Y, et al. Gadoteric acid-enhanced MRI versus triple-phase MDCT for the preoperative detection of hepatocellular carcinoma. *Am J Roentgenol* 2009;192:1675–81.
11. Fraser C, Harris E. Generation and application of data on biological variation in clinical chemistry. *Crit Rev Clin Lab Sci* 1989;27:409–37.
12. Gowans E, Fraser C. Biological variation of serum and urine creatinine and creatinine clearance: ramifications for interpretation of results and patient care. *Ann Clin Biochem* 1988;25:259–63.
13. Harris E. Effects of intra- and interindividual variation on the appropriate use of normal ranges. *Clin Chem* 1974;20:1535–42.
14. Kuwahara T, Sakai T, Majima Y, Hirai K, Tanikawa K. Serial changes in serum alpha-fetoprotein prior to detection of hepatocellular carcinoma in liver cirrhosis. *Hepatogastroenterol* 1993;40:347–51.
15. Group formed to establish 'Guidelines for evidence-based clinical practice for the treatment of liver cancer'. *Clinical Practice Guidelines for Hepatocellular Carcinoma (in Japanese)*. Tokyo, Kanehara, 2009.
16. Plebani M, Giacomini A, Beghi L, dePaoli M, Roveroni G, Galeotti F, et al. Serum tumor markers in monitoring patients: interpretation of results using analytical and biological variation. *Anticancer Res* 1996;16:2249–52.
17. Sölétormos G, Schiøler V, Nielsen D, Skovsgaard T, Dombrowsky P. Interpretation of results for tumor markers on the basis of analytical imprecision and biological variation. *Clin Chem* 1993;39:2077–83.

Novel method to measure serum levels of des-gamma-carboxy prothrombin for hepatocellular carcinoma in patients taking warfarin: A preliminary report

Hidenori Toyoda,^{1,4} Takashi Kumada,¹ Yukio Osaki,² Toshifumi Tada,¹ Yuji Kaneoka³ and Atsuyuki Maeda³

¹Department of Gastroenterology, Ogaki Municipal Hospital, Ogaki, Japan; ²Department of Gastroenterology, Osaka Red Cross Hospital, Osaka, Japan;

³Department of Surgery, Ogaki Municipal Hospital, Ogaki, Japan

(Received November 4, 2011/Revised January 30, 2012/Accepted January 31, 2012/Accepted manuscript online February 9, 2012)

Des-gamma-carboxy prothrombin (DCP) is a useful tumor marker for hepatocellular carcinoma (HCC), but its utility is limited in patients taking vitamin K antagonists. We evaluated the NX-DCP ratio, a newly developed method to measure serum DCP, for its ability to identify DCP elevation induced by HCC in this patient subpopulation. Conventional DCP measurements and the NX-DCP ratio were compared in patients with and without HCC, all of whom were taking the vitamin K antagonist warfarin. We found no differences in conventional DCP measurements between patients with and without HCC due to warfarin treatment. In contrast, the NX-DCP ratio was significantly higher in patients with HCC; the NX-DCP ratio in all patients without HCC was <1.50. When the cut-off was fixed at 1.50, sensitivity and specificity for HCC diagnosis were 60.0% and 100.0%, respectively, which are comparable to those of conventional DCP measurements in patients not taking warfarin. The novel NX-DCP ratio identifies patients on warfarin with elevated DCP due to HCC and is useful as a tumor marker for HCC in this patient subpopulation. (*Cancer Sci*, doi: 10.1111/j.1349-7006.2012.02232.x, 2012)

Prothrombin, or coagulation factor II, is a 71 600 Da protein that consists of three regions: fragment 1; fragment 2; and prothrombin. Fragment 1 consists of 156 amino acids, including 41 amino acids forming an *N*-terminal gamma-glutamic acid (Gla)-containing domain. Prothrombin is first synthesized in the liver as a precursor with 10 glutamic acid (Glu) residues, which are then modified to Gla residues by gamma-glutamylcarboxylase in the presence of vitamin K, O₂, and CO₂ before it is released into the bloodstream.

However, in the absence of vitamin K or in the presence of vitamin K antagonists, gamma-carboxylation is impaired, and prothrombin with the remaining Glu residues, which is inactive with respect to coagulation, is released into the bloodstream.⁽¹⁾ Prothrombin in this form is called des-gamma-carboxy prothrombin (DCP) or protein induced by vitamin K absence/antagonist-II (PIVKA-II). As the number of Glu residues unconverted to Gla varies, DCP is present as a mixture of prothrombin with various numbers of Glu residues, ranging from 1 to 10. In addition, because the Gla residue can bind to calcium, it is known that the 3-D protein structure of DCP will be different in the presence of calcium, and depends on the number of Glu residues.⁽²⁾

Des-gamma-carboxy prothrombin is frequently found in the blood of patients with hepatocellular carcinoma (HCC). Because DCP is elevated in many patients with HCC but not in patients with chronic hepatitis or cirrhosis without HCC,⁽³⁾ it has been routinely used as a tumor marker of HCC in clinical settings.⁽⁴⁻⁶⁾ However, serum DCP levels are also increased

in the absence of HCC when there is a shortage of vitamin K or in the presence of vitamin K antagonists.⁽⁷⁾ The value of DCP as a marker of HCC, therefore, is significantly reduced in patients who are taking vitamin K antagonists such as warfarin.

Previous studies reported differences in the number of Glu residues in DCP between patients with HCC and patients taking vitamin K antagonists.^(8,9) Conventionally, DCP is measured using a mAb produced by the cell line MU-3 (Picolumi PIVKA-II; EIDIA, Tokyo, Japan), which reportedly reacts predominantly with DCP with 9–10 Glu residues. MU-3 had lower affinity for DCP with one to five Glu residues.⁽¹⁰⁾ However, measuring DCP with this antibody alone can not differentiate between HCC-induced and vitamin K antagonist-associated elevations of DCP, making it difficult to evaluate whether rises in DCP are caused by HCC in patients taking vitamin K antagonists.

In the present study, we attempted to identify HCC-induced DCP in patients with HCC taking the vitamin K antagonist warfarin through the use of two mAbs against DCP, P-11 and P-16 (Sekisui Medical, Tokyo, Japan), which have a reactivity profile different from MU-3. We found clinical utility in DCP as a marker for HCC in patients taking warfarin when measured with the combination of MU-3, P-11, and P-16.

Materials and Methods

Preparation of electrochemiluminescence immunoassay (ECLIA) reagents with P-11 and P-16. Magnetic beads coated with P-16 mAb (Sekisui Medical) were prepared as follows: 1 mL of 30 mg/mL magnetic bead suspension (Dynabeads M-450 Epoxy; Life Technologies, Carlsbad, CA, USA) was placed into a test tube and the magnetic beads were trapped by a magnet to separate the supernatant. After the supernatant was discarded, 1 mL P-16 mAb (0.5 mg/mL in 0.15 mol/L PBS, pH 7.8) was added to the magnetic beads and stirred at 25°C for 18 h. After washing the magnetic beads, 2 mL of 1% BSA in 0.15 mol/L PBS (pH 7.8) were added and stirred at 25°C for 18 h to block the beads. These beads were diluted to 1 mg/mL using the bead dilution reagent (0.05 mol/L Tris buffer (pH 7.5), 0.15 mol/L NaCl, 0.01% Tween 20, 0.1% NaN₃, 10% normal rabbit serum, and 0.1% mouse serum) when in use.

Ruthenium (Ru)-conjugated P-11 mAb was prepared by the following procedure: 68 µL Ru-complex compounds (10 mg Ru (II) Tris (bipyridyl)-NHS ester in 1 mL DMSO) was added

[†]To whom correspondence should be addressed.
E-mail: tkumada@he.mirai.ne.jp

to 1 mL P-11 mAb (1 mg/mL in 0.15 mol/L PBS, pH 7.8) (Sekisui Medical) for conjugation, and stirred at 25°C for 30 min. Then, 50 μ L of 2 mol/L glycine was added to terminate the conjugation reaction, and Ru-conjugated P-11 mAb was isolated by collecting the Ru-bound protein fraction using Sephadex G-25 (previously equilibrated with 10 mmol/L PBS, pH 6.0). The Ru-conjugated P-11 mAb was then diluted to 1 μ g/mL using Ru dilution reagent (0.015 mol/L HEPES buffering solution [pH 7.8], 0.15 mol/L NaCl, 0.013 mol/L CaCl₂, 0.1% Tween 20, 0.1% NaN₃, 5% normal rabbit serum, and 0.1% mouse serum) when in use.

Measurement of conventional DCP (with MU-3 antibody), NX-DCP (with P-11 and P-16 antibodies), and NX-DCP ratio. Conventional DCP, which is measured using MU-3 antibody and is currently used in clinical settings, was measured with ECLIA using the Picolumi III automated analyzer (EIDIA). NX-DCP was measured by ECLIA. Briefly, 25 μ L magnetic beads coated with P-16 mAb (1 mg/mL) and 150 μ L Ru-conjugated P-11 mAb (1 μ g/mL) were added to samples at 30°C for 9 min to obtain the value of NX-DCP. The NX-DCP ratio was calculated by dividing the value of DCP measured using the conventional Picolumi method by the value of NX-DCP.

Reactivity of MU-3, P-11, and P-16 mAbs based on the time allowed for decarboxylation from prothrombin. We prepared DCP with varying numbers of Glu residues by applying different time intervals for decarboxylation from prothrombin (Enzyme Research Laboratories, Swansea, UK), according to the method of Bajah *et al.*⁽²⁾ Specifically, 0.78 mL ammonium bicarbonate solution (0.1 mol/L, pH 8.0) was applied to 4.6 mg/mL prothrombin solution overnight for dialysis against an ammonium bicarbonate solution at 4°C. Then 0.1 mol/L EDTA*2Na was applied to the solution after dialysis until a final concentration of 10 mmol/L was achieved, and the solution was allowed to stand at room temperature for 30 min. This solution was dialyzed again against an ammonium bicarbonate solution for 2 h at 4°C then aliquoted into six heat-resistant vials with a screw cap and lyophilized. The vials were then filled with nitrogen gas and heated to 110°C for 0, 30 min, 1, 2, 6, or 23 h to create six different samples. We coated microplates with 100 μ L of each sample at 0.1 μ g/mL, and tested reactivity of the MU-3, P-11, and P-16 mAbs in the presence of 4 mmol/L calcium chloride. The experiments were repeated three times and the average value was calculated.

Patients. A total of 338 patients were diagnosed with primary, non-recurrent HCC between January 2006 and December 2009 at Ogaki Municipal Hospital (Ogaki, Japan). Of these, 14 patients had been taking warfarin when HCC was diagnosed. Six patients at Osaka Red Cross Hospital (Osaka, Japan) who were diagnosed as having primary HCC during the same period and had been taking warfarin at the time of diagnosis were also enrolled in the study. We analyzed the stored serum samples from these 20 patients, obtained at the time of HCC diagnosis. The diagnosis of HCC was made by histological examination or appropriate imaging characteristics using criteria of the guidelines by the American Association for the Study of Liver Diseases.⁽¹¹⁾ Tumor stage on imaging findings was assessed according to the TNM classification of the Liver Cancer Study Group of Japan.⁽¹²⁾

Control samples were obtained from 56 patients with chronic liver disease without HCC who were followed up at Ogaki Municipal Hospital. Samples were collected during routine HCC surveillance during the same period. These patients had been taking warfarin when serum samples were collected and provided informed consent for their stored serum samples to be used for research. The diagnosis of chronic liver disease was made with histological examination in 45 patients, includ-

ing 10 with cirrhosis. The remaining 11 patients were diagnosed with cirrhosis based on imaging findings and biochemical tests. To ensure that controls did not have HCC, these patients were followed for 3 years after serum sampling by ultrasonography, CT, or MRI to ensure that none had developed HCC.

The protocol for the clinical part of this study was approved by the institutional review board of Ogaki Municipal Hospital and carried out in compliance with the Helsinki Declaration. Written informed consent was obtained from all study patients for the use of clinical and laboratory data and stored serum samples.

Statistical analyses. Differences in percentages between groups were analyzed using the χ^2 -test. Differences in mean quantitative values were analyzed by the Mann-Whitney *U*-test. Changes in the NX-DCP ratio with increases in HCC stage were analyzed with the Jonckheere-Terpstra test. Data analyses were carried out using JMP statistical software, version 6.0 (Macintosh version; SAS Institute, Cary, NC, USA). All *P*-values were derived from two-tailed tests, with *P* < 0.05 considered to indicate statistical significance.

Results

Reactivity of MU-3, P-11, and P-16 antibodies with DCP based on time allowed for decarboxylation from prothrombin. Figure 1 shows the reactivity of the MU-3, P-11, and P-16 antibodies according to the time allowed for decarboxylation from prothrombin. MU-3 did not react with prothrombin (0 min) and its reactivity increased as the heating time (time allowed for decarboxylation) increased, with maximum reactivity to the sample after 6 h of heating. In contrast, P-11 and P-16 showed maximum reactivity to the 1-h sample and reactivity decreased as the heating time (time allowed for decarboxylation) increased.

Patient characteristics and levels of conventional DCP, NX-DCP, and NX-DCP ratio. Warfarin was used to treat atrial fibrillation in 48 patients, a history of mitral or aortic valve replacement in 13 patients, and a history of cerebral infarction in 15 patients. Table 1 summarizes the characteristics of the patients with and without HCC. There were no differences in patient




Fig. 1. Reactivity of MU-3, P-11, and P-16 antibodies according to the time allowed for decarboxylation from prothrombin. MU-3 did not react with prothrombin (0 min) and its reactivity increased as the heating time (time allowed for decarboxylation) increased, with maximum reactivity to the sample after 6 h of heating. Both P-11 and P-16 showed maximum reactivity to the 1-h sample and reactivity decreased as the heating time increased.

Table 1. Background characteristics of study patients with and without hepatocellular carcinoma (HCC) (n = 76)

	Patients with HCC (n = 20)	Patients without HCC (n = 56)	P-value
Mean age ± SD, years (range)	72.4 ± 8.0 (46–83)	70.0 ± 9.8 (46–86)	0.3211
Sex, female/male	6 (30.0)/14 (70.0)	19 (33.9)/37 (66.1)	0.9651
Albumin, g/dL (mean ± SD)	3.82 ± 0.42	3.97 ± 0.51	0.2276
Total bilirubin, mg/dL (mean ± SD)	1.02 ± 0.65	0.82 ± 0.52	0.1288
Platelets (×10 ³ /μL)	158 ± 75	160 ± 49	0.4754
INR	1.75 ± 0.58	1.76 ± 0.58	0.7816
Mean tumor size ± SD, cm (range)	3.35 ± 1.84 (1.1–8.4)	–	–
Number of tumors, single/multiple	15 (75.0)/5 (25.0)	–	–
Tumor stage, I/II/III†	3 (15.0)/11 (55.0)/6 (30.0)	–	–

†According to the TNM classification of the Liver Cancer Study Group of Japan. Unless otherwise indicated, values in parentheses indicate percentages. INR, International normalized ratio.

age, sex, serum albumin, serum total bilirubin, platelet count, or prothrombin levels.

Figure 2 compares conventional DCP levels, NX-DCP levels, and NX-DCP ratios between patients with and without HCC. No differences were found in conventional DCP levels between patients with and without HCC (median, 2600.5 mAU/mL and range, 1060–96920 mAU/mL in patients with HCC versus median, 20550.5 mAU/mL and range, 1355–71783 mAU/mL in patients without HCC; $P = 0.7952$). In contrast, NX-DCP levels in patients with HCC (median, 34135.0 mAU/mL; range, 260–67581 mAU/mL) were significantly lower than in patients without HCC (median, 40708.0 mAU/mL; range, 5026–60443 mAU/mL; $P = 0.0291$). As a result, the NX-DCP ratio was significantly higher in patients with HCC (median, 1.92; range, 0.35–10.32) than in patients without HCC (median, 0.49; range, 0.12–1.33; $P < 0.0001$).

Sensitivity, specificity, and positive and negative predictive values of NX-DCP ratio for diagnosis of HCC. Figure 3(a) shows the receiver operating characteristic (ROC) curve of the NX-DCP ratio for the diagnosis of HCC. The area under the ROC curve was 0.8928. The highest Youden index was 0.68 when the cut-off was fixed as 0.65 and the highest accuracy was 89.5% when the cut-off was fixed as 1.50, based on the sensitivity and specificity analysis (Fig. 3b). When the cut-off was fixed as 0.65, sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), and accuracy were 95.0%, 73.2%, 55.9%, 97.6%, and 78.9%, respectively. When the cut-off was fixed as 1.50, sensitivity, specificity, PPV, NPV, and accuracy were 60.0%, 100.0%, 100.0%, 87.5%, and 89.5%, respectively.

Serum alpha-fetoprotein (AFP) and *Lens culinaris* agglutinin-reactive fraction of AFP (AFP-L3) levels in patients with HCC. Serum levels of AFP and AFP-L3 were measured⁽¹³⁾ in patients with HCC in the same serum for the measurement of NX-DCP ratio (AFP-L3 was not measured in five patients). The median (range) values were 18.4 ng/mL (0.8–68 470 ng/mL) for AFP and 3.6% (0–45.2%) for AFP-L3. When the cut-off levels of AFP and AFP-L3 were fixed as 20 ng/mL and 5%, respectively, according to previous reports,^(14,16) 10 of 20 patients (50.0%) showed elevation of AFP and seven of 15 patients (46.7%) showed elevation of AFP-L3. These two tumor markers were not increased in six of 15 patients (40.0%).

NX-DCP ratio and progression of HCC. Figure 4 shows the NX-DCP ratio in patients according to HCC stage. Despite the small number of patients, there was a gradual increase in the NX-DCP ratio as the stage increased ($P = 0.0315$).

Discussion

Hepatocellular carcinoma is the sixth most common cancer and the third most common cause of cancer-related death worldwide.^(17,18) In Japan, HCC is currently the third most common cause of death from cancer in men and the fifth in women.⁽¹⁹⁾ The incidence of HCC is also increasing in the US.^(20,21) Improvements of tumor markers specific for HCC contribute to early detection of HCC. Three markers for HCC are currently used clinically, AFP, AFP-L3, and DCP. The utility of each of these tumor markers for detection and diagnosis of HCC, for evaluation of tumor progression, and for determination of patient prognosis has been reported.^(4,22–24) Elevation




Fig. 2. Serum levels of conventional des-gamma-carboxy prothrombin (DCP), NX-DCP, and the NX-DCP ratio in patients with and without hepatocellular carcinoma (HCC) taking warfarin. (a) Serum levels of conventional DCP. No differences were found between two groups ($P = 0.7952$). (b) Serum levels of NX-DCP were significantly higher in patients without HCC compared to those with HCC ($P = 0.0291$). (c) The NX-DCP ratio was significantly higher in patients with HCC than in those without HCC, consequently ($P < 0.0001$).




Fig. 3. Receiver operating characteristic (ROC) analysis and the determination of cut-off level of the NX-DCP ratio for the diagnosis of hepatocellular carcinoma. (a) The area under the ROC curve was 0.8928. (b) The highest Youden index was 0.68 when the cut-off was fixed as 0.65 and the highest accuracy was 89.5% when the cut-off was fixed as 1.50.




Fig. 4. The NX-DCP ratio according to hepatocellular carcinoma (HCC) stage in 20 patients with HCC taking warfarin (box plot). There was a gradual increase in the NX-DCP ratio as the HCC stage increased ($P = 0.0315$).

of DCP is often observed in HCC patients without elevation of AFP or AFP-L3, and is useful as a complement to these other two markers in the diagnosis of HCC. In addition, elevation of DCP is reportedly associated with a high rate of portal vein invasion and poor prognosis.⁽²⁵⁾ Elevation of DCP is also associated with better outcomes when hepatectomy, rather than radiofrequency ablation, is carried out in patients treated with curative intent.^(26,27)

However, DCP loses its value as a tumor marker of HCC in patients taking warfarin.⁽⁷⁾ Due to the marked decrease in vitamin K level caused by warfarin intake, DCP levels significantly increase in individuals taking warfarin, even in the absence of HCC. Therefore, DCP has no clinical utility as a tumor marker for HCC in this patient subpopulation.

The present study evaluated the reactivity of new antibodies against DCP, antibodies P-11 and P-16, based on the number of Glu residues. The number of Glu residues increases as the time allowed for decarboxylation from prothrombin increases.⁽²⁾ Our results showed P-11 and P-16 have higher reactivity with DCP with fewer Glu residues than MU-3, the

antibody that is conventionally used for the measurement of DCP. The NX-DCP level that is measured by P-11 and P-16 antibodies, therefore, represents predominantly DCP caused by reduced vitamin K availability. Consequently, the elevation of the NX-DCP ratio calculated in the equation: conventional DCP/NX-DCP, reflects more specifically the elevation of DCP by HCC.

There were no differences in the conventional measurements of DCP between patients with and without HCC who are taking warfarin. The NX-DCP ratio was significantly lower in patients without HCC than in patients with HCC. The NX-DCP ratio varied in patients with HCC, as was conventional DCP in patients not taking warfarin, because the production of DCP by HCC is variable. In contrast, in all patients without HCC, the NX-DCP ratio was low despite high conventional DCP levels in the same patients; no patients had an elevated NX-DCP ratio. The results indicate that the NX-DCP ratio could pinpoint the elevation of DCP caused by HCC, thereby restoring the value of DCP as a marker for HCC in patients taking warfarin.

When the cut-off level was fixed at 1.5 on the basis of maximal accuracy, the sensitivity, specificity, PPV, and NPV were comparable to those of conventional DCP in the general population with normal vitamin K levels, as previously reported.⁽²⁸⁾ The NX-DCP ratio, therefore, seems to be useful as a marker for HCC in patients taking warfarin.

The elevation of other tumor markers for HCC, AFP and AFP-L3, were observed in only half of the patients with HCC taking warfarin. In addition, both AFP and AFP-L3 were negative in 40% of patients with HCC. Des-gamma-carboxy prothrombin is a complimentary marker of AFP/AFP-L3 for HCC. The elevation of DCP without the elevation of AFP and AFP-L3 was observed in 16.1% of patients (cut-off, 20 ng/mL for AFP, 10% for AFP-L3, and 40 mAU/mL for DCP)⁽²⁹⁾ and 24.8% of patients (cut-off, 400 ng/mL for AFP, 15% for AFP-L3, and 100 mAU/mL for DCP).⁽³⁰⁾ The measurement of the NX-DCP ratio, therefore, will be important for the detection and diagnosis of HCC even when AFP or AFP-L3 is measured simultaneously.

There are several limitations to this study. The most important limitation was the small number of study patients, especially patients with HCC. The number of patients with HCC taking warfarin is low, so it was difficult to increase the number of study patients. Consequently, it was difficult to evaluate the value of the NX-DCP ratio in indicating progression of

HCC, including tumor stage progression and portal vein invasion, and in predicting patient outcome. Further studies will be necessary to establish the value of the NX-DCP ratio as a tumor marker for HCC in patients taking warfarin. In addition, the value of the NX-DCP ratio was evaluated only in patients who were taking the vitamin K antagonist warfarin; its value was not evaluated in HCC patients in whom vitamin K is reduced or absent through other mechanisms such as heavy alcohol intake or nutritional deficiency. The value of the NX-DCP ratio as a marker for HCC should be confirmed for these subpopulations in the future.

In conclusion, the novel NX-DCP ratio identified elevation of DCP due to HCC in patients taking the vitamin K antago-

nist warfarin. Thus, by using this ratio, DCP can be used as a marker for HCC even in patients taking warfarin.

Acknowledgments

We thank Jun Nishimura and Masato Uehara (EIDIA, Tokyo, Japan) for their technical assistance.

Disclosure Statement

The authors have no conflicts of interest to declare.

References

- Blanchard RA, Furie BC, Jorgensen M, Kruger SF, Furie B. Acquired vitamin K-dependent carboxylation deficiency in liver disease. *N Engl J Med* 1981; **305**: 242–8.
- Bajaj SP, Price PA, Russell WA. Decarboxylation of gamma-carboxyglutamic acid residues in human prothrombin. *J Biol Chem* 1982; **257**: 3726–31.
- Kuromatsu R, Tanaka M, Shimauchi Y *et al*. Usefulness of ED036 kit for measuring serum PIVKA-II levels in small hepatocellular carcinoma. *J Gastroenterol* 1997; **32**: 507–12.
- Liebman HA, Furie BC, Tong MJ *et al*. Des-gamma-carboxy (abnormal) prothrombin as a serum marker of primary hepatocellular carcinoma. *N Engl J Med* 1984; **310**: 1427–31.
- Fujiyama S, Morishita T, Hashiguchi O, Sato T. Plasma abnormal prothrombin (des-gamma-carboxy prothrombin) as a marker of hepatocellular carcinoma. *Cancer* 1988; **61**: 1621–8.
- Mita Y, Aoyagi Y, Yanagi M, Suda T, Suzuki Y, Asakura H. The usefulness of determining des-gamma-carboxy prothrombin by sensitive enzyme immunoassay in the early diagnosis of patients with hepatocellular carcinoma. *Cancer* 1998; **82**: 1643–8.
- Deyashiki Y, Nishioka Y, Takahashi K, Kosaka Y, Suzuki K. Evaluation of des-gamma-carboxy prothrombin as a marker protein of hepatocellular carcinoma. *Cancer* 1989; **64**: 2546–51.
- Liebman HA. Isolation and characterization of a hepatoma-associated abnormal (des-gamma-carboxy) prothrombin. *Cancer Res* 1989; **49**: 6493–7.
- Uehara S, Gotoh K, Handa H, Tomita H, Senshuu M. Distribution of the heterogeneity of des-gamma-carboxyprothrombin in patients with hepatocellular carcinoma. *J Gastroenterol Hepatol* 2005; **20**: 1545–52.
- Naraki T, Kohno N, Saito H *et al*. Gamma-carboxyglutamic acid content of hepatocellular carcinoma-associated des-gamma-carboxy prothrombin. *Biochem Biophys Acta* 2002; **1586**: 287–98.
- Bruix J, Sherman M. Management of hepatocellular carcinoma. *Hepatology* 2005; **42**: 1208–36.
- Liver Cancer Study Group of Japan. *The General Rules for the Clinical and Pathological Study of Primary Liver Cancer*, English edn. Tokyo: Kanehara, 2003.
- Kagebayashi C, Yamaguchi I, Akinaga A *et al*. Automated immunoassay system for AFP-L3% using on-chip electrokinetic reaction and separation by affinity electrophoresis. *Anal Biochem* 2009; **388**: 306–11.
- Oka H, Tamori A, Kuroki T, Kobayashi K, Yamamoto S. Prospective study of alpha-fetoprotein in cirrhotic patients monitored for development of hepatocellular carcinoma. *Hepatology* 1994; **19**: 61–6.
- Koda M, Murawaki Y, Mitsuda A *et al*. Predictive factors for intrahepatic recurrence after percutaneous ethanol injection therapy for small hepatocellular carcinoma. *Cancer* 2000; **88**: 529–37.
- Toyoda H, Kumada T, Tada T *et al*. Clinical utility of highly sensitive Lens culinaris agglutinin-reactive alpha-fetoprotein in hepatocellular carcinoma patients with alpha-fetoprotein < 20 ng/mL. *Cancer Sci* 2011; **102**: 1025–31.
- Parkin DM, Bray F, Ferlay J, Pisani P. Global cancer statistics, 2002. *CA Cancer J Clin* 2002; **55**: 74–108.
- Befeler AS, DiBisceglie AM. Hepatocellular carcinoma: diagnosis and treatment. *Gastroenterology* 2002; **122**: 1609–19.
- Umemura T, Kiyosawa K. Epidemiology of hepatocellular carcinoma in Japan. *Hepatol Res* 2007; **37**: S95–100.
- El-Serag HB, Mason AC. Rising incidence of hepatocellular carcinoma in the United States. *N Engl J Med* 1999; **340**: 745–50.
- El-Serag HB, Davila JA, Petersen NJ, McGlynn KA. The continuing increase in the incidence of hepatocellular carcinoma in the United States: an update. *Ann Intern Med* 2003; **139**: 817–23.
- Tsukuma H, Hiyama T, Tanaka S *et al*. Risk factors for hepatocellular carcinoma among patients with chronic liver disease. *N Engl J Med* 1993; **328**: 1797–801.
- Taketa K, Sekiya C, Namiki N *et al*. Lectin-reactive profiles of alpha-fetoprotein characterizing hepatocellular carcinoma and related conditions. *Gastroenterology* 1990; **99**: 508–18.
- Fujiyama S, Tanaka M, Maeda S, Ashihara H, Hirata R, Tomita K. Tumor markers in early diagnosis, follow-up and management of patients with hepatocellular carcinoma. *Oncology* 2002; **62**: S57–63.
- Koike Y, Shiratori Y, Sato S *et al*. Des-gamma-carboxy prothrombin as a useful predisposing factor for the development of portal venous invasion in patients with hepatocellular carcinoma: a prospective analysis of 227 patients. *Cancer* 2001; **91**: 561–9.
- Toyoda H, Kumada T, Kaneoka Y *et al*. Prognostic value of pretreatment levels of tumor markers for hepatocellular carcinoma on survival after curative treatment of patients with HCC. *J Hepatol* 2008; **49**: 223–32.
- Kobayashi M, Ikeda K, Kawamura Y *et al*. High serum des-gamma-carboxy prothrombin level predicts poor prognosis after radiofrequency ablation of hepatocellular carcinoma. *Cancer* 2009; **115**: 571–80.
- Nomura F, Ishijima M, Kuwa K, Tanaka N, Nakai T, Ohnishi K. Serum des-gamma-carboxy prothrombin levels determined by a new generation of sensitive immunoassay in patients with small-sized hepatocellular carcinoma. *Am J Gastroenterol* 1999; **94**: 650–4.
- Toyoda H, Kumada T, Kiriya S *et al*. Prognostic significance of simultaneous measurement of three tumor markers in patients with hepatocellular carcinoma. *Clin Gastroenterol Hepatol* 2006; **4**: 111–7.
- Toyoda H, Kumada T, Osaki Y *et al*. Staging hepatocellular carcinoma by a novel scoring system (BALAD score) based on serum markers. *Clin Gastroenterol Hepatol* 2006; **4**: 1528–36.

Letter to the Editor

DOI:10.1111/j.1478-3223.2012.02763.x

Incidence of hepatocellular carcinoma and response to interferon therapy in HCV-infected patients: effect of factors associated with the therapeutic response and incidence of HCC

To the Editor:

Several previous studies reported a significantly lower incidence of hepatocellular carcinoma (HCC) in hepatitis C virus (HCV)-infected patients who showed sustained virological response (SVR) to or relapsed on antiviral therapy with interferon (IFN) or peginterferon (PEG-IFN), with or without ribavirin compared with no responders (NR; i.e. partial response, viral breakthrough, or null-response) (1, 2). The reduction in HCC incidence was especially marked in patients with SVR. These results have been taken as evidence that antiviral therapy has an effect of suppressing the development of HCC.

Recently reported viral and host factors that are strongly associated with response to anti-HCV therapy (3, 4) may also be associated with the pathogenesis of HCC. Amino acid substitution in the HCV core region, a viral factor reportedly associated with response to PEG-IFN and ribavirin therapy (3), is also associated with the development of HCC (5). Regarding host factors associated with response to anti-HCV therapy (4), genetic polymorphisms near the *IL28B* gene are reportedly associated with hepatic steatosis (6) and interact with amino acid substitutions in the HCV core region (7), both of which are associated with the development of HCC (5, 8).

We analysed the incidence of HCC in 448 patients who completed anti-HCV therapy with IFN or PEG-IFN and in whom the genetic polymorphisms near *IL28B* gene were analysed after the approval of the

hospital ethics committee and obtaining written informed consent. We found significant differences in the incidence of HCC between patients with SVR ($n = 247$), relapse ($n = 122$), and NR ($n = 79$) (Fig. 1A, $P < 0.0001$ by Log-rank test). However, the prevalence of patients having TT genotype at rs8099917 near the *IL28B* gene, which is associated with favourable response to anti-HCV therapy, was significantly lower in patients with NR (SVR, 85.8%; relapse, 80.3%; NR, 39.2%; $P < 0.0001$ by Chi-square test). In addition, we found significant differences in the incidence of HCC also according to the genotype of rs8099917 (Fig. 1B, $P = 0.0156$). Although multivariate analysis using Cox proportional hazard model including age, gender, HCV genotype, and the outcome of therapy, but not *IL28B* polymorphisms identified SVR ($P = 0.0083$) and relapse ($P = 0.0493$) as independent factors that were associated with lower incidence of HCC, it failed to detect an independent factor that was associated with the incidence of HCC when *IL28B* polymorphisms were included. These results suggested that the previously reported differences in the incidence of HCC by anti-HCV response may not have been due to the ability of antiviral therapy to suppress HCC, but rather simply they may reflect the ability of such treatment to better identify patients at high-risk for HCC based on response to anti-HCV therapy.

It is not elucidated whether the results in our present analyses were simply due to the small number of




Fig. 1. Cumulative incidence of hepatocellular carcinoma (HCC) after antiviral therapy with interferon or peginterferon. (A) Incidence of HCC is significantly lower in patients with sustained virological response (SVR), those with relapse, and those with no response (NR) that includes partial response, viral breakthrough, or null-response, in that order. (B) Incidence of HCC is significantly lower in patients with TT genotype at rs8099917 near the *IL28B* gene, which is associated with the favourable response to antiviral therapy.

patients analysed or the incidence of HCC after antiviral therapy is similar regardless of response, when they are stratified by host and viral factors. In addition, our present analyses failed to examine the association between amino acid substitutions in the HCV core region and the incidence of HCC due to the small number of patients in whom the information of this substitution was available. Nonetheless, with the emergence of factors that can be independently associated with both the response to antiviral therapy and the development of HCC, the effect of response to antiviral therapy with IFN or PEG-IFN on the incidence of HCC will require re-examination taking *IL28B* polymorphisms and amino acid substitutions in the HCV core region into consideration.

Hidenori Toyoda and Takashi Kumada
Department of Gastroenterology, Ogaki Municipal Hospital,
Ogaki, Japan

References

1. Imai Y, Kawata S, Tamura S, *et al.* Relation of interferon therapy and hepatocellular carcinoma in patients with chronic hepatitis C. *Ann Intern Med* 1998; **129**: 94–9.
2. Kasahara A, Hayashi N, Mochizuki K, *et al.* Risk factors for hepatocellular carcinoma and its incidence after interferon treatment in patients with chronic hepatitis C. *Hepatology* 1998; **27**: 1394–402.
3. Akuta N, Suzuki F, Sezaki H, *et al.* Association of amino acid substitution pattern in core protein of hepatitis C virus genotype 1b high viral load and non-virological response to interferon-ribavirin combination therapy. *Intervirology* 2005; **48**: 372–80.
4. Ge D, Fellay J, Thompson AJ, *et al.* Genetic variation in *IL28B* predicts hepatitis C treatment-induced viral clearance. *Nature* 2009; **461**: 399–401.
5. Akuta N, Suzuki F, Kawamura Y, *et al.* Amino acid substitutions in the hepatitis C virus core region are the important predictor of hepatocarcinogenesis. *Hepatology* 2007; **46**: 1357–64.
6. Tillmann HL, Patel K, Muir AJ, *et al.* Beneficial *IL28B* genotype associated with lower frequency of hepatic steatosis in patients with chronic hepatitis C. *J Hepatol* 2011; **55**: 1195–200.
7. Abe H, Ochi H, Maekawa T, *et al.* Common variation of *IL28B* affects gamma-GTP levels and inflammation of the liver in chronically infected hepatitis C virus patients. *J Hepatol* 2010; **53**: 439–43.
8. Pekow JR, Bhan AK, Zheng H, Chung RT. Hepatic steatosis is associated with increased frequency of hepatocellular carcinoma in patients with hepatitis C-related cirrhosis. *Cancer* 2007; **109**: 2490–6.