

phosphorylation, whereby it translocates to the nucleus to act as a transcriptional factor for positive regulatory domain (PRD) I of the IFN- $\beta$  promoter. IRF-3 activation also induces phosphorylation of Ser385 and Ser386 or the serine/threonine (Ser/Thr) cluster between amino acids (aa) 396 and 405 (located at the C-terminus of IRF-3), a step that is essential for dimerization and nuclear translocation.

A previous study by Foy et al. showed that the NS3/4A serine protease derived from a subgenomic replicon participates in the suppression of the cellular pathway that activates IRF-3 [12]. The NS3/4A protein of HCV disrupts signaling of the double-stranded RNA (dsRNA) receptors, retinoic acid-inducible gene-1 (RIG-1), and Toll-like receptor 3 (TLR3) by inducing proteolysis of interferon promoter stimulator-1 (IPS-1) [13–15] and Toll/interleukin-1 receptor (TIR) domain-containing adaptor protein inducing IFN- $\beta$  (TRIF) [16], as well as by suppressing the downstream activation of IFN- $\beta$  [13].


In view of the above observations and the emerging data on the role of HCV in regulating the IRF-3 pathway by additional

mechanisms, we aimed to investigate the effects of various HCV proteins on IRF-3 activation, and further elucidate the underlying novel mechanisms.

## 2. Material and methods

### 2.1. Transient expression of the HCV core proteins E1, E2, and NS3-4A

HepG2 cells were transfected to express E1, E2, or NS3-4A HCV core protein under the control of EF promoter (Invitrogen). The HCV core expression vectors were derived from HCR6 (genotype 1b), HCR24-12K (genotype 2a), or HCR24-12Q (genotype 2a). The E1, E2, and NS3 clones derived from HCR6 contained either the full-length cDNAs encoding the core protein or 1 of the 3 different deletions (deletion mutants), each of which lacked 1 of the 3 basic amino acid regions (BR), BR1 deletion (aa 4–14), BR2 deletion (aa 37–44), and BR3 deletion (aa 57–72) [17]. HepG2 cells were transfected with 4  $\mu$ g of the core cDNA (amino acids 1–191), E1 (amino



**Fig. 1.** (A) Structures of the conditional expression vectors for HCV RNAs and proteins. The cDNA clones that displayed highest level of homology to the consensus sequences among the 3 clones were used to construct *HCR6-Rz* (nt 1–9611). *HCR6-Fse* clone harbored a termination codon introduced at nucleotide 3606 and truncated *HCR6-Age* clone lacked nucleotides 155–1967. They were flanked with ribozyme (Rz) and hepatitis D virus ribozyme (HDV-Rz) sequences under the control of the CAG promoter in the Cre/loxP switching expression cassette, which consisted of the neomycin resistance gene, as a stuffer region flanked by the loxP sequence. (B) IFN- $\beta$  mRNA levels in the cell lines *HCR6-Rz*, *HCR6-Fse*, and *HCR6-Age* before and after the expression of the HCV protein. The results are expressed as copy numbers per microgram of total RNA, as quantified by RT-PCR. (C) Efficiency of NDV plaque formation. Plaque assays were performed on Vero cells for NDV infectivity in *HCR6-Rz*, *HCR6-Fse*, or *HCR6-Age* before and after the expression of the HCV protein. The plaque numbers were counted 3 days after NDV inoculation. The ratio indicates the plaque numbers after the expression of the HCV genome divided by the plaque numbers before the expression of the HCV genome. (D) Suppression of IFN- $\beta$  mRNA induction by HCV expression, 18 days after NDV inoculation. Day 0, before the expression of the HCV genome; Day 48, after the expression of the HCV genome. The results are expressed relative to the levels on Day 0 (100%) in each of the 3 HCV-expressing systems. p.i.h., post-inoculation hour.

acids 192–383), E2 (amino acids 384–809), NS3-4A (amino acids 1027–1711), or the core regions lacking the BR, in 35-mm dishes, by using Lipofectamine 2000 (Invitrogen) at 37 °C for 6 h. The medium was subsequently replaced with normal culture medium, and the cells were harvested after 48 h.

### 2.2. Newcastle disease virus (NDV) infection and addition of polyriboinosinic:polyribocytidylic acid [poly(I:C)]

NDV (Miyadera strain) was propagated from swabs by using the embryonated egg culture method, as described in the Supplementary methods.

Poly(I:C) (20 µg/well; GE Healthcare) was added to HepG2 cells in 35-mm dishes, 48 h after transfection with the core expression vector derived from HCR6 (genotype 1b).

### 2.3. Native PAGE of IRF-3 and phosphorylated IRF-3

Cells were lysed in 30 µL of lysis buffer (50 mM Tris-HCl [pH 8.0], 1% NP-40, 150 mM NaCl, 100 µg/mL leupeptin, 1 mM PMSF, 5 mM Na<sub>3</sub>VO<sub>4</sub>), mixed vigorously, and centrifuged at 15,000 rpm for 10 min; the supernatant was isolated. Total protein samples (10 µg) were electrophoresed on a 7.5% native PAGE gel (Bio-Rad Laboratories) and transferred onto a PVDF membrane. IRF-3 homodimer and its monomer and Ser386-phosphorylated IRF-3 were detected by western blotting using polyclonal rabbit antibodies, anti-human IRF-3 (1:1000) [18] and anti-human Ser386-phosphorylated IRF-3 (1:10,000), respectively. Detection was achieved by enhanced chemiluminescence (ECL; Amersham, UK) according to the manufacturer's instructions. The rabbit anti-human IRF-3 and anti-human Ser386-phosphorylated IRF-3 antibodies were described previously [19].

### 2.4. Quantification of IFN-β gene expression

IFN-β mRNA expression was quantified by using real-time PCR (RT-PCR) as described previously [20] and Supplementary methods using the following primers and probes: sense (5'-CCATCTATGATGATGCTCCAGAA-3'), antisense (5'-TTTTCTTCCAGACTGTCTTCA-GA-3') and probe (5'-AGCACTGGCTGGAATGAGACTATTGTG-3').

## 3. Results


### 3.1. Induction of IFN and IRF-3 by HCV-Rz

To evaluate the effect of HCV gene persistent expression, cell lines expressing the HCV genomes in a Cre/loxP expression system [20] were established by transfecting the full-genome HCV (*HCR6-Rz*); core, E1, E2, and NS2 (*HCR6-Fse*); and E2~NS5b (*HCR6-Age*) (Fig. 1A) clones into HepG2 cells. Of the 3 HCV expression systems, only *HCR6-Rz* transiently induced endogenous IFN-β expression (Fig. 1B). No endogenous IFN-β was detected by RT-PCR on Day 0 or Day 48 (long culture; LC) [21] in any of the 3 systems (Fig. 1B).


### 3.2. Effect of HCV expression on NDV infection and IFN-β induction

Further, we examined whether the persistent expression of HCV genome influenced the plaque formation activity of NDV in *HCR6-Rz*-, *HCR6-Fse*-, and *HCR6-Age*-expressing cell lines. Prior to the expression of HCV proteins, the plaque numbers were similar across all the cell lines (Fig. 1C). Expression of HCV proteins for more than 48 days [21] increased plaque numbers in the *HCR6-Rz*- and *HCR6-Fse*-expressing cells from 7- to 8-fold (Fig. 1C); however, plaque numbers in the *HCR6-Age*-expressing cells remained constant. These findings thus indicate that HCV structural proteins

interfered with the induction of IFN-β mRNA, after NDV infection (18 h post-infection). Therefore, we measured IFN-β mRNA levels in the 3 cell lines, *HCR6-Rz*, *HCR6-Fse*, and *HCR6-Age*, by RT-PCR before (Day 0) and after (Day 48) inoculation with NDV (Fig. 1D). The mRNA expression of IFN-β was not observed prior to NDV infection in any of the 3 cell lines (Day 0). Notably, after 18 h of NDV inoculation and prior to the expression of various HCV proteins (Day 0), the levels of IFN-β mRNA transcription were similar among *HCR6-Rz*-, *HCR6-Fse*-, and *HCR6-Age*-expressing cell lines (Fig. 1D). Following HCV protein expression, the induced IFN-β mRNA expression was reduced to 20% in both *HCR6-Rz*- and *HCR6-Fse*-expressing cell lines. However, IFN-β mRNA expression remained constant in the *HCR6-Age*-expressing cell line.



**Fig. 2.** (A) HCV inhibition of IRF-3 nuclear translocation. IRF-3 is a constitutively expressed transcriptional factor that localizes in the cytoplasm in a diffuse manner, when inactive. After NDV inoculation and prior to the expression of the HCV genome, IRF-3 translocated to the nucleus but was retained at a perinuclear site in the *HCR6-Rz*-expressing cells (Day 48). IRF-3 colocalized with the HCV core protein. (B–D) Suppression of IRF-3 dimerization by HCV expression. The dimeric and monomeric forms of IRF-3 were detected by western blotting on native PAGE gels. The influence of (B) *HCR6-Rz* expression, (C) *HCR6-Fse* expression, and (D) *HCR6-Age* expression on IRF-3 dimerization is shown. NDV was used to induce IRF-3 dimerization.



**Fig. 3.** (A) Structures of the HCR6 core, E1, and E2 expression vectors encoding the HCV core (aa 1–191), E1 (amino acids 192–383), and E2 (amino acids 384–809) proteins, respectively, under the control of the EF1 promoter. (B) Left panel, IRF-3 dimerization induced by NDV before transfection with the expression vectors; right panel, IRF-3 dimerization induced by NDV after transfection with the expression vectors. (C) IFN-β mRNA induction levels at 12 and 18 h after NDV inoculation into HepG2 cells transfected with the vector alone, core, E1, or E2 proteins. The results are expressed relative to the induction levels of IFN-β in HepG2 cells transfected with the vector alone (100%), for post-inoculation, each time. (D) IRF-3 dimerization on administration of 10 mg/mL poly(I:C) before the expression (Day 0), and at 2, 4, 8, and 12 h after the expression of the HCV core (pEF-Core) or vector plasmid DNA. (E) Effects of the expression of vector (lane 1), R6 core (lane 2), R24 core (lane 3), R24 core (Q12K) (lane 4), R6-NS3-4A (lane 5), N1-NS3-4A (lane 6), and mock control (lane 7) on IRF-3 dimerization (upper column a) and phosphorylation of serine residue at amino acid 386 in IRF-3 (lower column b), after infection with NDV for 18 h. (F) Effects of the expressions of R6-C, R24-C, R24-C(Q12K), and R6-NS3 on IFN-β induction, 18 h after NDV inoculation. The IFN-β mRNA levels were assayed by RT-PCR. The results are expressed relative to the induction levels of IFN-β in HepG2 cells transfected with the vector alone (100%).

### 3.3. Effect of HCV on IRF-3 localization, nuclear translocation, and dimerization by NDV

The effect of HCV expression on cellular localization of IRF-3 was analyzed in HCR6-Rz-expressing cells infected with NDV

before (Day 0) and after LC (Fig. 2A). Prior to NDV infection, IRF-3 was detected in the cytoplasm by immunofluorescence. Notably, after 18 h of NDV inoculation and prior to HCR6-Rz protein expression, when IFN-β induction and IRF-3 dimerization were maximal, a substantial amount of IRF-3 translocated to the nucleus.

However, this nuclear translocation was suppressed in the presence of HCV proteins (Fig. 2A) and resulted in the co-localization of the HCV core protein with IRF-3 at perinuclear sites (Fig. 2A, superimposed image of IRF-3 and core protein immunostaining).

To elucidate the mechanism underlying the suppression of IFN- $\beta$  mRNA in *HCR6-Rz*- and *HCR6-Fse*-expressing cells, we examined the effect of HCV expression on IRF-3 dimerization after NDV infection (Fig. 2B). Interestingly, the levels of IRF-3 dimerization peaked at 12–18 h after NDV infection in the 3 cell lines lacking HCV expression (Day 0; Fig. 2). However, in the *HCR6-Rz*- and *HCR6-Fse*-expressing cell lines, IRF-3 dimerization was found to be significantly reduced, (Fig. 2B and C) when compared to that in the *HCR6-Age*-expressing cells (Fig. 2D).

#### 3.4. Identification of the HCV genome region responsible for the inhibition of IRF-3 dimerization and IFN- $\beta$ induction

To identify the HCV genome region responsible for suppression of IRF-3 dimerization, HepG2 cells were transfected to express the HCV core regions derived from HCR6, E1, or E2 (genotype 1b; Fig. 3A). Protein expression was confirmed by western blotting (data not shown). The HCV core protein suppressed IRF-3 dimerization, but E1 and E2 expressions had no effect on the dimerization (Fig. 3B). Expression of E1, E2, or the vector alone did not alter the levels of IFN- $\beta$  mRNA induced by NDV infection in HepG2 cells (Fig. 3C), but significantly reduced IFN- $\beta$  mRNA levels at both 12 and 18 h after infection (Fig. 3C).

#### 3.5. Effect of HCV core protein expression on IRF-3 dimerization through TLR3

Among the synthetic dsRNAs, poly(I:C) is a potent inducer of IFN- $\beta$  through TLR3. Accordingly, HepG2 cells transfected with poly(I:C) and the vector control (pEF1-vector) showed IRF-3 dimerization (Fig. 3D). In contrast, IRF-3 dimerization was suppressed in HepG2 cells expressing the HCV core protein albeit the induction of IFN- $\beta$  mRNA following poly(I:C) expression (data not shown).

#### 3.6. Effect of the HCV core protein NS3 and core proteins derived from genotype 2a on IRF-3 dimerization, compared to proteins derived from genotype 1b

Further, we investigated whether the HCV core protein NS3 and core proteins derived from other genotypes exerted the same effects on IRF-3 after 18 h of NDV infection. The core proteins derived from genotypes 1b (R6) and 2a (R24-12Q and R24-12K) suppressed IRF-3 dimerization in cells infected with NDV (Fig. 3E(a)). In contrast, IRF-3 dimerization remained unaltered in the presence of R6 clone NS3 protein, but was suppressed by the N clone NS3 protein. Thus, NS3-4A protein of R6 clone suppressed IRF-3 dimerization to a relatively lesser extent compared to that of the N1 strain. Similar results were obtained for the phosphorylation at Ser386 in IRF-3 (Fig. 3E(b)). The IFN- $\beta$  mRNA transcription was quantified in HepG2 cells by RT-PCR after transfection with these expression vectors (Fig. 3F).

#### 3.7. Identification of the HCV core region responsible for suppressing IRF-3 dimerization

We sought to identify the region of the HCV core protein responsible for suppressing IRF-3 dimerization. Expression vectors encoding the entire HCV core or the core region lacking 1 of the 3 basic amino acid regions (BR) that influenced nuclear translocation [17] were transfected into HepG2 cells, and the effects on IRF-3 dimerization were examined (Fig. 4A). Protein




Fig. 4. (A) Structures of the HCR6 core, E1, and E2 expression vectors carrying the complete core, BR1 deletion (aa 4–14), BR2 deletion (aa 37–44), and BR3 deletion (aa 57–72). (B) Western blotting to confirm the expression of the mutated core proteins. (C) Effects of the expression of various mutated core proteins on IRF-3 dimerization and IRF-3 phosphorylation at Ser386, 18 h after NDV inoculation. (D) Effects of the expression of each type of core protein region on IFN- $\beta$  mRNA synthesis, 18 h after NDV inoculation. The results are expressed relative to the induction levels of IFN- $\beta$  in HepG2 cells transfected with the vector alone (100%). IFN- $\beta$  mRNA levels were assayed by RT-PCR.

expression of the core and the deletion mutants (BR1, BR2, and BR3) was confirmed by western blotting (Fig. 4B). IRF-3 dimerization, phosphorylation at Ser386 of IRF-3, and induction of IFN- $\beta$  mRNA were suppressed in HepG2 cells expressing the entire core, a deletion of BR2, or a deletion of BR3 (Fig. 4C), but not in cells expressing the BR1-deleted HCV core regions (Fig. 4C and D).

#### 4. Discussion

The present study indicates that the HCV core protein inhibits IRF-3 dimerization, IRF-3 phosphorylation at Ser386, and IFN- $\beta$  induction. In addition, our study showed that the effect of the core protein derived from genotype 1b was similar to that of the core protein derived from genotype 2a, indicating that the inhibitory effect of the core protein might be effective in several genotypes of HCV. These findings are corroborated by a previous study by Foy et al. [12] who showed that HCV NS3/4 disrupts virus-associated-kinase-mediated IRF-3 activation, which further results in the suppression of IRF-3 phosphorylation, nuclear translocation, and IRF-3-dependent ISRE/PRDI activation. These findings indicate that attenuation of the IFN system was achieved through NS3/4A proteins via the interference of IRF-3 activation, thus strengthening our results, which show the potential of HCV core protein to interfere with IRF-3 activation in promoting persistent infection.

Furthermore, the present study showed that the N-terminal region of the core protein and BR-1 domain in particular are responsible for inactivating IRF-3. The N-terminal region (amino acids 1–59) of the HCV core protein has been identified as the binding region for a DEAD box protein (DDX3) [22]. Human DDX3, a putative RNA helicase, is a member of the highly conserved DEAD box subclass that includes the expression of murine PL10, *Xenopus* An3, and yeast Ded 1 proteins. Recently, expression of DDX3 was found to enhance IFN- $\beta$  promoter induction by TBK1/IKK $\epsilon$ , whereas silencing of DDX3 inhibited IFN- $\beta$  promoter and virus- or dsRNA-induced IRF-3 activation [23]. It was shown that Vaccinia virus K7 protein also binds to DDX3 and inhibits pattern recognition receptor-induced IFN- $\beta$  induction by preventing TBK1/IKK $\epsilon$ -mediated IFN- $\beta$  induction via impaired TBK1/IKK $\epsilon$ -induced activation of IRF-3 [23]. A previous study by Oshiumi et al. showed that DDX3 C-terminal region (amino acids 622–662) directly binds to the IFN-beta promoter stimulator-1 (IPS-1) CARD-like domain [24] as well as the N-terminal HCV core protein [36]. The present study demonstrated that the expression of the core protein decreased the levels of DDX3 expression (data not shown). This is in agreement with the result of a previous study, which showed that DDX3 is downregulated in HCV-associated hepatocellular carcinoma (HCC) and silencing of DDX3 accelerates cell growth [25]. Collectively, these findings suggest that DDX3 may be the target of the core protein for inhibiting IRF-3 activation.

In conclusion, our study revealed a crucial region of the HCV core protein, basic amino acid region 1, to interfere with IRF-3 activation and thereby inhibit the IFN signaling cascades. Therefore, the inhibitory effects that result in the IRF-3 pathway impairment could be rescued by deleting the basic region 1 of core protein, thus suggesting that it might be an effective treatment for HCV infection. Future studies involving DDX3 modification by the HCV core protein may be interesting to explore the cell growth-dysregulation mechanisms.

#### Acknowledgments

This study was supported in part by a grant from the Ministry of Education, Culture, Sports, Science and Technology of Japan; a grant from the Ministry of Health, Labour and Welfare of Japan; and the Program for Promotion of Fundamental Studies in Health Sciences of the National Institute of Biomedical Innovation of Japan; and the Cooperative Research Project on Clinical and

Epidemiological Studies of Emerging and Re-emerging Infectious Diseases.

#### Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.bbrc.2012.10.079>.

#### References

- [1] N. Kato, M. Hijikata, Y. Ootsuyama, et al., Molecular cloning of the human hepatitis C virus genome from Japanese patients with non-A, non-B hepatitis, *Proc. Natl. Acad. Sci. USA* 87 (1990) 9524–9528.
- [2] L.B. Seeff, F.B. Hollinger, H.J. Alter, et al., Long-term mortality and morbidity of transfusion-associated non-A, non-B, and type C hepatitis: a National Heart, Lung, and Blood Institute collaborative study, *Hepatology* 33 (2001) 455–463.
- [3] H.B. El-Serag, Hepatocellular carcinoma and hepatitis C in the United States, *Hepatology* 36 (2002) S74–S83.
- [4] D. Thomas, The hepatitis C viruses, in: C. Hagedorn, C. Rice (Eds.), *Hepatitis C Epidemiology*, Springer-Verlag, Berlin, 1999, pp. 25–42.
- [5] G.R. Foster, Interferons in host defense, *Semin. Liver Dis.* 17 (1997) 287–295.
- [6] R. Kaufman, Translation control of genome expression, in: N. Sonenberg, J. Hershey, M. Matthews (Eds.), *The double strand RNA-activated protein kinase PKR*, Cold Spring Harbor, New York, 2000, pp. 503–527.
- [7] M.J. Gale Jr., M.J. Korth, N.M. Tang, et al., Evidence that hepatitis C virus resistance to interferon is mediated through repression of the PKR protein kinase by the nonstructural 5A protein, *Virology* 230 (1997) 217–227.
- [8] D.R. Taylor, S.T. Shi, P.R. Romano, et al., Inhibition of the interferon-inducible protein kinase PKR by HCV E2 protein, *Science* 285 (1999) 107–110.
- [9] M. Yoneyama, W. Suhara, Y. Fukuhara, et al., Direct triggering of the type I interferon system by virus infection: activation of a transcription factor complex containing IRF-3 and CBP/p300, *Embo. J.* 17 (1998) 1087–1095.
- [10] J. Hiscott, Triggering the innate antiviral response through IRF-3 activation, *J. Biol. Chem.* 282 (2007) 15325–15329.
- [11] E.F. Meurs, A. Breiman, The interferon inducing pathways and the hepatitis C virus, *World J. Gastroenterol.* 13 (2007) 2446–2454.
- [12] E. Foy, K. Li, C. Wang, et al., Regulation of interferon regulatory factor-3 by the hepatitis C virus serine protease, *Science* 300 (2003) 1145–1148.
- [13] E. Meylan, J. Curran, K. Hofmann, et al., Cardif is an adaptor protein in the RIG-I antiviral pathway and is targeted by hepatitis C virus, *Nature* 437 (2005) 1167–1172.
- [14] X.D. Li, L. Sun, R.B. Seth, et al., Hepatitis C virus protease NS3/4A cleaves mitochondrial antiviral signaling protein off the mitochondria to evade innate immunity, *Proc. Natl. Acad. Sci. USA* 102 (2005) 17717–17722.
- [15] R. Lin, J. Lacoste, P. Nakhaei, et al., Dissociation of a MAVS/IPS-1/VISA/Cardif-IKKEpsilon molecular complex from the mitochondrial outer membrane by hepatitis C virus NS3-4A proteolytic cleavage, *J. Virol.* 80 (2006) 6072–6083.
- [16] K. Li, E. Foy, J.C. Ferreon, et al., Immune evasion by hepatitis C virus NS3/4A protease-mediated cleavage of the Toll-like receptor 3 adaptor protein TRIF, *Proc. Natl. Acad. Sci. USA* 102 (2005) 2992–2997.
- [17] R. Suzuki, Y. Matsuura, T. Suzuki, et al., Nuclear localization of the truncated hepatitis C virus core protein with its hydrophobic C terminus deleted, *J. Gen. Virol.* 76 (Pt 1) (1995) 53–61.
- [18] W. Suhara, M. Yoneyama, I. Kitabayashi, et al., Direct involvement of CREB-binding protein/p300 in sequence-specific DNA binding of virus-activated interferon regulatory factor-3 holocomplex, *J. Biol. Chem.* 277 (2002) 22304–22313.
- [19] M. Mori, M. Yoneyama, T. Ito, et al., Identification of Ser-386 of interferon regulatory factor 3 as critical target for inducible phosphorylation that determines activation, *J. Biol. Chem.* 279 (2004) 9698–9702.
- [20] K. Tsukiyama-Kohara, S. Tone, I. Maruyama, et al., Activation of the CKI-CDK-Rb-E2F pathway in full genome hepatitis C virus-expressing cells, *J. Biol. Chem.* 279 (2004) 14531–14541.
- [21] T. Nishimura, M. Kohara, K. Izumi, et al., Hepatitis C virus impairs p53 via persistent overexpression of 3beta-hydroxysterol Delta24-reductase, *J. Biol. Chem.* 284 (2009) 36442–36452.
- [22] A.M. Owsianka, A.H. Patel, Hepatitis C virus core protein interacts with a human DEAD box protein DDX3, *Virology* 257 (1999) 330–340.
- [23] M. Schroder, M. Baran, A.G. Bowie, Viral targeting of DEAD box protein 3 reveals its role in TBK1/IKKEpsilon-mediated IRF activation, *EMBO J.* 27 (2008) 2147–2157.
- [24] H. Oshiumi, K. Sakai, M. Matsumoto, T. Seya, DEAD/H BOX 3 (DDX3) helicase binds the RIG-I adaptor IPS-1 to up-regulate IFN-beta-inducing potential, *Eur. J. Immunol.* 40 (4) (2010) 940–948.
- [25] P.C. Chang, C.W. Chi, G.Y. Chau, et al., DDX3, a DEAD box RNA helicase, is deregulated in hepatitis virus-associated hepatocellular carcinoma and is involved in cell growth control, *Oncogene* 25 (2006) 1991–2003.

# Hepatitis C Virus Promotes Expression of the $3\beta$ -Hydroxysterol $\Delta$ 24-Reductase Through Sp1

Makoto Saito,<sup>1</sup> Michinori Kohara,<sup>2</sup> and Kyoko Tsukiyama-Kohara<sup>1\*</sup>

<sup>1</sup>Department of Experimental Phylaxiology, Faculty of Life Sciences, Kumamoto University, Kumamoto, Japan

<sup>2</sup>Department of Microbiology and Cell Biology, Tokyo Metropolitan Institute of Medical Science, Tokyo, Japan

Hepatitis C virus (HCV) establishes chronic infection, which often causes hepatocellular carcinoma. Overexpression of  $3\beta$ -hydroxysterol  $\Delta$ 24-reductase (DHCR24) by HCV has been shown to impair the p53-mediated cellular response, resulting in tumorigenesis. In the present study, the molecular mechanism by which HCV promotes the expression of DHCR24 was investigated. A significant increase in DHCR24 mRNA transcription was observed in a cell line expressing complete HCV genome, whereas no significant difference in the expression of DHCR24 was seen in cell lines expressing individual viral proteins. The 5'-flanking genomic region of DHCR24 was characterized to explore the genomic region and host factor(s) involved in the transcriptional regulation of DHCR24. As a result, the HCV response element (–167/–140) was identified, which contains AP-2 $\alpha$ , MZF-1, and Sp1 binding motifs. The binding affinity of the host factor to this response element was increased in nuclear extracts from cells infected with HCV and corresponded with augmented affinity of Sp1. Both mithramycin A (Sp1 inhibitor) and small interfering RNA targeting Sp1 prevented the binding of host factors to the response element. Silencing of Sp1 also downregulated the increased expression of DHCR24. The binding affinity of Sp1 to the response element was augmented by oxidative stress, whereas upregulation of DHCR24 in cells expressing HCV was blocked significantly by a reactive oxygen species scavenger. Elevated phosphorylation of Sp1 in response to oxidative stress was mediated by the ATM kinase. Thus, activation of Sp1 by oxidative stress is involved in the promotion of expression of DHCR24 by HCV. *J. Med. Virol.* 84:733–746, 2012. © 2012 Wiley Periodicals, Inc.

**KEY WORDS:** HCV; DHCR24; Sp1; oxidative stress

## INTRODUCTION

Hepatitis C virus (HCV) causes chronic hepatitis and hepatocellular carcinoma [Koike, 2007]. The estimated worldwide prevalence of HCV infection is 2.2–3.0% (130–170 million people) [Lavanchy, 2009], and chronic HCV infection is a major global public health concern. The most effective current treatment for HCV infection comprises combination therapy with PEGylated interferon- $\alpha$  and ribavirin [Bruchfeld et al., 2001; Lu et al., 2008]. However, this therapy has limited clinical efficacy, as sustained virological responses develop in only about half of patients infected with HCV genotype 1 [Kohara et al., 1995; Nakamura et al., 2002]. Efforts to develop therapies to treat HCV are also hindered by the high level of viral variation and the capacity of HCV to cause chronic infection. Therefore, there is an urgent need to develop effective treatments against chronic HCV infection.

Additional supporting information may be found in the online version of this article.

Grant sponsor: Ministry of Health and Welfare of Japan; Grant sponsor: Ministry of Education, Culture, Sports, Science and Technology of Japan; Grant sponsor: Program for Promotion of Fundamental Studies in Health Sciences of the National Institute of Biomedical Innovation; Grant sponsor: Cooperative Research Project on Clinical and Epidemiological Studies of Emerging and Re-emerging Infectious Diseases.

Kyoko Tsukiyama-Kohara present address is Transboundary Animal Diseases Center, Faculty of Agriculture, Kagoshima University, 1-21-24 Korimoto, Kagoshima-city 890-0065, Japan.

\*Correspondence to: Kyoko Tsukiyama-Kohara, PhD, Department of Experimental Phylaxiology, Faculty of Life Sciences, Kumamoto University, 1-1-1 Honjo, Kumamoto City, Kumamoto 860-8556, Japan. E-mail: kkohara@kumamoto-u.ac.jp, kkohara@agri.kagoshima-u.ac.jp

Accepted 16 January 2012

DOI 10.1002/jmv.23250

Published online in Wiley Online Library  
(wileyonlinelibrary.com).

A cell line that expresses complete HCV genome (RzM6-LC) was established to investigate the effects of persistent expression of HCV on cell growth [Tsukiyama-Kohara et al., 2004]. A monoclonal antibody (2-152a mAb) against the RzM6-LC cell line was also developed to produce clones that recognize both cell surface and intracellular molecules. As a result, 3 $\beta$ -hydroxysterol  $\Delta$ 24-reductase (DHCR24) was identified as the target of 2-152a mAb [Nishimura et al., 2009].

DHCR24 is an oxidoreductase with a broad expression pattern and shares homology with a family of flavin-adenine dinucleotide-dependent reductases [Waterham et al., 2001]. In mammals, DHCR24 functions as an enzyme to catalyze the conversion of desmosterol to cholesterol in the post-squalene cholesterol biosynthetic pathway, and it is essential for normal tissue development and maintenance [Waterham et al., 2001; Crameri et al., 2006]. DHCR24 regulates cholesterol synthesis and promotes recruitment of domain components into detergent-resistant membrane fractions [Crameri et al., 2006]. An absence of DHCR24 leads to desmosterolosis—a rare disorder of cholesterol biosynthesis [Waterham et al., 2001]. Expression of DHCR24 is downregulated in areas of the brain affected by Alzheimer's disease [Greeve et al., 2000], suggesting that DHCR24 has alternative functions. Indeed, DHCR24 is also known as seladin-1 (the selective Alzheimer's disease indicator 1), reflecting the association between DHCR24/seladin-1 and the selective vulnerability of the neurons in the affected areas of the brain. High levels of DHCR24/seladin-1 exert protective effects, conferring resistance against oxidative stress and preventing apoptotic cell death [Greeve et al., 2000; Benvenuti et al., 2005; Di Stasi et al., 2005; Luciani et al., 2005; Lu et al., 2008]. Endogenous DHCR24/seladin-1 levels are upregulated in response to acute oxidative stress [Wu et al., 2004; Benvenuti et al., 2006; Kuehnle et al., 2008], whereas the expression declines to low levels upon chronic exposure [Benvenuti et al., 2006; Kuehnle et al., 2008]. Therefore, DHCR24/seladin-1 may be involved in integrating cellular responses to oxidative stress. DHCR24 also functions as a hydrogen peroxide scavenger [Lu et al., 2008]. Based on these findings, DHCR24 may play a crucial role in maintaining cellular physiology by regulating both cholesterol synthesis and cellular defense against oxidative stress.

HCV infection impairs apoptosis induced by oxidative stress and inhibits p53 function via overexpression of DHCR24 [Nishimura et al., 2009]. Augmented expression of DHCR24 also facilitates efficient replication of HCV [Takano et al., 2011b]. Since DHCR24 may play a significant role in viral replication and in the tumorigenicity of the hepatocellular carcinoma related to HCV, the molecular mechanism of overexpression of DHCR24 in response to HCV was examined in the present study.

*J. Med. Virol.* DOI 10.1002/jmv

## MATERIALS AND METHODS

### Cell Lines

The HepG2 hepatoblastoma cell line, the HepG2-derived RzM6 cell line, which is capable of conditional control of expression of HCV genome (genotype 1b) based on the Cre/*loxP* system (RzM6-0d, no switching; RzM6-LC, switching of full genome HCV induced by tamoxifen), and HepG2-derived CN5 cell line, in which all HCV proteins were expressed conditionally by cre adenovirus (CN5-Cre) [Tsukiyama-Kohara et al., 2004] were propagated in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS). Using a stable expression system based on lentiviral vectors, HepG2/Lenti cell lines (core, E1, E2, NS2, NS3/4A, NS4B, NS5A, and NS5B) were established [Takano et al., 2011a]. An additional cell line—HepG2-emp—was infected with an empty lentiviral vector. Cells from the human hepatoma HuH-7 cell line were maintained in DMEM supplemented with 10% FBS and 0.4% glucose. The cell lines harboring HCV replicon, namely, R6FLR-N (genotype 1b) and FLR3-1 (genotype 1b), which are derived from HuH-7 [Takano et al., 2011b], were maintained under selective pressure with G418 (500  $\mu$ g/ml for R6FLR-N and FLR3-1) in DMEM GlutaMAX (Invitrogen, Carlsbad, CA) containing 10% FBS. Cured/HuH-7 K4 cells—cured of HCV by interferon- $\alpha$  treatment—were maintained in DMEM GlutaMAX containing 10% FBS without G418. The JFH/K4 cell line, which shows persistent infection with the HCV JFH-1 strain, was maintained in DMEM containing 10% FBS. The human fetal hepatic cell line WRL68 was obtained from the American Type Culture Collection and maintained in DMEM supplemented with 10% FBS, 1 mM sodium pyruvate, and 0.1 mM nonessential amino acids. The human hepatoma cell line PLC/PRF/5 was obtained from the Cell Resource Center for Biomedical Research Institute of Development, Aging and Cancer, Tohoku University, and cultured in Eagle's minimum essential medium supplemented with 10% FBS.

### Construction of DHCR24 Promoter Reporter Plasmids

Genomic DNA was extracted from HepG2 cells, and the 5'-flanking sequence of the predicted transcription start site of *DHCR24* (~5 kb) was isolated. The genomic region was inserted upstream of the firefly luciferase gene in pGL3-Basic (Promega, Madison, WI). Deletion constructs of the *DHCR24* promoter region were constructed using restriction enzymes and PCR (sense primer for -4956, 5'-GATCCTCGAGCACTCC-TGCTCACCCTGAT-3'; sense primer for -2982, 5'-GATCCTCGAGGAGGCTCACATTGTAGAAAG-3'; antisense primer, 5'-GTAGTAGATATCGAAGATAAGC-GAGAGCGG-3') and cloned into pGL3-Basic at the *Xho*I and *Nco*I sites.



### Dual Luciferase Reporter Assay

HepG2 cells ( $1 \times 10^4$  cells/well in a 96-well plate) were transfected with each of the 3 *DHCR24* promoter reporter plasmids and their deletion constructs (0.25  $\mu\text{g}$ /well) using cationic lipid (Lipofectamine LTX, Invitrogen). Samples were analyzed with the Dual-Glo Luciferase Assay System (Promega) at 48 h post-transfection, and luminescence was measured using a TriStar LB941 microplate reader (Berthold, Bad Wildbad, Germany). To account for differences in transfection efficiency, the luminescence produced by firefly luciferase (FL) was normalized to that produced by Renilla luciferase (RL), which was expressed by co-transfection with pRL-TK (Promega; 0.025  $\mu\text{g}$ /well).

### Electrophoresis Mobility Shift Assay

Nuclear extracts were prepared from  $5 \times 10^6$  to  $1 \times 10^7$  cells as described previously [Dignam et al., 1983]. Electrophoresis mobility shift assays (EMSA) were performed by a nonradioactive method using the DIG Gel Shift Kit (Roche, Indianapolis, IN). Briefly, binding reactions were performed by mixing the following components: 1  $\mu\text{g}$  of poly[d(I-C)], 0.1  $\mu\text{g}$  of poly L-lysine, 40 fmol DIG-labeled double-stranded oligonucleotide probe (HCV response element -167/-140 [28-mer], 5'-CCCCGCCTCGCGCGGCGGGCGG-GGAGAA-3'; Sp1 consensus sequence [22-mer], 5'-ATTCGATCGGGGCGGGCGAGC-3'; MZF1.1-4 consensus sequence [21-mer], 5'-GATCTAAAAGTGGG-GAGAAA-3'; AP-2 $\alpha$  consensus sequence [26-mer], 5'-GATCGAACTGACCGCCCGCGGCCCGT-3'), and 10  $\mu\text{g}$  of the nuclear extract in binding buffer (10 mM Tris-HCl, pH 7.5; 50 mM NaCl; 5 mM MgCl<sub>2</sub>; 0.5 mM EDTA). Where indicated, reactions were supplemented with unlabeled/competitive oligonucleotide at a 50-fold molar excess concentration before addition of the probe. Mithramycin A (MMA; Sigma, St. Louis, MO), which blocks the binding of Sp1 to target sequences, was added at different final concentrations (2.5, 5, and 10  $\mu\text{M}$ ) and incubated at 4°C for 1 h. For supershift assays, 1  $\mu\text{g}$  of monoclonal anti-Myc Tag antibody (Upstate Biotechnology, Lake Placid, NY) was added 30 min prior to addition of the probe. Binding reactions were carried out at 25°C for 30 min and electrophoresed on 6% acrylamide-0.5 $\times$  TBE gels, transferred to positively charged nylon membranes, and detected by a chemiluminescence method (Roche) and a LAS1000 scanner (Fujifilm Co., Tokyo, Japan).

### Silencing of Sp1, HCV, and *DHCR24* by siRNA

SP1 Validated Stealth RNAi<sup>TM</sup> siRNA (VHS40867, Invitrogen) was designed with the BLOCK-iT RNAi designer to target the human Sp1 mRNA sequence. RzM6-0d and RzM6-LC cells ( $1.5 \times 10^6$  cells in a 100-mm dish) were transfected with Sp1 siRNA (final concentration, 30 nM) using Lipofectamine RNAiMAX (Invitrogen) in Opti-MEM (Invitrogen) and incubated for 48 h at 37°C. The siRNAs specific for *DHCR24* and

HCV were designed and utilized as described previously [Nishimura et al., 2009].

### Kinase Inhibitors

ATM kinase inhibitor KU55933 (Wako Pure Chemical Industry, Osaka, Japan; final concentration, 10  $\mu\text{M}$ ), PI3K inhibitor LY294002 (Cell Signaling Technology, Beverly, MA; 50  $\mu\text{M}$ ), and MEK1 inhibitor PD98059 (Cell Signaling Technology; 50  $\mu\text{M}$ ) were added to cell cultures, which were incubated for 8 h at 37°C.

### Western Blotting

Western blotting was performed as described previously [Tsukiyama-Kohara et al., 2004] with the following primary antibodies: rabbit monoclonal anti-*DHCR24*/Seladin-1 (C59D8; Cell Signaling Technology); rabbit polyclonal anti-Sp1, anti-phospho-Akt (Ser473), and mouse monoclonal anti-phospho-ERK (Santa Cruz Biotechnology, Inc., Santa Cruz, CA); mouse monoclonal anti-HCV core (clone 31-2), E1 (clone 384), E2 (clone 544), NS4A (c14II-2-1), NS5A (32-2), NS5B (14-5), rabbit polyclonal anti-NS2, NS3 (R212), and NS4B (RR10) [Tsukiyama-Kohara et al., 2004]. Phosphorylation of Sp1 was investigated by 5% SDS-PAGE and immunoblotting with a polyclonal antibody against Sp1 phosphorylated at Ser101 (Active Motif, Carlsbad, CA) or Thr453 (Abcam, Cambridge, MA). Detection of  $\gamma$ H2AX was performed by 15% SDS-PAGE and immunoblotting with mouse monoclonal anti-phospho-histone H2AX (Ser139) (JBW301; Upstate Biotechnology). Phosphorylated ATM (Ser1981) and ATR (Ser428) were detected by specific antibodies (Cell Signaling Technology). Monoclonal anti-actin (Sigma), anti-histone H1 (Santa Cruz Biotechnology, Inc.), anti-HAUSP (Calbiochem, San Diego, CA), and anti-heat shock protein 90 (Stressgen, Victoria, BC, Canada) primary antibodies were used for normalization of Western blotting. Bound antibody was detected with a horseradish peroxidase-conjugated secondary antibody and visualization using ECL reagents (GE Healthcare, Piscataway, NJ) and an LAS1000 scanner (Fujifilm). Densitometric analysis of protein bands was performed with Image Quant software (Molecular Dynamics, Sunnyvale, CA).

### Quantitative PCR and HCV Infection

Total RNA was extracted from cell lines using ISOGEN, and reverse transcription of total RNA (125 ng) was performed with SuperScript III Reverse Transcriptase and Random Primers (Invitrogen). Synthesized cDNA samples were subjected to a TaqMan gene expression assay (Applied Biosystems, Foster City, CA), and the level of expression of *DHCR24* mRNA in each sample was normalized to the level of expression of *GAPDH* mRNA and represented as a ratio of the control (Hep-emp, CN5-Hep, or RzM6-0d). Infection of the human hepatocytes from human liver-uPA/SCID chimeric mice with HCV was performed,



and HCV RNA, *DHCR24* mRNA, and 18S rRNA were measured by quantitative PCR (qPCR), as described previously [Takano et al., 2011a].

### Statistical Analysis

The Student's *t*-test was used to analyze the statistical significance of the results; *P* values < 0.05 were considered statistically significant.

## RESULTS

### *DHCR24* Expression Is Upregulated by the Complete HCV Genome But Not by Individual Viral Proteins

Overexpression of *DHCR24* in human hepatocytes from human liver-uPA/SCID chimeric mice has been

observed after HCV infection (Fig. 1A). The overexpression of *DHCR24* in cells expressing HCV decreased to a similar extent as that observed in control cells following treatment with HCV siRNA (Fig. 1B). Since these findings suggest that overexpression of *DHCR24* is associated with the expression or infection by HCV, the identity of the viral factor involved in the augmentation of expression of *DHCR24* was examined. The level of expression of *DHCR24* mRNA was measured by quantitative RT-PCR (Fig. 1C) in HepG2-derived cell lines that stably express individual HCV proteins (core, E1, E2, NS2, NS3/4A, NS4B, NS5A, or NS5B; Supplementary Fig. 1). The level of expression of *DHCR24* mRNA was slightly higher in the cells expressing NS4B and NS5A than in control cells; however, there was no significant difference in the expression of *DHCR24* mRNA among these cell




Fig. 1. *DHCR24* expression is induced in the presence of the complete HCV genome. A: The amount of HCV RNA in chimeric mice infected with HCV-R6 (genotype 1b) was quantified by qPCR (left panel). The amount of *DHCR24* mRNA was measured, and the ratio to the amount of 18S rRNA was calculated in the tissues (right panel). B: Western blotting of *DHCR24*, HCV core, and actin protein in RzM6-0d and LC cells following treatment with the indicated siRNA. C: Level of *DHCR24* mRNA expression in cell lines with stable expression of individual HCV proteins, the HCV open reading

frame, or the complete HCV genome. Total RNA from HepG2/Lenti cell lines (emp, core, E1, E2, NS2, NS3/4A, NS4B, NS5A, and NS5B), CN5 cell lines (CN5-Hep, CN5-Swa, and CN5-Cre), or RzM6 cell lines (RzM6-0d and RzM6-LC) were prepared, and reverse transcription was performed. Synthesized cDNA was subjected to quantitative PCR. The level of expression of *DHCR24* mRNA for each sample was normalized to that of *GAPDH* mRNA and represented as a ratio of HepG2-emp (\**P* < 0.05).

lines. No significant upregulation of *DHCR24* mRNA was observed in the CN5-Cre cell line, which expresses all HCV proteins and is negative for viral replication [Tsukiyama-Kohara et al., 2004]. In contrast, significant upregulation of *DHCR24* was observed in a cell line that expresses the complete HCV genome (RzM6-LC) compared with the expression in HCV-negative control cells (RzM6-0d). Thus, expression of viral proteins alone is insufficient to reproduce the augmentation of expression of *DHCR24* induced by HCV.

### ***DHCR24* Promoter Activity Is Potentiated by the Expression of HCV**

The 5'-flanking region contains a number of possible transcriptional regulatory elements, including three candidate-binding motifs for the endoplasmic reticulum

(ER) stress-responsive transcription factor, XBP1. Cellular ER stress is induced in response to the expression of the HCV gene and infection by HCV [Tardif et al., 2005]. Thus, to explore host factors involved in the transcriptional regulation of *DHCR24*, the 5'-flanking genomic region (~5 kb) of *DHCR24* was isolated. Subsequently, *DHCR24* promoter reporter plasmids that contain the 5'-flanking region of *DHCR24* and the firefly luciferase gene were constructed (Fig. 2A). Relevant regions of the promoter were defined by constructing deletion mutants of the 5'-flanking regions, which were analyzed by a dual luciferase reporter assay in the presence or absence of full-length HCV genome expression—resulting from transfection with pCA-Rz [Tsukiyama-Kohara et al., 2004] or the control pCAGGS vector, respectively (Fig. 2B). Progressive shortening of the 5'-flanking regions did not




Fig. 2. *DHCR24* promoter activity is augmented by the expression of HCV. **A**: The 5'-flanking region of *DHCR24* was cloned from HepG2 and inserted upstream of the firefly luciferase gene in pGL3 (-4976/+113). A deletion series of the 5'-flanking regions was also constructed (-2982/+113, -2293/+113, and -515/+113). The black boxes indicate potential binding sites for the ER stress-responsive transcription factor, XBP-1. TSS, transcription start site (+1). **B**: HepG2 cells ( $1 \times 10^4$  cells/well in a 96-well plate) were co-transfected

with each *DHCR24* promoter reporter plasmid (0.25  $\mu$ g/well), a Renilla luciferase expression vector (pRL-TK; 0.025  $\mu$ g/well), and either an expression vector containing the HCV full-length genome (pCA-Rz; 0.5  $\mu$ g/well) or an empty expression vector (pCAGGS). Luciferase activity at 48 h post-transfection is shown as the ratio of firefly luciferase (FL) to Renilla luciferase (RL). Data are shown as the mean  $\pm$  SD from 2 representative experiments performed in triplicate (\* $P < 0.05$ ).

result in significant differences in the basal promoter activity (Fig. 2B). The  $-515/+113$  construct also produced a significant response in the presence of full-length HCV genome expression.

Additional reporter deletion mutants were constructed to define the region in the *DHCR24* promoter that is responsive to HCV expression. To this aim, potential binding motifs for transcription factors were predicted in the minimized *DHCR24* promoter sequence (nucleotides  $-515/+113$ ; Fig. 3A), and a series of promoter mutants containing sequential 100-bp

deletions was constructed. As shown in Fig. 3A, while the promoter activity of  $-515/+113$ ,  $-400/+113$ ,  $-300/+113$ , and  $-200/+113$  constructs was increased significantly by expression of HCV ( $*P < 0.05$ ), the promoter activity of the  $-100/+113$  construct was unchanged. Therefore, an HCV-responsive sequence appears to be located in the upstream region ( $-200$  to  $-100$  bp) from the transcriptional start site of *DHCR24*, which includes sequences with similarity to the consensus-binding motifs for AP-2 $\alpha$ , Sp1, MZF1, Pax-4, and NF-Y.




Fig. 3. Responsiveness of the *DHCR24* promoter to the expression of HCV. **A:** A *DHCR24* promoter series with sequential 100-bp deletions was constructed and analyzed as described in the legend to Fig. 2B ( $*p < 0.05$ ). **B:** An additional deletion series ( $-167/+113$ ,  $-144/+113$ , and  $-120/+113$ ) was constructed and analyzed as described in (A).

A more detailed deletion series (-167/+113, -144/+113, and -120/+113) was constructed (Fig. 3B) to determine the minimum-binding motif that responds to HCV expression. The responsiveness to the expression of HCV was lost with the removal of the proximal portion (-167 to -145), which includes candidate-binding motifs for AP-2 $\alpha$ , Sp1, and MZF-1. Thus, the identified HCV response element in the *DHCR24* promoter represents the minimum element of DNA sequence required for the promotion of the expression of *DHCR24* induced by HCV.

**HCV Expression Augments the Interaction Between the HCV Response Element and the Binding Molecule(s)**

Transcription of *DHCR24* is upregulated significantly in RzM6-LC cells that show persistent expression of

HCV [Nishimura et al., 2009]. Therefore, the effect of expression of HCV on the interaction between the HCV response element and its related transcription factor(s) was examined. Nuclear extracts were prepared from RzM6-LC cells, and an electrophoretic mobility shift assay (EMSA) using a DIG-labeled double-stranded oligonucleotide corresponding to the response element (-167/-140, 28 bp; Fig. 4A) was performed. The interaction between the response element and the nuclear factor was increased significantly in nuclear extracts from RzM6-LC cells compared with that in RzM6-0d cells (Fig. 4B). Thus, the binding affinity or quantity of the nuclear factor may be increased by the expression of HCV. The shifted band corresponding to the Sp1 consensus sequence also increased in RzM6-LC cells compared with that in control RzM6-0d cells, whereas no difference was




Fig. 4. The HCV response element mediates the overexpression of *DHCR24* induced by HCV. A: The HCV response element (-167/-140) in the 5'-flanking region of *DHCR24* includes sequences with similarity to the consensus-binding motifs for AP-2 $\alpha$ , Sp1, and MZF-1. B: Nuclear extracts were prepared from RzM6-0d and RzM6-LC cells and subjected to an electrophoresis mobility shift assay (EMSA; 10  $\mu$ g/sample) using the DIG-labeled HCV response element (28-bp), Sp1 (22-bp), or MZF-1 (21-bp) probes. Cold probe indicates unlabeled

oligonucleotides. The arrowhead indicates the interaction between the binding factor(s) and each oligonucleotide. C: Cured HuH-7/K4 cells and JFH/K4 cells were co-transfected with each *DHCR24* promoter reporter plasmid (0.5  $\mu$ g/well) and pRL-TK (0.05  $\mu$ g/well) and analyzed as described in Fig. 2B (\* $P$  < 0.05). (D) Nuclear extracts prepared from cured HuH-7/K4 cells or JFH/K4 cells were subjected to EMSA (25  $\mu$ g/sample) using the DIG-labeled HCV response element, Sp1, or MZF-1 probes.

noted in the intensity of the shifted band for the MZF-1 sequence between the RzM6-LC and RzM6-0d cells (Fig. 4B). In contrast to Sp1, the ability of endogenous MZF-1 to bind to its target sequence (affinity and/or amount) in the RzM6-LC cells was approximately equivalent to that observed in the RzM6-0d cells. Thus, MZF-1 is not likely to be involved in the increase, mediated by HCV expression, in the shifted band corresponding to the response element.

### HCV Infection Upregulates the Transcriptional Activity of the DHCR24 Promoter Through HCV Response Element

An in vitro model of HCV infection that replicates the entire HCV life cycle [Wakita et al., 2005] was used to confirm that transcription of *DHCR24* was mediated through the HCV response element. JFH/K4 cells, which show persistent infection with the HCV JFH-1 strain [Wakita et al., 2005], and control cells (cured HuH-7/K4) were transfected with the *DHCR24* promoter reporter plasmids, and promoter activity was measured. While *DHCR24* promoter reporters that included the HCV response element (−515/+113, −200/+113, and −167/+113) displayed significantly higher activity in JFH/K4 cells than in control cells, no difference was seen between the JFH/K4 cells and control cells transfected with the reporter lacking the HCV response element (−144/+113; Fig. 4C). These results suggest that the transcriptional activity of the *DHCR24* promoter was upregulated by HCV infection in a manner dependent on the response elements. Furthermore, augmentation of complex formation with the response element and the Sp1 probe was confirmed by EMSA using nuclear extracts from JFH/K4 and cured HuH-7/K4 cells (Fig. 4D).

### Sp1 Binds to the HCV Response Element

The HCV response element (−167/−140) includes candidate-binding motifs for Sp1, MZF-1, and AP-2 $\alpha$  (Fig. 4A). However, expression of AP-2 is repressed in the HepG2 cell line from which RzM6-LC cells are derived [Williams et al., 1988]. Thus, binding of AP-2 $\alpha$  to the response element was investigated by a supershift assay using anti-Myc and nuclear extract from HepG2 cells transfected with a Myc-tagged AP-2 $\alpha$  expression vector (Fig. 5A). The mobility of the DNA-AP-2 $\alpha$  complex was supershifted by the addition of anti-Myc (lane 6) but not control IgG (lane 5), whereas an additional shifted band corresponding to the response element was not observed after addition of anti-Myc (lane 3). Therefore, although exogenous AP-2 $\alpha$  protein expressed in HepG2 cells binds to the AP-2 $\alpha$  consensus sequence, it does not bind to the HCV response element.

The ability of Sp1 to form a DNA-protein complex with the HCV response element was investigated by performing EMSAs in the presence of mithramycin A (MMA)—a GC-specific DNA-binding antibiotic that binds to the GC-box in the promoter to block binding

of Sp1 or other Sp family proteins [Blume et al., 1991]. As shown in Fig. 5B, MMA (2.5, 5.0, and 10  $\mu$ M) inhibited complex formation in a dose-dependent manner. In contrast, the formation of DNA-protein complexes with the MZF-1 probe was not affected by the addition of MMA, suggesting that the inhibition mediated by MMA was specific for the GC box-Sp1, and that complex formation with the response element requires the Sp1 binding site. A supershift assay using nuclear extract from HepG2 cells transfected with a Myc-tagged Sp1 expression vector and anti-Myc was also performed (Supplementary Fig. 2). The mobility of the HCV response element and the Sp1 consensus sequence was supershifted partially by addition of anti-Myc (lanes 3 and 6). The effect of silencing the expression of Sp1 with small interfering RNA (siRNA) was analyzed by EMSA using nuclear extracts from Sp1-knockdown RzM6-0d and RzM6-LC cells (Fig. 5C). DNA-protein complexes with the response element or the Sp1 probe were not observed (lanes 2, 4, 6, and 8); however, formation of DNA-MZF-1 complexes was not influenced by siRNA treatment (lanes 9–12). Immunoblotting was used to confirm efficient silencing of the Sp1 protein in cells used to generate the nuclear extracts (Fig. 5D). A significant decrease in the expression of *DHCR24* was observed in the cytosolic fraction from RzM6 cells transfected with siRNA specific for Sp1 (Fig. 5D). Thus, these results indicate that Sp1, but neither AP-2 $\alpha$  nor MZF-1, bound to the HCV response element, and that Sp1 may play an important role in the transcriptional regulation of *DHCR24*.

### Transcriptional Regulation of DHCR24 Through the HCV Response Element Is Mediated by Oxidative Stress

*DHCR24* functions as a mediator of the cellular response to oxidative stress [Greeve et al., 2000; Benvenuti et al., 2005; Di Stasi et al., 2005; Luciani et al., 2005; Lu et al., 2008] and is a hydrogen peroxide scavenger [Lu et al., 2008]. Expression of the *DHCR24* gene is also induced in response to oxidative stress [Wu et al., 2004; Benvenuti et al., 2006; Kuehnle et al., 2008]. Expression of the HCV gene elevates the level of reactive oxygen species (ROS) via dysregulation of ER-mediated calcium homeostasis, which results in oxidative stress [Tardif et al., 2005]. Therefore, the role of oxidative stress induced by HCV in the regulatory mechanism of the expression of *DHCR24* was examined. HepG2 cells were treated with hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and transfected with reporter plasmids containing the *DHCR24* promoter deletion mutants. Measurement of promoter activity revealed a significant increase in transcription in response to oxidative stress (H<sub>2</sub>O<sub>2</sub>) for *DHCR24* promoters containing the HCV response element (−4976/+113, −2982/+113, −515/+113, and −167/+113) but not for the promoter lacking the response element (−144/+113; Fig. 6A). Therefore, enhanced transcription in response to oxidative stress by reporter constructs




Fig. 5. Sp1 binds to the HCV response element. **A:** Nuclear extract was prepared from HepG2 cells transfected with pcDNA6-AP-2α-myc and subjected to EMSA (10 μg/sample) using the DIG-labeled HCV response element or AP-2α probes (26-bp). For a supershift analysis of myc-tagged AP-2α, anti-Myc, or control IgG was added to the binding reaction. The closed arrowhead indicates the interaction between the binding factor(s) and each oligonucleotide, and an additional interaction with antibody is indicated by an open arrowhead. **B:** Nuclear extract from HepG2 cells was pre-incubated at 4°C for 1 h

with different concentrations (2.5, 5, and 10 μM) of mithramycin A (MMA) and subjected to EMSA (10 μg/sample) using the DIG-labeled HCV response element, Sp1, or MZF-1 probes. **C:** Nuclear extracts were prepared from RzM6 cells transfected with Sp1 siRNA or control siRNA and subjected to EMSA (10 μg/sample) using the DIG-labeled HCV response element, Sp1, or MZF-1 probes. **D:** Expression of Sp1, DHCR24, and other proteins was detected in both the nuclear fraction (N), used for the EMSA shown in Fig. 4C, and in the cytosolic-membrane fraction (C).

containing the *DHCR24* promoter may be mediated through the HCV response element. The formation of complexes containing the response element or Sp1 probe was increased markedly in the nuclear extracts from the H<sub>2</sub>O<sub>2</sub>-treated HepG2 cells (Fig. 6B) or other hepatic cell lines (Supplementary Fig. 3), suggesting that oxidative stress enhances the binding affinity of Sp1 to the HCV response element.

### Overexpression of *DHCR24* in M6-LC Cells Is Blocked by an ROS Scavenger

The increase in the expression of *DHCR24* induced by oxidative stress can be blocked by treatment with an ROS scavenger, *N*-acetylcysteine (NAC) [Wu et al., 2004], which is a precursor of the potent biological antioxidant glutathione. The H<sub>2</sub>O<sub>2</sub>-induced overexpression




Fig. 6. Oxidative stress increases the transcription of *DHCR24* through the HCV response element and Sp1. A: HepG2 cells ( $1 \times 10^4$  cells/well in a 96-well plate) were co-transfected with individual *DHCR24* promoter reporter plasmids (0.5  $\mu$ g/well) and pRL-TK (0.05  $\mu$ g/well). Forty-four hours post-transfection, cells were treated with or without 1 mM H<sub>2</sub>O<sub>2</sub> for 4 h and analyzed as described in Fig. 2B ( $*P < 0.05$ ). B: Nuclear extracts prepared from H<sub>2</sub>O<sub>2</sub>-treated (1 mM, 4 h) or untreated HepG2 cells were subjected to EMSA (10  $\mu$ g/sample) using the DIG-labeled HCV response element, Sp1, or MZF-1 probes. Densitometric analysis of shifted bands was performed using the Image Quant software. Data are shown as the mean  $\pm$  SD from triplicate quantifications of two representative experiments ( $*P < 0.05$ ).

of *DHCR24* was inhibited by pre-treatment with NAC and blocked partially by NAC treatment after the induction of oxidative stress ( $\sim 50\%$  suppression; Fig. 7A). The enhanced expression of *DHCR24* in RzM6-LC cells decreased after 12 or 24 h of treatment with NAC without influencing the level of expression of HCV, suggesting that overexpression of *DHCR24* in cells expressing HCV is mediated through oxidative stress.

#### Overexpression and Enhanced Phosphorylation of Sp1 in the Cells Expressing HCV

Sp1 is a transcription factor that is activated in response to a variety of cellular stressors, including oxidative stress [Schafer et al., 2003; Chu and Ferro, 2006; Dasari et al., 2006; Qin et al., 2009; Lin et al., 2011]. Thus, Sp1 may play an important role in linking oxidative stress and augmentation of *DHCR24* transcription in cells expressing HCV. Sp1 was overexpressed significantly in RzM6-LC cells treated with H<sub>2</sub>O<sub>2</sub> compared with the control cells (Fig. 8A). Phosphorylation of Sp1 at Ser101 was also elevated

under oxidative stress. Both the basal level and phosphorylation status of nuclear Sp1 were higher in the presence of HCV (RzM6-LC cells) than in the absence of HCV (RzM6-0d cells; Fig. 8B).

Phosphorylation of Sp1 at Ser101 is a target of the DNA damage signaling pathway mediated by ATM (ataxia telangiectasia mutated) and ATR (ATM and Rad3-related) kinases [Olofsson et al., 2007; Iwahori et al., 2008]. As shown in Fig. 8C, phosphorylation of Sp1 at Ser101 was no longer detectable following pre-treatment with an ATM kinase inhibitor (KU55933) before exposure to H<sub>2</sub>O<sub>2</sub>. In contrast, phosphorylation was not affected by other kinase inhibitors (phosphatidylinositol-3 kinase inhibitor, LY294002 or MEK1 inhibitor, PD98059). Similarly, phosphorylation of Sp1 at Thr453, which is important for transcriptional activation of Sp1 [Milanini-Mongiati et al., 2002; D'Addario et al., 2006; Hsu et al., 2006; Lin et al., 2011], was not seen in response to oxidative stress following treatment with KU55933 (Fig. 8C). The induction of expression of *DHCR24* after H<sub>2</sub>O<sub>2</sub> exposure was suppressed significantly by treatment with KU55933 or NAC, which corresponds with inhibition of Sp1 phosphorylation. In the presence of MMA, the phosphorylation of Sp1 was not inhibited. However, since MMA blocks the binding of Sp1 [Blume et al., 1991], the induction of expression of *DHCR24* by H<sub>2</sub>O<sub>2</sub> was inhibited. Impairment of *DHCR24* induction by H<sub>2</sub>O<sub>2</sub> was also observed after treatment with siRNAs targeting ATM (Supplementary Fig. 4).

Studies on the relationship between HCV and ATM have reported that the interaction of NS3/4A with ATM results in delayed de-phosphorylation of both phosphorylated ATM and phosphorylated histone H2AX at Ser139 ( $\gamma$ H2AX), which acts as a substrate for ATM in response to DNA damage [Lai et al., 2008]. In the present study, delayed de-phosphorylation of  $\gamma$ H2AX was also observed in HCV replicon cells (Supplementary Fig. 5), which corresponded with increased phosphorylation of the H2AX Ser139 residue in cells expressing HCV (Fig. 8). Similarly, phosphorylation of ATM was sustained in HCV replicon cells (Supplementary Fig. 6). Therefore, DNA repair may be impaired in cells expressing or replicating HCV, resulting in sustained DNA damage. As a result, downstream substrates such as Sp1 Ser101 and Thr453 residues or the H2AX Ser139 residue may be phosphorylated to a greater extent in cells expressing HCV compared with control cells in the basal state or cells under oxidative stress (Fig. 8A and B).

Taken together, these results indicate that the oxidative stress induced by HCV may produce quantitative as well as qualitative activation of Sp1, thereby resulting in augmentation of *DHCR24* transcription.

#### DISCUSSION

HCV establishes chronic infection and induces persistent overexpression of *DHCR24* in human hepatocytes [Nishimura et al., 2009]. HCV also confers






Fig. 7. Overexpression of *DHCR24* in the cells expressing HCV is blocked by treatment with an oxidative stress scavenger. **A:** HepG2 cells were treated without (lanes 1, 4, and 6) or with (lanes 2 and 5) NAC (10 mM, 8 h). Cells treated with H<sub>2</sub>O<sub>2</sub> (1 mM, 4 h) were also treated with 10 mM NAC for 8 h either before (pre; lanes 3 and 4) or after (post; lanes 5 and 6) H<sub>2</sub>O<sub>2</sub> exposure. Whole-cell lysates (40  $\mu$ g/lane) were analyzed by 10% SDS-PAGE and immunoblotting using a *DHCR24*/Seladin-1 mAb. Immunoblotting with an actin mAb served as the internal loading control. The ratio of *DHCR24*/actin was normalized to that of untreated cells (lane 1). **B:** RzM6-LC cells were treated with NAC (10 mM) for 12 h (lane 2) or 24 h (lane 4). Whole-cell lysates were analyzed as described in (A). The ratio of HCV core to actin protein was also calculated. Experiments were performed three times, and representative results are shown.

resistance to the apoptosis induced by oxidative stress and suppresses p53 activity by blocking nuclear p53 acetylation and increasing the interaction between p53 and HDM2 (p53-specific E3 ligase) in the cytoplasm, which may be mediated by inhibition of p53 degradation. Thus, the augmentation of *DHCR24* by HCV reflects the tumorigenicity of hepatocytes. The present study identified the genomic region of *DHCR24* that is responsive to HCV, and showed that this response is mediated through the activation of Sp1 induced by oxidative stress. In general, expression of the HCV gene elevates the levels of ROS through dysregulation of ER-mediated calcium homeostasis [Tardif et al., 2005]. In healthy cells, ROS usually exist in equilibrium with antioxidants that scavenge ROS and prevent cellular injury. However, this critical balance may be disrupted in the cells infected with HCV, resulting in the accumulation of

ROS and the development of constitutive oxidative stress.

Sp1 is a member of the Sp/KLF family of transcription factors that bind to GC elements of promoters [Black et al., 2001; Kaczynski et al., 2003; Chu and Ferro, 2005; Li and Davie, 2010]. Under a variety of endogenous and exogenous stimuli—including oxidative stress and DNA damage—activation of Sp1 may be mediated via induction of expression of Sp1 and post-translational modifications such as acetylation, sumoylation, O-linked glycosylation, and phosphorylation. Sp1 is phosphorylated by several kinases, including DNA-dependent protein kinase, casein kinase II, and cyclin A/cdk2, which exert both positive and negative effects on transcription [Jackson et al., 1990; Armstrong et al., 1997; Fojas de Borja et al., 2001; Ryu et al., 2003]. Sp1 is the only Sp/KLF family member to contain putative consensus SQ/TQ cluster domains within the transactivation domains, which suggests that Sp1 is a substrate of the PI3K-related kinases, for example, ATM, DNA-dependent protein kinase, and ATR. Indeed, Sp1 is a target of the ATM-dependent DNA damage response pathway [Iwahori et al., 2007, 2008; Olofsson et al., 2007]. ATM plays a central role in orchestrating molecular events involved in double-strand break signaling, which is mediated via the phosphorylation of a variety of substrate proteins—including p53 and BRCA1 transcription factors—involved in the DNA damage response. As a result, these phosphorylation events lead to cell cycle checkpoint activation, DNA repair, altered gene expression patterns, and/or apoptosis [Shiloh, 2006].

Given the role of Sp1 in oxidative stress [Schafer et al., 2003; Chu and Ferro, 2006; Dasari et al., 2006; Rojo et al., 2006; Qin et al., 2009; Lin et al., 2010], Sp1 may be regulated by the oxidative stress induced by HCV and the subsequent phosphorylation, which depends on ATM. However, little is known regarding the regulation of Sp1 in response to DNA damage. Although the precise role of phosphorylation of Sp1 at Ser101 in the DNA damage response is unclear, the similar kinetics of Sp1 and  $\gamma$ H2AX phosphorylation [Olofsson et al., 2007] suggest that Sp1 is an early target of the DNA damage response pathway. Thus, Sp1 may be involved in modulating the cellular response to DNA damage to prevent cell death [Ryu et al., 2003]. Phosphorylation of Sp1 at Ser101 and histone H2AX, which occurs in parallel in response to oxidative stress, was enhanced in cells expressing HCV compared with that observed in control cells (Fig. 8A). Interestingly, augmentation of Sp1 phosphorylation in parallel with histone H2AX phosphorylation was also detected for cells expressing HCV in the basal state (Fig. 8A and B), which may be primarily due to the increase in endogenous Sp1 protein (Fig. 8A and B). In support of these results, enhanced phosphorylation of Ser101 on Sp1 occurs upon HSV-1 infection, and is mediated by ATM [Iwahori et al., 2007]. Thus, increased phosphorylation of Sp1 and  $\gamma$ H2AX in cells expressing HCV is likely to reflect the higher activity




Fig. 8. Overexpression and elevated phosphorylation of Sp1 in the cells expressing HCV. **A**: RzM6-0d and RzM6-LC cells were treated with or without H<sub>2</sub>O<sub>2</sub> (1 mM, 4 h). Whole-cell lysates (15 μg/lane) were analyzed by 15% SDS-PAGE and immunoblotting using phospho-H2AX (Ser139) (γH2AX) and HCV core mAbs. An actin mAb served as an internal loading control. Whole-cell lysates (25 μg/lane) were analyzed by 5% SDS-PAGE and immunoblotting using anti-Sp1 (phosphorylated Sp1 and native Sp1, as indicated) and anti-phospho-Sp1 (Ser101) was performed. **B**: RzM6-0d and RzM6-LC cells were fractionated to produce nuclear (N) and cytosolic-membrane fractions (C). Fractionated samples (15 μg/lane) were analyzed as described in (A). The ratio of phosphorylated Sp1 to Sp1 protein is indicated. Immunoblotting using anti-HAUSP served as a

high-molecular-weight loading control. **C**: RzM6-0d cells were pre-treated for 8 h with NAC (10 mM), MMA (10 μM), KU55933 (KU; 10 μM), LY294002 (LY; 50 μM), or PD98059 (PD; 50 μM) and incubated for 4 h in the absence or presence of H<sub>2</sub>O<sub>2</sub> (1 mM). Whole-cell lysates (40 μg/lane) were analyzed by 5% SDS-PAGE and immunoblotting using anti-phospho-Sp1 (Ser101), (Thr453), and polyclonal anti-Sp1 (white arrowhead, phosphorylated Sp1; black arrowhead, native Sp1). Detection of HAUSP was performed to confirm the quantity of loaded protein in each lane. Whole-cell lysates (25 μg/lane) were analyzed simultaneously by 10% SDS-PAGE and immunoblotting using anti-DHCR24/seladin-1 mAb, anti-phospho-Akt (Ser473), and anti-phospho-ERK antibodies.

of ATM, which may result from the accumulation and frequency of DNA damage caused by increased generation of endogenous ROS.

Oxidative stress is a common mechanism of liver injury [Loguercio and Federico, 2003] and is mediated by the direct effects of ROS on signal transduction pathways, including extracellular signal-regulated kinase 1/2 (ERK1/2), c-Jun N-terminal kinase (JNK), and p38 mitogen-activated protein kinases (MAPKs), which act as downstream kinases in the MAPK cascade to phosphorylate Sp1 Thr453/739 residues [Milanini-Mongiati et al., 2002; D'Addario et al., 2006; Hsu et al., 2006; Chuang et al., 2008; Lin et al., 2011]. These signal transduction pathways are also stimulated by oxidative stress in the hepatic cells expressing or

replicating HCV, [Qadri et al., 2004; Burdette et al., 2010; Lin et al., 2010]. Therefore, oxidative stress in response to HCV may induce downstream signaling pathways, such as ERK1/2, JNK, and p38 MAPK as well as ATM/ATR, to activate Sp1 via post-translational modifications.

Sp1 is a host factor activated by several viral proteins, including HIV-1 Vpr, and HTLV-1 Tax [Peng et al., 2003; Amini et al., 2004; Chang et al., 2005; Zhang et al., 2009]. The HCV core and NS5A proteins also activate Sp1 [Lee et al., 2001; Xiang et al., 2010]. The HCV core upregulates the DNA-binding activity and phosphorylation of Sp1 [Lee et al., 2001], and NS5A may also exert a similar effect on Sp1 activity. However, a physical interaction between these

proteins and Sp1 has not yet been demonstrated. Both HCV core and NS5A proteins have a high potential for oxidative stress induction [García-Mediavilla et al., 2005; Dionisio et al., 2009], which may mediate activation of Sp1. On the other hand, individual viral proteins were insufficient to increase the expression of *DHCR24* (Fig. 1A). Therefore, in addition to induction of oxidative stress by each viral protein, the persistence of the signaling pathways induced by oxidative stress, for example, ATM (Supplementary Fig. 6), may also be required for the Sp1-mediated increase in the expression of *DHCR24*.

The results of the present study revealed that knockdown of expression of Sp1 almost completely blocked the enhanced expression of *DHCR24*. Sp1 is expressed ubiquitously in various mammalian cells and is involved in regulating the transcriptional activity of genes implicated in many cellular processes [Black et al., 2001; Kaczynski et al., 2003; Chu and Ferro, 2005; Li and Davie, 2010]. Thus, Sp1 may represent an essential master regulator among the myriad of transcription factors involved in the direct regulation of *DHCR24* transcription.

In conclusion, HCV was shown to enhance the expression of *DHCR24* via the activation of Sp1, which may shed light on the mechanism of tumorigenesis associated with HCV.


#### ACKNOWLEDGMENTS

The authors would like to thank Ms. Ryoko Takehara and Yuri Kasama for their technical support, Yuko Tokunaga for her comments, and Dr. Chieko Kai for her support.

#### REFERENCES

- Amini S, Saunders M, Kelley K, Khalili K, Sawaya BE. 2004. Interplay between HIV-1 Vpr and Sp1 modulates p21(WAF1) gene expression in human astrocytes. *J Biol Chem* 279:46046–46056.
- Armstrong SA, Barry DA, Leggett RW, Mueller CR. 1997. Casein kinase II-mediated phosphorylation of the C terminus of Sp1 decreases its DNA binding activity. *J Biol Chem* 272:13489–13495.
- Benvenuti S, Luciani P, Vannelli GB, Gelmini S, Franceschi E, Serio M, Peri A. 2005. Estrogen and selective estrogen receptor modulators exert neuroprotective effects and stimulate the expression of selective Alzheimer's disease indicator-1, a recently discovered antiapoptotic gene, in human neuroblast long-term cell cultures. *J Clin Endocrinol Metab* 90:1775–1782.
- Benvenuti S, Saccardi R, Luciani P, Urbani S, Deledda C, Cellai I, Francini F, Squecco R, Rosati F, Danza G, Gelmini S, Greeve I, Rossi M, Maggi R, Serio M, Peri A. 2006. Neuronal differentiation of human mesenchymal stem cells: Changes in the expression of the Alzheimer's disease-related gene seladin-1. *Exp Cell Res* 312:2592–2604.
- Black AR, Black JD, Azizkhan-Clifford J. 2001. Sp1 and kruppel-like factor family of transcription factors in cell growth regulation and cancer. *J Cell Physiol* 188:143–160.
- Blume SW, Snyder RC, Ray R, Thomas S, Koller CA, Miller DM. 1991. Mithramycin inhibits SP1 binding and selectively inhibits transcriptional activity of the dihydrofolate reductase gene in vitro and in vivo. *J Clin Invest* 88:1613–1621.
- Bruchfeld A, Stahle L, Andersson J, Schvarcz R. 2001. Ribavirin treatment in dialysis patients with chronic hepatitis C virus infection—A pilot study. *J Viral Hepat* 8:287–292.
- Burdette D, Olivarez M, Waris G. 2010. Activation of transcription factor Nrf2 by hepatitis C virus induces the cell-survival pathway. *J Gen Virol* 91:681–690.
- Chang LK, Chung JY, Hong YR, Ichimura T, Nakao M, Liu ST. 2005. Activation of Sp1-mediated transcription by Rta of Epstein-Barr virus via an interaction with MCAF1. *Nucleic Acids Res* 33:6528–6539.
- Chu S, Ferro TJ. 2005. Sp1: Regulation of gene expression by phosphorylation. *Gene* 348:1–11.
- Chu S, Ferro TJ. 2006. Identification of a hydrogen peroxide-induced PP1-JNK1-Sp1 signaling pathway for gene regulation. *Am J Physiol Lung Cell Mol Physiol* 291:L983–992.
- Chuang JY, Wang YT, Yeh SH, Liu YW, Chang WC, Hung JJ. 2008. Phosphorylation by c-Jun NH<sub>2</sub>-terminal kinase 1 regulates the stability of transcription factor Sp1 during mitosis. *Mol Biol Cell* 19:1139–1151.
- Crameri A, Biondi E, Kuehne K, Lutjohann D, Thelen KM, Perga S, Dotti CG, Nitsch RM, Ledesma MD, Mohajeri MH. 2006. The role of seladin-1/DHCR24 in cholesterol biosynthesis, APP processing and Abeta generation in vivo. *EMBO J* 25:432–443.
- D'Addario M, Arora PD, McCulloch CA. 2006. Role of p38 in stress activation of Sp1. *Gene* 379:51–61.
- Dasari A, Bartholomew JN, Volonte D, Galbiati F. 2006. Oxidative stress induces premature senescence by stimulating caveolin-1 gene transcription through p38 mitogen-activated protein kinase/Sp1-mediated activation of two GC-rich promoter elements. *Cancer Res* 66:10805–10814.
- Di Stasi D, Vallacchi V, Campi V, Ranzani T, Daniotti M, Chiodini E, Fiorentini S, Greeve I, Prinetti A, Rivoltini L, Pierotti MA, Rodolfo M. 2005. *DHCR24* gene expression is upregulated in melanoma metastases and associated to resistance to oxidative stress-induced apoptosis. *Int J Cancer* 115:224–230.
- Dignam JD, Lebovitz RM, Roeder RG. 1983. Accurate transcription initiation by RNA polymerase II in a soluble extract from isolated mammalian nuclei. *Nucleic Acids Res* 11:1475–1489.
- Dionisio N, Garcia-Mediavilla MV, Sanchez-Campos S, Majano PL, Benedicto I, Rosado JA, Salido GM, Gonzalez-Gallego J. 2009. Hepatitis C virus NS5A and core proteins induce oxidative stress-mediated calcium signalling alterations in hepatocytes. *J Hepatol* 50:872–882.
- Fojas de Borja P, Collins NK, Du P, Azizkhan-Clifford J, Mudryj M. 2001. Cyclin A-CDK phosphorylates Sp1 and enhances Sp1-mediated transcription. *EMBO J* 20:5737–5747.
- García-Mediavilla MV, Sánchez-Campos S, González-Pérez P, Gómez-Gonzalo M, Majano PL, Lopez-Cabrera M, Clemente G, García-Monzon C, González-Gallego J. 2005. Differential contribution of hepatitis C virus NS5A and core proteins to the induction of oxidative and nitrosative stress in human hepatocyte-derived cells. *J Hepatol* 43:606–613.
- Greeve I, Hermans-Borgmeyer I, Brellinger C, Kasper D, Gomez-Isla T, Behl C, Levkau B, Nitsch RM. 2000. The human DIMINUTO/DWARF1 homolog seladin-1 confers resistance to Alzheimer's disease-associated neurodegeneration and oxidative stress. *J Neurosci* 20:7345–7352.
- Hsu MC, Chang HC, Hung WC. 2006. HER-2/neu represses the metastasis suppressor RECK via ERK and Sp transcription factors to promote cell invasion. *J Biol Chem* 281:4718–4725.
- Iwahori S, Shirata N, Kawaguchi Y, Weller SK, Sato Y, Kudoh A, Nakayama S, Isomura H, Tsurumi T. 2007. Enhanced phosphorylation of transcription factor sp1 in response to herpes simplex virus type 1 infection is dependent on the ataxia telangiectasia-mutated protein. *J Virol* 81:9653–9664.
- Iwahori S, Yasui Y, Kudoh A, Sato Y, Nakayama S, Murata T, Isomura H, Tsurumi T. 2008. Identification of phosphorylation sites on transcription factor Sp1 in response to DNA damage and its accumulation at damaged sites. *Cell Signal* 20:1795–1803.
- Jackson SP, MacDonald JJ, Lees-Miller S, Tjian R. 1990. GC box binding induces phosphorylation of Sp1 by a DNA-dependent protein kinase. *Cell* 63:155–165.
- Kaczynski J, Cook T, Urrutia R. 2003. Sp1- and Kruppel-like transcription factors. *Genome Biol* 4:206.
- Kohara M, Tanaka T, Tsukiyama-Kohara K, Tanaka S, Mizokami M, Lau JY, Hattori N. 1995. Hepatitis C virus genotypes 1 and 2 respond to interferon-alpha with different virologic kinetics. *J Infect Dis* 172:934–938.

- Koike K. 2007. Hepatitis C virus contributes to hepatocarcinogenesis by modulating metabolic and intracellular signaling pathways. *J Gastroenterol Hepatol* 22:S108–S111.
- Kuehnle K, Cramer A, Kalin RE, Luciani P, Benvenuti S, Peri A, Ratti F, Rodolfo M, Kulic L, Heppner FL, Nitsch RM, Mohajeri MH. 2008. Prosurvival effect of DHCR24/Seladin-1 in acute and chronic responses to oxidative stress. *Mol Cell Biol* 28:539–550.
- Lai CK, Jeng KS, Machida K, Cheng YS, Lai MM. 2008. Hepatitis C virus NS3/4A protein interacts with ATM, impairs DNA repair and enhances sensitivity to ionizing radiation. *Virology* 370:295–309.
- Lavanchy D. 2009. The global burden of hepatitis C. *Liver Int* 29:74–81.
- Lee S, Park U, Lee YI. 2001. Hepatitis C virus core protein transactivates insulin-like growth factor II gene transcription through acting concurrently on Egr1 and Sp1 sites. *Virology* 283:167–177.
- Li L, Davie JR. 2010. The role of Sp1 and Sp3 in normal and cancer cell biology. *Ann Anat* 192:275–283.
- Lin W, Tsai WL, Shao RX, Wu G, Peng LF, Barlow LL, Chung WJ, Zhang L, Zhao H, Jang JY, Chung RT. 2010. Hepatitis C virus regulates transforming growth factor beta1 production through the generation of reactive oxygen species in a nuclear factor kappaB-dependent manner. *Gastroenterology* 138:2509–2518.
- Lin HH, Lai SC, Chau LY. 2011. Heme oxygenase-1/carbon monoxide induces vascular endothelial growth factor expression via p38 kinase-dependent activation of Sp1. *J Biol Chem* 286:3829–3838.
- Loguercio C, Federico A. 2003. Oxidative stress in viral and alcoholic hepatitis. *Free Radic Biol Med* 34:1–10.
- Lu X, Kambe F, Cao X, Kozaki Y, Kaji T, Ishii T, Seo H. 2008. 3Beta-hydroxysteroid-delta24 reductase is a hydrogen peroxide scavenger, protecting cells from oxidative stress-induced apoptosis. *Endocrinology* 149:3267–3273.
- Luciani P, Gelmini S, Ferrante E, Lania A, Benvenuti S, Baglioni S, Mantovani G, Cellai I, Ammannati F, Spada A, Serio M, Peri A. 2005. Expression of the antiapoptotic gene seladin-1 and octreotide-induced apoptosis in growth hormone-secreting and non-functioning pituitary adenomas. *J Clin Endocrinol Metab* 90:6156–6161.
- Milanini-Mongiati J, Pouyssegur J, Pages G. 2002. Identification of two Sp1 phosphorylation sites for p42/p44 mitogen-activated protein kinases: Their implication in vascular endothelial growth factor gene transcription. *J Biol Chem* 277:20631–20639.
- Nakamura H, Ogawa H, Kuroda T, Yamamoto M, Enomoto H, Kishima Y, Yoshida K, Ito H, Matsuda M, Noguchi S. 2002. Interferon treatment for patients with chronic hepatitis C infected with high viral load of genotype 2 virus. *Hepatology* 49:1373–1376.
- Nishimura T, Kohara M, Izumi K, Kasama Y, Hirata Y, Huang Y, Shuda M, Mukaidani C, Takano T, Tokunaga Y, Nuriya H, Satoh M, Saito M, Kai C, Tsukiyama-Kohara K. 2009. Hepatitis C virus impairs p53 via persistent overexpression of 3beta-hydroxysteroid Delta24-reductase. *J Biol Chem* 284:36442–36452.
- Olofsson BA, Kelly CM, Kim J, Hornsby SM, Azizkhan-Clifford J. 2007. Phosphorylation of Sp1 in response to DNA damage by ataxia telangiectasia-mutated kinase. *Mol Cancer Res* 5:1319–1330.
- Peng H, He H, Hay J, Ruyechan WT. 2003. Interaction between the varicella zoster virus IE62 major transactivator and cellular transcription factor Sp1. *J Biol Chem* 278:38068–38075.
- Qadri I, Iwahashi M, Capasso JM, Hopken MW, Flores S, Schaack J, Simon FR. 2004. Induced oxidative stress and activated expression of manganese superoxide dismutase during hepatitis C virus replication: Role of JNK, p38 MAPK and AP-1. *Biochem J* 378:919–928.
- Qin K, Zhao L, Ash RD, McDonough WF, Zhao RY. 2009. ATM-mediated transcriptional elevation of prion in response to copper-induced oxidative stress. *J Biol Chem* 284:4582–4593.
- Rojo AI, Salina M, Salazar M, Takahashi S, Suske G, Calvo V, de Sagarra MR, Cuadrado A. 2006. Regulation of heme oxygenase-1 gene expression through the phosphatidylinositol 3-kinase/PKC-zeta pathway and Sp1. *Free Radic Biol Med* 41:247–261.
- Ryu H, Lee J, Zaman K, Kubilis J, Ferrante RJ, Ross BD, Neve R, Ratan RR. 2003. Sp1 and Sp3 are oxidative stress-inducible, anti-death transcription factors in cortical neurons. *J Neurosci* 23:3597–3606.
- Schafer G, Cramer T, Suske G, Kemmner W, Wiedenmann B, Hocker M. 2003. Oxidative stress regulates vascular endothelial growth factor-A gene transcription through Sp1- and Sp3-dependent activation of two proximal GC-rich promoter elements. *J Biol Chem* 278:8190–8198.
- Shiloh Y. 2006. The ATM-mediated DNA-damage response: Taking shape. *Trends Biochem Sci* 31:402–410.
- Takano T, Kohara M, Kasama Y, Nishimura T, Saito M, Kai C, Tsukiyama-Kohara K. 2011a. Translocase of outer mitochondrial membrane 70 expression is induced by hepatitis C virus and is related to the apoptotic response. *J Med Virol* 83:801–809.
- Takano T, Tsukiyama-Kohara K, Hayashi M, Hirata Y, Satoh M, Tokunaga Y, Tateno C, Hayashi Y, Hishima T, Funata N, Sudo M, Kohara M. 2011b. Augmentation of DHCR24 expression by hepatitis C virus infection facilitates viral replication in hepatocytes. *J Hepatol* 55:512–521.
- Tardif KD, Waris G, Siddiqui A. 2005. Hepatitis C virus, ER stress, and oxidative stress. *Trends Microbiol* 13:159–163.
- Tsukiyama-Kohara K, Tone S, Maruyama I, Inoue K, Katsume A, Nuriya H, Ohmori H, Ohkawa J, Taira K, Hoshikawa Y, Shibasaki F, Reth M, Minatogawa Y, Kohara M. 2004. Activation of the CKI-CDK-Rb-E2F pathway in full genome hepatitis C virus-expressing cells. *J Biol Chem* 279:14531–14541.
- Wakita T, Pietschmann T, Kato T, Date T, Miyamoto M, Zhao Z, Murthy K, Habermann A, Krausslich HG, Mizokami M, Bartenschlager R, Liang TJ. 2005. Production of infectious hepatitis C virus in tissue culture from a cloned viral genome. *Nat Med* 11:791–796.
- Waterham HR, Koster J, Romeijn GJ, Hennekam RC, Vreken P, Andersson HC, FitzPatrick DR, Kelley RI, Wanders RJ. 2001. Mutations in the 3beta-hydroxysteroid Delta24-reductase gene cause desmosterolosis, an autosomal recessive disorder of cholesterol biosynthesis. *Am J Hum Genet* 69:685–694.
- Williams T, Admon A, Luscher B, Tjian R. 1988. Cloning and expression of AP-2, a cell-type-specific transcription factor that activates inducible enhancer elements. *Genes Dev* 2:1557–1569.
- Wu C, Miloslavskaya I, Demontis S, Maestro R, Galaktionov K. 2004. Regulation of cellular response to oncogenic and oxidative stress by Seladin-1. *Nature* 432:640–645.
- Xiang Z, Qiao L, Zhou Y, Babiuk LA, Liu Q. 2010. Hepatitis C virus nonstructural protein-5A activates sterol regulatory element-binding protein-1c through transcription factor Sp1. *Biochem Biophys Res Commun* 402:549–553.
- Zhang L, Zhi H, Liu M, Kuo YL, Giam CZ. 2009. Induction of p21(CIP1/WAF1) expression by human T-lymphotropic virus type 1 Tax requires transcriptional activation and mRNA stabilization. *Retrovirology* 6:35.



## Short communication

## Translocase of outer mitochondrial membrane 70 induces interferon response and is impaired by hepatitis C virus NS3

Yuri Kasama<sup>a</sup>, Makoto Saito<sup>a</sup>, Takashi Takano<sup>b</sup>, Tomohiro Nishimura<sup>c</sup>, Masaaki Satoh<sup>a,d</sup>, Zhongzhi Wang<sup>a</sup>, Salem Nagla Elwy Salem Ali<sup>a,e</sup>, Shinji Harada<sup>e</sup>, Michinori Kohara<sup>f</sup>, Kyoko Tsukiyama-Kohara<sup>a,\*</sup>

<sup>a</sup> Department of Experimental Phylaxiology, Faculty of Life Sciences, Kumamoto University, 1-1-1 Honjo Kumamoto City, Kumamoto 860-8556, Japan

<sup>b</sup> Division of Veterinary Public Health, Nippon Veterinary and Life Science University, 1-7-1 Kyonan, Musashino, Tokyo 180-8602, Japan

<sup>c</sup> KAKETSUKEN, Kyokushi, Kikuchi, Kumamoto 869-1298, Japan

<sup>d</sup> Department of Virology I, National Institute of Infectious Diseases, Tokyo 162-8640, Japan

<sup>e</sup> Department of Medical Virology, Faculty of Life Sciences, Kumamoto University, Japan

<sup>f</sup> Department of Microbiology and Cell Biology, Tokyo Metropolitan Institute of Medical Science, 2-1-6 Kamikitazawa, Setagaya-ku, Tokyo 156-8506, Japan

## ARTICLE INFO

## Article history:

Received 5 September 2011

Received in revised form 13 October 2011

Accepted 13 October 2011

Available online 20 October 2011

## Keywords:

HCV  
Tom70  
MAVS  
IFN  
IRF-3  
NS3

## ABSTRACT

Hepatitis C virus (HCV) elevated expression of the translocase of outer mitochondrial membrane 70 (Tom70). Interestingly, overexpression of Tom70 induces interferon (IFN) synthesis in hepatocytes, and it was impaired by HCV. Here, we addressed the mechanism of this impairment. The HCV NS3/4A protein induced Tom70 expression. The HCV NS3 protein interacted in cells, and cleaved the adapter protein mitochondrial anti-viral signaling (MAVS). Ectopic overexpression of Tom70 could not inhibit this cleavage. As a result, IRF-3 phosphorylation was impaired and IFN- $\beta$  induction was suppressed. These results indicate that MAVS works upstream of Tom70 and the cleavage of MAVS by HCV NS3 protease suppresses signaling of IFN induction.

© 2011 Elsevier B.V. All rights reserved.

Type I interferon (IFN) induction is the front line of host defense against viral infection. Intracellular double-stranded RNA is a viral replication intermediate and contains pathogen-associated molecular patterns (PAMPS) (Saito et al., 2008) that are recognized by pathogen-recognition receptors (PRRs) to induce IFN. One PRR family includes the Toll-like receptors (TLRs), which are predominantly expressed in the endosome (Heil et al., 2004). Another route of IFN induction takes place in the cytosol through activation of specific RNA helicases, such as retinoic acid-inducible (RIG)-I and melanoma differentiation associated gene 5 (MDA5). The ligand for RIG-I is an uncapped 5' triphosphate RNA, which is found in viral RNAs of the *Flaviviridae* family, including hepatitis C virus (HCV), paramyxovirus, and rhabdoviruses (Kato et al., 2006). MDA5 recognizes viruses with protected 5' RNA ends, for example,

picornaviruses (Hornung et al., 2006). The adapter protein that links the RNA helicase to the downstream MAPK, NF- $\kappa$ B, and IRF-3 signaling pathways is referred to as the mitochondrial anti-viral signaling (MAVS) protein (Seth et al., 2005); alternative names include IPS-1, interferon-promoter stimulator 1; VISA, virus-induced signaling adaptor; and CARDIF, CARD adapter inducing IFN. HCV nonstructural protein 3 (NS3) possesses a serine protease domain at the N terminus (amino acids (aa) 1–180) and has been found to cleave adaptor proteins, MAVS at aa 508 (Meylan et al., 2005) and Toll/IL-1R domain-containing adapter inducing IFN- $\beta$ -deficient (TRIF at aa 372; Ferreon et al., 2005). These cleavages provoke abrogation of the induction of the IFN pathway.

The translocase of the outer membrane (TOM) is responsible for initial recognition of mitochondrial preproteins in the cytosol (Baker et al., 2007; Neupert and Herrmann, 2007). The TOM machinery consists of 2 import receptors, Tom20 and Tom70, and, along with several other subunits, comprises the general import pore (Abe et al., 2000). Recently, Tom70 was found to interact with MAVS (Liu et al., 2010). Ectopic expression or silencing of Tom70, respectively, enhanced or impaired IRF3-mediated gene expression and IFN- $\beta$  production. Sendai virus infection accelerated the

\* Corresponding author. Present address: Transboundary Animal Diseases Center, Faculty of Agriculture, Kagoshima University, 1-21-24 Korimoto Kagoshima-shi, Kagoshima 890-0065, Japan. Tel.: +81 99 285 3589/96 373 5560; fax: +81 99 285 3589/96 373 5562.

E-mail address: [kkohara@kumamoto-u.ac.jp](mailto:kkohara@kumamoto-u.ac.jp) (K. Tsukiyama-Kohara).