

## HEMATOPOIESIS AND STEM CELLS

### Polymorphic *Sirpa* is the genetic determinant for NOD-based mouse lines to achieve efficient human cell engraftment

Takuji Yamauchi,<sup>1</sup> Katsuto Takenaka,<sup>1</sup> Shingo Urata,<sup>1</sup> Takahiro Shima,<sup>1</sup> Yoshikane Kikushige,<sup>1</sup> Takahito Tokuyama,<sup>1</sup> Chika Iwamoto,<sup>1</sup> Mariko Nishihara,<sup>1</sup> Hiromi Iwasaki,<sup>2</sup> Toshihiro Miyamoto,<sup>1</sup> Nakayuki Honma,<sup>3</sup> Miki Nakao,<sup>4</sup> Takashi Matozaki,<sup>5</sup> and Koichi Akashi<sup>1,2</sup>

<sup>1</sup>Department of Medicine and Biosystemic Science, Kyushu University Graduate School of Medical Sciences, Fukuoka, Japan; <sup>2</sup>Center for Cellular and Molecular Medicine, Kyushu University Hospital, Fukuoka, Japan; <sup>3</sup>Innovative Drug Research Laboratories, Kyowa Hakko Kirin Co Ltd, Tokyo, Japan;

<sup>4</sup>Department of Bioscience and Biotechnology, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University, Fukuoka, Japan; and

<sup>5</sup>Division of Molecular and Cellular Signaling, Department of Biochemistry and Molecular Biology, Kobe University Graduate School of Medicine, Kobe, Japan

#### Key Points

- NOD-specific *Sirpa* polymorphism is the genetic determinant of highly efficient xenograft activity in NOD-based immunodeficient mouse models.

Current mouse lines efficient for human cell xenotransplantation are backcrossed into NOD mice to introduce its multiple immunodeficient phenotypes. Our positional genetic study has located the NOD-specific polymorphic *Sirpa* as a molecule responsible for its high xenograft efficiency: it recognizes human CD47 and the resultant signaling may cause NOD macrophages not to engulf human grafts. In the present study, we established C57BL/6.*Rag2<sup>null</sup>Il2rg<sup>null</sup>* mice harboring NOD-*Sirpa* (BRGS). BRGS mice engrafted human hematopoiesis with an efficiency that was equal to or even better than that of the NOD.*Rag1<sup>null</sup>Il2rg<sup>null</sup>* strain, one of the best xenograft models. Consequently, BRGS mice are free from other NOD-related abnormalities; for example, they

have normalized C5 function that enables the evaluation of complement-dependent cytotoxicity of antibodies against human grafts in the humanized mouse model. Our data show that efficient human cell engraftment found in NOD-based models is mounted solely by their polymorphic *Sirpa*. The simplified BRGS line should be very useful in future studies of human stem cell biology. (*Blood*. 2013;121(8):1316-1325)

#### Introduction

Immunodeficient mice are widely used to reconstitute human hematopoiesis by xenotransplantation of hematopoietic stem cells (HSCs).<sup>1,2</sup> This “humanized” mouse model provides a powerful tool with which to evaluate the biologic properties of human HSCs and progenitors in vivo.<sup>3,4</sup> Such xenotransplantation systems have also been used to study human cancer stem cells.<sup>5-8</sup>

Elimination of the lymphoid system is the first step to achieving reconstitution of human hematopoiesis. To deplete T and B cells, the *scid* mutation in the *Prkdc* gene<sup>9-11</sup> or disruption of the recombination activating gene 1 or 2 (*Rag1* and *Rag2*)<sup>12,13</sup> has been introduced into various mouse strains. In addition, to deplete natural killer (NK) cells or their functions, the IL-2 receptor common  $\gamma$  chain subunit (*Il2rg*)<sup>14-16</sup> or beta-2-microglobulin (*B2m*)<sup>17-19</sup> is disrupted.

However, depletion of lymphoid cells is not sufficient and it has been shown empirically that additional strain-specific factors modulate human hematopoietic engraftment in the xenotransplantation setting. For example, within the SCID strain, the SCID with the NOD background was the gold standard for the xenotransplantation assay based on its high efficiency.<sup>11</sup> In fact, recent studies have shown that among the lymphoid-depleted mouse strains, the NOD-*scid* *Il2rg<sup>null</sup>* (NSG/NOG)<sup>14,15</sup> and NOD.*Rag1<sup>null</sup>Il2rg<sup>null</sup>*

(NOD-RG)<sup>20</sup> strains are the most efficient; the BALB/c.*Rag2<sup>null</sup>Il2rg<sup>null</sup>* (BALB-RG) strain is the next efficient<sup>21,22</sup>; and the C57BL/6 strains with *scid*,<sup>23</sup> *Rag2<sup>null</sup>*, *Rag2<sup>null</sup>B2m<sup>null</sup>*, *Rag2<sup>null</sup>Prf<sup>null</sup>*,<sup>24</sup> or *Rag2<sup>null</sup>Jak3<sup>null</sup>*<sup>25</sup> mutations are unable to reconstitute human hematopoiesis. The NOD strain has multiple immune deficiencies, including defects of appropriate regulation of the T-lymphocyte repertoire, antigen presenting cell function, NK cell function,<sup>26</sup> and hemolytic complement (C5) and cytokine production from macrophages,<sup>27</sup> and these abnormalities are presumed to collaborate to cause the development of autoimmune diabetes and hemolytic anemia.<sup>26,28</sup> To establish xenotransplantation models, lymphoid-depleted strains have been backcrossed into the NOD/ShiLt-inbred strain multiple times to introduce such numerous NOD-specific abnormalities.<sup>14,15</sup> However, it was unknown whether we could select a genetic determinant(s) specially required to achieve the NOD-specific high engraftment capability for human cells.

Previously, we used positional genetics to characterize the molecular basis for this capability in the NOD strain by measuring the ability of mouse BM stromal layers to support hematopoietic long-term culture-initiating cell activity (LTC-IC) in vitro and identified the strain differences as the polymorphism of the *Sirpa* gene located within the insulin-dependent diabetes (*Idd-13*) locus.<sup>24</sup>

Submitted June 28, 2012; accepted December 9, 2012. Prepublished online as *Blood* First Edition paper, January 4, 2013; DOI 10.1182/blood-2012-06-440354.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked “advertisement” in accordance with 18 USC section 1734.

The online version of this article contains a data supplement.

© 2013 by The American Society of Hematology

Stroma cells from the NOD BM supports LTC-IC of human cells, but those from C57BL/6 could not. Enforced expression of the NOD-type SIRPA enabled C57BL/6 stroma cells to support human LTC-IC.<sup>24</sup> This in vitro finding is also applicable to the in vivo setting, as shown by another study in which a human SIRPA BAC transgene introduced into *Rag2<sup>null</sup>Il2rg<sup>null</sup>* mice on a mixed 129; BALB/c background significantly improved the efficiency of human hematopoietic engraftment.<sup>29</sup>

SIRPA is a transmembrane protein that contains 3 Ig-like domains within the extracellular region. It is expressed in macrophages, myeloid cells, and neurons, and interacts with its ligand CD47 through its respective IgV-like domains, where the NOD strain has specific polymorphism. CD47 is a member of the Ig superfamily that is ubiquitously expressed in hematopoietic and nonhematopoietic cells. The cytoplasmic region of SIRPA has immunoreceptor tyrosine-based inhibitory motifs, and binding cell-surface CD47 with SIRPA on macrophages provokes inhibitory signals through phosphorylation of these inhibitory motifs of SIRPA,<sup>30</sup> preventing their phagocytic activity.<sup>31-33</sup> A recent study also showed that transgenic expression of mouse CD47 into CD34<sup>+</sup>CD38<sup>-</sup> human fetal liver cells significantly enhanced the human cell engraftment into BALB-RG mice.<sup>34</sup> Based on these data, the binding of NOD-SIRPA with human CD47 might produce signals for mouse macrophages not to engulf human HSCs, which presumably makes the strain permissive for human HSC engraftment.<sup>24</sup>

The most important question was whether the NOD-specific highly efficient human cell engraftment in vivo could be explained solely by the NOD-*Sirpa* polymorphism. In the present study, we established a C57BL/6.*Rag2<sup>null</sup>Il2rg<sup>null</sup>* (C57BL/6-RG) mouse line harboring the NOD-type *Sirpa*. Our data show clearly that replacement of the C57BL/6-type *Sirpa* with the NOD-type *Sirpa* is sufficient for the C57BL/6-RG strain to be endowed with the xenotransplantation capability that is at least equal to NOD-RG mice. Therefore, we successfully segregated the genetic abnormality responsible for efficient human cell engraftment from multiple genetic abnormalities in the NOD strain. The simplified humanized mouse system established by the new C57BL/6.*Rag2<sup>null</sup>Il2rg<sup>null</sup>*NOD-*Sirpa* (BRGS) strain should be very useful in improving xenotransplantation strategies in future studies of human cell biology.

## Methods

### Mice

C57BL/6, C57BL/6.NOD-*Idd13*, NOD, NOD.CB17-*Prkdc<sup>scid</sup>* (NOD-*scid*), and NOD.Cg-*Rag1<sup>tm1Mom</sup>Il2rg<sup>tm1Wjl</sup>/Sz* (NOD-RG) mice were purchased from the Jackson Laboratory; C57BL/6.*Rag2<sup>tm1Fva</sup>Il2rg<sup>tm1Wjl</sup>* (C57BL/6-RG) mice were purchased from Taconic. All mice were bred and maintained in individual ventilated cages at the Kyushu University Animal Facility and fed with autoclaved food and water. BRGS mice were generated by breeding C57BL/6-RG and C57BL/6.NOD-*Idd13* mice and backcrossed with C57BL/6-RG mice. *Rag2* gene and *Sirpa* gene are located on chromosome 2 with 17.1 cM. First, we repeated the breeding of C57BL/6-RG and C57BL/6.NOD-*Idd13* mice, and after 10 breedings, we obtained the recombination between the *Rag2*<sup>-</sup> and the *Sirpa*<sup>NOD</sup> loci by chromosomal crossover. This was examined by genotyping by the microsatellite markers *D2Mit447* and *D2Mit338*, which are 0.63 cM apart on chromosome 2, during interbreeding. In addition, *Sirpa*, *Rag2*, and *Il2rg* were genetically typed by PCR and direct sequencing. In C57BL/6.NOD-(*D2Mit447*-*D2Mit338*) *Rag2<sup>null</sup>Il2rg<sup>null</sup>* mice, the region between *D2Mit447* and *D2Mit338* contains 33 genes, including *Sirpa*, but *Sirpa* is the only gene

within the *Idd13* locus that is expressed in BM stromal cells and macrophages and had coding sequence polymorphism between the NOD and other strains.<sup>24</sup> Therefore, we refer to our established mouse line as BRGS herein. Sequences of the oligonucleotide primers used are provided in supplemental Table 1 (available on the *Blood* Web site; see the Supplemental Materials link at the top of the online article). All experiments were conducted following the guidelines of the institutional animal committee of Kyushu University.

### Binding affinity of mouse macrophages to human CD47-Fc

Mouse macrophages were obtained by peritoneal lavage. Cells were stained with purified anti-mouse Sirpa (P84; BD Biosciences) conjugated with PE and anti-mouse CD11b (3A33; Beckman Coulter) conjugated with FITC. CD11b<sup>+</sup>SIRPA<sup>+</sup> cells were defined as mature macrophages. The binding between SIRPA and CD47 was assessed by staining with biotinylated human CD47-Fc conjugated with streptavidin-allophycocyanin (APC),<sup>35</sup> and analyzed with a FACSAria III cell sorter (BD Biosciences).

### In vitro mouse macrophage phagocytosis assays for human hematopoietic stem cells

Phagocytic activity of mouse macrophages against the human CD34<sup>+</sup>CD38<sup>-</sup> population that contains the majority of human HSCs was evaluated in vitro, as described previously.<sup>36</sup> In brief, mouse peritoneal-derived macrophages were incubated at  $1.0 \times 10^4$  cells in 200  $\mu$ L of RPMI 1640 medium in Falcon culture tubes (2058; BD Biosciences). Cells were opsonized with CD34 antibody (sc-19621; Santa Cruz Biotechnology), incubated with mouse IFN- $\gamma$  (100 ng/mL; R&D Systems) for 24 hours, and then lipopolysaccharide (0.3  $\mu$ g/ $\mu$ L) for 1 hour. Human cord blood (CB) HSCs were then added to the tubes. Two hours after coinubation with macrophages and target cells, the phagocytic index was calculated using the following formula: phagocytic index = number of ingested cells/(number of macrophages/100). At least 200 macrophages were counted by a blinded observer.

### Sensitivity of BRGS mice to irradiation


Cohorts of BRGS mice were exposed to varying doses of the whole-body irradiation from a <sup>137</sup>Cs  $\gamma$ -irradiator. The mice were examined daily and euthanized when moribund. Surviving mice were euthanized at 8 weeks after irradiation. NOG/NSG mice are highly radiosensitive because of the *scid* mutation. To examine the radiosensitivity of BRGS mice, 6- to 10-week-old BRGS mice were irradiated with 550-670 cGy, and monitored for 8 weeks. Early deaths were observed in the mouse group irradiated with more than 620 cGy, whereas those irradiated with 550-580 cGy survived at the end of 8 weeks. Based on these data, we irradiated BRGS mice at 580 cGy in all xenotransplantation experiments. The irradiation doses for experiments with NOD-RG (420 cGy) and C57BL/6-RG (670 cGy) were decided by radiosensitivity experiments.

### Transplantation of human HSCs into mice

CB cells were collected during normal full-term deliveries after obtaining informed consent in accordance with the Declaration of Helsinki (provided by the Kyushu Block Red Cross Blood Center, Japan Red Cross Society). Mononuclear cells were separated by Ficoll-Hypaque density-gradient centrifugation. Lineage-depleted CB cells were obtained magnetically using a lineage cell depletion kit (Miltenyi Biotec). A total of  $5 \times 10^3$  CD34<sup>+</sup>CD38<sup>-</sup> cells were injected intrafemorally into mice. Within an individual experiment, mice of each strain received CD34<sup>+</sup>CD38<sup>-</sup> cells purified from the same mixture of CB cells from multiple donors. After transplantation, mice were given sterile water containing prophylactic enrofloxacin (Baytril; Bayer HealthCare). Mice were killed 8, 16, or 24 weeks after transplantation.

### Antibodies, cell staining, and sorting

For the analyses of mouse T, B, and NK cells, mouse peripheral blood cells were stained with PE-conjugated anti-CD3 (145-2C11), FITC-conjugated



**Figure 1. BRGS mice lack lymphocytes and SIRPA recognizes human CD47-Fc.** (A) Frequencies of blood leukocytes, erythrocytes, hemoglobin, and platelets in BRGS mice. Leukocyte counts in BRGS ( $2.69 \pm 1.01 \times 10^6/\mu\text{L}$ ) and NOD-RG mice ( $2.23 \pm 0.7 \times 10^6/\mu\text{L}$ ) are significantly decreased compared with that in C57BL/6 mice ( $9.6 \pm 0.32 \times 10^6/\mu\text{L}$ ). BRGS mice have normal erythrocyte ( $9.63 \pm 0.63 \times 10^6/\mu\text{L}$ ), hemoglobin ( $14.0 \pm 0.6$  g/dL), and platelet ( $8.7 \pm 2.0 \times 10^9/\mu\text{L}$ ) counts. (B) Representative FACS plots of blood in C57BL/6 and BRGS mice. BRGS mice lacked T, B, and NK cells. (C) Binding activity of human CD47-Fc to SIRPA expressed in peritoneal macrophages derived from C57BL/6-RG, BRGS, or NOD-RG mice. Macrophages from BRGS and NOD-RG mice, but not those from C57BL/6-RG mice, were stained with human CD47-Fc on FACS. (D) Phagocytosis assay of C57BL/6-RG, BRGS, or NOD-RG macrophages against human CD34<sup>+</sup>CD38<sup>-</sup> CB HSCs ( $n = 3$ ). The phagocytic index was determined as the number of engulfed cells per 100 macrophages. Bars indicate mean  $\pm$  SD.

anti-CD19 (1D3), APC-conjugated anti-NK1.1 (PK136; BD Biosciences), and Pacific Blue-conjugated anti-Gr-1 (RB6-8C5; BioLegend). Sorting of CD34<sup>+</sup>CD38<sup>-</sup> subfractions was accomplished by staining lineage-depleted CB cells with FITC-conjugated anti-CD34 (581/CD34) and PE-conjugated anti-CD38 (HIT2; BD Biosciences). For analysis and sorting of human cells in the immunodeficient mice, FITC-conjugated anti-CD4 (RPA-T4), CD33 (HIM3-4), CD41a (HIP8), TCR  $\alpha\beta$  (WT31), TCR  $\gamma\delta$  (11F2), Ig $\lambda$  L chain (JDC-12), Ig $\kappa$  L chain (G20-193; BD Biosciences), anti-CD10 (SS2/36; Dako), PE-conjugated anti-CD8 (RPA-T8), CD20 (2H7), NKp46 (9E2; BD Biosciences), CD235a (JC159; Dako), PE-Cy7-conjugated anti-CD3 (SK7; BD Biosciences), CD19 (HIB19; BioLegend), APC-conjugated anti-CD45 (J33; Beckman Coulter), and PaB-conjugated anti-mouse CD45 (30-F11; BioLegend) monoclonal antibodies were used in addition to the antibodies described in the preceding paragraph. Nonviable cells were excluded by propidium iodide staining. The cells were analyzed and sorted with a FACS Aria cell sorter (BD Biosciences).

#### Complement-dependent hemolytic activity

To estimate the serum complement activity of mice, the peripheral blood of mice were collected in 1.5-mL tubes and allowed to stand at room temperature for 1 hour. The serum was collected after centrifugation of the blood at 200g for 15 minutes at 4°C and stored  $-80^\circ\text{C}$  until use. The mixtures of each diluted sera of mice,  $3.75 \times 10^6$  erythrocytes of sheep and 2.5  $\mu\text{g}$  of zymosan (Imgenex) were incubated 10 hours at 37°C. After incubation, the absorbance of each sample at 415 nm was measured.

#### In vivo antibody treatment in a disseminated lymphoma xenograft model

A total of  $8 \times 10^5$  Raji cells (Burkitt lymphoma cell line; American Type Culture Collection) were injected into BRGS or NOD-RG mice (6–10 weeks of age) via the tail vein. Raji cells proliferated predominantly in the BM. Ten days after injection, these mice were IP injected daily with 200  $\mu\text{g}$  of

rituximab or mouse IgG2a control for 1 week and then BM cells were collected and analyzed with the FACS Aria III.

#### Statistical analysis

Data are presented as means  $\pm$  SD. The significance of the differences between groups was determined via the Student  $t$  test. For comparison of complement-dependent hemolytic activity among the mouse strains, repeated-measures ANOVA was performed.


## Results

### Establishment of the BRGS mouse

The BRGS mouse line was established by breeding the C57BL/6-RG with the C57BL/6.NOD-*Idd13* mouse that is congenic for NOD-derived *Idd13* locus within which the *Sirpa* is the only gene that is polymorphic and is expressed in the BM stromal cells.<sup>24</sup> BRGS mice were all born healthy and displayed good fertility. They showed a median life span of 65 weeks without the development of lymphoma that usually occurs in the NOD-*scid* strain after the age of  $> 5$  months.<sup>11</sup>

As shown in Figure 1A, BRGS mice had normal levels of hemoglobin and platelets, but a low number of leukocytes. This is because of the lack of CD3<sup>+</sup> T cells, CD19<sup>+</sup> B cells, and NK1.1<sup>+</sup> NK cells (Figure 1B). IP macrophages from either C57BL/6-RG, BRGS or NOD-RG mice were evaluated for the binding to human CD47 on FACS. CD11b<sup>+</sup> peritoneal macrophages strongly expressed SIRPA in all of these strains. As shown in Figure 1C, both macrophages from the BRGS and those from the NOD-RG strain bound to the human CD47-Fc protein, whereas those from the

**Figure 2. BRGS mice show efficient engraftment of human HSCs comparable to NOD-RG mice.** In the BM, human HSC engraftment was examined by flow cytometric analysis 8 weeks after transplantation. C57BL/6-RG mice (■; n = 5), female BRGS mice (●; n = 17), male BRGS mice (○; n = 12), female NOD-RG mice (▲; n = 13), and male NOD-RG mice (△; n = 5) mice were analyzed. (A) Both BRGS and NOD-RG female mice showed excellent human CD45<sup>+</sup> reconstitution. BRGS male mice showed significantly better engraftment compared with NOD-RG male mice. (B) Frequencies of CD19<sup>+</sup> B cells, CD33<sup>+</sup> myeloid cells, and CD34<sup>+</sup>CD38<sup>-</sup> HSCs in BRGS and NOD-RG mice.



C57BL/6 strain did not, confirming that BRGS mice have the NOD-type SIRPA that can bind to human CD47. Consistent with these binding data, when macrophages of each strain were cultured with human CD34<sup>+</sup>CD38<sup>-</sup> cells, macrophages from C57BL/6-RG mice, but not those from BRGS or NOD-RG mice, actively engulfed human CD34<sup>+</sup>CD38<sup>-</sup> cells, as shown by the significant elevation of the phagocytic index in the C57BL/6-RG mice (Figure 1D).


**BRGS mice are capable of multilineage reconstitution of human hematopoiesis with efficiency at least equal to that of NOD-RG mice**

A recent study has shown that intrafemoral injection is more efficient than IV injection in the xenotransplantation setting.<sup>37</sup> We used intrafemoral injection into adult mice in the present study because our preliminary data also showed that human cell chimerisms of adult BRGS by intrafemoral injection was significantly better than those with IV injection (data not shown). We transplanted  $5 \times 10^3$  CD34<sup>+</sup>CD38<sup>-</sup> human CB cells intrafemorally into C57BL/6-RG, BRGS or NOD-RG mice at the age of 6-8 weeks. Before transplantation, C57BL/6-RG, BRGS, and NOD-RG mice were irradiated with 670, 580, and 420 cGy, respectively. Each dose was set by irradiation tolerance experiments (see the Methods).

At 8 weeks after transplantation, human CD45<sup>+</sup> cells were not detectable in C57BL/6-RG mice (Figure 2A). Both BRGS and

NOD-RG showed successful reconstitution and their average frequencies of human CD45<sup>+</sup> cells were 59.9% and 55.8%, respectively. Recent studies have shown that in the NSG strain,<sup>15</sup> female recipients better support the reconstitution of human hematopoiesis, although the underlying mechanism for this remains unclear.<sup>38,39</sup> As shown in Figure 2A, NOD-RG and BRGS female mice showed equally excellent human CD45<sup>+</sup> reconstitution at approximately 70% chimerism. NOD-RG male mice, however, showed significantly poor reconstitution (16.6% of human cell chimerism on average) compared with NOD-RG female mice. In contrast, the percentages of human cell chimerisms in BRGS male mice (approximately 45%) were only slightly lower than those in BRGS female mice and, as a result, BRGS male mice showed significantly better engraftment compared with NOD-RG male mice.

In the BM, the percentages of CD19<sup>+</sup> B cells, CD33<sup>+</sup> myeloid cells, and CD34<sup>+</sup>CD38<sup>-</sup> cells that contain the majority of human HSCs were almost equal between the BRGS and the NOD-RG strains irrespective of sex (Figure 2B). Representative FACS plots at 8 weeks after injection are shown in Figure 3A. In the spleen, small numbers of CD3<sup>+</sup> T cells and CD3<sup>-</sup>NKp46<sup>+</sup> NK cells, as well as CD41<sup>+</sup> megakaryocytes and CD235a<sup>+</sup> erythrocytes, were found in both BRGS and NOD-RG mice (Figure 3A) and there was no significant difference in the percentages of these cells between the 2 strains regardless of sex. The majority of human cells in the spleen were CD19<sup>+</sup> B cells (Figure 3B). Although BM human



**Figure 3. Multilineage human HSC reconstitution in BRGS mice.** (A) Representative FACS plots at 8 weeks after transplantation in the BM, spleen, and thymus. (B) Human hematopoietic reconstitution in the spleens of BRGS and NOD-RG recipients (●: BRGS female; ○: BRGS male; ▲: NOD-RG female; △: NOD-RG male). There were no significant differences in the percentages of human CD45<sup>+</sup> cells and human CD19<sup>+</sup> B cells between these mice. (C) Human hematopoietic reconstitution in the thymus of BRGS and NOD-RG recipients. There were no significant differences in the percentages of human CD45<sup>+</sup> cells and human CD3<sup>+</sup> T cells between these mice. Symbols are as in panel B.

CD19<sup>+</sup> cells were mainly CD10<sup>+</sup>CD20<sup>-</sup> immature B cells, the majority (approximately 90%) of human spleen CD19<sup>+</sup> cells were CD10<sup>-</sup>CD20<sup>+</sup> mature B cells (data not shown). Thymic T cells were found in both the BRGS and NOD-RG strains, and the majority of human CD3<sup>+</sup> T cells in the thymus were CD4<sup>+</sup>CD8<sup>+</sup> immature T cells (Figure 3A,C).

Figure 4 shows the analysis of reconstitution of human HSCs at 16 weeks after transplantation. In this analysis, we used only female BRGS and NOD-RG mice. In the BM, both BRGS and NOD-RG mice showed sustained human cell engraftment and the frequencies of human CD45<sup>+</sup> cells were 64.4% and 51.1% in average, respectively, which were comparable to their levels at 8 weeks after transplantation. The percentages of CD33<sup>+</sup> myeloid cells, CD19<sup>+</sup> B cells, and CD34<sup>+</sup>CD38<sup>-</sup> HSCs were comparable to those at 8 weeks after transplantation (Figure 4A).

In the thymus, the percentage of CD3<sup>+</sup> T cells was increased up to approximately 80% and approximately 60% in the BRGS and NOD-RG strains, respectively. In addition to CD4<sup>+</sup>CD8<sup>+</sup> thymic precursors, both CD4<sup>+</sup> and CD8<sup>+</sup> single-positive T cells were present and expressed surface TCR- $\alpha\beta$  or TCR- $\gamma\delta$ , suggesting that human T-cell maturation occurs in the BRGS thymus, as has been shown previously in the NOG, NSG, and NOD-RG mouse lines<sup>14-16,20</sup> (Figure 4B). The number of CD20<sup>+</sup> mature B cells in the spleen was increased and they expressed surface Ig light chain  $\lambda/\kappa$ , reflecting their normal maturation (Figure 4C).


### The BRGS mouse maintains self-renewal of human HSCs in the long term

Figure 5A shows the changes in human cell chimerism in female BRGS mice in the long term. The frequency of human CD45<sup>+</sup> cells was maintained at a high level at least until 24 weeks after transplantation. B-cell frequencies gradually declined, but human myeloid, T, and NK cells progressively increased after engraftment (Figure 5B). The delayed reconstitution of these lineages of human cells has also been reported in studies using NSG mice.<sup>40,41</sup>

Figure 5C shows the results of the serial transplantation analysis. After confirmation of human cell engraftment at 8 weeks after the first transplantation,  $1 \times 10^6$  human CD45<sup>+</sup> cells were purified from primary BRGS recipients. These cells were transplanted into irradiated secondary BRGS recipients by intraperitoneal injection and tested for engraftment after another 8 weeks. Four of 6 secondary BRGS recipients showed multilineage engraftment of human CD33<sup>+</sup>, CD19<sup>+</sup>, and CD3<sup>+</sup> cells (Figure 5C). These data strongly suggest that BRGS mice can support long-term reconstitution and self-renewal of human HSCs.

### The BRGS mouse is useful for experiments using CDC of antibodies in the xenotransplantation setting

One of the problems in NOD-based xenograft models is that the cytotoxic activities of antibodies are unable to be evaluated in vivo



**Figure 4. Human hematolymphoid reconstitution at 16 weeks after transplantation.** (A) In the BM, BRGS mice showed sustained multilineage engraftment of human hematopoiesis at a level comparable to that in NOD-RG mice (●: BRGS female; ▲: NOD-RG female). (B) In the thymus, CD3<sup>+</sup> T cells were developed and their frequencies were comparable in BRGS and NOD-RG mice. On FACS analysis, cells were differentiated into CD4<sup>+</sup> and CD8<sup>+</sup> single-positive T cells expressing the surface TCR-αβ chain. (C) In the spleen, CD10<sup>+</sup>CD19<sup>+</sup>CD20<sup>+</sup> mature B cells expressing surface Ig light chain λ or κ chain were present.


in humanized mice. First, antibody-dependent cell-mediated cytotoxicity (ADCC) does not operate efficiently in xenotransplantation experiments because these strains of mice are deficient in NK cells, the major player for ADCC. In addition, immunodeficient phenotypes of the NOD strain include complement-dependent hemolytic activity due to a deficiency of C5,<sup>42</sup> which is essential for antibodies to exert complement-dependent cytotoxicity (CDC). All NOD-based immunodeficient strains have this abnormality, whereas the BRGS strain does not because it has a C57BL/6 background except for the NOD-type SIRPA. We tested CDC activity in C57BL/6-based strains, including the C57BL/6, C57BL/6-RG, and BRGS mice, and in NOD-based strains such as NOD, NOD-*scid*, and NOD-RG. As shown in Figure 6, sera from all of the C57BL/6-based strains, including the BRGS strain, showed CDC activities on sheep RBCs, whereas this was not found in any of NOD-based strains. There were no significant differences in CDC activities among the C57BL/6, C57BL/6-RG, and BRGS strains.

To determine whether BRGS mice had restored CDC in vivo,  $8 \times 10^5$  cells of Raji, a Burkitt lymphoma cell line expressing human CD45, was injected into BRGS or NOD-RG mice. Ten days after transplantation, either rituximab, an anti-CD20 antibody that has both CDC and ADCC activities, or a control IgG2a antibody

was administered IP for 7 days (Figure 7A) and the effect of antibody injection on elimination of Raji cells was evaluated. Representative results are shown in Figure 7B. In mice injected with control IgG2a, Raji cells rapidly proliferated up to approximately 90% in the BM of both BRGS and NOD-RG mice. In contrast, by injection of rituximab, percentages of human CD45<sup>+</sup> Raji cells were significantly decreased in BRGS mice (15.1%), whereas the percentages of human CD45<sup>+</sup> cells in NOD-RG mice were only slightly reduced by rituximab treatment (79.2%). Representative FACS data are shown in Figure 7C. These data clearly show that the CDC activity of antibodies was able to operate in the BRGS strain.

## Discussion

The NOD/ShiLt inbred mouse strain, which was originally developed by selecting cataract-prone strains,<sup>27</sup> exhibits susceptibility to the spontaneous development of autoimmune insulin-dependent diabetes mellitus (IDDM) and many other autoimmune disorders. The susceptibility to IDDM is polygenic and genetic loci associated with susceptibility to IDDM have been identified through the



**Figure 5. Evaluation of self-renewal of human HSCs in the BRGS mouse model.** (A) Change in frequency of human CD45<sup>+</sup> cells after transplantation. The level of human CD45<sup>+</sup> cells was maintained at a high level until 24 weeks after transplantation (8 weeks, n = 29; 16 weeks, n = 17; and 24 weeks, n = 4). (B) Change in the frequency of human CD33<sup>+</sup> myeloid cells, B cells, T cells, and NK cells in the BM and spleen during the 24 weeks after transplantation (□: 8 weeks; ▨: 16 weeks; and ▩: 24 weeks). Note that the B-cell numbers gradually decreased and were compensated for by myeloid, T, and NK cells. (C) To test the self-renewal ability of human HSCs maintained in the first recipient mice, 1 × 10<sup>6</sup> human CD45<sup>+</sup> cells were sorted from first-recipient mice and injected into second-recipient mice. Only female mice were used as recipients. After another 8 weeks, 4 of 6 BRGS secondary recipients showed multilineage engraftment of human CD33<sup>+</sup>, CD19<sup>+</sup>, and CD3<sup>+</sup> cells. Representative FACS plots are shown.


development of congenic mouse strains. More than 20 *Idd* loci have been identified. NOD mice display multiple aberrant immunophenotypes, and introduction of these abnormalities into immunodeficient mouse lines by multiple backcrossing accelerated human cell engraftment in xenotransplantation assays.<sup>11,14,15</sup>

In the current study, we present formal proof that under disruption of T, B, and NK cells, NOD-specific *Sirpa* polymorphism could explain the efficient human cell engraftment in the NOD strain. We replaced the *Idd13* locus of C57BL/6-RG mice with that of B6. The NOD-*Idd13* mouse has the C57BL/6 background but is congenic for the NOD-derived 23cM segment of chromosome 2 extending from microsatellite marker *D2Mit274* through *D2Mit343*.<sup>43</sup> In a previous study, we resolved the sequence corresponding to the phenotype of support of human LTC-IC to a region of 960 kilobases, within which coding regions of 14 genes reside. *Sirpa* was the only gene within the *Idd13* locus expressed in BM stromal cells and macrophages and had coding sequence polymorphism between the NOD and other strains.<sup>24</sup> To determine whether the efficient human cell engraftment in the NOD strains was completely dependent on the NOD-type SIRPA polymorphism, we compared the engraftment efficiency of the BRGS mouse with the NOD-RG mouse as a control, because in both strains RAG and  $\gamma$ c genes are disrupted to disturb lymphoid cell development. The NOD-RG strain displays the excellent human

cell engraftment comparable to the NOG/NSG strain<sup>21</sup> in which the SCID mutation instead of RAG-1 disruption is introduced. Our data show that the reconstitution activity of human hematopoiesis in BRGS mice is at least equal to that in NOD-RG mice in terms of engraftment levels and multilineage reconstitution. Therefore, replacement of the C57BL/6-*Sirpa* with the NOD-*Sirpa* is sufficient for the C57BL/6-RG strain to gain the human cell engraftment capability equal to the NOD-RG strain. NOD-SIRPA is able to bind human CD47, signaling of which inhibits activation of host macrophages to engulf human HSCs (Figure 1D), and therefore this signaling might be able to inhibit xenograft rejection.<sup>24</sup>


The polymorphism of *Sirpa* could explain the strain-specific trend toward human cell acceptability in xenotransplantation experiments. There are 20 amino acid differences in the sequences of *Sirpa* IgV domain between the NOD and B6 strains. Among these, 5 amino acid residues are unique for NOD compared with C57BL/6, BALB/c, ICR, and C3H. By testing their binding affinity to human CD47 and their ability to support human LTC-IC, we found that the xenograft capability-related NOD-specific polymorphism can be aggregated to a single location of polymorphism (C.I., K.T., S.U., T.Y., K.I., J.K., T.M., K.A., The efficient engraftment of human hematopoiesis in the BALB/c strain is mounted by BALB/c-specific *Sirpa* polymorphism that enhances binding affinity to human CD47, manuscript in preparation). In

**Figure 6. BRGS mice had CDC activity.** Sera from BRGS and C57BL/6-based mice showed CDC activity, whereas none of the NOD-based strains did. Five mice were analyzed in each strain. There were no statistical differences in CDC activities among the C57BL/6-based strains.



addition, we found recently that Balb/c mice also have another polymorphism at the *Sirpa* IgV domain. Protein-binding assays show that C57BL/6-SIRPA never binds to human CD47, but Balb/c-SIRPA and NOD-SIRPA showed modest and very high binding affinity, respectively, correctly reflecting their strain-specific graft efficiencies.<sup>44</sup> Furthermore, a recent study has shown that the enforced expression of human SIRPA by a human BAC transgene enables the 129;Balb/c.*Rag1*<sup>null</sup>/*Il2rg*<sup>null</sup> mouse to engraft

human cells as efficiently as the NSG mouse.<sup>29</sup> Therefore, in xenograft models, the degree of SIRPA-CD47 interaction decided by *Sirpa* polymorphism is one of the most critical factors to achieve efficient human cell engraftment. Further study is required to understand how the different binding affinity between these mouse polymorphic SIRPAs and human CD47 is translated into cytoplasmic signaling that leads to respective efficiency for xenotransplantation capabilities.



**Figure 7. CDC activity of antibodies is evaluable in vivo in the BRGS xenogeneic model.** (A) Experimental scheme of this experiment. Raji cells were injected into mice at the tail vein. Ten days after the injection, either rituximab or control IgG2a antibody (200 µg each) was injected IP daily for 1 week. (B) Frequencies of human CD45<sup>+</sup> Raji cells in the BM of BRGS and NOD-RG mice with or without rituximab injection. A significant reduction of Raji cells was found only in BRGS mice injected with rituximab. (C) Representative FACS plots of the BM cells of BRGS and NOD-RG mice after injection of rituximab or control IgG.



Several recent studies have shown that, in xenograft models, female mice somehow present significantly better reconstitution than do male mice.<sup>38,39</sup> It remains unclear whether sex-related factors such as steroid hormones can affect the engraftment of human HSCs. In the present study, the human cell chimerism obtained in the BRGS strain was quite high, reaching > 90% in 9 of 17 BRGS female mice, but none of the 13 NOD-RG female mice achieved that level at 8 weeks after transplantation (Figure 2A). Furthermore, although BRGS male mice displayed lower levels of human chimerism (approximately 45%), NOD-RG male mice showed significantly lower levels than did NOD-RG female mice, reaching only < 20% of human cell chimerism in average on our conditions (Figure 2A). As a result, the human cell chimerism in BRGS male mice was significantly better than that in NOD-RG male mice. Therefore, the BRGS mice showed a trend toward higher levels of human chimerism in both the males and the females. These results may suggest that unknown genetic abnormalities antagonizing human cell reconstitution can exist outside of the *Idd13* locus in the NOD strain.

There remain many unknown factors that affect the efficiency of human cell reconstitution in mouse xenotransplantation models. For example, the BRGS model is capable of long-term, multilineage human hematopoietic reconstitution, but human myeloid, T, and NK cell reconstitution were significantly delayed compared with the B-cell lineage (Figure 5B). This pattern of reconstitution is commonly observed in other xenotransplantation models.<sup>40,41</sup> Since the introduction of human cytokines such as thrombopoietin and membrane-bound SCF into humanized mouse models,<sup>45,46</sup> myeloid reconstitution has been accelerated, so the delay could have been due to insufficient cross-reactivity of mouse cytokines with human cytokine receptors. It is also possible that the mouse hematopoietic microenvironment, including putative myeloid or lymphoid niches, is not appropriate for human HSC development. The elucidation of such unknown factors is necessary to develop further efficient xenotransplantation models for future studies.

We have also shown herein the usefulness of the BRGS line in testing the function of killing antibodies via CDC activity. Because rituximab has both ADCC and CDC activity<sup>47</sup> and because NK cells, the major player for ADCC, are absent in efficient xenograft models such as NOG,<sup>14</sup> NSG,<sup>15</sup> and NOD-RG<sup>20</sup> mice, the disappearance of Raji cells after rituximab injection in the BRGS system must have been dependent largely on its CDC activity. Selective cell depletion by killing antibodies should be very useful in xenograft experiments, for example, in targeting cancer stem cells,<sup>48</sup> and in removing specific human cell component(s) from reconstituted human hematolymphopoiesis *in vivo*. Therefore, the BRGS humanized mouse model is applicable to future, more sophisticated xenograft experiments.

In summary, in the present study, we selected NOD-type polymorphic *Sirpa* from multiple abnormalities within the NOD

background and introduced it into the common C57BL6 mouse line together with *Rag2<sup>null</sup>/Il2rg<sup>null</sup>* mutations. The xenograft efficiency of the BRGS line was equal to, or even better than, the NOD-RG line, which is currently one of the best xenograft models. This result formally proves that NOD-specific *Sirpa* polymorphism is the genetic determinant of highly efficient xenograft activity in NOD-based immunodeficient mouse models. Sparing other NOD-specific abnormalities in this model also resulted in normalized C5 function, which should help in future studies using CDC activity of antibodies *in vivo*. The use of the BRGS line should also save time in introducing other genes for further modification of the line, keeping the high efficiency corresponding to the NOD-based models without performing multiple backcrosses. Therefore, this simplified mouse model should be very useful in future xenotransplantation experiments using human cells.

## Acknowledgments

The authors thank Atsushi Odawara and Yasuyuki Okawa for purification of the CD47-Fc protein and the Kyushu Block Red Cross Blood Center for providing the CB samples.

This work was supported in part by a grant-in-aid from the Ministry of Education, Culture, Sports, Science and Technology of Japan (to K.A. and K.T.), a grant-in-aid from the Ministry of Health, Labor and Welfare of Japan (to K.A.), the Takeda Science Foundation (to K.T.), the Cell Science Research Foundation (to K.T.), the Sumitomo Foundation (to K.T.), the Japan Leukemia Research Fund (to K.T.), and the Uehara Memorial Foundation (to K.A.).

## Authorship

Contribution: T.Y., K.T., and S.U. coordinated the project, designed and performed the experiments, analyzed the data, and wrote the manuscript; T.S., Y.K., T.T., and C.I. performed the experiments; M. Nishihara managed the mice; H.I., T. Miyamoto, and K.A. designed the experiments, reviewed the data, and edited the manuscript; N.H. provided the antibodies and technical advice; M. Nakao performed the experiments and provided technical advice; and T. Matozaki provided the antibodies and technical advice.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Koichi Akashi, MD, PhD, Department of Medicine and Biosystemic Science, Kyushu University Graduate School of Medical Sciences, 3-1-1 Maidashi, Higashi-ku, Fukuoka 812-8582, Japan; e-mail: akashi@med.kyushu-u.ac.jp.

## References

1. Ishikawa F, Yoshida S, Saito Y, et al. Chemotherapy-resistant human AML stem cells home to and engraft within the bone-marrow endosteal region. *Nat Biotechnol*. 2007;25(11):1315-1321.
2. Manz MG, Di Santo JP. Renaissance for mouse models of human hematopoiesis and immunobiology. *Nat Immunol*. 2009;10(10):1039-1042.
3. Bhatia M, Wang JC, Kapp U, Bonnet D, Dick JE. Purification of primitive human hematopoietic cells capable of repopulating immune-deficient mice. *Proc Natl Acad Sci U S A*. 1997;94(10):5320-5325.
4. Guenechea G, Gan OI, Dorrell C, Dick JE. Distinct classes of human stem cells that differ in proliferative and self-renewal potential. *Nat Immunol*. 2001;2(1):75-82.
5. Clarke MF, Dick JE, Dirks PB, et al. Cancer stem cells—perspectives on current status and future directions: AACR Workshop on cancer stem cells. *Cancer Res*. 2006;66(19):9339-9344.
6. Ailles LE, Weissman IL. Cancer stem cells in solid tumors. *Curr Opin Biotechnol*. 2007;18(5):460-466.
7. Bonnet D, Dick JE. Human acute myeloid leukemia is organized as a hierarchy that originates from a primitive hematopoietic cell. *Nat Med*. 1997;3(7):730-737.
8. Kikushige Y, Ishikawa F, Miyamoto T, et al. Self-renewing hematopoietic stem cell is the primary target in pathogenesis of human chronic lymphocytic leukemia. *Cancer Cell*. 2011;20(2):246-259.
9. Greiner DL, Hesselton RA, Shultz LD. SCID mouse models of human stem cell engraftment. *Stem Cells*. 1998;16(3):166-177.
10. McCune JM, Namikawa R, Kaneshima H, Shultz LD, Lieberman M, Weissman IL. The SCID-hu mouse:

- murine model for the analysis of human hematolymphoid differentiation and function. *Science*. 1988; 241(4873):1632-1639.
11. Shultz LD, Schweitzer PA, Christianson SW, et al. Multiple defects in innate and adaptive immunologic function in NOD/LtSz-scid mice. *J Immunol*. 1995;154(1):180-191.
  12. Shultz LD, Lang PA, Christianson SW, et al. NOD/LtSz-Rag1null mice: an immunodeficient and radioresistant model for engraftment of human hematolymphoid cells, HIV infection, and adoptive transfer of NOD mouse diabetogenic T cells. *J Immunol*. 2000;164(5):2496-2507.
  13. Shultz LD, Banuelos S, Lyons B, et al. NOD/LtSz-Rag1nullPfpnull mice: a new model system with increased levels of human peripheral leukocyte and hematopoietic stem-cell engraftment. *Transplantation*. 2003;76(7):1036-1042.
  14. Ito M, Hiramatsu H, Kobayashi K, et al. NOD/SCID/gamma(c)(null) mouse: an excellent recipient mouse model for engraftment of human cells. *Blood*. 2002;100(9):3175-3182.
  15. Shultz LD, Lyons BL, Burzenski LM, et al. Human lymphoid and myeloid cell development in NOD/LtSz-scid IL2R gamma null mice engrafted with mobilized human hematopoietic stem cells. *J Immunol*. 2005;174(10):6477-6489.
  16. Ishikawa F, Yasukawa M, Lyons B, et al. Development of functional human blood and immune systems in NOD/SCID/IL2 receptor (gamma) chain-(null) mice. *Blood*. 2005;106(5):1565-1573.
  17. Christianson SW, Greiner DL, Hesselton RA, et al. Enhanced human CD4+ T cell engraftment in beta2-microglobulin-deficient NOD-scid mice. *J Immunol*. 1997;158(8):3578-3586.
  18. Kollet O, Peled A, Byk T, et al. beta2 microglobulin-deficient (B2m(null)) NOD/SCID mice are excellent recipients for studying human stem cell function. *Blood*. 2000;95(10):3102-3105.
  19. Ishikawa F, Livingston AG, Wingard JR, Nishikawa S, Ogawa M. An assay for long-term engrafting human hematopoietic cells based on newborn NOD/SCID/beta2-microglobulin(null) mice. *Exp Hematol*. 2002; 30(5):488-494.
  20. Pearson T, Shultz LD, Miller D, et al. Non-obese diabetic-recombination activating gene-1 (NOD-Rag1 null) interleukin (IL)-2 receptor common gamma chain (IL2r gamma null) null mice: a radioresistant model for human lymphohaematopoietic engraftment. *Clin Exp Immunol*. 2008;154(2): 270-284.
  21. Brehm MA, Cuthbert A, Yang C, et al. Parameters for establishing humanized mouse models to study human immunity: analysis of human hematopoietic stem cell engraftment in three immunodeficient strains of mice bearing the IL2rgamma(null) mutation. *Clin Immunol*. 2010;135(1):84-98.
  22. Gorantla S, Sneller H, Walters L, et al. Human immunodeficiency virus type 1 pathobiology studied in humanized BALB/c-Rag2-/-gamma(c)-/- mice. *J Virol*. 2007;81(6):2700-2712.
  23. Christianson SW, Greiner DL, Schweitzer IB, et al. Role of natural killer cells on engraftment of human lymphoid cells and on metastasis of human T-lymphoblastoid leukemia cells in C57BL/6J-scid mice and in C57BL/6J-scid bg mice. *Cell Immunol*. 1996;171(2):186-199.
  24. Takenaka K, Prasolava TK, Wang JC, et al. Polymorphism in Sirpa modulates engraftment of human hematopoietic stem cells. *Nat Immunol*. 2007;8(12):1313-1323.
  25. Ono A, Hattori S, Kariya R, et al. Comparative study of human hematopoietic cell engraftment into BALB/c and C57BL/6 strain of rag-2/jak3 double-deficient mice. *J Biomed Biotechnol*. 2011;2011:539748.
  26. Kataoka S, Satoh J, Fujiya H, et al. Immunologic aspects of the nonobese diabetic (NOD) mouse. Abnormalities of cellular immunity. *Diabetes*. 1983;32(3):247-253.
  27. Makino S, Kunimoto K, Muraoka Y, Mizushima Y, Katagiri K, Tochino Y. Breeding of a non-obese, diabetic strain of mice. *Jikken Dobutsu*. 1980; 29(1):1-13.
  28. Oldenborg PA, Gresham HD, Chen Y, Izui S, Lindberg FP. Lethal autoimmune hemolytic anemia in CD47-deficient nonobese diabetic (NOD) mice. *Blood*. 2002;99(10):3500-3504.
  29. Strowig T, Rongvaux A, Rathinam C, et al. Transgenic expression of human signal regulatory protein alpha in Rag2-/-gamma(c)-/- mice improves engraftment of human hematopoietic cells in humanized mice. *Proc Natl Acad Sci U S A*. 2011; 108(32):13218-13223.
  30. Oshima K, Ruhul Amin AR, Suzuki A, Hamaguchi M, Matsuda S. SHPS-1, a multifunctional transmembrane glycoprotein. *FEBS Lett*. 2002;519(1-3):1-7.
  31. Barclay AN, Brown MH. The SIRP family of receptors and immune regulation. *Nat Rev Immunol*. 2006;6(6):457-464.
  32. Matozaki T, Murata Y, Okazawa H, Ohnishi H. Functions and molecular mechanisms of the CD47-SIRPalpha signalling pathway. *Trends Cell Biol*. 2009;19(2):72-80.
  33. Tsai RK, Discher DE. Inhibition of "self" engulfment through deactivation of myosin-II at the phagocytic synapse between human cells. *J Cell Biol*. 2008;180(5):989-1003.
  34. Legrand N, Huntington ND, Nagasawa M, et al. Functional CD47/signal regulatory protein alpha (SIRP(alpha)) interaction is required for optimal human T- and natural killer- (NK) cell homeostasis in vivo. *Proc Natl Acad Sci U S A*. 2011; 108(32):13224-13229.
  35. Motegi S, Okazawa H, Ohnishi H, et al. Role of the CD47-SHPS-1 system in regulation of cell migration. *EMBO J*. 2003;22(11):2634-2644.
  36. Kuriyama T, Takenaka K, Kohno K, et al. Engulfment of hematopoietic stem cells caused by down-regulation of CD47 is critical in the pathogenesis of hemophagocytic lymphohistiocytosis. *Blood*. 2012;120(19):4058-4067.
  37. Yahata T, Ando K, Sato T, et al. A highly sensitive strategy for SCID-repopulating cell assay by direct injection of primitive human hematopoietic cells into NOD/SCID mice bone marrow. *Blood*. 2003;101(8):2905-2913.
  38. Notta F, Doulatov S, Dick JE. Engraftment of human hematopoietic stem cells is more efficient in female NOD/SCID/IL-2Rgc-null recipients. *Blood*. 2010;115(18):3704-3707.
  39. McDermott SP, Eppert K, Lechman ER, Doedens M, Dick JE. Comparison of human cord blood engraftment between immunocompromised mouse strains. *Blood*. 2010;116(2):193-200.
  40. Traggiai E, Chicha L, Mazzucchelli L, et al. Development of a human adaptive immune system in cord blood cell-transplanted mice. *Science*. 2004; 304(5667):104-107.
  41. Tanaka S, Saito Y, Kunisawa J, et al. Development of mature and functional human myeloid subsets in hematopoietic stem cell-engrafted NOD/SCID/IL2rgammaKO mice. *J Immunol*. 2012;188(12):6145-6155.
  42. Baxter AG, Cooke A. Complement lytic activity has no role in the pathogenesis of autoimmune diabetes in NOD mice. *Diabetes*. 1993;42(11): 1574-1578.
  43. Prochazka M, Serreze DV, Frankel WN, Leiter EH. NOR/Lt mice: MHC-matched diabetes-resistant control strain for NOD mice. *Diabetes*. 1992;41(1):98-106.
  44. Iwamoto C, Takenaka K, Urata S, et al. The efficient engraftment of human hematopoiesis in the Balb/c strain is mounted by Balb/c-specific SIRPA polymorphism that enhances binding affinity to human CD47 [abstract]. *Blood (ASH Annual Meeting Abstracts)*. 2011;118(21):4008.
  45. Rongvaux A, Willinger T, Takizawa H, et al. Human thrombopoietin knockin mice efficiently support human hematopoiesis in vivo. *Proc Natl Acad Sci U S A*. 2011;108(6):2378-2383.
  46. Takagi S, Saito Y, Hijikata A, et al. Membrane-bound human SCF/KL promotes in vivo human hematopoietic engraftment and myeloid differentiation. *Blood*. 2012;119(12):2768-2777.
  47. Chao MP, Alizadeh AA, Tang C, et al. Anti-CD47 antibody synergizes with rituximab to promote phagocytosis and eradicate non-Hodgkin lymphoma. *Cell*. 2010;142(5):699-713.
  48. Kikushige Y, Shima T, Takayanagi S, et al. TIM-3 is a promising target to selectively kill acute myeloid leukemia stem cells. *Cell Stem Cell*. 2010; 7(6):708-717.

## TRANSPLANTATION

### Quantitation of hematogones at the time of engraftment is a useful prognostic indicator in allogeneic hematopoietic stem cell transplantation

Takahiro Shima,<sup>1,2</sup> Toshihiro Miyamoto,<sup>1</sup> Yoshikane Kikushige,<sup>1,2</sup> Yasuo Mori,<sup>1,2</sup> Kenjiro Kamezaki,<sup>1</sup> Ken Takase,<sup>3</sup> Hideho Henzan,<sup>3</sup> Akihiko Numata,<sup>3</sup> Yoshikiyo Ito,<sup>4</sup> Katsuto Takenaka,<sup>1</sup> Hiromi Iwasaki,<sup>2</sup> Tomohiko Kamimura,<sup>4</sup> Tetsuya Eto,<sup>3</sup> Koji Nagafuji,<sup>1</sup> Takanori Teshima,<sup>2</sup> Koji Kato,<sup>1</sup> and Koichi Akashi<sup>1,2</sup>

<sup>1</sup>Department of Medicine and Biosystemic Science and <sup>2</sup>Center for Cellular and Molecular Medicine, Graduate School of Medical Sciences, Kyushu University Graduate School of Medicine, Fukuoka, Japan; <sup>3</sup>Department of Hematology, Harasanshin Hospital, Fukuoka, Japan; and <sup>4</sup>Department of Hematology, Hamanomachi General Hospital, Fukuoka, Japan

#### Key Points

- Quantitation of hematogones at engraftment is useful to predict prognosis of patients treated with allogeneic stem cell transplantation.

Transient marrow expansion of normal B-cell precursors, termed hematogones, is occasionally observed after hematopoietic stem cell transplantation (HSCT). To understand the clinical significance of this phenomenon, we enumerated hematogones in 108 consecutive patients who received allogeneic HSCT for the treatment of hematologic malignancies, including acute myelogenous leukemia, advanced myelodysplastic syndromes, acute lymphoblastic leukemia, and non-Hodgkin lymphoma. Hematogone quantitation was performed at the time of complete donor engraftment (median day 25 and 32 in patients who received bone marrow and cord blood cell transplants, respectively).

Hematogones were polyclonal B cells, and their frequencies correlated positively with blood B-cell numbers, and inversely with donors' but not recipients' age, suggesting that hematogones reflect cell-intrinsic B-cell potential of donor cells. Interestingly, patients developing hematogones that comprised > 5% of bone marrow mononuclear cells constituted a group with significantly prolonged overall survival and relapse-free survival, irrespective of their primary disease or donor cell source. In addition, patients with > 5% hematogones developed severe acute graft-versus-host diseases less frequently, which may contribute toward their improved survival. We therefore conclude that the amount of hematogones at the time of engraftment may be a useful tool in predicting the prognosis of patients treated with allogeneic HSCT. (*Blood*. 2013;121(5):840-848)

#### Introduction

Hematogones are transient increases in lymphoblast-looking cells in the bone marrow.<sup>1,2</sup> Because of the morphologic resemblance between residual leukemic clones and hematogones, expansion of hematogones during the recovery phase from chemotherapy and bone marrow transplantation occasionally causes diagnostic confusion.<sup>1,3</sup> Phenotypic analyses have demonstrated that hematogones are normal B-cell precursors, including pro-B, pre-B, and immature B cells coexpressing CD10 and CD19.<sup>1,2</sup> The fact that hematogones become prominent in the recovery phase after chemotherapy or hematopoietic stem cell transplantation (HSCT)<sup>1-6</sup> suggests that they could reflect active B-cell reconstitution. They are also sometimes seen in steady-state hematopoiesis, especially in healthy infants and young people.<sup>2,4,7,8</sup> Previous work demonstrated that in the recovery phase after chemotherapy, the percentage of hematogones in the bone marrow was inversely correlated with patients' age.<sup>1</sup> However, it is unclear whether the age-associated decline in hematogones frequency reflects cell-intrinsic defects of hematopoietic stem cell activity or cell-extrinsic defects such as aging of the bone marrow microenvironment.

Recent reports have shown that hematogone expansion correlates with favorable outcomes in acute myelogenous leukemia

(AML) patients treated with chemotherapy<sup>5</sup> or cord blood transplantation (CBT).<sup>6</sup> However, the precise number or frequency above which hematogones correlate with clinical significance has not been clarified. Previous reports<sup>1,5,6</sup> have reported hematogone frequency relative to bone marrow mononuclear cells (MNCs), total nuclear cells (TNCs), and frequencies of B-cell precursors, and as a result, hematogone expansion has been described with frequencies ranging from > 0% to 0.9%.<sup>1,5,6</sup>

To better understand the etiology and clinical significance of hematogones, we measured percentages of B-cell precursors in the bone marrow via flow cytometry in 108 consecutive patients with hematologic malignancies, including AML, advanced myelodysplastic syndromes (MDS), acute lymphoblastic leukemia (ALL), and lymphoma, who achieved successful engraftment after allogeneic HSCT at our institution. The analysis of hematogones was performed on the day of engraftment, defined as the day when circulating granulocytes reached  $> 0.5 \times 10^9/L$  for 3 consecutive days,<sup>9-12</sup> and the bone marrow showed complete donor-type chimerism via polymerase chain reaction (PCR) analysis. To minimize the effect of expanding granulocytes on hematogone frequencies, bone marrow MNCs were used for flow cytometric

Submitted February 7, 2012; accepted November 19, 2012. Prepublished online as *Blood* First Edition paper, December 11, 2012; DOI 10.1182/blood-2012-02-409607.

The online version of this article contains a data supplement.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2013 by The American Society of Hematology

**Table 1. Patients' characteristics**

	BMT: no. (%), [range]	CBT: no. (%), [range]	Total (%)	P
Recipient sex, male/female	35 (32)/24 (22)	26 (24)/23 (21)		NS
Recipient age, y	49.2 (mean) [20-66]	47.3 (mean) [19-67]		NS
Donor sex, male/female	37 (34)/22 (20)	24 (22)/25 (23)		NS
Donor age, y	36.7 (mean) [17-66]	0		< .001
No. of infused cell, /kg	$2.80 \times 10^8$ (mean) [0.92-4.02]	$0.28 \times 10^8$ (mean) [0.18-0.50]		< .001
<b>Primary disease</b>				NS
<b>AML/advanced MDS</b>				
CR	14 (13)	11 (10)	25 (23)	
non-CR	21 (19)	14 (13)	35 (32)	NS
Total	35 (32)	25 (23)	60 (56)	
<b>ALL</b>				
CR	3 (3)	3 (3)	6 (6)	
non-CR	2 (2)	10 (9)	12 (11)	NS
Total	5 (5)	13 (12)	18 (17)	
<b>Lymphoma</b>				
CR	7 (6)	3 (3)	10 (9)	
non-CR	12 (11)	8 (7)	20 (19)	NS
Total	19 (18)	11 (10)	30 (28)	
<b>Conditioning regimen</b>				< .01
TBI/CY	28 (26)	24 (22)		
BU/CY	14 (13)	0 (0)		
RIC	17 (16)	25 (23)		
<b>GVHD prophylaxis</b>				< .001
TAC + sMTX	51 (47)	6 (6)		
CSP + sMTX	8 (7)	28 (26)		
CSP + MMF	0 (0)	15 (14)		
<b>HLA disparity</b>				< .001
6/6	39 (36)	1 (1)		
5/6	20 (19)	7 (6)		
4/6	0 (0)	22 (20)		
3/6	0 (0)	19 (18)		
Days required for engraftment	25 (median) [15-32]	32 (median) [14-39]		< .01

ALL indicates acute lymphoblastic leukemia; AML, acute myelogenous leukemia; BMT, bone marrow transplantation; BU, busulfan; CBT, cord blood transplantation; CR, complete remission; CSP, cyclosporine; CY, cyclophosphamide; GVHD, graft-versus-host disease; MDS, myelodysplastic syndrome; MMF, mycophenolate mofetil; NS, not significant; RIC, reduced-intensity conditioning; sMTX, short-term methotrexate; TAC, tacrolimus; and TBI, total body irradiation.

analyses in all cases. Our data suggest that the number of hematogones generally reflects cell-intrinsic B-cell potential of donor hematopoietic stem cells (HSCs) and that this declines with aging. We also found that hematogone frequencies of > 5% of total MNCs is a useful cutoff line to distinguish patient groups with significantly better overall survival (OS) or with relapse-free survival (RFS), irrespective of their primary diseases or donor cell sources. We propose that the quantitation of hematogones at engraftment may be useful to predict the prognosis of patients treated with allogeneic HSCT.

## Methods

### Patients

From 2005 to 2010, 134 patients with high-grade hematologic malignancies were treated with allogeneic HSCT in Kyushu University Hospital. These patients included AML cases with high risk,<sup>13</sup> relapsed or refractory status, advanced MDS cases with intermediate-II or high risk on International Prognostic Scoring System classification,<sup>14,15</sup> ALL cases with high risk,<sup>16</sup> relapsed or refractory status, and relapsed non-Hodgkin lymphoma cases. Within these 134 patients, grafts were rejected in 5 cases and residual malignant cells proliferated soon after HSCT in 21 cases, without achieving successful engraftment. The remaining consecutive 108 cases, in which allogeneic HSCT was successful and complete donor-type chimerism was documented, were enrolled in this study. Fifty-nine and 49 patients received bone marrow transplantation (BMT) and CBT, respectively. Patients'

characteristics are summarized in Table 1. This study was approved by the institutional review board of Kyushu University Hospital and conducted in accordance with the Declaration of Helsinki.

### Evaluation of remission status before HSCT

Before HSCT, patients were intensively searched for residual malignant cells to define their pretransplantation remission status. In acute leukemia or advanced MDS cases, bone marrow samples were checked first by microscopic analysis, and were subjected to multicolor flow cytometric analysis.<sup>13,17</sup> Complete remission (CR) was diagnosed when percentages of cells of leukemia phenotype were < 0.5% in the bone marrow. Furthermore, 21 patients with acute leukemia or MDS had leukemia-specific genes such as BCR-ABL, FLT3-ITD, AML1-ETO, and MLL fusions, and PCR amplification of these genes were used to detect minimal residual disease (MRD).<sup>13</sup> Within these 21 patients, 17 patients were diagnosed as CR based on flow cytometric analyses. CR results for these 17 patients were also confirmed by PCR. In lymphoma patients, remission status was defined by evaluating the involvement of lymphoid organs using FDG-PET CT scan and/or MRI methods, and was also defined by evaluating the involvement of bone marrow by flow cytometry, as previously described.<sup>18</sup>

### Transplantation procedures

Patients' characteristics were not statistically different between BMT and CBT recipient groups in terms of sex, age, and primary disease (Table 1). Conditioning regimen consisted of total body irradiation/cyclophosphamide (CY) for 28 BMT and 24 CBT recipients, busulfan (BU)/CY for 14 BMT recipients, and fludarabine-based reduced-intensity conditioning<sup>19,20</sup> for 17 BMT and 25 CBT recipients, respectively (Table 1).

Prophylaxis for graft-versus-host disease (GVHD) was tacrolimus/short-term methotrexate (sMTX) for 51 BMT and 6 CBT recipients, cyclosporine (CSP)/sMTX for 8 BMT and 28 CBT recipients, and CSP/mycophenolate mofetil for 15 CBT recipients (Table 1). The mean number of donor cells transplanted was  $2.8 \times 10^8/\text{kg}$  in BMT recipients and  $0.28 \times 10^8/\text{kg}$  in CBT recipients. Bone marrow units were obtained from the Japan Marrow Donor Program or related donors, and cord blood units were obtained from the Japanese Cord Blood Bank Network.

### Evaluation for engraftment

The bone marrow sampling for the analysis of hematogones was performed when patients achieved successful engraftment. The standard criterion for engraftment was used according to previous studies.<sup>9-12</sup> Blood neutrophil numbers were checked daily after transplantation, and the successful engraftment was defined when neutrophils exceeded  $0.5 \times 10^9/\text{L}$  for 3 consecutive days. When patients met the criteria for engraftment, host/donor microchimerism analysis was performed (see the next section). If the analysis showed complete donor type chimerism, hematogones in the bone marrow were counted by multicolor flow cytometric analysis.

### Chimerism analysis

To analyze donor/recipient cell chimerism, PCR amplification of polymorphic short tandem repeats (STR) was performed to confirm engraftment of donor cells. PCR using synthesized oligonucleotide templates were performed using TAKARA Taq Reagent Kits and run in the Perkin Elmer GeneAmp PCR system 9600 or 2400. The donor-cell origin and recipient-cell origin PCR product mixture was loaded onto the 373A sequencer (Applied Biosystems) with a size marker, and the data were processed using the GeneScan software (Applied Biosystems) as described previously.<sup>21</sup>

### Flow cytometry analysis and cell sorting

The bone marrow mononuclear cells were prepared by the gradient centrifugation method as previously described.<sup>22,23</sup> Cells were stained with allophycocyanin-conjugated anti-CD34 (BD Pharmingen), biotin-conjugated anti-CD38 (Caltag Laboratories), FITC-conjugated anti-CD10 (DAKO), PE-conjugated anti-CD20 (BD Biosciences), PE-Cy7-conjugated anti-CD19 (BioLegend), and Cy5-PE-conjugated lineage (Lin) mixture (anti-CD3, -CD4, -CD8 (BD Pharmingen) -CD11b (Caltag Laboratories), -CD14, and -CD56 (Beckman Coulter)).<sup>22-25</sup> Streptavidin-conjugated Cy7-allophycocyanin (BD Pharmingen) was used for visualization of biotinylated antibodies. For analysis of mature B cells, peripheral blood (PB) cells were stained with FITC-conjugated anti-CD10 (DAKO), PE-conjugated anti-CD20 (BD Biosciences), PE-Cy7-conjugated anti-CD19 (BioLegend), and Cy5-PE-conjugated Lin mixture. Available PB cells at day 90 after HSCT could be obtained from 64 patients and evaluated. Cells were analyzed by using a FACSAria (BD Biosciences) or FACSCanto (BD Biosciences). Cell sorting was performed on a 5-color FACSAria (BD Biosciences). To minimize contamination, cells were collected after the second round of sorting using sorting gates identical to those used in the first-round sorting. Definition of hematogones is a series of normal B-lymphoid precursors, including CD34<sup>+</sup>CD38<sup>+</sup>CD10<sup>+</sup>CD19<sup>+</sup>Lin<sup>-</sup> pro-B cells, CD34<sup>-/lo</sup>CD38<sup>+</sup>CD10<sup>+</sup>CD19<sup>+</sup> pre-B cells, and CD34<sup>-</sup>CD38<sup>+</sup>CD10<sup>+</sup>CD19<sup>+</sup> CD20<sup>+</sup> immature B cells<sup>26-28</sup> in bone marrow MNCs. Isotype controls were used to define the cutoff of positivity of each antigen on a FACS.

### PCR analysis of IGH gene rearrangement

To analyze clonality of IGH gene rearrangements status of hematogones, DNA was obtained from 10 000 double-sorted cells<sup>22,23,29</sup> from all recipients presenting > 0.1% MNCs of hematogones on FACS. Then PCR amplification of DJ<sub>H</sub> and VDJ<sub>H</sub> gene rearrangement was performed as described previously.<sup>24,30</sup>

### Statistical analysis

Relationships of percentages of hematogones with age, the day of engraftment, and numbers of circulating B lymphocytes were analyzed with the

Spearman rank correlation analysis. Comparison between 2 groups or condition was tested with the Mann-Whitney *U* test. The categorical variables were analyzed with the 2-tailed  $\chi^2$  test. Survival was plotted with Kaplan-Meier curves, taking the interval from date of HSCT to death/relapse or last contact. Comparisons between each group were performed with the log-rank test and the Cox proportional hazards model. Univariate analysis was performed with logistic or exact logistic regression, and the parameters that present  $P < .20$  were reevaluated by multivariate analysis.<sup>31</sup> Multivariate analysis was performed with logistic regression applying Firth's bias reduction. A  $P$  value < .05 was considered to be statistically significant.

## Results

### Hematogones that appeared at the time of engraftment are polyclonal B-cell precursors

One hundred eight consecutive cases treated with successful allogeneic BMT or CBT were enrolled in this study. Hematogones in the bone marrow were counted on the day of engraftment by multicolor flow cytometric analysis. The successful engraftment was judged when neutrophils exceeded  $> 0.5 \times 10^9/\text{L}$  for 3 consecutive days.<sup>9-12</sup> At this phase, it is critical to exclude residual leukemic cells or host-derived B-cell precursors from a cell fraction of hematogones. To this end, polymorphic STR was amplified to test the host/donor microchimerism, and only when patients' bone marrow consisted of 100% donor-derived cells, the analysis of hematogones was performed. The complete donor-type chimerism verifies that host-derived normal hematopoietic cells and malignant leukemic cells have been eliminated below the sensitivity of FACS,<sup>32-34</sup> and therefore that phenotypically defined hematogones in these patients on FACS were donor-derived normal cells.


Hematogones were morphologically blastic cells (Figure 1A), and were identified by surface phenotype, according to the definition of pro-B, pre-B, and/or immature B cells that coexpress CD10 and CD19 on their cell surface<sup>1,26-28</sup> (Figure 1B). To minimize the effect of granulocytes on hematogone frequencies, we used MNCs instead of TNCs in our analysis. The frequencies of hematogones in MNCs are usually higher than those in TNCs (not shown), as reported previously.<sup>1</sup>

The time median to engraftment was 25 and 32 days in patients treated with BMT and CBT, respectively (Table 1). The time required for engraftment appears to be consistent with previous reports.<sup>34-39</sup> Percentages of B-cell precursors within the bone marrow MNCs at the time of complete donor-type engraftment were significantly higher in CBT recipients than in BMT recipients (6.37% vs 1.75%;  $P < .001$ ; Figure 1C). There was no significant relationship between the day of engraftment (the day of sampling) and the frequency of hematogones (Figure 1D).

In 106 of 108 patients who had > 0.1% of B-cell precursors in the bone marrow MNCs, B-cell precursors were purified by a multicolor FACS and were subjected to IGH rearrangement analysis. In all of these patients, B-cell precursors were polyclonal based on the rearrangement analysis of the IGH genes (Figure 1E).

### Hematogones generally represent B-cell recovery potential of the graft and their emergence is related to age of donors but not recipients

Because hematogones are normal B-cell precursors, we tested whether the presence of a high number of them could reflect the active B-cell recovery after HSCT. FACS analysis of circulating blood cells revealed that the frequency of bone marrow B-cell precursors was significantly correlated with the number of blood



**Figure 1. Detection of hematogones after allogeneic HSCT.** (A) Typical appearance of hematogones in the bone marrow after HSCT (Giemsa staining  $\times 1000$ ; OLYMPUS BH-2 microscope [Olympus]; ACT-2U imaging software [Nikon];  $27^{\circ}\text{C}$ ). (B) Evaluation of hematogones on a 5-color FACS. Hematogones are defined as MNCs coexpressing CD10 and CD19 in the bone marrow at engraftment. They include  $\text{CD}34^{+}\text{CD}38^{+}\text{CD}10^{+}\text{CD}19^{+}\text{Lin}^{-}$  pro-B cells,  $\text{CD}34^{\text{low}}\text{CD}38^{+}\text{CD}10^{+}\text{CD}19^{+}$  pre-B cells, and  $\text{CD}34^{-}\text{CD}38^{+}\text{CD}10^{+}\text{CD}19^{+}\text{CD}20^{+}$  immature B cells. (C) Percentage of hematogones in the bone marrow MNCs in patients who received BMT and CBT. CBT recipients presented much higher frequency of hematogones compared with BMT recipients ( $P < .001$ ). Solid bars indicate the median percentage of hematogones for each recipient; MNC, mononuclear cells; BMT, bone marrow transplantation; and CBT, cord blood transplantation. (D) The relationship between the day of engraftment and percentages of hematogones. There was no relationship between these parameters. (E) *IGH* rearrangement analysis of purified hematogones. B-cell precursors were polyclonal in all 106 recipients analyzed.


B cells at the time of engraftment ( $R = 0.47, P < .001$ ; Figure 2A), and with those even on day 90 after HSCT ( $R = 0.22, P < .01$ ; Figure 2B). These results suggest that expansion of hematogones reflects not only enhanced B-cell reconstitution potential of the graft, but also prolonged B cell-producing capability of donor HSCs.

The age of BMT donors ranged from 17 to 66 years old (median, 37 years; Table 1). Interestingly, there was a significant inversed correlation between donor age and percentage of bone marrow hematogones in patients treated with BMT ( $R = 0.32, P = .02$ ; Figure 2C blue line). When the age of CBT donors were defined as 0-year old, the significant inversed correlation between age and hematogone numbers was also found in all patients entered in this study ( $R = 0.42, P < .001$ ; Figure 2C black line). In contrast, recipients' age and hematogone numbers did not show any relationship (Figure 2D). Furthermore, as shown in Table 2, the time of engraftment was not affected by primary diseases of patients, or by their remission status at the time of HSCT. Thus, although the patients who fail to achieve CR are usually treated with higher total doses of chemotherapeutic drugs because of their refractory disorders, it did not affect the day of engraftment or the day of hematogone analysis for this study. These data strongly suggest that the number of hematogones after HSCT generally reflects the cell-intrinsic B-cell recovery potential of donor HSCs, which may decline by aging.

**The emergence of hematogones up to > 5% of MNCs in the bone marrow represents a good prognosis for patients treated with allogeneic HSCT**

It should be critical to draw a line of hematogone numbers to distinguish a group of patients with clinical significance. Therefore, we first compared the OS and RFS among patient subgroups with  $\leq 1\%$ ,  $1\%-2\%$ ,  $2\%-3\%$ ,  $3\%-4\%$ ,  $4\%-5\%$ , or  $> 5\%$  of hematogones in our study (Figure 3A-B). Strikingly, patients who developed hematogones up to  $> 5\%$  of MNCs showed significantly better 3-year OS (100%) and RFS (93.3%), compared with any other group. Patient groups with  $\leq 1\%$ ,  $1\%-2\%$ ,  $2\%-3\%$ , and  $3\%-4\%$  of hematogones showed similar 3-year OS and RFS that were 37%-53% and 22%-51%, respectively. Interestingly, patients with  $4\%-5\%$  hematogones appeared to show intermediate levels of OS (86%) and RFS (64%), although this is not statistically better than those in patients with  $\leq 1\%$  hematogones (Figure 3A-B). Based on these results, we hypothesized that the development of  $> 5\%$  of hematogones might be critical to distinguish a patient group with favorable prognosis.

According to this criteria, 43 patients developed  $> 5\%$  MNCs of hematogones ( $\text{HG}^{+}$ ) and the remaining 65 patients had  $\leq 5\%$  MNCs of hematogones ( $\text{HG}^{-}$ ). As shown in Figure 3A, in  $\text{HG}^{+}$  patients, 3-year OS was 100%, whereas in  $\text{HG}^{-}$  patients, it was 45% ( $P < .001$ ). The favorable OS in  $\text{HG}^{+}$  groups is at least



**Figure 2. Analysis of hematogones, and the correlation of their frequency compared with blood B-cell numbers and age of donors.** (A) A relationship between frequencies of hematogones and blood B cells at engraftment ( $P < .001$ ). (B) A relationship between frequencies of hematogones at engraftment and blood B cells on day 90 ( $P < .01$ ). (C) A relationship between frequency of hematogones and donor's age in patients who received BMT (blue line,  $P = .023$ ), and in all recipients treated with either BMT or CBT (black line,  $P < .001$ ). (D) No significant relationship was observed between frequency of hematogones and recipient age. NS indicates not significant.

because of the less frequent disease relapse. As shown in Figure 3B, significant association was observed between the presence of hematogones and 3-year RFS after HSCT: 3-year RFS was 93% and 37% in  $HG^+$  and  $HG^-$  patients, respectively ( $P < .001$ ). The association between the presence of  $> 5\%$  hematogones and favorable OS and RFS was also seen when the analysis was performed in patient subgroups that received either BMT or CBT (Figure 3C-D). These data strongly suggest that the emergence of hematogones is a useful predictor of favorable outcomes at least in terms of OS and RFS, irrespective of donor cell source.

**The emergence of hematogones ( $> 5\%$  of MNCs) marks favorable outcomes for allogeneic HSCT especially in patients who failed to achieve complete remission, irrespective of primary malignant disease**

We then analyzed whether the good prognosis designated by the emergence of hematogones is dependent on the primary malignant disorder. The OS and RFS were analyzed in each patient group with AML/advanced MDS, ALL, or non-Hodgkin lymphoma. As shown in Figure 4,  $HG^+$  patients always showed significantly better OS and RFS compared with  $HG^-$  patients, in any of these patients groups suffering from different primary diseases.

It is well known that the achievement of CR at the time of transplantation favorably affects the prognosis after allogeneic HSCT.<sup>13</sup> Interestingly, in AML/advanced MDS patients, the  $HG^+$  group showed significantly prolonged OS and RFS compared with the  $HG^-$  group, irrespective of their remission status at HSCT (Figure 5). The similar analysis was performed in ALL and lymphoma patient groups (supplemental Figure 1, available on the Blood Web site; see the Supplemental Materials link at the top of the online article). Although each group contained only a limited number of patients, statistically significant prolonged OS and RFS were also seen in patients who did not achieve CR at HSCT in both the ALL and the lymphoma patient groups.

Thus, the appearance of hematogones might mark favorable OS and RFS regardless of their primary malignancy.

**Expansion of hematogones is frequently observed in patients who did not develop infection or severe acute GVHD**


In this study, all 43  $HG^+$  patients are currently alive, although primary diseases have relapsed in 3 patients. In contrast, 32 of 65  $HG^-$  patients have died. The causes of death in these 32  $HG^-$  patients are shown in Table 3. Twenty-six patients died of their refractory primary disease, and 6 patients died of TRM, including

**Table 2. Time required for engraftment in patients grouped by their primary disease or complication of infection or acute GVHD**

Disease	Remission status	Time required for engraftment, d					
		BMT			CBT		
		No.	Mean, d	P	No.	Mean, d	P
AML and advanced MDS	CR/non-CR	14/21	27.0/24.4	.11	11/14	31.5/31.8	.37
ALL	CR/non-CR	3/2	23.7/29.5	.35	3/10	34.3/32.1	.45
Lymphoma	CR/non-CR	7/12	26.1/22.8	.10	3/8	33.0/29.3	.44
			Overall	.10		Overall	.53
<b>After HSCT</b>							
Infections	Yes/No	38/21	25.3/24.8	.66	22/27	30.8/32.2	.61
Acute GVHD	Grade II-IV/Grade 0-I	26/33	24.7/25.3	.25	14/35	32.3/31.3	.47

ALL indicates acute lymphoblastic leukemia; AML, acute myelogenous leukemia; BMT, bone marrow transplantation; CBT, cord blood transplantation; CR, complete remission; GVHD, graft-versus-host disease; HSCT, hematopoietic stem cell transplantation; and MDS, myelodysplastic syndrome.





\*:  $p < 0.05$  \*\*:  $p < 0.01$  \*\*\*:  $p < 0.001$

**Figure 3. Patients whose hematogones comprised > 5% bone marrow MNCs constitute a group with significantly improved survival, irrespective of HSC sources.** (A-B) The Kaplan-Meier estimates of (A) OS and (B) RFS among patient subgroups with  $\leq 1\%$  (gray line),  $1\%-2\%$  (green line),  $2\%-3\%$  (orange line),  $3\%-4\%$  (red line),  $4\%-5\%$  (purple line),  $\leq 5\%$  (black bold line), or  $> 5\%$  (blue bold line) hematogones in the bone marrow MNCs. Forty-three patients who developed  $> 5\%$  MNC hematogones ( $HG^+$ ) showed significantly better 3-year OS and RFS, compared with any of each group ( $P < .01$  and  $P < .05$ , respectively), as well as to 65 patients with  $\leq 5\%$  MNCs hematogones ( $HG^-$ ;  $P < .001$  for both). (C-D) The Kaplan-Meier estimates of (C) OS and (D) RFS in  $HG^+$  and  $HG^-$  groups that received transplants with BMT or CBT. The improved OS and RFS were seen in  $HG^+$  groups regardless of the source of HSC.  $HG^+$  indicates patients developed hematogones ( $> 5\%$  of bone marrow MNCs);  $HG^-$  indicates patients who failed to develop hematogones ( $\leq 5\%$  of bone marrow MNCs).

acute GVHD (2 patients) and viral infections (4 patients). Of the 26 patients who died of primary disease, 24 developed both acute GVHD and infections.

We analyzed the relationship between the emergence of documented hematogones in the bone marrow, and variables including sex of donor/recipient, days required for engraftment, primary diseases, times of intensive chemotherapy before HSCT, remission status, conditioning regimen, documented infectious disease, and episode of acute/chronic GVHD by using univariate and multivariate analysis. These analyses were performed in patient groups treated with BMT and CBT, respectively.

There were no correlations found between the emergence of  $> 5\%$  of hematogones and clinical factors such as the day of engraftment, primary disease, times of intensive chemotherapies, and remission status, in either univariate or multivariate analyses. As shown in Table 4, in univariate analysis, a hematogone increase up to  $> 5\%$  of MNCs was found more frequently in patients without viral infection (such as cytomegalovirus, human herpesvirus 6, and adenovirus; BMT:  $P = .03$ ; CBT:  $P < .01$ ), and those did not develop severe acute GVHD of grade II-IV (BMT:  $P < .01$ ; CBT:  $P < .01$ ). Time required for engraftment did not differ between patient groups with or without infections, or acute GVHD (Table 2). These data appear to be compatible with the analysis of causes of death in  $HG^-$  patients (Table 3). On the other hand, in multivariate analysis, severe acute GVHD of grade II-IV, but not infection was the significant risk factor for emergence of hematogones (BMT:  $P = .03$ ; CBT:  $P = .04$ ; Table 4). Based on


these analyses, the emergence of hematogones heralds less frequent development of severe acute GVHD.

## Discussion

Hematogones are immature B-cell precursors that reside mainly in the bone marrow of every normal individual,<sup>1,2,27,40</sup> and their numbers could reflect activity of normal B lymphopoiesis. Hematogones are occasionally seen in large numbers in healthy people, especially in infants and young children.<sup>2,4,7,8</sup> Interestingly, recent reports have suggested that the presence of detectable numbers of hematogones at the recovery phase from myelosuppression reflects better prognosis of patients with AML treated with chemotherapy<sup>5</sup> or CBT,<sup>6</sup> although the underlying mechanism of this phenomenon is unclear. The increase of hematogones may reflect eradication of leukemic cells that could inhibit normal hematopoiesis,<sup>1,5</sup> or rapid immune reconstitution that could suppress infection and severe acute GVHD in an allogeneic HSCT setting.<sup>6</sup>

In these reports, the presence of hematogones was documented when they were detectable at a low frequency:  $\geq 0.01\%$  of TNCs at a recovery phase<sup>5</sup> or  $> 0\%$  and  $> 0.9\%$  of MNCs on day 21 and 100, respectively.<sup>6</sup> In contrast, the patient cohort in our study received allogeneic HSCT, and the majority (106 of 108 cases) of patients had  $> 0.1\%$  of hematogones at engraftment by our multicolor flow cytometric analysis (Figure 1C). Therefore, it was critical to set an appropriate threshold value and timing of sampling





**Figure 4. Patients who developed > 5% hematogones as a fraction of their MNCs constitute a group with significantly improved survival, irrespective of their primary disease.** The Kaplan-Meier estimates of OS and RFS in HG<sup>+</sup> and HG<sup>-</sup> patients differentiated with their primary disease. In each group of patients with (A) AML or advanced MDS, (B) ALL, and (C) lymphoma, HG<sup>+</sup> groups showed significantly better OS and RFS, compared with the HG<sup>-</sup> group ( $P < .001$  for both).


to decide a clinically meaningful increase of hematogones in an allogeneic HSCT setting. Furthermore, previous studies were performed only in patients with AML,<sup>5,6</sup> but not in patients with lymphoid neoplasms, presumably because it was difficult to discriminate a small number of neoplastic lymphoid cells from hematogones.<sup>6</sup>

To accurately enumerate hematogones in patients with various clinical backgrounds and with different donor cell sources, we performed the analysis on the day when patients met the clinical criteria of engraftment<sup>9-12</sup> and displayed complete donor-type chimerism. The confirmation for donor-type chimerism allowed us to avoid miscounting neoplastic lymphoid cells as hematogones. Because these samples should be free from host-derived cells, we included patients with lymphoid malignancies in our study. We rigorously measured the frequencies of hematogones within bone marrow MNCs by 6-color flow cytometric analysis.

In our study, donor-derived hematogones were polyclonal, based on *IGH* rearrangement analysis in all cases, and therefore the presence of hematogones should be a snapshot of normal B lymphopoiesis at the recovery phase. In fact, the frequencies of hemato-

gones at engraftment were correlated with circulating B-cell numbers at least until day 90 (Figure 2). Importantly, we here show that the frequencies of hematogones were correlated with donors' age, but not with recipients' age, suggesting the age-dependent decline of B-cell potential of donor HSCs. This is compatible with previous mouse studies in which younger HSCs are capable of producing more abundant B cells.<sup>41-43</sup>

According to our criteria, the engraftment was seen on days 25 and 32 (median) in BMT and CBT groups (Figure 1D), respectively, consistent with previous studies.<sup>35-39,44</sup> Within each BMT or CBT group, the engraftment day was not significantly altered by the patients' primary disease or remission status at transplantation (Table 2). The bone marrow sampling for hematogone analysis was performed on the day of engraftment. As shown in Table 3, the timing of sampling (= the day of engraftment) was not significantly related to emergence of hematogones in univariate and multivariate analyses. Interestingly, however, our data suggest that when hematogones reach > 5% of MNCs at engraftment, it has a profound clinical impact on patients' OS and RFS (Figure 3A-B). It is of note that the emergence of documented hematogones



**Figure 5. The presence of hematogones marks a group with good prognosis in AML/advanced MDS patients.** The Kaplan-Meier estimates of OS and RFS in HG<sup>+</sup> and HG<sup>-</sup> patients in AML or advanced MDS differentiated with their remission status before HSCT. Significantly better OS and RFS were seen in HG<sup>+</sup> groups irrespective of their remission status.

**Table 3. Cause of death in patients with less than 5% hematogones**

Primary disease	Cause of death, no. (%)					
	Relapse of primary disease			Transplantation-related mortality		
	Infection	Acute GVHD (Grade II-IV)	Total	Infection	Acute GVHD (Grade II-IV)	Total
AML and advanced MDS	14/33 (42.4)	14/33 (42.4)	15/33 (45.4)	2/33 (6.1)	0/33 (0)	2/33 (6.1)
ALL	3/9 (33.3)	3/9 (33.3)	3/9 (33.3)	1/9 (11.1)	1/9 (11.1)	2/9 (22.2)
Lymphoma	7/23 (30.4)	7/23 (30.4)	8/23 (34.8)	1/23 (4.3)	1/23 (4.3)	2/23 (8.7)
Total	24/65 (36.9)	24/65 (36.9)	26/65 (40)	4/65 (6.2)	2/65 (3.1)	6/65 (9.2)

ALL indicates acute lymphoblastic leukemia; AML, acute myelogenous leukemia; GVHD, graft-versus-host disease; and MDS, myelodysplastic syndrome.

was not related to times of intensive chemotherapies or remission status of patients before transplantation in both univariate and multivariate analyses (Table 3). This result suggests that the emergence of hematogones was not affected by the potential damage of host microenvironment through multiple chemotherapies.

The improvement of OS and RFS in HG<sup>+</sup> patients was seen in all patient groups: those suffering from AML/advanced MDS, ALL, or lymphoma (Figure 4). Furthermore, this effect became more evident when patients who had failed to achieve CR before transplantation were analyzed (Figure 5). In this case, the appearance of hematogones clearly marks a subgroup with favorable OS and RFS, irrespective of their primary diseases. In contrast, in patients who had achieved CR before transplantation, prolonged OS and RFS were found only in patients with AML/advanced MDS, but not in patients with ALL or lymphoma (Figure 5). A larger study including higher number of patients should be performed to clarify the impact of hematogones on HSCT results in CR patients with lymphoid malignancies.

The analyses of risk factors for the appearance of > 5% MNCs of hematogones revealed that in both BMT and CBT patients, the less frequent occurrences of severe acute GVHD and infections were significantly correlated in univariate analyses, whereas the less frequent severe acute GVHD was the only risk factor in multivariate analyses (Table 3). As shown in Table 4, all 32 deaths occurred only in HG<sup>-</sup> patients, and 24 of these 32 patients developed both severe acute GVHD and infection before the relapse of the disease. In these patients, doses of immunosuppressive drugs were escalated to control acute GVHD, which might cause development of infections as well as recurrence of primary disease.<sup>6,45,46</sup> It is therefore possible that less frequent development of severe acute GVHD in HG<sup>+</sup> patients is one of the reasons for their better OS and RFS.

The rapid reconstitution of the immune system represented by a high number of hematogones should be able to prevent infection.<sup>6,45</sup> In turn, successful prevention of acute GVHD could result in proliferation of hematogones because acute GVHD itself may suppress hematopoietic recovery by targeting the bone marrow HSC niche<sup>47</sup> or by attacking directly B-lymphoid cells.<sup>46</sup> In addition, the fact that improvement of RFS is associated with the expansion of hematogones suggests an interesting possibility that

B cells play a role in the graft-versus-leukemia effect,<sup>48</sup> although this is still controversial.<sup>41</sup> Also in turn, it is possible that the successful eradication of neoplastic cells from the bone marrow by HSCT results simply in rapid expansion of hematogones.

Thus, our data suggest that the expansion of hematogones is a useful indicator to discriminate a patient group with improved OS and RFS after allogeneic HSCT. Based on rigorous evaluation of frequencies of hematogones after HSCT, we propose that 5% of MNCs is a threshold value for a clinically valuable increase of hematogones. The prognostic value of this definition should be tested by future studies in larger groups of patients.

### Acknowledgments

The authors thank the medical and nursing staff working on the Fukuoka Blood and Marrow Transplantation Group for providing patient information, and D. Dalma-Weiszhausz for critically reviewing the manuscript.

This work was supported, in part, by a Grant-in-Aid from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (K.A., T.M.).

### Authorship

Contribution: T.S. and T.M. coordinated the project, designed and performed the transplantation and experiments, analyzed the data, and wrote the manuscript; Y.K., Y.M., K. Kamezaki, K. Takenaka, H.I., K.N., T.T, and K. Kato performed the transplantation and provided technical advice; K. Takase, H.H., A.N., Y.I., T.K., and T.E. provided patient information, clinical samples, and technical advice; and K.A. designed the experiments, reviewed the data, and edited the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Koichi Akashi, Department of Medicine and Biosystemic Sciences, Kyushu University Graduate School of Medicine, 3-1-1 Maidashi, Higashi-ku, Fukuoka 812-0054, Japan; e-mail: akashi@med.kyushu-u.ac.jp.

**Table 4. Risk factors for development of more than 5% MNCs of hematogone based on univariate and multivariate analyses**

	BMT				CBT			
	Univariate		Multivariate		Univariate		Multivariate	
	Odds ratio (95% CI)	P	Odds ratio (95% CI)	P	Odds ratio (95% CI)	P	Odds ratio (95% CI)	P
Infections, Yes/No	0.19 (0.05-0.74)	.03	0.42 (0.02-1.58)	.23	0.16 (0.04-0.57)	< .01	0.37 (0.02-2.81)	.10
Acute GVHD, Grade II-IV/0-I	0.09 (0.01-0.73)	< .01	0.04 (0.00-0.85)	.03	0.12 (0.03-0.48)	< .01	0.04 (0.00-0.80)	.04

BMT indicates bone marrow transplantation; CBT, cord blood transplantation; CI, confidence interval; GVHD, graft-versus-host disease; and MNC, mononuclear cell.

## References

- McKenna RW, Washington LT, Aquino DB, Picker LJ, Kroft SH. Immunophenotypic analysis of hematogones (B-lymphocyte precursors) in 662 consecutive bone marrow specimens by 4-color flow cytometry. *Blood*. 2001;98(8):2498-2507.
- Sevilla DW, Colovai AI, Emmons FN, Bhagat G, Alobeid B. Hematogones: a review and update. *Leuk Lymphoma*. 2010;51(1):10-19.
- Muehleck SD, McKenna RW, Gale PF, Brunning RD. Terminal deoxynucleotidyl transferase (TdT)-positive cells in bone marrow in the absence of hematologic malignancy. *Am J Clin Pathol*. 1983;79(3):277-284.
- Davis RE, Longacre TA, Cornbleet PJ. Hematogones in the bone marrow of adults. Immunophenotypic features, clinical settings, and differential diagnosis. *Am J Clin Pathol*. 1994;102(2):202-211.
- Chantepie SP, Salaun V, Parienti JJ, et al. Hematogones: a new prognostic factor for acute myeloblastic leukemia. *Blood*. 2011;117(4):1315-1318.
- Honebrink T, Dayton V, Burke MJ, et al. Impact of bone marrow hematogones on umbilical cord blood transplantation outcomes in patients with acute myeloid leukemia. *Biol Blood Marrow Transplant*. 2012;18(6):930-936.
- Brady KA, Atwater SK, Lowell CA. Flow cytometric detection of CD10 (cALLA) on peripheral blood B lymphocytes of neonates. *Br J Haematol*. 1999;107(4):712-715.
- Longacre TA, Foucar K, Crago S, et al. Hematogones: a multiparameter analysis of bone marrow precursor cells. *Blood*. 1989;73(2):543-552.
- Dreger P, Kloss M, Petersen B, et al. Autologous progenitor cell transplantation: prior exposure to stem cell-toxic drugs determines yield and engraftment of peripheral blood progenitor cell but not of bone marrow grafts. *Blood*. 1995;86(10):3970-3978.
- Davies SM, Kollman C, Anasetti C, et al. Engraftment and survival after unrelated-donor bone marrow transplantation: a report from the national marrow donor program. *Blood*. 2000;96(13):4096-4102.
- Bensinger WI, Martin PJ, Storer B, et al. Transplantation of bone marrow as compared with peripheral-blood cells from HLA-identical relatives in patients with hematologic cancers. *N Engl J Med*. 2001;344(3):175-181.
- Laughlin MJ, Barker J, Bambach B, et al. Hematopoietic engraftment and survival in adult recipients of umbilical-cord blood from unrelated donors. *N Engl J Med*. 2001;344(24):1815-1822.
- Dohner H, Estey EH, Amadori S, et al. Diagnosis and management of acute myeloid leukemia in adults: recommendations from an international expert panel, on behalf of the European LeukemiaNet. *Blood*. 2010;115(3):453-474.
- Greenberg P, Cox C, LeBeau MM, et al. International scoring system for evaluating prognosis in myelodysplastic syndromes. *Blood*. 1997;89(6):2079-2088.
- Appelbaum FR, Anderson J. Allogeneic bone marrow transplantation for myelodysplastic syndrome: outcomes analysis according to IPSS score. *Leukemia*. 1998;12(Suppl 1):S25-S29.
- Goldman JM, Schmitz N, Niethammer D, Gratwohl A. Allogeneic and autologous transplantation for haematological diseases, solid tumours and immune disorders: current practice in Europe in 1998. Accreditation Sub-Committee of the European Group for Blood and Marrow Transplantation. *Bone Marrow Transplant*. 1998;21(1):1-7.
- Perea G, Lasa A, Aventin A, et al. Prognostic value of minimal residual disease (MRD) in acute myeloid leukemia (AML) with favorable cytogenetics [t(8;21) and inv(16)]. *Leukemia*. 2006;20(1):87-94.
- Cheson BD, Pfistner B, Juweid ME, et al. Revised response criteria for malignant lymphoma. *J Clin Oncol*. 2007;25(5):579-586.
- Uchida N, Wake A, Takagi S, et al. Umbilical cord blood transplantation after reduced-intensity conditioning for elderly patients with hematologic diseases. *Biol Blood Marrow Transplant*. 2008;14(5):583-590.
- Kim SW, Matsuo K, Fukuda T, et al. Reduced-intensity unrelated donor bone marrow transplantation for hematologic malignancies. *Int J Hematol*. 2008;88(3):324-330.
- Oda S, Oki E, Maehara Y, Sugimachi K. Precise assessment of microsatellite instability using high resolution fluorescent microsatellite analysis. *Nucleic Acids Res*. 1997;25(17):3415-3420.
- Kikushige Y, Yoshimoto G, Miyamoto T, et al. Human Flt3 is expressed at the hematopoietic stem cell and the granulocyte/macrophage progenitor stages to maintain cell survival. *J Immunol*. 2008;180(11):7358-7367.
- Mori Y, Iwasaki H, Kohno K, et al. Identification of the human eosinophil lineage-committed progenitor: revision of phenotypic definition of the human common myeloid progenitor. *J Exp Med*. 2009;206(1):183-193.
- Imamura R, Miyamoto T, Yoshimoto G, et al. Mobilization of human lymphoid progenitors after treatment with granulocyte colony-stimulating factor. *J Immunol*. 2005;175(4):2647-2654.
- Yoshimoto G, Miyamoto T, Jabbarzadeh-Tabrizi S, et al. FLT3-ITD up-regulates MCL-1 to promote survival of stem cells in acute myeloid leukemia via FLT3-ITD-specific STAT5 activation. *Blood*. 2009;114(24):5034-5043.
- Galy A, Travis M, Cen D, Chen B, Human T, B, natural killer, and dendritic cells arise from a common bone marrow progenitor cell subset. *Immunity*. 1995;3(4):459-473.
- LeBien TW. Fates of human B-cell precursors. *Blood*. 2000;96(1):9-23.
- LeBien TW, Tedder TF. B lymphocytes: how they develop and function. *Blood*. 2008;112(5):1570-1580.
- Yoshimoto G, Nagafuji K, Miyamoto T, et al. FLT3 mutations in normal karyotype acute myeloid leukemia in first complete remission treated with autologous peripheral blood stem cell transplantation. *Bone Marrow Transplant*. 2005;36(11):977-983.
- Kikushige Y, Ishikawa F, Miyamoto T, et al. Self-renewing hematopoietic stem cell is the primary target in pathogenesis of human chronic lymphocytic leukemia. *Cancer Cell*. 2011;20(2):246-259.
- Maldonado G, Greenland S. Simulation study of confounder-selection strategies. *Am J Epidemiol*. 1993;138(11):923-936.
- Baron F, Sandmaier BM. Chimerism and outcomes after allogeneic hematopoietic cell transplantation following nonmyeloablative conditioning. *Leukemia*. 2006;20(10):1690-1700.
- Matsuda K, Yamauchi K, Tozuka M, et al. Monitoring of hematopoietic chimerism by short tandem repeats, and the effect of CD selection on its sensitivity. *Clin Chem*. 2004;50(12):2411-2414.
- Boeck S, Hamann M, Pihusch V, et al. Kinetics of dendritic cell chimerism and T cell chimerism in allogeneic hematopoietic stem cell recipients. *Bone Marrow Transplant*. 2006;37(1):57-64.
- Laughlin MJ, Eapen M, Rubinstein P, et al. Outcomes after transplantation of cord blood or bone marrow from unrelated donors in adults with leukemia. *N Engl J Med*. 2004;351(22):2265-2275.
- Rocha V, Labopin M, Sanz G, et al. Transplants of umbilical-cord blood or bone marrow from unrelated donors in adults with acute leukemia. *N Engl J Med*. 2004;351(22):2276-2285.
- Takahashi S, Iseki T, Ooi J, et al. Single-institute comparative analysis of unrelated bone marrow transplantation and cord blood transplantation for adult patients with hematologic malignancies. *Blood*. 2004;104(12):3813-3820.
- Couban S, Simpson DR, Barnett MJ, et al. A randomized multicenter comparison of bone marrow and peripheral blood in recipients of matched sibling allogeneic transplants for myeloid malignancies. *Blood*. 2002;100(5):1525-1531.
- Vigorito AC, Azevedo WM, Marques JF, et al. A randomized, prospective comparison of allogeneic bone marrow and peripheral blood progenitor cell transplantation in the treatment of hematological malignancies. *Bone Marrow Transplant*. 1998;22(12):1145-1151.
- Nunez C, Nishimoto N, Gartland GL, et al. B cells are generated throughout life in humans. *J Immunol*. 1996;156(2):866-872.
- Rossi DJ, Bryder D, Zahn JM, et al. Cell intrinsic alterations underlie hematopoietic stem cell aging. *Proc Natl Acad Sci U S A*. 2005;102(26):9194-9199.
- Frasca D, Landin AM, Lechner SC, et al. Aging down-regulates the transcription factor E2A, activation-induced cytidine deaminase, and Ig class switch in human B cells. *J Immunol*. 2008;180(8):5283-5290.
- Allman D, Miller JP. The aging of early B-cell precursors. *Immunol Rev*. 2005;205:18-29.
- Stem Cell Trialists' Collaborative Group. Allogeneic peripheral blood stem-cell compared with bone marrow transplantation in the management of hematologic malignancies: an individual patient data meta-analysis of nine randomized trials. *J Clin Oncol*. 2005;23(22):5074-5087.
- Shimabukuro-Vornhagen A, Hallek MJ, Storb RF, von Bergwelt-Baildon MS. The role of B cells in the pathogenesis of graft-versus-host disease. *Blood*. 2009;114(24):4919-4927.
- Baker MB, Riley RL, Podack ER, Levy RB. Graft-versus-host-disease-associated lymphoid hypoplasia and B cell dysfunction is dependent upon donor T cell-mediated Fas-ligand function, but not perforin function. *Proc Natl Acad Sci U S A*. 1997;94(4):1366-1371.
- Shono Y, Ueha S, Wang Y, et al. Bone marrow graft-versus-host disease: early destruction of hematopoietic niche after MHC-mismatched hematopoietic stem cell transplantation. *Blood*. 2010;115(26):5401-5411.
- Miklos DB, Kim HT, Miller KH, et al. Antibody responses to H-Y minor histocompatibility antigens correlate with chronic graft-versus-host disease and disease remission. *Blood*. 2005;105(7):2973-2978.

# Contribution of Bone Marrow-Derived Hematopoietic Stem/Progenitor Cells to the Generation of Donor-Marker<sup>+</sup> Cardiomyocytes *In Vivo*

Mitsuhiro Fukata<sup>1\*</sup>, Fumihiko Ishikawa<sup>1,2\*</sup>, Yuho Najima<sup>2</sup>, Takuji Yamauchi<sup>1</sup>, Yoriko Saito<sup>2</sup>, Katsuto Takenaka<sup>1</sup>, Kohta Miyawaki<sup>1</sup>, Hideki Shimazu<sup>1</sup>, Kazuya Shimoda<sup>3</sup>, Takaaki Kanemaru<sup>4</sup>, Kei-ichiro Nakamura<sup>5</sup>, Keita Odashiro<sup>1</sup>, Koji Nagafuji<sup>6</sup>, Mine Harada<sup>1</sup>, Koichi Akashi<sup>1,7</sup>

**1** Department of Medicine and Biosystemic Science, Kyushu University Graduate School of Medical Science, Fukuoka, Japan, **2** Laboratory for Human Disease Models, RIKEN Research Center for Allergy and Immunology, Yokohama, Japan, **3** Department of Gastroenterology and Hematology, Faculty of Medicine, Miyazaki University, Miyazaki, Japan, **4** Morphology Core, Kyushu University, Fukuoka, Japan, **5** Second Department of Anatomy, Kurume University School of Medicine, Kurume, Japan, **6** Division of Hematology and Oncology, Department of Medicine, Kurume University School of Medicine, Kurume, Japan, **7** Center for Cellular and Molecular Medicine, Kyushu University Hospital, Fukuoka, Japan

## Abstract

**Background:** Definite identification of the cell types and the mechanism relevant to cardiomyogenesis is essential for effective cardiac regenerative medicine. We aimed to identify the cell populations that can generate cardiomyocytes and to clarify whether generation of donor-marker<sup>+</sup> cardiomyocytes requires cell fusion between BM-derived cells and recipient cardiomyocytes.

**Methodology/Principal Findings:** Purified BM stem/progenitor cells from green fluorescence protein (GFP) mice were transplanted into C57BL/6 mice or cyan fluorescence protein (CFP)-transgenic mice. Purified human hematopoietic stem cells (HSCs) from cord blood were transplanted into immune-compromised NOD/SCID/IL2r<sup>γ</sup><sup>null</sup> mice. GFP<sup>+</sup> cells in the cardiac tissue were analyzed for the antigenicity of a cardiomyocyte by confocal microscopy following immunofluorescence staining. GFP<sup>+</sup> donor-derived cells, GFP<sup>+</sup>CFP<sup>+</sup> fused cells, and CFP<sup>+</sup> recipient-derived cells were distinguished by linear unmixing analysis. Hearts of xenogeneic recipients were evaluated for the expression of human cardiomyocyte genes by real-time quantitative polymerase chain reaction. In C57BL/6 recipients, Lin<sup>-low</sup>CD45<sup>+</sup> hematopoietic cells generated greater number of GFP<sup>+</sup> cardiomyocytes than Lin<sup>-low</sup>CD45<sup>-</sup> mesenchymal cells (37.0±/−23.9 vs 0.00±/−0.00 GFP<sup>+</sup> cardiomyocytes per a recipient, P=0.0095). The number of transplanted purified HSCs (Lin<sup>-low</sup>Sca-1<sup>+</sup> or Lin<sup>-</sup>Sca-1<sup>+</sup>c-Kit<sup>+</sup> or CD34<sup>-</sup>Lin<sup>-</sup>Sca-1<sup>+</sup>c-Kit<sup>+</sup>) showed correlation to the number of GFP<sup>+</sup> cardiomyocytes (P<0.05 in each cell fraction), and the incidence of GFP<sup>+</sup> cardiomyocytes per injected cell dose was greatest in CD34<sup>-</sup>Lin<sup>-</sup>Sca-1<sup>+</sup>c-Kit<sup>+</sup> recipients. Of the hematopoietic progenitors, total myeloid progenitors generated greater number of GFP<sup>+</sup> cardiomyocytes than common lymphoid progenitors (12.8±/−10.7 vs 0.67±/−1.00 GFP<sup>+</sup> cardiomyocytes per a recipient, P=0.0021). In CFP recipients, all GFP<sup>+</sup> cardiomyocytes examined coexpressed CFP. Human troponin C and myosin heavy chain 6 transcripts were detected in the cardiac tissue of some of the xenogeneic recipients.

**Conclusions/Significance:** Our results indicate that HSCs resulted in the generation of cardiomyocytes via myeloid intermediates by fusion-dependent mechanism. The use of myeloid derivatives as donor cells could potentially allow more effective cell-based therapy for cardiac repair.

**Citation:** Fukata M, Ishikawa F, Najima Y, Yamauchi T, Saito Y, et al. (2013) Contribution of Bone Marrow-Derived Hematopoietic Stem/Progenitor Cells to the Generation of Donor-Marker<sup>+</sup> Cardiomyocytes *In Vivo*. PLoS ONE 8(5): e62506. doi:10.1371/journal.pone.0062506

**Editor:** Maurizio Pesce, Centro Cardiologico Monzino, Italy

**Received:** July 14, 2011; **Accepted:** March 26, 2013; **Published:** May 7, 2013

**Copyright:** © 2013 Fukata et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** The authors have no support or funding to report.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: mfukata-circ@umin.net (MF); f\_ishika@rcai.riken.jp (FI)

## Introduction

Modification of regenerative capacity in injured heart could be potentially alternative to conventional therapy for treating patients suffering from heart failure [1–7]. Based on the promising results in rodents [3,4], clinical trials of cellular therapy using bone marrow (BM) cells for ischemic heart disease patients have been designed. In many of clinical trials for improving the function of cardiac recovery, some favorable results were obtained following

injection of BM mononuclear cells (MNCs) [2,5–7]. However, careful examination needs to be performed in basic research because cell fate and the effects of transplanted cells are not fully unveiled [8].

BM contains heterogeneous cell populations including at least two distinct stem cells, hematopoietic stem cells (HSCs) and mesenchymal stem cells (MSCs) [9], and various progenitors of myeloid and lymphoid lineages. Both HSCs and MSCs have been reported to acquire the phenotype of cardiomyocytes in syngeneic