

Exacerbations of Chronic Heart Failure (OPTIMIZE-HF) study showed that the BNP level at discharge was the best predictor of 1-year mortality and/or re-hospitalization among older patients (≥ 65 years) hospitalized with HF.¹⁴ Increase in BNP is useful for predicting cardiac events in stable outpatients with non-ischemic chronic HF.¹⁵ Logeart et al found that when HF patients have a 70% prevalence of systolic dysfunction of LVEF ($< 45\%$), high BNP at pre-discharge assessment is a strong, independent marker of death or re-admission after decompensated congestive HF,¹⁶ which is also consonant with the present result that the prevalence of DCM with systolic dysfunction was 76.9%. In addition, Javaheri et al reported that low DBP was associated with worse survival in patients with HF and a decreased ejection fraction.¹⁷ Moreover, the Framingham study showed that DBP was negatively correlated with coronary heart disease and mortality in subjects > 60 years old,¹⁸ while DBP ≤ 60 mmHg was associated with higher CV mortality in patients with systolic hypertension aged > 70 years.¹⁹ Because low DBP impairs coronary blood flow during diastole, it could cause myocardial ischemia and the progression of HF. The present study provided similar results to these findings mentioned here.

In contrast, the OPTIMIZE-HF trial showed that a lower SBP at admission was 1 of the predictors for the combined endpoint of death or re-hospitalization within 60–90 days after discharge. DBP, however, was not included among the candidate variables.²⁰ The Canada community-based study indicated that a lower SBP was a predictor of mortality among elderly patients with HF.²¹ The reasons why we found DBP to be a predictor of re-hospitalization in the present analysis are as follows: (1) DBP was not assessed in some other studies; (2) the present patient cohort (aged 67.2 ± 15.2 years) was younger than the Canada HF patient cohort (aged 75.3 ± 11.8 years) and the OPTIMIZE-HF cohort (aged 73.1 ± 14.2 years); and (3) the mean SBP of the present patients at discharge (112.2 ± 16.3 mmHg) was lower than that of the Canada HF cohort (148 ± 34 mmHg) or the OPTIMIZE-HF cohort (124.3 ± 24 mmHg). The present finding that a low DBP may increase the re-hospitalization rate indicates that DBP can be an important predictor of re-hospitalization provided that SBP is not high. Although the reason for the association of DBP with re-hospitalization for HF still needs to be elucidated, careful BP monitoring is necessary in patients with HF.

We found that BUN, DBP and plasma BNP are important predictors of re-hospitalized HF, and further consideration of what this result means, is needed. First, the prevalence (76.9%) of DCM in the present study was greater than in the JCARE-CARD study (24.0%), suggesting that the degree of HF in the present study was higher than average. Severe HF causes renal dysfunction, which may increase BUN, suggesting that high BUN is linked to the severity of HF and, as suggested by cardio-renal syndrome, the correction of BUN may become a target for HF treatment using ACEI or ARB.^{22–25} In the BEST study, the use of high-dose loop diuretics (total daily dose ≥ 160 mg furosemide equivalents) was associated with significantly increased mortality in patients with BUN > 21.0 mg/dl.⁷ Patients with a more severe degree of HF generally received larger doses of loop diuretics and were required to evaluate volume management strategies. This is in agreement with the present study (Table 1). In addition, the measurement of plasma BNP has been useful to determine the severity of HF because BNP is secreted in response to ventricular wall stress.²⁶ Indeed, β -blockers are known to affect plasma BNP, and Matsumura et al indicated that low-dose regimens of ACEI/ARB or β -blockers have favorable effects on the prognosis of

Japanese patients with DCM.²⁷ The decreasing of plasma BNP would be an excellent strategy in the treatment of HF patients. In contrast, lower DBP may be the result of altered peripheral arterial capacitance or resistance and low cardiac output due to HF. Low DBP can be a surrogate marker of HF severity but does not appear to be an appropriate target or guide for HF treatment.

Study Limitations

The main limitations of this study were its retrospective design and observational nature. As is the case for all retrospective analyses, it is possible that unrecognized or recognized confounders influenced the results despite adjustment for various factors. Various factors that can alter BUN level, such as medication dosage, cachexia, a high protein diet, and muscle wasting, were not assessed. Thus, prospective studies are needed to establish the predictors of re-hospitalization and CV death such as BUN or DBP. We used serum Cr as a surrogate marker of renal function. The Cockcroft-Gault and Modification of Diet in Renal Disease (MDRD) formulae for eGFR incorporate age and gender to account for differences in muscle mass. Because these factors were included in the present regression model, the predictors might have cancelled each other out, so we used the serum Cr level as was done in the OPTIMIZE-HF trial. Also, Smilde et al compared GFR estimated using different formulae with measured GFR in patients with systolic HF and found that all formulae had inaccuracies,²⁸ therefore we used serum Cr instead of eGFR.

Conclusions

High serum BUN (≥ 22.5 mg/dl), high plasma BNP (≥ 250 pg/ml), and low DBP (< 60 mmHg) predict CV events in patients hospitalized for ADHF. These risk factors may be useful to distinguish low-risk patients from those at high risk who may benefit from closer monitoring and more aggressive treatment.

Disclosures

Sources of funding: this work was supported by grants-in-aids from the Japanese Ministry of Health, Labor, and Welfare, and the Japanese Ministry of Education, Culture, Sports, Science and Technology, and by Grants from the Japan Heart Foundation, and the Japan Cardiovascular Research Foundation. Conflict of interest: none.

References

- Krumholz HM, Parent EM, Tu N, Vaccarino V, Wang Y, Radford MJ, et al. Readmission after hospitalization for congestive heart failure among medicare beneficiaries. *Arch Intern Med* 1997; **157**: 99–104.
- Goldberg RJ, Ciampa J, Lessard D, Meyer TE, Spencer FA. Long-term survival after heart failure: A contemporary population-based perspective. *Arch Intern Med* 2007; **167**: 490–496.
- Cuffe MS, Califf RM, Adams KF Jr, Benza R, Bourge RC, Colucci WS, et al. Short-term intravenous milrinone for acute exacerbation of chronic heart failure: A randomized controlled trial. *JAMA* 2002; **287**: 1541–1547.
- Binanay C, Califf RM, Hasselblad V, O'Connor CM, Shah MR, Sopko G, et al; ESCAPE Investigators and ESCAPE Study Coordinators. Evaluation study of congestive heart failure and pulmonary artery catheterization effectiveness: The ESCAPE trial. *JAMA* 2005; **294**: 1625–1633.
- Shiba N, Nochioka K, Miura M, Kohno H, Shimokawa H; CHART-2 Investigators. Trend of westernization of etiology and clinical characteristics of heart failure patients in Japan: First report from the CHART-2 study. *Circ J* 2011; **75**: 823–833.
- Tsuchihashi-Makaya M, Hamaguchi S, Kinugawa S, Yokota T, Goto D, Yokoshiki H, et al; JCARE-CARD Investigators. Characteristics and outcomes of hospitalized patients with heart failure and reduced vs preserved ejection fraction: Report from the Japanese Cardiac Registry of Heart Failure in Cardiology (JCARE-CARD). *Circ J* 2009; **73**: 1893–1900.

7. Testani JM, Cappola TP, Brensinger CM, Shannon RP, Kimmel SE. Interaction between loop diuretic-associated mortality and blood urea nitrogen concentration in chronic heart failure. *J Am Coll Cardiol* 2011; **58**: 375–382.
8. Fonarow GC, Adams KF Jr, Abraham WT, Yancy CW, Boscardin WJ, ADHERE Scientific Advisory Committee, Study Group, and Investigators. Risk stratification for in-hospital mortality in acutely decompensated heart failure: Classification and regression tree analysis. *JAMA* 2005; **293**: 572–580.
9. Giamouzis G, Kalogeropoulos AP, Georgiopoulos VV, Agha SA, Rashad MA, Laskar SR, et al. Incremental value of renal function in risk prediction with the Seattle Heart Failure Model. *Am Heart J* 2009; **157**: 299–305.
10. Cauthen CA, Lipinski MJ, Abbate A, Appleton D, Nusca A, Varma A, et al. Relation of blood urea nitrogen to long-term mortality in patients with heart failure. *Am J Cardiol* 2008; **101**: 1643–1647.
11. Sands JM. Mammalian urea transporters. *Ann Rev Physiol* 2003; **65**: 543–566.
12. Cohn JN, Levine TB, Olivari MT, Garberg V, Lura D, Francis GS, et al. Plasma norepinephrine as a guide to prognosis in patients with chronic congestive heart failure. *N Engl J Med* 1984; **311**: 819–823.
13. Cheng V, Kazanagra R, Garcia A, Lenert L, Krishnaswamy P, Gardetto N, et al. A rapid bedside test for B-type peptide predicts treatment outcomes in patients admitted for decompensated heart failure: A pilot study. *J Am Coll Cardiol* 2001; **37**: 386–391.
14. Kociol RD, Horton JR, Fonarow GC, Reyes EM, Shaw LK, O'Connor CM, et al. Admission, discharge, or change in B-Type natriuretic peptide and long-term outcomes: Data from Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients With Heart Failure (OPTIMIZE-HF) linked to Medicare claims. *Circ Heart Fail* 2011; **4**: 628–636.
15. Nishiyama K, Tsutomoto T, Yamaji M, Kawahara C, Fujii M, Yamamoto T, et al. Biological variation of brain natriuretic peptide and cardiac events in stable outpatients with nonischemic chronic heart failure. *Circ J* 2011; **75**: 341–347.
16. Logeart D, Thabut G, Jourdain P, Chavelas C, Beyne P, Beauvais F, et al. Predischarge B-type natriuretic peptide assay for identifying patients at high risk of re-admission after decompensated heart failure. *J Am Coll Cardiol* 2004; **43**: 635–641.
17. Javaheri S, Shukla R, Zeigler H, Wexler L. Central sleep apnea, right ventricular dysfunction, and low diastolic blood pressure are predictors of mortality in systolic heart failure. *J Am Coll Cardiol* 2007; **20**: 2028–2034.
18. Franklin S, Larson MG, Khan SA, Wong ND, Leip EP, Kannel WB, et al. Does the relation of blood pressure to coronary heart disease risk change with aging? The Framingham Heart Study. *Circulation* 2001; **103**: 1245–1249.
19. Protogerou AD, Safar ME, Iaria P, Safar H, Le Dudal K, Filipovsky J, et al. Diastolic blood pressure and mortality in the elderly with cardiovascular disease. *Hypertension* 2007; **50**: 172–180.
20. O'Connor CM, Abraham WT, Albert NM, Clare R, Gattis Stough W, Gheorghide M, et al. Predictors of mortality after discharge in patients hospitalized with heart failure: An analysis from the Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients with Heart Failure (OPTIMIZE-HF). *Am Heart J* 2008; **156**: 662–673.
21. Lee DS, Austin PC, Rouleau JL, Liu PP, Naimark D, Tu JV. Predicting mortality among patients hospitalized for heart failure: Derivation and validation of a clinical model. *JAMA* 2003; **290**: 2581–2587.
22. Maschio G, Alberti D, Janin G, Locatelli F, Mann JF, Motolese M, et al. Effect of the angiotensin-converting enzyme inhibitor benazepril on the progression of chronic renal insufficiency: The Angiotensin-Converting-Enzyme Inhibition in Progressive Renal Insufficiency Study Group. *N Engl J Med* 1996; **334**: 939–945.
23. The SOLVD Investigators. Effect of enalapril on survival in patients with reduced left ventricular ejection fractions and congestive heart failure. *N Engl J Med* 1991; **325**: 293–302.
24. Granger CB, McMurray JJ, Yusuf S, Held P, Michelson EL, Olofsson B, et al; CHARM Investigators and Committees. Effects of candesartan in patients with chronic heart failure and reduced left ventricular systolic function intolerant to angiotensin-converting-enzyme inhibitors: The CHARM-Alternative trial. *Lancet* 2003; **362**: 772–776.
25. Packer M, Coats AJ, Fowler MB, Katus HA, Krum H, Mohacsi P, et al; Carvedilol Prospective Randomized Cumulative Survival Study Group. Effect of carvedilol on survival in severe chronic heart failure. *N Engl J Med* 2001; **344**: 1651–1658.
26. Jourdain P, Jondeau G, Funck F, Gueffet P, Le Helloco A, Donal E, et al. Plasma brain natriuretic peptide-guided therapy to improve outcome in heart failure: The STARS-BNP Multicenter Study. *J Am Coll Cardiol* 2007; **49**: 1733–1739.
27. Matsumura Y, Takata J, Kitaoka H, Kubo T, Baba Y, Hoshikawa E, et al. Long-term prognosis of dilated cardiomyopathy revisited: An improvement in survival over the past 20 years. *Circ J* 2006; **70**: 376–383.
28. Smilde TD, van Veldhuisen DJ, Navis G, Voors AA, Hillege HL. Drawbacks and prognostic value of formulas estimating renal function in patients with chronic heart failure and systolic dysfunction. *Circulation* 2006; **114**: 1572–1580.

ORIGINAL ARTICLE

Derivation of a mathematical expression for predicting the time to cardiac events in patients with heart failure: a retrospective clinical study

Akemi Yoshida¹, Masanori Asakura¹, Hiroshi Asanuma², Akira Ishii¹, Takuya Hasegawa¹, Tetsuo Minamino³, Seiji Takashima⁴, Hideaki Kanzaki¹, Takashi Washio⁵ and Masafumi Kitakaze¹

The prognoses for patients with certain diseases are estimated by averaging the results of clinical trials. To investigate the possibility of deriving a mathematical formula for the estimation of prognosis, we formulated the equation $\tau = f(x_1, \dots, x_p)$, where x_1, \dots, x_p are clinical features and τ represents the clinical outcome for heart failure (HF). We attempted to determine the function to mathematically formulate the relationship between clinical features and outcomes for these patients. We followed 151 patients (mean age: 68.6 ± 14.6 years; men: 61.6%) who were consecutively hospitalized and discharged as a result of acute decompensated HF (ADHF) between May 2006 and December 2009. The mathematical analysis was performed through a probabilistic modeling of the relational data by assuming a Poisson process for rehospitalization owing to HF and by linearly approximating the relationship between the clinical factors and the mean elapsed time to rehospitalization. The former assumption was validated by a statistical test of the data, and the contribution of each parameter was assessed based on the coefficients of the linear relation. Using a regularization method to analyze 402 clinical parameters, we identified 252 factors that substantially influenced the elapsed time until rehospitalization. With the probability model based on the Poisson process, the actual (X ; 388 ± 377 days) and estimated (Y ; 398 ± 381 days) elapsed times to rehospitalization were tightly correlated ($Y = 1.0076X + 6.5531$, $R^2 = 0.9879$, $P < 0.0001$). We established a mathematical formula that closely predicts the clinical outcomes of patients who are hospitalized with ADHF and discharged after appropriate treatment.

Hypertension Research (2013) 36, 450–456; doi:10.1038/hr.2012.200; published online 20 December 2012

Keywords: heart failure; mathematical model; prognosis; rehospitalization

INTRODUCTION

Studies show that numerous factors, including disease severity, treatment protocols and the environment, independently determine patients' prognoses. For example, in patients with chronic heart failure (CHF), many studies have shown that various independent indices of the severity of CHF, such as plasma B-type natriuretic peptide (BNP) level, left ventricular function, exercise tolerance or New York Heart Association (NYHA) functional class affect the time to hospitalization or cardiac death.^{1–5} However, because we could not identify the elapsed time until hospitalization in certain patients with CHF, we estimated this time using knowledge of the pathophysiology of CHF, our experience with previous comparable patients and Kaplan–Meier plots of their hospitalization in the clinical studies; we then explained our estimation to each patient. This procedure led us to conclude that estimating the elapsed time to rehospitalization is a type of problem that is specific to clinical medical science because the results and outcomes of biology or basic medical sciences can be

derived from mathematically formulated equations. Furthermore, other fields of basic science, such as physics and mathematics or applied sciences, such as mechanics, thermodynamics and fluid dynamics, are mathematically formulated; the observational phenomena in applied sciences other than medical science can be predicted by mathematical equations, for example, the law of universal gravitation.⁶ The most important issue in deriving a mathematical expression for relationships among two or more factors is the prediction of the future value of one variable based on the other factor(s). All phenomena, such as the severity of CHF and the patients' characteristics before the occurrence of clinical events, may therefore provide a mathematical equation for the clinical outcome if we can relate factors in the patient's clinical status to clinical outcomes such as rehospitalization.

To investigate this possibility, we sought to solve the equation $\tau = f(x_1, \dots, x_p)$, where x_1, \dots, x_p represent clinical features affecting the clinical outcome for CHF. We attempted to determine the

¹Department of Cardiovascular Medicine, National Cerebral and Cardiovascular Center, Suita, Japan; ²Cardiovascular Science and Technology, Kyoto Prefectural University of Medicine, Kyoto, Japan; ³Cardiovascular Medicine, Osaka University Graduate School of Medicine, Suita, Japan; ⁴Molecular Cardiology, Osaka University Graduate School of Medicine, Suita, Japan and ⁵The Institute of Scientific and Industrial Research, Osaka University, Suita, Japan
Correspondence: Dr M Kitakaze, Department of Cardiovascular Medicine, National Cerebral and Cardiovascular Center, 5-7-1 Fujishirodai, Suita, Osaka 565-8565, Japan.
E-mail: kitakaze@zf6.so-net.ne.jp

Received 12 September 2012; revised 7 October 2012; accepted 29 October 2012; published online 20 December 2012

function (f) to yield τ , the time to rehospitalization, from the clinical parameters (x_1, \dots, x_p) reflecting patient characteristics at the time of discharge.

METHODS

Ethics statement

This study was approved by National Cerebral and Cardiovascular Center Research Ethics Committee. The Committee decided that the acquisition of informed consent from the 151 subjects was not required according to the Japanese Clinical Research Guideline because this was a retrospective observational study. Instead, we made a public announcement in accordance with the request of the Ethics Committee and the Guideline.

Subjects and clinical parameters

A total of 486 patients with acute decompensated heart failure (ADHF) were admitted between May 2006 and December 2009. Because patients who were admitted for ADHF only once were excluded, the remaining 151 patients were included in this study. The oldest hospitalization was adopted regarding repeat patients during this study. The diagnosis of HF was confirmed by an expert team of cardiologists using the Framingham criteria.⁷ Careful history-taking, physical examinations, laboratory tests, chest X-rays, electrocardiograms, Doppler echocardiographic studies, coronary angiography and right heart catheterization were performed during the hospitalization. The timing of patient discharge was determined by the expert team of cardiologists in charge of the HF department; discharge was recommended when the patients presented no signs of decompensation, such as NYHA functional class <3, no sign of rales, no galloping rhythm, stable blood pressure and an improvement in renal function due to an optimal treatment that followed international guidelines.⁸ Rehospitalization for the enrolled patients was defined as hospitalization for decompensated HF. The primary end point was the first rehospitalization for decompensated HF.

Cardiac catheterization

Left ventricular pressure was recorded with a 5-F pigtail catheter. Left ventricular volume and ejection fraction were determined with left ventriculography with a contrast medium using Kennedy's formula. Right-sided catheterization was performed using a 7F Swan-Ganz catheter to measure pulmonary capillary wedge pressure, mean pulmonary artery pressure (PAP), right ventricular end-diastolic pressure and mean right atrial pressure. Cardiac output was measured using the estimated Fick principle and the Thermal dilution. Systemic vascular resistance and pulmonary vascular resistance were calculated using the established formulas: systemic vascular resistance = $80 \times (\text{mean pulmonary artery pressure} - \text{mean right atrial pressure}) / \text{cardiac output}$ and pulmonary vascular resistance = $80 \times (\text{mean pulmonary artery pressure} - \text{pulmonary capillary wedge pressure}) / \text{cardiac output}$.

Echocardiography

Echocardiographic examinations were performed with a Sonos-5500 (Philips Medical System, Andover, MA, USA), Alpha 10 (Hitachi-Aloka Medical, Tokyo, Japan), Vivid 7 Dimension (GE Healthcare, Buckinghamshire, UK), ACUSON Sequoia C256 (Mochida Simens Medical System, Tokyo, Japan) or Aplio XV (Toshiba Medical Systems, Tochigi, Japan) machine with a 2.5-MHz probe. Patients underwent a Doppler echocardiographic study for HF at admission and before discharge. Standard views were recorded, including the parasternal long-axis, short-axis and apical 4- and 2-chamber views, and cardiac chamber sizes and left atrial dimensions were evaluated according to the recommendations of the American Society of Echocardiography.⁹ The severity of valve regurgitation was quantified on a semicontinuous scale from none (0) to severe.⁴ Pulsed-wave Doppler examination and Doppler tissue imaging of the mitral annulus was performed. The peak mitral early diastolic inflow and atrial filling (E and A) velocities and the E -wave deceleration time were obtained. The sample volumes of the pulsed Doppler tissue imaging were determined at the septal and lateral margins of the mitral annulus. The peak

early mitral annular velocities were measured, and then the average values of the septal and lateral velocities were used as E' .

The mathematical model for the rehospitalization process

To construct a model for future rehospitalization using the basic clinical factors for the patients, we adopted two working assumptions for the practical rehospitalization process.

Assumption 1. A mean elapsed time τ_i from discharge to the rehospitalization of patient i depends on some of the given clinical factors $X^i = \{x_1^i, \dots, x_p^i\}$ of the patient, that is, a common subset $X_S^i \subseteq X^i$ over all patients. The dependency is primarily approximated by the following inverse linear relation:

$$\tau_i \cong \frac{1}{\sum_{x_j^i \in X_S^i} \beta_j x_j^i + \gamma} \quad (1)$$

where the denominator represents the expected frequency of cardiovascular rehospitalization per day, X_S^i is a set of values of the factors in X_S for patient i , β_j is the contributing weight of the j th factor to the frequency and γ is the intrinsic frequency for any patient.

Assumption 2. The clinical factors X_S^i of patient i are fairly stable between discharge and rehospitalization. Thus, the expectation value of the mean elapsed time τ_i remains nearly constant for patient i . As any event occurring with a constant frequency in a given time period is generated by a Poisson process,¹⁰ rehospitalization also occurs via this process under Assumption 2. Thus, the probability density $p_i(t)$ for the rehospitalization of patient i at an elapsed time t after discharge is represented by the following exponential formula:

$$p_i(t) = \frac{1}{\tau_i} \exp\left(-\frac{t}{\tau_i}\right) \quad (2)$$

The parameter τ_i is given by Equation (1) according to Assumption 1.

We next describe the assumption test. Assumption 1 is limited to the relationship between the parameter τ_i and the clinical factors X_S^i . If the accuracy of the approximation is insufficient, we can easily extend it to a nonlinear relation such as a higher-order polynomial. Assumption 2 essentially characterizes the process of the occurrence of rehospitalization and defines the formula for its probability density $p_i(t)$. Accordingly, before the modeling of the rehospitalization process based on a given data set, a test should be applied to verify that Assumption 2 actually holds true for the given data set.

With n samples in the data set $D = \{(X^i, \tau_i) | i = 1, \dots, n\}$, where X^i is the set of clinical factor values for patient i , and τ_i is the elapsed time at rehospitalization after discharge, we first compute a histogram of the rehospitalization occurrences over t , that is, the number of rehospitalization occurrences \hat{m}_k in each elapsed time interval $((k-1)\Delta t, k\Delta t)$ ($k = 1, \dots, q$) in the data set. The number of equal-width bins q into which to partition the sample range $[0, q\Delta t]$ is appropriately chosen to be $q = \sqrt{n}$. (Venables and Ripley)¹¹ We also expect a certain value of \hat{m}_k by Equation (2) under Assumption 2. The value \hat{m}_k computed from the data set and its value expected by Equation (2), m_k , should be consistent if Assumption 2 holds for the data set. Consistency with m_k and \hat{m}_k is evaluated by the following G-score:¹²

$$G = 2 \sum_{k=1}^q \hat{m}_k \ln \frac{\hat{m}_k}{m_k} \quad (3)$$

Because this G-score is known to follow a χ^2 distribution of degree $q-2$, we applied a χ^2 -test to the null hypothesis that the histogram of the given data set is consistent with Equation (2), that is, that Assumption 2 holds true for the data set. If the P -value of the test is less than a specific risk level α such as $\alpha = 0.05$, we conclude that Assumption 2 does not hold for the data set. This G-test is known to be more rigorous than the well-known Pearson's χ^2 -test.

Thus, our problem was to derive the expectation value m_k ($k = 1, \dots, q$) from Equation (2). We considered that τ_i of the patients in D are sampled from a common population distribution $p_\tau(\tau)$. Therefore, the total probability

distribution of the rehospitalization time $P(t)$ is expected to be a superposition of Equation (2) for various τ sampled from $p_\tau(\tau)$, as follows, where $p(t)$ is $p_\tau(t)$ in Equation (2) for a general τ :

$$P(t) = \int_0^\infty p_\tau(\tau)p(t)d\tau = \int_0^\infty p_\tau(\tau)\frac{1}{\tau}\exp\left(-\frac{t}{\tau}\right)d\tau$$

We use the following natural conjugate prior distribution for the unknown $p_\tau(\tau)$:

$$p_\tau(\tau) = \frac{\tau^{-n}\exp\left(-1/\tau\sum_{i=1}^n\tau_i\right)}{\int_0^\infty\tau^{-n}\exp\left(-1/\tau\sum_{i=1}^n\tau_i\right)d\tau}$$

where τ_i is given by the data set D . The selection of this parameter distribution is widely considered to be reasonable in Bayesian statistics because it preserves the exponential shape of the distribution of elapsed times t .¹³ After several manipulations, the following $P(t)$ is derived:

$$P(t) = \frac{(n+1)\left(\sum_{i=1}^n\tau_i\right)^{n+1}}{\left(\sum_{i=1}^n\tau_i+t\right)^{n+2}}$$

Accordingly, the expectation m_k is given by the accumulation of $P(t)$ over $((k-1)\Delta t, k\Delta t]$ as follows:

$$m_k = n \int_{(k-1)\Delta t}^{k\Delta t} P(t)dt = n \left(\frac{\sum_{i=1}^n\tau_i}{\sum_{i=1}^n\tau_i+(k-1)\Delta t} \right)^{n+1} - n \left(\frac{\sum_{i=1}^n\tau_i}{\sum_{i=1}^n\tau_i+k\Delta t} \right)^{n+1} \tag{4}$$

Using Equations (3) and (4), we tested the validity of Assumption 2 for the given data set D .

Finally, we describe the modeling algorithm. First, the value of every factor x_j^i for all patients $i=1, \dots, n$ in D was normalized to fit into the interval $[0,1]$ using the maximum and minimum values. This normalization to eliminate differences in the factor scales was necessary to allow for the measurement of the essential contribution of each factor's variation to τ_i . Subsequently, we applied Equations (1) and (2) to the normalized data set D_N to model the probabilistic rehospitalization process when Assumption 2 holds for the data set. We determined the model parameters β_j and γ in Equation (1) to maximize the following objective function:

$$L(\beta_1, \dots, \beta_p, \gamma) = \ln \left[\prod_{i=1}^n \left(\sum_{j=1}^p \beta_j x_j^i + \gamma \right) \exp \left\{ - \left(\sum_{j=1}^p \beta_j x_j^i + \gamma \right) \tau_i \right\} \right] - \lambda \left(\sum_{j=1}^p |\beta_j| + |\gamma| \right) \tag{5}$$

The first term is the log-likelihood of the model consisting of Equations (1) and (2) over D_N . The second term is called an $L1$ -regularization term, which penalizes the coefficients of negligible factors by setting them equal to zero when the larger hyper-parameter λ eliminates more factors.^{13,14} This term avoids the over-fitting of the model to the data set by selecting a set of effective factors X_j^i from a given X^i . In our study, λ is tuned to be 0.02 to maintain the largest value of Equation(5) similarly to the other parameters β_j and γ .

To seek the optimum parameter values of $\beta_1, \dots, \beta_p, \gamma$ that maximize the objective function $L(\beta_1, \dots, \beta_p, \gamma)$, we applied a simple greedy hill-climbing algorithm, in which the parameter values are iteratively modified toward their gradient direction $(\partial L/\beta_1, \dots, \partial L/\beta_p, \partial L/\gamma)$. When the improvement of L becomes nearly negligible, the resulting parameter values are taken as the optima. Because this process depends on the initial values of the parameters,

we repeated this optimization 100 times starting with random initial values and selected the result providing the maximum L .

RESULTS

Patients characteristics

Out of the 151 patients, 36 died of cardiovascular events after rehospitalization during the follow-up period. The remaining 115 patients were readmitted to our hospital at a median time of 296 days after discharge (range, 3–1891). Among these patients, the HF etiologies were valvular heart disease ($n=38$), dilated cardiomyopathy ($n=30$), hypertrophic cardiomyopathy ($n=22$), ischemic heart disease ($n=20$), hypertensive heart disease ($n=17$) and others. Their mean age was 68.6 ± 14.6 years (range, 19–93), and 38% of the patients were women. The clinical characteristics of the 151 patients are summarized in Table 1.

Validation of the formula

We hypothesized that the time-to-rehospitalization histogram for all patients (Figure 1) should be distributed exponentially if the mathematically estimated formula for the prognosis of each patient is regarded as a Poisson distribution. We therefore validated the assumptions of the model architecture. The goodness of fit was controlled by a χ^2 -test, considering that the incidence rates of rehospitalization or death differ depending on the patients. Thus, the null hypothesis that the observed frequency is a mixed Poisson process was tested, as explained in the Methods section. We chose an elapsed time to rehospitalization of 150 days, which is one-thirteenth of the range of the time interval $[1,1,950]$ according to the measure of $q = \sqrt{n} = \sqrt{151} \cong 13$. As a result, the P -value was 0.29, which was far larger than 0.05, and we confirmed that the null hypothesis was not rejected. Therefore, we concluded that the mathematically derived estimation formula for the rehospitalization of each patient was a mixed Poisson distribution.

Factors in rehospitalization for HF

We collected 402 clinical factors (Figures 2 and 3), and 150 out of 402 factors having small effects on the prognosis were automatically excluded by the regularization method described in the Methods section. Finally, we selected 252 factors for the analysis (Figures 2 and 3). The estimation results for the attribute coefficients are presented in bar graph form and numerically.

Regarding underlying diseases in HF, whereas dilated cardiomyopathy (-4.5), hypertrophic cardiomyopathy (-1.5) and hypertensive heart disease (-1.0) had better outcomes, valvular disease (7.4) and dilated phase hypertrophic cardiomyopathy (2.4) had poor prognoses. Ischemia (4.4) was the worst trigger of HF. Based on laboratory data, whereas elevated inflammatory response values, such as white blood cell counts ($-1.6/5.8$; at admission/at discharge) or C-reactive protein levels ($-2.2/8.1$; at admission/at discharge), did not indicate a poor prognosis at admission, these elevated inflammatory response values at discharge were associated with a poor prognosis. Increases in the levels of aspartate aminotransferase (6.6), alanine aminotransferase (3.2), uric acid (6.6) and BNP (4.8) at discharge also indicated a poor prognosis. Patients who received dopamine (11.9), isosorbide dinitrate (5.0) or diuretic (2.0) infusions in the acute management of HF showed worse prognoses. In contrast, the use of dobutamine (-2.5) or nitroglycerin (-2.5) drip infusions resulted in better prognoses.

Regarding oral medications at discharge, the angiotensin-converting enzyme alacepril (-4.2), the β -blocker carvedilol (-7.1 , the best response), the angiotensin receptor blocker telmisartan (-1.6), the diuretic furosemide (-4.2), the lipid-lowering drugs pitavastatin

Table 1 Patient characteristics

	Population (n = 151)
Age (years)*	68.6 ± 14.6
Gender, female, n (%)	58 (38)
Medical history	
Frequency of heart failure (time)*	3.2 ± 2.5
Hypertension	73 (48)
Diabetes mellitus	55 (36)
Hyperlipidemia	45 (30)
Signs at admission	
Elevated jugular venous pressure	84 (56)
S ₃ gallop	85 (56)
Lower extremity edema	76 (50)
NYHA functional class: II/III/IV	54/44/53
Clinical scenario: 1/2/3/4/5	28/77/34/0/12
Nohria—profile A	2 (1)
Nohria—profile B	108 (72)
Nohria—profile C	28 (19)
Nohria—profile L	13 (9)
Baseline characteristics at admission/at discharge	
Heart rate (beats min ⁻¹)*	84.4 ± 26.7/73.2 ± 58.3
Systolic BP (mm Hg)*	124.4 ± 31.8/ 111.0 ± 15.8
Diastolic BP (mm Hg)*	68.5 ± 17.5/59.4 ± 8.4
Body weight (kg)*	57.3 ± 13.5/52.3 ± 11.9
Δ Body weight (kg)*	4.6 ± 3.8
Laboratory factors at admission/at discharge	
Hemoglobin (g dl ⁻¹)*	12.4 ± 7.7/11.8 ± 2.0
Leukocytes (10 ⁹ l ⁻¹)*	6940 ± 2982/ 5968 ± 2464
Blood urea nitrogen (mg dl ⁻¹)*	28.6 ± 20.7/30.0 ± 19.7
Creatinine (mg dl ⁻¹)*	1.27 ± 0.90/1.24 ± 0.69
Sodium (mEq l ⁻¹)*	137.6 ± 3.9/136.8 ± 4.3
Uric acid (mg dl ⁻¹)*	7.5 ± 2.0/7.4 ± 2.1
T-bil (mg dl ⁻¹)*	0.92 ± 0.67/0.71 ± 0.42
C-reactive protein (mg dl ⁻¹)*	1.3 ± 2.8/0.7 ± 1.8
BNP (pg ml ⁻¹)*	920 ± 956/439 ± 548
Δ BNP (pg ml ⁻¹) (1 month after discharge-at discharge)*	78 ± 226
Echocardiographic factors at admission/at discharge	
Left ventricular end-diastolic dimension (mm)*	58.9 ± 13.3/58.3 ± 11.9
Left ventricular end-systolic dimension (mm)*	47.4 ± 15.2/45.8 ± 14.6
Fractional shortening (%)*	21.2 ± 11.5/23.1 ± 11.4
Ventricular septum thickness (mm)*	9.6 ± 2.9/9.6 ± 2.7
Posterior wall thickness (mm)*	9.8 ± 2.5/9.6 ± 2.0
Left atrial diastolic dimension (mm)*	49.9 ± 8.1/47.8 ± 9.3
Pressure across tricuspid valve (mm Hg)*	37.0 ± 16.3/25.4 ± 10.5
Medication at admission	
Use of dopamine, n (%)	10 (6)
Use of dobutamine, n (%)	33 (22)
Use of phosphodiesterase inhibitor, n (%)	13 (9)
Use of carperitide, n (%)	32 (21)
Use of nitroglycerin, n (%)	22 (15)
Use of diuretics, n (%)	60 (40)

Abbreviations: BNP, B-type natriuretic peptide; BP, blood pressure; NYHA, New York Heart Association; T-bil, total bilirubin.

*Plus or minus values are means ± s.d. Clinical profiles were classified as profile A (dry-warm), B (wet-warm), C (wet-cold) or L (dry-cold).

(−3.3), atorvastatin (−2.9) and ezetimibe (−2.2), the coronary dilator isosorbide dinitrate (−3.1), the antiallergic fexofenadine hydrochloride (−5.1), the sedative-hypnotic triazolam (−3.2), proton pump inhibitor lansoprazole (−0.9) and all antifatulents, except toughmac, led to better prognoses. However, Ca inhibitor nifedipine (9.4) resulted in the worst outcome, and all diabetes drugs, antiarrhythmic drugs, potassium agents, vitamins and purgatives, excluding senna, were associated with worse prognoses.




Figure 1 Time-to-rehospitalization histogram for all patients.

Fitting the model to clinical data

The mean actual value for rehospitalization (X) was 388 ± 377 days, whereas the mean estimated value calculated by the probability model based on a Poisson process (Y) was 398 ± 381 days; X and Y were very tightly correlated (Figure 4). The results showed that the mathematical formula for rehospitalization time is the dependent variable, and the clinical and personal factors before rehospitalization are the independent variables.

DISCUSSION

This study provided evidence that the values of numerous factors, including risk factors at one phase of disease, can be used to construct a mathematical equation to predict clinical outcomes. We were able to derive the equation $\tau = f(x_1, \dots, x_p)$, where τ is the time to a future clinical event and x_1, \dots, x_p are clinical factors observed before the event. In this case, τ represents the days until rehospitalization after discharge, and x_1, \dots, x_p are the clinical and personal factors for patients hospitalized for ADHF. This study provides evidence that the clinical outcome of τ in this context is a function of 252 significant factors such as plasma BNP levels at and soon after discharge. This study presents the time to rehospitalization as the dependent variable and the clinical and personal factors before rehospitalization as the independent variables.

This study suggests the novel idea that the time to clinical events, such as rehospitalization or death, can be mathematically formulated from clinical and personal factors, demonstrating that clinical medicine can engage in physical science. The novelty of this study is based on the fact that clinical outcomes have been thought to be determined mainly from medical knowledge and the experience of the physicians. It can be argued that the known effectiveness of drugs may determine the time course of clinical events. Although this is partially true,^{15–17} no one knows how one drug or the combination of several drugs affects patients with different degrees of severity of a given disease. It may also be argued that large-scale trials may better depict clinical outcomes; for example, the patients with BNP levels of < 170 pg/ml showed a 20% reduction of rehospitalization compared with the patients with BNP levels greater than 170 pg/ml.^{18,19} Evaluating such results by Kaplan–Meier analysis is common in clinical medicine; however, this analysis only provides the average tendency of the average patient to undergo rehospitalization and does not

Predictor variables	maximum value	coefficient	graph	Predictor variables	maximum value	coefficient	graph	Predictor variables	maximum value	coefficient	graph
Age	93.0	-0.578	I	Laboratory data on admission: platelet				Right heart catheterization: body surface area			
Gender	1.0	-4.455	II	Laboratory data on admission: albumin				Left heart catheterization: systolic aortic pressure			
Etiology of HF: dilated cardiomyopathy	1.0	-4.471	II	Laboratory data on admission: total bilirubin	6.7	-1.697	II	Left heart catheterization: diastolic aortic pressure			
Etiology of HF: dilated phase hypertrophic cardiomyopathy	1.0	2.409	II	Laboratory data on admission: AST	789.0	2.740	II	Left heart catheterization: aortic pressure mean	136.0	-1.159	II
Etiology of HF: hypertensive heart disease	1.0	-1.044	II	Laboratory data on admission: ALT	653.0	1.359	II	Left heart catheterization: LV ejection fraction	3.0	0.519	II
Etiology of HF: ischemic heart disease (ICM)				Laboratory data on admission: sodium				Left heart catheterization: LVEDVI	477.0	2.252	II
Etiology of HF: hypertrophic cardiomyopathy	1.0	-1.493	II	Laboratory data on admission: potassium				Left heart catheterization: LVESVI	432.0	0.772	II
Etiology of HF: cardiac sarcoidosis				Laboratory data on admission: creatinin				Prognosis: left ventricle assisting system	1.0	-3.224	II
Etiology of HF: myocarditis				Laboratory data on admission: blood urea nitrogen				Cardiac resynchronization therapy: this admission	1.0	-2.286	II
Etiology of HF: valvular heart disease	1.0	7.361	III	Laboratory data on admission: uric acid				Cardiac resynchronization therapy: prior admission	1.0	2.521	II
Etiology of HF: others	1.0	3.789	II	Laboratory data on admission: C-reactive protein	24.5	-2.160	II	Implantable cardioverter-defibrillator: this admission	1.0	-2.995	II
Etiology of HF: valvular heart disease + ICM	1.0	0.445	I	Laboratory data on admission: blood sugar				Implantable cardioverter-defibrillator: prior admission	1.0	1.881	II
Endomyocardial biopsy: with or without	1.0	2.475	II	Laboratory data on admission: hemoglobin A1c				Pacemaker: this admission			
Comorbidity: diabetes mellitus				Laboratory data on admission: BNP				Pacemaker: prior admission	1.0	4.092	II
Comorbidity: Hypertension	1.0	1.968	II	Laboratory data on admission: iron	421.1	-0.162	I	coronary artery bypass graft: this admission	1.0	0.976	II
Comorbidity: Hyperlipidemia	1.0	-1.886	II	Laboratory data on admission: UIBC	477.0	1.729	II	coronary artery bypass graft: prior admission	1.0	-2.455	II
Comorbidity: chronic atrial fibrillation	1.0	3.544	II	Laboratory data on admission: ferritin				Percutaneous coronary intervention: this admission	1.0	-4.455	II
Comorbidity: cerebrovascular disease	1.0	1.172	I	Laboratory data on admission: free T3	12.6	-1.623	II	Percutaneous coronary intervention: prior admission	1.0	-2.419	II
Comorbidity: chronic obstructive pulmonary disease	1.0	3.318	II	Laboratory data on admission: free T4				Vascular surgery: this admission	1.0	-0.825	II
Comorbidity: arteriosclerosis obliterans	1.0	-1.547	II	Laboratory data on admission: thyrotropin-stimulating hormone				Vascular surgery: prior admission	1.0	5.861	II
Family history of cardiovascular disease				Echocardiographic data on admission: LVdD	106.0	-1.205	I	Vascular disease: aneurysm	1.0	3.159	II
Frequency of HF				Echocardiographic data on admission: LVdS	95.0	-3.233	II	Ablation: this admission			
Number of living with family	6.0	0.386	I	Echocardiographic data on admission: %FS	81.0	5.205	III	Ablation: prior admission			
Partner: with or without	1.0	1.599	II	Echocardiographic data on admission: IVS	20.0	2.210	II	Other surgery: prior admission	1.0	-3.880	II
Alcohol intake				Echocardiographic data on admission: PW	21.0	3.576	II	Valvular surgery: this admission			
Onset type of HF: ADHF (de novo)	1.0	-1.627	II	Echocardiographic data on admission: LAD	98.0	-0.747	I	Valvular surgery: prior admission	1.0	-5.514	II
Onset type of HF: acute on chronic				Echocardiographic data on admission: TMF-E	269.0	-1.760	II	Mitral valve plasty: this admission			
Onset type of HF: others				Echocardiographic data on admission: TMF-A	152.0	-2.120	II	Mitral valve plasty: prior admission	1.0	-2.491	II
Trigger of ADHF: volume over	1.0	-2.806	II	Echocardiographic data on admission: TMF-DcT				Tricuspid annuloplasty or valve replacement: this admission			
Trigger of ADHF: arrhythmia	1.0	-0.271	I	Echocardiographic data on admission: TR grade	13.0	-3.414	II	Tricuspid annuloplasty or valve replacement: prior admission	1.0	2.126	II
Trigger of ADHF: infection				Echocardiographic data on admission: TRPG				Aortic valve replacement: this admission			
Trigger of ADHF: anemia	1.0	-3.122	II	Echocardiographic data on admission: PAEDP				Aortic valve replacement: prior admission			
Trigger of ADHF: others	1.0	1.114	I	Echocardiographic data on admission: MR grade	4.0	-2.910	II	Findings at discharge: systolic blood pressure			
Trigger of ADHF: afterload mismatch	1.0	2.375	II	Echocardiographic data on admission: MR grade	4.0	0.344	II	Findings at discharge: diastolic blood pressure			
Trigger of ADHF: ischemia	1.0	4.390	III	Echocardiographic data on admission: AS	1.0	0.936	II	Findings at discharge: heart rate	772.0	-2.456	II
Trigger of ADHF: missed drug	1.0	2.713	II	Echocardiographic data on admission: MS	1.0	5.126	III	Findings at discharge: body weight			
Trigger of ADHF: chronic change (unclear)				Medications on admission: beta-blocker	1.0	-3.031	II	Difference of body weight (on admission - at discharge)			
Nohria: cold	1.0	-2.750	II	Medications on admission: ACEI	1.0	3.098	II	Laboratory data at discharge: leukocyte	23500.0	5.780	III
Nohria: wet				Medications on admission: ARB	1.0	-2.150	II	Laboratory data at discharge: neutrophil			
Nohria: warm	1.0	1.553	II	Medications on admission: eplerenone	1.0	5.156	III	Laboratory data at discharge: lymphocyte	58.6	-0.270	II
Nohria: dry	1.0	-3.422	II	Medications on admission: other diuretics	1.0	8.603	III	Laboratory data at discharge: hemoglobin			
Clinical scenario: 1	1.0	-0.867	I	Medications on admission: spironolactone	1.0	3.804	II	Laboratory data at discharge: platelet			
Clinical scenario: 2	1.0	2.704	II	Medications on admission: amiodarone	1.0	3.860	II	Laboratory data at discharge: albumin	5.3	-1.356	II
Clinical scenario: 3	1.0	2.947	II	Medications on admission: warfarine	1.0	-0.196	I	Laboratory data at discharge: total bilirubin			
Clinical scenario: 5	1.0	-3.387	II	Medications on admission: statin	1.0	4.241	II	Laboratory data at discharge: AST	575.0	6.585	III
Findings on admission: NYHA	4.0	-4.070	III	Medications on admission: DM (oral drug)	1.0	1.750	II	Laboratory data at discharge: ALT	511.0	3.184	II
Findings on admission: systolic blood pressure				Medications on admission: DM (insulin)				Laboratory data at discharge: sodium			
Findings on admission: diastolic blood pressure				Medications on admission: digoxin				Laboratory data at discharge: potassium	8.5	0.345	II
Findings on admission: heart rate	200.0	0.447	I	Acute phase treatment: carperitide	1.0	1.177	II	Laboratory data at discharge: creatinin			
Findings on admission: body weight				Acute phase treatment: dopamine	1.0	11.918	III	Laboratory data at discharge: blood urea nitrogen			
Findings on admission: body height				Acute phase treatment: dobutamin	1.0	-2.537	II	Laboratory data at discharge: uric acid	16.4	6.567	III
Findings on admission: chest X-ray CTR	88.0	-3.346	II	Acute phase treatment: isosorbide dinitrate	1.0	5.039	III	Laboratory data at discharge: C-reactive protein	17.2	8.109	III
Findings on admission: congestion				Acute phase treatment: nitroglycerin	1.0	-2.537	II	Laboratory data at discharge: blood sugar			
Findings on admission: α_2 gallop	1.0	6.263	III	Acute phase treatment: diuretics venoclysis	1.0	1.993	II	Laboratory data at discharge: BNP	3832.6	4.770	II
Findings on admission: nocturnal dyspnea	1.0	5.619	III	Acute phase treatment: phosphodiesterase II inhibitor				Laboratory data one month after discharge: creatinin			
Findings on admission: elevated jugular venous pressure	1.0	0.224	I	Use of biphasic positive airway pressure				Laboratory data one month after discharge: BNP	2397.6	-3.767	II
Findings on admission: lower extremity edema	1.0	-3.961	II	Use of adaptive servo ventilator	1.0	0.228	I	Laboratory data: difference of BNP (1 month - at discharge)	1655.3	1.570	II
Findings on admission: coldness of limbs	1.0	-3.216	II	Use of assist device: IABP or PCPS	3.0	3.310	II	Echocardiographic data at discharge: LVdD			
Findings on admission: respiratory rate				Use of assist device:left ventricle assisting system	1.0	3.993	II	Echocardiographic data at discharge: LVdS			
Findings on admission: percutaneous oxygen saturation	100.0	-1.137	I	Use of blood transfusion				Echocardiographic data at discharge: %FS			
Findings on admission: fraction of inspired oxygen	100.0	-3.858	II	Right heart catheterization: pulmonary capillary wedge pressure				Echocardiographic data at discharge: IVS			
ECG (rhythm): sinus rhythm				Right heart catheterization: right atrium	18.0	-3.104	II	Echocardiographic data at discharge: PW	18.0	0.643	I
ECG (rhythm): atrial fibrillation or tachycardia or flutter	1.0	-0.745	I	Right heart catheterization: systolic right ventricle				Echocardiographic data at discharge: LAD	75.0	-6.889	II
ECG (rhythm): sick sinus syndrome				Right heart catheterization: diastolic right ventricle	20.0	-1.569	II	Echocardiographic data at discharge: AR	3.5	3.091	II
ECG (rhythm): pacemaker	1.0	-5.431	III	Right heart catheterization: systolic pulmonary artery				Echocardiographic data at discharge: MR	4.0	-0.457	I
ECG (rhythm): complete atrioventricular block				Right heart catheterization: diastolic pulmonary artery				Echocardiographic data at discharge: TR			
ECG (rhythm): others				Right heart catheterization: mean pulmonary artery				Echocardiographic data at discharge: TRPG	66.0	0.456	I
ECG: ventricular tachycardia or fibrillation	1.0	0.404	I	Right heart catheterization: cardiac output (c-Fick)	7.6	0.646	II	Echocardiographic data at discharge: IVC	1.0	-1.421	I
ECG: complete left bundle branch block	1.0	3.116	II	Right heart catheterization: cardiac index (c-Fick)	4.3	1.574	II	Echocardiographic data at discharge: TMF-E	230.0	0.980	I
Laboratory data on admission: leukocytes	26300.0	-1.619	II	Right heart catheterization: cardiac output (Thermo)	9.7	3.877	II	Echocardiographic data at discharge: TMF-A			
Laboratory data on admission: neutrophil				Right heart catheterization: cardiac index (Thermo)	6.3	4.170	II	Echocardiographic data at discharge: DcT			
Laboratory data on admission: lymphocyte				Right heart catheterization: systemic vascular resistance				Echocardiographic data at discharge: E/E'	55.0	5.962	II
Laboratory data on admission: hemoglobin				Right heart catheterization: pulmonary vascular resistance							

Figure 2 Factors influencing the estimation of rehospitalization for HF and the contribution of each parameter. All of the clinical and personal factors for the patients with HF. Predictor variables with coefficient indicate the factors selected after the application of the regularization method. Negative values indicate favorable impact on prognosis, whereas positive values indicate undesirable effect. HF, heart failure; ADHF, acute decompensated heart failure; NYHA, New York Heart Association; CTR, cardiothoracic ratio; ECG, electrocardiogram; AST, aspartate aminotransferase; ALT, alanine aminotransferase; BNP, B-type natriuretic peptide; UIBC, unsaturated iron-binding capacity; LVdD, left ventricular end-diastolic dimension; LVdS, left ventricular end-systolic dimension; FS, fractional shortening; IVS, interventricular septal thickness; PW, left ventricular posterior thickness; LAD, left atrial dimension; TMF-E, the peak mitral inflow early diastolic velocity; TMF-A, the peak mitral inflow atrial filling; DcT, deceleration time; TR PG, tricuspid regurgitation pressure gradient; PAEDP, pulmonary artery end-diastolic pressure; MR, mitral regurgitation; AR, aortic regurgitation; AS, aortic stenosis; MS, mitral stenosis; ACEI, angiotensin-converting enzyme inhibitors; ARB, angiotensin receptor blockers; DM, diabetes mellitus; IABP, intraaortic balloon pumping; PCPS, percutaneous cardio pulmonary support; EDVI, end-diastolic volume index; ESVI, end-systolic volume index; IVC, inferior vena cava respiratory change; E/E', ratio of peak mitral E-wave velocity to peak mitral annular velocity.

prospectively provide a future clinical outcome for each patient. Indeed, in the epidemiological study, many biomarkers, such as BNP levels or C-reactive protein levels in addition to the classical risk factors, such as hypertension or diabetes mellitus, are known to be related to cardiovascular events and death. However, Wang *et al.*²⁰ showed that although multiple biomarkers are associated with a high relative risk of adverse events, even in the combination of these factors they add only moderately to the prediction of risk in an individual person. This suggests that the occurrence of cardiovascular events may not be well predictable or mathematically formulated. On the other hand, using the formula developed in this study, we can identify the

day of a clinical event to within a small range, suggesting that we need more clinical data to predict the future outcomes or obtain the mathematical formula for the prediction than we expected.

It would be difficult to strictly prove that this mathematical formula is correct because no gold standard or correct answer is available in the medical literature. However, there are hints as to the correctness of this formula. First, we assume that the probability of rehospitalization follows a Poisson distribution; if this is true, a histogram of the day of rehospitalization after discharge should follow a Poisson distribution. We found that the present data for the actual day of rehospitalization are distributed as a Poisson distribution.




Figure 4 Correlation between the clinical data and the values calculated using the mathematical formula. The clinical data are in excellent agreement with the calculated times.

factors with strong effects on prognosis could have confounded each other, the results of this study are probably not weakened because we obtained a good fitting to the clinical outcome using these factors. When we consider the clinical and pathophysiological meaning of each factor, we need to pay attention to each factor independently.

The other main limitation of this study is that the patient population consists of a retrospective cohort. However, because we enrolled all of the patients who were admitted to our department during the entry period, the selection bias may be small. Furthermore, this is a single-center study, so the formula may be true only in our institute. However, because (1) approximately one-half of the patients who were hospitalized during this time were referred from other hospitals, (2) the nature and treatment of HF did not differ among the hospitals and (3) our hospital sets a high standard for CHF treatment and specializes in receiving CHF patients from all over Japan; we believe that the formula developed in this study may be generalized. We estimated the day of rehospitalization in this study; however, the important issue is the ability to make this prediction, which needs further investigation.

CONCLUSIONS

This study demonstrated that clinical medicine and practice can use a mathematical formula to predict clinical outcomes or events using current data. A prospective study is needed to test whether this formula predicts the day of rehospitalization in CHF patients who are admitted because of ADHF and discharged after treatment. The application of these risk factors to individual CHF patients may distinguish those patients who are at low risk from those who are at high risk and may benefit from closer monitoring and aggressive treatment.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We thank T Takayama, M Shishikura and Y Miyawaki for assisting in the collection of data. This work is supported by grants-in-aid from the Japanese Ministry of Health, Labor and Welfare (H23-Nanchi-Ippan-22 to MK) and the Japanese Ministry of Education, Culture, Sports, Science and Technology (21390251 to MK), and also by grants from the Japan Heart Foundation (PPAR to MK) and the Japan Cardiovascular Research Foundation (Stacin to MK).

- Braunwald E. Biomarkers in heart failure. *N Engl J Med* 2008; **358**: 2148–2159.
- Fonarow GC, Peacock WF, Phillips CO, Givertz MM, Lopatin M. Admission B-type natriuretic peptide levels and in-hospital mortality in acute decompensated heart failure. *J Am Coll Cardiol* 2007; **49**: 1943–1950.
- Abraham WT, Fonarow GC, Albert NM, Stough WG, Gheorghiadu M, Greenberg BH, O'Connor CM, Sun JL, Yancy CW, Young JB. Predictors of in-hospital mortality in patients hospitalized for heart failure: insights from the Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients with Heart Failure (OPTIMIZE-HF). *J Am Coll Cardiol* 2008; **52**: 347–356.
- Mancini DM, Eisen H, Kussmaul W, Mull R, Edmunds Jr LH, Wilson JR. Value of peak exercise oxygen consumption for optimal timing of cardiac transplantation in ambulatory patients with heart failure. *Circulation* 1991; **83**: 778–786.
- Itoh H, Taniguchi K, Koike A, Doi M. Evaluation of severity of heart failure using ventilatory gas analysis. *Circulation* 1990; **81**(Suppl): I131–I137.
- Newton I. *The Mathematical Principles of Natural Philosophy, Book I*. Benjamin Motte: London, 1687.
- McKee PA, Castelli WP, McNamara PM, Kannel WB. The natural history of congestive heart failure: the Framingham study. *N Engl J Med* 1971; **285**: 1441–1446.
- Hunt SA, Abraham WT, Chin MH, Feldman AM, Francis GS, Ganiats TG, Jessup M, Konstam MA, Mancini DM, Michl K, Oates JA, Rahko PS, Silver MA, Stevenson LW, Yancy CW 2009 Focused update incorporated into the ACC/AHA 2005 guidelines for the diagnosis and management of heart failure in adults: A Report of the American College of Cardiology Foundation/American heart association task force on practice guidelines: developed in collaboration with the International Society for Heart and Lung Transplantation. *Circulation* 2009; **119**: e391–e479.
- Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, Picard MH, Roman MJ, Seward J, Shanewise JS, Solomon SD, Spencer KT, Sutton MS, Stewart WJ. Recommendations for chamber quantification: a report from the American Society of Echocardiography's Guidelines and Standards Committee and the Chamber Quantification Writing Group, developed in conjunction with the European Association of Echocardiography, a branch of the European Society of Cardiology. *J Am Soc Echocardiogr* 2005; **18**: 1440–1463.
- Gullberg J. *Mathematics from the Birth of Numbers*. WW Norton: New York, 1997.
- Venables WN, Ripley BD. *Modern Applied Statistics with S*. Springer: Berlin, 2002.
- Sokal RR, Rohlf FJ. *Biometry: The Principles and Practice of Statistics in Biological Research*. Freeman: New York, 1994.
- Bishop CM. *Pattern Recognition and Machine Learning*. Springer: New York, 2006.
- Tibshirani R. Regression shrinkage and selection via the Lasso. *J R Stat Soc Ser B* 1996; **58**: 267–288.
- The Digitalis Investigation Group. The effect of digoxin on mortality and morbidity in patients with heart failure. *N Engl J Med* 1997; **336**: 525–533.
- Packer M, Bristow MR, Cohn JN, Colucci WS, Fowler MB, Gilbert EM, Shusterman NH. Carvedilol Heart Failure Study Group. The effect of carvedilol on morbidity and mortality in patients with chronic heart failure. *N Engl J Med* 1996; **334**: 1349–1355.
- Jong P, Yusuf S, Rousseau MF, Ahn SA, Bangdiwala SI. Effect of enalapril on 12-year survival and life expectancy in patients with left ventricular systolic dysfunction: a follow-up study. *Lancet* 2003; **361**: 1843–1848.
- Maeda K, Tsutomoto T, Wada A, Mabuchi N, Hayashi M, Tsutsui T, Ohnishi M, Sawaki M, Fujii M, Matsumoto T, Kinoshita M. High levels of plasma brain natriuretic peptide and interleukin-6 after optimized treatment for heart failure are independent risk factors for morbidity and mortality in patients with congestive heart failure. *J Am Coll Cardiol* 2000; **36**: 1587–1593.
- Daniels LB, Maisel AS. Natriuretic peptides. *J Am Coll Cardiol* 2007; **50**: 2357–2368.
- Wang TJ, Gona P, Larson MG, Tofler GH, Levy D, Newton-Cheh C, Jacques PF, Rifai N, Selhub J, Robins SJ, Benjamin EJ, D'Agostino RB, Vasan RS. Multiple biomarkers for the prediction of first major cardiovascular events and death. *N Engl J Med* 2006; **355**: 2631–2639.

Direct comparison of the diagnostic capability of cardiac magnetic resonance and endomyocardial biopsy in patients with heart failure

Akemi Yoshida¹, Hatsue Ishibashi-Ueda², Naoaki Yamada³, Hideaki Kanzaki¹, Takuya Hasegawa¹, Hiroyuki Takahama¹, Makoto Amaki¹, Masanori Asakura^{1,4}, and Masafumi Kitakaze^{1,4*}

¹Department of Cardiovascular Medicine, National Cerebral and Cardiovascular Center, Suita, Osaka, Japan; ²Department of Pathology, National Cerebral and Cardiovascular Center, Suita, Osaka, Japan; ³Department of Radiology, National Cerebral and Cardiovascular Center, Suita, Osaka, Japan; and ⁴Department of Clinical Research and Development, National Cerebral and Cardiovascular Center, Suita, Osaka, Japan

Received 19 July 2012; revised 11 September 2012; accepted 30 November 2012

Aims The diagnostic performance of cardiac magnetic resonance (CMR) has not been compared with that of other imaging modalities. Therefore, this study investigated the diagnostic capabilities of CMR and endomyocardial biopsy (EMB) in patients with heart failure (HF).

Methods and results We studied 136 patients with cardiomyopathy who underwent both CMR and EMB. Independent diagnoses were made according to the results of (i) CMR alone; (ii) EMB alone; (iii) clinical data plus echocardiogram; (iv) clinical data, echocardiogram, plus CMR; and (v) clinical data, echocardiogram, plus EMB. These diagnoses were then compared with the final diagnosis (gold standard) that was made using the complete clinical data, including EMB and CMR. The sensitivities of the diagnosis strategies of (i–v) relative to the final diagnosis were 67, 79, 86, 97, and 100%, respectively. CMR alone demonstrated better sensitivity for cardiac sarcoidosis and greater specificity for dilated cardiomyopathy than EMB alone. CMR also tended to show better sensitivity for hypertensive heart disease. There was no difference between the diagnostic capability of CMR and EMB for hypertrophic cardiomyopathy (HCM). However, CMR showed excellent sensitivity (100%) for apical and obstructive HCM, whereas EMB displayed better sensitivity for dilated HCM. Moreover, combined diagnosis with clinical data, echocardiogram, plus CMR achieved superior agreement with the final diagnosis in comparison with EMB alone.

Conclusion Non-invasive CMR demonstrated excellent diagnostic capability for patients with HF and was as effective as or superior to EMB. In particular, the use of CMR in combination with clinical data unrelated to EMB may provide excellent diagnostic accuracy for HF.

Keywords Heart failure • CMR • Endomyocardial biopsy • Diagnosis • Aetiology • Cardiomyopathy

Introduction

Heart failure (HF) is a common clinical syndrome caused by various cardiovascular diseases.¹ Despite the discovery, development, and adoption of novel therapies for HF, the mortality and morbidity resulting from this condition have remained high and

are currently increasing. Accordingly, accurate diagnosis of the underlying aetiology of HF is important for appropriate management and treatment. In addition to conventional clinical methods, gadolinium-enhanced cardiac magnetic resonance (CMR) and endomyocardial biopsy (EMB) are useful diagnostic modalities for identifying the aetiology of HF. EMB is considered

* Corresponding author. Departments of Clinical Research and Development, and Cardiovascular Medicine, National Cerebral and Cardiovascular Center, 5-7-1 Fujishiro-dai, Suita, Osaka 565-8565, Japan. Tel: +81 6 6833 5012, Fax: +81 6 6836 1120, Email: kitakaze@zf6.so-net.ne.jp

Published on behalf of the European Society of Cardiology. All rights reserved. © The Author 2013. For permissions please email: journals.permissions@oup.com.

to be the gold standard for diagnosing myocarditis as well as certain infiltrative cardiac diseases, such as amyloidosis, sarcoidosis, and haemochromatosis. In contrast, CMR is also required to identify patients with cardiomyopathy accurately, according to the Consensus Panel Report.² CMR is a non-invasive, accurate, and reproducible imaging technique that can be used to evaluate cardiac morphology and function, and provide valuable information for tissue characterization. In addition, several studies have suggested that CMR techniques using late-gadolinium enhancement (LGE) are useful for diagnosing various types of cardiomyopathies. Indeed, both CMR and EMB have demonstrated good performance in patients with troponin-positive acute chest pain but without coronary artery disease.³ However, there have been no reports directly comparing the diagnostic utility of CMR and EMB in patients with HF.

Therefore, we compared the diagnostic capability of CMR and EMB in HF patients and also assessed the diagnostic performance of the combined use of CMR and all clinical data in comparison with EMB alone.

Methods

Selection of patients

A total of 1034 consecutive patients with HF of unknown aetiology were evaluated between January 2007 and July 2009. Patients who were admitted to our institution for the management of HF, who had LV hypertrophy and/or LV dysfunction, and who had received EMB and LGE CMR were included in this study. Patients were excluded if they had one or more of the following conditions:

substantial valvular or ischaemic heart disease; congenital heart disease; constrictive pericarditis; idiopathic restrictive cardiomyopathy; an ambiguous final diagnosis; dilated cardiomyopathy (DCM) with an LVEF > 55%; poor-quality CMR; or an inadequate myocardial biopsy. Of the patients examined, 25 were given an ambiguous final diagnosis for the following reasons: 17 patients did not receive sufficient detailed investigations to reach the final diagnosis; 3 patients were suspected as having arrhythmogenic right ventricular cardiomyopathy (ARVC) but they did not fulfil the Task Force Criteria;⁴ 3 patients were suspected as having dilated hypertrophic cardiomyopathy (HCM) but hypertrophic stage was not detected; and 2 patients were diagnosed as LV non-compaction and we excluded these patients because EMB cannot diagnose this condition. As a result, we enrolled 136 patients in this study (Figure 1).

For all patients, a careful medical history was collected and physical examinations, laboratory tests, echocardiography, coronary angiography, and right heart catheterization were performed. EMB and CMR were also performed in all patients to evaluate evidence of HF. We identified the aetiology of HF using all possible diagnostic approaches in addition to EMB and CMR.

This study was approved by our Institutional Research Ethics Committee. The Committee decided that informed consent from the 136 subjects was not required according to the Japanese Clinical Research Guidelines because this was a retrospective, observational study. Instead, we made a public announcement as per the request of the Ethics Committee.

Aetiology of cardiomyopathy

According to the clinical data, echocardiogram, CMR, and EMB, six diagnoses were made for each patient, including (i) CMR diagnosis; (ii) EMB diagnosis; (iii) the combined diagnosis with clinical data




Figure 1 Study profile. Flow chart of the 1034 consecutive patients with heart failure of unknown aetiology admitted to our institution. The chart shows the immediate exclusion of cardiomyopathy due to significant coronary artery disease (CAD) and the further management of these patients. CMR, cardiac magnetic resonance; DCM, dilated cardiomyopathy; EMB, endomyocardial biopsy.

plus echocardiogram; (iv) the combined diagnosis with clinical data, echocardiogram, plus CMR; (v) the combined diagnosis with clinical data, echocardiogram, plus EMB; and (vi) the final diagnosis. The CMR and EMB diagnoses (i and ii) were established according to the results of CMR or EMB alone, and the investigators were blinded to all of the other data. Clinical data were defined as any method that could be used to diagnose HF other than echocardiography, CMR, or EMB, such as the collection of a patient's medical history, laboratory tests, scintigraphy, and coronary angiography. The final diagnosis (vi) was made prior to patient discharge by an expert team of cardiologists using all of the available data, including the results of EMB, CMR, and other diagnostic modalities. In addition, an expert team of cardiologists, who were not specialists in either CMR or EMB but could interpret these studies, was recruited. The final diagnoses were based on the recommendations of the 2008 European Society of Cardiology (ESC) report for the classification of cardiomyopathies.⁵ In patients with several causes of HF, the most significant cause was associated with the diagnosis.

Each diagnosis was assigned according to one of the following categories: DCM, HCM, hypertensive heart disease (HHD), ARVC, muscular dystrophy, infiltrative myocardial disease (i.e. amyloidosis and sarcoidosis), myocarditis, or other causes.

Cardiac magnetic resonance images and analysis

Images were acquired using a 1.5 T scanner (Sonata, Siemens Medical Solutions, Erlangen, Germany). The CMR protocol consisted of a cardiac functional study, spin-echo imaging, and LGE imaging, as previously described.⁶ For the cardiac functional study, three standard long-axis slices and a stack of contiguous short-axis slices (slice thickness, 6 mm; slice gap, 4 mm) were acquired as ECG-gated steady-state free-precession cine images with radial scans and breath-holding. T2-weighted spin-echo images were acquired using half-Fourier acquisition single shot turbo spin-echo (HASTE) before contrast injection with an echo time of 82 ms and fat saturation in the same position as the cine images. LGE images were acquired in the same positions as the cine images at 2, 5, 10, and 20 min after i.v. injection of 0.15 mmol/kg of gadolinium-diethyltriaminepentaacetic acid. The inversion delay time was 300 ms.

The cine and LGE images were evaluated by several observers who were blinded to the clinical data. The EF and volumes were measured quantitatively for the left and right ventricles according to the end-diastolic and end-systolic endocardial contours from a stack of short-axis cine images using ARGUS software. The LV mass (LVM) was calculated as the total myocardial volume multiplied by the specific gravity of the myocardium (1.05 g/mL). The ventricular end-diastolic and end-systolic volumes (EDV and ESV, respectively) and the LVM were standardized according to the body surface area (m²). The presence, location, and extent of LGE were determined using a standard 17 segment LV model.⁷ We classified the pattern of enhancement as subendocardial, midwall (longitudinal stripes), subepicardial, or transmural, as well as patchy (focal enhancement not following the coronary vascular territories) or diffuse.

Endomyocardial biopsy and analysis

Biopsy specimens were taken from the endocardium at the right inter-ventricular septum using Technowood disposable biopsy forceps (TONOKURA IKA KOGYO CO., LTD, Tokyo, Japan) via the right internal jugular vein or right femoral vein, as described elsewhere.⁸ Three to five specimens were obtained from each patient. No complications related to EMB were observed. Biopsy specimens were

immediately fixed in 15% formalin for 24 h, embedded in paraffin, and cut into 4 µm thick sections. The sections were stained with haematoxylin and eosin and Masson's trichrome. Some of the EMB specimens were frozen for polymerase chain reaction (PCR) analysis for the detection of enterovirus when myocarditis was suspected. Congo red staining was added when amyloidosis was suspected. Immunohistochemistry was performed in ARVC, myocarditis, amyloidosis, dystrophic cardiomyopathy, and some cases of HCM, as appropriate.

While EMB analysis at final diagnosis was made as above using all of the other data, EMB diagnosis was evaluated using only haematoxylin and eosin and Masson's trichrome stains by several cardiac pathologists who were not aware of the clinical features of the patients in this study.

Diagnosis of cardiomyopathy by cardiac magnetic resonance or endomyocardial biopsy

The diagnosis of cardiomyopathy by CMR and EMB was based on well-established and widely accepted definitions.^{9,10} A CMR diagnosis was made according to the dimensions, regional and global wall motion, wall thickness, and the presence and pattern of LGE,^{11–14} whereas an EMB diagnosis was made according to the report for classification of cardiomyopathies.^{4,5,15,16}

A histological diagnosis of DCM was performed by examining the following criteria: interstitial fibrosis, replacement fibrosis, inflammatory cell infiltrates, cellular hypertrophy, and myocardial cell degeneration.¹⁷ Histopathological criteria for HCM included severe myocyte hypertrophy, myocyte disarray > 10%, plexiform fibrosis, and nuclear hypertrophy. The diagnosis of HHD was made according to the presence of moderate myocyte hypertrophy,¹⁸ interstitial fibrosis, and the lack of myocyte disarray. The presence of non-caseating epithelioid granulomas with giant cells was considered indicative of cardiac sarcoidosis (CS).¹⁶ The diagnosis of myocarditis was based on the Dallas criteria modified by the Japanese Circulation Society Guidelines.¹⁹ Based on this modified version of the Dallas criteria, the immunohistochemistry was used to characterize the inflammatory infiltrates. The cut-off for mononuclear cell infiltrates was an inflammatory infiltrate count of at least 5/high power field. We confirmed the diagnoses of cardiac amyloidosis by electron microscopy and performed immunohistochemistry for amyloid typing. The histology diagnosis for ARVC was made according to the Task Force Criteria.⁴

The characteristics of DCM for CMR included dilation and impaired contraction of one or both ventricles and an LVEF < 55%.²⁰ Moreover, the wall thickness is normal or decreased. HCM is characterized by hypertrophy of the left ventricle and occasionally the right ventricle, normal or reduced LV volume, and normal LV contraction or hypercontraction. Apical HCM was regarded as hypertrophy of the apex, and hypertrophic obstructive cardiomyopathy (HOCM) was regarded as an obstruction to the LV outflow tract. We defined dilated HCM as an LVEF ≤ 50%^{21,22} and evidence of wall thickening prior to the study. Generally, dilated HCM is characterized by a relative wall thickness with a dilated LV cavity. LV hypertrophy is common in HHD, and additional common findings include a relative wall thickness with or without a dilated LV cavity. The use of CMR for CS can demonstrate certain characteristic features, such as septal thinning, ventricular dilatation, segmental systolic dysfunction, global systolic dysfunction, or ventricular aneurysm. We referred to the typical LGE pattern for diagnosis of DCM, HCM, HHD, and CS. The typical LGE pattern regarded a DCM LGE pattern as patchy or longitudinal midwall enhancement, a HCM LGE pattern as patchy and located at the LV–RV junction, a CS LGE pattern as a non-ischaemic pattern with enhancement of the midwall or epicardium at various sites, especially the anteroseptal

and inferolateral walls, and a HHD LGE pattern as similar to the DCM LGE pattern based upon a previous report.¹⁰ Myocarditis was diagnosed when subepicardial and midwall areas demonstrated an increased signal in the T2-weighted image or when the lateral and inferolateral walls demonstrated an LGE distribution in the epicardium toward the mid myocardial wall. ARVC is characterized by regional or global dysfunction, dilatation, and focal aneurysm of the right ventricle noted in the 2010 guideline,⁴ whereas amyloidosis is characterized by concentric hypertrophy with normal or reduced contractility, a thickened interatrial septum, bi-atrial dilation, and a circumferential pattern of LGE, preferentially involving the subendocardium but occasionally demonstrating a patchy transmural pattern. Dystrophic cardiomyopathy in LGE preserves the subendocardium and is more frequently located in the LV lateral wall.

Statistical analysis

Continuous variables were expressed as the mean \pm standard deviation (SD), whereas categorical variables were expressed as numbers and percentages. Comparisons between groups were performed using a two-sample *t*-test for normally distributed continuous variables and the Wilcoxon test for variables that did not demonstrate a normal distribution. For categorical variables, we used the χ^2 test and Fisher's exact test, as appropriate. For each type of cardiomyopathy, the sensitivity, specificity, diagnostic accuracy, positive predictive value (PPV), negative predictive value (NPV), and 95% confidence interval (CI) for the CMR diagnosis, EMB diagnosis, and combined diagnosis with clinical data, echocardiogram, plus CMR were calculated in comparison with the final diagnosis, which served as the gold standard. The PPV and NPV were computed using the following formulae: $PPV = \text{true positive}/(\text{true positive} + \text{false positive})$; and $NPV = \text{true negative}/(\text{true negative} + \text{false negative})$. Diagnostic accuracy was calculated using the following formulae: $\text{diagnostic accuracy} = (\text{true positive} + \text{true negative})/\text{total}$. A comparison of the diagnostic methods was performed using McNemar's test. The analyses were performed using JMP version 7 statistical software. All of the presented 95% CI are two.

Results

Study population and patient characteristics

A total of 136 patients were studied (Supplementary material, Table S1). The mean age of these patients was 52 ± 17 years (range 16–81 years): 83 of the patients were male, and 18 patients suffered from AF. EMB and CMR with LGE were performed in all patients, and none of the 136 patients was diagnosed with significant coronary artery disease. The most common diagnosis was DCM (54 patients, 40%), which was followed by HCM (36 patients, 26%). The remaining 46 patients were diagnosed with a secondary cardiomyopathy or HHD. The HCM patients included 4 cases of apical hypertrophy, 11 cases of HOCM, and 15 cases of HCM in the dilated phase.

The CMR results revealed asymmetric septal hypertrophy (septal/free wall thickness ratio ≥ 1.3) in 25 patients (18%), most of whom had either HCM (84%) or HHD (12%) detailed in the Supplementary material, Table S1.

The median patient follow-up period was 655 days (range 243–1143 days), and no diagnoses were changed during this time.

Comparison between cardiac magnetic resonance, endomyocardial biopsy, and the combined diagnosis

The sensitivity of EMB, CMR, and the combined diagnosis with clinical data plus echocardiogram, with clinical data, echocardiogram, plus CMR, and with clinical data, echocardiogram, plus EMB was 67, 79, 86, 97, and 100% relative to the final diagnosis, respectively. Table 1 shows the diagnostic performance of CMR, EMB, and the combined diagnosis with clinical data, echocardiogram, plus CMR. The use of CMR demonstrated a diagnostic capability comparable with EMB for all causes of HF. The highest level of sensitivity of EMB was for DCM (89%) followed by HCM (75%) and HHD (36%) (Table 2), whereas the greatest sensitivity of CMR was observed for DCM (83%) followed by HCM (81%) and CS (76%). Furthermore, to explore the relative merits of CMR vs. EMB, we investigated indications of EMB noted in the 2007 guidelines.²³ EMB demonstrated a better diagnostic yield for DCM and dilated HCM, whereas CMR demonstrated better diagnostic performance for cases of CS and HHD even when the indication for EMB was a class I. The diagnostic analysis is listed in Table 3. We gave six patients with dilated HCM an incorrect diagnosis of CS and also gave five patients with HHD an incorrect diagnosis of DCM using CMR. In contrast, we tended to misdiagnose CS and HHD as DCM and HCM as HHD when using EMB. Specifically, the six patients with HCM who were misdiagnosed for HHD by EMB included three patients with HOCM diagnoses, two with apical HCM diagnoses, and one with a diagnosis of dilated HCM. Table 2 shows the sensitivity and specificity for the use of EMB, CMR, and the combined diagnosis with clinical data, echocardiogram, plus CMR. Overall, CMR demonstrated increased specificity for DCM compared with EMB, and CMR also tended to be more sensitive for the diagnosis of CS and HHD. In contrast, EMB demonstrated lower sensitivity than CMR for most diagnoses, with the exception of DCM.

We also examined the diagnostic accuracy of CMR, EMB, and the combined diagnosis with clinical data, echocardiogram, plus CMR (Table 2). The sensitivity of the combined diagnosis with clinical data, echocardiogram, plus CMR was greater than that of EMB for the detection of HCM, CS, and HHD. The agreement of both CMR and EMB with a final diagnosis of DCM, HCM, CS, and HHD was noted to be 72, 58, 23, and 21%, respectively (Figure 2). Conversely, both CMR and EMB misdiagnosed 6, 3, 12, and 21% of patients with DCM, HCM, CS, and HHD, respectively. Importantly, all of the patients who received accurate diagnoses with EMB alone were also correctly diagnosed using the combined diagnosis with clinical data, echocardiogram, plus CMR.

Characteristics and details of cardiac magnetic resonance

We analysed the frequency of the use of the typical LGE pattern only in cases in which we diagnosed DCM, HCM, CS, and HHD. The CMR results revealed a DCM LGE pattern, HCM LGE pattern, CS LGE pattern, or HHD LGE pattern in 78, 53, 82, and 79% of the patients with DCM, HCM, CS, and HHD, respectively (Figure 3). In addition, the patients with typical LGE patterns were more likely to receive an accurate diagnosis. LGE in the papillary

Table 1 Agreement of endomyocardial biopsy, cardiac magnetic resonance, or combined diagnosis with clinical data, echocardiogram, plus cardiac magnetic resonance with final diagnosis in 136 patients based on endomyocardial biopsy indication

Final diagnoses, <i>n</i>	Number		EMB diagnosis, <i>n</i> (%)		CMR diagnosis, <i>n</i> (%)		Combined diagnosis, <i>n</i> (%)
	I	IIa/IIb	I	IIa/IIb	I	IIa/IIb	
DCM, 54	30	24	26 (87)	22 (92)	24 (80)	21 (88)	51 (94)
HCM, 36	11	25	7 (64)	20 (80)	6 (55)	23 (92)	35 (97)
Dilated HCM, 15	9	6	7 (78)	5 (83)	4 (44)	5 (83)	15 (100)
Obstructive HCM, 11	0	11	0 (0)	9 (82)	0 (0)	11 (100)	11 (100)
Apical HCM, 4	0	4	0 (0)	2 (50)	0 (0)	4 (100)	4 (100)
Sarcoidosis, 17	8	9	2 (25)	4 (44)	6 (75)	7 (78)	17 (100)
HHD, 14	5	9	2 (40)	3 (33)	5 (100)	4 (44)	14 (100)
Others							
ARVC, 5	1	4	0 (0)	0 (0)	1 (100)	4 (100)	5 (100)
Myocarditis, 4	4	0	3 (75)	0 (0)	1 (25)	0 (0)	4 (100)
Amyloidosis, 3	0	3	0 (0)	2 (67)	0 (0)	3 (100)	3 (100)
Dystrophic cardiomyopathy, 3	1	2	0 (0)	0 (0)	1 (100)	2 (100)	3 (100)
Total	60	76	40 (67)	51 (67)	44 (73)	64 (84)	132 (97)

ARVC, arrhythmogenic right ventricular cardiomyopathy; CMR, cardiac magnetic resonance; DCM, dilated cardiomyopathy; EMB, endomyocardial biopsy; HCM, hypertrophic cardiomyopathy; HHD, hypertensive heart disease.

muscle was frequently found in patients with HCM or sarcoidosis, while it was rarely or never seen in patients with DCM or HHD.

Discussion

This was the first study to compare the diagnostic performance of EMB and CMR in patients with HF. Non-invasive CMR, especially when combined with clinical data and echocardiogram, may provide an excellent diagnostic capacity for identifying the underlying aetiology in patients with HF, equal to or better than invasive EMB. Moreover, CMR is a powerful modality which in combination with clinical data including echocardiogram is sufficient for defining the pathophysiology of HF.

Although it is important to compare the diagnostic potential of EMB and CMR across a large number of patients with HF, comparisons using a large population have not been possible because it is extremely difficult to perform both EMB and CMR with LGE in sufficient patients. Our findings revealed that both the invasive EMB technique and the non-invasive CMR technique demonstrated good diagnostic performance (67% vs. 79%), whereas the use of CMR in combination with clinical data including echocardiogram unrelated to the EMB findings demonstrated excellent diagnostic performance (97%). Importantly, CMR alone could not surpass the diagnostic accuracy of EMB, which underscores the importance of EMB. However, the combined diagnosis was more accurate, which suggests that the use of CMR in combination with clinical data plus echocardiogram is the most reliable, non-invasive method for the diagnosis of HF in a routine clinical setting. Thus, we concluded that CMR is equal to or possibly superior to the use of EMB for the diagnosis of the underlying aetiology of HF,

especially in patients with sarcoidosis, HHD, others, and those with a class II indication for EMB.

These results suggest that CMR should be used more often than EMB for the initial diagnosis of HF. In addition, the cost of EMB is approximately three times greater than that of delayed enhancement CMR, and most patients can receive CMR at a clinic but would require a hospital stay to undergo EMB and perform EMB with right heart catheterization (which also contributes to the high cost of EMB). Indeed, the 2009 American College of Cardiology (ACC)/American Heart Association (AHA) chronic HF guidelines proposed that EMB should not be performed for the routine evaluation of patients with HF,²⁴ as EMB is often associated with sampling errors and complications. Therefore, although we do not deny the usefulness of EMB for the diagnosis of the underlying aetiology of HF, we suggest that CMR should be used more frequently for this type of diagnosis.

Diagnostic performance of cardiac magnetic resonance and endomyocardial biopsy

Although EMB provides suggestive findings in patients with DCM, HHD, and dystrophic cardiomyopathy, these findings are non-specific, and a definitive diagnosis cannot be made by EMB *per se*. In contrast, cardiac amyloidosis, CS, HCM, and myocarditis have specific histological characteristics and can be conclusively diagnosed using EMB alone if myocardial biopsy specimens contain these lesions (Figure 4). In our study, an accurate and conclusive diagnosis of such conditions could be reached using EMB alone in 38 out of 60 patients (Table 1), and we tended to misdiagnose CS and HHD as DCM, and HCM as HHD by EMB (Table 3).

Table 2 Sensitivity, specificity, positive predictive value, negative predictive value, accuracy of cardiac magnetic resonance diagnosis, and combined diagnoses with clinical data, echocardiogram, plus cardiac magnetic resonance vs. endomyocardial biopsy diagnosis

A. Cardiac magnetic resonance vs. endomyocardial biopsy										
	n	Sensitivity		rTPF	95% CI	n	Specificity		rFPF	95% CI
		CMR	EMB				CMR	EMB		
DCM	54	83%	89%	1.07	0.89–1.28	82	93%	69%	4.3	1.72–10.7
HCM	36	81%	75%	0.93	0.69–1.26	100	98%	94%	3	0.48–18.64
CS	17	76%	35%	0.46	0.20–1.07	119	92%	100%	0	–
HHD	14	64%	36%	0.56	0.22–1.43	122	96%	92%	2	0.59–6.81

B. Cardiac magnetic resonance vs. endomyocardial biopsy												
	n	PPV		rPPV	95% CI	n	NPV		rNPV	95% CI	Accuracy	
		CMR	EMB				CMR	EMB			CMR	EMB
DCM	54	88%	66%	0.75	0.60–0.93	82	89%	90%	1.01	0.89–1.15	89%	75%
HCM	36	94%	82%	0.87	0.71–1.08	100	93%	91%	0.98	0.89–1.07	93%	89%
CS	17	57%	100%	1.77	1.18–2.66	119	96%	92%	0.95	0.88–1.02	90%	92%
HHD	14	64%	33%	0.52	0.20–1.31	122	96%	93%	0.97	0.90–1.04	93%	86%

C. Combined procedure using clinical data with echocardiogram plus cardiac magnetic resonance vs. endomyocardial biopsy										
	n	Sensitivity		rTPF	95% CI	n	Specificity		rFPF	95% CI
		Combined	EMB				Combined	EMB		
DCM	54	94%	89%	0.94	0.82–1.08	82	100%	69%	–	–
HCM	36	97%	75%	0.77	0.61–0.97	100	99%	94%	6	0.54–67.19
CS	17	100%	35%	0.35	0.17–0.74	119	99%	100%	0	–
HHD	14	100%	36%	0.36	0.16–0.80	122	98%	92%	5	0.88–28.27

D. Combined procedure using clinical data with echocardiogram plus cardiac magnetic resonance vs. endomyocardial biopsy												
	n	PPV		rPPV	95% CI	n	NPV		rNPV	95% CI	Accuracy	
		Combined	EMB				Combined	EMB			Combined	EMB
DCM	54	100%	66%	0.65	0.52–0.81	82	96%	90%	0.94	0.85–1.03	98%	75%
HCM	36	97%	82%	0.84	0.69–1.02	100	99%	91%	0.92	0.86–0.99	99%	89%
CS	17	94%	100%	1.06	0.93–1.20	119	100%	92%	0.92	0.86–0.97	99%	92%
HHD	14	88%	33%	0.38	0.16–0.89	122	100%	93%	0.93	0.87–0.98	99%	86%

CI, confidence interval; CMR, cardiac magnetic resonance; CS, cardiac sarcoidosis; DCM, idiopathic dilated cardiomyopathy; EMB, endomyocardial biopsy; FPF, false positive fraction; HCM, hypertrophic cardiomyopathy; HHD, hypertensive heart disease; NPV, negative predictive value; PPV, positive predictive value; TPF, true positive fraction. 95% CIs were calculated according to the ratio of CMR diagnosis and combined diagnosis to EMB diagnosis. The bold values indicate a significant difference.

These misdiagnoses were attributed to non-specific changes in the biopsy specimens or the inappropriate sampling of sites separate from the lesions due to the patchy distribution of the lesions.^{17,25–27} However, there are merits in the classification of infiltrating inflammatory cells by either immunohistochemistry or a PCR method to guide treatment. The ESC 2012 guidelines also stated that the use of EMB may be needed to confirm the diagnosis in patients with suspected myocarditis, sarcoidosis, and amyloidosis.¹

Cardiac magnetic resonance is a safe procedure, and images of diagnostic quality can be obtained in $\geq 98\%$ of patients.²⁸ The use of CMR also allowed us to obtain detailed images of not

only functional and morphological abnormalities but also tissue pathology. In this study, EMB was superior to CMR for diagnoses of DCM and dilated HCM, whereas CMR demonstrated an improved diagnostic yield over EMB in cases of non-dilated HCM, CS, HHD, and other rare diseases (with the exception of myocarditis). Moreover, this tendency was the same independent of the EMB indication. In contrast to previous studies demonstrating a high level of sensitivity and specificity within only a limited study population, our study observed lower diagnostic agreement between methods. Because CMR was used to differentiate between a broad spectrum of diagnostic characteristics in HF patients, which resembles the clinical setting, CMR alone could

Table 3 Comparison of cardiac magnetic resonance diagnosis or endomyocardial biopsy diagnosis with final diagnosis

Final diagnosis	CMR diagnosis	DCM (n = 54)	HCM (n = 36)	CS (n = 17)	HHD (n = 14)	Others (n = 15)
DCM, n (%)		45 (83)	0	1	5 (36)	0
HCM, n (%)		1	29 (81)	1	0	0
CS, n (%)		2 (4)	6 (17)	13 (76)	0	2 (13)
HHD, n (%)		3 (5)	1	0	9 (64)	0
Others, n (%)		3 (5)	0	2 (12)	0	12 (80)

Final diagnosis	EMB diagnosis	DCM (n = 54)	HCM (n = 36)	CS (n = 17)	HHD (n = 14)	Others (n = 15)
DCM, n (%)		48 (89)	3 (8)	10 (59)	7 (50)	6 (40)
HCM, n (%)		3 (6)	27 (75)	0	2 (14)	1 (7)
CS, n (%)		0	0	6 (35)	0	0
HHD, n (%)		2 (4)	6 (17)	0	5 (36)	2 (13)
Others, n (%)		1	0	1	0	5 (33)

The bold values indicate diagnostic concordance between cardiac magnetic resonance diagnosis or endomyocardial biopsy diagnosis and final diagnosis.




Figure 2 Diagnostic capabilities of endomyocardial biopsy (EMB), cardiac magnetic resonance (CMR), and the combined diagnosis with CMR and EMB. DCM, idiopathic dilated cardiomyopathy; HCM, hypertrophic cardiomyopathy; CS, cardiac sarcoidosis; HHD, hypertensive heart disease. The ratio between EMB and CMR for the diagnosis of DCM, HCM, CS, and HHD.

not assign a correct diagnosis for 28 patients (21%) (Table 1). The use of CMR tended to misdiagnose HCM as CS, and HHD as DCM. Additionally, six HCM patients who were misdiagnosed with CS all had dilated HCM. A study by Hansen *et al.* suggested that the use of CMR in CS patients demonstrates similar results to those obtained in patients with HCM or idiopathic cardiomyopathy,²⁹ which is consistent with the present data (Table 3). In the five cases where HHD was misdiagnosed, they were consistently misdiagnosed as DCM due to the similarity of the images.¹⁰ However, in HCM patients, CMR demonstrated excellent

diagnostic performance (100%) for apical HCM and HOCM, which suggests that CMR has the ability to evaluate the heterogeneous appearance of HCM better than any other imaging modality,^{30,31} and this represents the main difference between CMR and EMB. We could not refer to the diagnostic accuracy of CMR in patients with myocarditis in our study because the number with myocarditis was too small. On the other hand, Marvorogeni *et al.* importantly concluded that both CMR and PCR prove useful for the detection of myocarditis, while CMR is important to detect the development of HF.³² Our data are consistent with the previous study³² showing that CMR and EMB have equivalent ability to reach the diagnosis and judge the pathophysiology.

Although the merits and demerits of CMR differ from those of EMB, its diagnostic capability was shown to be equivalent or even superior to that of EMB.

Superiority of the combined diagnosis

The combined diagnosis with non-invasive clinical data provides a sharp impact on an accurate diagnosis of HF.³³ Likewise, the combined diagnosis with clinical data, echocardiogram, plus CMR was shown to be very effective in the current study. Out of 54 DCM patients, 9 were misdiagnosed by CMR, but 6 of these 9 patients were correctly diagnosed using the combined diagnostic technique. Of 36 HCM patients, 7 were misdiagnosed by CMR, although 6 of these 7 patients were correctly assessed using the combined diagnosis (Tables 2 and 3). Moreover, the other misdiagnosed patients were also correctly assessed using the combined diagnosis.

Regarding the combined diagnosis with echocardiogram, the present study suggests that even the use of only clinical characteristics including echocardiogram can provide relatively high diagnostic performance compared with that in a previous report,³³ since our Department is specialized in diagnosing patients with non-ischaemic HF. However, the addition of CMR and EMB on top of the clinical information with echocardiogram increases the accuracy to 97% and 100%, respectively. Our original conclusion




Figure 3 Representative cardiac magnetic resonance (CMR) findings. (A) Dilated cardiomyopathy (DCM): midwall longitudinal thin late-gadolinium enhancement (LGE) in the anteroseptum wall without wall thickening (arrowheads). (B) Hypertensive heart disease (HHD): broad, ill-defined, and mild LGE in the midwall of the septum (arrowheads) with LV concentric hypertrophy. (C) Hypertrophic obstructive cardiomyopathy (HOCM): LGE in the LV–RV junctions of the anteroseptum and inferoseptum (arrowheads). Note: left atrial dilatation, LV asymmetric hypertrophy, papillary muscle hypertrophy, and LV outflow tract obstruction (arrow). (D) Dilated phase HCM: midwall patchy LGE in the anteroseptum and inferoseptum (arrowheads) and epicardial LGE in the anterior and posterior regions (arrow). (E) Cardiac sarcoidosis: subepicardial LGE of the anteroseptum with wall thinning and inferoseptum (arrowheads), subendocardial LGE of the lateral region (thin arrow), and LGE in the papillary muscle (thick arrow).

that CMR provides a diagnostic capability comparable with EMB seems to be true even with the clinical information including echocardiogram. Furthermore, all of the patients who were correctly diagnosed by EMB were correctly diagnosed using the combined technique, which indicates that the combined method was superior to the use of EMB in this study. Previous studies performed in populations with only one clinically suspected disease reported high diagnostic accuracy,^{11,12,34} and our results indicate that CMR would probably be available for a broad spectrum of HF patients, particularly those with a class II indication for EMB, for

differentiation between unknown aetiologies. Furthermore, the knowledge of diseases that are prone to misdiagnosis would increase the diagnostic performance for determining the aetiology of HF in a routine clinical setting. Although CMR is a non-invasive method, as is an echocardiogram, it is equal or superior to an echocardiogram because it can provide specific tissue characterization in addition to cardiac morphology and function. Accordingly, we suggest that it would be better initially to perform CMR in all patients, especially those with a class II indication for EMB, and diagnose the underlying aetiology of HF through the use of




Figure 4 Representative examples of histological findings from endomyocardial biopsies. (A) Dilated cardiomyopathy (DCM): the photomicrograph demonstrates replacement fibrosis (blue areas, black arrow) and moderate myocyte hypertrophy (Masson trichrome stain, bar = 50 μ m). (B) Hypertrophic cardiomyopathy (HCM): severe hypertrophy of myocytes, myocyte disarray, and bizarre nuclei are shown (Masson trichrome stain, bar = 50 μ m). (C) Cardiac sarcoidosis: non-caseating epithelioid granulomas with giant cells (white arrow) are shown (haematoxylin and eosin stain, bar = 30 μ m). (D) Cardiac amyloidosis: amorphous amyloid deposits (blue-grey) in the perimyocytes were consistent with amyloidosis in the interstitium of the myocardium (Masson's trichrome, bar = 50 μ m).

CMR and other non-invasive modalities. Then, if the combined diagnosis fails, EMB can be used as a second diagnostic modality.

Study limitations

This study had several limitations. First, all of the patients with HF of unknown aetiology were not assigned to receive both EMB and CMR; EMB was performed to reveal the underlying aetiology of HF, according to the Scientific Statement,²³ whereas CMR was performed in all patients without contraindications for CMR. Secondly, we included patients admitted to the Department with HF, and there were remarkably few patients in our Department who had coronary artery disease. However, even if such patients had been included, CMR would have probably been more useful to diagnose prior myocardial infarction due to spontaneous recanalization or coronary vasospasm than EMB. Thirdly, in the clinical setting, there are always cases with an ambiguous diagnosis despite a detailed investigation. We excluded these cases primarily on the premise that we would achieve a more precise diagnostic yield by avoiding these cases. Regarding EMB procedures, we took 3–5 biopsy specimens for each patient in our study, in accordance with the appropriate guidelines.²³ Additionally, all samples were taken from the right ventricle according to the protocol of our facility. In most patients, we took five samples to decrease sampling error, although the sampling number was decreased to three specimens in patients with both a pre-existing LBBB with a high risk for developing complete atrioventricular

block⁸ and obvious idiopathic DCM. The collection of samples from both ventricles may have increased the significance of the findings, but we collected the minimum requirement to decrease the procedural risk of EMB. Finally, this was also a retrospective study from a single centre. Our findings must be carefully interpreted and should be replicated in a prospective, large, multicentre investigation. Despite these limitations, our study has important strengths, such as the inclusion of a sufficient number of patients administered both EMB and CMR, a more precise final diagnosis using all available data, and broad clinical applications.

Supplementary material

Supplementary material is available at *European Journal of Heart Failure* online.

Funding

The Ministry of Health, Labor, and Welfare-Japan (grants-in-aid H23-Nanchi-Ippan-22 to M.K.), the Ministry of Education, Culture, Sports, Science and Technology-Japan (grants-in-aid 21390251 to M.K.), the Japan Heart Foundation (grants to M.K.), the Japan Cardiovascular Research Foundation (grants to M.K.).

Conflict of interest: none declared.

References

- McMurray JJ, Adamopoulos S, Anker SD, Auricchio A, Bohm M, Dickstein K, Falk V, Filippatos G, Fonseca C, Gomez-Sanchez MA, Jaarsma T, Kober L, Lip GY, Maggioni AP, Parkhomenko A, Pieske BM, Popescu BA, Ronnevik PK, Rutten FH, Schwitler J, Seferovic P, Stepinska J, Trindade PT, Voors AA, Zannad F, Zeheer A, Bax JJ, Baumgartner H, Ceconi C, Dean V, Deaton C, Fagard R, Funck-Brentano C, Hasdai D, Hoes A, Kirchhof P, Knuuti J, Kolh P, McDonagh T, Moulin C, Reiner Z, Sechtem U, Sirnes PA, Tendera M, Torbicki A, Vahanian A, Windecker S, Bonet LA, Avraamides P, Ben Lamin HA, Brignole M, Coca A, Cowburn P, Dargie H, Elliott P, Flachskampf FA, Guida GF, Hardman S, Jung B, Merkely B, Mueller C, Nanas JN, Nielsen OW, Orn S, Parissis JT, Ponikowski P. ESC guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: The Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC. *Eur J Heart Fail* 2012;**14**:803–869.
- Pennell DJ, Sechtem UP, Higgins CB, Manning WJ, Pohost GM, Rademakers FE, Rossum ACV, Shaw LJ, Yucel EK. Clinical indications for cardiovascular magnetic resonance (CMR): Consensus Panel report. *Eur Heart J* 2004;**25**:1940–1965.
- Baccouche H, Mahrholdt H, Meinhardt G, Merher R, Voehringer M, Hill S, Klingel K, Kandolf R, Sechtem U, Yilmaz A. Diagnostic synergy of non-invasive cardiovascular magnetic resonance and invasive endomyocardial biopsy in troponin-positive patients without coronary artery disease. *Eur Heart J* 2009;**30**:2869–2879.
- Marcus FI, McKenna WJ, Sherrill D, Basso C, Bauce B, Bluemke DA, Calkins H, Corrado D, Cox MG, Daubert JP, Fontaine G, Gear K, Hauer R, Nava A, Picard MH, Protonotarios N, Saffitz JE, Sanborn DM, Steinberg JS, Tandri H, Thiene G, Towbin JA, Tsatsopoulou A, Wichter T, Zareba W. Diagnosis of arrhythmogenic right ventricular cardiomyopathy/dysplasia: proposed modification of the Task Force Criteria. *Eur Heart J* 2010;**31**:806–814.
- Elliott P, Andersson B, Arbustini E, Bilinska Z, Cecchi F, Charron P, Dubourg O, Kuhl U, Maisch B, McKenna WJ, Monserrat L, Pankuweit S, Rapezzi C, Seferovic P, Tavazzi L, Keren A. Classification of the cardiomyopathies: a position statement from the European Society of Cardiology Working Group on Myocardial and Pericardial Diseases. *Eur Heart J* 2008;**29**:270–276.
- Kono AK, Yamada N, Higashi M, Kanzaki S, Hashimura H, Morita Y, Sakuma T, Noguchi T, Naito H, Sugimura K. Dynamic late gadolinium enhancement simply quantified using myocardium to lumen signal ratio: normal range of ratio and diffuse abnormal enhancement of cardiac amyloidosis. *J Magn Reson Imaging* 2011;**34**:50–55.
- Cerqueira MD, Weissman NJ, Dilsizian V, Jacobs AK, Kaul S, Laskey WK, Pennell DJ, Rumberger JA, Ryan T, Verani MS. Standardized myocardial segmentation and nomenclature for tomographic imaging of the heart: a statement for healthcare professionals from the Cardiac Imaging Committee of the Council on Clinical Cardiology of the American Heart Association. *Circulation* 2002;**105**:539–542.
- Holzmann M, Nicko A, Kuhl U, Noutsias M, Poller W, Hoffmann W, Morguet A, Witzensbichler B, Tschöpe C, Schultheiss HP, Pauschinger M. Complication rate of right ventricular endomyocardial biopsy via the femoral approach: a retrospective and prospective study analyzing 3048 diagnostic procedures over an 11-year period. *Circulation* 2008;**118**:1722–1728.
- Cunningham KS. An approach to endomyocardial biopsy interpretation. *J Clin Pathol* 2006;**59**:121–129.
- Karamitsos TD, Francis JM, Myerson S, Selvanayagam JB, Neubauer S. The role of cardiovascular magnetic resonance imaging in heart failure. *J Am Coll Cardiol* 2009;**54**:1407–1424.
- Patel MR, Cawley PJ, Heitner JF, Klem I, Parker MA, Jaroudi WA, Meine TJ, White JB, Elliott MD, Kim HW, Judd RM, Kim RJ. Detection of myocardial damage in patients with sarcoidosis. *Circulation* 2009;**120**:1969–1977.
- Vogelsberg H, Mahrholdt H, Deluigi CC, Yilmaz A, Kispert EM, Greulich S, Klingel K, Kandolf R, Sechtem U. Cardiovascular magnetic resonance in clinically suspected cardiac amyloidosis: noninvasive imaging compared to endomyocardial biopsy. *J Am Coll Cardiol* 2008;**51**:1022–1030.
- Abdel-Aty H, Boye P, Zagrosek A, Wassmuth R, Kumar A, Messroghli D, Bock P, Dietz R, Friedrich MG, Schulz-Menger J. Diagnostic performance of cardiovascular magnetic resonance in patients with suspected acute myocarditis: comparison of different approaches. *J Am Coll Cardiol* 2005;**45**:1815–1822.
- McCrohon JA, Moon JC, Prasad SK, McKenna WJ, Lorenz CH, Coats AJ, Pennell DJ. Differentiation of heart failure related to dilated cardiomyopathy and coronary artery disease using gadolinium-enhanced cardiovascular magnetic resonance. *Circulation* 2003;**108**:54–59.
- Magnani JW, Dec GW. Myocarditis: current trends in diagnosis and treatment. *Circulation* 2006;**113**:876–890.
- Diagnostic standard and guidelines for sarcoidosis. *Jpn J Sarcoidosis Granulomatous Disord* 2006;**26**:77–82. [in Japanese]
- Hughes SE. The pathology of hypertrophic cardiomyopathy. *Histopathology* 2004;**44**:412–427.
- Gradman AH, Alfayoumi F. From left ventricular hypertrophy to congestive heart failure: management of hypertensive heart disease. *Prog Cardiovasc Dis* 2006;**48**:326–341.
- JCS Joint Working Group. Guidelines for diagnosis and treatment of myocarditis (JCS 2009): digest version. *Circ J* 2011;**75**:734–743.
- Zimmermann O, Grebe O, Merkle N, Nusser T, Kochs M, Bienek-Ziolkowski M, Hombach V, Torzewski J. Myocardial biopsy findings and gadolinium enhanced cardiovascular magnetic resonance in dilated cardiomyopathy. *Eur J Heart Fail* 2006;**8**:162–166.
- Biagini E, Coccolo F, Ferlito M, Perugini E, Rocchi G, Bacchi-Reggiani L, Lofiego C, Boriani G, Prandstraller D, Picchio FM, Branzi A, Rapezzi C. Dilated-hypokinetic evolution of hypertrophic cardiomyopathy: prevalence, incidence, risk factors, and prognostic implications in pediatric and adult patients. *J Am Coll Cardiol* 2005;**46**:1543–1550.
- Harris KM, Spirito P, Maron MS, Zenovich AG, Formisano F, Lesser JR, Mackey-Bojack S, Manning WJ, Udelson JE, Maron BJ. Prevalence, clinical profile, and significance of left ventricular remodeling in the end-stage phase of hypertrophic cardiomyopathy. *Circulation* 2006;**114**:216–225.
- Cooper LT, Baughman KL, Feldman AM, Frustaci A, Jessup M, Kuhl U, Levine GN, Narula J, Starling RC, Towbin J, Virmani R. The role of endomyocardial biopsy in the management of cardiovascular disease: a scientific statement from the American Heart Association, the American College of Cardiology, and the European Society of Cardiology Endorsed by the Heart Failure Society of America and the Heart Failure Association of the European Society of Cardiology. *Eur Heart J* 2007;**28**:3076–3093.
- Hunt SA, Abraham WT, Chin MH, Feldman AM, Francis GS, Ganiats TG, Jessup M, Konstam MA, Mancini DM, Michl K, Oates JA, Rahko PS, Silver MA, Stevenson LW, Yancy CW. 2009 focused update incorporated into the ACC/AHA 2005 Guidelines for the Diagnosis and Management of Heart Failure in Adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines: developed in collaboration with the International Society for Heart and Lung Transplantation. *Circulation* 2009;**119**:e391–e479.
- Wu LA, Lapeyre AC 3rd, Cooper LT. Current role of endomyocardial biopsy in the management of dilated cardiomyopathy and myocarditis. *Mayo Clin Proc* 2001;**76**:1030–1038.
- Ardehali H, Howard DL, Hariri A, Qasim A, Hare JM, Baughman KL, Kasper EK. A positive endomyocardial biopsy result for sarcoid is associated with poor prognosis in patients with initially unexplained cardiomyopathy. *Am Heart J* 2005;**150**:459–463.
- Basso C, Ronco F, Marcus F, Abudurehman A, Rizzo S, Frigo AC, Bauce B, Maddalena F, Nava A, Corrado D, Grigoletto F, Thiene G. Quantitative assessment of endomyocardial biopsy in arrhythmogenic right ventricular cardiomyopathy/dysplasia: an *in vitro* validation of diagnostic criteria. *Eur Heart J* 2008;**29**:2760–2771.
- Bruder O, Schneider S, Nothnagel D, Dill T, Hombach V, Schulz-Menger J, Nagel E, Lombardi M, van Rossum AC, Wagner A, Schwitler J, Senges J, Sabin GV, Sechtem U, Mahrholdt H. EuroCMR (European Cardiovascular Magnetic Resonance) registry: results of the German pilot phase. *J Am Coll Cardiol* 2009;**54**:1457–1466.
- Hansen MW, Merchant N. MRI of hypertrophic cardiomyopathy: part 2, differential diagnosis, risk stratification, and posttreatment MRI appearances. *AJR Am J Roentgenol* 2007;**189**:1344–1352.
- Rickers C, Wilke NM, Jerosch-Herold M, Casey SA, Panse P, Panse N, Weil J, Zenovich AG, Maron BJ. Utility of cardiac magnetic resonance imaging in the diagnosis of hypertrophic cardiomyopathy. *Circulation* 2005;**112**:855–861.
- Moon JC, Fisher NG, McKenna WJ, Pennell DJ. Detection of apical hypertrophic cardiomyopathy by cardiovascular magnetic resonance in patients with non-diagnostic echocardiography. *Heart* 2004;**90**:645–649.
- Mavrogeni S, Spargias C, Bratis C, Kolovou G, Markussis V, Papadopolou E, Constadoulakis P, Papadimitropoulos M, Douskou M, Pavlides G, Cokkinos D. Myocarditis as a precipitating factor for heart failure: evaluation and 1-year follow-up using cardiovascular magnetic resonance and endomyocardial biopsy. *Eur J Heart Fail* 2011;**13**:830–837.
- Ardehali H, Qasim A, Cappola T, Howard D, Hruban R, Hare JM, Baughman KL, Kasper EK. Endomyocardial biopsy plays a role in diagnosing patients with unexplained cardiomyopathy. *Am Heart J* 2004;**147**:919–923.
- Friedrich MG, Sechtem U, Schulz-Menger J, Holmvang G, Alakija P, Cooper LT, White JA, Abdel-Aty H, Gutberlet M, Prasad S, Aletras A, Laissy JP, Paterson I, Filipchuk NG, Kumar A, Pauschinger M, Liu P. Cardiovascular magnetic resonance in myocarditis: a JACC White Paper. *J Am Coll Cardiol* 2009;**53**:1475–1487.

Circulation Research

JOURNAL OF THE AMERICAN HEART ASSOCIATION



Activation of Natural Killer T Cells Ameliorates Postinfarct Cardiac Remodeling and Failure in Mice

Mochamad Ali Sobirin, Shintaro Kinugawa, Masashige Takahashi, Arata Fukushima, Tsuneaki Homma, Taisuke Ono, Kagami Hirabayashi, Tadashi Suga, Putri Azalia, Shingo Takada, Masaru Taniguchi, Toshinori Nakayama, Naoki Ishimori, Kazuya Iwabuchi and Hiroyuki Tsutsui

Circ Res. 2012;111:1037-1047; originally published online August 10, 2012;
doi: 10.1161/CIRCRESAHA.112.270132

Circulation Research is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231
Copyright © 2012 American Heart Association, Inc. All rights reserved.
Print ISSN: 0009-7330. Online ISSN: 1524-4571

The online version of this article, along with updated information and services, is located on the
World Wide Web at:

<http://circres.ahajournals.org/content/111/8/1037>

Data Supplement (unedited) at:

<http://circres.ahajournals.org/content/suppl/2012/08/10/CIRCRESAHA.112.270132.DC1.html>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation Research* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the Permissions and Rights Question and Answer document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Circulation Research* is online at:
<http://circres.ahajournals.org/subscriptions/>