

Table 1

Clinical Sign - Summary

B120719

Day numbers relative to Start Date

Group	Sex	Clinical Sign	-4	-3	-2	-1
			a.m.	a.m.	a.m.	a.m.
1	m	ANIMALS ALIVE	3	3	3	3
		ANIMALS NORMAL	3	3	3	3
2	m	ANIMALS ALIVE	3	3	3	3
		ANIMALS NORMAL	3	3	3	3
3	m	ANIMALS ALIVE	3	3	3	3
		ANIMALS NORMAL	3	3	3	3
4	m	ANIMALS ALIVE	3	3	3	3
		ANIMALS NORMAL	3	3	3	3

- 51 -

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg Group 4 - 500 mg/kg

Time 10 , 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Table 1

Clinical Sign - Summary

B120719

		Day numbers relative to Start Date												
Group	Sex	Clinical Sign	1	1	2	2	3	3	4	4	5	5	6	6
			Before dosing	Time 10	Before dosing	Time 10	Before dosing	Time 10	Before dosing	Time 10	Before dosing	Time 10	Before dosing	Time 10
1	m	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	3	3	3	3	3	3	3	3	3	3	3
2	m	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	1	1	1	1	0	3	0	3	0	3	0
		Vomitus (Dosing formulation)	3	.	.	.	3
		Vomitus (Food)	.	.	2	.	2
		Vomitus (Frothy material)
		Vomitus (Dosing solution)	.	2	.	2	.	2	.	.	.	3	.	.
		Diarrhea	1
3	m	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	1	2	1	1	0	2	0	2	0	2	0
		Death
		Moribundity
		Decrease in locomotor activity
		Vomitus (Dosing formulation)	3	.	.	.	3
		Vomitus (Food)	.	.	1	.	2	.	1
		Vomitus (Frothy material)	.	.	1
		Vomitus (Dosing solution)	.	2	.	2	.	3	.	.	.	3	.	.
		Loose stool	1	1	1
		Diarrhea	1	1	.	.	1	.	.	.
No-stool	1	.		

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg

Time 10, 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Table 1

Clinical Sign - Summary

B120719

		Day numbers relative to Start Date												
Group	Sex	Clinical Sign	7	7	8	8	9	9	10	10	11	11	12	12
			Before dosing	Time 10	Before dosing	Time 10	Before dosing	Time 10	Before dosing	Time 10	Before dosing	Time 10	Before dosing	Time 10
1	m	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	3	3	3	3	3	3	3	3	3	3	3
2	m	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	0	3	0	3	2	2	2	2	0	2	1
		Vomitus (Dosing formulation)
		Vomitus (Food)	1	.	1	3	1	.
		Vomitus (Frothy material)	.	.	.	1	.	1
		Vomitus (Dosing solution)	.	3	.	2	.	.	.	1	.	.	.	2
		Diarrhea
3	m	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	1	0	2	0	2	2	1	1	2	0	2	2
		Death
		Moribundity
		Decrease in locomotor activity
		Vomitus (Dosing formulation)
		Vomitus (Food)	1	.	.	.	1	.	1	.	1	2	1	.
		Vomitus (Frothy material)	1	1
		Vomitus (Dosing solution)	.	3	.	3	.	.	.	2	.	1	.	1
		Loose stool
		Diarrhea	1	1	1	.
No-stool	.	.	1		

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg

Time 10, 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Table 1 Clinical Sign - Summary

B120719

		Day numbers relative to Start Date					
Group	Sex	Clinical Sign	13 Before dosing	13 Time 10	14 Before dosing	14 Time 10	15 a.m.
1	m	ANIMALS ALIVE	3	3	3	3	3
		ANIMALS NORMAL	3	3	3	3	3
2	m	ANIMALS ALIVE	3	3	3	3	3
		ANIMALS NORMAL	3	1	3	2	3
		Vomitus (Dosing formulation)
		Vomitus (Food)
		Vomitus (Frothy material)
		Vomitus (Dosing solution)	.	2	.	1	.
		Diarrhea
3	m	ANIMALS ALIVE	3	3	2	2	0
		ANIMALS NORMAL	1	1	0	0	.
		Death	.	.	1	.	.
		Moribundity	.	.	.	2	.
		Decrease in locomotor activity	.	.	.	1	.
		Vomitus (Dosing formulation)
		Vomitus (Food)
		Vomitus (Frothy material)
		Vomitus (Dosing solution)	1	2	.	1	.
		Loose stool
		Diarrhea	1	.	1	.	.
No-stool	1	.	1	.	.		

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg

Time 10, 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Table 1

Clinical Sign - Summary

B120719

Group	Sex	Clinical Sign	Day numbers relative to Start Date											
			1 Before dosing	1 Time 10	2 Before dosing	2 Time 10	3 Before dosing	3 Time 10	4 Before dosing	4 Time 10	5 Before dosing	5 Time 10	6 Before dosing	6 Time 10
4	m	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	
		ANIMALS NORMAL	3	0	1	0	1	0	1	0	1	0	2	0
		Death	
		Lateral position	
		Vomitus (Dosing formulation)	2	.	.	.	2	
		Vomitus (Food)	.	.	1	1	.	.	.	
		Vomitus (Frothy material)	.	.	2	.	.	1	
		Vomitus (Dosing solution)	.	3	.	3	.	3	.	.	3	.	.	
		Loose stool	.	.	1	.	1	.	1	.	1	.	.	
		Diarrhea	.	.	.	1	1	.	1	1	.	1	1	
		Watery diarrhea	1	
		No-stool	1	.	.	.	

Group 4 - 500 mg/kg

Time 10 , 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Table 1

Clinical Sign - Summary

B120719

Group	Sex	Clinical Sign	Day numbers relative to Start Date								
			7 Before dosing	7 Time 10	8 Before dosing	8 Time 10	9 Before dosing	9 Time 10	10 Before dosing	10 Time 10	11 Before dosing
4	m	ANIMALS ALIVE	2	2	2	2	2	2	2	2	2
		ANIMALS NORMAL	2	0	0	0	0	1	0	1	0
		Death	1
		Lateral position	1
		Vomitus (Dosing formulation)
		Vomitus (Food)	2	.	1	.	.
		Vomitus (Frothy material)	1	1	.	2
		Vomitus (Dosing solution)	1	2	.	2	.	.	.	1	.
		Loose stool
		Diarrhea	1	1	1	1	1	.	.	.	1
		Watery diarrhea
		No-stool	.	.	1	.	.	.	1	.	.

Group 4 - 500 mg/kg

Time 10 , 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Table 1

Clinical Sign - Summary

B120719

Group	Sex	Clinical Sign	Day numbers relative to Start Date											
			-52 a.m.	-51 a.m.	-50 a.m.	-49 a.m.	-48 a.m.	-47 a.m.	-46 a.m.	-45 a.m.	-44 a.m.	-43 a.m.	-42 a.m.	-41 a.m.
1	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	2	2	2	2	3	3	3	3	3	3	3
		Menses	.	1	1	1	1
2	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	3	2	2	2	2	3	3	3	3	3	3
		Menses	.	.	1	1	1	1
3	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	3	3	3	3	3	3	3	3	3	3	3
		Menses
4	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	3	3	3	3	3	3	3	3	3	3	3
		Menses

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg Group 4 - 500 mg/kg

Time 10 , 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Table 1

Clinical Sign - Summary

B120719

		Day numbers relative to Start Date												
		-40	-39	-38	-37	-36	-35	-34	-33	-32	-31	-30	-29	
		a.m.	a.m.	a.m.	a.m.	a.m.	a.m.	a.m.	a.m.	a.m.	a.m.	a.m.	a.m.	
Group	Sex	Clinical Sign												
1	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	3	3	3	3	3	3	3	3	3	3	3
		Menses
2	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	3	3	3	3	3	3	3	3	3	3	3
		Menses
3	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	3	3	3	3	3	3	3	3	3	3	3
		Menses
4	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	3	3	3	3	2	2	2	3	3	2	3
		Menses	1	1	1	.	.	1	.

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg Group 4 - 500 mg/kg

Time 10 , 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Table 1

Clinical Sign - Summary

B120719

Group	Sex	Clinical Sign	Day numbers relative to Start Date											
			-28 a.m.	-27 a.m.	-26 a.m.	-25 a.m.	-24 a.m.	-23 a.m.	-22 a.m.	-21 a.m.	-20 a.m.	-19 a.m.	-18 a.m.	-17 a.m.
1	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	3	3	3	3	3	2	2	2	3	3	3
		Menses	1	1	1	.	.	.
2	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	3	3	3	3	3	3	3	3	3	3	3
		Menses
3	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	3	3	3	3	3	3	3	3	3	3	3
		Menses
4	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	2	3	3	3	3	3	3	2	2	3	3	3
		Menses	1	1	1	.	.	.

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg Group 4 - 500 mg/kg

Time 10 , 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Table 1

Clinical Sign - Summary

B120719

		Day numbers relative to Start Date												
		-16	-15	-14	-13	-12	-11	-10	-9	-8	-7	-6	-5	
		a.m.	a.m.	a.m.	a.m.	a.m.	a.m.	a.m.	a.m.	a.m.	a.m.	a.m.	a.m.	
Group	Sex	Clinical Sign												
1	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	3	3	3	3	3	3	3	3	3	3	3
		Menses
2	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	3	3	3	3	3	3	3	3	3	3	3
		Menses
3	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	3	3	3	3	3	3	3	3	3	3	3
		Menses
4	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3
		ANIMALS NORMAL	3	3	3	3	3	3	3	3	3	3	3	3
		Menses

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg Group 4 - 500 mg/kg

Time 10 , 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Table 1

Clinical Sign - Summary

B120719

		Day numbers relative to Start Date				
Group	Sex	Clinical Sign	-4 a.m.	-3 a.m.	-2 a.m.	-1 a.m.
1	f	ANIMALS ALIVE	3	3	3	3
		ANIMALS NORMAL	3	3	3	3
		Menses
2	f	ANIMALS ALIVE	3	3	3	3
		ANIMALS NORMAL	3	2	1	1
		Menses	.	1	2	2
3	f	ANIMALS ALIVE	3	3	3	3
		ANIMALS NORMAL	3	3	3	3
		Menses
4	f	ANIMALS ALIVE	3	3	3	3
		ANIMALS NORMAL	3	3	3	3
		Menses

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg Group 4 - 500 mg/kg

Time 10 , 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Table 1

Clinical Sign - Summary

B120719

		Day numbers relative to Start Date													
Group	Sex	Clinical Sign	1	1	2	2	3	3	4	4	5	5	6	6	
			Before dosing	Time 10	Before dosing	Time 10	Before dosing	Time 10	Before dosing	Time 10	Before dosing	Time 10	Before dosing	Time 10	
1	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3	
		ANIMALS NORMAL	3	3	3	3	3	3	3	3	3	3	3	2	
		Menses	1
2	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3	
		ANIMALS NORMAL	3	0	3	0	3	2	3	0	3	0	2	0	
		Moribundity
		Decrease in locomotor activity
		Vomitus (Dosing formulation)	3	3
		Vomitus (Food)
		Vomitus (Frothy material)
		Vomitus (Dosing solution)	.	3	.	3	.	1	3	.	.
		Menses	1	1
		Diarrhea
No-stool		
3	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3	
		ANIMALS NORMAL	3	0	2	0	1	0	2	0	2	0	3	0	
		Death
		Vomitus (Dosing formulation)	3	3
		Vomitus (Food)	1
		Vomitus (Frothy material)
		Vomitus (Dosing solution)	.	3	.	3	.	3	3	.	.
		Loose stool
Diarrhea	.	.	1	.	2	.	1	1	1		

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg

Time 10, 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Table 1

Clinical Sign - Summary

B120719

		Day numbers relative to Start Date													
Group	Sex	Clinical Sign	7	7	8	8	9	9	10	10	11	11	12	12	
			Before dosing	Time 10	Before dosing	Time 10	Before dosing	Time 10	Before dosing	Time 10	Before dosing	Time 10	Before dosing	Time 10	
1	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	3	3	
		ANIMALS NORMAL	2	2	2	2	3	3	3	3	3	3	3	3	
		Menses	1	1	1	1	
2	f	ANIMALS ALIVE	3	3	3	3	3	3	3	2	2	2	2	2	
		ANIMALS NORMAL	1	1	1	1	1	3	1	1	2	0	0	0	
		Moribundity	1	
		Decrease in locomotor activity	1	
		Vomitus (Dosing formulation)	
		Vomitus (Food)	1	
		Vomitus (Frothy material)	1	.	1	
		Vomitus (Dosing solution)	.	1	.	2	.	.	.	1	.	2	.	.	1
		Menses	1	1	1	1
		Diarrhea	1	.	1	.	1	1	1
		No-stool	1	.	.	.	1	.	
3	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	2	2	
		ANIMALS NORMAL	3	0	3	0	1	3	1	0	2	0	2	0	
		Death	1	.	
		Vomitus (Dosing formulation)	
		Vomitus (Food)	1	.	1	.	2	.	.	.	
		Vomitus (Frothy material)	1	
		Vomitus (Dosing solution)	.	3	.	3	1	.	.	3	.	1	.	1	
		Loose stool	1	
Diarrhea	1	.	.	1			

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg

Time 10 , 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Table 1

Clinical Sign - Summary

B120719

		Day numbers relative to Start Date					
Group	Sex	Clinical Sign	13 Before dosing	13 Time 10	14 Before dosing	14 Time 10	15 a.m.
1	f	ANIMALS ALIVE	3	3	3	3	3
		ANIMALS NORMAL	3	3	3	3	3
		Menses
2	f	ANIMALS ALIVE	2	2	2	2	2
		ANIMALS NORMAL	0	1	2	2	0
		Moribundity
		Decrease in locomotor activity
		Vomitus (Dosing formulation)
		Vomitus (Food)
		Vomitus (Frothy material)
		Vomitus (Dosing solution)	.	1	.	.	.
		Menses
		Diarrhea	1
No-stool	1	.	.	.	2		
3	f	ANIMALS ALIVE	2	2	2	2	2
		ANIMALS NORMAL	1	0	1	0	1
		Death
		Vomitus (Dosing formulation)
		Vomitus (Food)	.	2	.	.	.
		Vomitus (Frothy material)
		Vomitus (Dosing solution)	.	2	.	2	.
		Loose stool
		Diarrhea	1	.	1	.	1

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg

Time 10, 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Table 1

Clinical Sign - Summary

B120719

Group	Sex	Clinical Sign	Day numbers relative to Start Date											
			1 Before dosing	1 Time 10	2 Before dosing	2 Time 10	3 Before dosing	3 Time 10	4 Before dosing	4 Time 10	5 Before dosing	5 Time 10	6 Before dosing	6 Time 10
4	f	ANIMALS ALIVE	3	3	3	3	3	3	3	3	3	3	2	2
		ANIMALS NORMAL	2	0	0	0	1	0	2	0	0	0	0	0
		Moribundity	1
		Lateral position	1
		Vomitus (Dosing formulation)	3	1
		Vomitus (Food)	.	.	1	.	1
		Vomitus (Frothy material)	.	.	2	.	.	1
		Vomitus (Dosing solution)	.	3	.	3	.	3	.	.	2	.	.	.
		Menses	1	1	1	1	1	1	1	1	1	.	1	1
		Loose stool	.	.	1	1
		Diarrhea	2	.	1	1	2	1	2	1
		No-stool	1

Group 4 - 500 mg/kg

Time 10 , 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Table 1

Clinical Sign - Summary

B120719

		Day numbers relative to Start Date									
Group	Sex	Clinical Sign	7	7	8	8	9	9	10	10	11
			Before dosing	Time 10	Before dosing	Time 10	Before dosing	Time 10	Before dosing	Time 10	Before dosing
4	f	ANIMALS ALIVE	2	2	2	1	1	1	1	1	1
		ANIMALS NORMAL	0	1	0	0	0	0	0	0	1
		Moribundity	.	.	1
		Lateral position	.	.	1
		Vomitus (Dosing formulation)
		Vomitus (Food)	.	.	1	.	1	.	1	.	.
		Vomitus (Frothy material)	1	.	.	.
		Vomitus (Dosing solution)	.	1	.	1	1	.	.	1	.
		Menses	1	1	1
		Loose stool	1	.	.
		Diarrhea	1	.	2	.	.	1	.	.	.
		No-stool

Group 4 - 500 mg/kg

Time 10 , 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Table 2 Body Weight - Summary

B120719

Bodyweight (kg)

Sex: Male		Day(s) Relative to Start Date							
		-52	-45	-42	-35	-28	-21	-14	-7
0 mg/kg	Mean	3.00	2.93	2.93	3.07	3.10	3.27	3.23	3.33
	SD	0.20	0.21	0.21	0.15	0.17	0.15	0.12	0.12
	N	3	3	3	3	3	3	3	3
50 mg/kg	Mean	3.07	3.07	3.10	3.23	3.20	3.33	3.37	3.40
	SD	0.29	0.29	0.26	0.23	0.26	0.32	0.29	0.35
	N	3	3	3	3	3	3	3	3
150 mg/kg	Mean	3.03	3.10	3.10	3.20	3.20	3.30	3.33	3.40
	SD	0.21	0.20	0.20	0.20	0.20	0.20	0.21	0.20
	N	3	3	3	3	3	3	3	3
500 mg/kg	Mean	3.00	3.00	2.97	3.07	3.07	3.23	3.20	3.27
	SD	0.46	0.40	0.35	0.45	0.45	0.50	0.50	0.50
	N	3	3	3	3	3	3	3	3

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Table 2 Body Weight - Summary

B120719

Bodyweight (kg)

Sex: Male		Day(s) Relative to Start Date					
		1	4	7	8	11	14
0 mg/kg	Mean	3.27	3.27	-	3.23	3.33	3.27
	SD	0.15	0.15	-	0.15	0.15	0.15
	N	3	3	-	3	3	3
50 mg/kg	Mean	3.43	3.40	-	3.33	3.23	3.17
	SD	0.32	0.44	-	0.40	0.35	0.25
	N	3	3	-	3	3	3
150 mg/kg	Mean	3.37	3.27	-	3.10	2.97	2.83
	SD	0.15	0.25	-	0.20	0.21	0.25
	N	3	3	-	3	3	3
500 mg/kg	Mean	3.27	3.10	2.50	3.20	3.00	-
	SD	0.50	0.60	-	0.42	0.28	-
	N	3	3	1	2	2	-

[Day 7 : Final Body Weight.]

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Table 2 Body Weight - Summary

B120719

Bodyweight (kg)

Sex: Female		Day(s) Relative to Start Date							
		-52	-45	-42	-35	-28	-21	-14	-7
0 mg/kg	Mean	3.00	2.97	2.97	3.03	3.03	3.13	3.13	3.13
	SD	0.36	0.31	0.31	0.25	0.25	0.25	0.32	0.32
	N	3	3	3	3	3	3	3	3
50 mg/kg	Mean	2.87	2.87	2.87	2.97	2.97	3.00	3.00	3.07
	SD	0.21	0.21	0.12	0.12	0.12	0.17	0.17	0.12
	N	3	3	3	3	3	3	3	3
150 mg/kg	Mean	2.77	2.73	2.77	2.83	2.80	2.93	2.90	2.90
	SD	0.25	0.21	0.15	0.21	0.26	0.31	0.30	0.30
	N	3	3	3	3	3	3	3	3
500 mg/kg	Mean	2.97	2.87	2.87	2.93	2.93	2.93	2.93	2.97
	SD	0.25	0.25	0.15	0.12	0.12	0.12	0.12	0.06
	N	3	3	3	3	3	3	3	3

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Table 2 Body Weight - Summary

B120719

Bodyweight (kg)

Sex: Female		Day(s) Relative to Start Date							
		1	4	5	8	10	11	12	14
0 mg/kg	Mean	3.17	3.13	-	3.07	-	3.17	-	3.07
	SD	0.38	0.42	-	0.47	-	0.38	-	0.38
	N	3	3	-	3	-	3	-	3
50 mg/kg	Mean	3.07	3.00	-	2.80	2.60	2.80	-	2.60
	SD	0.15	0.20	-	0.20	-	0.14	-	0.14
	N	3	3	-	3	1	2	-	2
150 mg/kg	Mean	2.90	2.83	-	2.77	-	2.70	2.80	2.40
	SD	0.30	0.31	-	0.32	-	0.26	-	0.28
	N	3	3	-	3	-	3	1	2
500 mg/kg	Mean	2.97	2.77	2.50	2.70	-	2.90	-	-
	SD	0.12	0.15	-	0.14	-	-	-	-
	N	3	3	1	2	-	1	-	-

[Day 5, Day 7 and 12 : Final Body Weight.]

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]