

Group	Sex	Animal	Clinical Sign	Site	Day numbers relative to Start Date										
					10 Before dosing	10 Time 10	11 Before dosing	11 Time 10	12 Before dosing	12 Time 10	13 Before dosing	13 Time 10	14 Before dosing		
2	f	50201	No Abnormalities Detected		
			Moribundity	X	
			Decrease in locomotor activity	1	
			Vomitus (Dosing formulation)	
			Vomitus (Frothy material)	
			Vomitus (Dosing solution)	
		50202	Diarrhea
			No-stool	X	
			No Abnormalities Detected	X	.	X	X	
			Vomitus (Dosing formulation)	
			Vomitus (Dosing solution)	.	X	.	X	.	X	.	X	.	X	.	
			No-stool	X	.	X	.	X	.	.	
		50203	No Abnormalities Detected	.	X	X	X	X	
			Vomitus (Dosing formulation)	
			Vomitus (Food)	
			Vomitus (Frothy material)	X	
			Vomitus (Dosing solution)	.	.	.	X	
			Menses	
Diarrhea	X	X	X	X	X	.	.				
No-stool				

- 231 -

Severity Codes: X = Present; 1 = Slight

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg Group 4 - 500 mg/kg

Time 10 ; 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Day numbers relative to Start Date

Group	Sex	Animal	Clinical Sign	Site	14	15
					Time 10	a.m.
2	f	50201	No Abnormalities Detected		.	.
			Moribundity		.	.
			Decrease in locomotor activity		.	.
			Vomitus (Dosing formulation)		.	.
			Vomitus (Frothy material)		.	.
		Vomitus (Dosing solution)		.	.	
		Diarrhea		.	.	
		No-stool		.	.	
		50202	No Abnormalities Detected		X	.
			Vomitus (Dosing formulation)		.	.
			Vomitus (Dosing solution)		.	.
			No-stool		.	X
		50203	No Abnormalities Detected		X	.
			Vomitus (Dosing formulation)		.	.
			Vomitus (Food)		.	.
Vomitus (Frothy material)			.	.		
Vomitus (Dosing solution)			.	.		
Menses			.	.		
Diarrhea			.	.		
No-stool			.	X		

Severity Codes: X = Present; 1 = Slight

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg Group 4 - 500 mg/kg

Time 10, 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Group	Sex	Animal	Clinical Sign	Site	Day numbers relative to Start Date										
					1 Before dosing	1 Time 10	2 Before dosing	2 Time 10	3 Before dosing	3 Time 10	4 Before dosing	4 Time 10	5 Before dosing		
3	f	50301	No Abnormalities Detected		X
			Vomitus (Dosing formulation)	1	.	.	
			Vomitus (Food)	X	
			Vomitus (Dosing solution)	.	X	.	X	.	X	.	X	.	.	.	
		50302	No Abnormalities Detected	X	.	X	.	X	.	X	.	X	.	X	
			Vomitus (Dosing formulation)	1	.	.	
			Vomitus (Food)	
		50303	No Abnormalities Detected	X	.	X	X	.	X	
			Death	
			Vomitus (Dosing formulation)	1	.	.	
			Vomitus (Food)	
			Vomitus (Frothy material)	
			Vomitus (Dosing solution)	.	X	.	X	.	X	
			Loose stool	
			Diarrhea	X	

Severity Codes: X = Present; 1 = Slight

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg Group 4 - 500 mg/kg

Time 10, 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Group	Sex	Animal	Clinical Sign	Site	Day numbers relative to Start Date										
					5 Time 10	6 Before dosing	6 Time 10	7 Before dosing	7 Time 10	8 Before dosing	8 Time 10	9 Before dosing	9 Time 10		
3	f	50301	No Abnormalities Detected		.	X	.	X	.	X	.	X	.	X	
			Vomitus (Dosing formulation)		.	.	1
			Vomitus (Food)	
			Vomitus (Dosing solution)		X	.	.	.	X	.	X
		50302	Diarrhea	
			No Abnormalities Detected		.	X	.	X	.	X	X
			Vomitus (Dosing formulation)		.	.	1
		50303	Vomitus (Food)	
			Vomitus (Dosing solution)		X	.	.	.	X	.	X	.	X	.	.
			No Abnormalities Detected		.	X	.	X	.	X	X
			Death	
			Vomitus (Dosing formulation)		.	.	1
			Vomitus (Food)		X	.	.
			Vomitus (Frothy material)	
			Vomitus (Dosing solution)		X	.	.	.	X	.	X	.	X	.	.
Loose stool		X	.			
Diarrhea				

Severity Codes: X = Present; 1 = Slight

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg Group 4 - 500 mg/kg

Time 10, 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Group	Sex	Animal	Clinical Sign	Site	Day numbers relative to Start Date										
					10 Before dosing	10 Time 10	11 Before dosing	11 Time 10	12 Before dosing	12 Time 10	13 Before dosing	13 Time 10	14 Before dosing		
3	f	50301	No Abnormalities Detected		X	.	X	.	X	
			Vomitus (Dosing formulation)	
			Vomitus (Food)	.	.	.	X	.	.	.	X	.	.		
			Vomitus (Dosing solution)	.	X	X	.	.		
		50302	Diarrhea	X	.	X	.	X	.	X	.
			No Abnormalities Detected	.	.	X	.	X	.	X	.	X	.	X	.
			Vomitus (Dosing formulation)
		50303	Vomitus (Food)	X	.	.	X	X	.	.
			Vomitus (Dosing solution)	.	X	.	.	.	X	.	.	X	.	.	
			No Abnormalities Detected
			Death	X	
			Vomitus (Dosing formulation)	
			Vomitus (Food)	
			Vomitus (Frothy material)	X	
			Vomitus (Dosing solution)	.	X	.	X	
Loose stool				
Diarrhea	.	.	X				

Severity Codes: X = Present; 1 = Slight

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg Group 4 - 500 mg/kg

Time 10, 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Day numbers relative to Start Date

Group	Sex	Animal	Clinical Sign	Site	14	15
					Time 10	a.m.
3	f	50301	No Abnormalities Detected		.	.
			Vomitus (Dosing formulation)		.	.
			Vomitus (Food)		.	.
			Vomitus (Dosing solution)		X	.
			Diarrhea		.	X
		50302	No Abnormalities Detected		.	X
			Vomitus (Dosing formulation)		.	.
			Vomitus (Food)		.	.
			Vomitus (Dosing solution)		X	.
		50303	No Abnormalities Detected		.	.
			Death		.	.
			Vomitus (Dosing formulation)		.	.
			Vomitus (Food)		.	.
			Vomitus (Frothy material)		.	.
			Vomitus (Dosing solution)		.	.
			Loose stool		.	.
			Diarrhea		.	.

Severity Codes: X = Present; 1 = Slight

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg Group 4 - 500 mg/kg

Time 10, 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Group	Sex	Animal	Clinical Sign	Site	Day numbers relative to Start Date									
					1 Before dosing	1 Time 10	2 Before dosing	2 Time 10	3 Before dosing	3 Time 10	4 Before dosing	4 Time 10	5 Before dosing	
4	f	50401	No Abnormalities Detected		X	X	.	X	.	.
			Moribundity
			Lateral position
			Vomitus (Dosing formulation)	1	.
			Vomitus (Food)	.	.	X
			Vomitus (Frothy material)	.	.	X
		50402	Vomitus (Dosing solution)	.	X	.	X	.	X	.	X	.	.	.
			Diarrhea	X
			Moribundity	X
			Lateral position	X
			Vomitus (Dosing formulation)	1	.	.
			Vomitus (Frothy material)	X	X	X	.	.	.
		50403	Vomitus (Dosing solution)	.	X	.	X	.	X	.	X	.	.	.
			Menses	X	X	X	X	X	X	X	X	X	X	X
			Diarrhea	X	.	X	X	X	.	.
			No-stool	X
			No Abnormalities Detected	X	X	.	.
			Vomitus (Dosing formulation)	1	.
			Vomitus (Food)	X
			Vomitus (Frothy material)	.	.	X
			Vomitus (Dosing solution)	.	X	.	X	.	X	.	X	.	.	.
Menses			
Loose stool			
Diarrhea	.	.	X	X	X			

Severity Codes: X = Present; 1 = Slight

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg Group 4 - 500 mg/kg

Time 10, 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Group	Sex	Animal	Clinical Sign	Site	Day numbers relative to Start Date								
					5 Time 10	6 Before dosing	6 Time 10	7 Before dosing	7 Time 10	8 Before dosing	8 Time 10	9 Before dosing	9 Time 10
4	f	50401	No Abnormalities Detected		X
			Moribundity		X	.	.	.
			Lateral position		X	.	.	.
			Vomitus (Dosing formulation)	
			Vomitus (Food)		X	.	.	.
			Vomitus (Frothy material)	
			Vomitus (Dosing solution)		X
			Diarrhea		.	X	X	X	.	X	.	.	.
		50402	Moribundity	
			Lateral position	
			Vomitus (Dosing formulation)	
			Vomitus (Frothy material)	
			Vomitus (Dosing solution)	
			Menses	
			Diarrhea	
			No-stool	
		50403	No Abnormalities Detected	
			Vomitus (Dosing formulation)		.	.	1
			Vomitus (Food)		X	.
			Vomitus (Frothy material)		X
			Vomitus (Dosing solution)		X	.	.	.	X	.	X	X	.
			Menses		.	X	X	X	X	X	.	.	.
			Loose stool	
			Diarrhea		X	X	.	.	.	X	.	.	X

Severity Codes: X = Present; 1 = Slight

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg Group 4 - 500 mg/kg

Time 10, 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Group	Sex	Animal	Clinical Sign	Site	Day numbers relative to Start Date										
					10 Before dosing	10 Time 10	11 Before dosing	11 Time 10	12 Before dosing	12 Time 10	13 Before dosing	13 Time 10	14 Before dosing		
4	f	50401	No Abnormalities Detected		
			Moribundity	
			Lateral position	
			Vomitus (Dosing formulation)	
			Vomitus (Food)	
			Vomitus (Frothy material)	
			Vomitus (Dosing solution)	
		50402	Diarrhea
			Moribundity
			Lateral position
			Vomitus (Dosing formulation)
			Vomitus (Frothy material)
			Vomitus (Dosing solution)
			Menses
		50403	Diarrhea
			No-stool
			No Abnormalities Detected	.	.	X
			Vomitus (Dosing formulation)
			Vomitus (Food)	X
			Vomitus (Frothy material)
			Vomitus (Dosing solution)	.	X
Menses			
Loose stool	X			
Diarrhea			

Severity Codes: X = Present; 1 = Slight

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg Group 4 - 500 mg/kg

Time 10, 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Day numbers relative to Start Date

Group	Sex	Animal	Clinical Sign	Site	14	15
					Time 10	a.m.
4	f	50401	No Abnormalities Detected		.	.
			Moribundity		.	.
			Lateral position		.	.
			Vomitus (Dosing formulation)		.	.
			Vomitus (Food)		.	.
			Vomitus (Frothy material)		.	.
			Vomitus (Dosing solution)		.	.
			Diarrhea		.	.
			50402	Moribundity		.
		Lateral position		.	.	
		Vomitus (Dosing formulation)		.	.	
		Vomitus (Frothy material)		.	.	
		Vomitus (Dosing solution)		.	.	
		Menses		.	.	
		Diarrhea		.	.	
		No-stool		.	.	
		50403	No Abnormalities Detected		.	.
		Vomitus (Dosing formulation)		.	.	
		Vomitus (Food)		.	.	
		Vomitus (Frothy material)		.	.	
		Vomitus (Dosing solution)		.	.	
Menses		.	.			
Loose stool		.	.			
Diarrhea		.	.			

Severity Codes: X = Present; 1 = Slight

Group 1 - 0 mg/kg Group 2 - 50 mg/kg Group 3 - 150 mg/kg Group 4 - 500 mg/kg

Time 10, 2h after dosing;

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Appendix 2 Body Weight

B120719

Sex: Male Bodyweight (kg)

0 mg/kg	Day(s) Relative to Start Date							
	-52	-45	-42	-35	-28	-21	-14	-7
10101	3.0	3.0	3.0	3.1	3.2	3.4	3.3	3.4
10102	3.2	3.1	3.1	3.2	3.2	3.3	3.3	3.4
10103	2.8	2.7	2.7	2.9	2.9	3.1	3.1	3.2

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Appendix 2 Body Weight

B120719

Sex: Male Bodyweight (kg)

50 mg/kg	Day(s) Relative to Start Date							
	-52	-45	-42	-35	-28	-21	-14	-7
10201	2.9	2.9	3.0	3.1	3.1	3.2	3.2	3.2
10202	3.4	3.4	3.4	3.5	3.5	3.7	3.7	3.8
10203	2.9	2.9	2.9	3.1	3.0	3.1	3.2	3.2

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Appendix 2 Body Weight

B120719

Sex: Male Bodyweight (kg)

150 mg/kg	Day(s) Relative to Start Date							
	-52	-45	-42	-35	-28	-21	-14	-7
10301	3.2	3.3	3.3	3.4	3.4	3.5	3.5	3.6
10302	2.8	2.9	2.9	3.0	3.0	3.1	3.1	3.2
10303	3.1	3.1	3.1	3.2	3.2	3.3	3.4	3.4

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Appendix 2 Body Weight

B120719

Sex: Male Bodyweight (kg)

500 mg/kg	Day(s) Relative to Start Date							
	-52	-45	-42	-35	-28	-21	-14	-7
10401	3.4	3.4	3.3	3.5	3.5	3.7	3.7	3.8
10402	3.1	3.0	3.0	3.1	3.1	3.3	3.2	3.2
10403	2.5	2.6	2.6	2.6	2.6	2.7	2.7	2.8

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Sex: Male Bodyweight (kg)

0 mg/kg	Day(s) Relative to Start Date					
	1	4	7	8	11	14
10101	3.4	3.4	-	3.4	3.5	3.4
10102	3.3	3.3	-	3.2	3.3	3.3
10103	3.1	3.1	-	3.1	3.2	3.1

[Day 7 : Final Body Weight.]

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Sex: Male Bodyweight (kg)

50 mg/kg	Day(s) Relative to Start Date					
	1	4	7	8	11	14
10201	3.2	3.1	-	3.1	2.9	2.9
10202	3.8	3.9	-	3.8	3.6	3.4
10203	3.3	3.2	-	3.1	3.2	3.2

[Day 7 : Final Body Weight.]

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Sex: Male Bodyweight (kg)

150 mg/kg	Day(s) Relative to Start Date					
	1	4	7	8	11	14
10301	3.5	3.3	-	3.1	2.9	2.8
10302	3.2	3.0	-	2.9	2.8	2.6
10303	3.4	3.5	-	3.3	3.2	3.1

[Day 7 : Final Body Weight.]

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Sex: Male Bodyweight (kg)

500 mg/kg	Day(s) Relative to Start Date					
	1	4	7	8	11	14
10401	3.8	3.7	-	3.5	3.2	-
10402	3.2	3.1	-	2.9	2.8	-
10403	2.8	2.5	2.5	-	-	-

[Day 7 : Final Body Weight.]

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Appendix 2 Body Weight

B120719

Sex: Female Bodyweight (kg)

0 mg/kg	Day(s) Relative to Start Date							
	-52	-45	-42	-35	-28	-21	-14	-7
50101	3.4	3.3	3.3	3.3	3.3	3.4	3.5	3.5
50102	2.9	2.9	2.9	3.0	3.0	3.1	3.0	3.0
50103	2.7	2.7	2.7	2.8	2.8	2.9	2.9	2.9

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]

Appendix 2 Body Weight

B120719

Sex: Female Bodyweight (kg)

50 mg/kg	Day(s) Relative to Start Date							
	-52	-45	-42	-35	-28	-21	-14	-7
50201	2.7	2.7	2.8	2.9	2.9	2.9	2.9	3.0
50202	2.8	2.8	2.8	2.9	2.9	2.9	2.9	3.0
50203	3.1	3.1	3.0	3.1	3.1	3.2	3.2	3.2

[Dose level was changed from 500 mg/kg to 250 mg/kg on Day 5.]