

- Belaguli, N.S., Sepulveda, J.L., Nigam, V., Charron, F., Nemer, M., Schwartz, R.J., 2000. Cardiac tissue enriched factors serum response factor and GATA-4 are mutual coregulators. *Mol. Cell Biol.* 20, 7550–7558.
- Beltrami, A.P., Barlucchi, L., Torella, D., Baker, M., Limana, F., Chimenti, S., et al., 2003. Adult cardiac stem cells are multipotent and support myocardial regeneration. *Cell* 114, 763–776.
- Benezra, R., Davis, R.L., Lockshon, D., Turner, D.L., Weintraub, H., 1990. The protein Id: a negative regulator of helix-loop-helix DNA binding proteins. *Cell* 61, 49–59.
- Benson, C.C., Zhou, Q., Long, X., Miano, J.M., 2011. Identifying functional single nucleotide polymorphisms in the human CARome. *Physiol. Genomics* 43, 1038–1048.
- Bergmann, O., Bhardwaj, R.D., Bernard, S., Zdunek, S., Barnabe-Heider, F., Walsh, S., et al., 2009. Evidence for cardiomyocyte renewal in humans. *Science* 324, 98–102.
- Bernstein, E., Kim, S.Y., Carmell, M.A., Murchison, E.P., Alcorn, H., Li, M.Z., et al., 2003. Dicer is essential for mouse development. *Nat. Genet.* 35, 215–217.
- Berridge, M.J., Bootman, M.D., Roderick, H.L., 2003. Calcium signalling: dynamics, homeostasis and remodelling. *Nat. Rev. Mol. Cell Biol.* 4, 517–529.
- Bers, D.M., Guo, T., 2005. Calcium signaling in cardiac ventricular myocytes. *Ann. N. Y. Acad. Sci.* 1047, 86–98.
- Bhalla, S.S., Robitaille, L., Nemer, M., 2001. Cooperative activation by GATA-4 and YY1 of the cardiac B-type natriuretic peptide promoter. *J. Biol. Chem.* 276, 11439–11445.
- Bhavsar, P.K., Dellow, K.A., Yacoub, M.H., Brand, N.J., Barton, P.J., 2000. Identification of cis-acting DNA elements required for expression of the human cardiac troponin I gene promoter. *J. Mol. Cell Cardiol.* 32, 95–108.
- Biressi, S., Tagliafico, E., Lamorte, G., Monteverde, S., Tenedini, E., Roncaglia, E., et al., 2007a. Intrinsic phenotypic diversity of embryonic and fetal myoblasts is revealed by genome-wide gene expression analysis on purified cells. *Dev. Biol.* 304, 633–651.
- Biressi, S., Molinaro, M., Cossu, G., 2007b. Cellular heterogeneity during vertebrate skeletal muscle development. *Dev. Biol.* 308, 281–293.
- Bismuth, K., Relaix, F., 2010. Genetic regulation of skeletal muscle development. *Exp. Cell Res.* 316, 3081–3086.
- Black, B., Olson, E.N., 1998. Transcriptional control of muscle development by myocyte enhancer factor-2 (MEF2) proteins. *Annu. Rev. Cell Dev. Biol.* 14, 167–196.
- Bladt, F., Riethmacher, D., Isenmann, S., Aguzzi, A., Birchmeier, C., 1995. Essential role for the c-met receptor in the migration of myogenic precursor cells into the limb bud. *Nature* 376, 768–771.
- Blank, R.S., McQuinn, T.C., Yin, K.C., Thompson, M.M., Takeyasu, K., Schwartz, R.J., et al., 1992. Elements of the smooth muscle alpha-actin promoter required in cis for transcriptional activation in smooth muscle. Evidence for cell type-specific regulation. *J. Biol. Chem.* 267, 984–989.
- Bober, E., Franz, T., Arnold, H.H., Gruss, P., Tremblay, P., 1994. Pax-3 is required for the development of limb muscles: a possible role for the migration of dermomyotomal muscle progenitor cells. *Development* 120, 603–612.
- Bopp, D., Burri, M., Baumgartner, S., Frigerio, G., Noll, M., 1986. Conservation of a large protein domain in the segmentation gene paired and in functionally related genes of *Drosophila*. *Cell* 47, 1033–1040.
- Borsani, G., DeGrandi, A., Ballabio, A., Bulfone, A., Bernard, L., Banfi, S., et al., 1999. EYA4, a novel vertebrate gene related to *Drosophila* eyes absent. *Hum. Mol. Genet.* 8, 11–23.
- Bottaro, D.P., Ruin, J., Faletto, D.L., Chan, A.M.L., Kmiecik, T.E., Vande Woude, G.F., et al., 1991. Identification of the hepatocyte growth factor receptor as the c-met protooncogene product. *Science* 251, 802–804.

- Braun, T., Gautel, M., 2011. Transcriptional mechanisms regulating skeletal muscle differentiation, growth and homeostasis. *Nat. Rev. Mol. Cell Biol.* 12, 349–361.
- Braun, T., Buschhausen-Denker, G., Bober, E., Tannich, E., Arnold, H.H., 1989. A novel human muscle factor related to but distinct from MyoD1 induces myogenic conversion in 10T1/2 fibroblasts. *EMBO J.* 8, 701–709.
- Brennecke, J., Hipfner, D.R., Stark, A., Russell, R.B., Cohen, S.M., 2003. bantam encodes a developmentally regulated microRNA that controls cell proliferation and regulates the proapoptotic gene hid in *Drosophila*. *Cell* 113, 25–36.
- Brewer, A., Pizzey, J., 2006. GATA factors in vertebrate heart development and disease. *Expert Rev. Mol. Med.* 8, 1–20.
- Brohmann, H., Jagla, K., Birchmeier, C., 2000. The role of Lbx1 in migration of muscle precursor cells. *Development* 127, 437–445.
- Brown, C.O., 3rd., Chi, X., Garcia-Gras, E., Shirai, M., Feng, X.H., Schwartz, R.J., 2004. The cardiac determination factor, Nkx2-5, is activated by mutual cofactors GATA-4 and Smad1/4 via a novel upstream enhancer. *J. Biol. Chem.* 279, 10659–10669.
- Bruneau, B.G., Logan, M., Davis, N., Levi, T., Tabin, C.J., Seidman, J.G., et al., 1999. Chamber-specific cardiac expression of Tbx5 and heart defects in Holt-Oram syndrome. *Dev. Biol.* 211, 100–108.
- Bruneau, B.G., Nemer, G., Schmitt, J.P., Charron, F., Robitaille, L., Caron, S., et al., 2001. A murine model of Holt-Oram syndrome defines roles of the T-box transcription factor Tbx5 in cardiogenesis and disease. *Cell* 106, 709–721.
- Buckingham, M., 1992. Making muscle in mammals. *Trends Genet.* 8, 144–148.
- Buckingham, M.E., Meilhac, S.M., 2011. Tracing cells for tracking cell lineage and clonal behaviour. *Dev. Cell* 21, 394–409.
- Buckingham, M., Vincent, S.D., 2009. Distinct and dynamic myogenic populations in the vertebrate embryo. *Curr. Opin. Genet. Dev.* 19, 444–453.
- Buckingham, M., Bajard, L., Chang, T., Daubas, P., Hadchouel, J., Meilhac, S., et al., 2003. The formation of skeletal muscle: from somite to limb. *J. Anat.* 202, 59–68.
- Buckingham, M., Meilhac, S., Zaffran, S., 2005. Building the mammalian heart from two sources of myocardial cells. *Nat. Rev. Genet.* 6, 826–835.
- Burri, M., Tromvoukis, Y., Bopp, D., Frigerio, G., Noll, M., 1989. Conservation of the paired domain in metazoans and its structure in three isolated human genes. *EMBO J.* 8, 1183–1190.
- Buskin, J.N., Hauschka, S.D., 1989. Identification of a myocyte-specific nuclear factor which binds to the muscle-specific enhancer of the mouse muscle creatine kinase gene. *Mol. Cell. Biol.* 9, 2627–2640.
- Cai, C.L., Liang, X., Shi, Y., Chu, P.H., Pfaff, S.L., Chen, J., et al., 2003. Isl1 identifies a cardiac progenitor population that proliferates prior to differentiation and contributes a majority of cells to the heart. *Dev. Cell* 5, 877–889.
- Cai, C.L., Zhou, W., Yang, L., Bu, L., Qyang, Y., Zhang, X., et al., 2005. T-box genes coordinate regional rates of proliferation and regional specification during cardiogenesis. *Development* 132, 2475–2487.
- Cai, C., Martin, J., Sun, Y., Cui, L., Wang, L., Ouyang, K., et al., 2008. A myocardial lineage derives from tbx18 epicardial cells. *Nature* 454, 104–108.
- Cai, X., Nomura-Kitabayashi, A., Cai, W., Yan, J., Christoffels, V.M., Cai, C.L., 2011. Myocardial Tbx20 regulates early atrioventricular canal formation and endocardial epithelial-mesenchymal transition via Bmp2. *Dev. Biol.* 360, 381–390.
- Callis, T.E., Pandya, K., Seok, H.Y., Tang, R.H., Tatsuguchi, M., Huang, Z.P., et al., 2009. MicroRNA-208a is a regulator of cardiac hypertrophy and conduction in mice. *J. Clin. Invest.* 119, 2772–2786.
- Calmont, A., Ivins, S., Van Bueren, K.L., Papangelis, I., Kyriakopoulou, V., Andrews, W.D., et al., 2009. Tbx1 controls cardiac neural crest cell migration during arch artery

- development by regulating Gbx2 expression in the pharyngeal ectoderm. *Development* 136, 3173–3183.
- Cao, Y., Yao, Z., Sarkar, D., Lawrence, M., Sanchez, G.J., Parker, M.H., et al., 2010. Genome-wide MyoD binding in skeletal muscle cells: a potential for broad cellular reprogramming. *Dev. Cell* 18, 662–674.
- Cartwright, E.J., Mohamed, T., Oceandy, D., Neyses, L., 2011. Calcium signaling dysfunction in heart disease. *Biofactors* 37, 175–181.
- Carvajal, J.J., Cox, D., Summerbell, D., Rigby, P.W.J., 2001. A BAC transgenic analysis of the Mrf4/Myf5 locus reveals interdigitated elements that control activation and maintenance of gene expression during muscle development. *Development* 128, 1857–1868.
- Carvajal, J.J., Keith, A., Rigby, P.W.J., 2008. Global transcriptional regulation of the locus encoding the skeletal muscle determination genes Mrf4 and Myf5. *Genes Dev.* 22, 265–276.
- Carvalho, B.M., Bassani, R.A., Franchini, K.G., Bassani, J.W., 2006. Enhanced calcium mobilization in rat ventricular myocytes during the onset of pressure overload-induced hypertrophy. *Am. J. Physiol. Heart Circ. Physiol.* 291, H1803–H1813.
- Cesana, M., Cacchiarelli, D., Legnini, I., Santini, T., Sthandier, O., Chinappi, M., et al., 2011. A long noncoding RNA controls muscle differentiation by functioning as a competing endogenous RNA. *Cell* 147, 358–369.
- Chang, C.P., Neilson, J.R., Bayle, J.H., Gestwicki, J.E., Kuo, A., Stankunas, K., et al., 2004. A field of myocardial-endocardial NFAT signaling underlies heart valve morphogenesis. *Cell* 118, 649–663.
- Chapman, D.L., Garvey, N., Hancock, S., Alexiou, M., Agulnik, S.I., Gibson-Brown, J.J., et al., 1996. Expression of the T-box family genes, Tbx1–Tbx5, during early mouse development. *Dev. Dyn.* 206, 379–390.
- Charron, F., Nemer, M., 1999. GATA transcription factors and cardiac development. *Semin. Cell Dev. Biol.* 10, 85–91.
- Charron, F., Paradis, P., Bronchain, O., Nemer, G., Nemer, M., 1999. Cooperative interaction between GATA-4 and GATA-6 regulates myocardial gene expression. *Mol. Cell Biol.* 19, 4355–4365.
- Chen, J., Kitchen, C.M., Streb, J.W., Miano, J.M., 2002. Myocardin: a component of a molecular switch for smooth muscle differentiation. *J. Mol. Cell. Cardiol.* 34, 1345–1356.
- Chen, J.F., Mandel, E.M., Thomson, J.M., Wu, Q., Callis, T.E., Hammond, S.M., et al., 2006. The role of microRNA-1 and microRNA-133 in skeletal muscle proliferation and differentiation. *Nat. Genet.* 38, 228–233.
- Chen, J.F., Murchison, E.P., Tang, R., Callis, T.E., Tatsuguchi, M., Deng, Z., et al., 2008. Targeted deletion of Dicer in the heart leads to dilated cardiomyopathy and heart failure. *Proc. Natl. Acad. Sci. USA* 105, 2111–2116.
- Chen, H., Wang, D., Bai, C., Wang, X., 2010. Proteomics-based biomarkers in chronic obstructive pulmonary disease. *J. Proteome Res.* 9, 2798–2808.
- Cheng, G., Hagen, T.P., Dawson, M.L., Barnes, K.V., Menick, D.R., 1999. The role of GATA, CARG, E-box, and a novel element in the regulation of cardiac expression of the Na⁺-Ca²⁺ exchanger gene. *J. Biol. Chem.* 274, 12819–12826.
- Chi, N., Epstein, J.A., 2002. Getting your Pax straight: Pax proteins in development and disease. *Trends Genet.* 18, 41–47.
- Chiang, C.S., Huang, C.H., Chieng, H., Chang, Y.T., Chang, D., Chen, J.J., et al., 2009. The Ca(v)3.2 T-type Ca(2⁺) channel is required for pressure overload-induced cardiac hypertrophy in mice. *Circ. Res.* 104, 522–530.
- Chong, S.-W., Korzh, V., Jiang, Y.-J., 2009. Myogenesis and molecules—insights from zebrafish *Danio rerio*. *J. Fish Biol.* 74, 1693–1755.

- Christ, B., Brand-Saberi, B., 2002. Limb muscle development. *Int. J. Dev. Biol.* 46, 905–914.
- Christoffels, V.M., Hoogaars, W.M., Tessari, A., Clout, D.E., Moorman, A.F., Campione, M., 2004. T-box transcription factor Tbx2 represses differentiation and formation of the cardiac chambers. *Dev. Dyn.* 229, 763–770.
- Clabby, M.L., Robison, T.A., Quigley, H.F., Wilson, D.B., Kelly, D.P., 2003. Retinoid X receptor alpha represses GATA-4-mediated transcription via a retinoid-dependent interaction with the cardiac-enriched repressor FOG-2. *J. Biol. Chem.* 278, 5760–5767.
- Clark, I.B., Boyd, J., Hamilton, G., Finnegan, D.J., Jarman, A.P., 2006. D-six4 plays a key role in patterning cell identities deriving from the *Drosophila* mesoderm. *Dev. Biol.* 294, 220–231.
- Clark, K.A., McElhinny, A.S., Beckerle, M.C., Gregorio, C.C., 2002. Striated muscle cytoarchitecture: an intricate web of form and function. *Annu. Rev. Cell Dev. Biol.* 18, 637–706.
- Cohen, E.D., Wang, Z., Lepore, J.J., Lu, M.M., Taketo, M.M., Epstein, D.J., et al., 2007. Wnt/beta-catenin signaling promotes expansion of Isl-1-positive cardiac progenitor cells through regulation of FGF signaling. *J. Clin. Invest.* 117, 1794–1804.
- Constantinides, P.G., Jones, P.A., Gevers, W., 1977. Functional striated muscle cells from non-myoblast precursors following 5-azacytidine treatment. *Nature* 267, 364–366.
- Cooper, S.J., Trinklein, N.D., Nguyen, L., Myers, R.M., 2007. Serum response factor binding sites differ in three human cell types. *Genome Res.* 17, 136–144.
- Cordes, K.R., Sheehy, N.T., White, M.P., Berry, E.C., Morton, S.U., Muth, A.N., et al., 2009. miR-145 and miR-143 regulate smooth muscle cell fate and plasticity. *Nature* 460, 705–710.
- Dai, Y.S., Markham, B.E., 2001. p300 functions as a coactivator of transcription factor GATA-4. *J. Biol. Chem.* 276, 37178–37185.
- Dai, Y.S., Cserjesi, P., Markham, B.E., Molkentin, J.D., 2002. The transcription factors GATA4 and dHAND physically interact to synergistically activate cardiac gene expression through a p300-dependent mechanism. *J. Biol. Chem.* 277, 24390–24398.
- Dalton, S., Treisman, R., 1992. Characterization of SAP-1, a protein recruited by serum response factor to the c-fos serum response element. *Cell* 68, 597–612.
- Davis, R.L., Weintraub, H., Lassar, A.B., 1987. Expression of a single transfected cDNA converts fibroblasts to myoblasts. *Cell* 51, 987–1000.
- Debrus, S., Rahbani, L., Marttila, M., Delorme, B., Paradis, P., Nemer, M., 2005. The zinc finger-only protein Zfp260 is a novel cardiac regulator and a nuclear effector of alpha1 adrenergic signaling. *Mol. Cell Biol.* 25, 8669–8682.
- de la Pompa, J.L., Timmerman, L.A., Takimoto, H., Yoshida, H., Elia, A.J., Samper, E., et al., 1998. Role of the NF-ATc transcription factor in morphogenesis of cardiac valves and septum. *Nature* 392, 182–186.
- De Santa, F., Barozzi, I., Mietton, F., Ghisletti, S., Polletti, S., Tusi, B.K., et al., 2010. A large fraction of extragenic RNA pol II transcription sites overlap enhancers. *PLoS Biol.* 8, e1000384.
- Desjardins, P.R., Burkman, J.M., Shrager, J.B., Allmond, L.A., Stedman, H.H., 2002. Evolutionary implications of three novel members of the human sarcomeric myosin heavy chain gene family. *Mol. Biol. Evol.* 19, 375–393.
- Deutsch, U., Dressler, G.R., Gruss, P., 1988. Pax 1, a member of a paired box homologous murine gene family, is expressed in segmented structures during development. *Cell* 53, 617–625.
- Di Lisi, R., Millino, C., Calabria, E., Altruda, F., Schiaffino, S., Ausoni, S., 1998. Combinatorial cis-acting elements control tissue-specific activation of the cardiac troponin I gene in vitro and in vivo. *J. Biol. Chem.* 273, 25371–25380.

- Diehl, A.G., Zarepari, S., Qian, M., Khanna, R., Angeles, R., Gage, P.J., 2006. Extraocular muscle morphogenesis and gene expression are regulated by Pitx2 gene dose. *Invest. Ophthalmol. Vis. Sci.* 47, 1785–1793.
- Dinger, M.E., Amaral, P.P., Mercer, T.R., Pang, K.C., Bruce, S.J., Gardiner, B.B., et al., 2008. Long noncoding RNAs in mouse embryonic stem cell pluripotency and differentiation. *Genome Res.* 18, 1433–1445.
- Dong, F., Sun, X., Liu, W., Ai, D., Klysiak, E., Lu, M.-F., 2006. Pitx2 promotes development of splanchnic mesoderm-derived branchiomeric muscle. *Development* 133, 4891–4899.
- Doran, P., Donoghue, P., O'connell, K., Gannon, J., Ohlendieck, K., 2007. Proteomic profiling of pathological and aged skeletal muscle fibres by peptide mass fingerprinting (Review). *Int. J. Mol. Med.* 19, 547–564.
- Du, M., Zhao, J.X., Yan, X., Huang, Y., Nicodemus, L.V., Yue, W., et al., 2011. Fetal muscle development, mesenchymal multipotent cell differentiation, and associated signaling pathways. *J. Anim. Sci.* 89, 583–590.
- Duband, J.L., Gimona, M., Scatena, M., Sartore, S., Small, J.V., 1993. Calponin and SM 22 as differentiation markers of smooth muscle: spatiotemporal distribution during avian embryonic development. *Differentiation* 55, 1–11.
- Dyer, L., Kirby, M., 2009. Sonic hedgehog maintains proliferation in secondary heart field progenitors and is required for normal arterial pole formation. *Dev. Biol.* 330, 305–317.
- Eisenberg, I., Alexander, M.S., Kunkel, L.M., 2009. miRNAs in normal and diseased skeletal muscle. *J. Cell. Mol. Med.* 13, 2–11.
- Epstein, D.J., Vekemans, M., Gros, P., 1991. Spotch (Sp2H), a mutation affecting development of the mouse neural tube, shows a deletion within the paired homeodomain of Pax3. *Cell* 67, 767–774.
- Epstein, J., Cai, J., Glaser, T., Jepeal, L., Maas, R., 1994. Identification of a paired domain recognition sequence and evidence for DNA-dependent conformational changes. *J. Biol. Chem.* 269, 8355–8361.
- Epstein, J.A., Shapiro, D.N., Chang, J., Lam, P.Y., Maas, R.L., 1996. Pax3 modulates expression of the c-met receptor during limb muscle development. *Proc. Natl. Acad. Sci. USA* 93, 4213–4218.
- Erz˘en, I., Sketelj, J., Agbulut, O., Angaut-Petit, D., Butler-Browne, G.S., 2001. Coexpression of myosin heavy chain 2b with myosin heavy chain 1—fact or artefact? *Basic Appl. Myol.* 11, 119–125.
- Evans, T., Felsenfeld, G., 1989. The erythroid-specific transcription factor Eryf1: a new finger protein. *Cell* 58, 877–885.
- Fan, X., Brass, L.F., Poncz, M., Spitz, F., Maire, P., Manning, D.R., 2000. The alpha subunits of Gz and Gi interact with the eyes absent transcription cofactor Eya2, preventing its interaction with the six class of homeodomain-containing proteins. *J. Biol. Chem.* 275, 32129–32134.
- Firulli, A.B., 2003. A HANDful of questions: the molecular biology of the heart and neural crest derivatives (HAND)-subclass of basic helix-loop-helix transcription factors. *Gene* 312, 27–40.
- Firulli, A.B., Firulli, B.A., Wang, J., Rogers, R.H., Conway, S.J., 2010. Gene replacement strategies to test the functional redundancy of basic helix-loop-helix transcription factor. *Pediatr. Cardiol.* 31, 438–448.
- Fischbach, K.F., Technau, G., 1984. Cell degeneration in the developing optic lobes of the sine oculis and small optic-lobes mutants of *Drosophila melanogaster*. *Dev. Biol.* 104, 219–239.
- Fischer, A., Klattig, J., Kneitz, B., Diez, H., Maier, M., Holtmann, B., et al., 2005. Hey basic helix-loop-helix transcription factors are repressors of GATA4 and GATA6 and restrict expression of the GATA target gene ANF in fetal hearts. *Mol. Cell Biol.* 25, 8960–8970.

- Fowler, M.R., Naz, J.R., Graham, M.D., Bru-Mercier, G., Harrison, S.M., Orchard, C.H., 2005. Decreased Ca^{2b} extrusion via $\text{Na}^b/\text{Ca}^{2b}$ exchange in epicardial left ventricular myocytes during compensated hypertrophy. *Am. J. Physiol. Heart Circ. Physiol.* 288 , H2431–H2438.
- Franz, T., Kothary, R., Surani, M.A.H., Halata, Z., Grim, M., 1993. The splotch mutation interferes with muscle development in the limbs. *Anat. Embryol.* 187, 153–160.
- Fulcoli, F.G., Huynh, T., Scambler, P.J., Baldini, A., 2009. *Tbx1* regulates the BMP *smad1* pathway in a transcription independent manner. *PLoS One* 4, e6049.
- Gabella, G., 2002. Development of visceral smooth muscle in vertebrate myogenesis. In: Brand-Saberi, B. (Ed.), *Results and Problems in Cell Differentiation*, vol. 38. Springer, Freiburg, pp. 1–37.
- Galili, N., Davis, R.J., Fredericks, W.J., Mukhopadhyay, S., Rauscher, F.R., Emanuel, B.S., et al., 1993. Fusion of a fork head domain gene to PAX3 in the solid tumor alveolar rhabdomyosarcoma. *Nat. Genet.* 5, 230–235.
- Galliot, B., de Vargas, C., Miller, D., 1999. Evolution of homeobox genes: Q50 Paired-like genes founded the Paired class. *Dev. Genes Evol.* 209, 186–197.
- Garg, V., Yamagishi, C., Hu, T., Kathiriya, I.S., Yamagishi, H., Srivastava, D., 2001. *Tbx1*, a DiGeorge syndrome candidate gene, is regulated by sonic hedgehog during pharyngeal arch development. *Dev. Biol.* 235, 62–73.
- Garg, V., Kathiriya, I.S., Barnes, R., Schluterman, M.K., King, I.N., Butler, C.A., et al., 2003. GATA4 mutations cause human congenital heart defects and reveal an interaction with TBX5. *Nature* 424, 443–447.
- Giordani, J., Bajard, L., Demignon, J., Daubas, P., Buckingham, M., Maire, P., 2007. Six proteins regulate the activation of *Myf5* expression in embryonic mouse limbs. *Proc. Natl. Acad. Sci. USA* 104, 11310–11315.
- Goddeeris, M.M., Schwartz, R., Klingensmith, J., Meyers, E.N., 2007. Independent requirements for Hedgehog signaling by both the anterior heart field and neural crest cells for outflow tract development. *Development* 134, 1593–1604.
- Goddeeris, M.M., Rho, S., Petiet, A., Davenport, C.L., Johnson, G.A., Meyers, E.N., et al., 2008. Intracardiac septation requires hedgehog-dependent cellular contributions from outside the heart. *Development* 135, 1887–1895.
- Gorza, L., 1990. Identification of a novel type 2 fiber population in mammalian skeletal muscle by combined use of histochemical myosin ATPase and anti-myosin monoclonal antibodies. *J. Histochem. Cytochem.* 38, 257–265.
- Goulding, M.D., Chalepakis, G., Deutsch, U., Erselius, J.R., Gruss, P., 1991. Pax-3, a novel murine DNA binding protein expressed during early neurogenesis. *EMBO J.* 10 , 1135–1147.
- Goulding, M., Lumsden, A., Paquette, A.J., 1994. Regulation of Pax-3 expression in the dermomyotome and its role in muscle development. *Development* 120, 957–971.
- Graef, I.A., Chen, F., Chen, L., Kuo, A., Crabtree, G.R., 2001. Signals transduced by $\text{Ca}^{(2b)}$ /calcineurin and NFATc3/c4 pattern the developing vasculature. *Cell* 105 , 863–875.
- Gre´pin, C., Dagnino, L., Robitaille, L., Haberstroh, L., Antakly, T., Nemer, M., 1994. A hormone-encoding gene identifies a pathway for cardiac but not skeletal muscle gene transcription. *Mol. Cell Biol.* 14, 3115–3129.
- Greulich, F., Rudat, C., Kispert, A., 2011. Mechanisms of T-box gene function in the developing heart. *Cardiovasc. Res.* 91, 212–222.
- Grifone, R., Kelly, R.G., 2007. Heartening news for head muscle development. *Trends Genet.* 23, 365–369.
- Grifone, R., Laclef, C., Spitz, F., Lopez, S., Demignon, J., Guidotti, J.E., et al., 2004. Six1 and *Eyal* expression can reprogram adult muscle from the slow-twitch phenotype into the fast-twitch phenotype. *Mol. Cell. Biol.* 24, 6253–6567.

- Grifone, R., Demignon, J., Houbron, C., Souil, E., Niro, C., Seller, M., et al., 2005. Six1 and Six4 homeoproteins are required for Pax3 and MRF expression during myogenesis in the mouse embryo. *Development* 132, 2235–2249.
- Grifone, R., Demignon, J., Giordani, J., Niro, C., Souil, E., Bertin, F., et al., 2007. Eya1 and Eya2 proteins are required for hypaxial somitic myogenesis in the mouse embryo. *Dev. Biol.* 302, 602–616.
- Gros, J., Manceau, M., Thome, V., Marcelle, C., 2005. A common somitic origin for embryonic muscle progenitors and satellite cells. *Nature* 435, 954–958.
- Gross, M.K., Moran-Rivard, L., Velasquez, T., Nakatsu, M.N., Jagla, K., Goulding, M., 2000. Lbx1 is required for muscle precursor migration along a lateral pathway into the limb. *Development* 127, 413–424.
- Gruss, P., Walther, C., 1992. Pax in development. *Cell* 69, 719–722.
- Gualberto, A., LePage, D., Pons, G., Mader, S.L., Park, K., Atchison, M.L., et al., 1992. Functional antagonism between YY1 and the serum response factor. *Mol. Cell. Biol.* 12, 4209–4214.
- Gudjonsson, T., Adriance, M.C., Sternlicht, M.D., Petersen, O.W., Bissell, M.J., 2005. Myoepithelial cells: their origin and function in breast morphogenesis and neoplasia. *J. Mammary Gland Biol. Neoplasia* 10, 261–272.
- Guettler, S., Vartiainen, M.K., Miralles, F., Larijani, B., Treisman, R., 2008. RPEL motifs link the serum response factor cofactor MAL but not myocardin to Rho signaling via actin binding. *Mol. Cell. Biol.* 28, 732–742.
- Guevel, L., Lavoie, J.R., Perez-Iratxeta, C., Rouger, K., Dubreil, L., Feron, M., et al., 2011. Quantitative proteomic analysis of dystrophic dog muscle. *J. Proteome Res.* 10, 2465–2478.
- Guñler, I., Russell, A.P., 2010. MicroRNAs in skeletal muscle: their role and regulation in development, disease and function. *J. Physiol.* 588 (Pt 21), 4075–4087.
- Guo, L., Nakamura, K., Lynch, J., Opas, M., Olson, E.N., Agellon, L.B., et al., 2002. Cardiac-specific expression of calcineurin reverses embryonic lethality in calreticuline-deficient mouse. *J. Biol. Chem.* 277, 50776–50779.
- Guo, H., Ingolia, N.T., Weissman, J.S., Bartel, D.P., 2010. Mammalian microRNAs predominantly act to decrease target mRNA levels. *Nature* 466, 835–840.
- Gygi, S.P., Rochon, Y., Franza, R., Aebersold, R., 1999. Correlation between protein and mRNA abundance in yeast. *Mol. Cell. Biol.* 19, 1720–1730.
- Habets, P.E., Moorman, A.F., Clout, D.E., van Roon, M.A., Lingbeek, M., van Lohuizen, M., et al., 2002. Cooperative action of Tbx2 and Nkx2.5 inhibits ANF expression in the atrioventricular canal: implications for cardiac chamber formation. *Genes Dev.* 16, 1234–1246.
- Haddad, F., Bodell, P.W., Qin, A.X., Giger, J.M., Baldwin, K.M., 2003. Role of antisense RNA in coordinating cardiac myosin heavy chain gene switching. *J. Biol. Chem.* 278, 37132–37138.
- Haddad, F., Bodell, P.W., Qin, A.X., Jiang, W., Giger, J.M., Baldwin, K.M., 2008. Intergenic transcription and developmental regulation of cardiac myosin heavy chain genes. *Am. J. Physiol. Heart Circ. Physiol.* 294, H29–H40.
- Harrelson, Z., Kelly, R.G., Goldin, S.N., Gibson-Brown, J.J., Bollag, R.J., Silver, L.M., et al., 2004. Tbx2 is essential for patterning the atrioventricular canal and for morphogenesis of the outflow tract during heart development. *Development* 131, 5041–5052.
- Hasegawa, K., Lee, S.J., Jobe, S.M., Markham, B.E., Kitsis, R.N., 1997. cis-Acting sequences that mediate induction of beta-myosin heavy chain gene expression during left ventricular hypertrophy due to aortic constriction. *Circulation* 96, 3943–3953.

- Hasty, P., Bradley, A., Morris, J.H., Edmondson, D.G., Venuti, J.M., Olson, E.N., et al., 1993. Muscle deficiency and neonatal death in mice with a targeted mutation in the myogenin gene. *Nature* 364, 501–506.
- Heanue, T.A., Reshef, R., Davis, R.J., Mardon, G., Oliver, G., Tomarev, S., et al., 1999. Synergistic regulation of vertebrate muscle development by Dach2, Eya2, and Six1, homologs of genes required for *Drosophila* eye formation. *Genes Dev.* 13, 3231–3243.
- Herbert, S.P., Stainer, D.Y.R., 2011. Molecular control of endothelial cell behaviour during blood vessel morphogenesis. *Nat. Rev. Mol. Cell Biol.* 12, 551–564.
- Hoogaars, W.M., Tessari, A., Moorman, A.F., de Boer, P.A., Hagoort, J., Soufan, A.T., et al., 2004. The transcriptional repressor Tbx3 delineates the developing central conduction system of the heart. *Cardiovasc. Res.* 62, 489–499.
- Hooper, S.L., Thuma, J.B., 2005. Invertebrate muscles: muscle specific genes and proteins. *Physiol. Rev.* 85, 1001–1060.
- Horst, D., Ustanina, S., Sergi, C., Mikuz, G., Juergens, H., Braun, T., et al., 2006. Comparative expression analysis of Pax3 and Pax7 during mouse myogenesis. *Int. J. Dev. Biol.* 50, 47–54.
- Hu, T., Yamagishi, H., Maeda, J., McAnally, J., Yamagishi, C., Srivastava, D., 2004. Tbx1 regulates fibroblast growth factors in the anterior heart field through a reinforcing autoregulatory loop involving forkhead transcription factors. *Development* 131, 5491–5502.
- Hu, P., Geles, K.G., Paik, J.H., DePinho, R.A., Tjian, R., 2008. Codependent activators direct myoblast-specific MyoD transcription. *Dev. Cell* 15, 534–546.
- Huang, T., Lin, A.E., Cox, G.F., Golden, W.L., Feldman, G.L., Ute, M., et al., 2002. Cardiac phenotype in chromosome 4q- syndrome with and without a deletion of the dHAND gene. *Genet. Med.* 4, 464–467.
- Hutcheson, D.A., Zhao, J., Merrell, A., Haldar, M., Kardon, G., 2009. Embryonic and fetal limb myogenic cells are derived from developmentally distinct progenitors and have different requirements for b-catenin. *Genes Dev.* 23, 997–1013.
- Hutson, M.R., Kirby, M.L., 2007. Model systems for the study of heart development and disease. Cardiac neural crest and conotruncal malformations. *Semin. Cell Dev. Biol.* 1 (8), 101–110.
- Hutson, M.R., Zhang, P., Stadt, H.A., Sato, A.K., Li, Y.X., Burch, J., et al., 2006. Cardiac arterial pole alignment is sensitive to FGF8 signaling in the pharynx. *Dev. Biol.* 295, 486–497.
- Huxley, H.E., 1969. The mechanism of muscular contraction. *Science* 164, 1356–1366.
- IHGSC, 2004. Finishing the euchromatic sequence of the human genome. *Nature* 431, 931–945.
- Ip, H.S., Wilson, D.B., Heikinheimo, M., Tang, Z., Ting, C.N., Simon, M.C., et al., 1994. The GATA-4 transcription factor transactivates the cardiac muscle-specific troponin C promoter-enhancer in nonmuscle cells. *Mol. Cell. Biol.* 14, 7517–7526.
- Ito, M., Dabrowska, R., Guerriero, V., Hartshornem, D.J., 1989. Identification in turkey gizzard of an acidic protein related to the C-terminal portion of smooth muscle myosin light chain kinase. *J. Biol. Chem.* 264, 13971–13974.
- Ivey, K.N., Muth, A., Arnold, J., King, F.W., Yeh, R.F., Fish, J.E., et al., 2008. MicroRNA regulation of cell lineages in mouse and human embryonic stem cells. *Cell Stem Cell* 2, 219–229.
- Jerome, L.A., Papaioannou, V.E., 2001. DiGeorge syndrome phenotype in mice mutant for the T-box gene, Tbx1. *Nat. Genet.* 27, 286–291.
- Jesudason, E.C., 2009. Airway smooth muscle: an architect of the lung? *Thorax* 64, 541–545.

- Johansen, F.-E., Prywes, R., 1995. Serum response factor: transcriptional regulation of genes induced by growth factors and differentiation. *Biochim. Biophys. Acta* 1242, 1–10.
- Jostes, B., Walter, C., Gruss, P., 1990. The murine paired box gene, Pax7, is expressed specifically during the development of the nervous and muscular system. *Mech. Dev.* 33, 27–37.
- Kaikkonen, M.U., Lam, M.T.Y., Glass, C.K., 2011. Non-coding RNAs as regulators of gene expression and epigenetics. *Cardiovasc. Res.* 90, 430–440.
- Kakita, T., Hasegawa, K., Morimoto, T., Kaburagi, S., Wada, H., Sasayama, S., 1999. p300 protein as a coactivator of GATA-5 in the transcription of cardiac-restricted atrial natriuretic factor gene. *J. Biol. Chem.* 274, 34096–34102.
- Kathiriya, I.S., King, I.N., Murakami, M., Nakagawa, M., Astle, J.M., Gardner, K.A., et al., 2004. Hairy-related transcription factors inhibit GATA-dependent cardiac gene expression through a signal-responsive mechanism. *J. Biol. Chem.* 279, 54937–54943.
- Kawakami, K., Ohto, H., Ikeda, K., Roeder, R.G., 1996. Structure, function and expression of a murine homeobox protein AREC3, a homologue of *Drosophila sine oculis* gene product, and implication in development. *Nucleic Acids Res.* 1996, 303–310.
- Kelly, R.G., Jerome-Majewska, L.A., Papaioannou, V.E., 2004. The del22q11.2 candidate gene *Tbx1* regulates branchiomic myogenesis. *Hum. Mol. Genet.* 13, 2829–2840.
- Kim, S., Ip, H.S., Lu, M.M., Clendenin, C., Parmacek, M.S., 1997. A serum response factor-dependent transcriptional regulatory program identifies distinct smooth muscle cell sublineages. *Mol. Cell Biol.* 17, 2266–2278.
- Kim, T.G., Chen, J., Sadoshima, J., Lee, Y., 2004. *Jumonji* represses atrial natriuretic factor gene expression by inhibiting transcriptional activities of cardiac transcription factors. *Mol. Cell Biol.* 24, 10151–10160.
- Kim, H.K., Lee, Y.S., Sivaprasad, U., Malhotra, A., Dutta, A., 2006. Muscle-specific microRNA miR-206 promotes muscle differentiation. *J. Cell Biol.* 174, 677–687.
- Kirby, R.J., Hamilton, G.M., Finnegan, D.J., Johnson, K.J., Jarman, A.P., 2001. *Drosophila* homolog of the myotonic dystrophy-associated gene, *six5*, is required for muscle and gonad development. *Curr. Biol.* 11, 1044–1049.
- Kitano, H., 2002. Systems biology: a brief overview. *Science* 295, 1662–1664.
- Klaus, A., Saga, Y., Taketo, M.M., Tzahor, E., Birchmeier, W., 2007. Distinct roles of Wnt/beta-catenin and Bmp signaling during early cardiogenesis. *Proc. Natl. Acad. Sci. USA* 104, 18531–18536.
- Ko, J.L., Engel, J.D., 1993. DNA-binding specificities of the GATA transcription factor family. *Mol. Cell Biol.* 13, 4011–4022.
- Kodo, K., Yamagishi, H., 2011. A decade of advances in the molecular embryology and genetics underlying congenital heart defects. *Circ. J.* 75, 2296–2304.
- Kodo, K., Nishizawa, T., Furutani, M., Arai, S., Yamamura, E., Joo, K., et al., 2009. GATA6 mutations cause human cardiac outflow tract defects by disrupting semaphorin-plexin signaling. *Proc. Natl. Acad. Sci. USA* 106, 13933–13938.
- Kontaridis, M.I., Swanson, K.D., David, F.S., Barford, D., Neel, B.G., 2006. PTPN11 (*Shp2*) mutations in LEOPARD syndrome have dominant negative, not activating effects. *J. Biol. Chem.* 281, 6785–6792.
- Kraus, F., Haenig, B., Kispert, A., 2001. Cloning and expression analysis of the mouse T-box gene *Tbx18*. *Mech. Dev.* 100, 83–86.
- Kuang, S., Rudnicki, M.A., 2008. The emerging biology of satellite cells and their therapeutic potential. *Trends Mol. Med.* 14, 82–91.
- Kuang, S., Charge, S.B., Seale, P., Huh, M., Rudnicki, M.A., 2006. Distinct roles for Pax7 and Pax3 in adult regenerative myogenesis. *J. Cell Biol.* 172, 103–113.
- Kuang, S., Kuroda, K., Le Grand, F., Rudnicki, M.A., 2007. Asymmetric self-renewal and commitment of satellite stem cells in muscle. *Cell* 129, 999–1010.

- Kumar, J.P., 2009. The sine oculis homeobox (SIX) family of transcription factors as regulators of development and disease. *Cell. Mol. Life Sci.* 66, 565–583.
- Kuo, H., Chen, J., Ruiz-Lozano, P., Zou, Y., Nemer, M., Chien, K.R., 1999. Control of segmental expression of the cardiac-restricted ankyrin repeat protein gene by distinct regulatory pathways in murine cardiogenesis. *Development* 126, 4223–4234.
- Kuwahara, K., Wang, Y., McAnally, J., Richardson, J.A., Bassel-Duby, R., Hill, J.A., et al., 2006. TRPC6 fulfills a calcineurin signaling circuit during pathologic cardiac remodeling. *J. Clin. Invest.* 116, 3114–3126.
- Kwon, C., Arnold, J., Hsiao, E.C., Taketo, M.M., Conklin, B.R., Srivastava, D., 2007. Canonical wnt signaling is a positive regulator of mammalian cardiac progenitors. *Proc. Natl. Acad. Sci. USA* 104, 10894–10899.
- Kwon, C., Qian, L., Cheng, P., Nigam, V., Arnold, J., Srivastava, D., 2009. A regulatory pathway involving notch1/beta-catenin/isl1 determines cardiac progenitor cell fate. *Nat. Cell Biol.* 11, 951–957.
- Laclef, C., Hamard, G., Demignon, J., Souil, E., Houbron, C., Maire, P., 2003a. Altered myogenesis in Six1 deficient mice. *Development* 130, 2239–2252.
- Laclef, C., Souil, E., Demignon, J., Maire, P., 2003b. Thymus, kidney and craniofacial abnormalities in Six 1 deficient mice. *Mech. Dev.* 120, 669–679.
- Lagha, M., Sato, T., Bajard, L., Daubas, P., Esner, M., Montarras, D., et al., 2008. Regulation of skeletal muscle stem cell behavior by Pax3 and Pax7. *Cold Spring Harb. Symp. Quant. Biol.* 73, 1 – 9.
- Lagos-Quintana, M., Rauhut, R., Lendeckel, W., Tuschl, T., 2001. Identification of novel genes coding for small expressed RNAs. *Science* 294, 853–858.
- Lagos-Quintana, M., Rauhut, R., Yalcin, A., Meyer, J., Lendeckel, W., Tuschl, T., 2002. Identification of tissue-specific microRNAs from mouse. *Curr. Biol.* 12, 735–739.
- Larsen, W.J., Bleyl, S.B., Brauer, P.R., Francis-West, P.H., Schoenwolf, G.C., 2009. *Human embryology*. Philadelphia, PA: Churchill Livingstone/Elsevier.
- Latinkic, B.V., Cooper, B., Smith, S., Kotecha, S., Towers, N., Sparrow, D., et al., 2004. Transcriptional regulation of the cardiac-specific MLC2 gene during *Xenopus* embryonic development. *Development* 131, 669–679.
- Laugwitz, K.L., Moretti, A., Lam, J., Gruber, P., Chen, Y.H., Woodard, S., et al., 2005. Postnatal isl1⁺ cardioblasts enter fully differentiated cardiomyocyte lineages. *Nature* 433, 647–653.
- Le Grand, F., Rudnicki, M.A., 2007. Skeletal muscle satellite cells and adult myogenesis. *Curr. Opin. Cell Biol.* 19, 628–633.
- Lee, R.C., Feinbaum, R.L., Ambros, V., 1993. The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell* 75, 843–854.
- Lepore, J.J., Mericko, P.A., Cheng, L., Lu, M.M., Morrissey, E.E., Parmacek, M.S., 2006. GATA-6 regulates semaphorin 3C and is required in cardiac neural crest for cardiovascular morphogenesis. *J. Clin. Invest.* 116, 929–939.
- Lepper, C., Conway, S.J., Fan, C.-M., 2009. Adult satellite cells and embryonic muscle progenitors have distinct genetic requirements. *Nature* 460, 627–631.
- Lepper, C., Fan, C.-M., 2010. Inducible lineage tracing of Pax7-descendant cells reveals embryonic origin of adult satellite cells. *Genesis* 48, 424–436.
- Lescroart, F., Kelly, R.G., Le Garrec, J.-F., Nicolas, J.-F., Meilhac, S.M., Buckingham, M., 2010. Clonal analysis reveals common lineage relationships between head muscles and second heart field derivatives in the mouse embryo. *Development* 137, 3269–3279.
- Lewis, B.P., Burge, C.B., Bartel, D.P., 2005. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* 120, 15–20.
- Li, S., Wang, D., Wang, Z., Richardson, J., Olson, E.N., 2003. The serum response factor coactivator myocardin is required for vascular smooth muscle development. *Proc. Natl. Acad. Sci. USA* 100, 9366–9370.

- Li, S., Czubyrt, M.P., McAnally, J., Bassel-Duby, R., Richardson, J.A., Wiebe, F.F., et al., 2005. Requirement for serum response factor for skeletal muscle growth and maturation revealed by tissue-specific gene deletion in mice. *Proc. Natl. Acad. Sci. USA* 102, 1082–1087.
- Li, S., Chang, S., Qi, X., Richardson, J.A., Olson, E.N., 2006. Requirement of a myocardin-related transcription factor for development of mammary myoepithelial cells. *Mol. Cell Biol.* 26, 5797–5808.
- Lien, C.L., Wu, C., Mercer, B., Webb, R., Richardson, J.A., Olson, E.N., 1999. Control of early cardiac-specific transcription of *Nkx2-5* by a GATA-dependent enhancer. *Development* 126, 75–84.
- Lin, L., Bu, L., Cai, C.L., Zhang, X., Evans, S., 2006. *Isl1* is upstream of sonic hedgehog in a pathway required for cardiac morphogenesis. *Dev. Biol.* 295, 756–763.
- Lin, L., Cui, L., Zhou, W., Dufort, D., Zhang, X., Cai, C.L., et al., 2007. Beta-catenin directly regulates *Isl1* expression in cardiovascular progenitors and is required for multiple aspects of cardiogenesis. *Proc. Natl. Acad. Sci. USA* 104, 9313–9318.
- Lin, X., Huo, Z., Liu, X., Zhang, Y., Li, L., Zhao, H., et al., 2010. A novel GATA6 mutation in patients with tetralogy of Fallot or atrial septal defect. *J. Hum. Genet.* 55, 662–667.
- Lints, T.J., Parsons, L.M., Hartley, L., Lyons, I., Harvey, R.P., 1993. *Nkx-2.5*: a novel murine homeobox gene expressed in early heart progenitor cells and their myogenic descendants. *Development* 119, 419–431.
- Liu, W., Yasui, K., Ophhof, T., Ishiki, R., Lee, J.K., Kamiya, K., et al., 2002. Developmental changes of Ca(2p) handling in mouse ventricular cells from early embryo to adulthood. *Life Sci.* 71, 1279–1292.
- Liu, W., Selever, J., Wang, D., Lu, M.F., Moses, K.A., Schwartz, R.J., et al., 2004. *Bmp4* signaling is required for outflow-tract septation and branchial-arch artery remodeling. *Proc. Natl. Acad. Sci. USA* 101, 4489–4494.
- Liu, N., Williams, A.H., Kim, Y., McAnally, J., Bezprozvannaya, S., Sutherland, L.B., et al., 2007. An intragenic MEF2-dependent enhancer directs muscle-specific expression of microRNAs 1 and 133. *Proc. Natl. Acad. Sci. USA* 104, 20844–20849.
- Liu, C., Shen, A., Li, X., Jiao, W., Zhang, X., Li, Z., 2008a. T-box transcription factor *TBX20* mutations in Chinese patients with congenital heart disease. *Eur. J. Med. Genet.* 51, 580–587.
- Liu, N., Bezprozvannaya, S., Williams, A.H., Qi, X., Richardson, J.A., Bassel-Duby, R., et al., 2008b. microRNA-133a regulates cardiomyocyte proliferation and suppresses smooth muscle gene expression in the heart. *Genes Dev.* 22, 3242–3254.
- Lu, M.F., Pressman, C., Dyer, R., Johnson, R.L., Martin, J.F., 1999. Function of Rieger syndrome gene in left-right asymmetry and craniofacial development. *Nature* 401, 276–278.
- Lu, J.R., Bassel-Duby, R., Hawkins, A., Chang, P., Valdez, R., Wu, H., 2002. Control of facial muscle development by *MyoR* and *capsulin*. *Science* 298, 2378–2381.
- Lye, C.M., Fasano, L., Woolf, A.S., 2010. Ureter myogenesis: putting *Teashirt* into context. *J. Am. Soc. Nephrol.* 21, 24–30.
- Lynch, J., Guo, L., Gelebart, P., Chilibeck, K., Xu, J., Molkenin, J.D., et al., 2005. Calreticulin signals upstream of calcineurin and MEF2C in a critical Ca(2p)- dependent signaling cascade. *J. Cell Biol.* 170, 37–47.
- Machida, M., Takagaki, Y., Matsuoka, R., Kawaguchi, N., 2011. Proteomic comparison of spherical aggregates and adherent cells of cardiac stem cells. *Int. J. Cardiol.* 153, 296–305.
- Maeda, J., Yamagishi, H., McAnally, J., Yamagishi, C., Srivastava, D., 2006. *Tbx1* is regulated by forkhead proteins in the secondary heart field. *Dev. Dyn.* 235, 701–710.
- Maitra, M., Koenig, S.N., Srivastava, D., Garg, V., 2010. Identification of GATA6 sequence variants in patients with congenital heart defects. *Pediatr. Res.* 68, 281–285.

- Majesky, M.W., 2003. Vascular smooth muscle diversity: insights from developmental biology. *Curr. Atheroscler. Rep.* 5, 208–213.
- Makuch, R., Birukov, K., Shirinsky, V., Dabrowska, R., 1991. Functional interrelationship between calponin and caldesmon. *Biochem. J.* 15, 33–38.
- Mansouri, A., Hallonet, M., Gruss, P., 1996a. Pax genes and their roles in cell differentiation and development. *Curr. Opin. Cell Biol.* 8, 851–857.
- Mansouri, A., Stoykova, A., Torres, M., Gruss, P., 1996b. Dysgenesis of cephalic neural crest derivatives in Pax7^{-/-} mutant mice. *Development* 122, 831–838.
- Mansouri, A., Goudreau, G., Gruss, P., 1999. Pax genes and their role in organogenesis. *Cancer Res.* 59, 1707 s–1710s.
- Marieb, E., Hoehn, K., 2007. *Human Anatomy & Physiology*, seventh ed. Pearson Benjamin Cummings, San Francisco p. 317.
- Marin, T.M., Keith, K., Davies, B., Conner, D.A., Guha, P., Kalaitzidis, D., et al., 2011. Rapamycin reverses hypertrophic cardiomyopathy in a mouse model of Leopard syndrome-associated PTPN11 mutation. *J. Clin. Invest.* 121, 1026–1043.
- Maroto, M., Reshef, R., Muñsterberg, A.E., Koester, S., Goulding, M., Lassar, A.B., 1997. Ectopic Pax-3 activates MyoD and Myf-5 expression in embryonic mesoderm and neural tissue. *Cell* 89, 139–148.
- Martínez-Estrada, O.M., Lettice, L.A., Essafi, A., Guadix, J.A., Slight, J., Velecela, V., et al., 2010. Wt1 is required for cardiovascular progenitor cell formation through transcriptional control of snail and E-cadherin. *Nat. Genet.* 42, 89–93.
- Marx, S.O., Reiken, S., Hisamatsu, Y., Jayaraman, T., Burkhoff, D., Rosemlit, N., et al., 2000. PKA phosphorylation dissociates FKBP12.6 from the calcium release channel (ryanodine receptor): defective regulation in failing hearts. *Cell* 101, 365–376.
- Matkovich, S.J., Van Booven, D.J., Eschenbacher, W.H., Dorn, G.W.I.I., 2011. RISC RNA sequencing for context-specific identification of in vivo microRNA. *Circ. Res.* 108, 18–26.
- McBride, K., Charron, F., Lefebvre, C., Nemer, M., 2003. Interaction with GATA transcription factors provides a mechanism for cell-specific effects of c-Fos. *Oncogene* 22, 8403–8412.
- McCarthy, J.J., 2008. MicorRNA-206: the skeletal muscle-specific myomiR. *Biochim. Biophys. Acta* 1779, 682–691.
- McFadden, D.G., Charite, J., Richardson, J.A., Srivastava, D., Firulli, A.B., Olson, E.N., 2000. A GATA-dependent right ventricular enhancer controls dHAND transcription in the developing heart. *Development* 127, 5331–5341.
- McFadden, D.G., Barbosa, A.C., Richardson, J.A., Schneider, M.D., Srivastava, D., Olson, E.N., 2004. The Hand1 and Hand2 transcription factors regulate expansion of the embryonic cardiac ventricles in a gene dosage-dependent manner. *Development* 132, 189–201.
- McGregot, E., Dunn, M.J., 2006. Proteomics of the heart: unraveling disease. *Circ. Res.* 98, 309–321.
- McGrew, M.J., Bogdanova, N., Hasegawa, K., Hughes, S.H., Kitsis, R.N., Rosenthal, N., 1996. Distinct gene expression patterns in skeletal and cardiac muscle are dependent on common regulatory sequences in the MLC1/3 locus. *Mol. Cell Biol.* 16, 4524–4534.
- McGuigan, K., Phillips, P.C., Postlethwait, J.H., 2004. Evolution of sarcomeric myosin heavy chain genes: evidence from fish. *Mol. Biol. Evol.* 21, 1042–1056.
- McHugh, K.M., 1996. Molecular analysis of gastrointestinal smooth muscle development. *J. Pediatr. Gastroenterol. Nutr.* 23, 379–394.
- McKinnell, I.W., Ishibashi, J., Le Grand, F., Punch, V.G., Addicks, G.C., Greenblatt, J.F., et al., 2008. Pax7 activates myogenic genes by recruitment of a histone methyltransferase complex. *Nat. Cell Biol.* 10, 77–84.

- McLin, V.A., Henning, S.J., Jamrich, M., 2009. The role of the visceral mesoderm in the development of the gastrointestinal tract. *Gastroenterology* 136, 2074–2091.
- Merika, M., Orkin, S.H., 1993. DNA-binding specificity of GATA family transcription factors. *Mol. Cell. Biol.* 13, 3999–4010.
- Mesaeli, N., Nakamura, K., Zvaritch, E., Dickie, P., Dziak, E., Krause, K.H., et al., 1999. Calreticulin is essential for cardiac development. *J. Cell Biol.* 144, 857–868.
- Mesbah, K., Harrelson, Z., Theveniau-Ruissy, M., Papaioannou, V.E., Kelly, R.G., 2008. Tbx3 is required for outflow tract development. *Circ. Res.* 103, 743–750.
- Miano, J.M., 2003. Serum response factor: toggling between disparate programs of gene expression. *J. Mol. Cell. Cardiol.* 35, 577–593.
- Miano, J.M., Long, X., Fujiwara, K., 2007. Serum response factor: master regulator of the actin cytoskeleton and contractile apparatus. *Am. J. Physiol. Cell Physiol.* 292, C70–C81.
- Minty, A., Kedes, L., 1986. Upstream regions of the human cardiac actin gene that modulate its transcription in muscle cells: presence of an evolutionarily conserved repeated motif. *Mol. Cell. Biol.* 6, 2125–2136.
- Miralles, F., Posern, G., Zaromytidou, A.I., Treisman, R., 2003. Actin dynamics control SRF activity by regulation of its coactivator MAL. *Cell* 113, 329–342.
- Miwa, T., Kedes, L., 1987. Duplicated CArG box domains have positive and mutually dependent regulatory roles in expression of the human α -cardiac actin gene. *Mol. Cell. Biol.* 7, 2803–2813.
- Molkentin, J.D., 2006. Dichotomy of Ca²⁺ in the heart: contraction versus intracellular signaling. *J. Clin. Invest.* 116, 623–626.
- Molkentin, J.D., Lu, J.R., Antos, C.L., Markham, B., Richardson, J., Robbins, J., et al., 1998. A calcineurin-dependent transcriptional pathway for cardiac hypertrophy. *Cell* 93, 215–228.
- Molkentin, J.D., Kalvakolanu, D.V., Markham, B.E., 1994. Transcription factor GATA-4 regulates cardiac muscle-specific expression of the α -myosin heavy-chain gene. *Mol. Cell Biol.* 14, 4947–4957.
- Molkentin, J.D., 2000. The zinc finger-containing transcription factors GATA-4, -5, and -6. Ubiquitously expressed regulators of tissue-specific gene expression. *J. Biol. Chem.* 275, 38949–38952.
- Molkentin, J.D., Lin, Q., Duncan, S.A., Olson, E.N., 1997. Requirement of the transcription factor GATA4 for heart tube formation and ventral morphogenesis. *Genes Dev.* 11, 1061–1072.
- Mootosamy, R.C., Dietrich, S., 2002. Distinct regulatory cascades for head and trunc myogenesis. *Development* 129, 573–583.
- Moretti, A., Caron, L., Nakano, A., Lam, J.T., Bernhausen, A., Chen, Y., et al., 2006. Multipotent embryonic isl1⁺ progenitor cells lead to cardiac, smooth muscle, and endothelial cell diversification. *Cell* 127, 1151–1165.
- Morikawa, Y., Cserjesi, P., 2008. Cardiac neural crest expression of Hand2 regulates outflow and second heart field development. *Circ. Res.* 103, 1422–1429.
- Morin, S., Charron, F., Robitaille, L., Nemer, M., 2000. GATA-dependent recruitment of MEF2 proteins to target promoters. *EMBO J.* 19, 2046–2055.
- Morrissey, E.E., Ip, H.S., Lu, M.M., Parmacek, M.S., 1996. GATA-6: a zinc finger transcription factor that is expressed in multiple cell lineages derived from lateral mesoderm. *Dev. Biol.* 177, 309–322.
- Morrissey, E.E., Tang, Z., Sigrist, K., Lu, M.M., Jiang, F., Ip, H.S., et al., 1998. GATA6 regulates HNF4 and is required for differentiation of visceral endoderm in the mouse embryo. *Genes Dev.* 12, 3579–3590.

- Morrissey, E.E., Ip, H.S., Tang, Z., Lu, M.M., Parmacek, M.S., 1997. GATA-5: a transcriptional activator expressed in a novel temporally and spatially-restricted pattern during embryonic development. *Dev. Biol.* 183, 21–36.
- Murphy, A.M., Thompson, W.R., Peng, L.F., Jones, L., 2nd., 1997. Regulation of the rat cardiac troponin I gene by the transcription factor GATA-4. *Biochem. J.* 322 (Pt 2), 393–401.
- Murphy, M., Kardon, G., 2011. Origin of vertebrate limb muscle: the role of progenitor and myoblast populations. *Curr. Top. Dev. Biol.* 96, 1–32.
- Murre, C., McCaw, P.S., Vaessin, H., Caudy, M., Jan, L.Y., Jan, Y.N., et al., 1989. Interactions between heterologous helix-loop-helix proteins generate complexes that bind specifically to a common DNA sequence. *Cell* 58, 537–544.
- Nabeshima, Y., Hanaoka, K., Hayasaka, M., Esumi, E., Li, S., Nonaka, I., et al., 1993. Myogenin gene disruption results in perinatal lethality because of severe muscle defect. *Nature* 364, 532–535.
- Naiche, L.A., Harrelson, Z., Kelly, R.G., Papaioannou, V.E., 2005. T-box genes in vertebrate development. *Annu. Rev. Genet.* 39, 219–239.
- Nakamura, S., Hayashi, K., Iwasaki, K., Fujioka, T., Egusa, H., Yatani, H., et al., 2010. Nuclear import mechanism for myocardin family members and their correlation with vascular smooth muscle cell phenotype. *J. Biol. Chem.* 285, 37314–37323.
- Nakayama, H., Bodi, I., Maillet, M., DeSantiago, J., Domeier, T.L., Mikoshiba, K., et al., 2010. The IP3 receptor regulates cardiac hypertrophy in response to select stimuli. *Circ. Res.* 107, 659–666.
- Nakazawa, M., Uchida, K., Aramaki, M., Kodo, K., Yamagishi, C., Takahashi, T., et al., 2011. Inositol 1,4,5-trisphosphate receptors are essential for the development of the second heart field. *J. Mol. Cell. Cardiol.* 51, 58–66.
- Natarajan, A., Yamagishi, H., Ahmad, F., Li, D., Roberts, R., Matsuoka, R., et al., 2001. Human eHAND, but not dHAND, is down-regulated in cardiomyopathies. *J. Mol. Cell. Cardiol.* 33, 1607–1614.
- Natarajan, A., Lemos, D.R., Rossi, F.M.V., 2010. Fibro-adipogenic progenitors. *Cell Cycle* 9, 2045–2046.
- Needham, D.M., 1926. Red and white muscles. *Physiol. Rev.* 6, 1–27.
- Nemer, G., Nemer, M., 2003. Transcriptional activation of BMP-4 and regulation of mammalian organogenesis by GATA-4 and -6. *Dev. Biol.* 254, 131–148.
- Nicholas, S.B., Philipson, K.D., 1999. Cardiac expression of the Na(p)/Ca(2p) exchanger NCX1 is GATA factor dependent. *Am. J. Physiol.* 277 (1Pt 2), H324–H330.
- Nie, X., Deng, C.X., Wang, Q., Jiao, K., 2008. Disruption of smad4 in neural crest cells leads to mid-gestation death with pharyngeal arch, craniofacial and cardiac defects. *Dev. Biol.* 316, 417–430.
- Niu, Z., Li, A., Zhang, S.X., Schwartz, R.J., 2007. Serum response factor micromanaging cardiogenesis. *Curr. Opin. Cell Biol.* 19, 618–627.
- Noden, D.M., Francis-West, P., 2006. The differentiation and morphogenesis of craniofacial muscles. *Dev. Dyn.* 235, 1194–1218.
- Noden, D.M., Marcucio, R., Borycki, A.G., Emerson, C.P., Jr., 1999. Differentiation of avian craniofacial muscles: I. Patterns of early regulatory gene expression and myosin heavy chain synthesis. *Dev. Dyn.* 216, 96–112.
- Norman, C., Runswick, M., Pollock, R., Treisman, R., 1988. Isolation and properties of cDNA clones encoding SRF, a transcription factor that binds to the c-fos serum response element. *Cell* 55, 989–1003.
- O'Rourke, J.R., Georges, S.A., Seay, H.R., Tapscott, S.J., McManus, M.T., Goldhamer, D.J., et al., 2007. Essential role for Dicer during skeletal muscle development. *Dev. Biol.* 311, 359–368.

- Oceandy, D., Cartwright, E.J., Emerson, M., Prehar, S., Baudoin, F.M., Zi, M., et al., 2007. Neuronal nitric oxide synthase signaling in the heart is regulated by the sarcolemmal calcium pump 4b. *Circulation* 115, 83–492.
- Oh, J., Richardson, J.A., Olson, E.N., 2005. Requirement of myocardin-related transcription factor-B for remodeling of branchial arch arteries and smooth muscle differentiation. *Proc. Natl. Acad. Sci. USA* 102, 15122–15127.
- Ohlendieck, K., 2010. Proteomics of skeletal muscle glycolysis. *Biochim. Biophys. Acta* 1804, 2089–2101.
- Ohlendieck, K., 2011. Skeletal muscle proteomics: current approaches, technical challenges and emerging technique. *Skelet. Muscle* 1, 6–15.
- Ohto, H., Kamada, S., Tago, K., Tominaga, S.I., Ozaki, H., Sato, S., et al., 1999. Cooperation of six and eya in activation of their target genes through nuclear translocation of Eya. *Mol. Cell. Biol.* 19, 6815–6824.
- Oliver, G., Wehr, R., Jenkins, N.A., Copeland, N.G., Cheyette, B.N., Hartenstein, V., et al., 1995. Homeobox genes and connective tissue patterning. *Development* 121, 693–705.
- Olivey, H.E., Svensson, E.C., 2010. Epicardial–myocardial signaling directing coronary vasculogenesis. *Circ. Res.* 106, 818–832.
- Olson, E.N., 1990. MyoD family: a paradigm for development. *Genes Dev.* 4, 1454–1461.
- Olson, E.N., 1993. Regulation of muscle transcription by the MyoD family: the heart of the matter. *Circ. Res.* 72, 1 – 8.
- Olson, E.N., Klein, W.H., 1994. bHLH factors in muscle development: dead lines and commitments, what to leave in and what to leave out. *Genes Dev.* 8, 1 – 8.
- Olson, E.N., Nordheim, A., 2010. Linking actin dynamics and gene transcription to drive cellular motile functions. *Nat. Rev. Mol. Cell Biol.* 11, 353–365.
- Olson, E.N., Srivastava, D., 1996. Molecular pathways controlling heart development. *Science* 272, 671–676.
- Olson, E.N., Arnold, H.H., Rigby, P.W.J., Wold, B.J., 1996. Know your neighbours: three phenotypes in null mutants of the myogenic bHLH gene MRF4. *Cell* 85, 1 – 5.
- rom, U.A., Shiekhatar, R., 2011. Noncoding RNAs and enhancers: complications of a long-distance relationship. *Trends Genet.* 27, 433–439.
- rom, U.A., Derrien, T., Beringer, M., Gumireddy, K., Gardini, A., Bussotti, G., et al., 2010. Long noncoding RNAs with enhancer-like function in human cells. *Cell* 143, 46–58.
- Oustanina, S., Hause, G., Braun, T., 2004. Pax7 directs postnatal renewal and propagation of myogenic satellite cells but not their specification. *EMBO J.* 23, 3430–3439.
- Owens, G.K., 1995. Regulation of differentiation of vascular smooth muscle cells. *Physiol. Rev.* 75, 487–517.
- Owens, G.K., Kumar, M.S., Wamhoff, B.R., 2004. Molecular regulation of vascular smooth muscle cell differentiation in development and disease. *Physiol. Rev.* 84, 767–801.
- Ozaki, H., Watanabe, Y., Takahashi, K., Kitamura, K., Tanaka, A., Urase, K., et al., 2001. Six4, a putative myogenin gene regulator, is not essential for mouse embryonal development. *Mol. Cell. Biol.* 21, 3343–3350.
- Pandur, P., Lasche, M., Eisenberg, L.M., Kuhl, M., 2002. Wnt-11 activation of a noncanonical Wnt signalling pathway is required for cardiogenesis. *Nature* 418, 636–641.
- Papaioannou, V.E., 2001. T-box genes in development: from hydra to humans. *Int. Rev. Cytol.* 207, 1–70.
- Park, E., Watanabe, Y., Smyth, G., Miyagawa-Tomita, S., Meyers, E., Klingensmith, J., et al., 2008. An FGF autocrine loop initiated in second heart field mesoderm regulates morphogenesis at the arterial pole of the heart. *Development* 135, 3599–3610.
- Parmacek, M.S., 2010. Myocardin-related transcription factor-A: mending a broken heart. *Circ. Res.* 107, 168–170.

- Pe´rez-Pomares, J.M., Gonza´les-Rosa, J.M., Mu˜oz-Chapuli, R., 2009. Building the vertebrate heart—an evolutionary approach to cardiac development. *Int. J. Dev. Biol.* 53, 1427–1443.
- Pignoni, F., Hu, B., Zavitz, K.H., Xiao, J., Garrity, P.A., Zipursky, S.L., 1997. The eye-specification proteins *So* and *Eya* form a complex and regulate multiple steps in *Drosophila* eye development. *Cell* 91, 881–891.
- Plageman, T.F., Jr., Yutzey, K.E., 2005. T-box genes and heart development: putting the “T” in heart. *Dev. Dyn.* 232, 11–20.
- Ponting, C.P., Oliver, P.L., Reik, W., 2009. Evolution and functions of long noncoding RNAs. *Cell* 136, 629–641.
- Porter, G.A., Jr., Makuck, R.F., Rivkees, S.A., 2003. Intracellular calcium plays an essential role in cardiac development. *Dev. Dyn.* 227, 280–290.
- Pownall, M.E., Gustafsson, M.K., Emerson, C.P., 2002. Myogenic regulatory factors and the specification of muscle progenitors in vertebrate embryos. *Annu. Rev. Cell Dev. Biol.* 18, 747–783.
- Prall, O.W.J., Menon, M.K., Solloway, M.J., Watanabe, Y., Zaffran, S., Bajolle, F., et al., 2007. An *Nkx2-5/bmp2/smad1* negative feedback loop controls heart progenitor specification and proliferation. *Cell* 128, 947–959.
- Puri, P.L., Sartorelli, V., Yang, X.J., Hamamori, Y., Ogryzko, V.V., Howard, B.H., et al., 1997. Differential roles of p300 and PCAF acetyltransferases in muscle differentiation. *Mol. Cell* 1, 35–45.
- Ranger, A.M., Grusby, M.J., Hodge, M.R., Gravallese, E.M., de la Brousse, F.C., Hoey, T., et al., 1998. The transcription factor NF-ATc is essential for cardiac valve formation. *Nature* 392, 186–190.
- Rangrez, A.Y., Massey, Z.A., Meuth, V.M., Metzinger, L., 2011. miR-143 and miR-145. *Circ. Cardiovasc. Genet.* 4, 197–205.
- Rao, P.K., Kumar, R.M., Farkhondeh, M., Baskerville, S., Lodish, H.F., 2006. Myogenic factors that regulate expression of muscle-specific microRNAs. *Proc. Natl. Acad. Sci. USA* 103, 8721–8726.
- Rao, P.K., Toyama, Y., Chiang, H.R., Gupta, S., Bauer, M., Medvid, R., et al., 2009. Loss of cardiac microRNA-mediated regulation leads to dilated cardiomyopathy and heart failure. *Circ. Res.* 105, 585–594.
- Reamon-Buettner, S.M., Ciribilli, Y., Inga, A., Borlak, J., 2008. A loss-of-function mutation in the binding domain of *HAND1* predicts hypoplasia of the human hearts. *Hum. Mol. Genet.* 17, 1397–1405.
- Reiter, J.F., Alexander, J., Rodaway, A., Yelon, D., Patient, R., Holder, N., et al., 1999. *Gata5* is required for the development of the heart and endoderm in zebrafish. *Genes Dev.* 13, 2983–2995.
- Relaix, F., Rocancourt, D., Mansouri, A., Buckingham, M., 2004. Divergent functions of murine *Pax3* and *Pax7* in limb muscle development. *Genes Dev.* 18, 1088–1105.
- Relaix, F., Rocancourt, D., Mansouri, A., Buckingham, M., 2005. A *Pax3/Pax7*-dependent population of skeletal muscle progenitor cells. *Nature* 435, 948–953.
- Relaix, F., Montarras, D., Zaffran, S., Gayraud-Morel, B., Rocancourt, D., Tajbakhsh, S., et al., 2006. *Pax3* and *Pax7* have distinct and overlapping functions in adult muscle progenitor cells. *J. Cell Biol.* 172, 91–102.
- Reznikoff, C.A., Brankow, D.W., Heidelberger, C., 1973. Establishment and characterization of a cloned line of C3H mouse embryo cells sensitive to post confluence inhibition of division. *Cancer Res.* 33, 3231–3238.
- Rivkees, S.A., Chen, M., Kulkarni, J., Browne, J., Zhao, Z., 1999. Characterization of the murine *A1* adenosine receptor promoter, potent regulation by *GATA-4* and *Nkx2.5*. *J. Biol. Chem.* 274, 14204–14209.

- Rochais, F., Mesbah, K., Kelly, R.G., 2009. Signaling pathways controlling second heart field development. *Circ Res.* 104, 933–942.
- Roderick, H.L., Higazi, D.R., Smyrnias, I., Fearnley, C., Harzheim, D., Bootman, M.D., 2007. Calcium in the heart: when it's good, it's very very good, but when it's bad, it's horrid. *Biochem. Soc. Trans.* 35, 957–961.
- Rosemblit, N., Moschella, M.C., Ondrias, E., Gutstein, D.E., Ondrias, K., Marks, A.R., 1999. Intracellular calcium release channel expression during embryogenesis. *Dev. Biol.* 206, 163–177.
- Rosenberg, P., Hawkins, A., Stiber, J., Shelton, J.M., Hutcheson, K., Bassel-Duby, R., et al., 2004. TRPC3 channels confer cellular memory of recent neuromuscular activity. *Proc. Natl. Acad. Sci. USA* 101, 9387–9392.
- Rosoff, M.L., Nathanson, N.M., 1998. GATA factor-dependent regulation of cardiac m2 muscarinic acetylcholine gene transcription. *J. Biol. Chem.* 273, 9124–9129.
- Rudnicki, M.A., Braun, T., Hinuma, S., Jaenisch, R., 1992. Inactivation of MyoD in mice leads to up-regulation of the myogenic HLH gene Myf-5 and results in apparently normal muscle development. *Cell* 71, 383–390.
- Rudnicki, M.A., Schnegelsberg, P.N., Stead, R.H., Braun, T., Arnold, H.H., Jaenisch, R., 1993. MyoD or Myf-5 is required for the formation of skeletal muscle. *Cell* 75, 1351–1359.
- Saccone, V., Puri, P.L., 2010. Epigenetic regulation of skeletal myogenesis. *Organogenesis* 6, 48–53.
- Sambasivan, R., Gayraud-Morel, B., Dumas, G., Cimper, C., Paisant, S., Kelly, R.G., et al., 2009. Distinct regulatory cascades govern extraocular and pharyngeal arch muscle progenitor cell fates. *Dev. Cell* 16, 810–821.
- Sambasivan, R., Kuratani, S., Tajbakhsh, S., 2011. An eye on the head: the development and evolution of craniofacial muscles. *Development* 138, 2401–2415.
- Sartorelli, V., Puri, P.L., Hamamori, Y., Ogryzko, V., Chung, G., Nakatani, Y., et al., 1999. Acetylation of MyoD directed by PCAF is necessary for the execution of the muscle program. *Mol. Cell* 4, 725–734.
- Sasazuki, T., Sawada, T., Sakon, S., Kitamura, T., Kishi, T., Okazaki, T., et al., 2002. Identification of a novel transcriptional activator, BSAC, by a functional cloning to inhibit tumor necrosis factor-induced cell death. *J. Biol. Chem.* 277, 28853–28860.
- Schiaffino, S., Reggiani, C., 2011. Fiber types in mammalian skeletal muscles. *Physiol. Rev.* 91, 1447–1531.
- Schleiffarth, J.R., Person, A.D., Martinsen, B.J., Sukovich, D.J., Neumann, A., Baker, C.V., et al., 2007. Wnt5a is required for cardiac outflow tract septation in mice. *Pediatr. Res.* 61, 386–391.
- Schultheiss, T.M., Burch, J.B., Lassar, A.B., 1997. A role for bone morphogenetic proteins in the induction of cardiac myogenesis. *Genes Dev.* 11, 451–462.
- Schwarz-Sommer, Z., Huijser, P., Nacken, W., Saedler, H., Sommer, H., 1990. Genetic control of flower development by homeotic genes in *Antirrhinum majus*. *Science* 250, 931–936.
- Scott, W., Stevens, J., Bindr-Macleod, S.A., 2001. Human skeletal muscle fiber type classifications. *Phys. Ther.* 81, 1810–1816.
- Seale, P., Sabourin, L.A., Girgis-Gabardo, A., Mansouri, A., Gruss, P., Rudnicki, M.A., 2000. Pax7 is required for the specification of myogenic satellite cells. *Cell* 102, 777–786.
- Seo, H.C., Curtiss, J., Mlodzik, M., Fjose, A., 1999. Six class homeobox genes in *Drosophila* belong to three distinct families and are involved in head development. *Mech. Dev.* 83, 127–139.

- Sepulveda, J.L., Vlahopoulos, S., Iyer, D., Belaguli, N., Schwartz, R.J., 2002. Combinatorial expression of GATA4, Nkx2-5, and serum response factor directs early cardiac gene activity. *J. Biol. Chem.* 277, 25775–25782.
- Serva, A., Claas, C., Starkuviene, V., 2011. A potential of micorRNAs for high-content screening. *J. Nucleic Acids* 2011, 870903.
- Shapiro, D.N., Sublett, J.E., Li, B., Downing, J.R., Naeve, C.W., 1993. Fusion of PAX3 to a member of the Forkhead family of transcription factors in human alveolar rhabdomyosarcoma. *Cancer Res.* 53, 5108–5112.
- Shaw, P.E., Schroter, H., Nordheim, A., 1989. The ability of a ternary complex to form over the serum response element correlates with serum inducibility of the human c-fos promoter. *Cell* 56, 563–572.
- Shih, H.P., Gross, M.K., Kioussi, C., 2008. Muscle development: forming the head and trunk muscles. *Acta Histochem.* 110, 97–108.
- Shimizu, R., Yamamoto, M., 2005. Gene expression regulation and domain function of hematopoietic GATA factors. *Semin. Cell Dev. Biol.* 6, 129–136.
- Shore, P., Sharrocks, A.D., 1995. The MADS-box family of transcription factors. *Eur. J. Biochem.* 229, 1–13.
- Small, E.M., Olson, E.N., 2011. Pervasive roles of microRNAs in cardiovascular biology. *Nature* 469, 336–342.
- Small, E.M., Frost, R.J.A., Olson, E.N., 2010. MicorRNAs add a new dimension to cardiovascular disease. *Circulation* 121, 1022–1032.
- Sorrentino, R.P., Gajewski, K.M., Schulz, R.A., 2005. GATA factors in Drosophila heart and blood cell development. *Semin. Cell Dev. Biol.* 16, 107–116.
- Spitz, F., Demignon, J., Porteu, A., Kahn, A., Concordet, J.P., Daegelen, D., et al., 1998. Expression of myogenin during embryogenesis is controlled by Six/sine oculis homeoproteins through a conserved MEF3 binding site. *Proc. Natl. Acad. Sci. USA* 95, 14220–14225.
- Srivastava, D., 2006. Making or breaking the heart: from lineage determination to morphogenesis. *Cell* 126, 1037–1048.
- Srivastava, D., Cserjesi, P., Olson, E.N., 1995. A subclass of bHLH proteins required for cardiac morphogenesis. *Science* 270, 1995–1999.
- Srivastava, D., Thomas, T., Lin, Q., Kirby, M.L., Brown, D., Olson, E.N., 1997. Regulation of cardiac mesodermal and neural crest development by the bHLH transcription factor, dHAND. *Nat. Genet.* 16, 154–160.
- Strobeck, M., Kim, S., Zhang, J.C., Clendenin, C., Du, K.L., Parmacek, M.S., 2001. Binding of serum response factor to CArG box sequences is necessary but not sufficient to restrict gene expression to arterial smooth muscle cells. *J. Biol. Chem.* 11, 16418–16424.
- Stuart, E.T., Kioussi, C., Gruss, P., 1993. Mammalian Pax genes. *Ann. Rev. Genet.* 28, 219–238.
- Sun, Q., Chen, G., Streb, J.W., Long, X., Yang, Y., Stoeckert, C.J., Jr., et al., 2006. Defining the mammalian CArGome. *Genome Res.* 16, 197–207.
- Svensson, E.C., Tufts, R.L., Polk, C.E., Leiden, J.M., 1999. Molecular cloning of FOG-2: a modulator of transcription factor GATA-4 in cardiomyocytes. *Proc. Natl. Acad. Sci. USA* 96, 956–961.
- Svensson, E.C., Huggins, G.S., Dardik, F.B., Polk, C.E., Leiden, J.M., 2000. A functionally conserved N-terminal domain of the friend of GATA-2 (FOG-2) protein represses GATA4-dependent transcription. *J. Biol. Chem.* 275, 20762–20769.
- Tajbakhsh, S., 2005. Skeletal muscle stem and progenitor cells-reconciling genetics and lineage. *Exp. Cell Res.* 306, 364–372.
- Tajbakhsh, S., Buckingham, M., 1994. Mouse limb muscle is determined in the absence of the earliest myogenic factor myf-5. *Proc. Natl. Acad. Sci. USA* 91, 747–751.

- Tajbakhsh, S., Rocancourt, D., Cossu, G., Buckingham, M., 1997. Redefining the genetic hierarchies controlling skeletal myogenesis: Pax-3 and Myf-5 act upstream of MyoD. *Cell* 89, 127–138.
- Takahashi, K., Yamanaka, S., 2006. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* 126, 663–676.
- Takahashi, K., Abe, M., Hiwada, K., Kokubu, T., 1988. A novel troponinT-like protein (calponin) in vascular smooth muscle: interaction with tropomyosin paracrystals. *J. Hypertens. Suppl.* 6, S40–S43.
- Takahashi, K., Tanabe, K., Ohnuki, M., Narita, M., Ichisaka, T., Tomoda, K., et al., 2007. Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 131, 861–872.
- Tam, O.H., Aravin, A.A., Stein, P., Girard, A., Murchison, E.P., Cheloufi, S., et al., 2008. Pseudogene-derived small interfering RNAs regulate gene expression in mouse oocytes. *Nature* 453, 534–538.
- Tanaka, M., Chen, Z., Bartunkova, S., Yamasaki, N., Izumo, S., 1999. The cardiac homeobox gene *Csx/Nkx2.5* lies genetically upstream of multiple genes essential for heart development. *Development* 126, 1269–1280.
- Taylor, S.M., Jones, P.A., 1979. Multiple new phenotypes induced in 10T1/2 and 3T3 cells treated with 5-azacytidine. *Cell* 17, 771–779.
- Tevosian, S.G., Deconinck, A.E., Cantor, A.B., Rieffm, H.I., Fujiwaram, Y., Corfas, G., Orkin, S.H., 1999. FOG-2: a novel GATA-family cofactor related to multitype zinc-finger proteins Friend of GATA-1 and U-shaped. *Proc. Natl. Acad. Sci. USA* 96, 950–955.
- The´veniau-Ruissy, M., Dandonneau, M., Mesbah, K., Ghez, O., Mattei, M.G., Miquerol, L., et al., 2008. The del22q11.2 candidate gene *tbx1* controls regional outflow tract identity and coronary artery patterning. *Circ. Res.* 103, 142–148.
- Thomas, T., Yamagishi, H., Overbeek, P.A., Olson, E.N., Srivastava, D., 1998. The bHLH factors, dHAND and eHAND, specify pulmonary and systemic cardiac ventricles independent of left-right sidedness. *Dev. Biol.* 196, 228–236.
- Thomas, N.A., Koudijs, M., van Eeden, F.J., Joyner, A.L., Yelon, D., 2008. Hedgehog signaling plays a cell-autonomous role in maximizing cardiac developmental potential. *Development* 135, 3789–3799.
- Thuerauf, D.J., Hanford, D.S., Glembotski, C.C., 1994. Regulation of rat brain natriuretic peptide transcription. A potential role for GATA-related transcription factors in myocardial cell gene expression. *J. Biol. Chem.* 269, 17772–17775.
- Thum, T., Catalucci, D., Bauersachs, J., 2008. MicorRNAs: novel regulator in cardiac development and disease. *Cardiovasc. Res.* 79, 562–570.
- Tian, Y., Yuan, L., Goss, A.M., Wang, T., Yang, J., Lepore, J.J., et al., 2010. Characterization and in vivo pharmacological rescue of a *wnt2-gata6* pathway required for cardiac inflow tract development. *Dev. Cell* 18, 275–287.
- Tirosh-Finkel, L., Elhanany, H., Rinon, A., Tzahor, E., 2006. Mesoderm progenitor cells of common origin contribute to the head musculature and the cardiac outflow tract. *Development* 133, 1943–1953.
- Togi, K., Yoshida, Y., Matsumae, H., Nakashima, Y., Kita, T., Tanaka, M., 2006. Essential role of *Hand2* in interventricular septum formation and trabeculation during cardiac development. *Biochem. Biophys. Res. Commun.* 343, 144–151.
- Tohsato, Y., Baba, T., Baba, T., Mazaki, Y., Ito, M., Wanner, B.L., et al., 2010. Environmental dependency of gene knockouts on phenotype microarray analysis in *Escherichia coli*. *J. Bioinform. Comput. Biol.* 8 (Suppl. 1), 83–99.
- Treisman, R., 1986. Identification of a protein-binding site that mediates transcriptional response of the *c-fos* gene to serum factors. *Cell* 46, 567–574.

- Treisman, R., 1987. Identification and purification of a polypeptide that binds to the c-fos promoter serum response element. *EMBO J.* 6, 2711–2717.
- Trusolino, L., Bertotti, A., Comoglio, P.M., 2010. MET signalling: principles and functions in development, organ regeneration and cancer. *Nat. Rev. Mol. Cell Biol.* 11, 834–848.
- Tsai, S.F., Martin, D.I., Zon, L.I., D'Andrea, A.D., Wong, G.G., Orkin, S.H., 1989. Cloning of cDNA for the major DNA-binding protein of the erythroid lineage through expression in mammalian cells. *Nature* 339, 446–451.
- Tsuchihashi, T., Maeda, J., Shin, C.H., Ivey, K.N., Black, B.L., Olson, E.N., et al., 2011. Hand2 function in second heart field progenitors is essential for cardiogenesis. *Dev. Biol.* 351, 62–69.
- Tzahor, E., 2009. Heart and craniofacial muscle development: a new developmental theme of distinct myogenic fields. *Dev. Biol.* 327, 273–279.
- Uchida, K., Aramaki, M., Nakazawa, M., Yamagishi, C., Makino, S., Fukuda, K., et al., 2010. Gene knock-outs of inositol 1,4,5-trisphosphate receptors types 1 and 2 result in perturbation of cardiogenesis. *PLoS One* 5, e12500.
- Uchimura, T., Komatsu, Y., Tanaka, M., McCann, K., Mishina, Y., 2009. Bmp2 and bmp4 genetically interact to support multiple aspects of mouse development including functional heart development. *Genesis* 47, 374–384.
- Unwin, R.D.P., Whetton, A.D., 2006. Systematic proteome and transcriptome analysis of stem cell populations. *Cell Cycle* 5, 1587–1591.
- Van der Loop, F.T.L., Schaart, G., Timmer, E.D.J., Ramaekers, F.C.S., van Eys, G.J.J.M., 1996. Smoothelin, a novel cytoskeletal protein specific for smooth muscle cells. *J. Cell Biol.* 134, 401–411.
- van Rooij, E., Sutherland, L.B., Qi, X., Richardson, J.A., Hill, J., Olson, E.N., 2007. Control of stress-dependent cardiac growth and gene expression by a microRNA. *Science* 316, 575–579.
- van Rooij, E., Liu, N., Olson, E.N., 2008. MicroRNAs flex their muscles. *Trends Genet.* 24, 159–166.
- Van Wijk, B., van den Berg, G., Abu-Issa, R., Barnett, P., Van Der Velden, S., Schmidt, M., et al., 2009. Epicardium and myocardium separate from a common precursor pool by crosstalk between bone morphogenetic protein- and fibroblast growth factor-signaling pathways. *Circ. Res.* 105, 431–441.
- Vincent, S.D., Buckingham, M.E., 2010. How to make a heart: the origin and regulation of cardiac progenitor cells. *Curr. Top. Dev. Biol.* 90, 1–41.
- Vincentz, J.W., McWhirter, J.R., Murre, C., Baldini, A., Furuta, Y., 2005. Fgf15 is required for proper morphogenesis of the mouse cardiac outflow tract. *Genesis* 41, 192–201.
- Vindis, C., D'Angelo, R., Mucher, E., Ne'gre-Salvayre, A., Parini, A., Mialet-Perez, J., 2010. Essential role of TRPC1 channels in cardiomyoblasts hypertrophy mediated by 5HT2A serotonin receptors. *Biochem. Biophys. Res. Commun.* 391, 979–983.
- Wada, H., Hasegawa, K., Morimoto, T., Kakita, T., Yanazume, T., Sasayama, S., 2000. A p300 protein as a coactivator of GATA-6 in the transcription of the smooth musclemyosin heavy chain gene. *J. Biol. Chem.* 275, 25330–25335.
- Waldo, K.L., Kumiski, D.H., Wallis, K.T., Stadt, H.A., Hutson, M.R., Platt, D.H., et al., 2001. Conotruncal myocardium arises from a secondary heart field. *Development* 128, 3179–3188.
- Walther, C., Guenet, J.L., Simon, D., Deutsch, U., Jostes, B., Goulding, M.D., et al., 1991. Pax: a murine multigene family of paired box-containing genes. *Genomics* 11, 424–434.
- Wang, D.Z., Olson, E.N., 2004. Control of smooth muscle development by the myocardin family of transcriptional coactivators. *Curr. Opin. Genet. Dev.* 14, 558–566.
- Wang, D., Chang, P.S., Wang, Z., Sutherland, L., Richardson, J.A., Small, E., et al., 2001. Activation of cardiac gene expression by myocardin, a transcriptional cofactor for serum response factor. *Cell* 105, 851–862.