

山本経之(共著)	第I編, 1 創薬に向けての動物モデルの役割と問題点:実践行動薬理学	日本薬理学会、金芳堂、京都、p3-10	2010
山本経之(共著)	第II編, 第4章, 4 脳内報酬系における内因性/外因性カンナビノイドの作用:実践行動薬理学	日本薬理学会、金芳堂、京都、p282-290	2010
特許	山本経之 出願人 九州大学/藤本製薬) 薬物依存症治療剤 Therapeutic agent ((-)-bpap) for drug dependence EP20050809728 特許公報 EP1815853(ヨーロッパ15ヶ国) <span style="float: right;">2009年4月22日</span>		
星薬科大学薬品毒性学教室・教授		鈴木 勉	
Mori T, et al.	Inhibitory effects of SA4053 on the rewarding effects of abused drugs.	Addict. Biol.	In press
Ise Y, et al.	Rewarding effects of ethanol combined with low doses of morphine through dopamine D1 receptors.	J Nippon Med Sch.	In press
森友久, 吉澤一巳, 芝崎 真裕, 鈴木 勉	シグマ1受容体の生体機能と薬物依存における関与	日本アルコール薬物医学雑誌 (in press)	In press
森友久, 鈴木勉	オピオイドの定義	ペインクリニック	In press
Kazuhiro T, et al.	Usefulness of Olanzapine as an Adjunct to Opioid Treatment and for the Treatment of Neuropathic Pain.	Anesthesiology 116(1), 159-169	2012
Torigoe K, et al.	Olanzapine Suppresses the Rewarding and Discriminative Stimulus Effects Induced by Morphine.	Synapse, 66(2), 174-179 <sup>⑨</sup>	2012
Mori T, et al.	Sigma-1 receptor function is critical for both the discriminative stimulus and aversive effects of the kappa-opioid receptor agonist U-50488H.	Addict Biol 17(4), 717-724	2012
Mori T, et al.	Involvement of the Arachidonic Acid Cascade in the Hypersusceptibility to Pentylentetrazole-Induced Seizure during Diazepam Withdrawal.	Biol. Pharm. Bull. , 35(12), 2243-2246	2012
Mori T, et al.	Effects of dronabinol on morphine-Induced dopamine-related behavioral effects in animals.	Synapse, 66(11): 931-937 <sup>⑩</sup>	2012
Yoshizawa K, et al.	Role of dopamine D2 and D3 receptors in mediating the U-50,488H discriminative cue: Comparison with methamphetamine and cocaine.	Addict Biol. 17, 949-955	2012

Kuzumaki N, et al.	Multiple Analyses of G-Protein Coupled receptor(GPCR) Expression in the Development of Gefitinib-Resistance in Transforming Non-Small-Cell Lung Cancer.	PLos One 7(10) e44368	2012
Mori T, et al.	Discriminative stimulus effects of hallucinogenic drugs: a possible relation to reinforcing and aversive effects.	J. Pharmacol. Sci. 120, 70-76	2012
森友久, 鈴木勉	A. オピオイドとは何か 1. オピオイドについて.	ペインクリニック 33, s231-s238	2012
芝崎真裕, 森友久, 鈴木勉	オピオイドと GABA 作動性ニューロン.	Clinical Neuroscience. 30, 12	2012
鈴木勉	オピオイド鎮痛薬のやくりがく No.6	Pharma Tribune 4 (2)	2012
鈴木 勉	疾病と病態生理 改訂第3版	南江堂 3, 426-427 東京	2012
鈴木 勉	薬学用語辞典	日本薬学会東京化学同人刊、東京	2012
鳥越一宏, 吉澤一巳, 成田 年, 鈴木勉(分担執筆)	オピオイド 基礎を知って臨床で使いこなす	垣花 学, 成田 年 克誠堂出版 1, 3-21 東京	2012
Yoshizawa, K, et al.	Enhancement of glutamatergic transmission in the cingulate cortex in response to mild noxious stimuli under a neuropathic pain-like state	Synapse 65(7), 424-432	2011
Takemura Y, et al.	Up-regulation of bradykinin receptors is implicated in the pain associated with caerulein-induced acute pancreatitis.	Synapse 65(7), 608-616	2011
Takemura Y, et al.	Effects of gabapentin on brain hyperactivity related to pain and sleep disturbance under a neuropathic pain-like state using fMRI and brain wave analysis.	Synapse 65(7), 668-676	2011
Rahmadi M, et al.	Sleep disturbance associated with an enhanced orexinergic system induced by chronic treatment with paroxetine and milnacipran.	Synapse 65(7), 652-657	2011
Narita M, et al.	Sleep disturbances in a neuropathic pain-like condition in the mouse are associated with a ltered GABAergic transmission in the cingulate cortex.	Pain 152(6), 1358-1372	2011
Yoshizawa K, et al.	Activation of extracellular signal-regulated kinase is critical for the discriminative stimulus effects induced by U-50488H.	Synapse 65(10), 1052-1061	2011

Imai S, et al.	Possible involvement of prolonging spinal $\mu$ -opioid receptor desensitization in the development of antihyperalgesic tolerance to $\mu$ -opioids under a neuropathic pain-like state.	Synapse 65(9), 962-966	2011
Imai, S.et al.	Change in microRNAs associated with neuronal adaptive responses in the nucleus accumbens under neuropathic pain.	J. Neurosci. 31(43), 15294-15299	2011
Shibasaki, M, et al.	Role of actin depolymerizing factor in the development of methamphetamine-induced place preference in mice.	Eur. J. Pharmacol. 671, 70-78	2011
Tsukahara-Ohsu mi Y, et al.	The kappa opioid receptor agonist SA14867 has antinociceptive and weak sedative effects in models of acute and chronic pain.	Eur. J. Pharmacol. 671, 53-60, 2011	2011
Yamamizu K, et al.	The $\kappa$ opioid system regulates endothelial cell differentiation and pathfinding in vascular development.	Blood 118, 775-586	2011
Shibata K, et al.	The Astrocyte-Targeted Therapy by Bushi for the Neuropathic Pain in Mice.	PLoS One 6(8) e23511	2011
Torigoe, K. et al.	Olanzapine suppresses the rewarding and discriminative stimulus effects induced by morphine.	Synapse, 66(3) 66(2), 174-179 ⑩	2011
Torigoe K, et al.	Usefulness of olanzapine as an adjunct to opioid treatment and for the treatment of neuropathic pain	Anesthesiology. 116(1):159-169	2011
Furuta S ,et al.	Subdiaphragmatic vagotomy induces a functional change in visceral A $\delta$ primary afferent fibers in rats.	Synapse, 66(2), 95-105	2011
成田年	「第20回日本臨床精神神経薬理学会・第40回日本神経薬理学会合同年会」印象記	精神医学 53: 302-303	2011
荒川和彦, ほか	抗がん剤による末梢神経障害の特徴とその作用機序.	日本緩和医療薬学雑誌, 4: 1-13	2011
吉澤一巳, ほか	慢性疼痛治療におけるオピオイド鎮痛薬の適正使用	臨床整形外科 46: 317-325	2011
吉澤一巳, ほか	3.オピオイドの副作用とその対策	ライフ・サイエンス 5: 33-42	2011
鳥越一彦, ほか	神経障害性疼痛の治療・対策.	神経障害性疼痛の治療・対策. 31: 71-75	2011

鈴木 勉	New 薬理学 改訂第6版	田中千賀子/加藤隆一、 南光堂 372-379	2011
鈴木 勉(監修)	毒学教室 より深くより楽しく	伊藤哲朗、学研教育出版	2011
Niikura K, et al.	Neuropathic and chronic pain stimuli downregulate central $\mu$ -opioid and dopaminergic transmission.	Trends Pharmacol Sci, 31,299-305 (2010)	2010
Ikegami D, et al.	Epigenetic modulation at the CCR2 gene correlates with the maintenance of behavioral sensitization to methamphetamine.	Addict Biol, 15, 358-361 ⑫	2010
Narita M, et al.	Implication of dopaminergic projection from the ventral tagmental area to the anterior cingulate cortex in $\mu$ -opioid-induced place preference.	Addict Biol, 15, 434-447	2010
今井哲司, 成田年, 鈴木 勉	側坐核.	分子精神医学, 10, 54-58 (2010)	2010
今井哲司, 成田年, 池上大悟, 田村理絵, 佐伯麻衣, 葛巻直子, 鈴木勉	覚せい剤の依存性形成機序.	医薬ジャーナル, 46, 73-77 (2010)	2010
吉澤一巳, 成田年, 新倉慶一, 今井哲司, 葛巻直子, 鈴木 勉	疼痛下におけるモルヒネの精神依存形成抑制とその機序.	ペインクリニック, 31, 1434-1439 (2010)	2010
鈴木 勉	乱用防止と適正使用.	ファルマシア, 46, 825	2010
富山大学大学院医学薬学研究部・教授		新田淳美	
Yan Y., et al.	Intrastratial gene delivery of GDNF persistently attenuates methamphetamine self-administration and relapse in mice.	Int. J. Neuropsychopharmacol. ⑬	In press
Alkam T., et al.	Evaluation of emotional behaviors in young offspring of C57BL/6J mice after gestational and/or perinatal exposure to nicotine in six different time-windows.	Behav. Brain Res., 239: 80-89.	2013
Yan Y., et al.	Dissociable role of tumor necrosis factor alpha gene deletion in methamphetamine self-administration and cue-induced relapsing behavior in mice.	Psychopharmacology., 221: 427-436. ⑬	2012
Furukawa-Hibi Y., et al.	Absence of SHATI/Nat8l reduces social interaction in mice.	Neurosci Lett., 526: 79-84. ⑭	2012

入江徹美, 新田 淳美, 赤池昭紀:	国立大学法人における模擬患者養成及 び問題立脚型チュートリアル学習の現状.	薬学雑誌, 132; 337-363.	2012
Alkam T, et al.	Evaluation of object-based attention in mice.	Behav Brain Res. 220, 185-193.	2011
Furukawa-Hibi Y, et al.	The hydrophobic dipeptide Leu-Ile inhibits immobility induced by repeated forced swimming via the induction of BDNF.	Behav Brain Res. 220, 271-280.	2011
Furukawa-Hibi Y, et al.	Butyrylcholinesterase inhibitors ameliorate cognitive dysfunction induced by amyloid- $\beta$ in mice.	Behav Brain Res. 225, 222-229.	2011
Nakatani M, et al.	Periocular injection of in situ hydrogels containing Leu-Ile, an inducer for neurotrophic factors, promotes retinal ganglion cell survival after optic nerve injury.	Exp Eye Res. 93, 873-879. ⑮	2011
Oyagia A, et al.	Heparin-binding EGF-like growth factor is required for synaptic plasticity and memory formation.	Brain Res. 1419, 97-104.	2011
新田 淳美	－依存症の現状－アルコールおよび向精 神薬の乱用について	ファルマシア, 47 巻, 839-843.	2011
Ibi D, et al.	Combined effect of neonatal immune activation and mutant DISC1 on phenotypic changes in adulthood.	Behav Brain Res. 206:32-37.	2010
Ibi D, et al.	Piccolo knockdown-induced impairments of spatial learning and long-term potentiation in the hippocampal CA1 region.	Neurochem Int. 56:77-83.	2010
Ohki M, et al.	Tissue type plasminogen activator regulates myeloid-cell dependent neovascularization during tissue regeneration.	Blood. 115:4302-4312.	2010
Katsuno M, et al.	Disrupted TGF-beta signaling in spinal and bulbar muscular atrophy.	J Neurosci. 30:5702-5712.	2010
Ariyannur PS, et al.	Methamphetamine-induced neuronal protein NAT8L is the NAA biosynthetic enzyme: implications for specialized acetyl coenzyme A metabolism in the CNS.	Brain Res. 1335:1-13.	2010
Alkam T, et al.	Oral supplementation with Leu-Ile, a hydrophobic dipeptide, prevents the impairment of memory induced by amyloid beta in mice via restraining the hyperphosphorylation of extracellular signal-regulated kinase.	Behav Brain Res. 210:184-190. ⑯	2010

Yun J, et al.	Chronic restraint stress impairs neurogenesis and hippocampus-dependent fear memory in mice: possible involvement of a brain-specific transcription factor Npas4.	J Neurochem. 114:1840-1851.	2010
Furukawa-Hibi Y, et al.	Over expression of Piccolo C2A domain induces depression-like behavior in mice.	Neuro Report 21:1177-1181.	2010
新田 淳美 日比陽子 宮本嘉明 鍋島俊隆	薬物依存におけるピッコロの役割	日本アルコール・薬物医学会雑誌 45 巻 6 号, 525-529	2010
特許	新田淳美、宇野恭介 精神疾患関連遺伝子 shati/nat81 の遺伝子修飾による精神疾患診断方法 2013年2月19日		
	新田淳美、丹羽美苗、鍋島俊隆 精神障害関連遺伝子及びその利用 2012年3月1日	取得特許 (特願 2007-505883)	
	新田淳美 精神障害の診断方法および診断薬キット (特願 2011-126100) 2011年6月6日		
京都大学大学院医学研究科メディカルイノベーションセンター・特定准教授		疋田 貴俊	
Hikida T, et al.	Pathway-specific modulation of nucleus accumbens in reward and aversive behavior via selective transmitter receptors.	Proc Natl Acad Sci USA. 110:342-347 ⑰	2013
NiwaM, et al.	Adolescent stress-induced epigenetic control of dopaminergic neurons via glucocorticoids.	Science. 339:335-339	2013
Yawata S, et al.	Pathway-specific control of reward learning and its flexibility via selective dopamine receptor in the nucleus accumbens.	Proc Natl Acad Sci USA. 109:12764-12769 ⑱	2012
Hikida T, Gamo NJ, Sawa A:	DISC1 as a therapeutic target for mental illnesses	Expert Opin Ther Targets. 16:1151-1160	2012
Kimura K, et al.	Pathway-specific engagement of ephrinA5-EphA4/EphA5 system of the substantia nigra pars reticulata in cocaine-induced responses.	Proc Natl Acad Sci USA, 108: 9981-9986 ⑲	2011
Hikida, T., et al.	Distinct roles of synaptic transmission in direct and indirect striatal pathways to reward and aversive behavior.	Neuron, 66: 896-907 ⑳	2010
疋田貴俊	大脳基底核の直接路と間接路の機能分離: Annual Review 神経 2013 鈴木則宏、祖父江元、荒木信夫、宇川義	中外医学社、東京、pp.10-16	2013

	一、川原信隆編、		
疋田貴俊 ほか	精神疾患モデル動物の作製と治療法スクリーニング	実験医学増刊 30: 2074-2079	2012
疋田貴俊	精神疾患の分子遺伝学：最近の知見。	脳神経外科速報, 21: 1250-1254	2011
疋田貴俊、神谷篤	精神疾患モデル動物の可能性 – 遺伝子から神経回路へ。	実験医学, 28: 2205-2210	2010
特許	疋田貴俊・中西重忠 大脳基底核神経回路の神経伝達を解析する方法 特願 2012-512542 (国内移行)出願日 2012.10.22 2010年4月30日国際出願(PCT/JP2010/003089)		
名城大学薬学部・助教		間宮隆吉	
Koseki T, et al.	Galantamine attenuates reinstatement of cue-induced methamphetamine-seeking behavior in mice.	Addict Biol.	In press
Alkam T, et al.	Evaluation of emotional behaviors in young offspring of C57BL/6J mice after gestational and/or perinatal exposure to nicotine in six different time-windows.	Behav Brain Res. 239:80-89.	2013
Mouri A, et al.	Mouse strain differences in phencyclidine-induced behavioural changes.	Int J Neuropsychopharmacol. 15(6):767-779.	2012
Koseki T, et al.	Exposure to enriched environments during adolescence prevents abnormal behaviours associated with histone deacetylation in phencyclidine-treated mice.	Int J Neuropsychopharmacol. 15(10):1489-1501.	2012
Toriumi K, et al.	Prenatal NMDA receptor antagonism impaired proliferation of neuronal progenitor, leading to fewer glutamatergic neurons in the prefrontal cortex.	Neuropsychopharmacology. 37(6):1387-1396. ①	2012
Mouri A, et al.	MAGE-D1 regulates expression of depression-like behavior through serotonin transporter ubiquitylation.	J Neurosci. 32(13):4562-4580.	2012
Imai S, et al.	Ubiquitin-specific peptidase 46 (Usp46) regulates mouse immobile behavior in the tail suspension test through the GABAergic system.	PLoS One. 7(6):e39084.	2012
Lu L, et al.	Prenatal exposure to PCP produces behavioral deficits accompanied by the overexpression of GLAST in the prefrontal cortex of postpubertal mice.	Behav Brain Res. 220(1):132-139 ②	2011

Lu L, et al.	Genetic animal models of schizophrenia related with the hypothesis of abnormal neurodevelopment.	Biol Pharm Bull.34, 1358-1363 ㉓	2011
特許など	小鹿一、間宮隆吉、鍋島俊隆 「新規ステロイド誘導体及びその製造方法、並びにその新規ステロイド誘導体を含有する医薬」 出願番号:PCT/JP2010/052656		
東北大学大学院医学系研究科・教授		曾良一郎	
Yamashita M, et al.	Impaired Cliff Avoidance Reaction in Dopamine Transporter Knockout Mice	Psychopharmacology	in press
Ohara A, et al.	Exclusive expression of VMAT2 in noradrenergic neurons increases viability of homozygous VMAT2 knockout mice.	Biochemical and Biophysical Research Communications	in press
Nishizawa D, et al.	Genome-wide association study identifies a potent locus associated with human opioid sensitivity	Molecular Psychiatry	in press
Arime Y, et al.	Cortico-subcortical neuromodulation involved in the amelioration of prepulse inhibition deficits in dopamine transporter knockout mice.	Neuropsychopharmacology 37(11): 2522-2530 ㉔	2012
Uchiumi O, et al.	Serotonergic involvement in the amelioration of behavioral abnormalities in dopamine transporter knockout mice by nicotine.	Neuropharmacology 64(1): 348-356	2012
榊原泰史, 曾良一郎.	脳に働く“くすり”のメカニズム.	教育と医学. 60(10):31-38 ㉕	2012
Kishi T, et al	Serotonin 6 receptor gene is associated with methamphetamine-induced psychosis in a Japanese population.	Drug Alcohol Depend. 113(1):1-7	2011
Kobayashi H, et al.	Association analysis of the Adenosine A1 receptor gene polymorphisms in patients with methamphetamine dependence/psychosis.	Current Neuropharmacology 9:137-142	2011
Kobayashi H, et al.	Association analysis of the Tryptophan Hydroxylase 2 gene polymorphisms in patients with methamphetamine dependence/psychosis.	Current Neuropharmacology 9:176-182	2011
Ide S, et al.	(-)-Pentazocine induces visceral chemical antinociception, but not thermal, mechanical, or somatic chemical antinociception, in mu-opioid receptor knockout mice.	Molecular Pain 7(1):23	2011
Yoshimura T, et al.	Association analysis of the GDNF gene with methamphetamine use disorder in a Japanese population.	Prog Neuropsychopharmacol Biol Psychiatry. 35(5):1268-72	2011

Yamamoto H, et al	Mop reduction during long-term methamphetamine withdrawal was restored by chronic post-treatment with fluoxetine.	Current Neuropharmacology 9:73-78	2011
Hagino Y, et al.	Effect of MDMA on extracellular dopamine and serotonin levels in mice lacking dopamine and/or serotonin transporters.	Current Neuropharmacology 9:91-95	2011
Kasai S, et al.	Quantitative detection of m opioid receptor: Western blot analyses using m opioid receptor knockout mice.	Current Neuropharmacology 9:219-222	2011
Komatsu H, et al.	Decreased response to social defeat stress in $\mu$ -opioid-receptor knockout mice.	Pharmacology Biochemistry & Behavior 99: 676-682	2011
Sogawa C, et al.	Methylone and monoamine transporters: correlation with toxicity.	Current Neuropharmacology 9: 58-62	2011
Okahisa Y, et al.	Association between the Regulator of G-protein Signaling 9 Gene and Patients with Methamphetamine Use Disorder and Schizophrenia.	Curr Neuropharmacol 9(1):190-194	2011
Okahisa Y, et al.	Association Study of Two Cannabinoid Receptor Genes, CNR1 and CNR2, with Methamphetamine Dependence.	Curr Neuropharmacol 9(1):183-189	2011
Yokobayashi E, et al.	Association study of serine racemase gene with methamphetamine psychosis.	Curr Neuropharmacol 9(1):169-175	2011
Ujike H, et al.	Association Between 5HT1b Receptor Gene and Methamphetamine Dependence.	Curr Neuropharmacol 9(1):163-168 <sup>(26)</sup>	2011
Tsunoka T, et al.	No Association Between GRM3 and Japanese Methamphetamine-Induced Psychosis.	Curr Neuropharmacol 9(1):160-162	2011
Okumura T, et al.	Genetic Association Analysis of NOS1 and Methamphetamine-Induced Psychosis Among Japanese.	Curr Neuropharmacol 9(1):155-159	2011
Okochi T, et al.	Genetic Association Analysis of NOS3 and Methamphetamine-Induced Psychosis Among Japanese.	Curr Neuropharmacol 9(1):151-154	2011
Kishi T, et al.	Lack of association between prokineticin 2 gene and Japanese methamphetamine dependence.	Curr Neuropharmacol 9(1):133-136	2011
Kishi T, et al.	Association Analysis of Nuclear Receptor Rev-erb Alpha Gene (NR1D1) and Japanese Methamphetamine Dependence.	Curr Neuropharmacol 9(1):129-132	2011
Kishi T, et al.	No significant association between SIRT1 gene and methamphetamine-induced psychosis in the Japanese population.	Hum Psychopharmacol 2011 26(7):445-450	2011
Arime Y, et al.	Animal models of attention-deficit/hyperactivity disorder.	Biol Pharm Bull. 34(9),1373-1376.	2011
曾良一郎.	カテコールアミン神経伝達。認知症学 (上) その解明と治療の最新知見	日本臨床 69 増刊号 8; 224-227	2011

曾良一郎.	シンポジウム特集「認知機能障害に対する治療をどう評価するか」.	日本神経精神薬理学雑誌 31;239	2011
曾良一郎.	ドーパミン受容体 Dopamine receptor. ストレス科学辞典.	日本ストレス学会/(財)パブリックヘルスリサーチセンター監修:764, 実務教育出版, 東京	2011
曾良一郎, 氏家寛.	物質依存の神経化学.	脳とこころのプライマリケア第8巻 依存(福居顯二編集):50-59, シナジー, 東京	2011
曾良一郎.	LSD-25. 現代精神医学事典	加藤敏, 神庭重信, 中谷陽二, 武田雅俊, 鹿島晴雄, 狩野力八郎, 市川宏伸編:116, 弘文堂, 東京	2011
曾良一郎.	逆耐性現象. 現代精神医学事典	加藤敏, 神庭重信, 中谷陽二, 武田雅俊, 鹿島晴雄, 狩野力八郎, 市川宏伸編):215, 弘文堂, 東京	2011
曾良一郎.	受容体. 現代精神医学事典	加藤敏, 神庭重信, 中谷陽二, 武田雅俊, 鹿島晴雄, 狩野力八郎, 市川宏伸編:470, 弘文堂, 東京	2011
曾良一郎.	ドーパミン. ドーパミン仮説, 現代精神医学事典	加藤敏, 神庭重信, 中谷陽二, 武田雅俊, 鹿島晴雄, 狩野力八郎, 市川宏伸編:772-773, 弘文堂, 東京	2011
曾良一郎.	モノアミン仮説. 現代精神医学事典	加藤敏, 神庭重信, 中谷陽二, 武田雅俊, 鹿島晴雄, 狩野力八郎, 市川宏伸編):1021-1022, 弘文堂, 東京	2011
Frye CA, et al.	Progesterone reduces hyperactivity of female and male dopamine transporter knockout mice. (2010)	Behav Brain Res. 209(1):59-65	2010
Kishi T, et al.	Serotonin 1A receptor gene is associated with Japanese methamphetamine-induced psychosis patients.	Neuropharmacology. 58(2):452-456 ㉗	2010
Li B, et al.	Impaired spatial working memory and decreased frontal cortex BDNF protein level in dopamine transporter knock out mice. (2010)	Eur J Pharmacol. 25:628(1-3):104-107	2010
Ide S, et al.	Antidepressant-like effect of venlafaxine is abolished in $\mu$ -opioid receptor-knockout mice.	Journal of Pharmacological Sciences. 114(1):107-110	2010
Arai M, et al.	Enhanced carbonyl stress in a subpopulation of schizophrenia. (2010)	Arch Gen Psychiatry. 67(6):589-97	2010

Tsunoka T, et al.	Association analysis of GRM2 and HTR2A with methamphetamine-induced psychosis and schizophrenia in the Japanese population.	Prog Neuropsychopharmacol Biol Psychiatry. 34(4):639-44.	2010
Okuyama K, et al.	The Involvement of micro-Opioid Receptors in the Central Nervous System in the Worsening of Allergic Airway Inflammation by Psychological Stress in Mice.(2010)	Int Arch Allergy Immunol. 152(4):342-352	2010
Kobayashi H, et al.	The adenosine A2A receptor is associated with methamphetamine dependence/psychosis in the Japanese population	Behav Brain Funct. 6:50(2010)	2010
Kishi T, et al.	PROKR2 is associated with methamphetamine dependence in the Japanese population.	Prog Neuropsychopharmacol Biol Psychiatry.34(6):1033-6	2010
Ide S, et al.	Combination of cell culture assays and knockout mouse analyses for the study of opioid partial agonism.	Pain and Analgesia.Methods Mol Biol. 617:363-74	2010
Kishi T, et al.	Serotonin 6 receptor gene is associated with methamphetamine-induced psychosis in a Japanese population.	Drug Alcohol Depend. 113(1):1-7	2011
Kobayashi H, et al.	Association analysis of the Tryptophan Hydroxylase 2 gene polymorphisms in patients with methamphetamine dependence / psychosis.	Current Neuropharmacology 9:176-182	2011
曾良一郎.	薬物依存の脆弱性要因.	実験医学 28(5):38-42	2010
曾良一郎, 佐々木一益.	痛みがあると薬物依存になりにくいというのは本当か?.	Clinical Neuroscience 別冊 28(5); 581	2010
曾良一郎, 内海修.	シナプスの病態 薬物依存症.	Clinical Neuroscience 28(8); 982-930	2010
笠原好之, 有銘預世布, 福井麻美, 内海修, 曾良一郎.	コカイン依存研究の動向. 薬物依存の臨床各論—最新動向—,	日本臨床 68(8); 1479-1485	2010
千葉大学大学院医学研究院・教授		伊豫 雅臣 (研究協力者 関根吉統、大宮宗一郎)	
Kimura H, et al.	Potential treatment strategy of risperidone in long-acting injectable form for schizophrenia with dopamine supersensitivity psychosis.	Schizophr Res.	In press
Kanahara N, et al.	Orbitofrontal cortex abnormality and deficit schizophrenia.	Schizophr Res. 143(2-3):246-252.	2013

Okamoto, H., et al.	Orengedoku-to augmentation in cases showing partial response to yokukan-san treatment: a case report and literature review of the evidence for use of these Kampo herbal formulae.	Neuropsychiatr Dis Treat. 9, 151-155.	2013
Tadokoro S, et al.	Chronic treatment with aripiprazole prevents development of dopamine supersensitivity and potentially supersensitivity psychosis.	Schizophr Bull. 38(5):1012-1020. ㉘	2012
Chen, H., et al.	Protective effects of the antioxidant sulforaphane on behavioral changes and neurotoxicity in mice after the administration of methamphetamine.	Psychopharmacology (Berl). 222, 37-45. ㉙	2012
Yoshida T, et al.	Decreased serum levels of mature brain-derived neurotrophic factor (BDNF), but not its precursor proBDNF, in patients with major depressive disorder.	PLoS One. 7(8):e42676.	2012
Fujita, Y., et al.	The antibiotic minocycline prevents methamphetamine-induced rewarding effects in mice.	Pharmacol Biochem Behav. 101, 303-306.	2012
Sasaki T, et al.	Treatment of refractory catatonic schizophrenia with low dose aripiprazole.	Ann Gen Psychiatry. 11(1):12.	2012
Niitsu T, et al..	A randomized, double-blind, placebo-controlled trial of fluvoxamine in patients with schizophrenia: a preliminary study.	J Clin Psychopharmacol. 32(5):593-601.	2012
Niitsu, T, Iyo, M, Hashimoto, K. (REVIEW)	Sigma-1 receptor agonists as therapeutic drugs for cognitive impairment in neuropsychiatric diseases.	Curr Pharm Des. 18, 875-883.	2012
大宮宗一郎 ほか	未成年の物質使用のリスクとなりうるパーソナリティ.	日本アルコール薬物医学会雑誌. 46, 287-297.	2012
Niitsu, T., et al.	Associations of serum brain-derived neurotrophic factor with cognitive impairments and negative symptoms in schizophrenia. Prog Neuropsychopharmacol.	Biol Psychiatry. 35, 1836-1840.	2011
Okamoto H, et al..	A suspected case of somatoform disorder successfully treated with an herbal medicine.	J Altern Complement Med. 17(2):171-173.	2011
Shiina A, et al.	Expert consensus on hospitalization for assessment: a survey in Japan for a new forensic mental health system.	Ann Gen Psychiatry. 8;10:11.	2011
Tadokoro, S., et al.	Fluvoxamine may prevent onset of psychosis: a case report of a patient at ultra-high risk of psychotic disorder.	Ann Gen Psychiatry. 10:26	2011

Yoshida T, et al.	Criterion and construct validity of the CogState Schizophrenia Battery in Japanese patients with schizophrenia.	PLoS One. 6(5):e20469.	2011
Arakawa, S. et al.	Minocycline produced antidepressant-like effects on the learned helplessness rats with alterations in levels of monoamine in the amygdala and no changes in BDNF levels in the hippocampus at baseline.	Pharmacol Biochem Behav. 100, 601-606.	2011
Haraguchi T, et al.	Attitudes of Japanese psychiatrists toward forensic mental health as revealed by a national survey.	Psychiatry Clin Neurosci. 65(2):150-157.	2011
Fukami G, et al.	Effects of etizolam and ethyl loflazepate on the P300 event-related potential in healthy subjects.	Ann Gen Psychiatry. 9:37.	2010
Ishii D, et al.	D-serine enhances extinction of auditory cued fear conditioning via ERK1/2 phosphorylation in mice.	Prog Neuropsychopharmacol Biol Psychiatry., 34, 639-644.	2010
Niitsu T, et al.	Fluvoxamine improved cognitive impairments in a patient with schizophrenia.	Prog Neuropsychopharmacol Biol Psychiatry., 34, 1345-1346.	2010
Tanibuchi Y, et al.	A case of methamphetamine use disorder treated with the antibiotic drug minocycline.	Gen Hosp Psychiatry., 32, 559.e1-3. ㊦	2010
Tanibuchi Y, et al.	Characterization of [(3)H]CHIBA-1001 binding to alpha7 nicotinic acetylcholine receptors in the brain from rat, monkey, and human.	Brain Res., 1348, 200-208.	2010
東京医科歯科大学大学院・教授		西川 徹	
Umezaki, Y., et al	Brain perfusion asymmetries in patients with oral somatic delusions.	Eur Arch Psychiatry Clin Neurosci.	in press
Ishiwata, S., et al	Modulation of extracellular d-serine content by calcium permeable AMPA receptors in rat medial prefrontal cortex as revealed by in vivo microdialysis.	Int J Neuropsychopharmacol	in press
Ishiwata, S., et al	Increasing effects of S-methyl-L-cysteine on the extracellular D-serine concentrations in the rat medial frontal cortex.	Amino Acids.	in press
Kurumaji, A. et al.	An angiogenic drug, FG 7142, induced an increase in mRNA of Btg2 and Adamts1 in the hippocampus of adult mice.	Behav Brain Funct., 8, 8, 43	2012
Uezato, A., et al.	Further evidence for a male-selective genetic association of synapse-associated protein 97 (SAP97) gene with schizophrenia.	Behavioral and Brain Functions. , 8, 2	2012

Hayashi, H., et al.	Sorting nexin 27 interacts with multidrug resistance-associated protein 4 (MRP4) and mediates the internalization of MRP4.	J Biol Chem., 287,15054-15065	2012
Uezato, A., et al..	Improvement of asymmetrical temporal blood flow in refractory oral somatic delusion after successful electroconvulsive therapy.	J ECT. , 28, 50-51 ㉑	2012
Sasaki, T., et al.	Quantification of Dopamine Transporter in Human Brain Using Positron Emission Tomography with 18F-FE-PE2I.	J Nuc Med., 53, 1065-1073 ㉒	2012
Ozaki, A., et al.	Donepezil-induced sleep spindle in a patient with dementia with Lewy bodies.	Psychogeriatrics., 2, 255-228	2012
武田充弘, 川俣光太郎, 石橋哲, 西川徹, 車地暁生.	パーキンソン病の幻覚・妄想状態に aripiprazole が著効した一症例.	精神科, 20, 343-349	2012
Nishida M, et al.	Memory deficits due to brain injury: unique PET findings and dream alterations.	BMJ Case Reports, doi:10.1136/bcr.09.2011.4845	2011
Kurumaji A, et al.	Effects of novelty stress on hippocampal gene expression, corticosterone and motor activity in mice.	Neuroscience Research 71:161-167.	2011
Nishikawa T.	Analysis of free D-serine in mammals and its biological relevance.	J Chromatogr B Analyt Technol Biomed Life Sci. 879:3169-3183. ㉓	2011
吉池卓也, ほか	Aripiprazole のせん妄に対する有用性.	精神医学. 53: 543-549.	2011
西川 徹.	統合失調症の臨床と基礎の最前線. Medical Science Digest, 37:14(402)-17(405).	日本神経精神薬理学雑誌 27: 53-56	2011
上里 彰仁, 西川 徹.	統合失調症の病態メカニズム.	ファルマシア 47: 824-828.	2011
治徳 大介, 西川 徹, 吉川武男.	統合失調症における遺伝と環境要因.	精神科治療学 26:1355-1362.	2011
西川 徹	薬の作用メカニズムから見た統合失調症の病態. 脳(ブレイン)バンクー精神疾患の謎を解くために一.	加藤忠史&ブレインバンク委員会(編集)、光文社、東京、pp. 184-193.	2011
Okamoto N, et al	Rapid antidepressant effect of ketamine anesthesia during electroconvulsive therapy of treatment-resistant depression-Open-label trial comparing ketamine and propofol anesthesia.	J ECT. 26:223-227	2010
Oshima K, et al	Reliability and diagnostic validity for schizophrenia of Japanese version of the Bonn Scale for Assessment of Basic Symptoms (BSABS).	J Med Dent Sci. 57:83-94	2010

Kuroda Y, et al.	Chronic repetitive transcranial magnetic stimulation failed to change dopamine synthesis rate: Preliminary L-[b-11C]DOPA positron emission tomography study in patients with depression.	Psychiatry and Clinical Neurosciences. 64:659-662	2010
Shioiri A, et al.	White matter abnormalities as a risk factor for postoperative delirium revealed by diffusion tensor imaging.	Am J Geriatr Psychiatry. 18:743-753 ㉔	2010
佐々木健至, 行実知昭, 熱田英範, 石川洋世, 吉池卓也, 竹内崇, 大島一成, 山本直樹, 車地暁生, 西川 徹	幻覚妄想状態を呈したチアミン欠乏症の1例—チアミン血中濃度と治療反応性—	精神神経学雑誌. 112(2):97-110	2010
竹内 崇, 西川 徹	Perospirone と気分安定薬の併用により病相予防が可能となった双極性障害の2症例.	精神科治療学. 25(9):1257-1262	2010
藤田宗久, 上里彰仁, 川上礼子, 竹内 崇, 西川 徹, 車地暁生	松果体・鞍上部 germinoma 治療経過中に緊張病、その後躁状態を繰り返した1症.	精神科. 17(6):643-647	2010
西川 徹	NMDA 受容体-D-セリン系を標的とした新規統合失調症治療薬の開発. 特集: 統合失調症グルタミン酸系治療薬の臨床開発と基礎研究	日本神経精神薬理学雑誌 29:201-206	2010
西多昌規, 西川 徹	神経診察法の基本とピットフォール 精神症状と兆候	Clinical Neuroscience. 28:1340-1341	2010
成島健二, 西川 徹	新薬展望 2010 第 III 部 治療における最近の新薬の位置付け(薬効別) —新薬の広場— 統合失調症治療薬・抗精神病薬	医薬ジャーナル増刊号 新薬展望 46(S-1):330(540)-341(551)	2010
西川 徹	Ⅲ精神薬理学 3. D-セリンと統合失調症: 専門医のための精神科臨床リュミエール 16 脳科学エッセンシャル—精神疾患の生物学的理解のために	神庭重信・加藤忠史(編) 中山書店 東京 209-211	2010
公益財団法人東京都医学総合研究所・参事研究員		池田 和隆	
Ohara A, et al.	Exclusive expression of VMAT2 in noradrenergic neurons increases viability of homozygous VMAT2 knockout mice.	Biochem Biophys Res Commun	in press.
Kasai S, et al.	Reduced supraspinal nociceptive responses and distinct gene expression profile in CXBH recombinant inbred mice.	J Pain	in press.
Shinohara M, et al.	ADORA2A polymorphism predisposes children to encephalopathy with febrile status epilepticus.	Neurology	in press.
Nishizawa D, et al.	Genome-wide association study identifies a potent locus associated with human opioid sensitivity.	Mol Psychiatry ㉕	in press.

菅谷渚, ほか	依存重症度評価ツールの臨床応用.	精神神経学雑誌	in press.
Moriyama A, et al.	Association between genetic polymorphisms of the beta1-adrenergic receptor and sensitivity to pain and fentanyl in patients undergoing painful cosmetic surgery.	J Pharmacol Sci 121:48-57.	2012
Sato A, et al.	Rapamycin reverses impaired social interaction in mouse models of tuberous sclerosis complex.	Nat Commun ncomms 2295. (DOI: 10.1038)	2012
Ella E, et al.	Association between dopamine beta hydroxylase rs5320 polymorphism and smoking behaviour in elderly Japanese.	J Hum Genet 57:385-390.	2012
Sugaya N, et al.	Influence of GIRK channel inhibition on relapse risk in Japanese alcohol-dependent inpatients.	Jpn J Neuropsychopharmacol 32:165-167. ㊟	2012
沼尻真貴, ほか	Beta-アドレナリン受容体遺伝子多型と性格特性との関連研究.	日本神経精神薬理学雑誌 32:227-231.	2012
沼尻真貴, ほか	双極性障害のリチウム感受性に対する GSK-3Beta 遺伝子多型の影響.	日本神経精神薬理学雑誌 32:161-163.	2012
林田真和, ほか	遺伝子多型とオピオイド感受性—mu-オピオイド受容体遺伝子 A118G 多型を中心に—.	臨床麻酔 36(11):1591-1598.	2012
林田真和, ほか	遺伝子多型が疼痛治療におけるオピオイド必要量に及ぼす影響—mu-オピオイド受容体遺伝子の A118G 多型を中心として.	Practice of Pain Management 3:174-177.	2012
菅谷渚, 池田和隆	依存症. In: 脳科学辞典 加藤忠史編, <a href="http://bsd.neuroinf.jp/wiki/依存症">bsd.neuroinf.jp/wiki/依存症</a> .	MediaWiki.	2012
菅谷渚, 池田和隆	覚せい剤. In: 脳科学辞典 加藤忠史編, <a href="http://bsd.neuroinf.jp/wiki/覚せい剤">bsd.neuroinf.jp/wiki/覚せい剤</a> .	MediaWiki.	2012
池田和隆	幻覚剤あれこれ. In: TEXT 精神医学: 14 精神作用物質による精神および行動の障害 加藤進昌, 神庭重信, 笠井清登 編	南山堂, 東京, pp211.	2012
Ogai Y, Chin F, Ikeda K.:	Assessment of craving to predict relapse in patients with substance abuse/dependence. In: Advances in Psychology Research. Vol. 88 Alexandra M. Columbus, ed	Nova Science Publishers, New York, pp225-234.	2012

Berrocoso E, et al.	Active behaviours produced by antidepressants and opioids in the mouse tail suspension test.	Int J Neuropsychopharmacol. 5:1-12.	2012
井手聡一郎, 南雅文, 池田和隆	痛み感受性の遺伝的要因.	ペインクリニック. 33(1):67-74.	2012
Fukuda K, et al.	Diversity of opioid requirements for postoperative pain control following oral surgery - Is it affected by polymorphism of the mu-opioid receptor?	Anesth Prog. 57:145-149.	2011
Nishizawa D, et al.	Identification of selective agonists and antagonists to G protein-activated inwardly rectifying potassium channels: candidate medicines for drug dependence and pain.	Curr Neuropharmacol. 9(1):113-117.	2011
Kasai S, et al.	Quantitative detection of mu opioid receptor: Western blot analyses using mu opioid receptor knockout mice.	Curr Neuropharmacol. 9(1):219-222.	2011
Kobayashi T, et al.	Inhibition of G protein-activated inwardly rectifying K <sup>+</sup> channels by phencyclidine.	Curr Neuropharmacol. 9(1):244-246.	2011
Takamatsu Y, et al.	The selective serotonin reuptake inhibitor paroxetine, but not fluvoxamine, decreases methamphetamine conditioned place preference in mice.	Curr Neuropharmacol. 9(1):68-72. ③	2011
Yamamoto H, et al.	MOP reduction during long-term methamphetamine withdrawal was restored by chronic post-treatment with fluoxetine.	Curr Neuropharmacol. 9(1):73-78.	2011
Hagino Y, et al.	Effects of MDMA on extracellular dopamine and serotonin levels in mice lacking dopamine and/or serotonin transporters.	Curr Neuropharmacol. 9(1):91-95.	2011
Takamatsu Y, et al.	Enhanced hyperthermia induced by MDMA in parkin knockout mice.	Curr Neuropharmacol. 9(1):96-99.	2011
Sugaya N, et al.	Family dysfunction differentially affects alcohol dependence and drug dependence: a view from the Addiction Severity Index in Japan.	Int J Environ Res Public Health. 8:3922-3937.	2011
Ide S, et al.	(-)-Pentazocine induces visceral chemical antinociception, but not thermal, mechanical, or somatic chemical antinociception, in mu-opioid receptor knockout mice.	Molecular Pain. 7:23.	2011
Han W, et al.	Inhibitory role of inducible cAMP early repressor (ICER) in methamphetamine-induced locomotor sensitization.	PLoS ONE. 6:e21637.	2011

Kobayashi T, et al.	Inhibition of G protein-activated inwardly rectifying K <sup>+</sup> channels by different classes of antidepressants.	PLoS ONE. 6:e28208.	2011
Kasai S, et al.	Pharmacogenomics of the human mu-opioid receptor.	Pharmacogenomics. 12:1305-1320.	2011
Aoki J, et al.	Association study on catechol-O-methyltransferase (COMT) Val158Met gene polymorphism and NEO-FFI.	Psychiatry Res. 187:312-313.	2011
池田和隆	依存性薬物作用の解明が拓く新しい精神医学.	精神医学. 53(12):1189-1194.	2011
池田和隆	脳内報酬系の分子メカニズム.	日本神経精神薬理学雑誌 31:263-266.	2011
Han W, et al.	Reduced locomotor sensitization induced by methamphetamine and altered gene expression in ICER overexpressing mice.	日本神経精神薬理学雑誌. 31:79-80.	2011
笠井慎也, ほか	野生由来近交系マウス系統における Oprm1 遺伝子多型とモルヒネ感受性の関連性.	日本神経精神薬理学雑誌. 31:87-88.	2011
Ogai Y, et al.	Influence of GIRK channel inhibition on alcohol abstinence and relapse risk in Japanese alcohol-dependent outpatients.	日本神経精神薬理学雑誌. 31:95-96.	2011
菅谷渚, 池田和隆	報酬系における GIRK チャンネルの役割.	日本生物学的精神医学会誌. 22(4):263-268.	2011
池田和隆	薬物療法. In: 脳とこころのプライマリケア8 依存	福居顯二編, 株式会社シナジー, 東京, pp464-475.	2011
Narita M, et al.	Implication of dopaminergic projection from the ventral tegmental area to the anterior cingulate cortex in mu-opioid-induced place preference.	Addict Biol 15:434-447.	2010
Sora I, et al.	Transgenic mice in the study of drug addiction and the effects of psychostimulant drugs.	Ann. N.Y. Acad. Sci. 1187:218-246.	2010
Ide S, et al.	Combination of cell culture assays and knockout mouse analyses for the study of opioid partial agonism. In: Methods in Molecular Biology	Arpad Szallasi, ed, The humana press Inc, Totowa, pp363-374.	2010
Nishizawa D, et al.	Genetic polymorphisms and human sensitivity to opioid analgesics. In: Methods in Molecular Biology	Arpad Szallasi, ed, The humana press Inc, Totowa, pp395-420.	2010

Aoki J, et al.	The association between personality, pain threshold and a single nucleotide polymorphism (rs3813034) in the 3'-untranslated region of the serotonin transporter gene (SLC6A4).	J Clin Neurosci 17:574-578.	2010
Shiotsuki H, et al.	A rotarod test for evaluation of motor skill learning.	J Neurosci Methods 189:180-185.	2010
Ide S, et al.	Reduced emotional and corticosterone responses to stress in mu-opioid receptor knockout mice.	Neuropharmacol 58:241-247.	2010
Kobayashi T, et al.	Inhibition of G protein-activated inwardly rectifying K <sup>+</sup> channels by the selective norepinephrine reuptake inhibitors atomoxetine and reboxetine.	Neuropsychopharm acology 35:1560-1569.	2010
Aoki J, et al.	Association between 5-hydroxytryptamine 2A receptor gene polymorphism and postoperative analgesic requirements after major abdominal surgery.	Neurosci Lett 479:40-43.	2010
Hagino Y, et al.	Essential role of NMDA receptor channel epsilon4 subunit (GluN2D) in the effects of phencyclidine, but not methamphetamine.	PLoS ONE 5:e13722.	2010
Ide S, et al.	Antidepressant-like effect of venlafaxine is abolished in mu-opioid receptor-knockout mice.	Pharmacol Sci 114:107-110.	2010
Koide T, et al.	Advantage of using wild-derived mouse strains for a variety of pain-related studies: Genetic diversity and new genetic tools. In: Acute Pain	Sam D' Alonso, Katherine L. Grasso, ed, Nova Science Publishers, New York, pp79-99.	2010
Kobayashi D,	Association between analgesic requirements after major abdominal surgery and polymorphisms of the opioid metabolism-related gene ABCB1. In: Acute Pain	Sam D' Alonso, Katherine L. Grasso, ed, Nova Science Publishers, New York, pp101-110.	2010
山本秀子, 萩野洋子, 池田和隆	MDMA などの違法薬物の依存形成機序.	医薬ジャーナル 46(7):95-98.	2010
林田眞和, 池田和隆	ミューオピオイド受容体遺伝子とオピオイド感受性-癌性疼痛オピオイド治療の将来へ向けて. In: 癌性疼痛	花岡一雄編, 克誠堂 出版, 東京, pp90-93.	2010
曾良一郎, 石原佳奈, 笠原好之, 山本秀子, 池田和隆	中枢刺激薬の分子標的としてのモノアミントランスポーター. In: 実験薬理学 実践行動薬理学	社団法人日本薬理学会編, 株式会社金芳堂, 東京, pp263-271.	2010
池田和隆	人によって違う痛みと鎮痛. In: 緩和医療: 痛みの理解から心のケアまで	小川節郎, 鈴木勉, 池田和隆, 下山直人, 松島英介, 笠井慎也 編, 東京大学出版会, 東京, pp83-120.	2010

小林徹, 池田和隆	GIRK チャネル.	生体の科学 61(5):416-417.	2010
池田和隆	心の分子メカニズムの探索:気持ちよさの生まれ方. In: こころの働きと病・覚醒剤	NPO 法人脳の世紀推進会議編, 株式会社クパプロ, 東京, pp7-44.	2010
池田和隆	痛みと鎮痛における個人差の遺伝子メカニズム. In: 別冊・医学のあゆみ 原始感覚と情動-生体防御系としての情動機構とその破綻	福土審編, 医歯薬出版株式会社, 東京, p38-42.	2010
特許	池田和隆, 笠井慎也, 林田眞和, 樋口進 POMC 遺伝子解析による薬物感受性の評価方法 European Patent Office, 2151496		2012年 04月 25日【成立】
	池田和隆, 西澤大輔, 福田謙一 各種鎮痛関連遺伝子解析による薬物感受性の評価方法 特願 2011-288940		2011年 12月 28日【出願】
	池田和隆, 笠井慎也, 林田眞和, 樋口進 POMC 遺伝子解析による薬物感受性の評価方法 European Patent Office, 109055 (20080425 国内 PCT 出願:PCT-JP2008-058083) (20091023 国内特願:2009-511918)		2011年 10月 17日【成立】
	池田和隆, 西澤大輔, 福田謙一 サイクリック AMP 応答配列結合タンパク質遺伝子解析による薬物感受性および疾患脆弱性の評価方法 特願 2011-217104		2011年 9月 30日【出願】
	池田和隆, 西澤大輔, 福田謙一 アドレナリン受容体遺伝子解析による薬物感受性および疾患脆弱性の評価 特願 2011-191861		2011年 9月 2日【出願】
	池田和隆, 韓文華, 西澤大輔, 福田謙一 電位依存性カルシウムチャンネル遺伝子解析による薬物感受性の評価方法 特願 2010-270630		2010年 12月 3日
	池田和隆, 井手聡一郎, 曾良一郎 ミューオピオイド受容体遺伝子解析による薬物感受性の評価方法 特許登録第 4580924 号		2010年 9月 3日
	池田和隆, 西澤大輔, 福田謙一 アドレナリン受容体遺伝子解析による薬物感受性の評価方法 特願 2010-198319		2010年 9月 3日